Title: Strong environmental memory revealed by experimental evolution in static and fluctuating environments

Authors: Clare I. Abreu[1*], Shaili Mathur[1], Dmitri Petrov[1*]

Affiliations: [1]Department of Biology, Stanford University; Stanford CA, USA.
*Corresponding authors. Emails cabreu@stanford.edu, dpetrov@stanford.edu

Evolution in a static environment, such as a laboratory setting with constant and uniform
conditions, often proceeds via large-effect beneficial mutations that may become maladaptive in
other environments. Conversely, natural settings require populations to endure fluctuating
conditions. A naïve assumption is that the fitness of a lineage in a fluctuating environment is the
time-average of its fitness over the sequence of static conditions it encounters. However,
transitions between conditions may pose entirely new challenges, which could cause deviations
from this time-average. To test this, we tracked hundreds of thousands of barcoded yeast lineages
evolving in static and fluctuating conditions and subsequently isolated 900 mutants for pooled
fitness measurements in 15 environments. We find that fitness in fluctuating environments
indeed often deviates from the expectation based on static components, leading to fitness non-
additivity. Moreover, closer examination reveals that fitness in one component of a fluctuating
environment is often strongly influenced by the previous component. We show that this
environmental memory is especially common for mutants with high variance in fitness across
tested environments, even if the components of the focal fluctuating environment are excluded
from this variance. We employ a simple mathematical model and whole-genome sequencing to
propose mechanisms underlying this effect. Our results demonstrate that transitions in fluctuating
environments have large impacts on fitness and suggest a framework for interpreting change in
unpredictable conditions.

Introduction

The increasing scale of experimental evolution, thanks to new levels of throughput and
parallelism, provides unprecedented opportunities to understand evolutionary dynamics and
mechanisms. Unifying principles have emerged about evolution in a single environment,
including that replicate populations display parallel adaptation in recurrent genes, operons, or
pathways[1–5], with successive mutations having a saturating effect on fitness, resulting in
declining adaptability over time[6,7]. However, laboratory studies have mostly examined static
environments, which have relatively temporally constant and spatially uniform conditions, and
which often select for mutations that exhibit costly fitness tradeoffs if conditions change[8–12].
Such tradeoffs are relevant in natural settings where populations must endure distinct and
intermittently harsh conditions, unlike in continually replenished laboratory environments.
11
Experimental evolution in fluctuating environments, while relatively uncharted, is making strides
toward bridging this gap. One review found that costly fitness tradeoffs arise less often in
heterogeneous than homogenous environments[13]. Evidence also suggests that fluctuating
environments produce more surprising and unpredictable results. At the population scale,
fluctuations have been shown to lead to more variation across evolving replicates[14,15] . At the
individual level, a mutant's fate can be sensitive to the timescale[16] and ordering[17] of fluctuations,

and the molecular basis and fitness of mutations emerging from fluctuating environments can be
different from those in static environments[18,19]. These findings align with theory that predicts that
evolution is less deterministic in fluctuating environments[20,21].
21
The variety of evolutionary outcomes seen in fluctuating environments raises the possibility that
fluctuations render an environment fundamentally different from the sum of its parts. If fitness is
additive in a fluctuating environment, the fluctuations might be decomposed into static
components, and fitness in each component simply averaged to predict the overall fitness in the
fluctuating environment. However, a theoretical study demonstrated a counterintuitive finding
that temporal fluctuations do not always time-average fitness effects[20]. Supporting this result,
there is growing empirical evidence of non-additive fitness in fluctuating environments. For
example, a study of yeast gene-deletion mutants in two fluctuating environments found many
instances of non-additivity, due to both to under- and over-performance in fluctuating
environments[16]. Another study highlighted the fallibility of fitness measurements across
environmental gradients in predicting fitness in fluctuating conditions for a range of organisms,
finding inconsistency between parameters obtained in static environments and performance in
fluctuating environments[22]. While these results suggest that fluctuating environments may be
inherently different from static environments, the mechanisms of non-additivity in fluctuating
environments remain undefined.
37
One possible explanation for why fitness in a fluctuating environment cannot always be
predicted by time-averaging fitness in static environments could be the presence of different
transitions in the fluctuating environment. For example, a batch culture experiment with
fluctuations might transition from condition A to B and from B to A, while the same experiment
with static conditions would transition from A to A and from B to B. In a new environment, the
challenge of transitioning from a state of dormancy to growth manifests as a lag time. Lag times
between growth on different carbon sources were observed in classical studies of diauxic shifts[23],
and have more recently been associated with growth tradeoffs[24–26] and specificity of gene
expression[27]. Even static environments with a single carbon source can introduce transitions,
such as in batch culture experiments where microbes cease growth after depleting the resource,
after which a small amount of culture is diluted into fresh medium, and growth begins again after
the lag time. Experimental evolution of yeast in this sort of environment led to the emergence of
adaptive mutants that gained fitness advantage in the lag phase, and the magnitude of this
advantage was determined by the growth parameters of the previous cycle[28]. Their fitness
advantage was thus accrued during one cycle but realized during the subsequent one. This result
suggests that a mutant's lag time can depend on its pre-conditioning environment and raises the
question of how fitness would change in a fluctuating environment, where the pre-conditioning
environment differs from the environment in the following cycle.
56
Here, we test the hypothesis that non-additivity of fitness in fluctuating environments is due to
differences between transitions in static and fluctuating environments using high-throughput
genome barcoding. We first evolved clonal populations of *Saccharomyces cerevisiae*, each
containing ~500,000 unique barcoded lineages, in static environments, including stressors and
alternate carbon sources, as well as in temporally fluctuating pairwise combinations of these
environments. We tracked barcoded evolutionary trajectories and subsequently selected a pool of
889 mutants for follow-up fitness measurements in all environments. By comparing fitness in

fluctuating environments to the average fitness in the component static environments, we see a
spectrum of non-additivity. Furthermore, frequency trajectories of individual mutants reveal
striking changes in the components of fluctuating environments compared to corresponding
static environments, which are often hidden in the overall fitness measurements to some extent
due to these changes partially canceling each other. We show that these changes reveal a pattern
based in fitness differences across static environments, whereby mutants with larger variance in
fitness are more sensitive to transitions and exhibit more environmental memory, or fitness
change in fluctuating environments. We employ a simple mathematical model and whole-
genome sequencing to propose mechanisms underlying this environmental sensitivity, including
that mutations that cause higher fitness variance are environment-specific and may shift lag
times to a greater extent in fluctuating environments. Our results demonstrate that environmental
transitions alter fitness and fundamentally differentiate static from fluctuating environments.
76

Definitions

Non-additivity: Difference between overall fitness of a mutant in a fluctuating environment and its time-averaged fitness in the component static environments.

Environmental memory: Difference between fitness in an individual component of a fluctuating environment and fitness in a static environment with this component only.

Results

Large dataset acquired through experimental evolution and fitness measurements
To study how evolution and fitness depend on the environment, we began by evolving barcoded
yeast in static and fluctuating environments. We propagated clonal populations containing
~500,000 uniquely barcoded lineages, as developed in a previous study[29], for ~168 generations
in serial batch culture. We carried out evolutions in five static environments and five fluctuating
pairwise combinations of these static environments (**Fig. 1A**, **Table 1**, Methods; we included the
remaining five pairwise combinations in later fitness measurements). Static environments
contained either glucose (*Glu*), *S. cerevisiae*'s preferred carbon source, or less-optimal alternate
carbon sources (*Gal*: galactose, *Lac*: lactate), and in two environments we supplemented the
glucose condition with stressors known to affect *S. cerevisiae* fitness[30,31] (*NaCl*: sodium chloride,
H_2O_2: hydrogen peroxide) (**Table 1**).
12
We tracked changing barcode diversity during evolution and subsequently chose timepoints for
reviving and isolating mutants (Fig. S1). We created a pool of 889 uniquely barcoded mutants
for fitness measurements in all static and fluctuating environments. We added the pool at 5%
relative abundance to a preculture containing the ancestor strain, giving each barcoded lineage a
starting frequency of ~$5*10^{-5}$. We measured lineage frequencies over several cycles (with eight
generations per cycle) to determine fitness relative to the ancestor (**Fig. 1B-D**; Methods). Briefly,
we define fitness as log frequency change, corrected by mean population fitness:
20

$$f_{i+1} = f_i e^{(s-\bar{s})} \tag{1}$$

Figure 1: **Fitness measurements test for non-additivity in mutants evolved from static and fluctuating environments.** **A** We evolved the same barcoded library in five static conditions and five fluctuating pairs of these static conditions. We then isolated 889 mutants from the evolution experiments and performed pooled fitness measurements in all environments, plus five additional fluctuating environments. **B-C** After correcting for mean population fitness, we estimate a mutant's fitness from the slope of its log frequency. **D** The null hypothesis of additive fitness in a fluctuating environment is represented by the average of the slopes in panels B-C. Non-additive fitness might result if transitions in the fluctuating environment alter fitness of a mutant. **E** Non-additivity can be visualized by plotting observed fitness in a fluctuating environment vs. the null hypothesis. Mutants with non-additive fitness will appear above or below the one-to-one line.

23 Here, f_{i+1} represents frequency of a lineage at timepoint $i + 1$, which changes exponentially
24 from timepoint i if its fitness s differs from mean population fitness $\bar{s}$, which we measure with
25 barcoded neutral strains. We infer fitness in fluctuating environments separately for each
26 component and define the overall fitness as the average of the two components.
27
To check the reproducibility of fitness measurements, we included three biological replicates of
each condition. Replicates are strongly correlated (Figs. S2-S3), but we exclude one of the ten
fluctuating environments, *Lac/H₂O₂*, in analyses summarizing environments, because replicates
in this environment are weakly correlated (Fig. S3, Methods), but include it individually (Figs.
S3, S6-S7). To focus on mutants adaptive in their home evolution environments, we exclude
those with fitness less than 0.05 in their home environment, or per-cycle fitness 5% greater than
ancestral fitness, but results do not change for a higher threshold or for no threshold (Fig. S4).
Unsurprisingly, most mutants are adaptive in their home environment: excluding mutants below
the adaptive threshold brings the size of the pool to 695 (78% of the initial pool). Adaptive
mutants evolved in fluctuating environments often have lower fitness in their home environment
compared to mutants evolved in static environments (Fig. S5A). There are exceptions, such as

the Glu/H_2O_2 environment, but highly adaptive mutants often evolved in Lac, H_2O_2, and $NaCl$,
suggesting that static environments with stressors or a non-fermentable carbon source lead to
large fitness gains.

In fluctuating environments, non-additive fitness and memory are common
To determine whether fitness in fluctuating environments is additive, we compare it to fitness in
static environments, where we observe a diversity of behaviors within the pool (Fig. S5). The
null hypothesis of additive fitness in a fluctuating environment follows from the expectation that
mutants will increase or decrease their frequency in a given condition regardless of the previous
condition from which they were diluted. If mutants instead have non-additive fitness, this may
indicate that transitions between different conditions have unique effects on mutants' growth
compared to transitions between the same condition (**Fig. 1D**). Indeed, previous studies have
shown that environmental fluctuations can lead to changes in fitness, either due to changes in the
lag phase of growth[26,27] or other mechanisms[16,32] (**Fig. 1E**).

We find that non-additivity, or deviation from the time-average, is extremely common in our
mutant pool (**Fig. 2A**). For brevity, we highlight three of the ten fluctuating environments, which
include all five static environment components, and which contain examples of mutants both
over-performing and under-performing their time-averaged fitness (see Fig. S6 for the remaining
seven). To distinguish non-additivity from the effects of measurement noise, we compare
observed non-additivity to that expected based on the standard error of fitness in static
environments across all timepoints of biological replicates (a less conservative approach uses just
the standard error across replicates; see Methods). We find that 82% of fitness measurements in
fluctuating environments deviate from the time-average more than expected from measurement
noise. The magnitude of non-additivity is greater than 0.1, or a 10% change in fitness relative to
the ancestor, for 7% of clones, showing that fitness often deviates from additivity by a similar
order of magnitude as median fitness in static environments (0.25).

Even more striking than non-additive behaviors, mutants show large changes in fitness within
components of fluctuating environments that may not be captured by non-additivity. To fully
quantify the difference between static and fluctuating environments, we define *environmental*
memory as the difference between observed fitness in each component of a fluctuating
environment and fitness in the corresponding static environment. We expect zero memory if the
fitness is fully defined by growth during the current cycle, but not if fitness is influenced by (i.e.
'remembers') the previous cycle. We highlight several mutants with varying degrees of non-
additivity that have dramatically different behaviors from expectations based on static
environment fitness, leading to environmental memory (**Fig. 2B-D**). Observed frequency
trajectories in the fluctuating environment (solid lines), excluding the first two cycles, are plotted
against null expectation based on fitness in the component static environments (dashed lines). In
these plots, non-additivity appears as deviation from the overall slope of the dashed line, while
memory appears as deviation from segments of the dashed line in individual components.

For example, the green plots in panel **B** (corresponding to green dots in panel **A**) show examples
of mutants that have nearly additive fitness but exhibit memory, or slight changes within each
component that are not fully captured by non-additivity. Panel **C**, on the other hand, shows

Figure 2: **Non-additivity is common, but it masks even greater fitness changes**. **A** Plotting fitness of the mutant pool in three fluctuating environments against the null hypothesis of the average of their fitness in the static environments reveals many cases of non-additivity, or differences between these two quantities. We highlight particular cases in green, magenta, and blue. **B** Frequency trajectories (mean of three replicates) show that the static null prediction (also calculated from replicate mean) is inaccurate, and that overall non-additivity masks the totality of fitness change, or environmental memory. **C** Frequency trajectories of non-additive mutants show reversals of the outcome expected based on static environment fitness. **D** Frequency trajectories of additive mutants also show reversals, even if the overall trajectory and fitness is similar to what was expected. (Note the finer scale in these plots compared to those in panel **C**.) **E** Non-additivity, or the difference between overall fluctuating fitness and the additive prediction, is plotted for all measurements in nine fluctuating environments (orange). Sampling from prediction uncertainty (Methods) shows that the measurement exceeds noise. **F** Memory, or the difference between fitness in fluctuating environment components and corresponding static environments, is greater in magnitude than non-additivity.

mutants with highly non-additive fitness. In the leftmost plot, a mutant with high fitness in

glucose and low fitness in galactose dramatically reverses its fitness in the fluctuating
components, decreasing in frequency during the glucose cycle and increasing in frequency
during the galactose cycle. The other plots in panel **C** show similarly dramatic fitness reversals.
Fitness reversals even occur for mutants that have additive fitness (panel **D**; note the finer scales
in these plots). In this case, overall trajectories reach the expected frequency, hiding memory in
the overall measurement, since over-performance in one half of the environment cancels under-
performance in the other half. These results show that non-additivity in fluctuating environments
is not only common, but that even in cases where the overall fitness across the whole fluctuating
experiment is similar to the expectation, the component fitnesses often deviate sharply from the
null expectation.
95
Unlike non-additivity, environmental memory includes the difference between observed and null
in both halves of the fluctuating environment. Non-additivity is significant compared to
measurement noise (**Fig. 2E**) and has smaller magnitude than memory (**Fig. 2F**), which also
exceeds measurement noise and extends to roughly twice the range. As before, to estimate
measurement noise, we sample from a normal distribution with a standard deviation equal to the
standard error of predictions (Methods), finding that memory exceeds that expected by
measurement noise in 87% of fitness measurements in fluctuating environments (compared to
82% of non-additivity). The magnitude of memory in either component of a fluctuating
environment is greater than 0.1 for 30% of clones (this is true of non-additivity for 7% of
clones), illustrating that transitions in fluctuating environments lead to radical and widespread
changes in fitness, making fluctuating environments inherently different from a sequence of
static environments.
108
Fitness variance in static environments correlates with memory in fluctuating
environments
Overall patterns of environmental memory reveal a tradeoff in which mutants gain fitness in one
half of a fluctuating environment but lose fitness in the other half. To visualize memory of the
pool in each fluctuating environment, we first plot fitness in pairs of static environments,
coloring mutants by fitness difference (**Fig. 3A**; distance from the equal-fitness dashed line). As
shown in **Fig. 3B-D**, darker shades of blue denote higher fitness in the first environment, and
darker shades of red indicate higher fitness in the second environment. We preserved this
shading when plotting memory, or difference in fitness between each component of a fluctuating
environments and the corresponding static environment. As shown in **Fig. 3E**, the null
hypothesis, or no memory, sits at the origin of these plots. Data not on the origin can still
represent additive fitness if it sits on the $y = -x$ line, because in this case, fitness change in one
half of a fluctuating environment exactly cancels change in the other half. We observe that not
only do the measurements generally deviate from the origin, but that they tend to cluster around
the $y = -x$ line, showing that fitness gains occur in one component of a fluctuating environment
and losses occur in the other and that they often match each other well in magnitude (**Fig. 3F-H**).
This negative trend appears in all ten fluctuating environments (Fig. S6), indicating that
opposing fitness changes that are not fully captured by non-additivity are widespread and
generic. Furthermore, mutants tend to gain fitness in the component corresponding to the static
environment in which they do worse and lose fitness in the component corresponding to the
static environment in which they excel, indicated by the recurrent pattern of blue dots in the

Figure 3: **Fitness in one component of a fluctuating environment influences fitness in the other component**.
A Plotting fitness across a pair of static environments indicates mutants that have higher fitness in the first environment (blue) and mutants that have higher fitness in the second environment (red). **B-D** Fitness of the pool is shown in three pairs of static environments. **E** Plotting the difference of fitness in one component of a fluctuating environment and fitness in the corresponding static environment shows memory, or how fitness changes in fluctuating environments. The $y = -x$ line indicates mutants that change their fitness but are still additive. **F-H** The change in fitness in the two components of fluctuating environments is negatively correlated, indicating that mutants gain fitness in one component and lose fitness in the other. The presence of blue dots in the upper left quadrants and red dots in the lower right quadrants indicates that fitness gains often occur in the environment with lower static fitness and fitness losses often occur in the environment with higher static fitness, suggesting that in a fluctuating environment, fitness is influenced by the previous condition. **I** We define fitness difference between two static environments as the distance from the equal-fitness line. **J-L** Mutants with high mean fitness difference across all ten pairs are farther from the origin in these plots, meaning that they have more environmental memory.

upper left quadrant and red dots in the lower right quadrant. This surprising pattern suggests that
memory is indeed caused by influence of the previous condition.

A possible explanation for one cycle of a fluctuating environment influencing fitness in the next
is that a mutant may evolve to optimize transitions in a particular environment, whether static or
fluctuating. In that case, a lineage's physiological state at the end of one cycle might determine
its transition to lag phase and growth in the following cycle, and thus to different consequences if
the environment in the following cycle is the same or different. For mutants with bigger fitness
differences across static environments, transitions between environments could foreseeably
become more severe, leading to more changes in behavior in a fluctuating environment. Indeed,
the color arrangement in **Fig. 3F-H** suggests that fitness difference across a pair of static
environments may correlate with fitness change when those two environments fluctuate. For
example, dark red and dark blue dots, representing mutants with bigger fitness differences, tend
to sit farther from the origin, indicating that they have more memory. There are some exceptions

to this pattern, such as the dark red and blue dots clustered near the origin in **Fig. 3F**. These are
mutants with large fitness differences in static *Glu* compared to static *Gal*, and which change
their fitness in *Glu/Gal* relatively less than other mutants with similar fitness differences.
However, calculating global mean fitness difference, rather than just local *Glu/Gal* fitness
difference, reveals that these mutants have low global mean fitness difference. Overall, lower
global mean fitness difference translates to less memory in fluctuating environments (or
placement closer to the origin in these plots) (**Fig. 3I-L**).
151
As a more rigorous test of whether global mean fitness difference correlates with memory in a
fluctuating environment regardless of local fitness difference, we calculate partial correlations of
memory in each fluctuating environment with global mean fitness difference, controlling for and
excluding local fitness difference. The partial correlations are positive and significant in nine out
of ten fluctuating environments (all except for *Lac*/H_2O_2), even if static environments
corresponding to the components of the focal fluctuating environment are excluded from the
mean fitness difference (Fig. S7, Methods). This result shows that mutants with high fitness
variance are uniquely sensitive to transitions in fluctuating environments and that a particular
setting is not necessary to identify this characteristic sensitivity.
161
The result that bigger fitness differences in static environments propagate to bigger fitness
changes in fluctuating environments can be summarized by plotting mean fitness difference
across all pairs of static environments against mean fitness change across all fluctuating
environments (**Fig. 4A**), with memory and non-additivity plotted separately in different colors.
We confirm that fitness difference is an appropriate parameter to compare to fitness change by
decomposing it into two independent parameters, both of which contribute toward explaining
variation in fitness change in a general additive model (Fig. S8). While both memory and non-
additivity correlate with mean fitness difference, memory has a nearly one-to-one correlation,
meaning that in fluctuating environments, a mutant will on average change its fitness by an
amount nearly equal to its average fitness difference across static environments. While this is an
average effect, it could manifest in a particular fluctuating environment as a mutant reversing its
fitness, as we saw in the trajectories in **Fig. 2C-D**. Additionally, the negative trends in **Fig. 3F-H**
suggest that mutants compensate for fitness increases in one component of a fluctuating
environment with fitness decreases in the other component, suggesting that fitness reversals
could be common.
177
Mutants reverse their fitness often in fluctuating environments
To determine how often fitness reversals occur, we asked which of the two static environments is
a better predictor for fitness in a component of a fluctuating environment (for example, is static
Glu fitness or static *Gal* fitness better at predicting the *Glu* component of fitness in fluctuating
Glu/Gal?). We categorize the results as the same static environment being a better predictor (e.g.
static *Glu* better predicts fluctuating *Glu*), the other static environment being a better predictor
(e.g. static *Gal* better predicts fluctuating *Glu*), or a borderline case if the prediction and result
vary by less than 0.05 (we choose this threshold since the median standard error of fluctuating
component fitness is 0.046). Overall, the other static environment is a better predictor than the
same static environment in 29% of cases. In the top 10% of mutants ranked by mean fitness
difference, this statistic increases to 39%, while for the bottom 10% it decreases to 11% (**Fig.
4B**). As a control, we predict fitness in static environment replicates with either other replicates

of the same static environment or with replicates of another static environment; as a second
control, we predict fitness in fluctuating environment component replicates with other replicates
either from the same component or the other component (**Fig. 4C**). We expect both controls to be
noisy, since they use replicates rather than averages of replicates, and the second control to be
especially noisy, since fluctuating environment component measurements contain half as many
timepoints as static environment measurements. Indeed, the second control represents the
noisiest upper bound (13%) as opposed to the first control (2%). In either case, fitness reversals
in fluctuating environments occur more often than would be expected due to measurement error.
Most strikingly, mutants with the biggest fitness differences reverse their fitness most often,
despite these reversals requiring greater change.
200

Figure 4: **Fitness reversals in fluctuating environments are common and recapitulated by a simple model of lag time evolution. A** Mean memory, or mean absolute difference between fitness in a static environment and fitness in the corresponding fluctuating component, correlates strongly with mean fitness difference across static environments. This nearly one-to-one correlation indicates that on average, mutants change their fitness in a fluctuating environment by an amount nearly equal to their fitness difference across static environments. Mean non-additivity, or mean difference between fluctuating fitness and average static fitness, is a lower bound of mean memory. **B** The frequency of fitness reversals in fluctuating environments is quantified by the fraction of the time that the other static environment fitness is a better predictor of fitness in half of a fluctuating environment, or 29% of overall cases. The top 10% of mutants by mean fitness difference more often reverse their fitness. **C** Fitness in one replicate of a static environment is better predicted by a replicate of another static environment than another replicate of the same static environment 2% of the time. The same is true for replicates of fluctuating environment components in 13% of cases, a noisier control because it contains half as many timepoints. **D** A simple model reproduces the correlation pattern shown in panel **A**. **E** The model reproduces the difference between mutants with high and low mean fitness difference shown in panel **B**. **F** The ancestor dominates growth in the model, where mutants' fitness is determined by comparative lag (dis)advantages, which shift toward the mean of their distribution across static environments when the environment fluctuates.

To explain potential mechanisms of fitness change and reversal in fluctuating environments, we
employ a simple mathematical model describing our fitness measurements (**Fig. 4D-F**, Methods,
Supplementary Note 1). We assume that due to a starting frequency orders of magnitude higher
than the mutants, the ancestor dominates growth and sets the time to resource depletion, before
which mutants increase or decrease their frequencies (**Fig. 4F**). We assume that sensitivity to
transitions is more likely to affect lag than growth[33], and that mutants' fitness changes are
primarily due to changes in lag time compared to the ancestor, but that they retain the ancestor's
growth rate for a given environment. Mutants with more variance in fitness across static
environments have a wider range of lag (dis)advantages. Guided by partial correlations between
memory and global fitness differences as described above (Fig. S7), we also assume that in a
fluctuating environment, mutants shift their lag (dis)advantages proportionately to the width of
the distribution across all static environments, regardless of local fitness difference in the pair of
fluctuating environments. Additionally, we eliminate differences in yield across environments as
a cause of non-additivity and memory (Supplementary Note 1). These assumptions recapitulate
the result that fitness change in fluctuating environments increases with mean fitness difference
across static environments, and that memory exceeds non-additivity (**Fig. 4D**). We fit the overall
result that the other static environment is a better predictor for fitness in a fluctuating
environment component in 29% of cases by setting the shift in lag (dis)advantage to 80% of the
distance toward the mean of a mutant's distribution. The model also recapitulates that mutants
with more variance in fitness reverse their fitness more often, since their larger fitness variance
means the results are less likely to be borderline cases (**Fig. 4E**). These results suggest that lag
time evolution could be a cause of fitness reversals in fluctuating environments.
223

Environment	Composition	Evolved	Fitness measured	Multi-hit genes (no. of hits)	
Static:				**Homozygous SNPs**	**Copy number variation**
Glu	SC + 1.5% glucose	•	•	ZAP1 (2)	
Gal	SC + 1.5% galactose	•	•	ISN1 (2)	
Lac	SC + 0.7% lactate	•	•	-	
H_2O_2	SC + 1.5% glucose + 2.5mM hydrogen peroxide	•	•	-	
NaCl	SC + 1.5% glucose + 0.5M sodium chloride	•	•	tT(AGU)B (2), PAU8 (2)	ENA1 / ENA2 / ENA5 (18)
Fluctuating:					
Glu/Gal		•	•	ISN1 (2)	
Glu/Lac		•	•	ARN2 (2), GRX3 (2), ISN1 (2)	
Glu/H_2O_2		•	•	AFT1 (14), BOL2 (7), SNF5 (2), ZRT3 (2), RRP5 (2), ISN1 (2)	
Glu/NaCl		•	•	-	ENA1 / ENA2 / ENA5 (7)

H₂O₂/NaCl		•	•	-	ENA1 / ENA2 / ENA5 (4)
Gal/Lac			•	no evolution	no evolution
Gal/ H₂O₂			•	no evolution	no evolution
Gal/NaCl			•	no evolution	no evolution
Lac/ H₂O₂			•	no evolution	no evolution
Lac/NaCl			•	no evolution	no evolution

Table 1: **Descriptions of static and fluctuating environments.**

Environment-specific mutations may increase fitness variance and memory
In search of a genetic mechanism for significant lag time changes in mutants with high fitness
variance in fluctuating environments, we whole-genome sequenced a subset of the mutant pool,
obtaining 352 sequences (Methods). Most of our mutants contain only unique mutations, but
several of the environments produced mutants with mutations in the same genes. Of such multi-
hit genes, only five experienced more than five hits (**Table 1**). These five genes include:
ENA1/ENA2/ENA5, which in tandem encode a sodium efflux pump known to be necessary for
resuming growth following transfer into high salt[34] and here displayed repeated copy number
amplifications during evolution in high salt (18/43 *NaCl* mutants, 7/27 *Glu/NaCl* mutants, 4/34
H₂O₂/NaCl mutants); additionally, the genes AFT1 and BOL2, part of an iron-sensing and
uptake pathway[35,36], had SNPs in 28/72 mutants evolved in *Glu/H₂O₂* (including heterozygous
SNPs). We also assayed 359 mutants for ploidy (Methods), as the DNA barcoding process
sometimes induces self-diploidization[3], categorizing 161/359 mutants as diploid. To narrow this
list to pure diploids lacking other adaptive mutations, we eliminated the iron- and salt-related
mutations from the identified diploids, leaving 148 "pure" diploids (note that we do not eliminate
diploids with low-frequency mutations, so we cannot be sure how much of their fitness is due to
these mutations vs. diploidy alone). In summary, highly repeated mutations fall into three
groups: two that appear to have evolved to sense particular environments, and environment-
nonspecific diploids.
244

We categorize these groups of mutations by plotting their mean fitness difference and mean
memory. In agreement with a previous study showing that pure diploids benefit from low-
variance, modest gains in fitness across a range of environments[12], we find that diploids have
moderate fitness variance and memory (**Fig. 5A**). Iron-sensing mutants have high fitness
variance and memory, while salt-sensing mutants have moderate fitness variance and memory.
However, when the latter group is subdivided into those with high- and low-copy number
amplifications, we see that more copy number amplifications lead to comparatively higher
fitness variance and memory (**Fig. 5B**), showing that more extreme forms of this environment-
sensing mutation increase fitness variance and memory. These results offer a mechanistic
explanation of the adaptive strategy for a subset of mutants: having an ability to sense iron or salt
in their environment gives them a high fitness advantage in the relevant environment but likely
comes with a costly tradeoff, such as an increased lag, when the environment changes.

Figure 5: **Environment-sensing mutations are associated with higher fitness variance and memory. A** Three classes of mutants are plotted by mean fitness difference and mean memory (error bars are standard error of the mean). Diploids have moderate mean fitness variance and memory. Iron-sensing mutants have higher fitness variance and memory, while salt mutants do not. **B** Subdividing the pool of salt pump mutants into high and low copy number amplification reveals that more amplification leads to more higher mean fitness variance and memory. A single SNP mutant was observed in the same set of genes as the copy number variants.

Discussion

Natural environments are consistently perturbed by environmental fluctuations, and the
amplitude and frequency of some fluctuations are expected to increase[37]. An additional concern
is the ability of emerging viruses, drug-resistant bacteria, and mutating cancer cells to adapt to
ever-shifting host environments. Predicting evolution in fluctuating environments is therefore
important, but experiments have thus far mostly focused on adaptation to static conditions. Our
study addressed the question of how static environments change when fluctuations are added by
comparing evolution and fitness in both types of environments. We find that fitness in a
fluctuating environment can be profoundly altered by transitions that are lacking in static
environments, and that these changes are at least partially concealed by measurements of fitness
across the complete cycle of fluctuating environments, due to a tradeoff that causes fitness to
increase in one component of a fluctuating environment and decrease in the other component.
We show that mutants with high fitness variance across environments are uniquely sensitive to
transitions, reversing their fitness most often in fluctuating environments, even when this
variance is measured in conditions unseen in the fluctuating environment.
15
The observation that fitness in one component of a fluctuating environment influences fitness in
the other component suggests that the physiological state of a population at the end of its growth
cycle influences its growth in the following cycle. This result agrees with previous measurements
of yeast in static environments, which showed that fitness accrued in the final stage of the growth
cycle is realized during the lag phase of the following cycle for a set of highly adaptive nutrient-
sensing pathway mutants evolved from a minimal glucose-limited medium[28]. Subsequent
measurements revealed that these mutants are deleterious when the media is supplemented with

0.5M sodium chloride[12]. Based on our findings in the current study, we would expect that these
glucose-sensing mutants' large fitness variance would lead to environmental memory in
fluctuations. To test this expectation, we grew those mutants in an environment fluctuating
between optimal and high-salt conditions, indeed finding that their frequencies crashed in
optimal conditions after exposure to high-salt conditions (Fig. S9). These results echo the
environmental memory presented in this study, where we evolved and measured mutants in a
complex instead of minimal medium. The agreement across differing types of media and
fluctuations (both stressors and alternate carbon sources) suggests a general mechanism that
causes strains, particularly those with environment-sensing mutations, to change their lag phase
in one condition after exposure to another. As a result, predicting evolution in fluctuating
environments based on static environments may be difficult.
34
While extrapolating from static to fluctuating environments may be difficult, our study also
suggests that changes in fluctuating environments may be governed by a global structure
whereby a strain's fitness variance in static environments predicts its fitness change in
fluctuating environments. Another phenomenon apparently detrimental to predicting evolution,
epistatic interactions between mutations, was elucidated by global patterns. Originally observed
in empirical studies[6,38,39], these patterns illuminated a structure whereby a strain's background
fitness predicts the fitness effect of a new mutation, leading to unifying models[40] that were
recently applied to a novel context of predicting ecological function through species
interactions[41]. The success of global patterns in interpreting interactions between mutations and
between species bodes well for the problem outlined here of interpreting interactions between
environments.
46
To further generalize the trend that we demonstrate between fitness variance in static
environments and fitness change in fluctuating environments, more investigation is needed
beyond our comparison of a set of recently evolved mutants to their ancestral strain. For
example, other systems such as ecological communities approaching stable states should be
examined for the existence of this pattern. We grew the mutants in our pool from very low
relative frequencies in the presence of a majority of ancestor cells to prevent frequency-
dependent effects, but studies of frequency-dependent interactions in communities have also
shown evidence of additive[42,43] and non-additive[44,45] effects. Our study also did not address
whether the timing and order of fluctuations affects the final results, nor did we examine
environments fluctuating between more than two conditions. In all cases, mathematical models
offer a promising tool for interpreting how outcomes may be altered by fluctuation- and
frequency-dependent effects[46].
59
Predicting the effects of increasing environmental fluctuations is an important challenge in
ecology and evolutionary biology. Here we have demonstrated that the widespread existence of
memory in fluctuating environments makes them qualitatively different settings than just the sum
of their component static environments. This memory manifests as fitness changes in fluctuating
environments, likely because the growth environment in one cycle influences lag phase in the
following cycle. While transitions from one cycle to the next occur in both static and fluctuating
environments, the latter impose more severe shifts, especially for mutants with greater fitness
variance across static environments, possibly due to their environment-sensing mutations. This
trend reveals that while null expectations about fitness in fluctuating environments are

inadequate, their failure illuminates biological patterns and opens avenues for future
investigation.

Methods

Evolution experiments
<u>Founding populations</u>
The barcoded yeast population used in this study was constructed in a previous study[29] and
contains around half a million uniquely barcoded transformants. The library was created from an
S288C derivative, BY4709, in which a 'landing pad' was inserted into a neutral location in the
yeast genome that allows for high-frequency, site-specific genomic integration of plasmids via
the Cre–loxP recombination system. We created new stocks of this library and sequenced the
barcode region to confirm a high diversity of ~500,000 barcodes before beginning batch culture
experiments.
10
<u>Evolution batch cultures</u>
We evolved the population in ten evolution conditions (Table 1), which are composed of static or
fluctuating regimes of five environments: *Glu, Gal, Lac, H₂O₂, NaCl*. We began evolution
experiments by growing a single overnight 15-mL preculture containing the founding population
in synthetic complete (SC) medium with 1.5% glucose (*Glu* media). We then transferred 400ul
of this pre-culture to all evolution environments, containing 100mL of various media types. All
cultures were evolved by serial batch culture in 500-ml Delong flasks (Bellco) at 30°C and 223
r.p.m. We diluted cultures every 48 hours 250X by inoculating 400ul of culture into 100 ml fresh
medium. The SC base medium (per liter: 2g US Biological Drop-out Mix Complete, 6.7g RPI
yeast nitrogen base) was supplemented with either 1.5% D-glucose (RPI) (*Glu*), 1.5% D-
galactose (RPI) (*Gal*), or 0.7% sodium DL-lactate (Sigma) (*Lac*). In the remaining two
environments, we supplemented the *Glu* medium with either 0.5M sodium chloride (RPI) (*NaCl*)
or 2.5mM hydrogen peroxide (Ward's) (*H₂O₂*). All media was prepared in the weeks prior to
experiments and stored at 4°C until the final days before use, when we pre-warmed it in a 30°C
incubator. We tested hydrogen peroxide concentration in the *H₂O₂* media after weeks of storage
using the MQuant Peroxid-Test kit (for 1-3-10-30-100 mg/l) and found that the hydrogen
peroxide had reduced to ~60% of the original concentration. After learning of this effect, we
added hydrogen peroxide directly to media during the dilution/transfer step, rather than during
media preparation, in the fitness measurements (described below). We evolved in all fluctuating
environments with two replicates except for *Glu*/*Gal*, which had one replicate, and we evolved in
static environments with one replicate, except for *Glu* and *NaCl*, which both had two replicates.
We propagated the cultures for 20-23 growth cycles and stored 2 ml saturated culture by
aliquoting 1 ml cells into 1 ml 50% glycerol in two Eppendorf tubes and storing at -80°C. The
remainder of the cell culture was centrifuged and the cell pellet was resuspended in 2 ml sorbitol
solution (0.9 M sorbitol, 0.1 M Tris-HCL pH 7.5, 0.1 M EDTA pH 8.0), aliquoted into two
Eppendorf tubes and stored at −20°C for genomic extraction.
37
Barcode amplicon and whole-genome sequencing
<u>DNA extraction</u>
We extracted genomic DNA from each sample using the Norgen Fungi/Yeast Genomic DNA
Isolation 96-Well Kit, supplemented with lyticase. In collaboration with Norgen sales

representatives and research scientists, we modified this kit to use 96-well plates for all steps.
We extracted DNA using ~10% of our spun-down samples, or about 10^9 cells. Final DNA yield
was assayed with a Biotek fluorescent plate reader and averaged ~4ug per sample.
45
<u>Barcode amplicon library preparation and sequencing</u>
To track changes in barcoded lineage frequency over the course of evolution, we conducted PCR
amplification of the barcoded region with a two-step protocol. The primers used in the first step
of amplification are the same as described in (Kinsler et al), but with a different amplification
protocol. In both steps of PCR, we performed three reactions per sample in the standard Q5
master mix (per 50ul reaction):
•   10ul 5X Q5 Reaction Buffer
•   1ul 10 mM dNTPs
•   2.5ul 10uM forward primer
•   2.5ul 10uM reverse primer
•   0.5ul Q5 High-Fidelity DNA Polymerase
•   DNA
•   Nuclease-free water (to 50ul)
In the first step, we aimed for 100ng of genomic DNA template (or approximately 1/1000 of
total DNA in a saturated 100-ml experiment) per reaction, but since we added equal volumes to
all reactions, actual amount was sometimes as low as 25ng or as high as 250ng. For the first
cycle, we amplified the barcode region with the following program:
1. 98°C for 30 seconds
2. 98°C for 10 seconds
3. 62°C for 20 seconds
4. 72°C for 30 seconds
5. Repeat steps 2-4 29X for a total of 30 cycles
6. 72°C for 3 minutes
7. 4°C hold
After checking amplification for ~25-50 randomly chosen reactions on a gel, we then diluted the
step 1 product 500X into the second reaction, by diluting 25X into nuclease-free water, and then
20X into the same master mix listed above. The only difference between step 1 and 2 master mix
was the primers—we used IDT for Illumina UD Indexes in the second step (https://support-
docs.illumina.com/SHARE/AdapterSeq/Content/SHARE/AdapterSeq/Nextera/UDIndexesSeque
ncesNXT.htm), with unique pairs of forward and reverse primers for each reaction to avoid error
from index hopping or template switching (Kinsler et al 2023). We then ran the following
program to attach the IDT indices to barcode amplicons:
1. 98°C for 30 seconds
2. 98°C for 10 seconds
3. 62°C for 20 seconds
4. 72°C for 30 seconds
5. Repeat steps 2-4 11X for a total of 12 cycles
6. 72°C for 3 minutes
7. 4°C hold
We again checked for amplification of randomly chosen reactions on a gel and then pooled step
2 product across the three reaction replicates, and then pooled samples in groups of 96 using 10-
15ul per sample. We then performed a standard PCR purification protocol in one column per

sample, eluting into 100ul nuclease-free water, and measured concentration with a Qubit
fluorometer. Final concentration ranged from ~100-200 ng/ul. Our samples were then sent to
Admera Health (https://www.admerahealth.com/) for quality control (qPCR and either
Bioanalyzer or TapeStation) and sequencing. We used 2x150 paired-end sequencing along with
index sequencing reads on Illumina machines using patterned flow cells (either HiSeq X or
NovaSeq X). All amplicon samples were sequenced with at least 20% genomic DNA spiked in
(either whole genomes or phi-X) to ensure adequate diversity on the flow cell.

<u>Whole-genome library preparation and sequencing</u>
We prepared libraries with the Illumina DNA Prep kit (ILMN DNA LP (M) Tagmentation (96
Samples, IPB)). We followed the manufacturer's protocol, with the exception that we used 1/5
the normal volume of reagents, allowing us to use the kit for five times the normal amount of
samples, based on lab members' and others' success with this modification to the protocol. We
sent our libraries to Admera Health (https://www.admerahealth.com/) for quality control (qPCR
and either Bioanalyzer or TapeStation) and sequencing. We used 2x150 paired-end sequencing
along with index sequencing reads on NovaSeq X Illumina machines.

Tracking lineage dynamics during evolution
<u>Processing amplicon sequencing data</u>
Using BarcodeCounter2[47], sequencing reads from evolution timepoints were first aligned to the
reference barcode region using Bowtie2. Based on this alignment, both condition and lineage
barcodes were extracted. Reads were split into different files based on their multiplexing tags,
which were unique for each sample. We then clustered reads using Bartender[48] with Hamming
distance 2 and seed length 8. Examples of evolution trajectories can be seen in Fig. S1 and full
barcode count data for all evolution experiments is available on Dryad (link TBD). To isolate
mutants, we chose a single timepoint from each experiment with a high diversity of adaptive
trajectories, or where the most adaptive lineages reached a frequency of ~10^{-3}, for thawing and
cell-sorting.

Isolation, selection, and pooling of mutants for further analysis
<u>Cell-sorting of thawed stocks</u>
After choosing timepoints from each evolution experiment, yeast clones were isolated from
frozen samples. Frozen stock (~50ul, containing ~5×10^6 cells) was diluted into 500 µl PBS plus
1ul propidium iodide and used for flow cytometry sorting. Single cells were sorted into 96-well
plates with 100ul YPD medium in each well. Five plates of cells (480 cells) were sorted for each
evolutionary replicate (80 plates in total). Sorted cells were grown at 30°C for three days without
shaking, and reached saturation by day three. Saturated cell culture (~5ul) was removed from
each well and inoculated into a different 96-well plate, with 100ul fresh YPD medium in each
well. These replicated plates were grown at 30°C for two days without shaking to reach
saturation, and used for subsequent barcode identification. The remainder of the saturated cell
culture was mixed with 100ul 50% glycerol and stored at −80°C.

<u>DNA barcode identification by Metagrid</u>
High-throughput barcode sequencing of the plates of isolated and grown cells was performed as
in (Li et al), with the exception that cell lysis was performed by boiling 20ul of saturated cell
culture in 20mM NaOH. After sequencing, some wells were eliminated due to low coverage, and

of those remaining, we eliminated those with less than 60% of reads mapping to the top barcode
and those where the number of reads of the top barcode were equal to less than 1.5 times the
number of reads of the next barcode.
137
<u>Selection and pooling of mutants</u>
From the ~3700 wells in which we had identified single barcodes per well, we selected 889
clones for our mutant pool. We chose mutants with high fitness in their evolutionary trajectories
based on Fitmut2[49] inferences and ensured that no overlap existed between barcode sequences.
We selected mutants meeting any of the following criteria: (i) their inferred fitness from FitMut2
was greater than 0.51 (ii) the probability that they were adaptive was greater than 0.5, (iii) they
reached a final frequency of 10^{-5} (iv) they reached a frequency of 10^{-5} at the 6 cycles before the
final one (to account for environments where a small number of barcoded lineages took over the
population, reaching frequencies close to 1). Additionally, we added 10 randomly selected
weakly adaptive mutants (defined to have probability adaptive between 0.125 and 0.5). On
average, 89 clones were selected from each evolution condition, but counts ranged from a
minimum of 40 (H_2O_2) to 120 (*Glu*, *Glu*/H_2O_2 and *Glu*/*NaCl*) clones per condition. We revived
cultures from frozen glycerol replicate plates and grew for two days in fresh SC medium before
pipetting ~10ul of each selected barcode into subpools, which we then consolidated into a final
pool, weighting volumes to aim for equal frequency of all barcodes.
153
Fitness measurement experiments
Fitness measurement experiments were performed as described previously[3,12,28], where growth
competitions were set up between a pool of barcoded mutants and a reference strain. The change
in the frequency of each barcode over time reflects the fitness of the adaptive mutant possessing
that barcode, relative to the reference strain, although certain protocol details were changed. We
grew three separate overnight pre-cultures in 25 mL SC + 1.5% glucose for each round of
measurements: 1) ancestor reference strain, 2) mutant barcode pool, 3) barcoded neutral strains
with fitness equal to the ancestor. We then mixed these cultures at 93:5:2 ratio (correcting for
measured OD). This ratio allows for mutants to compete against the ancestor rather than
competing against each other, helps to minimize the change in average fitness throughout the
competition experiment, and reduces the effect of any frequency-dependent fitness effects as
well as any fitness-affecting interactions among the strains that may occur. We then inoculated
400 mL of this mixed culture (~$5*10^7$ cells) into 100 mL of fresh media in 500 mL DeLong
flasks. As in the evolution experiments, this culture was then grown at 30°C in an incubator
shaking at 223 RPM for 48 hr. After 48 hr of growth, 400ul of saturated culture was transferred
into fresh media of the same type. In static environments, we transferred the serial dilution four
more times, yielding five cycles over which to measure the rate at which a barcode's frequency
changed, and in fluctuating environments, we transferred five more times for six cycles.
172
173   After each transfer of 400ul, the left-over culture was frozen for sequencing barcodes later. To
174   prepare this culture for freezing, it was transferred to 50 mL conicals, spun down at 3000 rpm for
175   5 min, resuspended in 2 mL of sorbitol freezing solution (0.9 M sorbitol, 0.1 M Tris-HCL pH
176   7.5, 0.1 M EDTA pH 8.0), aliquoted into two wells of a 2-mL 96-well plate, and stored at -20°C.
177
We performed DNA extraction and library preparation with the same protocol described above
for evolution experiments, with one exception: we used restriction digest to cut the ApaLI

restriction site in the middle of the reference strain's barcode region. This step prevents the
ancestor from being sequenced, affording greater sequencing coverage to the mutant pool. We
performed the digest after DNA extraction and again after the second step of PCR, adding 1-2ul
ApaLI (NEB #R0507L) and 5ul 10X Cutsmart to 43-ul samples and incubating at 37°C for at
least 2 hr (or up to overnight).
185
186   **Fitness inferences and data analysis**
187   Processing amplicon sequencing data
188   As with the sequencing data from evolution experiments, we first extracted barcodes with
189   BarcodeCounter2[47]. We then used BarcodeCounter2 to map extracted barcodes to our known list
190   of mutants in our pool.
191
192   Fitness inference
193   We inferred fitness of mutants by solving Eqn. 1 for all mutants at each timepoint for each
194   condition and replicate: $f_{i+1} = f_i e^{(s-\bar{s})}$, where f_{i+1} represents frequency of a lineage at
195   timepoint $i + 1$, which changes exponentially from timepoint i if its fitness s differs from mean
196   population fitness $\bar{s}$.
197
198   We measure mean population fitness at each timepoint using ~50 uniquely barcoded neutral
199   strains, which were isolated and identified as neutral, or having fitness equal to the ancestor, in
200   ref TBD. Since we set the ancestor's fitness to zero, we can solve for mean fitness by plugging
201   the neutral strains' frequencies n into Eqn. 1:
202

$$n_{i+1} = n_i e^{(0-\bar{s})} \tag{2}$$

204
205   Solving for mean population fitness $\bar{s}$ gives:
206

$$\bar{s} = -\ln\left(\frac{n_{i+1}}{n_i}\right) \tag{3}$$

208
209   We take the average of Eqn. 3 over all neutral lineages at each timepoint $i + 1$ of each condition
210   and replicate, and plug the result into Eqn. 1 to determine fitness of each mutant:
211

$$s = \ln\left(\frac{f_{i+1}}{f_i}\right) + \bar{s} = \ln\left(\frac{f_{i+1}}{f_i}\right) - \ln\left(\frac{n_{i+1}}{n_i}\right) \tag{4}$$

213
We then average this fitness across all timepoints for a given mutant/replicate/condition. We
exclude the first timepoint, corresponding to the initial mixture of ancestor, neutrals, and mutants
described above, from all estimates. Static environment fitness measurement experiments include
five subsequent timepoints, while fluctuating environment fitness measurement experiments
include six subsequent timepoints, in order grow the population in three cycles of each condition.
In fluctuating environments, we exclude the second timepoint in addition to the first, assuming
the first two cycles are a sort of 'pre-culture,' with the exception of two fluctuating environments
(*Glu-H₂O₂* and *Lac-H₂O₂*) in which the barcoded pool reached a frequency greater than 50% in
the final timepoint (Fig. S10). In those conditions, we excluded the final timepoint and included

the second timepoint. Additionally, we excluded the final timepoint from the static environment
H_2O_2, for the same reason.

In static environment inferences, we average fitness as calculated by Eqn. 4 across all timepoints,
while in fluctuating environment inferences, we infer fitness separately for each of the two
components and define the overall fitness as the average of the two components. Each
component consists of two different timepoints, and we average these two estimates. In both
static and fluctuating environment inferences, we then average fitness across all relevant
timepoints in all three replicates to get a mean fitness estimate for each mutant.

We calculate standard deviation of each mutant's fitness by assuming normally distributed errors
and minimizing the negative log likelihood of a Gaussian distribution using the `minimize`
function from the `scipy.optimize` module in Python, with the `L-BFGS-B` (Limited-memory
Broyden-Fletcher-Goldfarb-Shanno with Box constraints) optimization method, setting initial
guess for mean fitness to 0 and standard deviation to 0.1. To calculate standard error of each
mutant's fitness, we bootstrapped this minimization with 250 trials, sampling with replacement
from all fitness measurements across all timepoints.

In rare cases, some mutants were not detected at all timepoints, either because they went extinct
or just because no counts were sampled. We exclude such timepoints from our fitness inferences.
In the case of static environments, if a mutant reached a detectable frequency at a later timepoint,
we include this timepoint, dividing Eqn. 4 by the number of cycles between detectable
frequencies. This is not possible in fluctuating environments because inferring fitness separately
for the two fluctuating environment components requires measuring frequency change between
consecutive timepoints only.

<u>Defining adaptive fitness</u>
In order to focus our analyses on adaptive mutants, we chose an adaptive threshold of 0.05 and
excluded mutants with fitness below this threshold in their home evolution environment. This
threshold was based on the flattening of the curve in Fig. S4A. However, no change to the final
results happens if no threshold is used, or if an adaptive threshold of 0.1 is used (Fig. S4B).

<u>Excluding data based on replicate-to-replicate correlations</u>
We compared replicate-to-replicate correlations of fitness to identify highly variable mutants or
conditions (Figs. S2-3). Correlations are strong, with Pearson r averaging 0.92 for static
environments and 0.93 for fluctuating environments. We exclude the Lac/H_2O_2 environment
from this average, because fitness is weakly correlated in this environment—during experiments,
growth in H_2O_2 after transfer from Lac was not visible for >24 hours, suggesting long lag times
or mortality. We exclude the Lac/H_2O_2 environment from analyses summarizing environments,
but include it individually (Figs. S3, S7-S9). Some mutants have highly variable fitness across
replicates, such as the extreme differences seen across $NaCl$ replicates (Fig. S2). We exclude 12
mutants with mean replicate-to-replicate fitness difference greater than 0.2 from analyses
summarizing environments in **Fig. 2** and **Fig. 4**, but results do not change if they are included
(Fig. S11).

<u>Comparison of measurement noise to non-additivity and memory</u>

To distinguish non-additivity and memory from the effects of measurement noise, we compared
observed non-additivity and memory to that expected based on measurement uncertainty alone,
as shown in **Fig. 2E-F**. To calculate the percentage of measurements of non-additivity (or
memory) with magnitude greater than would be expected from measurement noise alone, we
looped over all measurements of non-additivity (or memory), comparing the observed fitness to a
sample from a normal distribution with mean equal to the fitness based on static environments
and standard deviation equal to the standard error of this static environment fitness. The larger
the error, the more likely that sampling this distribution will produce larger values of non-
additive fitness (or memory) due to error alone. We repeated this loop 500 times and found the
average percentage of cases in which the magnitude of non-additivity (or memory) is greater
than the noise sample magnitude. In the more conservative version of this comparison (discussed
in the main text), we used the bootstrapped standard error of fitness across all timepoints within
all replicates, whereas a less conservative version of this comparison uses the standard error
across replicates. The latter comparison has lower uncertainty (88% of cases of non-additivity
have magnitude greater than the noise-induced sample magnitude vs. 82% in the former
comparison; 92% of cases of memory have magnitude greater than the noise-induced sample
magnitude vs. 87% in the former comparison), because the standard error across replicates
averages over an entire frequency trajectory rather than including noise within each trajectory.
287
<u>Partial correlation of fitness differences and fitness change</u>
We calculated partial correlations of memory in a given fluctuating environment with mean
fitness difference, controlling for local fitness difference in the relevant pair of static
environments, using the `partial_corr` method in the `pingouin` library in Python. We first
calculated mean fitness difference across nine out of ten pairs of static environments, excluding
the relevant pair. For example, in the case of memory in the *Glu*/*Gal* fluctuating environment, we
excluded fitness difference between static *Glu* and static *Gal*. The partial correlation is positive
and significant in nine out of ten fluctuating environments (Fig. S7A; the exception is *Lac*/H_2O_2,
where biological replicates are very weakly correlated).
297
We then excluded more pairs to see if memory in a given fluctuating environment is correlated
with fitness difference across unrelated pairs of environments, controlling for fitness difference
across the local pair of static environments. For example, we calculated the partial correlation of
Glu/*Gal* memory with mean fitness difference across pairs composed of *Lac*, H_2O_2, and *NaCl*,
controlling for *Glu*/*Gal* fitness difference. The correlations remain positive and significant in this
case (Fig S9B), showing that we can expect more memory in unseen environments based on
greater variance in fitness measured elsewhere.
305
Mathematical model of fitness measurements
Fig. 4D-E shows the results of a mathematical model of our fitness measurements. In this model,
which is derived in Supplementary Note 1, we assume that the ancestor dominates the population
by having a starting frequency many orders of magnitude higher than the mutants, and so sets the
time to resource depletion, before which mutants can increase in frequency. Mutants have
positive fitness if they grow faster than the ancestor or during the ancestor's lag time (Eqn. 9,
Supp. Note 1):
313

$$s_i = r_i \Delta T_L + \left(\frac{r_i - r_a}{r_a} \right) \ln(\text{DF}) \tag{5}$$

Here, s_i is the fitness of mutant i, r_i its exponential growth rate, ΔT_L the difference between
mutant's lag time and the ancestor's lag time (if this quantity is positive, the mutant has a lag
advantage over the ancestor), r_a the ancestor's exponential growth rate, and DF is the dilution
factor at each timepoint (or 250 in our experiments).

We further assume that mutants retain the ancestor's exponential growth rate in a given
environment, and fitness is determined only by lag (dis)advantage:

324
$$s_i = r_i \Delta T_L \tag{6}$$

326 Mutants with high fitness variance have bigger differences in fitness across environments and so
327 have wider distributions of lag (dis)advantage across environments.

329 Memory in a fluctuating environment component, or the difference between fitness in fluctuating
330 and static conditions, is determined by the difference in lag (dis)advantage between conditions:

332
$$s_{if} - s_{is} = r_i \left(\Delta T_{Lf} - \Delta T_{Ls} \right) \tag{7}$$

The mutant will have environmental memory if its lag (dis)advantage shifts in the fluctuating
environment. Since the ancestor's lag time likely changes in fluctuating conditions, memory will
happen if the mutant's lag time changes differently than the ancestor's in fluctuating conditions.
For example, if the ancestor's lag is four hours in the static *Glu* and six hours in the *Glu*
component of *Glu*/*Gal*, while the mutant's lag is three hours and five hours, respectively, there is
no memory. But if the mutant's lag is three hours in both conditions, memory is nonzero and
positive, meaning the mutant overperforms in fluctuating *Glu* compared to static *Glu* fitness.

To simulate fitness measurements in static and fluctuating environments, we first sampled
ancestor growth rates across five static environments from a random uniform distribution [0.3,
0.5]. We then sampled lag advantages for a pool of 700 mutants by first sampling each mutant's
"home" (or evolution condition origin) environment lag advantage from a random uniform
distribution [0.5, 3] and setting the mean and standard deviation of its lag (dis)advantage
distribution in all five static environments equal to 0.5 times its home advantage. This choice
was based on the fact that we observe that mutants with higher mean fitness difference also have
higher than average home environment fitness (Fig. S5E-F).

We then simulated fitness in fluctuating environments by shifting a mutant's lag (dis)advantage
away from its static environment value and toward the mean of the mutant's distribution. Guided
by our finding that memory correlates with global fitness difference when controlling for local
fitness difference (Fig. S7), we shift proportionately to the width of the mutant's distribution
across all static environments. We fit the result that the wrong static environment is a better
predictor for fitness in fluctuating environment components in 39% of cases by setting this shift
to 80% of the distance from the local lag (dis)advantage to the mean of the distribution.

Whole-genome sequencing analysis
We selected 359 mutants for sequencing, including as many mutants with high mean memory as possible. Two samples were thrown out because their barcoded regions did not match the appropriate barcode sequence, possibly due to contamination, and an additional five were discarded due to low (<20X) coverage, leaving 352 whole-genome sequences.

FASTQ processing
For each sample, we received two fastq files, one for each read of the paired-end sequencing (fastqR1 and fastqR2). Reads were mapped using bwa to *Saccharomyces cerevisiae* S288C reference genome R64-1-1 (https://downloads.yeastgenome.org/sequence/S288C_reference/genome_releases/) and converted to BAM format and sorted using samtools. Read groups were fixed using the Picard tool AddOrReplaceReadGroups. Duplicates were marked using the Picard tool MarkDuplicates and the BAM file indexed using samtools.

Single-nucleotide polymorphism and small indel variant calling
Single-nucleotide polymorphism (SNP) and small indel variants were called using the HaplotypeCaller tool from GATK. GVCF files were then merged using GenomicsDBImport and genotyping was performed on the combined gVCF data with GenotypeGVCFs to produce a single VCF file with the called variants for all samples. Average coverage for each sample was calculated using samtools, keeping only those with at least 20X coverage. Variants were then filtered to exclude ancestral genotypes, mitochondrial variants, and using GATK default filters for other filtering. Results were checked visually in IGV. We discarded variants that appeared erroneous due to poor alignment or coverage, particularly in repetitive regions, and marked true variants as homozygous or heterozygous, based on whether the variant appeared in closer to 50% or 100% of reads. Finally, variants were annotated using snpEff.

Copy number variant calling
BAM files were searched for high-coverage regions with samtools and bedtools, selecting genome sites with more than 3X average coverage. To eliminate ancestral copy number variation, all selected sites were compared and results appearing in more than 15 samples were dropped. The remaining results were checked visually in IGV for copy number variation.

Ploidy assay
We conducted a ploidy test as in ref [3]. Saturated cultures from the 96-well plates also used for whole-genome sequencing were mixed 5ul was pipetted onto YPD + 20 ug/ml benomyl (in DMSO) on rectangular agar plates, grown at room temperature for several days and then imaged. Under these conditions, diploid growth was strongly inhibited by benomyl but haploid growth was less affected. We pipetted onto two agar plates per 96-well plate, using every other well, in order to avoid effects of nearest neighbors on colony growth.

Acknowledgements

We are grateful to Grant Kinsler for patiently training us in the experimental system, as well as for valuable advice about PCR and sequencing, data analysis, and whole-genome sequencing analysis. We thank members of the Petrov Lab for helpful comments and suggestions. We thank Anastasia Lyulina for assistance with preliminary experiments, Katherine Xue, Katja Schwartz, and Jason Tarkington for advice on library preparation protocols, and Aditya Mahadevan for advice on whole-genome sequencing analysis. For additional helpful discussions, we thank Yuping Li, Benjamin Good, Daniel Wong, Jonas Cremer, Jeff Gore, Martina Dal Bello, and Maitreya Dunham. Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number F32GM145148.

References

1. Woods, R., Schneider, D., Winkworth, C. L., Riley, M. A. & Lenski, R. E. Tests of parallel molecular evolution in a long-term experiment with Escherichia coli. *Proc. Natl. Acad. Sci.* **103**, 9107–9112 (2006).

2. Tenaillon, O. *et al.* The Molecular Diversity of Adaptive Convergence. *Science* **335**, 457–461 (2012).

3. Venkataram, S. *et al.* Development of a Comprehensive Genotype-to-Fitness Map of Adaptation-Driving Mutations in Yeast. *Cell* **166**, 1585-1596.e22 (2016).

4. Good, B. H., McDonald, M. J., Barrick, J. E., Lenski, R. E. & Desai, M. M. The dynamics of molecular evolution over 60,000 generations. *Nature* **551**, 45–50 (2017).

5. Rudman, S. M. *et al.* Direct observation of adaptive tracking on ecological time scales in Drosophila. *Science* **375**, eabj7484 (2022).

6. Kryazhimskiy, S., Rice, D. P., Jerison, E. R. & Desai, M. M. Global epistasis makes adaptation predictable despite sequence-level stochasticity. *Science* **344**, 1519–1522 (2014).

7. Couce, A. & Tenaillon, O. A. The rule of declining adaptability in microbial evolution experiments. *Front. Genet.* **6**, 99 (2015).

8. Kover, P. X. *et al.* Pleiotropic effects of environment-specific adaptation in Arabidopsis thaliana. *New Phytol.* **183**, 816–825 (2009).

9. Schick, A., Bailey, S. F. & Kassen, R. Evolution of Fitness Trade-Offs in Locally Adapted Populations of Pseudomonas fluorescens. *Am. Nat.* **186**, S48–S59 (2015).

10. Li, Y., Petrov, D. A. & Sherlock, G. Single nucleotide mapping of trait space reveals Pareto fronts that constrain adaptation. *Nat. Ecol. Evol.* **3**, 1539–1551 (2019).

11. Jerison, E. R., Nguyen Ba, A. N., Desai, M. M. & Kryazhimskiy, S. Chance and necessity in the pleiotropic consequences of adaptation for budding yeast. *Nat. Ecol. Evol.* **4**, 601–611 (2020).

12. Kinsler, G., Geiler-Samerotte, K. & Petrov, D. A. Fitness variation across subtle environmental perturbations reveals local modularity and global pleiotropy of adaptation. *eLife* **9**, e61271 (2020).

13. Bono, L. M., Smith, L. B., Pfennig, D. W. & Burch, C. L. The emergence of performance trade-offs during local adaptation: insights from experimental evolution. *Mol. Ecol.* **26**, 1720–1733 (2017).

14. Cooper, T. F. & Lenski, R. E. Experimental evolution with E. coli in diverse resource environments. I. Fluctuating environments promote divergence of replicate populations. *BMC Evol. Biol.* **10**, 11 (2010).

15. Rodríguez-Verdugo, A. & Ackermann, M. Rapid evolution destabilizes species interactions in a fluctuating environment. *ISME J.* **15**, 450–460 (2021).

16. Salignon, J., Richard, M., Fulcrand, E., Duplus-Bottin, H. & Yvert, G. Genomics of cellular proliferation in periodic environmental fluctuations. *Mol. Syst. Biol.* **14**, e7823 (2018).

17.   Boyer, S., Hérissant, L. & Sherlock, G. Adaptation is influenced by the complexity of environmental change during evolution in a dynamic environment. *PLOS Genet.* **17**, e1009314 (2021).

18.   Alto, B. W., Wasik, B. R., Morales, N. M. & Turner, P. E. STOCHASTIC TEMPERATURES IMPEDE RNA VIRUS ADAPTATION. *Evolution* **67**, 969–979 (2013).

19.   Fasanello, V. J., Liu, P., Fay, J. C. & Botero, C. A. Fluctuating selection facilitates the discovery of broadly effective but difficult to reach adaptive outcomes in yeast. 2023.06.17.545402 Preprint at https://doi.org/10.1101/2023.06.17.545402 (2023).

20.   Cvijović, I., Good, B. H., Jerison, E. R. & Desai, M. M. Fate of a mutation in a fluctuating environment. *Proc. Natl. Acad. Sci.* **112**, E5021–E5028 (2015).

21.   Sæther, B.-E. & Engen, S. The concept of fitness in fluctuating environments. *Trends Ecol. Evol.* **30**, 273–281 (2015).

22.   Ketola, T. & Kristensen, T. N. Experimental Approaches for Testing if Tolerance Curves Are Useful for Predicting Fitness in Fluctuating Environments. *Front. Ecol. Evol.* **5**, (2017).

23.   Monod, J. The growth of bacterial cultures. *Annu. Rev. Microbiol.* **3**, 371–394 (1949).

24.   Siegal, M. L. Shifting Sugars and Shifting Paradigms. *PLOS Biol.* **13**, e1002068 (2015).

25.   Escalante-Chong, R. *et al.* Galactose metabolic genes in yeast respond to a ratio of galactose and glucose. *Proc. Natl. Acad. Sci.* **112**, 1636–1641 (2015).

26.   Basan, M. *et al.* A universal trade-off between growth and lag in fluctuating environments. *Nature* **584**, 470–474 (2020).

27.   Balakrishnan, R., de Silva, R. T., Hwa, T. & Cremer, J. Suboptimal resource allocation in changing environments constrains response and growth in bacteria. *Mol. Syst. Biol.* **17**, e10597 (2021).

28.   Li, Y. *et al.* Hidden Complexity of Yeast Adaptation under Simple Evolutionary Conditions. *Curr. Biol.* **28**, 515-525.e6 (2018).

29.   Levy, S. F. *et al.* Quantitative evolutionary dynamics using high-resolution lineage tracking. *Nature* **519**, 181–186 (2015).

30.   Zakrzewska, A. *et al.* Genome-wide analysis of yeast stress survival and tolerance acquisition to analyze the central trade-off between growth rate and cellular robustness. *Mol. Biol. Cell* **22**, 4435–4446 (2011).

31.   Dhar, R., Sägesser, R., Weikert, C. & Wagner, A. Yeast Adapts to a Changing Stressful Environment by Evolving Cross-Protection and Anticipatory Gene Regulation. *Mol. Biol. Evol.* **30**, 573–588 (2013).

32.   Ketola, T. & Saarinen, K. Experimental evolution in fluctuating environments: tolerance measurements at constant temperatures incorrectly predict the ability to tolerate fluctuating temperatures. *J. Evol. Biol.* **28**, 800–806 (2015).

33.   Vermeersch, L. *et al.* On the duration of the microbial lag phase. *Curr. Genet.* **65**, 721–727 (2019).

34.   McDaniel, E. A., Stuecker, T. N., Veluvolu, M., Gasch, A. P. & Lewis, J. A. Independent Mechanisms for Acquired Salt Tolerance versus Growth Resumption Induced by Mild Ethanol Pretreatment in Saccharomyces cerevisiae. *mSphere* **3**, e00574-18 (2018).

35.   Rivers, W. Characterizing Aft1/2-Grx3/4 Interaction and the Role of Bol2 During Iron Regulation in Saccharomyces cerevisiae.

36.   Martínez-Pastor, M. T., Perea-García, A. & Puig, S. Mechanisms of iron sensing and regulation in the yeast Saccharomyces cerevisiae. *World J. Microbiol. Biotechnol.* **33**, 75 (2017).

37.   Christensen, J. H. *et al.* Climate phenomena and their relevance for future regional climate change. in *Climate Change 2013 the Physical Science Basis* vol. 9781107057999 1217–1308 (Cambridge University Press, 2013).

38.   Khan, A. I., Dinh, D. M., Schneider, D., Lenski, R. E. & Cooper, T. F. Negative Epistasis Between Beneficial Mutations in an Evolving Bacterial Population. *Science* **332**, 1193–1196 (2011).

39.   Chou, H.-H., Chiu, H.-C., Delaney, N. F., Segrè, D. & Marx, C. J. Diminishing Returns Epistasis Among Beneficial Mutations Decelerates Adaptation. *Science* **332**, 1190–1192 (2011).

40.   Reddy, G. & Desai, M. M. Global epistasis emerges from a generic model of a complex trait. *eLife* **10**, e64740 (2021).

41.   Diaz-Colunga, J., Skwara, A., Vila, J. C. C., Bajic, D. & Sánchez, Á. Global epistasis and the emergence of ecological function. 2022.06.21.496987 Preprint at https://doi.org/10.1101/2022.06.21.496987 (2023).

42.   Violle, C., Pu, Z. & Jiang, L. Experimental demonstration of the importance of competition under disturbance. *Proc. Natl. Acad. Sci.* **107**, 12925–12929 (2010).

43.   Abreu, C. I., Woltz, V. L. A., Friedman, J. & Gore, J. Microbial communities display alternative stable states in a fluctuating environment. *PLOS Comput. Biol.* **16**, e1007934 (2020).

44.   Yi, X. & Dean, A. M. Bounded population sizes, fluctuating selection and the tempo and mode of coexistence. *Proc. Natl. Acad. Sci.* **110**, 16945–16950 (2013).

45. Letten, A. D., Dhami, M. K., Ke, P.-J. & Fukami, T. Species coexistence through simultaneous fluctuation-dependent mechanisms. *Proc. Natl. Acad. Sci.* **115**, 6745–6750 (2018).

46. Chesson, P. & Huntly, N. The Roles of Harsh and Fluctuating Conditions in the Dynamics of Ecological Communities. *Am. Nat.* **150**, 519–553 (1997).

47. Venkataram, S. BarcodeCounter2. (2021).

48. Zhao, L., Liu, Z., Levy, S. F. & Wu, S. Bartender: a fast and accurate clustering algorithm to count barcode reads. *Bioinformatics* **34**, 739–747 (2018).

49. Li, F., Mahadevan, A. & Sherlock, G. An improved algorithm for inferring mutational parameters from bar-seq evolution experiments. *BMC Genomics* **24**, 246 (2023).