1 **Dendritic *atoh1a*+ cells serve as transient intermediates during zebrafish Merkel cell**
2 **development and regeneration**
3

4 Evan W. Craig[1], Erik C. Black[1,2], Camille E.A. Goo[1], Avery Angell Swearer[1,2], Nathaniel G. Yee[1],
5 Jeffrey P. Rasmussen[1,3,*]
6
[1] Department of Biology, University of Washington, Seattle, WA, USA
[2] Molecular and Cellular Biology Program, University of Washington, Seattle, WA, USA
[3] Institute for Stem Cell and Regenerative Medicine, University of Washington, Seattle,
WA, USA
* Correspondence: rasmuss@uw.edu

Keywords: Ectodysplasin, epidermis, cell motility, somatosensory system, skin organogenesis,
piezo2, tp63, microvilli

ABSTRACT

Sensory cells often adopt specific morphologies that aid in the detection of external stimuli.
Merkel cells encode gentle touch stimuli in vertebrate skin and adopt a reproducible shape
characterized by spiky, actin-rich microvilli that emanate from the cell surface. The mechanism
by which Merkel cells acquire this stereotyped morphology from basal keratinocyte progenitors
is unknown. Here, we establish that dendritic Merkel cells (dMCs) express *atonal homolog 1a*
(atoh1a), extend dynamic filopodial processes, and arise in transient waves during zebrafish
skin development and regeneration. We find that dMCs share molecular similarities with both
basal keratinocytes and Merkel cells, yet display mesenchymal-like behaviors, including local
cell motility and proliferation within the epidermis. Furthermore, dMCs can directly adopt the
mature, microvilliated Merkel cell morphology through substantial remodeling of the actin
cytoskeleton. Loss of Ectodysplasin A signaling alters the morphology of dMCs and Merkel cells
within specific skin regions. Our results show that dMCs represent an intermediate state in the
Merkel cell maturation program and identify Ectodysplasin A signaling as a key regulator of
Merkel cell morphology.

INTRODUCTION

Organ development and function require that constituent cells adopt precise shapes. For example, epithelial cells often develop elaborate actin-based membrane protrusions integral to organ function, such as the brush border of intestinal enterocytes or the stereocilia of inner ear hair cells (reviewed by Sharkova et al., 2023). Defects in the morphogenesis of actin-based protrusions are linked to a variety of diseases, including colorectal cancer and deafness. Thus, elucidating the mechanisms of how epithelial cells adopt specific shapes is relevant to understanding both the basis for organ function and the etiology of human pathologies.

Merkel cells (MCs) are mechanosensory epidermal cells that interact with somatosensory neurites to form the MC-neurite complex, which mediates detection of gentle touch stimuli (reviewed by Woo et al., 2015). MCs populate diverse vertebrate skin compartments, including human plantar and palmar skin, mouse hairy and glabrous skin, and teleost facial and trunk skin (reviewed by Hartschuh et al., 1986; Whitear, 1989). MCs have a remarkably consistent morphology across these diverse vertebrate skin types. The core morphological features of MCs include a small and spherical cell body, a high nuclear-to-cytoplasmic ratio, and the presence of neurosecretory granules (Hartschuh et al., 1986; Whitear, 1989). But perhaps the most striking morphological feature of MCs are the numerous actin-rich microvilli that emanate off the cell surface, giving MCs a "mace-like" morphology (Hartschuh et al., 1986; Lane and Whitear, 1977; Sekerková et al., 2004; Takahashi-Iwanaga, 2003; Takahashi-Iwanaga and Abe, 2001; Toyoshima et al., 1998; Whitear and Lane, 1981).

The precise and reproducible cell shape of MCs has been proposed to impact their sensory function (Garant et al., 1980; Takahashi-Iwanaga, 2003; Takahashi-Iwanaga and Abe, 2001; Toyoshima et al., 1998; Yamashita et al., 1993); however, it is unknown how MCs acquire this stereotyped morphology. Cre-based lineage tracing experiments indicate that MCs derive from basal keratinocyte precursors in both mammalian and zebrafish skin (Brown et al., 2023; Morrison et al., 2009; Van Keymeulen et al., 2009). While these experiments begin to address MC progenitor identity, they fail to shed light on how cells within this lineage lose basal keratinocyte characteristics and adopt the hallmark MC morphology. A deeper understanding of the MC lineage may shed light on the origins of Merkel cell carcinoma, an aggressive skin cancer of unclear cellular origin (reviewed by Becker et al., 2017; Harms et al., 2018).

Interestingly, several histological observations support the presence of cellular intermediates in the MC lineage. First, ultrastructural studies identified "transitional cells" that exhibit characteristics of both keratinocytes and MCs, including fine cytoplasmic filaments and dense-core granules, respectively (English, 1974; English, 1977; Garant et al., 1980; Tachibana and Ishizeki, 1981; Tachibana and Nawa, 1980). Second, immunostaining for two MC markers—cytokeratins 18 and 20—revealed cells with oblong or dendritic morphologies in developing human plantar skin, human and rodent oral mucosa, and murine touch domes (Kim and Holbrook, 1995; Moll et al., 1984; Nakafusa et al., 2006; Tachibana et al., 1997; Tachibana et al., 1998). Although these heterogeneously shaped cells have been referred to as dendritic MCs (dMCs), it is unknown whether dMCs represent a developmentally immature MC state or a

75   functionally distinct subpopulation of neuroendocrine cells (reviewed by Xiao et al., 2014). Live-
76   cell imaging of MC development would be the most direct way to distinguish these possibilities.
77   However, to date, technical limitations of mammalian systems have precluded visualization of
78   dMC or MC dynamics in living animals.
79
We recently established zebrafish as an *in vivo* model for MC studies (Brown et al., 2023)*. Here,*
we leverage the optical accessibility of zebrafish skin to directly visualize individual cell
behaviors during the MC maturation process. By visualizing key stages of skin organogenesis
and regeneration using an F-actin reporter expressed in MCs, we describe a morphologically
distinct population of epidermal cells that shares characteristics with transitional cells and dMCs
observed in other types of vertebrate skin. Importantly, we document the direct maturation of
dMCs into MCs through cytoskeletal rearrangements. Furthermore, we show that Ectodysplasin
A (Eda) signaling is required for MCs to develop microvilli within trunk skin. Together, our results
provide the first *in vivo* characterizations of MC precursor states and identify Eda signaling as a
key regulator of zebrafish MC morphogenesis.
90
91   **RESULTS**
92
93   ***A transient, morphologically distinct population of keratinocyte-derived* atoh1a+ *cells***
94   ***emerges during skin development***
95
Staging of zebrafish post-embryonic development relies on standard length (SL) in millimeters,
with sexual maturity attained ~18 mm SL (Parichy et al., 2009). Cells with the prototypical MC
morphology—spherical with actin-rich, microvillar protrusions of ~1-2 microns in length—
populate several adult zebrafish skin compartments, including trunk skin (**Figure 1A,B**; (Brown
et al., 2023)). *Atonal homolog 1* (*Atoh1*) encodes a transcription factor necessary and sufficient
for murine MC development (Morrison et al., 2009; Ostrowski et al., 2015; Van Keymeulen et
al., 2009). Zebrafish *atoh1a* is an ortholog of murine *Atoh1*, and zebrafish MCs express
reporters inserted into the upstream region of the *atoh1a* locus (Brown et al., 2023). Using a
nuclear localized *atoh1a* reporter to label MCs, we previously found that trunk MCs begin to
appear during the onset of squamation (scale formation) at approximately 9 mm SL and rapidly
increase in density from 10-15 mm SL (Brown et al., 2023). As this nuclear reporter did not
allow for visualization of cell shapes, it is unclear whether morphological heterogeneity exists
within developing zebrafish MCs, including the possible presence of dMCs.
109
To assess MC morphology during squamation, we conducted a similar developmental staging
time course using *Tg(atoh1a:lifeact-egfp)*, which expresses a filamentous actin (F-actin) reporter
in MCs, allowing detailed visualization of the MC actin cytoskeleton *in vivo* (**Figure 1C**; (Brown
et al., 2023)). To simultaneously visualize scales, we stained animals with Alizarin Red S, which
labels the calcified scale matrix. Like our previous observations in adults, we identified cells with
the prototypical MC morphology at late juvenile and adult stages (**Figure 1F-H,** arrowheads).
Interestingly, we also observed a second population of *atoh1a+* cells with weaker transgene
signal and highly variable morphologies, characterized by ovoid cell bodies and long, filopodial-
like actin-rich protrusions (**Figure 1D-G,** arrows). For consistency with the literature (Boot et al.,

1992; Kim and Holbrook, 1995; Moll et al., 1984; Nakafusa et al., 2006; Tachibana et al., 1997;
Tachibana et al., 1998), we hereafter refer to these cells as dMCs, although we note that
whether dMCs are related to MCs or represent an alternative fate has not been established.
Manual cell counting from our imaging dataset revealed dMCs appeared in highest frequency
and density at 10-15 mm SL, when scales expand and total MC density rapidly increases
(**Figure 1I-K**). Next, we quantified *atoh1a* cell shapes at two different developmental stages by
assigning circularity and roundness values to thresholded images from our dataset. We defined
"circular" cells as those having circularity and roundness values >0.7. We found that 95.3% of
atoh1a+ cells in adult skin fell into the circular category, likely representing the mature,
microvilliated MC type (**Figure 1M,N**). By contrast, only 71.1% of *atoh1a+* cells in juvenile skin
fell into the circular category, with numerous *atoh1a+* cells adopting more oblong and less round
shapes. (**Figure 1L,N**). Thus, our results indicate that a morphologically distinct population of
atoh1a+ cells transiently populates the trunk skin during squamation.
132
Using a tamoxifen-inducible Cre driver expressed in basal keratinocytes [*TgBAC(ΔNp63:Cre-*
ERT2); (Brown et al., 2023)] and a quasi-ubiquituous Cre reporter transgene [*Tg(actb2:LOXP-*
BFP-LOXP-DsRed); (Kobayashi et al., 2014)], we previously found that zebrafish MCs derive
from *ΔNp63*-expressing basal keratinocytes (Brown et al., 2023). To determine if basal
keratinocytes also give rise to dMCs, we treated *TgBAC(ΔNp63:Cre-ERT2); Tg(actb2:LOXP-*
BFP-LOXP-DsRed); Tg(atoh1a:lifeact-egfp) embryos with 4-OHT to induce Cre-ERT2 activity at
1 day post-fertilization (dpf). This resulted in permanent DsRed expression in basal
keratinocytes and their cellular derivatives. We then raised animals to squamation stages. Due
to transgene mosaicism and/or incomplete Cre-ERT2 activation, not all basal keratinocytes
expressed DsRed (**Figure 1O**). Along with DsRed+ MCs (**Figure 1O'''**), we observed that a
subset of dMCs in juvenile skin were DsRed+ (**Figure 1O'-O''**). Interestingly, we often identified
DsRed+ dMCs at a distance from the nearest DsRed+ basal keratinocyte (**Figure 1O**). These
findings demonstrate a basal keratinocyte origin for dMCs and suggest they may be motile.
146
147   ***dMCs are the primary* atoh1a+ *cell type during early stages of skin regeneration***
148
Our identification of dMCs during squamation prompted us to hypothesize that dMCs may also
appear during specific stages of scale regeneration. To induce scale regeneration, we
performed scale "plucking" with fine forceps to remove both the bony scale and overlying
epidermis containing MCs (**Figure 2A**). Scale removal triggers a wound healing response from
neighboring keratinocytes, which migrate from surrounding, non-injured scales to cover the
denuded area within hours (Richardson et al., 2016). Over the course of several days,
keratinocytes then proliferate and reestablish a stratified epidermis (Richardson et al., 2016),
while dermal osteoblasts proliferate and undergo hypertrophy to regenerate the bony scale
(Aman et al., 2018; Cox et al., 2018; De Simone et al., 2021; Iwasaki et al., 2018; Sire et al.,
1997).
159
To determine whether MCs populated the regenerating scale epidermis, we plucked scales from
animals expressing *Tg(atoh1a:Lifeact-egfp)* and *Tg(sp7:mCherry)* (Singh et al., 2012), a
reporter for scale-forming osteoblasts (**Figure 2A,B**). As expected, scales underwent

substantial regeneration within the first 7 days post-pluck (dpp) (**Figure 2B-E**). *atoh1a*+ cells
appeared above regenerating scales beginning at 2 dpp and rapidly increased in density until 7
dpp, after which the density plateaued (**Figure 2F-J**). Strikingly, dMCs comprised ~95% of the
atoh1a+ cells present at 2 dpp (**Figure 2G,G',K**) and ~45% of the *atoh1a*+ cells at 3 dpp
(**Figure 2H,K**). After 3 dpp, the proportion of dMCs gradually decreased, with MCs becoming
the predominant *atoh1a*+ cell type at later stages of regeneration (**Figure 2I,K**)
169
dMCs and MCs on regenerating scales could arise from either *de novo* production or movement
of pre-existing *atoh1a*+ cells from surrounding, uninjured regions of epidermis. To distinguish
between these possibilities, we irreversibly photoconverted *atoh1a*+ cells expressing a nuclear
localized version of the photoconvertible protein Eos [*Tg(atoh1a:nls-Eos);* (Pickett et al., 2018)]
and then plucked scales (**Figure 2L,M**). In this experimental paradigm, pre-existing *atoh1a*+
cells contain both photoconverted and non-photoconverted nls-Eos, whereas newly
differentiating *atoh1a*+ cells contain only non-photoconverted nls-Eos. Notably, at 2 and 3 dpp
the epidermis contained many newly differentiated cells with oval nuclear morphologies that
likely represented dMCs **(Figure 2N,O,Q)**. By 7 dpp, nearly all *atoh1a*+ cells were newly derived
(**Figure 2P,Q**). Based on these results, we concluded that skin injury triggers *de novo*
production of dMCs and MCs within the trunk epidermis and that early stages of epidermal
regeneration are associated with a high proportion of dMCs. Thus, dMCs are a transient
atoh1a+ cell type during both skin development and regeneration.
183
184   ***dMCs exhibit molecular features of both basal keratinocytes and MCs***
185
To characterize the molecular and cellular properties of regenerating dMCs, we focused on 5
dpp, a timepoint that contains a mix of dMCs and MCs (**Figure 2K**). We first assayed
expression of Tp63, a transcription factor critical in epidermal development and highly
expressed in basal keratinocytes (Bakkers et al., 2002; Guzman et al., 2013; Lee and
Kimelman, 2002; Rangel-Huerta et al., 2021) (**Figure 3B, asterisks**). Immunostaining revealed
that MCs did not express detectable levels of Tp63 (**Figure 3A-A'**). By contrast, dMCs exhibited
significantly higher levels of Tp63 staining than MCs, albeit at lower levels than basal
keratinocytes, with ~75% of dMCs staining positive for Tp63 (**Figure 3B-C**). Thus, dMCs
express Tp63 at intermediate levels relative to basal keratinocytes and MCs.
195
Given our finding that dMCs expressed *atoh1a*, we postulated that dMCs might share additional
molecular properties with MCs. MCs express markers of neuroendocrine and mechanosensory
function, such as synaptic vesicle glycoprotein 2 (SV2) and the mechanosensitive cation
channel Piezo2, respectively. Consistent with previous observations during ontogeny (Brown et
al., 2023), we found that regenerated MCs expressed SV2 (**Figure 3D-D'**). SV2 staining of
dMCs in regenerating skin revealed a mix of SV2+ and SV2– dMCs (**Figure 3E-E'**). We
quantified the SV2 fluorescence intensity from dMCs with detectable signal and found dMCs
displayed lower SV2 levels than MCs (**Figure 3F**). Hybridization chain reaction (HCR) staining
for *piezo2* labeled both MCs (**Figure 3G-G', arrowhead**) and dMCs (**Figure 3G-G', arrow**) in
regenerating skin. Analysis of the *piezo2* signal revealed slightly lower levels in dMCs compared
to MCs (**Figure 3H**). In summary, regenerating dMCs display molecular properties that overlap

with both basal keratinocytes and MCs, suggesting they may represent a transitional or
immature MC.

MC and dMC protrusions are dynamic and differentially polarized

Apart from previous reports that tracked MC turnover over the course of days or weeks by
imaging cytoplasmic or nuclear reporters (Brown et al., 2023; Wright et al., 2017), little is known
about the *in vivo* dynamics of MCs. Given the differences in cell shapes between dMCs and
MCs during development and regeneration, we sought to compare the dynamics and polarities
of the actin cytoskeleton in these two populations. To this end, we mounted and intubated
Tg(atoh1a:lifeact-egfp) animals and performed live-cell confocal microscopy of fully intact skin
over several hours. Imaging of MCs in adult zebrafish revealed strong Lifeact-EGFP signal in
microvilli as expected (**Figure 4A, arrowhead**). We captured numerous microvillar reorientation,
retraction, and extension events occurring in all MCs (**Figure 4A-A''''; Movie 1**), suggesting
previously unappreciated MC microvillar dynamics. MC microvilli rarely extended >3 microns
from the membrane and often appeared to coalesce or merge together, although we lacked the
resolution to reliably characterize these merging events. Despite these microvillar dynamics, the
MC cortex remained spherical during imaging (**Figure 4A-A''''**). By analyzing reconstructed
cross-sections of MC microvilli in animals also expressing a Cdh1 (E-cadherin) knock-in reporter
to label keratinocyte membranes [Cdh1-tdTomato; (Cronan et al., 2018)] (**Figure 4C**), we found
the majority of MC microvilli extended in basal keratinocyte-facing or lateral directions, whereas
periderm-facing processes were less common (**Figure 4D,D',F,G**).

In contrast to MC microvilli, imaging of dMCs in juveniles revealed long, thin Lifeact-EGFP+
protrusions that routinely exceeded 10 microns in length and in extreme cases exceeded 25
microns (**Figure 4B-B'''', arrowheads**). dMCs dynamically rearranged both their protrusions
and cell cortex (**Figure 4B-B''''; Movie 2**). dMC protrusions tended to coalesce and extend from
one end of the cell, with the opposite side of the dMC having few or no protrusions (**Figure
4B''**). The majority of dMC protrusions extended in lateral or periderm-facing directions,
whereas basal keratinocyte-facing processes were less common (**Figure 4E,E',F,G).**
Reconstructed cross-sections also revealed that MCs tended to reside just below the periderm,
whereas dMCs were commonly in lower strata (**Figure 4D',E',H**). Thus, our observations
indicate MCs and dMCs occupy different strata of the epidermis and utilize dynamic actin-based
protrusions that are of distinct sizes and polarities.

dMCs are mesenchymal-like cells that directly mature into MCs

We wondered how MCs and dMCs behaved during periods of development when MC density
greatly increases in the trunk skin (10-15 mm SL, **Figure 1J,K**). Based on their association with
somatosensory neurites and desmosome-like contacts with keratinocytes, we hypothesized that
MCs would be relatively immotile. By contrast, our Cre lineage tracing data (**Figure 1O**) and
observations of dMC protrusion dynamics and polarity (**Figure 4**), led us to hypothesize that
dMCs may migrate within the epidermis.

To characterize MC and dMC motility, we intubated *Tg(atoh1a:lifeact-egfp)* juveniles and
imaged fields of view containing MCs and dMCs over 6 h (**Figure 5A, Movie 3**). Consistent with
our hypotheses, cell tracking revealed that MCs were largely immotile, whereas dMCs migrated
laterally within the epidermis (**Figure 5A, insets**). dMC motility was highly variable: On average,
dMCs migrated 2.4 µm/hr during imaging, but some moved upwards of 8 µm/hr, while others
remained stationary (**Figure 5B**). By contrast, MCs were immotile, with an average
displacement of 0.8 µm/hr, which likely reflects imaging drift (**Figure 5B**). dMCs were most
motile when adopting elongated, ovoid cell bodies with long, unipolar protrusions at one end of
the cell (**Figure 5E, pink arrowheads**). By contrast, dMCs with multipolar protrusions were
largely immotile (**Figure 5E, orange arrowheads**). Although the majority of dMCs traveled in a
persistent or linear manner (**Figure 5C**), dMCs had a significantly reduced track displacement
compared to total distance traveled (**Figure 5D**), suggesting that dMCs migrate directionally but
often switch directions. Consistent with this notion, dMC-dMC or dMC-MC contacts resulted in
lateral dMC movement away from the contact (**Movie 4**). Together these observations suggest
that MCs are immotile, epithelial-like cells, whereas dMCs are motile, mesenchymal-like cells
that undergo contact inhibition upon encountering another *atoh1a+* cell.
267
MCs are generally considered post-mitotic (Merot and Saurat, 1988; Moll et al., 1996; Vaigot et
al., 1987; Weber et al., 2023; Wright et al., 2017). Indeed, we did not observe instances of MC
cell division in our live-imaging of either homeostatic juvenile or regenerating adult skin. By
contrast, dMCs divided in both contexts (**Figure S1A-C**). During cell division, dMCs resorbed
their protrusions, developed a smooth actin-rich membrane, and underwent cytokinesis to
generate two daughter dMCs, which established protrusions orthogonal to the division plane
and migrated away from each other (**Figure S1A-C; Movie 5)**. Due to its experimental
tractability, we quantified dMC and MC cell division frequency during skin regeneration. To this
end, we injected 5-ethynyl-2´-deoxyuridine (EdU), a thymidine analog incorporated into DNA
during active DNA synthesis, at 4 dpp and fixed and stained scales at 5 dpp (**Figure S1D**). In
this experimental paradigm, no MCs stained positive for EdU, whereas 65% of dMCs stained
positive for EdU (**Figure S1E,F**). Thus, in addition to exhibiting differences in motility, dMCs and
MCs show contrasting cell cycle states.
281
Our findings that dMCs exhibited characteristics of both basal keratinocytes and MCs (**Figure 3**)
suggested they may represent an intermediate cellular state between these two cell types.
Indeed, during our live-imaging of juvenile and regenerating adult skin, we observed a small
subset of dMCs (*n*>10) withdraw their long protrusions, round up their cell body, and rapidly
extend microvilli reminiscent of the mature "mace-like" MC morphology (**Figure 5F; Movies
6,7**). These observations are consistent with the direct maturation of a dMC into a MC. In rare
instances (*n*=2), we observed the reciprocal process, i.e., the reversion of a MC into a dMC-like
morphology (**Figure 5G; Movie 8**), suggesting that some MCs can switch between cell states.
Thus, based on these observations, we conclude that dMCs are capable of cell migration, cell
division, and direct maturation into MCs (**Figure 5H**).
292
Genetic loss of Eda results in altered dMC and MC morphologies in trunk skin

Eda signaling promotes the development of diverse vertebrate skin appendages (reviewed by
Sadier et al., 2014), including zebrafish scales (Harris et al., 2008). We previously reported that
zebrafish homozygous for a presumptive null *eda* allele (*eda^{dt1261/dt1261}*, hereafter *eda^{-/-}*)
displayed reduced MC density within the trunk, but not facial, epidermis (Brown et al., 2023).
We postulated that Eda signaling may also promote the morphological maturation of MCs. To
address this hypothesis, we incrossed *eda^{+/-}* adults and used *Tg(atoh1a:lifeact-egfp)* localization
to compare MC morphology between homozygous mutants and sibling controls. As expected,
control trunk MCs developed a typical "mace-like" morphology decorated by microvilli (**Figure**
6A-A''',G). By contrast, we observed a striking and highly penetrant loss of MC microvillar
structures along the trunk of *eda^{-/-}* mutants (**Figure 6B-B''',G**). In *eda^{-/-}* trunk MCs, Lifeact-
EGFP signal accumulated near the membrane in a ring-like fashion (**Figure 6B', arrowhead**),
reminiscent of the morphology of dMCs during cell division (**Figure S1A-C**). The few MC
microvilli that formed in *eda^{-/-}* trunk MCs were short and thin, making them difficult to resolve
with confocal microscopy. To determine whether Eda globally regulated MC morphology
throughout the skin, we imaged MCs in the corneal epidermis, which is not squamated. This
analysis revealed corneal MCs with similar microvilli-rich morphologies in *eda^{-/-}* mutants and
siblings (**Figure 6C-D,G**). Finally, to determine whether loss of Eda signaling also impacted
dMC morphology, we compared the maximum protrusion length of dMCs in *eda^{-/-}* mutants and
controls and found that mutant dMCs had significantly longer protrusions (**Figure 6E,F,H;** *eda^{-/-}*
dMC mean lengths = 16.3 μm, siblings = 8.7 μm). In extreme cases, dMC protrusions reached
almost 40 microns in length in *eda^{-/-}* trunk epidermis (**Figure 6F**). Together, these results
indicate that Eda is necessary for the normal morphologies of dMCs and MCs specifically within
the trunk skin compartment.
317
318
DISCUSSION
The defining morphological features of MCs allows for their identification across diverse types of
vertebrate skin (Hartschuh et al., 1986; Whitear, 1989). How MCs acquire this precise
morphology during skin development or regeneration is unknown. While basal keratinocytes
have been identified as MC progenitors in mammalian and zebrafish skin, intermediate steps in
the MC lineage progression remain poorly understood. Previous histology-based experiments
noted transitional cells that could serve as cellular intermediates during MC maturation (English,
1974; English, 1977; Garant et al., 1980; Tachibana and Ishizeki, 1981; Tachibana and Nawa,
1980). However, these studies lacked the temporal resolution to rigorously establish a
relationship between transitional cells and MCs. Here, we directly address this knowledge gap
by using the optical accessibility of zebrafish skin to track the emergence of MCs during
development and regeneration. Our results support a model in which dMCs emerge in lower
epidermal strata and migrate locally within the epidermis. Their migration occurs both laterally,
presumably due to their lateral-facing actin-rich membrane protrusions, and vertically within the
epidermis. As they mature in superficial strata, dMCs reorganize their actin cytoskeleton to
elaborate polarized microvilli, thereby adopting the terminal mace-like morphology characteristic
of MCs (**Figure 5H**).

A transient cell state during development and regeneration

We observed that dMCs appeared transiently during a period of skin organogenesis associated
with dermal appendage growth along the trunk. dMC frequency peaked during a time of rapid
increase in MC density. These findings closely mirror previous studies in other vertebrates. For
instance, the "dendritic:globular" MC ratio sharply decreased in embryonic and human fetal
palmar skin prior to the peak of MC density (Kim and Holbrook, 1995). Furthermore, our lineage
tracing identified basal keratinocytes as dMC precursors, supporting previous ultrastructural
findings that suggested basal keratinocytes give rise to transitional cells (English, 1974; English,
1977; Garant et al., 1980; Tachibana and Ishizeki, 1981; Tachibana and Nawa, 1980). However,
since those TEM studies did not reconstruct transitional cell morphology, any relationship
between transitional cells and dMCs has remained unclear. Based on our lineage tracing and
morphological data, we propose that dMCs and transitional cells likely represent similar cellular
intermediates. However, dMCs may represent just one of several intermediate states between
basal keratinocytes and MCs, each with their own characteristics as previously suggested
(Tachibana and Ishizeki, 1981).

Severe skin wounds, such as burns, can alter sensory perception, thereby degrading patient
quality of life (reviewed by Girard et al., 2017). Human MC regeneration after injury appears
poor (Stella et al., 1994), although engineered skin grafts present a promising avenue (Hahn et
al., 2019). Studies of the mechanisms of MC regeneration in model organisms may inform
improved wound healing treatments. Previous work in mammalian models found that MCs
regenerate after peripheral nerve crush (Burgess et al., 1974; Nurse et al., 1984), cauterization
of touch domes (Horch, 1982), full thickness wounds (Tachibana and Ishizeki, 1981), or
repeated skin shaving (Weber et al., 2023; Wright et al., 2017). We found that scale plucking
induced a high proportion of dMCs during the early phase of scale regeneration and that MC
density ultimately recovered to pre-injury levels. Using a photoconvertible reporter, we
established that MCs in regenerated epidermis form predominantly due to *de novo* regeneration
rather than migration from surrounding epidermal reservoirs. Due to the rapid kinetics, scale
pluck presents a tractable model for reliably triggering dMC/MC production after cutaneous
injury.

We used scale pluck as an approach to reliably induce dMCs and quantitatively characterize
their molecular properties during regeneration. We found that dMCs expressed the basal
keratinocyte marker Tp63 at intermediate levels relative to MCs (low) and basal cells (high). We
found that dMCs frequently expressed the MC markers *piezo2* and SV2, albeit at levels lower
than MCs. Together these data lend further support to the notion that transitional cells and
dMCs are one and the same, and suggest that dMCs may represent MC precursors.
Additionally, *piezo2* expression in dMCs suggests they may have the capacity for
mechanosensory function. Examining whether mechanosensation impacts any of the dMC
behaviors described here is an exciting area for future study.

Dynamic cellular behaviors of dMCs and MCs

The microvilli of several cell types, including cultured renal and immune cells, exhibit dynamic
motility (Cai et al., 2017; Meenderink et al., 2019). However, to our knowledge, no live-cell
imaging of MC microvilli has been reported. Our *in vivo* imaging revealed that MC microvilli were

dynamic and frequently extended, rearranged, or retracted. Our observations are consistent
with the lack of desmosomes along MC microvilli, a feature noted in a wide range of vertebrates
and skin compartments (Düring and Andres, 1976; Garant et al., 1980; Halata, 1975;
Hashimoto, 1972; Iggo and Muir, 1969; Landmann and Halata, 1980; Takahashi-Iwanaga and
Abe, 2001). We further found that MC microvilli were polarized in lateral- or basal keratinocyte-
facing orientations. Intriguingly, lamprey microvilli face both superficial and basal strata
(Daghfous et al., 2020; Takahashi-Iwanaga and Abe, 2001), whereas MC microvilli within cat
touch domes face superficial strata (Iggo and Muir, 1969). These observations suggest
malleability of MC microvilli polarity across vertebrates. If microvilli are involved in MC function
as previously proposed (Garant et al., 1980; Takahashi-Iwanaga, 2003; Takahashi-Iwanaga and
Abe, 2001; Toyoshima et al., 1998; Yamashita et al., 1993), reconciling microvillar dynamics
and polarity with MC function will be an interesting area for future exploration.

Our live imaging further revealed a number of previously unrecognized behaviors of dMCs,
many of which are properties of mesenchymal cells. First, we found that dMCs elaborate long,
thin actin-rich protrusions, reminiscent of protrusions previously observed with cytokeratin 20
staining in gerbil palate (Tachibana et al., 1997). These protrusions extended from lateral-facing
membranes and interdigitated between basal and suprabasal keratinocytes. Second, in contrast
to the immotile MCs, we observed that dMCs migrated laterally within the epidermis. dMC
migration correlated with the presence of unipolar protrusions, whereas dMCs with multipolar
protrusions were largely stationary. Intriguingly, dMCs with similar shapes (termed unipolar and
globular) were previously noted in adult human oral mucosa (Tachibana et al., 1998). Future
work is needed to dissect the molecular composition of dMC protrusions to determine whether
they are similar to filopodial/filopodial-like protrusions observed in other contexts. Third, we
observed that dMC-dMC or dMC-MC contacts resulted in repulsive behaviors, akin to contact
inhibition. As zebrafish MCs are exclusively solitary cells, we speculate that this behavior may
promote cell dispersion. Fourth, we observed that dMCs could divide, generating two daughter
cells with similar morphologies and behaviors. This suggests that at least a subset of dMCs may
be transient amplifying cells. Finally, and perhaps most significantly, we observed in rare
instances dMCs could mature into MCs by retracting their protrusions, adopting a spherical cell
body, and elaborating microvilli. This provides the first direct evidence that dMCs can serve as
MC precursors. Although we do not know yet what cues this maturation, it appears to involve a
transition from a mesenchymal-like to an epithelial-like state.

Ectodysplasin and MC maturation

In contrast to the well-studied stereocilia of hair cells (reviewed by Barr-Gillespie, 2015), virtually
nothing is known about molecular control of MC microvilli assembly. Nevertheless, both hair cell
stereocilia and MC microvilli contain a core of parallel actin bundles and share expression of at
least one actin bundling protein (Sekerková et al., 2004; Sekerková et al., 2006). We previously
found that Eda signaling was partially necessary for MC development along the trunk skin and
that *eda* mutants had a reduced rate of MC production (Brown et al., 2023). Here, we extended
these studies by examining the morphology of dMCs and MCs in *eda* loss of function mutants.
Surprisingly, we found that most MCs completely lacked microvilli in *eda* mutants, a novel MC
phenotype to our knowledge. This observation indicates Eda functions upstream of a MC

microvillar program; dissecting the connections between Eda signaling, MC microvilli, and MC

function represent important areas for further study, and *eda* mutants offer a promising system

to test MC sensory ability in the absence of microvilli. Furthermore, we found that dMCs

developed hyperextended protrusions in *eda* mutants. One interpretation of these data is that

dMC protrusion depends on local dMC and/or MC density. Addressing this possibility will require

rigorous spatial analysis and experimental manipulation independent of Eda signaling.

In summary, we use the live imaging advantages of zebrafish to describe the dynamic cell

behaviors of dMCs and MCs during development and regeneration. We establish that dMCs

represent an intermediate cell state between basal keratinocytes and MCs. We further identify

Eda signaling as essential for the regional morphogenesis of MCs. Finally, we speculate that the

mesenchymal-like behaviors of dMCs may have relevance for understanding the genesis of

Merkel cell carcinoma.

MATERIALS AND METHODS

Animals

Zebrafish and developmental staging

Zebrafish were housed at 26-27℃ on a 14/10 h light cycle. Zebrafish of either sex were used. Strains used: AB (Wild-Type), *(cdh1-tdTomato)^{xt18}* (Cronan et al., 2018), *Tg(atoh1a:nls-Eos)^{w214Tg}* (Pickett et al., 2018), *Tg(atoh1a:lifeact-EGFP)^{w259Tg}, TgBAC(ΔNp63:Cre-ERT2)^{w267Tg}* (Brown et al., 2023), *Tg(actb2:LOXP-BFP-LOXP-DsRed)^{sd27Tg}* (Kobayashi et al., 2014), *Tg(Ola.Sp7:mCherry-Eco.NfsB)^{pd46Tg}* [referred to as *Tg(sp7:mCherry)*] (Singh et al., 2012), and *eda^{dt1261}* (Harris et al., 2008). All zebrafish experiments were approved by the Institutional Animal Care and Use Committee at the University of Washington (protocol #4439-01).

To control for differences in growth rates, zebrafish post-embryonic development was staged based on standard length (SL) (Parichy et al., 2009). SL of fish was measured using the IC Measure software (The Imaging Source) on images captured on a Stemi 508 stereoscope (Zeiss) equipped with a DFK 33UX264 camera (The Imaging Source). *eda* mutants and siblings were sorted by visible phenotype starting at 7 mm SL. Mutants were grown separately from siblings.

Induction of TgBAC(ΔNp63:Cre-ERT2) with 4-OHT

To activate recombination with Cre-ERT2, 1 dpf *TgBAC(ΔNp63:Cre-ERT2)*; *Tg(actb2:LOXP-BFP-LOXP-DsRed); Tg(atoh1a:lifeact-EGFP)* embryos were treated with 10 μM 4-OHT for 24 hr and screened for successful recombination at 3-5 dpf as evidenced by DsRed+ epidermal cells. 4-OHT was prepared as described (Felker et al., 2016).

Scale pluck

A scale pluck protocol was used either to collect scales for immunofluorescence or to induce scale regeneration. To prepare fish for scale pluck, fish were anesthetized in 0.006-0.012% buffered MS-222 (MilliporeSigma, Cat #: E10521) diluted in system water until a surgical plane of anesthesia was reached. Anesthetized fish were moved under a dissecting microscope and placed on the lid of a petri dish. Dumont #5 forceps were used to pluck scales from the trunk in a posterior to anterior procession. Fish were then returned to system water and monitored for full recovery.

Imaging

Scale regeneration time course

Scales were plucked from adult *Tg(atoh1a:lifeact-egfp);Tg(sp7:mcherry)* zebrafish as described above. To facilitate subsequent imaging, the two rows along the midline were plucked. Plucked fish were returned to the recirculating system and imaged using confocal microscopy at the indicated dpp. 1.5x and 8x images were collected for quantification and downstream analyses.

Whole animal photoconversion

476 Prior to scale removal, *Tg(atoh1a:nls-Eos)* zebrafish were exposed to light from a UV LED
477 flashlight (McDoer) for 15 min in a reflective chamber constructed from a styrofoam box lined
478 with aluminum foil. A similar lateral region of the trunk was imaged over subsequent days
479 identified by approximate body position below the dorsal fin and relative to underlying pigment
480 stripes.
481
Confocal imaging
Short-term live imaging of juvenile and adult zebrafish was achieved by anesthetizing fish in
0.006-0.012% MS-222 diluted in system water for ~5 min. Once the fish was immobilized, it was
mounted in a custom imaging chamber and the fish body was secured by carefully adding
molten 1% agarose in system water. Agarose was not applied to the trunk region where imaging
occurred or near the gills and mouth of the fish to ensure survival. Agarose embedded fish were
then covered with MS-222 solution and placed under an A1R MP+ scanhead mounted on a Ni-
E upright microscope (Nikon). Except where noted otherwise, a 16x water dipping objective
(N.A. 0.8) was used for z-stack acquisition and images were post-processed using the
Denoise.ai function in NIS-Elements (Nikon). For short-term imaging, fish were taken off the
microscope after a maximum 25 min of image collection and returned to system water for
recovery.
494
Long-term live imaging of juvenile zebrafish was achieved through an intubation-based protocol
that delivered tricaine-water to immobilized zebrafish using a peristaltic-pump, similar to Xu et
al. (2015). In brief, fish were anesthetized in MS-222 for 8-10 min until gill movement became
very slow. Fish were transferred to a custom imaging chamber and embedded with 1% agarose
as described above. Continuous delivery of MS-222 was achieved by using forceps to gently
insert polyethylene tubing (Becton Dickinson #427421 or #427400) into the fish's mouth. This
delivery line was held in place by modeling clay and 0.08% MS-222 was delivered to the fish by
a peristaltic pump at a flow rate of 2-3 ml per min. Multipoint time lapse imaging was achieved
through Nikon elements software by manually setting large z-stacks for each field of view to
account for sample drift during imaging. Depending on the experiment, time lapse images were
acquired every 3-5 mins for up to 6 h at room temperature. To revive fish after multi-hour time
lapse imaging, the peristaltic pump line was transferred to a bottle containing system water and
fish were closely monitored for recovery. Data acquired from zebrafish that did not survive the
intubation session were excluded from further analysis.
509
510 **Staining**
511
Alizarin Red S staining
Alizarin Red S stains calcium deposits allowing visualization of osteoblast-derived zebrafish
scales. To visualize zebrafish scale development, live animals were stained for 20 min in a
solution of 0.01% (wt/vol) Alizarin Red S (ACROS Organics, Cat #: 400480250) dissolved in
system water and shielded from light, rinsed 3x, 5 mins in system water, then transferred back
into fresh system water as described (Bensimon-Brito et al., 2016).
518
Immunofluorescence

Zebrafish were anesthetized in a solution of 0.012% MS-222 in system water for 2 min.
Approximately 30 scales were plucked and transferred to 1.5 mL tubes containing 375 µl
1xPBS. For fixation, 125 µl of 16% paraformaldehyde (PFA) was added to achieve a final 4%
PFA solution. Scales were incubated for 20 min at room temperature on a gently rotating
platform. PFA was washed out by carefully removing the fixation solution and replacing it with
400 µl of 0.2% PBST (1xPBS + 0.2% Triton X-100). Scales were washed 3x, 5 min with PBST
then blocked with blocking solution (10% normal goat serum in PBST) for 2-3 h at room
temperature. Blocking solution was removed, and 200 µl of primary antibody solution made up
in blocking solution was added. Primary antibodies used: Mouse monoclonal anti-SV2 (DSHB
Cat #: SV2, RRID:AB_2315387) at 1:50; or Rabbit polyclonal anti-GFP (Thermo Fisher
Scientific, Cat #: A11122, RRID:AB_221569) at 1:500. Scales were incubated in primary
antibody overnight at 4℃, protected from light. The next day, scales were washed 4x, 15 min
with PBST before adding secondary antibody made up in blocking solution. Secondary
antibodies used: Goat anti-Mouse Alexa Fluor 647 (Thermo Fisher Scientific, Cat #: A32728,
RRID:AB_2633277) at 1:500; or Goat anti-Rabbit Alexa Fluor 488 (Thermo Fisher Scientific, Cat
#: A32731, RRID:AB_2633280) at 1:1000. Scales were incubated for 2 h at room temperature,
protected from light. Secondary solution was washed out 4x, 15 min using PBST. To visualize
nuclei, 5 ng/µl DAPI (MilliporeSigma, Cat #: 508741) was added. DAPI was washed out 4x, 5
min using PBST. Scales were mounted epidermis-side up between a microscope slide and
coverslip in ProLong Gold (Thermo Fisher Scientific, Cat #: P36930). Imaging was performed
with a 40x (NA 1.3) oil immersion objective.
541
542   *Hybridization chain reaction*
543   HCR on adult zebrafish scales using a custom *piezo2* probe set (Molecular Instruments;
544   accession #: XM_021468270.1; set size: 20; amplifier: B3) was performed as described (Brown
545   et al., 2023). Scales were imaged with a 25x water immersion objective (N.A. 1.1)
546
EdU labeling during scale regeneration
Scales were plucked from adult *Tg(atoh1a:lifeact-egfp);Tg(sp7:mCherry)* zebrafish to induce
scale regeneration. Fish were returned to their tanks for recovery prior to EdU injection.
Regenerating fish were anesthetized in MS-222 and placed ventral side up on a sponge
situated under a dissecting microscope. 10 µl of 10 mM EdU (Thermo Fisher Scientific, Cat #:
C10640) was injected intraperitoneally in the region between the pelvic fins. Fish were
transferred back to system water for recovery. Approximately 20 h later, these fish had their
regenerating scales plucked and subjected to EdU staining following the manufacturer's
instructions. If downstream immunofluorescence was required, this was performed after EdU
detection. Stained scales were imaged with a 40x (NA 1.3) oil immersion objective.
557
558   **Image and statistical analysis**
559
Image processing
Image processing was performed using FIJI/ImageJ (Schindelin et al., 2012) or Imaris (Oxford
Instruments). Unless otherwise indicated, all images were gathered through z-stack acquisition
and displayed as maximum intensity projections. Throughout the figures, single channel images

of *Tg(atoh1a:lifeact-egfp)* are color-coded according to signal intensity using the "Green Fire
Blue" lookup table in FIJI/ImageJ.

Cell counting & shape analysis
atoh1a+ cells were classified as dMCs or MCs as follows: Confocal z-stacks were max
projected and the ImageJ "Cell Counter" function was used to manually classify *atoh1a+* cells
as dMC or MC based on cell morphology. For shape analysis, max projected images were
thresholded using Huang's algorithm. Using the "Analyze Particles…" function, circularity and
roundness were calculated. In Figure 1N, "circular" cells were defined as those having circularity
and roundness values >0.7.

573
574 *Fluorescence intensity analysis*
575 Fluorescence intensity of MCs and dMCs was calculated from confocal z-stacks acquired using
576 identical settings by Corrected Total Cell Fluorescence (CTCF) in ImageJ. CTCF = Integrated
577 Density – (Area of selected cell X Mean fluorescence of background readings).

578
579 *z-depth analysis*
580 For the z-depth analysis in Figure 5H, confocal z-stacks were acquired using a 25x water
581 dipping objective (N.A. 1.1) and a z-step of 0.3 μm. The outer surface of the periderm was
582 defined as a depth of 0. The distance from the center of the cell bodies of dMCs and MCs to the
583 periderm surface was measured on reconstructed yz slices using the line tool in ImageJ.

584
Cell tracking
To quantify cell motility metrics, the trackmate plugin in ImageJ was used. z-stack acquired
timelapse images were imported and made into max projections. To correct drift encountered
during long term imaging acquisition, the Imagej plugin "correct 3D drift" was used to correct xy
drift. Trackmate was opened and manual detection was performed by tracking individual cells of
interest. The metrics calculated from trackmate (Total distance traveled, confinement ratio, etc)
are explained https://imagej.net/plugins/trackmate/algorithms.

592
Statistical analysis
Statistical analysis and graphing were performed using R (R Core Team, 2023), MATLAB
(MathWorks) or GraphPad (Prism). Statistical tests used and sample numbers are described in
the corresponding figure legend.

ACKNOWLEDGEMENTS
We thank the LSB Aquatics staff for animal care and Dr. David Tobin for sharing zebrafish. The
authors are grateful to all members of the Rasmussen lab for discussion, technical assistance,
and support. The SV2 monoclonal antibody used in this study was obtained from the
Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at
The University of Iowa, Department of Biology, Iowa City, IA 52242.

AUTHOR CONTRIBUTIONS
Evan W. Craig: Conceptualization, Investigation, Methodology, Visualization, Writing - Original
Draft, Funding acquisition; **Erik C. Black**: Investigation, Methodology; **Camille E.A. Goo**:

Investigation, Methodology; **Avery Angell Swearer**: Investigation, Methodology; **Nathaniel G.**
Yee: Investigation, Methodology; **Jeffrey P. Rasmussen**: Conceptualization, Investigation,
Methodology, Validation, Supervision, Visualization, Writing - Original Draft, Funding acquisition.
611

COMPETING INTERESTS

612 No competing interests declared.

613
614

FUNDING

This work was funded in part by a Graduate Research Fellowship (DGE-2140004) from the
National Science Foundation to EWC, an ISCRM Graduate Fellowship to ECB, R01-HD107108
from the Eunice Kennedy Shriver National Institute of Child Health and Human Development to
JPR, and a New Investigator Award from the Fred Hutch/University of Washington/Seattle
Children's Cancer Consortium, which is supported by the NIH/NCI Cancer Center Support Grant
P30 CA015704, to JPR. JPR is a Washington Research Foundation Distinguished Investigator.
622

DATA AVAILABILITY

624 All data needed to evaluate the conclusions in the paper are present in the paper and/or the
625 Supplementary Materials.

FIGURES AND FIGURE LEGENDS

Figure 1 - Craig et al.

Figure 1. Identification of a transient, morphologically distinct population of keratinocyte-derived *atoh1a*+ cells during skin development

(A) Schematic depicting live-imaging methodology for visualizing MCs (green) within the epidermis atop scales along the lateral zebrafish trunk skin. **(B)** Illustration showing organization of adult zebrafish trunk skin in cross section. Note the stratified layers of keratinocytes above

the bony scale. Merkel cells are shown in green near the outermost epidermal layer. **(C)** The
Tg(atoh1a:lifeact-egfp) reporter allows visualization of the MC F-actin cytoskeleton, including
short and spiky microvilli structures. **(D-H)** Representative confocal micrographs of *atoh1a+*
MCs (green; *Tg(atoh1a:lifeact-egfp)*)) and scales (magenta; Alizarin Red S staining) along the
lateral trunk in juvenile to adult zebrafish of the indicated stages. White dashed boxes indicate
regions magnified in (D'-H'). **(D'-H')** *Tg(atoh1a:lifeact-egfp)* signal intensity is color-coded using
the "Green Fire Blue" lookup table. MCs denoted by arrowheads, dMCs denoted by arrows. **(I)**
The percentage of *atoh1a+* cells with dMC morphology out of total number of *atoh1a+* cells
plotted against SL. Each dot represents the mean dMC percentage calculated from at least *n*=3
animals of that size range. For each individual zebrafish, multiple confocal micrographs were
acquired from the trunk region at 1.5x magnification and *atoh1a+* cells were manually
categorized as "MC" or "dMC" based on morphology and grouped for input data to calculate the
dMC ratio shown. **(J)** Plot of dMC density relative to SL from the dataset analyzed in (I). **(K)** Plot
of total *atoh1a+* cell density (dMCs and MCs) relative to SL from the dataset analyzed in (I).
(L,M) *atoh1a+* cell shape analysis for animals of the indicated stages. Black dots represent
individual *atoh1a+* cells that were assigned values for circularity and roundness after confocal
micrographs of *Tg(atoh1a:lifeact-egfp)* were thresholded to acquire *atoh1a+* cell shape outlines.
Top right quadrant indicates cells with the most round and circular morphologies, indicative of
mature MCs. Cells falling outside of the top right quadrant possess less round and circular
morphologies, indicative of dMCs. *n*=100 randomly selected cells displayed for each stage. **(N)**
Stacked barcharts of *atoh1a+* cell shape analysis performed in L-M. *atoh1a+* cells in adult skin
have significantly more circular morphologies than *atoh1a+* cells in juvenile skin (****, P<0.0001;
Fisher's Exact Test). **(O-O''')** Confocal micrographs of the trunk epidermis from a
TgBAC(ΔNp63:Cre-ERT2); Tg(actb2:LOXP-BFP-LOXP-DsRed); Tg(atoh1a:lifeact-egfp) juvenile
treated with 4-OHT at 1 dpf. Note the mosaic DsRed expression (magenta) in basal
keratinocytes and derivatives. DsRed+ dMCs shown in O'-O'', along with a DsRed+ MC in O'''
indicate both MC-types derive from a shared basal keratinocyte progenitor population. nm,
neuromasts containing clusters of *atoh1a+* hair cells. Error bars in (I,J,K) display SD of mean.
Scale bars: 50 μm (D-H), 20 μm (O), and 5 μm (D'-H',O'-O''').

Figure 2. dMCs are the primary *atoh1a*+ cell type during the early stages of skin regeneration

(A) Illustration of the zebrafish scale pluck regeneration model. Forceps are used to physically "pluck" scales from the adult zebrafish trunk to induce injury and trigger regeneration of dermal scales and overlying epidermis. **(B-E)** Representative confocal micrographs of scale-forming osteoblasts (magenta; *Tg(sp7:mCherry)*) at the indicated stages of regeneration. **(F-I)** Representative confocal micrographs of *atoh1a*+ cell shapes within the scale epidermis at the indicated stages of scale regeneration. Yellow arrows denote MCs, green arrows denote dMCs, white arrow denotes an autofluorescent pigment cell. White dashed boxes indicate regions magnified in (F',G'). **(J)** Plot of total *atoh1a*+ cell density (dMCs and MCs) over the course of scale regeneration. Each dot represents confocal images collected and analyzed from multiple zebrafish of the corresponding timepoint (2-7 dpp, *n*=8-13; 9-14 dpp, *n*=2-4). **(K)** Quantification of dMC frequency during scale regeneration. Each dot represents the mean dMC frequency from the dataset analyzed in (J). **(L)** Illustration depicting the design of the *atoh1a*+ cell

photoconversion experiment. *Tg(atoh1a:nls-Eos)* expresses a nuclear-localized and
photoconvertible Eos protein in *atoh1a+* cells (dMCs and MCs). *atoh1a+* cells (green) in
uninjured scales are irreversibly photoconverted from green to magenta using UV light. Scales
containing photoconverted cells are then plucked to induce regeneration. Pre-existing *atoh1a+*
cells will contain photoconverted nls-Eos (magenta) in the new scale region whereas new
atoh1a+ cells will contain only non-photoconverted nls-Eos (green). **(M-P)** Representative
confocal micrographs of the photoconverted *Tg(atoh1a:nls-Eos)* scale epidermis pre-scale pluck
(M) and post-pluck (N-P). Arrowheads denote pre-existing cells (containing photoconverted nls-
Eos), arrows denote *de novo* generated cells (containing only non-photoconverted nls-Eos). **(Q)**
Quantification of pre-existing (magenta line) and new (green line) *atoh1a+* cells from
photoconversion experiment at the indicated stages of regeneration. Each dot represents the
mean *atoh1a+* cell density from *n*=3-4 fish. Scale bars, 100 µm (B-E), 10 µm (F-I), 5 µm (F', G'),
and 10 µm (M-P).

Figure 3. dMCs share molecular features with keratinocytes and MCs

(**A-B)** Representative micrographs of MCs and dMCs within the 5 dpp regenerating scale
epidermis of a *Tg(atoh1a:lifeact-egfp)* adult visualized with anti-Tp63 (red) and anti-GFP (green)
antibodies. Mature MCs are Tp63 negative (A) whereas dMCs are often weakly Tp63 positive

(B). Dashed lines outline the nuclei. Asterisks indicate basal keratinocyte nuclei. **(C)** Violin plots
of Tp63 staining intensity in MCs, dMCs, and basal keratinocytes. Confocal micrographs
acquired with identical imaging parameters were analyzed for Corrected Total Cell
Fluorescence (CTCF) measured in arbitrary units. Each dot represents an individual cell (*n*=9
MCs, *n*=16 dMCs, *n*=14 basal keratinocytes). A one-way ANOVA with post-hoc Tukey HSD test
was used to compare Tp63 staining levels between cell types. **(D-E)** Representative
micrographs showing MCs and dMCs within the 5 dpp regenerating scale epidermis of a
Tg(atoh1a:lifeact-egfp) adult stained with anti-SV2 (magenta) and anti-GFP (green) antibodies.
MCs are always SV2 positive (D), whereas some dMCs (E) are SV2 positive. Dashed lines
outline the cell bodies. **(F)** Violin plots of SV2 staining intensity quantified using CTCF in MCs
and dMCs. Each dot represents an individual cell (*n*=16 MCs, *n*=22 dMCs). A non-parametric
Mann-Whitney test was used to compare SV2 staining levels between cell types. **(G)**
Representative micrographs of MCs and dMCS within the 5 dpp regenerating scale epidermis of
a *Tg(atoh1a:lifeact-egfp)* adult stained with an anti-*piezo2* HCR probeset (magenta) and an anti-
GFP (green) antibody. Dashed box indicates region magnified in inset. Both MCs and dMCs are
piezo2 positive. **(H)** Violin plots of *piezo2* HCR staining intensity in MCs and dMCs quantified
using CTCF. Each dot represents an individual cell (*n*=19 MCs, *n*=22 dMCs). A non-parametric
Mann-Whitney test was used to compare *piezo2* staining levels between cell types. *, P<0.05;
****, P<0.0001. Scale bars, 5 µm (A-B,D-E,G inset) and 20 µm (G).

Figure 4. MC and dMC protrusions are dynamic and differentially polarized in the skin

(A-A'''') Still images of a timelapse of a single MC expressing *Tg(atoh1a:lifeact-egfp)*. White
arrowheads point to microvilli extension, retraction, or merging events. **(B-B'''')** Still images of a
timelapse of a single dMC visualized with *Tg(atoh1a:lifeact-egfp)*, image acquisition as
described in (A). Note the longer, filopodial-like protrusions (white arrows) and highly
amorphous cell body. **(C-E)** Representative maximum intensity projection (C) or individual z-
slices (D,E) of the scale epidermis in an animal expressing *Tg(atoh1a:lifeact-egfp)* to label
dMCs and MCs (green) and *(cdh1-tdTomato)* to label lateral cell membranes (magenta). **(D',E')**
Reconstructed xz slices along the yellow lines in D and E. Dashed lines indicate the outer and
inner epidermal margins. Note that the MCs in D' are located near the upper, periderm layer of
the skin with basal facing protrusions (arrowhead), whereas the dMC in E' is located near the
lower, basal layer of the skin with a lateral facing protrusion (arrow). Brightness and contrast
were adjusted in (E,E') to better illustrate the dMC morphology. **(F)** Polar histograms of MC and
dMC protrusion angles. Lifeact-EGFP+ protrusions that could be individually resolved in 3D
were measured relative to the z-axis of the epidermis using the coordinate system diagrammed

in (G). Shaded regions indicate the number of protrusions falling in that angle range. Plots show

n=104 protrusions from 10 MCs and *n*=32 protrusions from 11 dMCs. **(G)** Stacked bar charts

depicting results in (F), with protrusions binned based on orientation. A χ^2 test was used to

compare between cell types (****, P<0.0001; χ^2 statistic, 18.9903). **(H)** Violin plots of z-depths of

MCs and dMCs measured relative to the periderm surface in the scale epidermis of *n*=130 MCs

and *n*=80 dMCs from 5 fish (11-12 mm SL). The average depth of the basal surface of basal

keratinocytes in the data set is indicated by a horizontal dashed line (-16.4 µm). A non-

parametric Mann-Whitney test was used to compare between cell types (****, P<0.0001). Scale

bars, 5 µm (A-C).

Figure 5. dMCs are motile cells that can directly mature into MCs in developing zebrafish skin

(A) Cell tracks (magenta) of individual MCs and dMCs traced over time. **(B)** Dot plot of cell speed (µm/hour) of individually tracked cells (*n*=60 MCs and *n*=74 dMCs). A non-parametric Mann-Whitney test was used to compare speed between cell types (****, P<0.0001). **(C)** Cell tracks scored for confinement ratio where values near 0 are indicative of confined movement and values near 1 are indicative of a linear trajectory (*n*=74 dMCs). **(D)** Dot plot of dMC track displacement, which measures the distance between the starting and ending point of each cell track, and total distance traveled (*n*=74 dMCs). A paired Mann-Whitney test was used to compare displacement and distance (****, P<0.0001). In (B-D), horizontal lines indicate the mean and error bars indicate the SD. **(E)** Representative timelapse stills from *Tg(atoh1a:lifeact-egfp)* expressing juvenile zebrafish skin. Magenta arrowheads highlight dMCs in their migratory configuration with unipolar protrusions. Orange arrowheads highlight dMCs in their exploratory or immotile configuration with multipolar protrusions. Note that individual cells can switch between the unipolar and multipolar configurations. **(F)** Representative timelapse stills of a dMC

to MC maturation event. Arrow indicates dMC protrusion retraction and arrowhead indicates

formation of microvilli. Note the cell body switches from an ovoid to spherical shape during this

transition. **(G)** Representative timelapse stills of a MC to dMC reversion event. Arrowhead

indicates microvilli and arrow indicates filopodial-like protrusion extension. **(H)** Schematic

depicting proposed model of dMC maturation events described in this study. (1) dMCs emerge

in lower epidermal strata from *ΔNp63+* basal keratinocyte progenitors; (2) dMCs are motile cells

that migrate in the direction of their protrusions and can undergo contact inhibition upon

encountering another dMC; (3) dMCs can directly adopt the mature MC morphology in upper

epidermal strata; and (4) in rare instances, MCs can revert to a dMC morphology. Scale bars,

10 μm (A), 5 μm (A, insets), 10 μm (E, E, insets), and 5 μm (F,G).

Figure 6 - Craig et al.

Figure 6. Genetic loss of Eda results in altered dMC and MC morphologies in trunk skin

(A,B) Representative confocal images of MCs within juvenile trunk epidermis of animals of the indicated genotypes and stages. Dashed boxes indicate cells magnified in (A',A'',B',B''). Arrowheads indicate microvilli and asterisks indicate intense Lifeact-EGFP signal forming a smooth, ring-like cortical structure evident in cross section in the $eda^{-/-}$ mutant epidermis. **(C,D)** Representative confocal images of MCs within corneal epidermis of animals of the indicated genotypes. Dashed boxes indicate cells magnified in (C',C'',D',D''). Arrowheads indicate microvilli. **(E,F)** Representative confocal images of dMCs within the trunk epidermis of animals of the indicated genotypes and stages. Yellow arrow in E indicates the longest protrusion on a dMC in eda sibling epidermis. White arrow in F indicates the longest protrusion on a dMC in $eda^{-/-}$ mutant epidermis. **(G)** Quantification of percentage of MCs with discernable microvilli in trunk or corneal epidermis of animals of the indicated genotypes. Each dot indicates a juvenile (9-20 mm SL) zebrafish of the respective genotype where a collection of images were analyzed (sibling trunk, n=386 cells from 3 fish; $eda^{-/-}$ trunk, n=235 cells from 4 fish; sibling cornea: n=169 cells from 4 fish; $eda^{-/-}$ cornea, n=332 cells from 4 fish). Fisher's exact test shows a significant difference between genotypes in the trunk but not cornea (****, P<0.0001; n.s., P=0.3413). **(H)** Violin plots of the longest dMC Lifeact-EGFP+ protrusion within the trunk epidermis of juveniles of the indicated genotypes. Each dot represents an individual cell (siblings, n=22 dMCs; $eda^{-/-}$, n=47 dMCs). A non-parametric Mann-Whitney test (****, P<0.0001) was used to compare between cell types. Scales bars, 10 µm (A-F), and 5 µm (A'-A''', B'-B''', C'-C'', D'-D'').

 SUPPLEMENTAL FIGURE AND LEGEND

Figure S1 - Craig et al.

Figure S1: Differences in cell cycle state of dMCs and MCs

(A-C) Still images from timelapses showing cell divisions of dMCs expressing *Tg(atoh1a:lifeact-egfp)* in the juvenile (A) or regenerating scale epidermis (B,C). Arrowheads denote cytokinetic furrows and arrows denote daughter cells. **(D)** Experimental design of EdU administration and imaging during scale regeneration. **(E,F)** Representative confocal images of EdU and anti-GFP staining in regenerating *Tg(atoh1a:lifeact-egfp)* scales. White dashed lines indicate nuclear outlines. Scale bars, 10 μm (A-C) and 5 μm (E-F).

SUPPLEMENTAL MOVIE LEGENDS

Movie 1. Microvilli dynamics in zebrafish MCs. Timelapse movie of MCs expressing *Tg(atoh1a:lifeact-egfp)* in juvenile zebrafish skin. Note spike-like microvilli that extend, retract, and coalesce. Scale bar, 5 µm.

Movie 2. Actin dynamics in zebrafish dMCs. Timelapse movie of a dMC expressing *Tg(atoh1a:lifeact-egfp)* in juvenile zebrafish skin. Note thread-like filopodial protrusions that are highly dynamic. Scale bar, 10 µm.

Movie 3. Long-term imaging reveals dMC motile behaviors. 6 hour timelapse movie of MCs and dMCs expressing *Tg(atoh1a:lifeact-egfp)* in an intubated juvenile zebrafish. Manual cell tracks (white traces) show dMCs crawling in the direction of their protrusions while MCs remain largely stationary. Scale bar, 10 µm.

Movie 4. dMC-dMC contact inhibition. Timelapse of a dMC-dMC contact (indicated by arrow at time 0 min) that results in both cells migrating in opposing directions. Scale bars, 10 µm.

Movie 5. *In vivo* dMC cell division in juvenile zebrafish skin. Timelapse movie of a dMC expressing *Tg(atoh1a:lifeact-egfp)* undergoing cell division. In all dMC division events recorded, dividing dMCs retract a single unipolar protrusion, form a circular and actin-rich membrane, and undergo cytokinesis. Daughter cells then rapidly extend filopodial protrusions and migrate opposite from one another. Scale bar, 10 µm.

Movie 6. dMC to MC maturation event. Timelapse of single dMC expressing *Tg(atoh1a:lifeact-egfp)* that withdraws its filopodial protrusions, rounds its cell cortex, and extends small microvillar extensions as observed in mature MCs. Scale bar, 5 µm.

Movie 7. dMC behaviors during regeneration. Timelapse of the regenerating scale epidermis at 3 dpp in a *Tg(atoh1a:lifeact-egfp)* zebrafish. Labels indicate different dMC behaviors including maturation, motility, and cell death. Scale bar, 10 µm.

Movie 8. MC to dMC reversion event. Timelapse of developing *Tg(atoh1a:lifeact-egfp)* expressing zebrafish skin with a mix of MCs and dMCs in the field of view. Note a cell with a MC morphology begins to transition into a dMC-like cell with a long unipolar protrusion. Scale bar, 10 µm.

REFERENCES

Aman, A. J., Fulbright, A. N. and Parichy, D. M. (2018). Wnt/β-catenin regulates an ancient
signaling network during zebrafish scale development. *Elife* **7**,.

Bakkers, J., Hild, M., Kramer, C., Furutani-Seiki, M. and Hammerschmidt, M. (2002).
Zebrafish DeltaNp63 is a direct target of Bmp signaling and encodes a transcriptional
repressor blocking neural specification in the ventral ectoderm. *Dev Cell* **2**, 617–627.

Barr-Gillespie, P.-G. (2015). Assembly of hair bundles, an amazing problem for cell biology.
Mol Biol Cell **26**, 2727–2732.

Becker, J. C., Stang, A., DeCaprio, J. A., Cerroni, L., Lebbé, C., Veness, M. and Nghiem, P.
(2017). Merkel cell carcinoma. *Nat Rev Dis Primers* **3**, 17077.

**Bensimon-Brito, A., Cardeira, J., Dionísio, G., Huysseune, A., Cancela, M. L. and Witten,
P. E.** (2016). Revisiting in vivo staining with alizarin red S--a valuable approach to
analyse zebrafish skeletal mineralization during development and regeneration. *BMC
Dev Biol* **16**, 2.

Boot, P. M., Rowden, G. and Walsh, N. (1992). The distribution of Merkel cells in human fetal
and adult skin. *Am J Dermatopathol* **14**, 391–396.

**Brown, T. L., Horton, E. C., Craig, E. W., Goo, C. E. A., Black, E. C., Hewitt, M. N., Yee, N.
G., Fan, E. T., Raible, D. W. and Rasmussen, J. P.** (2023). Dermal appendage-
dependent patterning of zebrafish atoh1a+ Merkel cells. *Elife* **12**, e85800.

Burgess, P. R., English, K. B., Horch, K. W. and Stensaas, L. J. (1974). Patterning in the
regeneration of type I cutaneous receptors. *J Physiol* **236**, 57–82.

**Cai, E., Marchuk, K., Beemiller, P., Beppler, C., Rubashkin, M. G., Weaver, V. M., Gérard,
A., Liu, T.-L., Chen, B.-C., Betzig, E., et al.** (2017). Visualizing dynamic microvillar
search and stabilization during ligand detection by T cells. *Science* **356**, eaal3118.

Cox, B. D., De Simone, A., Tornini, V. A., Singh, S. P., Di Talia, S. and Poss, K. D. (2018). In
Toto Imaging of Dynamic Osteoblast Behaviors in Regenerating Skeletal Bone. *Curr Biol*
28, 3937-3947.e4.

**Cronan, M. R., Matty, M. A., Rosenberg, A. F., Blanc, L., Pyle, C. J., Espenschied, S. T.,
Rawls, J. F., Dartois, V. and Tobin, D. M.** (2018). An explant technique for high-
resolution imaging and manipulation of mycobacterial granulomas. *Nat Methods* **15**,
1098–1107.

**Daghfous, G., Auclair, F., Blumenthal, F., Suntres, T., Lamarre-Bourret, J., Mansouri, M.,
Zielinski, B. and Dubuc, R.** (2020). Sensory cutaneous papillae in the sea lamprey
(Petromyzon marinus L.): I. Neuroanatomy and physiology. *J Comp Neurol* **528**, 664–
686.

**De Simone, A., Evanitsky, M. N., Hayden, L., Cox, B. D., Wang, J., Tornini, V. A., Ou, J.,
Chao, A., Poss, K. D. and Di Talia, S.** (2021). Control of osteoblast regeneration by a
train of Erk activity waves. *Nature* **590**, 129–133.

Düring, M. V. and Andres, K. H. (1976). The ultrastructure of taste and touch receptors of the
frog's taste organ. *Cell Tissue Res* **165**, 185–198.

English, K. B. (1974). Cell types in cutaneous type I mechanoreceptors (Haarscheiben) and
their alterations with injury. *Am J Anat* **141**, 105–126.

English, K. B. (1977). Morphogenesis of Haarscheiben in rats. *J Invest Dermatol* **69**, 58–67.

**Felker, A., Nieuwenhuize, S., Dolbois, A., Blazkova, K., Hess, C., Low, L. W. L., Burger, S.,
Samson, N., Carney, T. J., Bartunek, P., et al.** (2016). In vivo performance and
properties of tamoxifen metabolites for CreERT2 control. *PLoS One* **11**, e0152989.

Garant, P. R., Feldman, J., Cho, M. I. and Cullen, M. R. (1980). Ultrastructure of Merkel cells
in the hard palate of the squirrel monkey (Saimiri sciureus). *Am J Anat* **157**, 155–167.

Girard, D., Laverdet, B., Buhé, V., Trouillas, M., Ghazi, K., Alexaline, M. M., Egles, C.,

Misery, L., Coulomb, B., Lataillade, J.-J., et al. (2017). Biotechnological Management
of Skin Burn Injuries: Challenges and Perspectives in Wound Healing and Sensory
Recovery. *Tissue Eng Part B Rev* **23**, 59–82.

Guzman, A., Ramos-Balderas, J. L., Carrillo-Rosas, S. and Maldonado, E. (2013). A stem
cell proliferation burst forms new layers of P63 expressing suprabasal cells during
zebrafish postembryonic epidermal development. *Biol. Open* **2**, 1179–1186.

Hahn, J. M., Combs, K. A., Lloyd, C. M., McFarland, K. L., Boyce, S. T. and Supp, D. M.
(2019). Identification of Merkel cells associated with neurons in engineered skin
substitutes after grafting to full thickness wounds. *PLoS One* **14**, e0213325.

Halata, Z. (1975). The mechanoreceptors of the mammalian skin ultrastructure and
morphological classification. *Advances In Anatomy, Embryology and Cell Biology* **50**, 3–
77.

Harms, P. W., Harms, K. L., Moore, P. S., DeCaprio, J. A., Nghiem, P., Wong, M. K. K.,
Brownell, I., and International Workshop on Merkel Cell Carcinoma Research
(IWMCC) Working Group (2018). The biology and treatment of Merkel cell carcinoma:
current understanding and research priorities. *Nat Rev Clin Oncol* **15**, 763–776.

Harris, M. P., Rohner, N., Schwarz, H., Perathoner, S., Konstantinidis, P. and Nüsslein-
Volhard, C. (2008). Zebrafish eda and edar mutants reveal conserved and ancestral
roles of ectodysplasin signaling in vertebrates. *PLoS Genet* **4**, e1000206.

Hartschuh, W., Weihe, E. and Reinecke, M. (1986). The Merkel Cell. In *Biology of the*
Integument: 2 Vertebrates (ed. Bereiter-Hahn, J.), Matoltsy, A. G.), and Richards, K. S.),
pp. 605–620. Berlin, Heidelberg: Springer Berlin Heidelberg.

Hashimoto, K. (1972). Fine structure of Merkel cell in human oral mucosa. *J Invest Dermatol*
58, 381–387.

Horch, K. (1982). The influence of mechanoreceptor structures on regenerating sensory axons
after cutaneous nerve transection in the cat. *Neurosci Lett* **32**, 281–284.

Iggo, A. and Muir, A. R. (1969). The structure and function of a slowly adapting touch
corpuscle in hairy skin. *J Physiol* **200**, 763–796.

Iwasaki, M., Kuroda, J., Kawakami, K. and Wada, H. (2018). Epidermal regulation of bone
morphogenesis through the development and regeneration of osteoblasts in the
zebrafish scale. *Dev Biol* **437**, 105–119.

Kim, D. K. and Holbrook, K. A. (1995). The appearance, density, and distribution of Merkel
cells in human embryonic and fetal skin: their relation to sweat gland and hair follicle
development. *J Invest Dermatol* **104**, 411–416.

Kobayashi, I., Kobayashi-Sun, J., Kim, A. D., Pouget, C., Fujita, N., Suda, T. and Traver, D.
(2014). Jam1a-Jam2a interactions regulate haematopoietic stem cell fate through Notch
signalling. *Nature* **512**, 319–23.

Landmann, L. and Halata, Z. (1980). Merkel cells and nerve endings in the labial epidermis of
a lizard. *Cell Tissue Res* **210**, 353–357.

Lane, E. B. and Whitear, M. (1977). On the occurrence of Merkel cells in the epidermis of
teleost fishes. *Cell and Tissue Research* **182**, 235–46.

Lee, H. and Kimelman, D. (2002). A dominant-negative form of p63 is required for epidermal
proliferation in zebrafish. *Dev Cell* **2**, 607–616.

Meenderink, L. M., Gaeta, I. M., Postema, M. M., Cencer, C. S., Chinowsky, C. R.,
Krystofiak, E. S., Millis, B. A. and Tyska, M. J. (2019). Actin Dynamics Drive
Microvillar Motility and Clustering during Brush Border Assembly. *Dev Cell* **50**, 545-
556.e4.

Merot, Y. and Saurat, J. H. (1988). Proliferation of Merkel cells in the skin. *Acta Derm Venereol*
68, 366–367.

Moll, R., Moll, I. and Franke, W. W. (1984). Identification of Merkel cells in human skin by
specific cytokeratin antibodies: changes of cell density and distribution in fetal and adult

plantar epidermis. *Differentiation* **28**, 136–154.

Moll, I., Zieger, W. and Schmelz, M. (1996). Proliferative merkel cells were not detected in
human skin. *Arch Dermatol Res* **288**, 184–187.

Morrison, K. M., Miesegaes, G. R., Lumpkin, E. A. and Maricich, S. M. (2009). Mammalian
Merkel cells are descended from the epidermal lineage. *Developmental Biology* **336**, 76–
83.

Nakafusa, J., Narisawa, Y., Shinogi, T., Taira, K., Tanaka, T., Inoue, T. and Misago, N.
(2006). Changes in the number of Merkel cells with the hair cycle in hair discs on rat
back skin. *Br J Dermatol* **155**, 883–889.

Nurse, C. A., Macintyre, L. and Diamond, J. (1984). Reinnervation of the rat touch dome
restores the Merkel cell population reduced after denervation. *Neuroscience* **13**, 563–
571.

Ostrowski, S. M., Wright, M. C., Bolock, A. M., Geng, X. and Maricich, S. M. (2015). Ectopic
Atoh1 expression drives Merkel cell production in embryonic, postnatal and adult mouse
epidermis. *Development* **142**, 2533–2544.

Parichy, D. M., Elizondo, M. R., Mills, M. G., Gordon, T. N. and Engeszer, R. E. (2009).
Normal table of postembryonic zebrafish development: staging by externally visible
anatomy of the living fish. *Developmental Dynamics* **238**, 2975–3015.

Pickett, S. B., Thomas, E. D., Sebe, J. Y., Linbo, T., Esterberg, R., Hailey, D. W. and Raible,
D. W. (2018). Cumulative mitochondrial activity correlates with ototoxin susceptibility in
zebrafish mechanosensory hair cells. *Elife* **7**,.

R Core Team (2023). *R: A Language and Environment for Statistical Computing*. Vienna,
Austria: R Foundation for Statistical Computing.

Rangel-Huerta, E., Guzman, A. and Maldonado, E. (2021). The dynamics of epidermal
stratification during post-larval development in zebrafish. *Dev Dyn* **250**, 175–190.

Richardson, R., Metzger, M., Knyphausen, P., Ramezani, T., Slanchev, K., Kraus, C.,
Schmelzer, E. and Hammerschmidt, M. (2016). Re-epithelialization of cutaneous
wounds in adult zebrafish combines mechanisms of wound closure in embryonic and
adult mammals. *Development* **143**, 2077–88.

Sadier, A., Viriot, L., Pantalacci, S. and Laudet, V. (2014). The ectodysplasin pathway: from
diseases to adaptations. *Trends Genet* **30**, 24–31.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. *Nature Methods* **9**, 676–82.

Sekerková, G., Zheng, L., Loomis, P. A., Changyaleket, B., Whitlon, D. S., Mugnaini, E.
and Bartles, J. R. (2004). Espins are multifunctional actin cytoskeletal regulatory
proteins in the microvilli of chemosensory and mechanosensory cells. *J Neurosci* **24**,
5445–5456.

Sekerková, G., Zheng, L., Loomis, P. A., Mugnaini, E. and Bartles, J. R. (2006). Espins and
the actin cytoskeleton of hair cell stereocilia and sensory cell microvilli. *Cell Mol Life Sci*
63, 2329–2341.

Sharkova, M., Chow, E., Erickson, T. and Hocking, J. C. (2023). The morphological and
functional diversity of apical microvilli. *J Anat* **242**, 327–353.

Singh, S. P., Holdway, J. E. and Poss, K. D. (2012). Regeneration of amputated zebrafish fin
rays from de novo osteoblasts. *Developmental Cell* **22**, 879–86.

Sire, J.-Y., Quilhac, A., Bourguignon, J. and Allizard, F. (1997). Evidence for participation of
the epidermis in the deposition of superficial layer of scales in zebrafish (Danio rerio): A
SEM and TEM study. *J Morphol* **231**, 161–174.

Stella, M., Calcagni, M., Teich-Alasia, S., Ramieri, G., Cellino, G. and Panzica, G. (1994).
Sensory endings in skin grafts and scars after extensive burns. *Burns* **20**, 491–495.

Tachibana, T. and Ishizeki, K. (1981). Merkel cell development in the wound healing in the

labial mucosa of adult rabbits. *Arch Histol Jpn* **44**, 151–165.

Tachibana, T. and Nawa, T. (1980). Merkel cell differentiation in the labial mucous epithelium
of the rabbit. *J Anat* **131**, 145–155.

**Tachibana, T., Yamamoto, H., Takahashi, N., Kamegai, T., Shibanai, S., Iseki, H. and
Nawa, T.** (1997). Polymorphism of Merkel cells in the rodent palatine mucosa:
immunohistochemical and ultrastructural studies. *Arch Histol Cytol* **60**, 379–389.

Tachibana, T., Kamegai, T., Takahashi, N. and Nawa, T. (1998). Evidence for polymorphism
of Merkel cells in the adult human oral mucosa. *Arch Histol Cytol* **61**, 115–124.

Takahashi-Iwanaga, H. (2003). The Three-Dimensional Microanatomy of Merkel Cells. In *The
Merkel Cell* (ed. Baumann, K. I.), Halata, Z.), and Moll, I.), pp. 7–12. Berlin, Heidelberg:
Springer Berlin Heidelberg.

Takahashi-Iwanaga, H. and Abe, K. (2001). Scanning electron microscopic observation of
Merkel cells in the lamprey epidermis. *Kaibogaku Zasshi* **76**, 375–380.

Toyoshima, K., Seta, Y., Takeda, S. and Harada, H. (1998). Identification of Merkel cells by an
antibody to villin. *Journal of Histochemistry & Cytochemistry* **46**, 1329–34.

Vaigot, P., Pisani, A., Darmon, Y. M. and Ortonne, J. P. (1987). The majority of epidermal
Merkel cells are non-proliferative: a quantitative immunofluorescence analysis. *Acta
Derm Venereol* **67**, 517–520.

**Van Keymeulen, A., Mascre, G., Youseff, K. K., Harel, I., Michaux, C., De Geest, N.,
Szpalski, C., Achouri, Y., Bloch, W., Hassan, B. A., et al.** (2009). Epidermal
progenitors give rise to Merkel cells during embryonic development and adult
homeostasis. *Journal of Cell Biology* **187**, 91–100.

Weber, M., Nguyen, M. B., Li, M. Y., Flora, P., Shuda, M. and Ezhkova, E. (2023). Merkel
Cell Polyomavirus T Antigen-Mediated Reprogramming in Adult Merkel Cell Progenitors.
J Invest Dermatol S0022-202X(23)02120–6.

Whitear, M. (1989). Merkel cells in lower vertebrates. *Archivum Histologicum Japonicum* **52
Suppl**, 415–22.

Whitear, M. and Lane, E. B. (1981). Fine structure of Merkel cells in lampreys. *Cell and Tissue
Research* **220**, 139–51.

Woo, S.-H., Lumpkin, E. A. and Patapoutian, A. (2015). Merkel cells and neurons keep in
touch. *Trends Cell Biol* **25**, 74–81.

**Wright, M. C., Logan, G. J., Bolock, A. M., Kubicki, A. C., Hemphill, J. A., Sanders, T. A.
and Maricich, S. M.** (2017). Merkel cells are long-lived cells whose production is
stimulated by skin injury. *Dev Biol* **422**, 4–13.

Xiao, Y., Williams, J. S. and Brownell, I. (2014). Merkel cells and touch domes: more than
mechanosensory functions? *Exp Dermatol* **23**, 692–695.

Xu, C., Volkery, S. and Siekmann, A. F. (2015). Intubation-based anesthesia for long-term
time-lapse imaging of adult zebrafish. *Nature Protocols* **10**, 2064–73.

Yamashita, Y., Toida, K. and Ogawa, H. (1993). Observation of Merkel cells with scanning
electron microscopy. *Neurosci Lett* **159**, 155–158.