<u>Title:</u> Range expansion is both slower and less predictable across a spatial gradient in

temperature

<u>Authors:</u> Takuji Usui[1], Amy L. Angert[1,2]

<u>Author affiliations:</u>

[1]Department of Botany, University of British Columbia, Vancouver, British Columbia, V6T 1Z4,

Canada (usuitakuji@gmail.com)

[2]Department of Zoology, University of British Columbia, Vancouver, British Columbia, V6T

1Z4, Canada (angert@mail.ubc.ca)

<u>Corresponding author:</u> Takuji Usui (Department of Botany, University of British Columbia,

2122 Main Mall, Vancouver, BC, V6T 1Z4, Canada; email: usuitakuji@gmail.com)

<u>Running title:</u> Range expansion across abiotic gradients

<u>Keywords:</u> biological invasion; demography; dispersal; eco-evolutionary dynamics;

environmental gradient; experimental evolution; priority effects; range limit; variability

<u>Article type:</u> Letter

Word counts: Abstract (149), Main text (4989)

Number of references: 52

Number of figures: 4

<u>Statement of authorship:</u> TU conceived of this study with guidance from ALA. TU collected the

data. TU analysed the data, developed the figures, and wrote the first draft of the manuscript with

feedback from ALA. TU and ALA contributed to edits in preparation for manuscript submission.

ABSTRACT

Rapid evolution in colonizing populations can alter our ability to predict future range expansions. Recent studies suggest that how evolution influences the predictability of range expansion speed depends on the balance of genetic drift and selection at the range front, with factors increasing selection proposed to produce greater consistency, and hence increased predictability, in range expansion speed. Here, we test whether selection from environmental gradients across space produces more predictable range expansion speeds, using the experimental evolution of replicate, duckweed populations colonizing landscapes with and without a temperature gradient. While range expansion across a temperature gradient was slower, with range-front populations displaying higher population densities and genetic signatures consistent with greater selection, we found that range expansion speed was more variable and less predictable across replicate populations. Our results therefore challenge current theory, highlighting that selection and spatial priority effects could together govern the predictability of range expansion speeds.

INTRODUCTION

Predicting the movement of species ranges is a fundamental goal of ecology and is of increasing importance as biological invasions and climate change accelerate range expansion (Chen et al. 2011; Iseli et al. 2023; Nadeau & Urban 2019). Classic work on the ecological dynamics of range expansion has demonstrated that higher values of two key life-history traits – dispersal and reproductive output – increase the speed of range expansion (Kot et al. 1996; Skellam 1951). More recent work incorporating rapid evolution has shown that not only can dispersal and reproduction evolve to increase the average speed of range expansion, but that additionally,

evolution can directly influence the amount of variability in the speed of population spread

across replicate, range-expanding populations (Miller et al. 2020; Ochocki & Miller 2017;

Williams et al. 2016; Wiliams et al. 2019). However, when and in what direction evolution

modifies the variability, and hence predictability, of range expansion remains unclear.

On one hand, eco-evolutionary studies of range expansion predict that evolution should increase

variability in range expansion speed across replicate populations. Specifically, serial founding

events during range expansion can result in the fixation of maladaptive alleles at the range front

and consequent reduction in fitness ('expansion load') through a spatial analogue of genetic drift

('gene surfing') (Excoffier et al. 2009; Klopfstein et al. 2006; Perrier et al. 2020). Gene surfing

in small, range-expanding populations leads to the fixation of random amounts of expansion load

across replicate populations, thus generating variability in range expansion speed (Miller et al.

2020; Phillips 2015; Williams et al. 2019). The implication then, is that any single realisation of

range expansion in nature is drawn from a distribution of possible outcomes, making predictions

of range expansion more difficult.

In contrast, some recent experimental work testing the predictability of range expansions shows

that evolution can also reduce variance in range expansion speed (Miller et al. 2020; Williams et

al. 2019). This is predicted to occur when directional selection (e.g., for traits conferring greater

dispersal or environmental tolerance; Chuang & Peterson 2016; Freedman et al. 2020; Lancaster

et al. 2015) reduces trait variance at the range front, hence increasing consistency in the average

speed of range expansion (Phillips 2015; Williams et al. 2019). Overall, how evolution modifies

the variability in range expansion speed may therefore depend on the relative strength of gene

surfing and selection in range-front populations, and the factors that tip this balance one way or

another (Miller et al. 2020; Urquhart-Cronish et al. 2022; Williams et al. 2019).

One key factor that could tip this balance between the variance-generating and variance-reducing

effects of evolution is environmental variation in space. Specifically, species' ranges in nature

exist across broad-scale spatial abiotic gradients. Importantly, changing abiotic conditions across

space could lead to slower range expansion due to the need for populations to adapt to the

underlying abiotic gradient. This initial maladaptation at the range front can allow more time for

individuals to arrive from the range core, consequently increasing population size and genetic

variation at the range front over time (Fronhofer et al. 2017; Gilbert et al. 2017). In these

'steeper' range boundaries, directional selection favouring traits that confer tolerance to changing

abiotic conditions could weaken gene surfing and its genetic and demographic consequences

(Gilbert et al. 2017; Williams et al. 2019). Thus, selection from underlying abiotic gradients in

space could make the speed of range expansion among replicate populations less variable and

more predictable. Although there is a need to incorporate realistic sources of environmental

variation, experimentally testing how selection from abiotic gradients in space modifies the

predictability of replicate range expansions in nature remains difficult (Miller et al. 2020).

Experimental range expansions in "miniaturised" landscapes (Larsen & Hargreaves 2020;

Lustenhower et al. 2023) can provide a powerful bridge between theory and range expansion in

natural landscapes. First, these landscapes differ from traditional microcosms in that replicate

populations can disperse naturally across linear landscapes, such that realistic genetic and

demographic spatial structure arises during range expansions. Second, through replicated

landscapes we can track a distribution of ecological and evolutionary outcomes in colonising

populations. This allows experimental insights into the factors that contribute to variability in

range expansion that would not be possible without replication. Third, through experimental

evolution we can track the eco-evolutionary dynamics of range expansion unfolding over

multiple generations in real time, while field studies have mostly been retrospective, inferring

past demographic and evolutionary processes based on current spatial patterns. Despite the

increasing utility of these landscapes (Larsen & Hargreaves 2020), we still require experiments

that test the dynamics of range expansion across landscapes that incorporate selection from

spatial environmental variation.

Here, we use experimental populations of duckweeds in the *Lemna* species complex to

experimentally test how selection from spatial abiotic gradients modifies the predictability (i.e.,

among-replicate variability) of range expansion speed. Using replicate landscapes with and

without a temperature gradient in space, we predicted that: (i) range expansion speed should be

more predictable (i.e., less variable among replicates) when populations move across landscapes

with a spatial temperature gradient than without. We predicted that this would occur as: (ii) range

expansion across temperature gradients would be on average slower, resulting in higher

population densities and steeper range boundaries. Consequently, we predicted that there should

be stronger signatures of selection over genetic drift in populations expanding across temperature

gradients. If selection was stronger in populations expanding across temperature gradients in

space, then we predict: (iii) less among-replicate variability in genotypic composition for

gradient than uniform landscapes; and (iv) directional selection for traits relevant to thermal

tolerance in these populations.

MATERIALS & METHODS

Lemna species collection and culturing

Duckweeds are small, freshwater angiosperms with a rapid generation time of 2–5 days under ideal conditions (Landolt 1986). Although sexual reproduction occurs sporadically in nature, reproduction during experiments occurs exclusively through the clonal budding of daughter fronds (Docauer 1983; Landolt 1986). Multiple fronds can stay connected via a stipule to form rafts of ~3-5 fronds until the stipule breaks and daughter fronds disperse. We sampled 20 unique accessions of duckweeds in the *Lemna* species complex from 20 sites around the Pacific Northwest region of North America (Table S1). Upon sampling, we immersed fronds in a dilute bleach solution (10% v:v) to create axenic plants of each accession. We maintained each of our 20 accessions in 250 mL erlenmeyer flasks with 100 mL of artificial pond media (Appenroth et al. 1996), inside a temperature-controlled laboratory (20°C) and under full-spectrum LED lighting (SunBlaster 6400K; 16:8 hour light:dark cycle). To obtain sufficient numbers of individuals for the experiment, we grew out each accession for 4 weeks inside 20 L buckets at the UBC Botany greenhouse, with 17 L of water, 4 L of autoclaved potting soil (MiracleGro Potting Mix), 1.32 mM of KNO_3, 0.16 mM of $Ca(NO_3)_2$, and 0.02 mM of KH_2PO_4 as fertilisers.

Duckweeds in the *Lemna* species complex are thought to be comprised of *L. minor*, *L. turionifera*, and *L. japonica*. These three lineages are ecologically and genetically cryptic, and species delineation is currently a challenge despite the development of various molecular markers (Braglia et al. 2021a; Braglia et al. 2021b). Our barcoding approach based on Tubulin-Based Polymorphism (TBP) fingerprinting (Braglia et al. 2021a) and microsatellite genotyping

of each accession detected that our accessions were composed of 6 unique genotypes of *L. minor*

and 5 genotypes of *L. japonica* (Appendix S1; Tables S1-S3). While some studies have proposed

that *L. japonica* is a hybrid between *L. minor* and *L. turionifera* (Braglia et al. 2021a; Braglia et

al. 2021b; Volkova et al. 2023), we note that taxonomic delineation between these groups is at

present unstable. In our experiment, we therefore describe evolution as occurring through the

sorting of unique genotypes within the *Lemna* species complex.

Experimental landscapes and temperature treatment

We set up 50 experimental landscapes across five benches inside the greenhouse, with each

landscape constructed from aluminium gutters (150 cm x 12.5 cm; L x W) and filled with 13.5 L

of water and 2.5 L of potting soil. To ensure nutrients and water-levels did not deplete over time,

we added 0.13 mM of KNO_3, 0.02 mM of $Ca(NO_3)_2$, and 0.002 mM of KH_2PO_4 every 4 days to

each landscape, and continuously regulated water-levels using a float-valve connected to a sump

pump. We connected drip emitters to the sump pump so that water refilling occurred slowly and

did not disrupt the spatial structure of populations. We set up a temperature treatment (Fig. 1a)

such that range expansion occurred across landscapes with a spatial gradient in temperature

("gradient landscapes" hereafter; $N = 25$) or with spatially uniform temperatures ("uniform

landscapes" hereafter; $N = 25$). The position of gradient and uniform landscapes was randomised

within each bench. To establish a temperature gradient, we used a submersible heater (150W JBJ

aquarium heaters) set to 37°C and placed at one end of each gradient landscape. To maintain

uniform temperatures, we placed two heaters set to 24°C at both ends of each uniform landscape.

These set temperatures were chosen as they are close to the critical thermal maxima (T_{max}) and

thermal optima (T_{opt}), respectively, of the *Lemna* genotypes used (Fig. S1). We monitored water

temperatures using three underwater temperature loggers (HOBO pendants) placed at equal

spacing within each landscape. For gradient and uniform landscapes, the imposed temperature

treatment resulted in increasing or uniform temperatures across space, respectively (Fig. 1b).

Although our initial aim for gradient landscapes was to set up a stress gradient such that

populations experienced increasing heat stress as they dispersed, we note that: (i) average

temperatures at the experimental range limit of gradient landscapes were still relatively benign

(31.6°C; SD = 1.9°C; Fig. 1b); and (ii) as average temperatures across space were similar

between gradient and uniform landscapes (27.4°C; SD = 3.85°C and 26.9°C; SD = 1.53°C,

respectively), temperatures at the start of gradient landscapes were relatively cooler than in

uniform landscapes (24.3°C; SD = 2.6°C and 27.6°C; SD = 1.4°C, respectively). However, we

note that our temperature treatment still reflects the presence and absence of a temperature

gradient across space in gradient and uniform landscapes, respectively.

Range expansion

To initiate range expansion, we established genetically diverse *Lemna* populations, with each

population consisting of an even density of 20 accessions ($N = 20$ individuals per accession for a

founding population size of $N = 400$ individuals). We seeded each population at one end of each

landscape, allowing 2 days for plants to acclimate before range expansion. In gradient

landscapes, we seeded populations at the cooler end of the landscape and allowed range

expansion into increasingly warmer temperatures (Fig. 1a). To allow all populations the

opportunity to colonize the full spatial extent of the landscape, the experiment ran for 112 days.

During our experiment, we lost one replicate due to structural damage and lost 10 replicates on

one bench due to a disturbance event that disrupted population spread. We also lost 6 replicates

due to heavy algal growth on the water surface that obstructed population spread. In total, we

retained 33 experimental landscapes across four benches ($N = 16$ gradient landscapes; $N = 17$

uniform landscapes).

Quantifying range expansion speed

We quantified the mean and variability in range expansion speed across landscapes for the initial

24 days (i.e., ~5–12 generations) of range expansion. This timescale represents population spread

during the exponential growth phase and thus is most consistent with theoretical models of range

expansion (Kot et al. 1996; Skellam 1951). Also, some populations had reached the range limit

of landscapes by day 24. These replicates would not be able to spread further and their inclusion

after 24 days would artificially reduce the average and among-replicate variability in range

expansion speed (i.e., the ceiling effect).

To quantify range expansion speed, we took aerial photographs of each landscape every 4 days

(Nikkon D750; AF-S Nikkor 24-120 mm f/4G ED VR). From these images, we first quantified

the maximum distance travelled (cm) for each population by measuring the distance reached by

the furthest individual every 4 days. We then quantified the mean and among-replicate variability

in the distance travelled at each treatment level and time point. To estimate the difference in

among-replicate variability of distance travelled for gradient and uniform landscapes, we

calculated the log coefficient of variation ratio (lnCVR; Nakagawa et al. 2015) as:

212
$$lnCVR = \frac{lnCV_E}{lnCV_C} + \frac{1}{2(n_E - 1)} - \frac{1}{2(n_C - 1)} \qquad (1)$$

where subscripts E and C refer to the experimental (i.e., gradient) and control (i.e., uniform)

landscapes, CV is the coefficient of variation, and n is the sample size. We calculated lnCVR

such that positive values correspond to larger variability in distance travelled among gradient

compared to uniform landscapes. We use lnCVR because CV accounts for the linear and positive

mean-variance relationship on the log scale (Fig. S2; Taylor 1961). To report differences in CV,

we back-transformed estimates of lnCVR.

Quantifying range-front population density and steepness

We quantified spatiotemporal changes in population density at the range front by estimating the

percent cover of live (i.e., green) fronds on the water surface. We did this by colour thresholding

images obtained every 4 days using ImageJ (v1.53) and estimating percent cover in 2 cm x 12.5

cm grids at the range front (defined as the first 12 cm from the furthest travelled individual). We

then fit a logistic model of percent cover across space for each landscape to quantify the

maximum population density at the range front at each time point. We also quantified the

steepness of the range boundary (i.e., the rate of increase in population density) by fitting an

exponential model to the first three estimates of percent cover at the range front of each

landscape. Steepness was quantified as the exponent obtained from each model (Legault et al.

2020).

Quantifying range-front evolution

We quantified evolution as the change in genotype frequencies due to genotype sorting from the

founding population. Post-experiment, we haphazardly sampled 12 individual rafts of duckweeds

from each replicate range front. Each sample was then genotyped using four microsatellite

markers in a single multiplex reaction (Appendix S1; Table S2; Hart et al. 2017). We

successfully obtained genotype sequences for 272 individuals (88% of all genotyped samples).

To balance the number of samples obtained within each replicate and treatment, we rarefied

samples and dropped replicates represented by less than $N = 6$ individuals due to DNA extraction

or genotyping failure. For all downstream analyses, we therefore obtained a total of $N = 6$

individuals per replicate, for 10 and 17 replicates of gradient and uniform landscapes,

respectively. We note a higher extraction and genotyping failure for samples obtained from

gradient landscapes possibly due to greater damage to plant tissue from heat stress.

To investigate whether genotype sorting in the experiment led to consistent trait evolution, we

also measured a suite of genotype-specific traits *ex situ* in incubators and across a range of

temperatures (Appendix S2; Fig. S1). We parameterized traits hypothesised to be ecologically

relevant to range expansion across thermally variable landscapes, including: maximum growth

rate (r_{max}); thermal optimum (T_{opt}); critical thermal maximum (CT_{max}); critical thermal minimum

(CT_{min}); thermal breadth (i.e., the range of temperatures for which growth rate is within 80% of

the maximum; Padfield et al. 2021); thermal tolerance (i.e., the difference between CT_{max} and

CT_{min}); mean specific-leaf area (SLA; cm^2/g); mean root-shoot ratio (cm/mg); and plasticity in

both SLA and root-shoot ratio, as quantified by its CV across temperature (see Table S10 for

trait estimates at the genotype level). Before analyses, we scaled and centred each trait, and

estimated mean trait values at the population level for both our founding and evolved populations

by weighting trait values by the observed genotype frequencies of each population.

Statistical analyses

We tested whether temperature treatment led to differences in the ecological dynamics of range

expansion. First, we tested for treatment differences in mean range expansion speed by fitting a

linear mixed-effects model using the *lme4* package (Bates et al. 2015). We fit maximum distance

travelled (cm) as the response variable and an interaction between time (day) and temperature

treatment (gradient *vs* uniform) as fixed predictors, where mean range expansion speed is

modelled as slopes (cm/day). We also included bench ID as a fixed covariate and fitted a random

slope (i.e., speed) for each replicate. Second, we tested for treatment differences in among-

replicate variability of distance travelled by fitting a linear model with lnCVR as the response

variable and time as the predictor. We modelled the temporal trend in lnCVR as visual

inspections of CV in distance travelled showed changes over time for both gradient and uniform

landscapes (Fig. 2b). For sensitivity analyses, we also conducted a linear meta-regression of

lnCVR to explicitly model the sampling variation around each observation (Appendix S3). Third,

we tested for treatment differences in range-front population density (i.e., carrying capacity) and

steepness of the range boundary (i.e., exponent) by fitting separate linear regressions with either

population density or steepness as the response variable, respectively, and time, temperature

treatment, and their interaction as predictors. For sensitivity analyses, we also conducted a

separate meta-regression of population density and steepness by including the standard error

around each observation using the package *metafor* (Viechtbauer 2010).

We also tested whether temperature treatment led to differences in the evolutionary dynamics of

range expansion. First, we tested for treatment differences in the amount of genotypic or trait

change, as estimated by the Euclidean distance between the initial genotype or trait frequency in

the founding population and the post-experimental genotype or trait frequency for each

landscape. We fit separate linear regressions with Euclidean distance of genotype or trait

frequencies as the response variable, temperature treatment as the predictor, and bench ID as a

fixed covariate. Second, we tested for treatment differences in the final genotype or trait

composition in range-front populations. Here, we quantified compositional dissimilarity of

genotypes or traits across all pairwise populations within each treatment using Euclidean

dissimilarity matrices. We then conducted separate PERMANOVAs using *adonis2* in the *vegan*

package (Okansen et al. 2022) with genotype or trait dissimilarity among replicates as the

response variable, temperature treatment as the predictor, and bench ID as a fixed covariate.

Upon observing differences in genotype composition, we tested whether certain genotypes arose

to higher frequencies than predicted by chance by conducting a post-hoc binomial test on the

mean frequency of each genotype for each temperature treatment. Third, we tested whether

temperature treatment led to differences in the amount of among-replicate variability found in

the genotype composition of range-front populations. To quantify the amount of observed

among-replicate variability, we first averaged genotype frequencies across replicates within each

treatment. For each temperature treatment, we then calculated the Euclidean distance between

this mean genotype composition and the observed genotype composition of each replicate. As a

measure of the observed among-replicate variability in genotype composition, we then summed

these Euclidean distances across all replicates within each treatment. We tested for treatment

differences in among-replicate variability through a permutation test, with Euclidian distances

permuted randomly across treatment 100,000 times. Finally, we tested whether the observed

among-replicate variability in each treatment differed from that predicted by random sampling.

To quantify among-replicate variability predicted by random sampling, we first pooled

genotypes across replicates within each temperature treatment. From this pool, we randomly

drew 6 individuals with replacement (i.e., representing the 6 genotypes in our observed data) and

repeated this N times for each treatment, where N is the number of replicates for gradient ($N =$

10) and uniform landscapes ($N = 17$). We then estimated the mean genotype composition for

each temperature treatment and calculated the Euclidean distance between the simulated mean

genotype composition and the simulated genotype composition of each replicate. We summed

these Euclidean distances for each treatment as a measure of the among-replicate variability in

genotype composition predicted by random sampling. We repeated this simulation a total of

100,000 times to estimate the probability of obtaining the observed among-replicate variability in

genotype composition for each treatment based on random sampling. All model-fitting was

conducted in R (R Core Team 2020).

RESULTS

Temperature treatment on range expansion speed

On average, we found that range expansion speed across gradient landscapes was slower than

across uniform landscapes (time by treatment effect: $P = 0.016$; Table S4; Fig. 2a), consistent

with prediction (ii). Over 24 days of range expansion, the presence of a spatial temperature

gradient reduced the average speed of range expansion by 50.9% (95% CI: 36.0% to 53.8%)

compared to uniform landscapes (Table S4). We also found that the among-replicate variability

in distance travelled differed between gradient and uniform landscapes (Fig. 2b and 2c). At the

initial stages of range expansion, we found that there was less among-replicate variability (CV)

in distance travelled across gradient compared to uniform landscapes (e.g., a 40.3% [95% CI:

28.6% to 56.8%] reduction in CV across gradient compared to uniform landscapes on day 4;

Table S5), consistent with prediction (i). However, we found a significant increase in lnCVR

over time (lnCVR slope: $P = 0.002$) such that by day 24, CV in distance travelled was 78.2%

(95% CI: 28.6% to 151.0%) greater across gradient than uniform landscapes (Fig. 2c), contrary

to prediction. This result was consistent when including sampling variation around lnCVR in the

linear model (Table S5).

Temperature treatment on range-front density and steepness

We found that the population density reached at the range front differed over time and between

gradient and uniform landscapes (Fig. 2d-f). Although range-front densities were initially similar

between gradient and uniform landscapes, population density increased more rapidly over time

for gradient landscapes (treatment by time effect: $P = 0.002$; Fig. 2f), consistent with prediction

(ii). By day 24, population density at the range front was 88.1% (95% CI: 76.5% to 104.0%)

higher in gradient compared to uniform landscapes (Table S6). Along with increased population

density, we also found range boundaries to be steeper in gradient landscapes (Table S7; Figure

S3), consistent with prediction (ii). These results were consistent in sensitivity models that

included sampling error in estimates of range-front population density and steepness (Tables S6

and S7).

Temperature treatment on evolutionary change

We found that evolution in range-front populations (i.e., change in genotype frequencies from

the founder population) occurred in both gradient and uniform landscapes (Intercept: $P < 0.001$;

Table S8). We found partial support for prediction (iii) that selection is greater in gradient

landscapes. On average, the observed among-replicate variability in genotype composition was

greater in uniform compared to gradient landscapes (Fig. 4), although permutation tests revealed

that this treatment difference was non-significant ($P = 0.304$). Importantly, we found that the

observed among-replicate variability in genotype composition was greater than predicted by

random sampling for both gradient ($P = 0.003$) and uniform ($P < 0.001$) landscapes (Fig. 4). We

also found that genotype composition of range-front populations differed between gradient and

uniform landscapes after range expansion (PERMANOVA: $P = 0.002$; Table S9; Fig. 3a).

Specifically, range-front populations in gradient landscapes were often dominated by one

genotype (G11; Fig. 3c), with post-hoc binomial tests showing that this genotype consistently

increased in frequency in gradient landscapes ($P < 0.001$) but not in uniform landscapes ($P =$

0.869), suggesting selection for this genotype in the former (Fig. 3e).

Trait evolution (i.e., change in trait frequencies from the founder population) occurred in both

gradient and uniform landscapes (Intercept: $P = 0.004$; Table S8; Fig. S4). However, contrary to

prediction (iv), out of the 11 traits assessed we did not find significant treatment differences in

the extent of trait change (Treatment effect: $P = 0.345$; Table S8) or in trait composition of

range-front populations (PERMANOVA: $P = 0.088$; Table S9; Fig. S4).

DISCUSSION

Recent studies have highlighted that rapid evolution can complicate our ability to predict the eco-

evolutionary dynamics of range expansions (Miller et al. 2020; Shaw et al. 2023; Williams et al.

2019; Zilio et al. 2023). Here, we used replicate range-expanding populations to empirically test

whether the presence of selection from an abiotic gradient across landscapes could modify the

among-replicate variability, and hence predictability, of range expansion speed. Notably, we

found that over time range expansion speed became more variable, and hence less predictable, in

landscapes with a spatial temperature gradient than without (Fig. 2c). This occurred despite

gradient landscapes having slower range expansions on average, greater range-front population

densities (Fig. 2), and genetic signatures consistent with increased selection (Fig. 3)– conditions

that would otherwise predict less gene surfing and thus more consistency in the speed of range

expansions across replicate populations (Miller et al. 2020; Phillips 2015; Williams et al. 2019).

Our results therefore challenge current theory and highlight the complexities of predicting range

expansions even in highly standardised, replicate mesocosms with simple environmental

variation.

We found that the greater variability in range expansion speed observed for gradient compared to

uniform landscapes over time is driven by a combination of rapidly decreasing among-replicate

variability (CV) in distance travelled for uniform landscapes, and slightly increasing CV for

gradient landscapes over time (Fig. 2b). In uniform landscapes, the decrease in CV of distance

travelled over time occurred as mean range expansion speed increased more quickly than the

variance observed across replicates (Fig. 2a). A rapid increase in mean range expansion speed

across uniform landscapes is consistent with previous experimental evolution work on range

expansions, which found that range expansion speed can increase under constant environments

due to spatial sorting for dispersal phenotypes and low-density selection for increased

reproductive output at the range front (Ochocki & Miller 2017; Wagner et al. 2017; Weiss-

Lehman et al. 2017). In contrast, for gradient landscapes, the slight increase in CV of distance

travelled over time occurred as the variance in speed among replicates increased more quickly

than the mean range expansion speed (Fig. 2a). This occurred as range expansion remained slow

for most gradient landscapes but rapidly sped up in a few replicates (Fig. 2a), consequently

elevating the among-replicate variability observed over time.

It is possible that the faster range expansion seen in some of these gradient landscapes was partly

driven by selection from the temperature gradient (Moerman et al. 2022; Phillips et al. 2006;

Szücs et al 2017; Williams et al. 2016). We found that gradient landscapes had consistently

higher frequencies of one genotype (G11) at the range front (Fig. 3c and 3d). In contrast, this

genotype did not achieve frequencies >0.5 in any range-front population for uniform landscapes

(Fig. S5). We note that the evolutionary dynamics described above reflect longer term range

expansion and selection after 112 days (~22-56 generations), while the population dynamics

reflect shorter term range expansion in the first 24 days (~4-12 generations). Even so,

comparison to genetic data obtained at an earlier point in time from the fastest range-expanding

replicates confirm that this same genotype had reached high frequencies at the range-front of

gradient landscapes but not in uniform landscapes (Fig. S6), suggesting that rapid evolution in

the former occurred as early as within the first 36 days (i.e., 7–18 generations) of range

expansion.

While we did not find trait composition to differ significantly between gradient and uniform

landscapes within the 11-dimensional trait space (Fig. S4), post-hoc comparison of univariate

trait means revealed that this high-frequency genotype in gradient landscapes has both the

highest mean estimate of thermal breadth and the lowest maximum growth rates among all

genotypes (Table S10). The importance of maintaining performance across a broader range of

temperatures through thermal breadth evolution has been long emphasised in studies of range

expansion across latitudinal gradients in climate (Ackerly 2003; Lancaster 2016; Lancaster 2022)

and is also consistent with studies that report a positive relationship between range size and niche

breadth (Slatyer et al. 2013). While it could be surprising that range-expanding populations are

also fixed for the genotype with the lowest maximum growth rate, this could be explained by

generalist-specialist trade-offs between sustaining growth rates across a wider range of

temperatures and obtaining high growth rates at the temperature optimum (Huey & Hertz 1984).

Additionally, models of range expansion that incorporate density-dependence and life-history

trade-offs also predict that higher population densities and steeper range boundaries, as observed

in our gradient landscapes, can favour K-selected traits (e.g., increased competitive ability) at the

range front even if this comes at a cost of r-selected traits (e.g., increased reproductive output

and growth rates; Burton et al. 2010; Pachepsky & Levine 2011; Williams et al. 2016).

However, while the consistently high frequency of this genotype suggest that selection was

stronger at the range front of gradient over uniform landscapes, we also found in both landscapes

that the among-replicate variability in genetic composition was greater than predicted by random

sampling (Fig. 4). Such a pattern could result if the eco-evolutionary dynamics of range

expansion in landscapes with environmental variation is governed by a combination of selection

and spatial priority effects, such that the genotypes available to respond to selection in range-

front populations is shaped by individuals that initially occupy the range front through chance

dispersal (Baym et al. 2016; Hallatschek et al. 2007; Hewitt 2000; Williams et al. 2016). Taken

together, we propose that spatial priority effects and its variance-generating effects at the range

front, combined with directional selection from abiotic gradients in space, could lead on average

to slower but more variable speeds during range expansion, relative to range expansion across

uniform environments.

Our results highlight that even in ecological conditions under which the effects of gene surfing

are reduced, the speed of range expansion can be variable and difficult to predict. Previous

experimental and theoretical work suggest the predictability of range expansion speed depends

on the balance of evolutionary processes that generate and reduce variability (i.e., gene surfing

and selection, respectively) and the ecological factors that tip this balance (Miller et al. 2020;

Williams et al. 2019). We additionally suggest that predicting the dynamics of range expansion

may require an understanding of how selection interacts with spatial priority effects, and whether

priority effects could hamper the response to selection and increase among-replicate variability

in the speed of range expansion. As populations traverse across environmentally variable and

complex landscapes, studies of natural range expansions will need to incorporate how selection

from underlying environmental variation in space can influence our ability to predict future

range expansions.

ACKNOWLEDGEMENTS

We thank H. Ahmed, M. Catalani, A. Doherty, K. Kubeck, E. Menchions, T. Scriber, and P.

Seppala for their help with plant and greenhouse maintenance, and for providing assistance with

experimental setup. We thank M. Chen, E. Lam, D. Moxley and A. Wong for their generous

assistance with genetic work. We are grateful to R. Germain, S. Otto, M. Tseng, and J. Williams

for their comments on earlier versions of the manuscript. This work was supported by the UBC

International Doctoral Fellowship and the SSE Greg Lewontin Award to TU, and a Discovery

Grant from the Natural Sciences and Engineering Research Council of Canada to ALA.

REFERENCES

Ackerly, D.D. (2003). Community assembly, niche conservatism, and adaptive evolution in

changing environments. *Int. J. Plant Sci.*, 164, S165–S184.

Appenroth, K.J., Teller, S. & Horn, M. (1996). Photophysiology of turion formation and

germination in *Spirodela polyrhiza*. *Biol. Plantarum*, 38, 343–351.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *J. Stat. Softw.*, 67, 1–48.

Baym, M., Lieberman T.D., Kelsic, E.D., Remy, C., Gross, R., Yelin, I. & Kishony, R. (2016).

Spatiotemporal microbial evolution on antibiotic landscapes. *Science*, 353, 1147–1151.

Braglia, L., Breviario, D., Gianì, S., Gavazzi, F., De Gregori, J. & Morello, L. (2021a). New

insights into interspecific hybridization in *Lemna* L. Sect. *Lemna* (Lemnaceae Martinov).

Plants, 10, 2767.

Braglia, L., Lauria, M., Appenroth, K.J., Bog, M., Breviario, D., Grasso, A., *et al*. (2021b).

Duckweed species genotyping and interspecific hybrid discovery by Tubulin-Based

Polymorphism Fingerprinting. *Front. Plant. Sci.*, 12, 625670.

Burton, O.J., Phillips, B.L. & Travis, J.M.J. (2010). Trade-offs and the evolution of life-histories

during range expansion: Evolution during range expansion. *Ecol. Lett.*, 13, 1210–1220.

Chen, I.C., Hill, J.K., Ohlemüller, R., Roy, D.B. & Thomas, C.D. (2011). Rapid range shifts of

species associated with high levels of climate warming. *Science*, 333, 1024–1026.

Chuang, A. & Peterson, C.R. (2016). Expanding population edges: Theories, traits, and trade-

offs. *Glob. Change Biol.*, 22, 494–512.

Docauer, D. M. (1983). *A nutrient basis for the distribution of the Lemnaceae*. Dissertation.

University of Michigan, Ann Arbor, Michigan, USA.

Excoffier, L., Foll, M. & Petit, R.J. (2009). Genetic consequences of range expansions. *Annu.*

Rev. Ecol. Evol. S., 40, 481–501.

Freedman, M.G., Dingle, H., Strauss, S.Y., & Ramírez, S.R. (2020). Two centuries of monarch

butterfly collections reveal contrasting effects of range expansion and migration loss on

wing traits. *Proc. Natl. Acad. Sci. U.S.A.*, 117, 28887–28893.

Fronhofer, E.A., Nitsche, N. & Altermatt, F. (2017). Information use shapes the dynamics of

range expansions into environmental gradients. *Glob. Ecol. Biogeogr.,* 26, 400–411.

Gilbert, K.J., Sharp, N.P., Angert, A.L., Conte, G.L., Draghi, J.A., Guillaume, F., *et al.* (2017).

Local adaptation interacts with expansion load during range expansion: Maladaptation

reduces expansion load. *Am. Nat.*, 189, 368–380.

Hallatschek, O., Hersen, P., Ramanathan, S. & Nelson, D.R. (2007). Genetic drift at expanding

frontiers promotes gene segregation. *Proc. Natl., Acad. Sci. U.S.A.*, 104, 19926–19930.

Hart, S.P., Turcotte, M.M. & Levine, J.M. (2019). Effects of rapid evolution on species

coexistence. *Proc. Natl., Acad. Sci. U.S.A.*, 116, 2112–2117.

Hewitt, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, 405, 907–913.

Huey, R.B. & Hertz, P.E. (1984). Is a Jack-of-All-Temperatures a Master of None? *Evol.,* 38,

441–444.

Iseli, E., Chisholm, C., Lenoir, J., Haider, S., Seipel, T., Barros, A., *et al.* (2023). Rapid upwards

spread of non-native plants in mountains across continents. *Nat. Ecol. Evol.*, 7, 405–413.

Klopfstein, S., Currat, M. & Excoffier, L. (2006). The fate of mutations surfing on the wave of a

range expansion. *Mol. Biol. Evol.*, 23, 482–490.

Kot, M., Lewis, M.A. & Van Den Driessche, P. (1996). Dispersal data and the spread of

invading organisms. *Ecol.*, 77, 2027–2042.

Lancaster, L.T., Dudaniec, R.Y., Hansson, B. & Svensson, E.I. (2015). Latitudinal shift in

thermal niche breadth results from thermal release during a climate-mediated range

expansion. *J. Biogeogr.*, 42, 1953–1963.

Lancaster, L.T. (2016). Widespread range expansions shape latitudinal variation in insect

thermal limits. *Nat. Clim. Change*, 6, 618–621.

Lancaster, L.T. (2022). On the macroecological significance of eco-evolutionary dynamics: The

range shift–niche breadth hypothesis. *Philos. T. R. Soc. B.*, 377, 20210013.

Landolt, E. (1986). *Biosystematic investigations in the family of duckweeds, Lemnaceae: the*

family of Lemnaceae, a monographic study. Volume 2: Morphology, karyology, ecology,

geographic distribution, systematic position, nomenclature descriptions.

Veroeffentlichungen des Geobotanischen Institutes der ETH, Stiftung Rubel, Zurich,

Switzerland.

Larsen, C.D. & Hargreaves, A.L. (2020). Miniaturizing landscapes to understand species

distributions. *Ecography*, 43, 1625–1638.

Legault, G., Bitters, M.E., Hastings, A. & Melbourne, B.A. (2020). Interspecific competition

slows range expansion and shapes range boundaries. *Proc. Natl. Acad. Sci. U.S.A.*, 117,

26854–26860.

529 Lustenhouwer, N., Moerman, F., Altermatt, F., Bassar, R.D., Bocedi, G., Bonte, D., *et al.* (2023).

530   Experimental evolution of dispersal: Unifying theory, experiments and natural systems. *J.*

531   *Anim. Ecol.*, 92, 1113–1123.

532 Miller, T.E.X., Angert, A.L., Brown, C.D., Lee-Yaw, J.A., Lewis, M., Lutscher, F., *et al.* (2020).

533   Eco-evolutionary dynamics of range expansion. *Ecol.*, 101. e03139.

534 Moerman, F., Fronhofer, E.A., Altermatt, F. & Wagner, A. (2022). Selection on growth rate and

535   local adaptation drive genomic adaptation during experimental range expansions in the

536   protist *Tetrahymena thermophila*. *J. Anim. Ecol.*, 91, 1088–1103.

537 Nadeau, C.P. & Urban, M.C. (2019). Eco-evolution on the edge during climate change.

538   *Ecography*, ecog.04404.

539 Nakagawa, S., Poulin, R., Mengersen, K., Reinhold, K., Engqvist, L., Lagisz, M., *et al.* (2015).

540   Meta-analysis of variation: Ecological and evolutionary applications and beyond. *Methods*

541   *Ecol. Evol.*, 6, 143–152.

542 Ochocki, B.M. & Miller, T.E.X. (2017). Rapid evolution of dispersal ability makes biological

543   invasions faster and more variable. *Nat. Commun.*, 8, 14315.

544 Oksanen, J., Simpson, G.L., Blanchet, F.G., Kindt, R., Legendre, P. & Minchin, P.R. (2022).

545   Vegan: Community ecology package (v 2.6–2).

546 Pachepsky, E. & Levine, J.M. (2011). Density dependence slows invader spread in fragmented

547   landscapes. *Am. Nat.*, 177, 18–28.

548 Padfield, D., O'Sullivan, H. & Pawar, S. (2021). rTPC and nls.multstart: A new pipeline to fit

549   thermal performance curves in R. *Methods Ecol. Evol.*, 12, 1138–1143.

550 Perrier, A., Sánchez-Castro, D. & Willi, Y. (2020). Expressed mutational load increases toward

551   the edge of a species' geographic range. *Evol.*, 74, 1711–1723.

Phillips, B.L. (2015). Evolutionary processes make invasion speed difficult to predict. *Biol.*

Invasions, 17, 1949–1960.

Phillips, B.L., Brown, G.P., Webb, J.K. & Shine, R. (2006). Invasion and the evolution of speed

in toads. *Nature*, 439, 803–803.

R Core Team (2020). *R: A language and environment for statistical computing*. R Foundation for

Statistical Computing, Vienna, Austria.

Shaw, A.K., Lutscher, F. & Popovic, L. (2023). Pushed to the edge: Spatial sorting can slow

down invasions. *Ecol. Lett.*, 26, 1293–1300.

Skellam, J.G. (1951). Random Dispersal in Theoretical Populations. *Biometrika*, 38, 196–218.

Slatyer, R.A., Hirst, M. & Sexton, J.P. (2013). Niche breadth predicts geographical range size: A

general ecological pattern. *Ecol. Lett.*, 16, 1104–1114.

Szűcs, M., Vahsen, M.L., Melbourne, B.A., Hoover, C., Weiss-Lehman, C. & Hufbauer, R.A.

(2017). Rapid adaptive evolution in novel environments acts as an architect of population

range expansion. *Proc. Natl. Acad. Sci. U.S.A.*, 114, 13501–13506.

Urquhart-Cronish, M., Angert, A.L., Otto, S.P. & MacPherson, A. (2022). Density-dependent

selection during range expansion affects expansion load in life-history traits. *bioRxiv*,

2022.11.08.515702.

Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *J. Stat.*

Softw., 3, 1–48.

Volkova, P.A., Nachatoi, V.A. & Bobrov, A.A. (2023). Hybrid between *Lemna minor* and *L.*

turionifera (*L. × japonica*, Lemnaceae) in East Europe is more frequent than parental

species and poorly distinguishable from them. *Aquat. Bot.*, 184, 103593.

Wagner, N.K., Ochocki, B.M., Crawford, K.M., Compagnoni, A. & Miller, T.E.X. (2017).

Genetic mixture of multiple source populations accelerates invasive range expansion. *J.*

Anim. Ecol., 86, 21–34.

Weiss-Lehman, C., Hufbauer, R.A. & Melbourne, B.A. (2017). Rapid trait evolution drives

increased speed and variance in experimental range expansions. *Nat. Commun.*, 8, 14303.

Williams, J.L., Kendall, B.E. & Levine, J.M. (2016). Rapid evolution accelerates plant

population spread in fragmented experimental landscapes. *Science*, 353, 482–485.

Williams, J.L., Hufbauer, R.A. & Miller, T.E.X. (2019). How evolution modifies the variability

of range expansion. *Trends Ecol. Evol.*, 34, 903–913.

Zilio, G., Krenek, S., Gougat-Barbera, C., Fronhofer, E.A. & Kaltz, O. (2023). Predicting

evolution in experimental range expansions of an aquatic model system. *Evol. Lett.*, 7, 121–

131.

Figure legends

Figure 1. Experimental range expansion across landscapes with and without a spatial gradient in

temperature. (a) Replicate, mixed-genotype populations of *Lemna* expand their experimental

ranges across increasingly warmer temperatures (blue to red) from the range core to the range

front in gradient landscapes, or across benign and constant temperatures in uniform landscapes.

(b) Observed temperatures across gradient (red) and uniform (blue) landscapes during the

experiment. Lines represent the slopes of linear models of temperature by distance. We found a

significant and positive temperature slope in gradient landscapes ($\beta_{gradient}$ = 0.027, 95% CI:

0.027 to 0.028 °C/cm; t = 126.49; P < 0.001) and a slightly negative slope in uniform landscapes

($\beta_{uniform}$ = –0.001, 95% CI: –0.002 to 0.000 °C/cm; t = 68.50; P < 0.01).

Figure 2. Population dynamics of range expansion across landscapes with (red) and without

(blue) a temperature gradient across space. (a) Each line represents the maximum distance

travelled over time for a single, replicate range expansion (mean values shown by thick lines).

(b) Among-replicate variability (CV) in distance travelled over time for gradient and uniform

landscapes. (c) Slope and 95% CI of changes in the log coefficient of variation ratio (lnCVR)

over time, where estimates of lnCVR below zero (dashed grey line) indicate that there is less

among-replicate variability (CV) across gradient than uniform landscapes. (d-e) Population

densities at the range front (i.e., the first 0 to 12 cm) during range expansion. Lines show model

predictions from a logistic regression of density across space, with increasing colour intensity

representing changes over time from day 4 to 24. (f) Slope and 95% CI obtained from a linear

regression of range-front density over time.

Figure 3. Evolutionary dynamics of range expansion across landscapes with and without a

temperature gradient across space. (a) Violin plot for the magnitude of evolution observed in

gradient (red) and uniform (blue) landscapes, as quantified as the Euclidean distance of genotype

frequencies between founder and evolved populations at the range front. Points show the amount

of change in genotype frequencies for each replicate, while dashed lines represent the mean

change. (b) Genotype composition of range-front populations in gradient (red) and uniform

(blue) landscapes. The first two axes of a Principal Coordinates Analysis (PCoA) are shown,

accounting for 43.7% of the total variation in genotype composition. Filled circles are treatment

centroids, while clear symbols are replicate populations in each treatment. Ellipses represent one

standard deviation around treatment centroids, and lines represent distance from the centroid to

each replicate population. (c) Mean genotype frequencies observed across range-front

populations in gradient (left) and uniform (right) landscapes, with different colours representing

unique genotypes. (d) Genotype frequency for gradient (left) and uniform (right) landscapes.

Coloured points and error bars represent observed genotype frequencies and their 95% CIs in

evolved range-front populations. The initial genotype frequency in the founder population is

denoted by 'x'.

Figure 4. Observed versus simulated among-replicate variability in range-front genotype

composition in gradient (top) and uniform (bottom) landscapes. Histograms represent the

predicted probability distribution of among-replicate variability in genotype composition due to

random sampling alone, with black dotted lines representing the 95% confidence interval.

Vertical dashed red and blue lines represent the mean observed among-replicate variability in

genotype composition for gradient and uniform landscapes, respectively. Among-replicate

variability in genotype composition was quantified as the summed Euclidean distance between

the genotypic composition of each replicate population and the mean genotypic composition of

each treatment.

671 **Figure 1**

(a)

(b)

Figure 2

Figure 3

686   **Figure 4**

687