Structural and dynamic changes in P-Rex1 upon activation by PIP$_3$ and inhibition by IP$_4$

Sandeep K. Ravala[1], Sendi Rafael Adame-Garcia[2], Sheng Li[3], Chun-Liang Chen[1], Michael A. Cianfrocco[4], J. Silvio Gutkind[2], Jennifer N. Cash*[5], and John J. G. Tesmer*[1]

[1]Departments of Biological Sciences and of Medicinal Chemistry and Molecular Pharmacology, Purdue University, West Lafayette, Indiana 47907, United States

[2]Department of Pharmacology and Moores Cancer Center, University of California, San Diego, San Diego, CA 92093, USA

[3]Department of Medicine, University of California San Diego, La Jolla, CA 92093, USA.

[4]Department of Biological Chemistry, University of Michigan, Ann Arbor, Michigan 48109, United States

[5]Department of Molecular and Cellular Biology, University of California-Davis, Davis, CA, 95616, USA.

*To whom correspondence should be addressed: jcash@ucdavis.edu and jtesmer@purdue.edu

Abstract

PIP$_3$-dependent Rac exchanger 1 (P-Rex1) is abundantly expressed in neutrophils and plays central roles in chemotaxis and cancer metastasis by serving as a guanine-nucleotide exchange factor (GEF) for Rac. The enzyme is synergistically activated by PIP$_3$ and the heterotrimeric Gβγ subunits, but mechanistic details remain poorly understood. While investigating the regulation of P-Rex1 by PIP$_3$, we discovered that Ins(1,3,4,5)P$_4$ (IP$_4$) inhibits P-Rex1 activity and induces large decreases in backbone dynamics in diverse regions of the protein. Cryo-electron microscopy analysis of the P-Rex1·IP$_4$ complex revealed a conformation wherein the pleckstrin homology (PH) domain occludes the active site of the Dbl homology (DH) domain. This configuration is stabilized by interactions between the first DEP domain (DEP1) and the DH domain and between the PH domain and a 4-helix bundle (4HB) subdomain that extends from the C-terminal domain of P-Rex1. Disruption of the DH–DEP1 interface in a DH/PH-DEP1 fragment enhanced activity and led to a more extended conformation in solution, whereas mutations that constrain the occluded conformation led to decreased GEF activity. Variants of full-length P-Rex1 in which the DH–DEP1 and PH–4HB interfaces were disturbed exhibited enhanced activity during chemokine-induced cell migration, confirming that the observed structure represents the autoinhibited state in living cells. Interactions with PIP$_3$-containing liposomes led to disruption of these interfaces and increased dynamics protein-wide. Our results further suggest that inositol phosphates such as IP$_4$ help to inhibit basal P-Rex1 activity in neutrophils, similar to their inhibitory effects on phosphatidylinositol-3-kinase.

Introduction

Localized activation of signaling is required for proper cell migration. Phosphatidylinositol 3,4,5-trisphosphate (PIP_3)-dependent Rac exchanger 1 (P-Rex1) is a Rho guanine-nucleotide exchange factor (RhoGEF) abundantly expressed in neutrophils that mediates chemotaxis and the generation of reactive oxygen species via activation of Rac GTPases (Dorseuil et al., 1992). The protein is comprised of a catalytic Dbl homology (DH) domain followed by a pleckstrin homology (PH) domain, two DEP domains (DEP1 and DEP2), two PDZ domains (PDZ1 and PDZ2), and a C-terminal inositol polyphosphate-4-phosphatase-like (IP4P) domain (Fig. 1A).

P-Rex1 exhibits low basal activity until it becomes activated via direct interaction with membrane-bound regulators PIP_3 and Gβγ which act synergistically (Cash et al., 2019, 2016; Mayeenuddin et al., 2006; Welch et al., 2002), indicating that they use distinct modes of regulation. Although relatively little is known about how P-Rex1 transitions to an activated state, recent structural studies have defined their docking sites. Gβγ engages a scaffold composed of an amalgamation of the DEP2-PDZ1-PDZ2-IP4P domains and likely helps recruit P-Rex1 to the cell membrane (Cash et al., 2019). In contrast, PIP_3 binds to the PH domain (Hill et al., 2005) in a basic pocket (Cash et al., 2016), but this is not necessary for its recruitment to the cell membrane, implying that PIP_3 instead induces a conformational change that activates the enzyme (Cash et al., 2016). Because domains C-terminal to the catalytic DH domain are well known to be involved in autoinhibition (Chávez-Vargas et al., 2016; Hill et al., 2005; Ravala et al., 2020; Urano et al., 2008), the allosteric change induced by PIP_3 must defeat interdomain contacts and render the catalytic DH domain accessible to its substrate.

Here, we used hydrogen-deuterium exchange mass spectrometry (HDX-MS), cryo-electron microscopy (cryo-EM) single particle analysis (SPA), and small-angle X-ray scattering

(SAXS) along with functional studies and live cell experiments to show that activation of P-Rex1

involves disruption of two different inhibitory interfaces between domains across the length of the

protein. Surprisingly, we found that the PIP_3 headgroup analog IP_4 can reduce P-Rex1 activity by

stabilizing the autoinhibited conformation of the enzyme at physiologically relevant

concentrations, suggesting a previously unknown, additional mechanism of regulation. Our

experiments further suggest that P-Rex1 binding to PIP_3-containing membranes induces

conformational changes that unwind P-Rex1 into a fully active state.

Results

IP_4 induces protection from deuterium incorporation on regions of P-Rex1 distal from the PIP_3-

binding site.

Previous work suggested that PIP_3 binding to the PH domain activates P-Rex1 purely through an

allosteric mechanism (Cash et al., 2016). Thus, we anticipated that binding of the soluble

headgroup of PIP_3, IP_4, to full-length P-Rex1 could also lead to conformational changes

characteristic of the activated state. To test this, we analyzed P-Rex1 in the presence and absence

of IP_4 using HDX-MS. We observed strong protection from deuterium incorporation in the PIP_3-

binding site on the PH domain in the presence of IP_4 (Fig. 1A&B, Sup. Data 1). However, we also

observed strong protection in other regions of the protein: namely on the surface of the PH domain,

particularly in the β5/β6 loop, and in several regions within an extension of the C-terminal IP4P

domain that was not visualized in the P-Rex1–Gβγ complex (Fig. 1A&C) (Cash et al., 2019). We

speculated that these diverse regions form more stable long-range interactions in the presence of

IP_4.

IP$_4$ allosterically inhibits P-Rex1.

Based on our HDX-MS data, we hypothesized that IP$_4$ could inhibit activity of full-length P-Rex1.

Using an *in vitro* GEF activity assay on soluble Cdc42 in the presence of liposomes, we observed

that IP$_4$ inhibits PIP$_3$-mediated activation of P-Rex1 with an IC$_{50}$ value of 1.4 μM (Fig. 1D).

Competition was not observed with Ins(1,4,5)P$_3$, indicating that inhibition is dependent on the 3-

phosphate, which is critical for PIP$_3$ binding to the P-Rex1 PH domain (Cash et al., 2016).

However, IP$_4$ did not affect the activity of the P-Rex1 DH/PH or DH/PH-DEP1 fragments (Sup.

Fig. 1A&B). Collectively, these results indicated that IP$_4$ inhibits P-Rex1 allosterically and that

this inhibition is dependent on long range interactions between the regions shown to be protected

by IP$_4$ in the Gβγ-binding scaffold (DEP2-PDZ1-PDZ2-IP4P) and in the DH/PH-DEP1 module.

Negatively charged liposomes (containing PC/PS), including those that also contain PIP$_3$, inhibit

the GEF activity of the DH/PH-DEP1 and DH/PH fragments (Sup. Fig. 1C). Because full-length

P-Rex1 is not affected by PC/PS liposomes, this suggests that the observed inhibition represents a

non-productive interaction of the DH/PH-DEP1 and DH/PH fragments with negatively charged

surfaces in our assay. The lack of activation of DH/PH-DEP1 by PIP$_3$ prevents us from testing in

this assay whether IP$_4$ can inhibit via direct competition with PIP$_3$ at the PH domain.

IP$_4$ stabilizes long-range interactions mediated by the P-Rex1 DEP1 and PH domains.

To understand the molecular basis of IP$_4$-mediated stabilization and inhibition, we analyzed full-

length P-Rex1 with and without IP$_4$ using cryo-EM SPA (Sup. Fig. 2A). Initial datasets were

collected using a Glacios transmission electron microscope and then processed through 2D

classification (Sup. Fig. 2B-D). In both datasets, most classes showed only the Gβγ-binding core

of P-Rex1 (Sup. Fig. 2C&D). A few classes contained particles with additional mass to the side of

the core, close to the PDZ or DEP domains (Sup. Fig. 2C&D, orange boxes). This mass could

represent either the N-terminal domains or another P-Rex1 particle in proximity. Only in the

sample containing IP_4 could we observe class averages with additional mass next to the core

opposite the side that binds Gβγ (Sup. Fig. 2D, green boxes). Based on its size, location, our HDX-

MS data (Fig. 1), and low-resolution maps of P-Rex1 generated in a previous study (Cash et al.,

2019), this mass most likely corresponded to the N-terminal DH/PH-DEP1 domains interacting

with an elongated subdomain extending from the IP4P domain.

We next collected much larger datasets on the P-Rex1·IP_4 complex using a Krios

transmission electron microscope and determined the structure of this complex at an average

resolution of 4.1 Å (Fig. 2A-B, Table 1, and Sup. Fig. 3). Similar to the P-Rex1–Gβγ complex

(Cash et al., 2019), this sample exhibited a preferred orientation on grids, necessitating the addition

of data collected on a tilted sample (Sup. Fig. 3A,C). The resulting 3D reconstruction clearly shows

the Gβγ-binding core, composed of DEP2, PDZ1, PDZ2, and the majority of the IP4P domain,

from which there are two extensions of density that contact one another to form a loop-like

structure (Fig. 2A and Sup. Fig. 3D). One extension corresponds to a large insertion in the IP4P

domain that contains IP_4-stabilized regions (Fig. 1) and that was disordered in the P-Rex1–Gβγ

complex (Cash et al., 2019). The ordered elements of the insertion form a long 4-helix bundle

(4HB) most similar in fold to focal adhesion targeting (FAT) domains (Hayashi et al., 2002), which

are found in other peripheral membrane proteins involved in cell adhesion and migration. The

other extension corresponds to the DH/PH-DEP1 domains. Based on its distinct shape, the PH

domain (Cash, 2016) was fit into density along the side of the 4HB. Here, IP_4 is observed bound

to the PH domain PIP_3-binding site (Fig. 4B). The position of the DH domain was also obvious,

but individual helices of the DH domain were lower in resolution and more dynamic relative to

the rest of the structure (Sup. Fig. 3D). The relative positions of the DH and PH domains mandate

a severe bend in the helix connecting the DH and PH domains (Fig. 2A&B), resulting in a jack-

knifed configuration of the DH/PH module that blocks access to the GTPase binding site on the

DH domain. The remaining mass, immediately adjacent to the end of the DH domain opposite its

N-terminus, corresponds to DEP1. Using the required connectivity of its N- and C-termini to the

PH and DEP2 domains, respectively, DEP1 was docked in a manner that complemented residues

on the DH domain and corroborated the HDX-MS data (Fig. 4A). Weak density corresponding to

a long 5-turn extension of the αC helix of the PH domain connects to the N-terminus of DEP1, but

its C-terminal connection to DEP2 is disordered, likely explaining lower local resolution in the

DEP1 region (Sup. Fig. 3D). Overall, the conformation of the P-Rex1 Gβγ-binding core is

essentially the same as in the P-Rex1–Gβγ complex (RMSD deviation of 1.1 Å for 701 Cα atoms).

The contact between the PH domain and the 4HB is primarily mediated via the β1 and β2

strands and β5/β6 loop of the PH domain (Fig. 2C&D). In all previous crystal structures including

the P-Rex1 PH domain, this same surface formed extensive protein-protein lattice contacts (Cash

et al., 2016; Lucato et al., 2015). The residues directly involved in the interface are among the most

strongly protected in the presence of IP$_4$ as measured by HDX-MS (Fig. 1A-C, Fig. 4A). The PH

domain β3/β4 loop, which we previously showed to be a nonspecific anionic membrane-binding

loop, remains unstructured and is situated near a loop at the tip of the 4HB that is also unstructured

(residues 1109-1209, Fig. 2C) and contains known phosphorylation sites, some of which regulate

activity (Barber et al., 2012). At the interface, surface hydrophobic residues Leu279 and Ile286 on

the PH β1 and β2 strands and Tyr353 on the β5/β6 loop interact with a surface of 4HB including

Tyr1096 and His1224 (Fig. 2D). Charge complementarity is formed between Lys302 in the PH

domain and Asp1216 in 4HB, and between the 1-phosphate of IP$_4$ and Lys1217. Otherwise, IP$_4$

does not make direct contact with the 4HB. However, because different PH domain ligands

uniquely perturb the conformation of these regions in the PH domain (Cash et al., 2016), IP_4 could

indirectly stabilize this interface by trapping the β1 and β2 strands and β5/β6 loop in a

conformation with higher affinity for 4HB.

The jack-knife in the helix between the DH and PH domains is required to allow the PH

domain to interact with the 4HB and is stabilized by the DEP1 domain docking to the DH domain

(Fig. 2B). DH domain residues Leu173, Leu177, and Leu178 form a hydrophobic interface with

DEP1 residues Ile409, Ile457, Leu466, and Ala469 (Fig. 3A). Leu173 and Leu177 were previously

noted to be conspicuously exposed in structures of the DH/PH tandem (Cash et al., 2016). Thus,

the DEP1 domain stabilizes an inactive DH/PH tandem that is further stabilized by interaction with

the 4HB of the IP4P domain.

While conducting these studies, the structure of human P-Rex1 in the absence of IP_4 was

reported (PDB entry 7SYF) (Chang et al., 2022), allowing a comparison between the IP_4 and IP_4-

free states of autoinhibited P-Rex1 (Sup. Fig. 5). Overall, the domain organization is very similar,

but there is an ~3° rotation of the Gβγ-binding core in the IP_4 complex relative to the PH–4HB

interface such that DEP1 and DEP2 move closer together. It is possible that the binding of IP_4 at

this interface drives this conformational change. A caveat is that the protein used for the 7SYF

structure contained a T4 lysozyme domain inserted into the β3/β4 loop of the PH domain. Although

this domain was not visible in the reconstruction, its proximity to the IP_4 binding site may influence

the global conformation of P-Rex1. The similarities between P-Rex1±IP_4 also suggest that the IP_4-

binding site in the PH domain is freely accessible in the autoinhibited state. PIP_3 would, however,

not have access because the 4HB domain would block binding to a membrane surface.

The DH–DEP1 interface contributes to autoinhibition in vitro.

To test the contribution of the observed DH–DEP1 interface to autoinhibition, the interface was

disrupted by site-directed mutagenesis in the context of the DH/PH-DEP1 fragment. This fragment

is 5- to 10-fold less active (depending on assay conditions) than the DH/PH tandem alone (Sup.

Fig. 1C) (Ravala et al., 2020), confirming a specific role for DEP1 in autoinhibition (Fig. 3C).

Single point mutations in the interface profoundly affected GEF activity on soluble Cdc42. The

L173A variant had ~2.5-fold higher activity, whereas L177E and L178E exhibited 4- to 5-fold

higher activity, similar to the activity of DH/PH without the DEP1 domain. In the DEP1 domain,

I409A and L466A mutations resulted in ~2- and 4-fold higher activity, respectively. The α1/α2

loop (residues 77-90) of the DH domain, although ordered in previous structures of DH/PH bound

to GTPases (Cash et al., 2016; Lucato et al., 2015), is disordered in our structure (Fig. 3B).

However, modeling the DH domain from previous DH/PH crystal structures suggests that Lys89

and Arg78 would be close enough to form a bipartite ionic interaction with Glu456 in the DEP1

domain. Consistent with this hypothesis, the E456K variant was ~2.5-fold more active.

Surprisingly, one of the intended disruptive mutations, A170K, instead inhibited GEF activity by

~50% (Fig. 3C). Based on the available structures, Lys170 in the DH domain could form

complementary electrostatic interactions with Glu411 along with additional non-polar interactions

with the backbone and side chains of residues 409, 411, and 412 in the DEP1 domain, thereby

stabilizing the DH–DEP1 interface (Fig. 3B).

The DH—DEP1 interface stabilizes DH/PH-DEP1 and decreases flexibility.

Because disruption of the DH–DEP1 interface led to increased activity, we predicted that these

variants would have a more extended DH/PH module that is likely more dynamic. To test this,

wild-type (WT) DH/PH-DEP1 and its variants were assessed using a ThermoFluor assay to

determine their melting temperatures (T_m). Indeed, variants with increased activity also had lower

T_m values (Fig. 3D, Table 2). Conversely, A170K, which was less active than WT, showed a higher

T_m. We also analyzed these variants using size exclusion chromatography coupled to small-angle

X-ray scattering (SEC-SAXS) (Sup. Fig. 6A&B). In previous SAXS analyses, we observed that

DH/PH-DEP1 exhibited a more compact state with a smaller conformational landscape in solution

relative to DH/PH (Ravala et al., 2020). We hypothesized that mutations which disrupt the DH–

DEP1 interface would likewise lead to more elongated ensembles. Compared to WT DH/PH-

DEP1, which had a radius of gyration (R_g) of 30 ± 0.3 Å, the L177E variant (the most active

DH/PH-DEP1 variant tested) had an R_g of 31 ± 0.2 Å, suggestive of expansion (Fig. 3E and Table

3). Variants A170K and I409A had R_g values similar to that of WT. Kratky plots indicated that all

the samples had heterogeneous conformations (Sup. Fig. 6C). The shapes of the P(r) functions

were similar for all variants except a longer tail for L177E and I409A, consistent with a higher

proportion of extended conformations in solution (Sup. Fig. 6D).

Because the samples exhibited a high degree of heterogeneity in solution, the

conformational distribution of these variants was assessed using the ensemble optimization method

(EOM) (Fig. 3E&F, Sup. Fig. 6A) (Tria et al., 2015). The resulting ensemble for WT shows

predominant conformations with R_g values ~ 28 Å and a small fraction of extended conformations

with R_g ~ 39 Å (Fig. 3E). The selected ensemble for A170K had R_g values similar to WT (~29 Å),

however the A170K peak is broader in comparison, suggesting conformational heterogeneity and

structural changes. The L177E variant exhibited a larger shift to higher R_g, with an average R_g ~

30 Å, and a second significant population with R_g > 32 Å (Fig. 3E). Similar to the distribution of

R_g values, the D_{max} function distribution shows that the L177E variant had the most extended

conformation of the variants tested (Fig. 3F). Other analyses were also consistent with L177E

being the most flexible and extended variant (see methods).

Flexibility of the hinge in the α6-αN helix of the DH/PH module is important for autoinhibition.

One of our initial goals in this project was to determine a high-resolution structure of the

autoinhibited DH/PH-DEP1 core by X-ray crystallography. To this end, we started with the

DH/PH-DEP1 A170K variant, which was more inhibited than wild-type but still dynamic, and then

introduced S235C/M244C and K207C/E251C double mutants to completely constrain the hinge

in the α6-αN helix via disulfide bond formation in a redox sensitive manner. Single cysteine

variants K207C and M244C were generated as controls. The S235C/M244C variant performed as

expected, decreasing the activity of the A170K variant to nearly background in the oxidized but

not the reduced state (Sup. Fig. 4). However, the M244C single mutant exhibited similar effects,

suggesting that it forms disulfide bonds with cysteine(s) other than S235C. Indeed, the side chains

of Cys200 and Cys234 are very close to that of M244C. The K207C/E251C mutant was similar to

S235C/M244C under oxidized conditions, but ~15-fold more active (similar to WT DH/PH levels,

see Fig. 3C) under reducing conditions. The K270C variant, on the other hand, exhibited higher

activity than A170K on its own under oxidizing conditions, but similar activity to all the variants

except K207C/E251C when reduced. These results suggest that K207C/E251C in a reduced state

and K270C in an oxidized state favor a configuration where the DEP1 domain is less able to engage

the DH domain and maintain the kinked state. The mechanism for this is not known. Regardless,

these data show that perturbation of contacts between the kinked segments of the α6-αN helix can

have profound consequences on the activity of the DH/PH-DEP1 core.

Interactions at the P-Rex1 DH–DEP1 and PH–4HB interfaces contribute to autoinhibition in cells.

To evaluate the roles of the P-Rex1 DH–DEP1 and PH–4HB interfaces in living cells, we utilized

SRE luciferase-gene reporter assays as a read out of full-length P-Rex1 activity in HEK293T cells.

Mutations at the DH–DEP1 interface had a strong effect on activity, with L177E and L466E (Fig.

3A) exhibiting ~10-fold and ~4-fold higher activity relative to WT, respectively (Fig. 5A).

Perturbation of the PH–4HB interface (Fig. 2D) also increased activity in that Y1096A was ~5-

fold more active than WT (Fig. 5A). Mutation of other residues in the PH–4HB interface (Fig. 2D)

also increased activity, although to a lesser extent.

The most affected variants (L177E, L466E, and Y1096A) were then tested for their effects

on cell migration in response to chemokine gradients (Fig. 5B&C, Sup. Fig. 7A&B). For this, the

endogenous P-Rex1 gene in HeLa cells was first knocked out by CRISPR-Cas9 (Sup. Fig. 7C) and

the resulting cells were transfected with various P-Rex1 constructs (Sup. Fig. 7D). Cell migration

was then evaluated in the presence or absence of CXCL12 (upstream of P-Rex1 and Gβγ signaling)

and epidermal growth factor (EGF). CXCL12-induced chemotaxis was dependent on the

expression of P-Rex1 (Fig. 5B&C and Sup. Fig. 7), and all three variants caused a significantly

larger number of cells to migrate. However, EGF-induced chemotaxis in HeLa cells, which is not

dependent on P-Rex1, was unaffected by P-Rex1 expression. These data support that the DH–

DEP1 and PH–4HB interfaces of P-Rex1 mediate autoinhibition that specifically modulates

chemokine-induced cell migration.

P-Rex1 binding to PIP$_3$-containing model membranes induces a more open, dynamic

conformation.

Given the known binding site for Gβγ (Cash et al., 2019), the position of membrane-binding loops

such as the β1/β2 loop of the DEP1 domain (Ravala et al., 2020), and the position of the 4HB

domain, it does not seem that the PH domain would be able to interact with PIP_3 in a cell membrane

while P-Rex1 is in its autoinhibited conformation. To better understand the molecular

consequences of PIP_3 binding, HDX-MS measurements were taken on P-Rex1 in the presence of

liposomes ± PIP_3. Without PIP_3, the most notable changes in P-Rex1 in response to liposomes

were increases in protection of the lipid binding elements of both DEP domains (primarily their

β1/β2 loops; Sup. Fig. 8A, Sup. Data 2), which suggests that they directly interact with lipid

bilayers. Indeed, the isolated DEP1 can independently bind negatively charged liposomes (Ravala

et al., 2020). Within the PH domain, there was deprotection of the αC helix, but there was no

deprotection at the PH–4HB interface, suggesting that it remained intact. In contrast, membranes

with PIP_3 caused deprotection of the entire DH domain, the α6-αN linker and the helices at the

DH–DEP1 interface (Fig. 6A, Sup. Fig. 8B, Sup. Data 2). Although the core of the PH domain

showed, as expected, an increase in protection in the presence of PIP_3, the structural elements

contacting 4HB became deprotected, as did the PH-binding regions of 4HB. Collectively, these

data are consistent with loosening of interdomain contacts and unraveling of at least some fraction

of P-Rex1 onto the surface of the liposome.

Discussion

Here, we showed that IP_4 binding to the PIP_3-binding site in full-length P-Rex1 stabilizes

a closed, autoinhibited conformation of P-Rex1 by enhancing long-range contacts across the length

of the protein. DH–DEP1 and PH–4HB interactions are relieved upon P-Rex1 binding to PIP_3-

containing liposomes, leading to its activation. Canonically, PIP_3 signaling is negatively regulated

by PIP$_3$ phosphatases such as PTEN. Additionally, inositol phosphates, including IP$_7$ and IP$_4$, can

compete with PIP$_3$ binding to PH domains (Jia et al., 2007), representing another form of negative

regulation. Ins(1,3,4,5)P$_4$ is a major isoform of IP$_4$ in neutrophils (Stuart et al., 1994) where its

concentration is estimated to be 4 μM (French et al., 1991) and where P-Rex1 is highly expressed

(Welch et al., 2002). Because we can measure significant inhibition of P-Rex1 by IP$_4$ at > 1 μM

(Fig. 1D), IP$_4$ may exert biologically relevant control of Rac activation, at least in neutrophils.

Since P-Rex1 can bind membranes in the absence of PIP$_3$ or Gβγ, IP$_4$ may serve to suppress P-

Rex1 activity until a threshold concentration of PIP$_3$ is generated, allowing rapid activation of

already membrane-associated P-Rex1. It is worth noting that this regulation by specific inositol

phosphates may depend on the intracellular distribution of the enzymes responsible for their

synthesis (Gokhale et al., 2011).

Another key result of our study was to provide a molecular explanation for how the PH

and DEP1 domains contribute to P-Rex1 regulation. Nearly two decades ago, it was first reported

that domains located C-terminal to the DH domain contribute to P-Rex1 autoinhibition (Hill et al.,

2005). Deletion of the PH domain in P-Rex1 resulted in a large increase in activity in the context

of the full-length enzyme. However, the DH/PH fragment has higher activity than DH/PH-DEP1

and larger P-Rex1 fragments, indicating that the PH domain itself is not intrinsically inhibitory, as

it is in some other RhoGEF DH/PH tandems (Bandekar et al., 2019; Chen et al., 2020)(Ravala and

Tesmer, 2023). This apparent contradiction can now be explained by the fact that in DH/PH-DEP1

and larger fragments, the DH/PH module can jack-knife and position the PH domain in a manner

that blocks GTPase binding. This model of regulation was corroborated by recent structural studies

of P-Rex1 without IP$_4$ (Chang et al., 2022). The isolated DEP1 domain was also shown to play an

autoinhibitory role based on the relatively low activity of DH/PH-DEP1 relative to DH/PH (Ravala

et al., 2020). This can now be explained by its interaction with the DH domain, which,

consequently, positions the PH domain to block access of GTPases. Because the PH and DEP1

domains can mediate inhibition in the contexts of both DH/PH-DEP1 and the full-length enzyme,

and because there is little GEF activity when the DH–DEP1 interface is intact, the DH/PH-DEP1

module can be thought of as the core signaling circuit in P-Rex1. In support of this, our mutations

in the DH–DEP1 interface were, in general, more activating than those in the PH–4HB interface

in cells, although we note that our mutagenesis was not exhaustive (Fig. 5).

Using our functional data along with the known structures of P-Rex1, we assembled a

model for the activation of P-Rex1 by PIP_3 and Gβγ (Fig. 6B). P-Rex1 in its basal state may be

associated with IP_4 and exist in equilibrium at the membrane or in the cytosol. Indeed, in our HDX-

MS studies, the DEP1 and DEP2 domains show protection in the presence of liposomes even

without PIP_3 (Sup. Fig. 8, Sup. Data 2). In its autoinhibited configuration, the known membrane

anchoring elements (the GTPase binding site of the DH domain, the PIP_3-binding site of the PH

domain, the β3/β4 loop of the PH domain, the β1/β2 loops of DEP1 and DEP2, and the Gβγ

binding site in the C-terminal domains) cannot engage a common membrane plane (Fig. 6B). The

clear outlier is the PH domain which would not be able to engage the membrane along with the

other domains without unwrapping of the autoinhibited conformation. PIP_3 is not major driver of

membrane anchoring on its own (Barber et al., 2007; Cash et al., 2016), and nor is Gβγ, but they

do so synergistically in combination (Barber et al., 2007). In prior HDX-MS studies, Gβγ binding

did not have a large effect on the regions now known to be involved in autoinhibition and only

caused protection in regions of direct contact (Cash et al., 2019). This supports the idea that its

role in activation may be primarily related to translocation. This is consistent with the observation

that the ΔPH variant of P-Rex1 (which cannot form the DEP1–DH or PH–4HB interfaces or bind

PIP$_3$) is activated to the same extent by Gβγ as WT P-Rex1 (Hill et al., 2005). In autoinhibited P-

Rex1, the Gβγ binding site is readily accessible whereas that of PIP$_3$ is sequestered.

We propose that, after stimulation of GPCRs in neutrophils, Gβγ likely binds first and, with

the assistance of membrane binding elements in the DEP1 and 2 domains and possibly the β3/β4

loop of the PH domain, promotes loosening of the autoinhibited state (Fig. 6B). Generation of PIP$_3$

by PI3K then releases the DH/PH module from the 4HB and DEP1 domain, displacing any bound

IP$_4$. Because neither PIP$_3$ nor IP$_4$ can activate DH/PH-DEP1 GEF activity on a soluble GTPase

(Sup. Fig. 1A&B), we speculate that this unwrapping at the membrane with multiple points of

engagement across the protein is necessary for full activation of P-Rex1 (Fig. 6B). Indeed, our

HDX-MS and SAXS studies here support that fully activated P-Rex1 at the membrane will be

more extended and dynamic, making the DH domain more accessible to Rac1. However, what

remains unknown is the mechanism by which the PH domain is able to access PIP$_3$ at the

membrane, even in a "loosened" autoinhibited state. Furthermore, additional layers of P-Rex1

regulation exist that remain underexplored. For example, phosphorylation of the lipid binding loop

of DEP1 by PKA is known to inhibit P-Rex1 (Chávez-Vargas et al., 2016; Ravala et al., 2020).

Also, a potential interaction may occur between the basic β3/β4 loop of the PH domain and the

loop at the end of the 4HB (Fig. 2C). Both loops, consistent with their extended and dynamic

nature, have predicted and confirmed phosphorylation sites (Barber et al., 2012) and thus could

potentially modulate P-Rex1 activity if they interact. Phosphorylation of the basic β3/β4 loop

might be expected to inhibit activity based on the fact that it binds and localizes the protein to the

negatively charged plasma membrane, consistent with dephosphorylation of the loop leading to

activation (Montero et al., 2016, 2013).

Materials and Methods

Cloning and site-directed mutagenesis

Full-length human P-Rex1, Cdc42 and DH/PH-DEP1 expression constructs were described previously (Cash et al., 2019, 2016; Ravala et al., 2020). Mutations in DH/PH-DEP1 were created using QuikChange (Qiagen) and confirmed by DNA sequencing. Mutations in the pCEFL-HA-HaloTag-P-Rex1 WT construct were created by QuikChange II site-directed mutagenesis (Agilent 200523). All constructs were confirmed by sequencing and expression was tested by immunoblot.

Protein purification

Full-length P-Rex1 was transiently expressed in Freestyle 293-F cells and purified as discussed previously (Cash et al., 2019). Briefly, 48 h after transfection, the cells were harvested and lysed with Cell Lytic M (Sigma). After ultracentrifugation to remove the insoluble fraction, the protein was purified using glutathione agarose resin (Gold Biotechnology Inc.). The protein was subjected to TEV cleavage to remove the GST tag and then further purified using a Mono Q 5/50 GL anion exchange column (GE Healthcare Life Sciences). Finally, the protein was purified over an affinity column generated by conjugating human Rac1 to Affi-Gel 10 resin, although for Krios cryo-EM and HDX-MS experiments, this step was omitted.

P-Rex1 DH/PH-DEP1 proteins were expressed and purified as described previously (Ravala et al., 2020). Briefly, His-tagged protein was expressed in *E. coli* BL21(DE3) cells which were then lysed using an Avestin Emulsiflex-C3 high-pressure homogenizer. The cell lysate was clarified with high-speed centrifugation, the supernatant was collected, and protein was purified using Ni-NTA resin. The protein was subjected to TEV cleavage to remove the tag. The protein was then purified using a HiTrap SP sepharose column, concentrated, and subjected to size

exclusion chromatography on a Superdex S75 column (GE Healthcare) column. Cdc42 was

produced in an unprenylated form in *E. coli* and purified as previously described (Cash et al.,

2016).

Hydrogen-deuterium exchange mass spectrometry

HDX-MS experiments were performed as previously described (Cash et al., 2019). Briefly,

samples were mixed with D_2O buffer to initiate the HDX reaction and, at various time points, the

reaction was quenched with ice cold quench buffer and the samples immediately frozen on dry ice.

Samples were thawed at 4 °C and subjected to enzymatic digestion on an immobilized pepsin

column followed by LC separation and MS analysis. Data were analyzed using HDExaminer

(Sierra Analytics, LLC, Modesto, CA). Each sample was analyzed twice by HDX-MS, and the

data shown in graphs represent the average of these experiments. Each image in Supplemental

Data 1 and 2 shows one experiment (rainbow plots) or a difference analysis from those experiments

(red to blue plots). Only one of the two sets of experiments performed for each condition (+/-

liposomes or +/- IP4) is shown here. For each peptide, the average of all five time points was

calculated and used to plot the difference data onto the coordinates. Coordinates are colored using

a range of -20% (darkest blue, protection) to 20% (darkest red, deprotection). P-Rex1 was used at

a concentration of 1.7 mg/ml. IP_4 (Cayman Chemical) was added at a concentration of 100 μM.

For experiments with liposomes, liposomes were added at a molar ratio of 1 P-Rex1 to 4000 total

lipids. Liposomes were composed of 80:80:1 POPC:POPS:PIP_3 and prepared as previously

described (Cash et al., 2019).

Guanine-nucleotide exchange assays

Proteins were evaluated for their GEF activity using a fluorescence-based assay (Cash et al., 2019).

Briefly, N-methyl-anthraniloyl-GDP (mant-GDP) loaded soluble Cdc42 was used as a substrate

GTPase (2 µM) in a buffer containing 20 mM HEPES pH 8, 100 mM NaCl, 0.5 mM $MgCl_2$, 100

µM GTP, and reactions were started by addition of P-Rex1 (100 nM). The loss of fluorescence

was measured over time at 10 s intervals on a Flexstation 3 plate reader for 40 min. The data was

fit to the one-phase exponential decay model in GraphPad Prism with the span (Y_o-plateau) shared

among samples. For IP_4 competition curves, GEF assays were carried out in the presence of

liposomes containing 2.5 µM PIP_3, as indicated, and 200 µM each of POPC and POPS, prepared

as described previously (Cash et al., 2019).

Cryo-EM grid preparation and data collection

For cryo-EM sample preparation, P-Rex1 was used at a final concentration of 3 µM and *n*-dodecyl-

β-D-maltoside (DDM) was added to a final concentration of 0.08 mM. For samples with IP_4, a

final concentration of 40 µM IP_4 was added. A sample of 4 µl was applied to a glow-discharged

Quantifoil (1.2/1.3) 300-mesh grid which was then blotted with filter paper and plunge-frozen into

liquid ethane cooled with liquid nitrogen using a Vitrobot Mark IV (Thermo Fisher Scientific) set

to 4 °C, 100% humidity, 4 second blot, and a force of 10. Micrographs were collected either using

Leginon (Suloway et al., 2005) on a Glacios transmission electron microscope (Thermo Fisher

Scientific) operating at 200 keV and a K2 Summit direct electron detector (Gatan, Inc.) in counting

mode (0.98 Å/pixel) at a nominal magnification of 45,000x or using EPU (Thermo Fisher

Scientific) on a Titan Krios transmission electron microscope (Thermo Fisher Scientific) operating

at 300 keV and a K3 direct electron detector (Gatan, Inc.) in counting mode (1.054 Å/pixel) at a

nominal magnification of 81,000x. On the Krios, datasets were collected on both untilted and 30°

tilted grids (Table 1).

Cryo-EM data processing

To overcome the severe preferred orientation problem of our sample on grids, we collected data on 0˚ and 30˚ tilted samples on a Krios electron microscope and processed these datasets separately up through 2D classification (Table 1). For each dataset, micrograph assessment, particle picking, and contrast transfer function estimation were performed using Warp (Tegunov and Cramer, 2019). Particle stacks were taken into CryoSPARC (Punjani et al., 2020, 2017) and extensively cleaned using 2D classification. A final merged particle stack was used for *ab initio* reconstruction into one class followed by non-uniform refinement to obtain a map at an overall 4.1 Å resolution for the P-Rex1·IP$_4$ complex. For Glacios datasets, data were processed only through 2D classification.

Model building and refinement

Initial model building relied on docking existing atomic models for the DH and PH domains of P-Rex1 (PDB entries 5FL1 and 5D3Y) and the Gβγ-binding scaffold (PDB entry 6PCV). 5D3Y was used for the PH domain because the maps were consistent with IP$_4$ bound to the PIP$_3$ site of the PH domain. The DEP1 domain was placed using a non-domain swapped atomic model derived from PDB entry 6VSK. The linker between the PH and DEP1 domains and the 4HB domain were built by hand. A Dali search (Holm and Laakso, 2016) using backbone helices of 4HB revealed its topology to be similar to FAT domains, which was then used to adjust the register of each of its four helices. When Alphafold2 (Jumper et al., 2021) became available, it was used to further adjust the modeling of the 4HB domain and associated structural elements in the IP4P domain. Finally, when the cryo-EM structure of P-Rex1 (PDB entry 7SYF) and the crystal structure of the DH/PH-DEP1 module (PDB entry 7RX9) became available, they were used to confirm less certain regions.

Final rounds of real space refinement iterating with manual building were performed in Phenix

(Liebschner et al., 2019). Final structure statistics are given in Table 1, and the structure and

associated maps were deposited as PDB entry 8TUA and EMDB entry EMD-41621. Raw data

were deposited as EMPIAR entry EMPIAR-11967.

Structure visualization

UCSF ChimeraX (Pettersen et al., 2021) was used to make figures showing cryo-EM maps.

PyMOL (The PyMOL Molecular Graphics System, Version 2.5.5 Schrödinger, LLC) was used to

create all other structure images.

Size exclusion chromatography coupled to small-angle X-ray scattering (SEC-SAXS)

For in solution characterization of DH/PH-DEP1 and its variants, SEC-SAXS was conducted at

the BioCAT beamline (Sector18) at the Advanced Photon Source, Argonne National Laboratory

using an AKTA Pure FPLC and a Pilatus3 X 1M detector. Purified proteins were injected onto a

Superdex 200 Increase 10/300 column at a final concentration of 3-5 mg/ml in 20 mM HEPES,

pH 7, 300 mM NaCl, 2% glycerol, and elution from this column flowed into SAXS flow cell for

X-ray scattering. Data were collected every 1 s with 0.5 s exposure times at room temperature

using 12 KeV X-rays (1.033 Å wavelength) and a 3.67 m sample-to-detector distance. The

achievable q range for this experimental setup was (0.0043-0.3546 Å).

SAXS Analysis

The scattering data were processed using BioXTAS RAW 1.6.3 software (Hopkins et al., 2017)

(Sup. Fig. 6A) and used to determine the forward scattering I(0) and the radius of gyration, R_g via

Guinier analysis (Sup. Fig. 6B). The Kratky plot showed proteins to be flexible (Sup. Fig. 6C)

leading to unsuccessful rigid modeling efforts. However, use of the ensemble optimization method

(EOM) within ATSAS/3.0.5-2 aided in generating ensembles representing distinct conformational

states of the DH/PH-DEP1 fragment and its variants A170K, L177E, and I409A in solution at

equilibrium (Ravala et al., 2020). For EOM, the models were generated using crystallographic

coordinates from their respective crystal structures: PDB 5FI1 and PDB 6VSK. EOM generated

50,000 possible profiles for the full pool using default settings and native-like structures. From

these profiles, a sub ensemble that matches the experimental scattering data is selected by a genetic

algorithm run 10 times using default settings to verify the stability of the results (results from 1

run are shown in Fig. 3E&F). The pair distance distribution function P(r), which provides

maximum particle dimension of each protein, was calculated using GNOM (DI, 1992) from the

ATSAS 2.8.4 package (Franke et al., 2017) (Sup. Fig. 6D).

For quantitative analyses of the flexibility of the selected ensembles, R_{flex} and R_{σ} metrics

were derived from EOM data. WT DH/PH-DEP1 and A170K show R_{flex} values smaller than those

of the pools, which indicates that these proteins do not exhibit a fully flexible conformation. L177E

and I409A variants exhibit R_{flex} values close to those of the pool, suggestive of being highly

flexible. Since Guinier analyses and the normalized residual fits of the proteins show that SAXS

data quality is good (Sup. Fig. 6B), the R_{σ} value >1 is due to flexibility in the protein, consistent

with the EOM analysis (Fig. 3E&F, Sup. Fig. 6A). Results of the SAXS analysis are presented in

Table 3 in accordance with the revised guidelines for publishing SAXS data (Trewhella et al.,

2017). The SAXS data are deposited in SASBD (https://www.sasbdb.org/) with accession codes

SASDUF2 for DH/PH-DEP1 WT, SASDUG2 for DH/PH-DEP1 A170K, SASDUH2 for DH/PH-

DEP1 L177E and SASDUJ2 for DH/PH-DEP1 I409A.

Differential scanning fluorimetry

ThermoFluor experiments were performed on a QuantStudio 5 Real-Time PCR system in duplicate

with n =3 independent experiments. Purified DH/PH-DEP1 and its variants were incubated at 1

mg/ml in a buffer containing 20 mM HEPES pH 7.0, 300 mM NaCl, and 2 mM DTT with 2.5x

Sypro Orange dye in a final volume of 10 µl in a 384-well PCR plate. Fluorescence was monitored

as a function of temperature, and the T_m was determined by fitting the fluorescence data to a

sigmoidal curve and calculating the inflection point in GraphPad Prism.

Luciferase-gene reporter assay

HEK293T cells seeded in 12-wells plates coated with poly-D-lysine were transfected with 500 ng

of empty vector pHTN HaloTag CMV-neo (Promega G7721) or pCEFL-HaloTag-P-Rex1

constructs and co-transfected with 500 ng of SRE-firefly luciferase and 50 ng of Renilla luciferase

plasmids. Thirty-six hours after transfection, the cells were serum-starved overnight and then

luminescence signal was measured using Dual-Glo assay system (Promega E2920) according to

the manufacturer's instructions. Firefly-luminescence reads were normalized with Renilla-

luminescence signal and adjusted to the negative control.

Preparation of P-Rex1 KO in HeLa cells and cell migration assays

HeLa cells were lentiviral transduced with pLentiCRISPRv2 P-Rex1 guide RNA3 -

AGGCATTCCTGCATCGCATC (Genscript SC1678). Forty-eight hours after transduction, HeLa

cells were selected with puromycin [3 µg/mL] for 7 days. P-Rex1 KO was confirmed by western

blot. Chemotactic migration was measured by trans-well assays (Thermoscientific 140629). Inserts

of 24-well plates were pre-treated with Fibronectin [50 µg/mL] for 3 hours at 37 °C. Subsequently,

$5x10^4$ HeLa cells prepared in serum-free DMEM were plated on the inserts at the upper chamber.

Human CXCL12 [50 ng/mL] and EGF [50 ng/mL] (Sigma Aldrich SRP3276 and SRP6253) were

prepared in serum-free DMEM and used as chemoattractant in the lower chamber. Serum-free

DMEM was used as negative control. The plates with the inserts were incubated in a humidified

atmosphere at 37 °C and 5% CO_2 for 6 h. After incubation, the cells at the upper surface of the

membrane insert were carefully removed and the cells attached to the lower surface were gently

washed with PBS and then fixed with 4% paraformaldehyde for 15 min. After fixation, the cells

were gently washed with PBS followed by staining with 0.5% crystal violet for 20 min. Excess of

crystal violet was removed by gentle PBS washes. Migrating cells were imaged using an inverted

microscope. Quantification of particles corresponding to migrating cells was performed with FIJI

software.

Western blot

Protein samples prepared in Laemmli buffer were separated with SDS-PAGE using 4-12%

gradient gels followed by transfer to PVDF membranes. The membranes were blocked using 5%

non-fat milk in TBS-0.05% tween20 (TBST) and incubated overnight at 4 °C with primary

antibodies against P-Rex1 and GAPDH (Cell signaling technology #13168 and #5174

respectively). The membranes were washed three times with TBST and then incubated with

secondary antibodies in blocking solution for 2 h at room temperature. After washing three times

with TBST, the reactive bands were visualized using ECL detection reagents and CL-X-posure

films.

Statistical analysis

GEF assays described in this study were performed with n>=3 replicates. Statistical significance

was determined using one-way ANOVA test with a post hoc Dunnett's test for multiple

comparisons.

Acknowledgements

We thank Dr. Srinivas Chakravarthy, Beamline Scientist, for performing SEC-SAXS at BioCAT,

Chicago, IL and Dr. Jesse B. Hopkins, Deputy Director, BioCAT for help with SAXS data analysis

and interpretation. This research used resources of the Advanced Photon Source,

a U.S. Department of Energy (DOE) Office of Science User Facility operated for the DOE Office

of Science by Argonne National Laboratory under Contract No. DE-AC02-06CH11357. BioCAT

was supported by grant P30 GM138395 from the National Institute of General Medical Sciences

of the National Institutes of Health. We thank Dr. Thomas Klose in the Purdue Cryo-EM Facility

for technical assistance in cryo-EM data collection. Research reported in this publication was

supported by National Institutes of Health grants CA254402, CA221289, HL071818,

P30CA023168 (J.J.G.T.), by the Walther Cancer Foundation (J.J.G.T.), and by the National

Institute of General Medical Sciences of the National Institutes of Health grant R35GM146664

(J.N.C.).

 Table 1. Cryo-EM data collection, refinement, and validation statistics

	Structure: P-Rex1–IP$_4$ (EMDB: EMD-41621) (PDB: 8TUA) (EMPIAR: EMPIAR-11967)	
	Untilted	*Tilted*
Data collection		
Grids	Carbon Quantifoil	Carbon Quantifoil
Vitrification method	FEI Vitrobot	FEI Vitrobot
Microscope	Titan Krios	Titan Krios
Magnification	81000	81000
Voltage (kV)	300	300
Stage tilt (°)	0	30
Detector	K3 DED	K3 DED
Recording mode	Counting	Counting
Total electron exposure (e–/Å^2)	57.8	57.8
Number of frames	40	40
Defocus range (μm)	0.2 – 2.0	0.2 – 2.0
Pixel size (Å)	1.054	1.054
Data processing		
Number of micrographs	2,127	3,069
Initial particle images (no.)	806,067	1,620,545
Final particle images (no.)	89,450	119,739
Initial particle images merged (no.)	209,189	
Final total particle images (no.)	187,734	
Symmetry	C1	
Map resolution (Å)	4.1	
Refinement		
Initial model used (PDB code)	6PCV, 6VSK, 5D3X, 5FI1, 7RX9	
Model resolution (Å)	4.1	
FSC threshold	0.143	
Map sharpening *B* factor (Å^2)	-176	
Model composition		
Non-hydrogen atoms	10,693	
Hydrogens	10,720	
Protein residues	1,330	
Ligands	1 (4IP)	
B factors (Å^2; min/max/mean)		
Protein	30.9/202/123	
Ligand	143/143/143	
R.m.s. deviations		
Bond lengths (Å)	0.003	
Bond angles (°)	0.505	
Validation		
MolProbity score	2.03	
Clashscore	13.88	
Rotamer outliers (%)	0.94	
CaBLAM outliers (%)	2.2	
Ramachandran plot (%)		
Favored	94.5	
Allowed	5.5	
Outliers	0	

Model vs. Data	
CC mask	0.67
CC box	0.73
CC peaks	0.58
CC volume	0.66
Mean CC for ligand	0.67

Table 2. Thermofluor measurements of DH/PH-DEP1 variants

DH/PH-DEP1	T_m (°C)
WT	44.0 ± 0.2
A170K	45.4 ± 0.3 (p<0.0001)
L173A	43.0 ± 0.06 (p<0.0001)
L177A	43.4 ± 0.4 (p=0.0076)
L177E	43.4 ± 0.3 (p=0.0041)
L178A	43.1 ± 0.1 (p<0.0001)
L178E	42.9 ± 0.1 (p<0.0001)
I409A	41.7 ± 0.8 (p<0.0001)
E411K	43.9 ± 0.1 ns[b]
K415A	43.2 ± 0.5 (p=0.0047)
L451A	41.3 ± 0.5 (p<0.0001)
E456K	ND[a]
L466A	ND

From 2 independent experiments performed in triplicate.
P values are from one-way Anova comparisons with WT.
[a] ND: Not determined because
inflection point not observed.
[b] ns: nonspecific.

Table 3. SAXS parameters for DH/PH-DEP1 variants

	WT	A170K	L177E	I409A
Guinier Analysis				
$I(0)^a$	0.0081 ± 0.00004	0.0021 ± 0.003	0.0092 ± 0.00004	0.0034 ± 0.00002
R_g (Å)	30 ± 0.3	29 ± 0.08	31 ± 0.02	30 ± 0.4
Q_{min} (Å^{-1})	0.0047	0.005	0.0047	0.0047
Q_{max} (Å^{-1})	0.353	0.353	0.353	0.353
P(r) Analysis				
D_{max} (Å)	97	90	110	104
Volume (Å^3)	73900	75000	72500	74900
$MM_{exp}(MM_{cal})$ (kDa)	54 (54)	56 (54)	54 (54)	55 (55)
EOM Analysis				
Crystal Structures	5FI1;6VSK	5FI1;6VSK	5FI1;6VSK	5FI1;6VSK
q-Range (Å^{-1})	.00475-0.353	0.00446-0.353	.00475-0.353	00475-0.353
R_{flex}	70.1% (82.6%)	71.9% (85.3%)	82.9% (84.9%)	79.1% (84.9%)
R_σ	1.18	0.60	1.08	1.56
Skewness	2.39 / 0.41	1.11/ 0.40	0.85 / 0.42	1.50/ 0.40
Kurtosis	4.86/ -0.08	2.09 / -0.14	-0.27 / -0.12	0.77 / -0.14

[a] SAXS parameters $I(0)$, R_g, D_{max}, q_{min}, q_{max}, MM_{exp}, MM_{cal}, R_{flex}, R_σ are the experimentally determined intensity at zero scattering angle, radius of gyration, maximum particle dimension, minimum scattering angle, maximum scattering angle, molecular mass calculated from scattering data, molecular mass calculated based on amino acid sequence, flexibility metric of ensemble in comparison (pool value in parentheses), and ratio of standard deviation for the distribution of selected ensemble to that of pool, respectively. The values for EOM analysis are from the last run of the genetic algorithm.

References

Bandekar SJ, Arang N, Tully ES, Tang BA, Barton BL, Li S, Gutkind JS, Tesmer JJG. 2019. Structure of the C-terminal guanine nucleotide exchange factor module of Trio in an autoinhibited conformation reveals its oncogenic potential. *Sci Signal* **12**. doi:10.1126/scisignal.aav2449

Barber MA, Donald S, Thelen S, Anderson KE, Thelen M, Welch HCE. 2007. Membrane Translocation of P-Rex1 Is Mediated by G Protein beta Subunits and Phosphoinositide 3-Kinase. *Journal of Biological Chemistry* **282**:29967–29976. doi:10.1074/jbc.m701877200

Barber MA, Hendrickx A, Beullens M, Ceulemans H, Oxley D, Thelen S, Thelen M, Bollen M, Welch HCE. 2012. The guanine-nucleotide-exchange factor P-Rex1 is activated by protein phosphatase 1α. *The Biochemical journal* **443**:173–183. doi:10.1042/bj20112078

Cash JN, Davis EM, Tesmer JJG. 2016. Structural and Biochemical Characterization of the Catalytic Core of the Metastatic Factor P-Rex1 and Its Regulation by PtdIns(3,4,5)P 3. *Structure* **24**:730–740. doi:10.1016/j.str.2016.02.022

Cash JN, Urata S, Li S, Ravala SK, Avramova LV, Shost MD, Gutkind JS, Tesmer JJG, Cianfrocco MA. 2019. Cryo–electron microscopy structure and analysis of the P-Rex1–Gβγ signaling scaffold. *Sci Adv* **5**:eaax8855. doi:10.1126/sciadv.aax8855

Chang Y-G, Lupton CJ, Bayly-Jones C, Keen AC, D'Andrea L, Lucato CM, Steele JR, Venugopal H, Schittenhelm RB, Whisstock JC, Halls ML, Ellisdon AM. 2022. Structure of the metastatic factor P-Rex1 reveals a two-layered autoinhibitory mechanism. *Nat Struct Mol Biol* 1–7. doi:10.1038/s41594-022-00804-9

Chávez-Vargas L, Adame-García SR, Cervantes-Villagrana RD, Castillo-Kauil A, Bruystens JGH, Fukuhara S, Taylor SS, Mochizuki N, Reyes-Cruz G, Vázquez-Prado J. 2016. Protein Kinase A (PKA) Type I Interacts with P-Rex1, a Rac Guanine Nucleotide Exchange Factor. *Journal of Biological Chemistry* **291**:6182–6199. doi:10.1074/jbc.m115.712216

Chen M, Pan H, Sun L, Shi P, Zhang Y, Li L, Huang Y, Chen J, Jiang P, Fang X, Wu C, Chen Z. 2020. Structure and regulation of human epithelial cell transforming 2 protein. *Proc Natl Acad Sci* **117**:1027–1035. doi:10.1073/pnas.1913054117

Dorseuil O, Vazquez A, Lang P, Bertoglio J, Gacon G, Leca G. 1992. Inhibition of superoxide production in B lymphocytes by rac antisense oligonucleotides. *J Biol Chem* **267**:20540–20542. doi:10.1016/s0021-9258(19)36716-x

Franke D, Petoukhov MV, Konarev PV, Panjkovich A, Tuukkanen A, Mertens HDT, Kikhney AG, Hajizadeh NR, Franklin JM, Jeffries CM, Svergun DI. 2017. ATSAS 2.8: a

comprehensive data analysis suite for small-angle scattering from macromolecular solutions.
J Appl Crystallogr **50**:1212–1225. doi:10.1107/s1600576717007786

French PJ, Bunce CM, Stephens LR, Lord JM, McConnell FM, Brown G, Creba JA, Michell
RH. 1991. Changes in the levels of inositol lipids and phosphates during the differentiation of
HL60 promyelocytic cells towards neutrophils or monocytes. *Proc R Soc Lond Ser B: Biol Sci*
245:193–201. doi:10.1098/rspb.1991.0109

Gokhale NA, Zaremba A, Shears SB. 2011. Receptor-dependent compartmentalization of
PPIP5K1, a kinase with a cryptic polyphosphoinositide binding domain. *Biochem J* **434**:415–
426. doi:10.1042/bj20101437

Hayashi I, Vuori K, Liddington RC. 2002. The focal adhesion targeting (FAT) region of focal
adhesion kinase is a four-helix bundle that binds paxillin. *Nat Struct Biol* **9**:101–106.
doi:10.1038/nsb755

Hill K, Krugmann S, Andrews SR, Coadwell WJ, Finan P, Welch HCE, Hawkins PT, Stephens
LR. 2005. Regulation of P-Rex1 by phosphatidylinositol (3,4,5)-trisphosphate and
Gbetagamma subunits. *The Journal of biological chemistry* **280**:4166–4173.

Holm L, Laakso LM. 2016. Dali server update. *Nucleic Acids Res* **44**:W351–W355.
doi:10.1093/nar/gkw357

Hopkins JB, Gillilan RE, Skou S. 2017. BioXTAS RAW: improvements to a free open-source
program for small-angle X-ray scattering data reduction and analysis. *J Appl Crystallogr*
50:1545–1553. doi:10.1107/s1600576717011438

Jia Y, Subramanian KK, Erneux C, Pouillon V, Hattori H, Jo H, You J, Zhu D, Schurmans S,
Luo HR. 2007. Inositol 1,3,4,5-Tetrakisphosphate Negatively Regulates Phosphatidylinositol-
3,4,5- Trisphosphate Signaling in Neutrophils. *Immunity* **27**:453–467.
doi:10.1016/j.immuni.2007.07.016

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates
R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-
Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M,
Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW,
Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with
AlphaFold. *Nature* **596**:583–589. doi:10.1038/s41586-021-03819-2

Liebschner D, Afonine PV, Baker ML, Bunkóczi G, Chen VB, Croll TI, Hintze B, Hung L-W,
Jain S, McCoy AJ, Moriarty NW, Oeffner RD, Poon BK, Prisant MG, Read RJ, Richardson
JS, Richardson DC, Sammito MD, Sobolev OV, Stockwell DH, Terwilliger TC, Urzhumtsev
AG, Videau LL, Williams CJ, Adams PD. 2019. Macromolecular structure determination
using X-rays, neutrons and electrons: recent developments in Phenix. *Acta Crystallogr Sect D*
75:861–877. doi:10.1107/s2059798319011471

Lucato CM, Halls ML, Ooms LM, Liu H-J, Mitchell CA, Whisstock JC, Ellisdon AM. 2015. The
Phosphatidylinositol (3,4,5)-trisphosphate-dependent Rac Exchanger 1:Ras-related C3
Botulinum Toxin Substrate 1 (P-Rex1:Rac1) Complex Reveals the Basis of Rac1 Activation
in Breast Cancer Cells. *Journal of Biological Chemistry*. doi:10.1074/jbc.m115.660456

Mayeenuddin LH, McIntire WE, Garrison JC. 2006. Differential sensitivity of P-Rex1 to
isoforms of G protein betagamma dimers. *The Journal of biological chemistry* **281**:1913–
1920. doi:10.1074/jbc.m506034200

Montero JC, Seoane S, García-Alonso S, Pandiella A. 2016. Multisite phosphorylation of P-
Rex1 by protein kinase C. *Oncotarget* **7**:77937–77949. doi:10.18632/oncotarget.12846

Montero JC, Seoane S, Pandiella A. 2013. Phosphorylation of P-Rex1 at serine 1169 participates
in IGF-1R signaling in breast cancer cells. *Cellular Signalling* **25**:2281–2289.
doi:10.1016/j.cellsig.2013.07.018

Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, Morris JH, Ferrin TE.
2021. UCSF ChimeraX: Structure visualization for researchers, educators, and developers.
Protein Sci **30**:70–82. doi:10.1002/pro.3943

Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA. 2017. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**:290–296.
doi:10.1038/nmeth.4169

Punjani A, Zhang H, Fleet DJ. 2020. Non-uniform refinement: adaptive regularization improves
single-particle cryo-EM reconstruction. *Nat Methods* **17**:1214–1221. doi:10.1038/s41592-
020-00990-8

Ravala SK, Hopkins JB, Plescia CB, Allgood SR, Kane MA, Cash JN, Stahelin RV, Tesmer JJG.
2020. The first DEP domain of the RhoGEF P-Rex1 autoinhibits activity and contributes to
membrane binding. *J Biol Chem* **295**:12635–12647. doi:10.1074/jbc.ra120.014534

Stuart JA, Anderson KL, French PJ, Kirk CJ, Michell RH. 1994. The intracellular distribution of
inositol polyphosphates in HL60 promyeloid cells. *Biochem J* **303**:517–525.
doi:10.1042/bj3030517

Suloway C, Pulokas J, Fellmann D, Cheng A, Guerra F, Quispe J, Stagg S, Potter CS, Carragher
B. 2005. Automated molecular microscopy: The new Leginon system. *J Struct Biol* **151**:41–
60. doi:10.1016/j.jsb.2005.03.010

Tegunov D, Cramer P. 2019. Real-time cryo–EM data pre-processing with Warp. *Nat methods*
16:1146–1152. doi:10.1038/s41592-019-0580-y

Trewhella J, Duff AP, Durand D, Gabel F, Guss JM, Hendrickson WA, Hura GL, Jacques DA,
Kirby NM, Kwan AH, Pérez J, Pollack L, Ryan TM, Sali A, Schneidman-Duhovny D,
Schwede T, Svergun DI, Sugiyama M, Tainer JA, Vachette P, Westbrook J, Whitten AE.

2017. 2017 publication guidelines for structural modelling of small-angle scattering data from
biomolecules in solution: an update. *Acta Crystallogr Sect D* **73**:710–728.
doi:10.1107/s2059798317011597

Tria G, Mertens HDT, Kachala M, Svergun DI. 2015. Advanced ensemble modelling of flexible
macromolecules using X-ray solution scattering. *IUCrJ* **2**:207–217.
doi:10.1107/s205225251500202x

Urano D, Nakata A, Mizuno N, Tago K, Itoh H. 2008. Domain-domain interaction of P-Rex1 is
essential for the activation and inhibition by G protein betagamma subunits and PKA.
Cellular Signalling **20**:1545–1554. doi:10.1016/j.cellsig.2008.04.009

Welch HCE, Coadwell WJ, Ellson CD, Ferguson GJ, Andrews SR, Erdjument-Bromage H,
Tempst P, Hawkins PT, Stephens LR. 2002. P-Rex1, a PtdIns(3,4,5)P3- and Gbetagamma-
regulated guanine-nucleotide exchange factor for Rac. *Cell* **108**:809–821.

Figure 1. IP₄ binding causes dynamic changes in multiple domains of P-Rex1 and inhibits PIP₃-induced activation. A) Difference HDX-MS data plotted onto the domain layout of P-Rex1. Blue regions indicate less deuterium uptake upon IP₄ binding. Graphs show the exchange over time for select regions in the P-Rex1 (B) PH domain and (C) an IP4P region that was disordered in the P-Rex1–Gβγ structure. The average of two experiments is plotted with the bars representing the range of each time point. D) *In vitro* GEF activity of P-Rex1 evaluated on liposomes containing 2.5 μM PIP₃ in the presence of varying IP₄ concentrations (0-100 μM). Data were fit to exponentials to get rate constants by constraining the span to be shared. The resulting rates for each experiment were normalized by averaging two PIP₃ data points and two PC/PS data points to represent the top and bottom of the binding curve. The resulting normalized rates (min⁻¹) were fit with a one-phase binding curve wherein the top and bottom were constrained to 1 and 0, respectively, and the Hill coefficient fixed at -1. The resulting IC₅₀ was 1.4 μM with a confidence interval of 0.81 to 2.3. Data represent 4-5 independent experiments. Error bars represent the mean ± S.D.

Figure 2. Structure of the P-Rex1·IP₄ complex in an autoinhibited conformation. A) Cryo-EM reconstruction with atomic model superimposed. The kink between the DH and PH domains and the GTPase binding site are labeled. B) Atomic model without the cryo-EM map. C) The PH–4HB interface primarily involves the β1/β2 and β5/β6 loops of the PH domain, which were previously shown to be involved in protein-protein interactions in crystal structures (Cash et al., 2016), and the 4HB₁ and 4HB₂ helices of the 4HB domain. Flexible loops, including the basic β3/β4 loop of the PH domain involved in membrane binding (Cash et al., 2016), are shown as dashed lines. We speculate that this loop could interact with phosphorylated residues in the adjacent 4HB unstructured loop. D) Side chains in the PH–4HB interface. The 3-, 4-, and 5-position phosphates of bound IP₄ are labeled. Note that PIP₃ could not bind to the PH domain in this state due to steric blockade by the 4HB domain. The area of focus in (C) and (D) is circled in transparent grey in (B).

Figure 3. Mutations at the DH–DEP1 interface alter stability, conformation, and activity of DH/PH-DEP1.
A) Side chains that contribute to the hydrophobic interface formed between the DH and DEP1 domains. B) Electrostatic interactions contributing the DH–DEP1 interface. The dotted line indicates a disordered region on the DH domain containing positively charged residues that may interact with Glu456. The A170K mutant is expected to form a salt bridge with Glu411 and strengthen the interface. C) Fluorescence based *in vitro* GEF activity assay on soluble Cdc42 with variants of the purified DH/PH-DEP1 fragment. GEF activity in this experiment was fit to a one phase exponential decay normalized to that of DH/PH-DEP1 (WT). ****, P<0.0001. D) Representative Thermofluor analyses showing that mutations that disrupt the DH–DEP1 interface also destabilize the protein, as evidenced by decreased T_m values for each variant (see **Table 2**). Data are normalized from 0-100% representing lowest and highest fluorescence values. Note that A170K, which inhibits activity in panel C, increases stability. E, F) EOM analysis of SAXS data collected from mutations disrupting the DH–DEP1 interface indicate that these variants exhibit more extended conformations (see **Table 3**). EOM analyses provide the R_g and D_{max} distributions derived from selected ensembles. The gray curves correspond to the R_g and D_{max} distributions for the pool of structures used for each analysis.

Figure 4. HDX-MS and cryo-EM data support that IP$_4$ stabilizes a closed conformation of P-Rex1. A) Difference HDX-MS data plotted onto the structure of the P-Rex1 bound to IP$_4$. Blue regions indicate more protection upon IP$_4$ binding whereas red regions indicate less. See also Supplemental Data 1. B) Map representing IP$_4$ bound in the PIP$_3$-binding site of the PH domain. The 3-, 4-, and 5-phosphates of IP$_4$ are reasonably well-ordered.

Figure 5. Disruption of the DH–DEP1 and PH–4HB interfaces leads to increased P-Rex1 activity in cells. A) SRE luciferase-gene reporter assays. Mutations were cloned into full-length P-Rex1 in the pCEFL-HA-HaloTag vector, and these constructs, along with luciferase reporter genes, were co-transfected into HEK293T cells. Results depicted here are representative of three independent experiments, and error bars represent S.D. Non-transfected control (C) and empty vector transfected control (Halo) are shown. B) Mutations which led to enhanced P-Rex1 activity in luciferase reporter assays were evaluated for their effect on chemotaxis of HeLa cells with endogenous P-Rex1 knocked out (HeLa P-Rex1 KO; see Supplemental Figure 7). P-Rex1 constructs were transfected into HeLa P-Rex1 KO cells, and cell migration was evaluated in a trans-well migration assay upon stimulation with CXCL12 (50 ng/ml) or EGF (50 ng/ml). Data is presented as mean ± S.D. Significance (brackets) was determined using multiple comparison ANOVA followed by Šidák statistic test.

Figure 6. HDX-MS supports that P-Rex1 undergoes long range conformational changes when binding PIP₃-containing liposomes. A) HDX-MS of P-Rex1 in the presence of PIP₃-containing liposomes. A model of P-Rex1 in an open conformation bound to a membrane containing PIP₃ was created and is shown colored according to difference HDX-MS data plotted onto the coordinates. HDX-MS data were collected in the presence of liposomes containing PIP₃ and compared to data collected on P-Rex1 alone. Blue and red regions indicate less and more protection, respectively, upon PIP₃-containing liposome binding. These changes occur specifically in the presence of PIP₃ (see Supplemental Figure 8). The black line at the top represents a membrane surface and the dashed lines represent covalent lipid modifications. Using available structural information, Gβγ and Rac1 were docked into this model (although neither were present in this HDX-MS experiment). B) Cartoon schematic of our model of the steps involved in the activation of P-Rex1.

Supplemental Figure 1. P-Rex1 GEF activity assays show that truncated constructs are not inhibited by IP$_4$ and are not activated by PIP$_3$-containing liposomes. A and B) Fluorescence-based *in vitro* GEF activity assays with purified P-Rex1 DH/PH and DH/PH-DEP1 proteins (100 nM). Proteins were incubated with varying concentrations (0-5 μM) of IP$_4$ for 10 min at RT before adding soluble Cdc42 to measure activity. C) *In vitro* GEF activity of full-length P-Rex1, DH/PH, and DH/PH-DEP1 in the absence or presence of liposomes (± 2.5 μM PIP$_3$). The proteins were incubated with the liposomes for 20 min at RT before adding soluble Cdc42 to measure activity.

Supplemental Figure 2. Samples of P-Rex1 ± IP$_4$ imaged on a Glacios microscope highlight a conformation of P-Rex1 that is stabilized by IP$_4$. A) SDS-PAGE of a sample of purified P-Rex1 that was frozen on grids shown with a molecular weight marker. B) Representative micrographs of P-Rex1 samples. C) Particles of unliganded P-Rex1. These 2D classes represent the best particles (35,302 total) in this data set. D) Particles of P-Rex1·IP$_4$. These 2D classes represent the best particles (79,378 total) in this data set. 2D classification was performed similarly for the two datasets to select particles that showed the highest resolution features. Pixel size was 1.054 Å, and box size was 288 pixels. Classes that were obviously centered between two separate particles were excluded. Approximately twice as many micrographs were collected for the P-Rex1·IP$_4$ sample. The number of particles in each class is shown in green text in the box with each class. Orange boxes indicate particles with extra mass to the side of the Gβγ-binding core while green boxes indicate particles with extra mass opposite the Gβγ-binding site on the core.

Supplemental Figure 3. Cryo-EM 2D classification and 3D reconstruction. A) 2D class averages of all particles included in the cryo-EM reconstruction. B) Fourier shell correlation (FSC) curve for the final 3D reconstruction. C) Euler angle distribution of the final cryo-EM structure. D) Cryo-EM map of the P-Rex1·IP$_4$ complex colored according to local resolution calculated by CryoSPARC using a 0.143 FSC cutoff. E) Map-to-model FSC calculated between the atomic model and 3D reconstruction using Phenix showing that the refined model represents the 3D reconstruction.

Supplemental Figure 4. Introduction of a disulfide bridge in the DH/PH hinge reduces DH/PH-DEP1 activity. A) Atomic structure of the hinge between the DH and PH domains. Side chains of residues mutated to cysteine are shown. B) GEF activity of DH/PH-DEP1 Cys variants in the background of the A170K mutation. The rate of nucleotide exchange on soluble Cdc42 was normalized to that of the A170K DH/PH-DEP1 variant. Residues were mutated and activities measured under reducing (+DTT) and non-reducing (-DTT) conditions. Similar results were obtained when using $FeCN_6^-$ as an alternative oxidation strategy (data not shown). Data are from at least three independent experiments with error bars representing the mean ± S.D. ****, P<0.0001.

Supplemental Figure 5. The overall structure of the P-Rex1·IP₄ complex is similar to that of unliganded P-Rex1. Alignment of the P-Rex1·IP₄ complex with the previously determined structure of unliganded P-Rex1 (grey, PDB entry 7SYF). Here, the αH helix of the IP4P domain (residues 1482-1514) in each structure was aligned.

Supplemental Figure 6. SAXS analyses of DH/PH-DEP1 variants. (A) SAXS data (points) and EOM fit (line) for WT DH/PH-DEP1 and variants with mutations at the DH–DEP1 interface. A170K, L177E, and I409A are the variants. Normalized fit residuals are shown in the bottom panel. (B) Guinier analysis with residuals of fit (below). (C) Dimensionless Kratky plots suggest that all the samples have significant degrees of heterogeneity that will not permit rigid body fits. (D) Pair distance distribution (P(r)) functions, normalized by I(0). All curves were nongaussian, indicating the presence of some flexibility or extended conformations.

Supplemental Figure 7. P-Rex1 is required for CXCL12-stimulated cell migration downstream of the CXCR4 chemokine receptor. A) Crystal violet staining of trans-well migration inserts showing HeLa cell migration. Chemotaxis was evaluated under conditions without stimulation and after stimulation with CXCL12 (50 ng/mL) or EGF (50 ng/mL) and compared to cells transfected with a WT P-Rex1 construct. B) Quantification of (A), representing results from at least three independent experiments. Error bars show S.D. Significance was determined using Šidák statistic test. C) Western blot analysis of P-Rex1 protein levels in HeLa P-Rex1 KO cells and with wild-type P-Rex1 transfected back in. D) Western blot analysis of P-Rex1 wild-type and variant expression levels in HeLa P-Rex1 KO cells. GAPDH expression level was used as an internal control. P-Rex1 and GAPDH were detected using anti-P-Rex1 and anti-GAPDH antibodies, respectively.

Supplemental Figure 8. Regions across the length of P-Rex1 become more exposed upon binding liposomes specifically containing PIP₃. A) HDX-MS data collected in the presence of liposomes without PIP$_3$ and compared to data collected on P-Rex1 alone. Blue regions indicate less dynamic behavior or protection upon liposome binding while red regions indicate more. B) HDX-MS data was collected in the presence of liposomes containing PIP$_3$ and compared to data collected on P-Rex1 alone. Blue regions indicate less dynamic behavior or protection upon PIP$_3$-containing liposome binding while red regions indicate more. See also Supplemental Data 2.