Structural and mechanistic insights into human choline transport

Tsai-Hsuan Weng[1], Ainara Claveras Cabezudo[2,3], Wiebke Jösting[1], Andre Bazzone[4], Sonja Welsch[5], Gonca Gursu[1], Gerhard Hummer[2,6], Di Wu[1]*, Schara Safarian[1,7,8]*

Affiliations

[1] Department of Molecular Membrane Biology, Max Planck Institute of Biophysics, D-60438 Frankfurt/Main, Germany

[2] Department of Theoretical Biophysics, Max Planck Institute of Biophysics, D-60438 Frankfurt/Main, Germany

[3] IMPRS on Cellular Biophysics, D-60438 Frankfurt/Main, Germany

[4] Nanion Technologies GmbH, Ganghoferstr. 70a, 80339, Munich, Germany

[5] Central Electron Microscopy Facility, Max Planck Institute of Biophysics, D-60438 Frankfurt am Main, Germany

[6] Institute of Biophysics, Goethe University Frankfurt, D-60438 Frankfurt/Main, Germany

[7] Department of Microbiology and Immunology, School of Biomedical Sciences, University of Otago, Dunedin, NZ-9054, New Zealand

[8] present address: Fraunhofer Institute for Translational Medicine and Pharmacology ITMP Frankfurt, D-60438 Frankfurt/Main, Germany

*Correspondence and requests for materials should be addressed to Di Wu (di.wu@biophys.mpg.de) and Schara Safarian (schara.safarian@biophys.mpg.de).

Abstract

Human feline leukaemia virus subgroup C receptor-related proteins 1 and 2 (FLVCR1 and 2) are major facilitator superfamily transporters from the solute carrier family 49. Dysregulation of these ubiquitous transporters has been linked to various haematological and neurological disorders. While both FLVCRs were initially proposed to hold a physiological function in heme transport, subsequent studies questioned this notion. Here, we used structural, computational and biochemical methods and conclude that these two FLVCRs function as human choline transporters. We present cryo-electron microscopy structures of FLVCRs in different inward- and outward-facing conformations, captured in the apo state or in complex with choline in their translocation pathways. Our findings provide insights into the molecular framework of choline coordination and transport, largely mediated by conserved cation-π interactions, and further illuminate the conformational dynamics of the transport cycle. Moreover, we identified a heme binding site on the protein surface of the FLVCR2 N-domain, and observed that heme actively drives the conformational transitions of the protein. This auxiliary binding site might indicate a potential regulatory role of heme in the FLVCR2 transport mechanisms. Our work

40   resolves the contested substrate specificity of the FLVCRs, and sheds light on the process of maintaining

41   cellular choline homeostasis at the molecular level.

Introduction

The feline leukaemia virus subgroup C receptor (FLVCR) family, a member of the major facilitator superfamily (MFS) of secondary active transporters, consists of four paralogues encoded by the human *SLC49* gene group[1]. FLVCR1 (SLC49A1) was initially identified as the cell receptor for feline leukaemia virus (FeLV), a retrovirus that causes profound anaemia in cats by interfering with erythropoiesis[2]. FLVCR2 (SLC49A2), another major member of the family, shares 60% sequence identity with FLVCR1 in the transmembrane domain but does not bind to the feline leukaemia virus subgroup C envelope protein[3]. Both transporters exhibit ubiquitous tissue distribution in humans and have significant haemato- and neuropathological implications[1,4]. Dysfunction of FLVCR1 caused by germline mutations is associated with posterior column ataxia with retinitis pigmentosa (PCARP)[5], while its alternatively spliced isoforms have been linked to Diamond-Blackfan anaemia (DBA)[6]. Similarly, truncation and missense mutations in the gene encoding for FLVCR2 are associated with the autosomal-recessive cerebral proliferative vasculopathy known as Fowler syndrome[7,8]. Furthermore, both FLVCR variants are suggested to play a key role in cell development and differentiation, including angiogenesis and tumorigenesis[9–12].

Here, we studied the molecular architecture, the conformational landscape, and the contested substrate specificity of human FLVCR1 and FLVCR2 using single-particle analysis cryogenic electron microscopy (SPA cryo-EM) and molecular dynamics (MD) simulations. Our findings indicate that FLVCR1 and FLVCR2 both represent previously uncharacterized, ubiquitous human choline transporters.

Results

Overall architecture of FLVCR1 and FLVCR2

We stably integrated the human *SLC49A1* and *SLC49A2* genes into human embryonic kidney cells (HEK293), and recombinantly produced wildtype FLVCR1a (hereafter referred to as FLVCR1) and FLVCR2 variants with C-terminal flag-tag modifications, respectively. Affinity-purified samples of each transporter were subjected to SPA cryo-EM (Supplementary Fig. 1).

We determined the respective structures of apo FLVCR1 and FLVCR2 at 2.9 Å resolution, captured in inward-facing conformations (Fig. 1a,b and Extended Data Fig. 1 and 2). Both FLVCR paralogs share a common MFS-type architecture composed of 12 transmembrane α-helices (TM) arranged into two pseudo-symmetrical domains[13] (Fig. 1c). In both inward-facing structures, the N-domain (TM1-6) and the C-domain (TM7-12) are connected by a long and flexible loop containing two short horizontal helices (H1 and H2) on the inner side of the membrane (Fig. 1). On the inner side of FLVCR2, we resolved a short helical segment at the C-terminus (H3), while the density of the FLVCR1 C-terminus was less pronounced (Extended Data Fig. 3). Further, we identified an N-linked glycosylation site at N265 of FLVCR1 which locates within the external loop connecting TMs 5 and 6 (EL5-6; Fig. 1a). This glycosylation pattern is absent in the analogous site of the FLVCR2 structure.

The inward-facing conformations of FLVCR1 and FLVCR2 exhibit a close resemblance to each other, with a root mean square deviation (Cα r.m.s.d.) of 0.993 Å. Both structures feature a wedge-shaped solvent-accessible cavity that is mainly created by the separation of TMs 4 and 5 of the N-domain from TMs 10 and 11 of the C-domain (Extended Data Fig. 4a). This space extends halfway across the membrane with a similar depth of around 23 Å in both (Fig. 2c and Extended Data Fig. 4b). On the level of the outer leaflet, TMs 1, 2, and 5 of the N-domain and TMs 7, 8, and 11 of the C-domain pack tightly against each other and thus shield the central cavity from the external space (Extended Data Fig. 4a). Notably, TMs 1, 4, and 7 exhibit disordered regions at their C-terminal ends, followed by short kinked helical motifs designated TMs 1b, 4b, and 7b (Fig. 1c and Extended Data Fig. 3).

Conformational difference between inward- and outward-facing structures of FLVCR2

From the as-isolated FLVCR2 samples we were able to determine not only the inward-facing

conformations but also the outward-facing conformation from class averages of even particle

distribution (Fig. 2a and Extended Data Fig. 2a). The outward-facing conformation of FLVCR2 is

characterized by a cavity that is accessible from the external space created by a rocker-switch rigid-

body motion of the two domains (Fig. 2b,c and Supplementary Video 1). This motion shifts the outer

halves of all TMs away from the central axis, while the inner halves adopt an inward movement to

obstruct the exit (Fig. 2b). The cavities in outward- and inward-facing conformations are 7.4 Å and 8.6

Å wide at their respective openings (Fig. 2c). The wedge-like volume of both cavities is surrounded by

an uncharged and hydrophobic protein environment, lined by mostly conserved residues (Extended

Data Fig. 5). Compared to the inward-facing cavity, the outward-facing cavity demonstrates a more

restricted pathway within its central region. In this narrowed segment of the cavity, the channel ends

at W102 and Y325, two residues that are highly conserved in both FLVCR transporters (Fig. 2c and

Extended Data Fig. 4b and 5).

In line with an alternating-access model, the inward-facing conformation of FLVCR2 features a tightly-

sealed external gate. This is mainly achieved by the juxtaposition of TM1b (N-domain) and TM7b (C-

domain; Fig. 2c,d). The inter-domain interaction between these two motifs is stabilized by a hydrogen

bonding network consisting of two pseudo-symmetry-related asparagine residues (N110-N332), as well

as E343 (TM8) and N239 (EL5-6; Supplementary Fig. 2). Furthermore, we identified a stable inter-

domain salt bridge between D124 (TM2) and R333 (TM7b), presumably reinforcing the external

occlusion (Fig. 2c,d, and Supplementary Fig. 2). FLVCR1 appears to employ a similar gating mechanism

on its external side, given that all the residues involved in the hydrogen bonding network of FLVCR2 are

found to be highly conserved between these two paralogs (Extended Data Fig. 4b,c). However, the

D124-R333 salt bridge in FLVCR2 is not conserved in FLVCR1. While the location of the D147 residue in

FLVCR1 aligns with that of D124 in FLVCR2, FLVCR1 possesses a glutamine taking the position of the

arginine (Extended Data Fig. 4c).

During the transition from inward- to outward-facing conformation of FLVCR2, interactions contributing to the external gate become disrupted, while the formation of the internal gate occludes the central cavity from the internal side. TM4b of the N-domain moves in proximity to the N-terminal end of TM11 of the C-domain to establish a first level of occlusion (Fig. 2b,c). An interaction network consisting of several hydrogen bonds and a salt bridge is found within this region (Fig. 2e). E435 (TM11) plays a central and versatile role by stably forming a hydrogen bond with S203 (TM4b) and a concurrent salt bridge with R200 (TM4b; Fig. 2e and Supplementary Fig. 2). An additional inter-domain hydrogen bonding site is identified between S199 (TM4b) and S439 (TM11; Fig. 2e and Supplementary Fig. 2). In the peripheral region, K372 and R374 (EL8-9) approach N209 (EL4-5) and S212 (TM5) to form hydrogen bond pairs and thus block the lateral accessibility of the cavity (Fig. 2e and Supplementary Fig. 2). A second level of occlusion was observed beneath the internal ends of TMs 4, 10 and 11, where H1 and H3 are positioned in close proximity (Fig. 2c,f). Here, the backbone carbonyl group and amide nitrogen of A283 (H1) form stable hydrogen bonds with N497 (H3) and Y431 (IL10-11), respectively (Fig. 2f and Supplementary Fig. 2). Together with the loop connecting TMs 10 and 11, the two helical motifs H1 and H3 serve as a latch to secure the closure of the two domains on the internal side.

 Substrate-specificity and auxiliary binding sites

The transport substrates of both FLVCR1 and FLVCR2 have been contested for decades[14–17]. Both

FLVCR1 and FLVCR2 were previously proposed to hold a function in heme transport. With the goal of

assessing their heme-binding capabilities and localizing the putative heme-binding site, we performed

cryo-EM experiments using heme as a potential substrate.

A central observation upon determining the structure of heme-supplemented FLVCR2 was that the

presence of heme resolved the conformational heterogeneity, fully driving the transporter into the

outward-facing conformation (Fig. 3a and Extended Data Fig. 2b). Surprisingly, no density features of

heme were noticeable within the central cavity. Instead, we observed a density corresponding to heme

in the vicinity of the N-terminus of TM1 on the internal side, where it replaces the density we assigned

as a lipid molecule in the apo outward-facing structure (Fig. 3a,b). The protein surface in this particular

region displays a patch of positive charges, suggesting the heme molecule to be held in place by the

formation of electrostatic interactions between its two propionate groups and the side chains of R81,

R82, and K273 (Fig. 3b and Extended Data Fig. 5d). The map density of bound heme in our structure

indicates local mobility in the heme position due to the absence of macrocycle rigidifying axial

coordination (Fig. 3b). By contrast, cryo-EM studies on FLVCR1 in the presence of heme did not reveal

any heme binding or notable conformational changes. It is to note that neither the cavity dimensions

nor the residues lining the central cavities of FLVCR1 and FLVCR2 suggest the existence of a heme-

binding pocket.

To further characterize the molecular nature of heme binding in FLVCR2, we carried out atomistic MD

simulations, focusing on FLVCR2 in a lipid bilayer composed of 75% palmitoyloleoyl

phosphatidylethanolamine (POPE) and 25% palmitoyloleoyl phosphatidylglycerol (POPG). To probe

possible binding into the transport pathway, we first placed a heme close to the cavity opening in the

outward-facing conformation. In the subsequent MD simulations, the heme immediately crossed the

solvent layer and periodic boundary to interact with the loop region on the internal side for the

remainder of the simulation (Supplementary Video 2). We then positioned heme on the inner side of

the bilayer near the binding site observed by cryo-EM at the N-terminus of TM1 (Fig. 2c). In all replicas,

the contact between heme and the R82 residue was lost early in the simulations as the heme

reoriented and was absorbed into the lipid bilayer. However, the contact between one of the

propionate groups and R83 was consistently maintained. Upon membrane insertion, the heme formed

additional contacts with other residues (Supplementary Fig. 3). Stable interactions were observed

between the methyl group of heme and I270 (TM6), as well as one of the propionate groups and K273

(TM6). Our MD simulation data, together with our structural findings, point to the existence of a heme-

binding site at the N-domain of FLVCR2, located near the inner surface of the membrane. This finding

is in line with previously reported results demonstrating heme-binding at the N-terminal segment of

FLVCR2[18].

Subsequently, we focused our attention on choline as a potential substrate of FLVCR1 and FLVCR2 based

on the characteristic tryptophan-tyrosine clusters observed at the presumed substrate-binding cavities

of FLVCR1 and FLVCR2, respectively. This chemical environment is found analogously in the prokaryotic

choline transporter LicB, as well as predicted models of eukaryotic choline transporters[19–21] (Extended

Data Fig. 6). Indeed, we obtained cryo-EM maps of choline-bound structures of FLVCR1 and FLVCR2 at

a resolution of 2.6 Å and 2.8 Å, respectively (Fig. 3d,e and Extended Data Fig. 2b and 3c). Both structures

were captured in the inward-facing conformation, suggesting that choline is also capable of resolving

the conformational heterogeneity in FLVCR2 by trapping it in an inward-facing substrate-bound state.

Intriguingly, this effect is opposite to that of heme described above. The choline-bound structures

exhibit a Cα r.m.s.d. of 0.688 Å (FLVCR1) and 1.002 Å (FLVCR2) to their respective apo inward-facing

structures. The choline-bound inward-facing structures of FLVCR1 and FLVCR2 resemble each other and

their substrate binding sites are found in analogous locations within the transporter scaffold. The

bound choline molecule is caged inside the central binding site between the two domains of the FLVCRs,

surrounded mainly by TMs 1, 2, 4, 5, 7, and 11 (Fig. 3d,e). The binding sites in the two transporters are

formed by a suite of conserved residues. The central coordinating tryptophan residue (W102[FLVCR1] and

W125[FLVCR2]) of TM1 is located above the choline, constraining the diffusion of the molecule towards the

external gate. Two additional aromatic residues of TM7, one tyrosine (Y325 [FLVCR1] and Y349[FLVCR2]) and

one phenylalanine (F348[FLVCR1] and F324[FLVCR2]), line the peripheral space of the binding site and restrict

the movement of the choline within the pocket. Our MD simulations demonstrate that the quaternary

ammonium group of choline stably interacts with the conserved tryptophan residue (W125[FLVCR1] and

W102[FLVCR2]), forming concomitant cation-π interactions with both the tryptophan and tyrosine residues

(Fig. 3f,g). The hydroxyl group of the choline molecule showed more versatile interactions. Two

asparagine residues identified near the hydroxyl group formed transient hydrogen bonds with choline

during our simulations. In addition, at least two interacting water molecules were consistently present

near the OH-group of the bound choline, which was predominantly oriented away from the conserved

tryptophan towards the solvent phase (Supplementary Fig. 4 and Supplementary Video 3). It is

noteworthy that all these coordinating residues are fully conserved in mammals, suggesting a

specialized role of FLVCRs in recognizing and transporting choline across species (Fig. 3h).

To gain further insights into the spectrum of substrate selectivity of FLVCR1 and FLVCR2, we performed

nano differential scanning fluorimetry (nanoDSF) using choline, phosphocholine, and betaine as

candidate substrates (Fig. 3i and Extended Data Fig. 7). For both transporters, strong quenching of

tryptophan fluorescence could be observed by the addition of choline, but not its structurally related

metabolites (Fig. 3i). We determined choline dissociation rate constants for FLVCR1 at 25.10 ± 1.26 μM

and FLVCR2 at 125.22 ± 4.87 μM[22]. Subsequently, we attempted to assess the kinetic properties of both

FLVCRs via a solid-supported membrane-based electrophysiological (SSME) approach (Fig. 3j). For both

transporters we observed a rapid increase in current upon application of choline, followed by a fast

current decay, which can be attributed to an electrogenic substrate binding induced pre-steady state

(PSS) event. However, only for FLVCR2, a second current phase was observed, revealing a much slower

decay time constant. This current phase is typical for transport events that slowly charge the membrane

capacitor in SSME experiments[23,24]. Despite the structural similarities described above, FLVCRs seem to

have different kinetic fingerprints that might either originate from subtle structural differences

identified at the termini of the two transporters, or specific requirements to their surrounding

environment. Further, our data indicate that FLVCR-facilitated choline transport displays no immediate

pH dependency within the physiological range tested in our assay (Extended Data Fig. 8). Thus, we

conclude that FLVCRs most likely represent choline-driven uniporters. We then determined choline

binding affinities based on the PSS peaks and obtained values of 0.52 ± 0.07 mM for FLVCR1 and 3.28

± 0.31 mM for FLVCR2. While the absolute affinities deduced from the SSME experiments are weaker

than those from the more direct DSF measurements, which may reflect an underlying transport phase,

the relative values consistently demonstrate a similar 5- to 6-fold difference in choline affinity between

the two transporters[25]. We note that our DSF-derived binding affinities more likely reflect values that

are realistic in a biological setting, as plasma choline concentrations are relatively stable at 5-10 μM[26].

Translocation pathway of choline in FLVCRs

MFS transporters typically cycle between inward- and outward-facing conformations, facilitating

substrate translocation in an alternating-access manner. Substrate binding often plays a pivotal role in

eliciting the conformational transitions[27]. Our cryo-EM data support this mechanism for at least FLVCR2,

where we see a full transition from the outward-facing to inward-facing state upon choline binding.

Our MD simulations map the route for substrate entry along the pathway in the outward-facing

conformation of FLVCR2. After spontaneously diffusing into the opening, choline initially interacts with

several residues near the protein surface, mainly D124 (Supplementary Fig. 5). It sequentially

approaches the deeper recesses of the cavity, primarily engaging with conserved aromatic residues

W102 and Y325. In a particular entry event, we observed choline moving to a position below the W102

residue within the binding site, consistent with our observations in the choline-bound structure of

FLVCR2 (Supplementary Video 4).

The substantial global conformational changes triggered by choline further alter the local arrangement

of substrate-coordinating residues within the translocation pathway. The rearrangement of the binding

site results in a more constricted pocket, promoted by the inward movement of the conserved residues

closer to the choline molecule (Extended Data Fig. 9a). The repositioning of these residues, especially

of the aromatic side chains, restricts the accessibility of the binding pocket and promotes choline

capture and engulfment.

The choline-bound inward-facing conformations likely represent the pre-release phase during the

translocation. In their choline-bound structures, the cavities of both FLVCRs share common

characteristics, exhibiting a neutral interior but a negatively charged surface at the exit (Extended Data

Fig. 5b,e). FLVCR1 features a slightly smaller cavity volume of 513 Å^3 compared to 579 Å^3 of FLVCR2

(Extended Data Fig. 9b). In an inter-domain interaction site at the cavity opening, a serine residue of

TM4 (S222[FLVCR1] and S199[FLVCR2]) and a glutamate side chain of TM11 (E459[FLVCR1] and E435[FLVCR2]) remain

in close proximity even with the internal gate open (Extended Data Fig. 9c,d). While these conserved

residues may interact to limit pathway accessibility, a peripheral solvent-accessible channel emerging

from the internal space to the binding site reveals a semi-open translocation pathway in the choline-

bound inward-facing conformations of both transporters (Extended Data Fig. 9b).

In our MD simulations, we captured a spontaneous choline release event from FLVCR2 through the

semi-open pathway that would complete the translocation process (Supplementary Video 5). The

choline spontaneously left the binding site after ~700 ns, transited in steps through the half channel,

and then escaped into the solvent at ~900 ns. Our apo inward-facing structures plausibly illustrate the

following post-release states. We speculate that an event occurs during the substrate release process

in both FLVCRs, resulting in the dissipation of the conserved serine-glutamate interactions and causing

the full opening of the internal cavity. This is suggested by our observations that the distance between

the S-E residues expands in the apo inward-facing structures relative to the choline-bound states

(Δd_{FLVCR1} = 3.3 Å and Δd_{FLVCR2} = 5.2 Å; Fig. 2e, Extended Data Fig. 4d and 9c,d). A subtle conformational

transition takes place upon the release of the choline in both FLVCRs, manifested by a rigid-body shift

between two domains that moves the inner halves slightly away from the central axis perpendicular to

the membrane plane (Extended Data Fig. 9e,f). The rearrangement of the TMs results an increased

angle between the N-domain and C-domain, widening by 0.9° in FLVCR1 compared to 4.3° in FLVCR2.

Consequently, the central cavities become less compact, enlarging from volumes of 513 Å^3 and 579 Å^3

for FLVCR1 and FLVCR2, to 771 Å^3 and 1046 Å^3, respectively. Meanwhile, the molecular architecture of

the external gate remains largely unchanged in the post-release inward-facing states of both FLVCRs

(Fig. 2d, Extended Data Fig. 4c and 9c).

Choline binding and release events drive additional local rearrangements in the translocation pathway

of FLVCR2. The inward-facing apo structure of FLVCR2 reveals a distinct lateral opening of the pathway,

directed towards the inner membrane leaflet of the lipid bilayer (Extended Data Fig. 10). This channel,

located between TM5 (N-domain) and TM8 (C-domain), accommodates a lipid/detergent-like density

in the apo state, as observed in our cryo-EM results (Extended Data Fig. 10b). Yet, when choline is

present in the central cavity, residue F220 seals the opening and prevents the lipid/detergent-like

density from accessing the channel (Extended Data Fig. 10c). The relevance of this site for

environmental interactions was further supported during our simulations with FLVCR2, where lipid-

entering events were observed (Extended Data Fig. 10a). The displacement of F220 allowed one of the

lipid's acyl tails to enter and reside within this channel. Whether or not this might be connected to a

putative allosteric regulatory mechanism linked to lipid binding in FLVCR2 as observed in other MFS-

type transporters remains to be further investigated[28,29].

Discussion

FLVCR transporters were considered to be human heme transporters. In this study, we have determined cryo-EM structures of FLVCR1 and FLVCR2 in distinct apo, heme and choline-bound states, providing valuable insights into architecture, substrate interaction and conformational dynamics of these two MFS transporters. Our data support previous studies regarding the heme-binding properties of FLVCR2, yet clearly demonstrate that choline is the transport substrate of both FLVCRs. Based on our structural findings, we suggest a rocker-switch alternating-access mechanism for the transport cycle of choline import (Fig. 4). As the resting state, the apo outward-facing FLVCR2 conformation is set up for choline binding from the external space. Substrate-induced conformational changes drive FLVCR2 towards its inward-facing state. Finally, the choline will be released to the cytosol or the mitochondrial matrix, depending on the sub-cellular localization of FLVCR1 and FLVCR2[18]. Subsequent to choline release, FLVCRs will transition into their outward-facing state in order to re-initiate the transport cycle. Our structural data did not reveal fully occluded conformations that are key features of alternating access. Hence, we suspect that these occluded conformations exist only transiently and rapidly convert towards either the inward- or outward-facing conformations.

Our findings suggest that FLVCRs act as uniporters, conducting facilitated diffusion across the membrane. Uniporters conventionally utilize the chemical potential of substrates as the only energy source driving the transport process thermodynamically[30]. This mechanism appears to match the kinetic fingerprints of FLVCRs observed in the SSM-based assays. Our cryo-EM data also reveal that choline-binding prompts the conformational transition in FLVCR2 from the outward-facing to the inward-facing state without requiring additional energy inputs. Corroborating this finding, our MD simulation of FLVCR2 captured several spontaneous entry events of choline into the cavity in the outward-facing conformation, including a traverse below the central coordinating tryptophan residue, reminiscent of the local binding environment in the inward-facing choline-bound state (Supplementary Fig. 5 and Supplementary Video 4). Furthermore, we observed a release event of choline from the inward-facing conformation of FLVCR2 (Supplementary Video 5).

While choline is the primary transport substrate for FLVCR2, it should be highlighted that heme can

also induce conformational changes, albeit leading to the opposing outward-facing state. This concurs

with the previous reports that FLVCR2 is capable of binding heme with the N-domain yet not

transporting it[16,18,31]. As labile cytoplasmic heme is readily degraded by heme oxygenase, we anticipate

that the heme interaction of FLVCR2 will be restricted to mitochondria where heme biogenesis takes

place. We performed matrix targeting sequences (MTSs) predictions to identify the targeting sequence

of FLVCR2[32]. The N-terminal region of FLVCR2 exhibits a propensity score comparable to that of proteins

known to be trafficked to the mitochondria (Supplementary Fig. 6). Correspondingly, the endogenous

FLVCR2 protein, originating from its chromosomal locus, has been previously reported to localize within

mitochondria[18]. We conclude that the physiological role of FLVCR2 is most likely to act as a

mitochondrial choline importer that may be regulated by the availability of heme within the

mitochondrial matrix. In addition, our MD simulations revealed that a physiological mitochondrial

membrane potential influences the rates of choline entry and release in FLVCR2, favoring inward-

directed translocation of this positively charged substrate (Supplementary Fig. 7 and Supplementary

Table 1). Choline import in rat kidney mitochondria was previously found to depend on the membrane

potential, which matches with our observations[33]. In contrast to FLVCR2, FLVCR1 showed no binding or

response to heme, and obtained a very low propensity score in the MTSs prediction (Supplementary

Fig. 6). Combined with insights from genome-wide association study data, we presume that FLVCR1

acts as a plasma membrane-localized choline transporter[34].

The successful elucidations of FLVCR structures in their choline-bound state establish the SLC49 family

as a new class of human choline transporters[20]. Despite the positive charge of choline, the fully

conserved binding site in FLVCRs exhibits a neutral charge environment. As a result, the coordination

of bound choline is predominantly mediated by the cation-π interaction formed between the

trimethylammonium group of choline and conserved aromatic residues. In a similar manner, the

bacterial choline transporter LicB features a binding site composed of six aromatic and one amide side

chains, including W17 of TM1 that corresponds to the critical coordinating tryptophan residues in FLVCRs[19] (Extended Data Fig. 6a). We performed a comparative analysis using the AlphaFold2 structures of other putative human choline transporters (Extended Data Fig. 6b-d). Our examination of these structures reveals binding pockets containing aromatic residues. In the high-affinity choline transporter SLC5A7, the putative binding pocket encompasses four tryptophan and two tyrosine residues, potentially forming a cation-π trap. The CTL (SLC44) family members feature a unique fold and the exact position of their binding site remains unclear due to the absence of high-resolution structures. However, the putative pocket of human SLC44A1 suggests the involvement of a tyrosine residue in choline coordination[35]. The SLC22 family members share the same MFS fold with FLVCRs, and their putative binding site with a broad substrate spectrum also contains aromatic and amide side chains that are potential participants in choline coordination[36]. Our structural findings and analyses suggest that cation-π interactions may be a common principle for choline coordination. While the cation-π interaction appears to be highly specific and selective for substrate coordination in FLVCRs, our data suggest that the hydroxyl group of choline preferentially forms promiscuous and dynamic interactions either with the surrounding protein environment, such as amide-containing residues, or with water molecules.

Choline, an indispensable quaternary ammonium cation with water-soluble vitamin-like characteristics, is vital for a myriad of biological processes[37]. Dietary choline uptake mediated by certain transporters is necessary as our bodies have limited capacity for *de novo* synthesis[38]. Choline deficiency is linked to a plethora of undesired health consequences, for instance, skeletal muscle atrophy and neurodegenerative diseases[39]. Additionally, abnormal choline metabolism is emerging as a hallmark for oncogenesis and pathological inflammatory conditions[20,40,41]. Disorders including PCARP and DBA associated with FLVCR1 and the Fowler syndrome related to FLVCR2 might stem from choline deficiency that leads to developmental or degenerative complications. In conjunction with the known physiological importance of choline and our current findings, a link between defective FLVCR function and the associated pathologies becomes evident. Choline stands as the precursor for essential

compounds, such as the membrane phospholipid phosphatidylcholine (PC), the neurotransmitter

acetylcholine, and the methyl group donor betaine[39]. Altered choline metabolism is apt to cause

abnormal cell proliferation and differentiation conditions, impacting especially the nervous and

hematologic systems due to disturbances in neurotransmitter synthesis and methylation activities[42,43].

Beyond these impacts, choline metabolism in macrophages also affects mitochondrial morphology and

metabolism[41]. The disruption of PC synthesis results in the destruction of mitochondrial cristae,

compelling cells to rely only on substrate-level phosphorylation for ATP production. A handful of animal

studies have further underscored the interplay between choline supplementation and brain

development, particularly in angiogenesis[44–47]. Choline deficiency has been linked to decreased DNA

methylation within the promoters of genes that are important regulators of angiogenesis[44]. Moreover,

dysfunctional mutations or complete loss of choline transporters SLC5A7 or SLC44A1 have been

implicated in certain neurodegenerative conditions, including but not limited to distal hereditary motor

neuronopathy, congenital myasthenic syndrome, and childhood-onset neurodegeneration[48–52]. The

precise molecular mechanisms linking choline transport to these disorders are not yet fully deciphered

and require further study given their clinical implications.

Online Methods

Generation of Inducible HEK293 Stable Cell Lines.

The complementary DNAs of full-length wildtype FLVCR1 (human SLC49A1, NCBI Reference Sequence

NM_014053) and FLVCR2 (human SLC49A2, NCBI Reference Sequence: NM_017791) were cloned into

pcDNA5/FRT/TO (Invitrogen) vectors, respectively. The gene for both FLVCRs was modified by a C-

terminal FLAG fusion tag. Further details are found in sequence data provided as supplementary

information (Supplementary Tables 2 and 3). The recombinant Flp-In T-REx293-FLVCR1 and Flp-In T-

REx293-FLVCR2 cell lines were generated by using a tetracycline-inducible and commercially available

Flp-In T-REx host-cell line system from Invitrogen. Flp-In T-REx293 cells were cultured in high-glucose

Dulbecco's Modified Eagle's Medium (DMEM; Sigma-Aldrich) supplemented with 10% fetal bovine

serum (FBS; Gibco), 1% Pen/Strep (Gibco), 1 μg ml^{-1} Zeocin (Thermo Fisher), and 15 μg ml^{-1} blasticidin

S hydrochloride (AppliChem) at 37 °C in an atmosphere of 5% CO_2. Cells were not tested for

mycoplasma contamination. For stable integration, the pcDNA5/FRT-FLVCR1-FLAG and pcDNA5/FRT-

FLVCR2-FLAG vectors were cotransfected with the Flp recombinase encoding expression vector pOG44

(Thermo Fisher) at a 1:13 mass ratio, respectively. All transfection procedures were performed with

Lipofectamine™ 2000 reagent according to the manufacturer's instructions (Thermo Fisher). To select

stable clones, transfected cells were cultivated with growth medium containing 100 μg ml^{-1} hygromycin

B (AppliChem).

Production and Purification of the human FLVCR1 and FLVCR2.

For protein production, the Flp-In T-REx293-FLVCR1 and Flp-In T-REx293-FLVCR2 cell lines were cultured

in roller bottles (Greiner Bio-One) in growth media containing 100 μg ml^{-1} hygromycin B for 14 d under

the above-mentioned conditions. Gene expression was induced at 100% confluence by adding a final

concentration of 2 μg ml^{-1} doxycycline hydrochloride. After 72 h, cells were harvested with Accutase

solution (Sigma-Aldrich) and stored at −80 °C until further use. Harvested cells were suspended in cold

lysis buffer containing 25 mM Tris pH 7.4, 150 mM NaCl, and 0.1 g ml^{-1} SigmaFast

ethylenediaminetetraacetic acid (EDTA)-free protease inhibitor (Sigma-Aldrich) and disrupted by

stirring under high-pressure nitrogen atmosphere (750 MPa) for 45 min at 4 °C in a cell-disruption

vessel (Parr Instrument). The cell lysate was centrifuged at 8,000g at 4 °C for 15 min. Subsequently, the

low-velocity supernatant was centrifuged at 220,000g at 4 °C for 60 min. Pelleted membranes were

resuspended and stored in a storage buffer containing 25 mM Tris pH 7.4, 150 mM NaCl, 10% glycerol

(v/v), and 0.1 g ml^{-1} SigmaFast EDTA-free protease inhibitor (Sigma-Aldrich).

All purification steps of both FLVCRs were performed at 4 °C. Isolated membranes were solubilized with

1% (w/v) lauryl maltose neopentyl glycol (LMNG; GLYCON Biochemicals) with gentle stirring for 1 h.

The insoluble membrane fraction was removed via ultracentrifugation at 220,000g for 1 h.

Subsequently, the supernatant was incubated with ANTI-FLAG® M2 Affinity Gel resin (Merck) for 1 h.

The resin was preequilibrated with a buffer containing 50 mM Tris pH 7.4, 150 mM NaCl, and 0.02%

LMNG (w/v). The washing step was performed using 20 column volumes (CVs) of wash buffer [50 mM

Tris pH 7.4, 150 mM NaCl, 5% (v/v) glycerol, and 0.02% LMNG). The protein was eluted from the M2

resin with 10 CVs of the same buffer supplemented with 4 mM FLAG® Peptide (Merck). The eluted

sample was concentrated and subjected to a Superdex 200 Increase 10/300 column (GE Healthcare)

equilibrated with size exclusion chromatography (SEC) buffer [50 mM Tris pH 7.4, 150 mM NaCl, and

0.001% (w/v) LMNG]. Peak fractions were pooled, concentrated to 1.5 mg ml^{-1} using an Amicon 50-kDa

cut-off concentrator (Merck Millipore), and stored for further analysis.

For the dodecyl maltoside (DDM) and cholesterol hemisuccinate (CHS) solubilized samples used for

proteoliposome reconstitution, the proteins were purified based on the same protocol. The membrane

solubilization buffer contains 1% (w/v) DDM/0.2% (w/v) CHS instead of LMNG. Similarly, the wash

buffer contains 0.1% (w/v) DDM/0.02% (w/v) CHS, and the SEC buffer contains 0.02% (w/v) DDM/0.004%

(w/v) CHS.

Immunoblotting

Affinity-purified proteins were subjected to SDS-PAGE and immunoblotting. FLAG-tagged FLVCR1 and

FLVCR2 were detected using anti-FLAG (F3165, Sigma-Aldrich) antibodies at 1:1,000 dilution. Anti-

mouse IgG antibody conjugated with alkaline phosphatase (A9316, Sigma-Aldrich) was used as

secondary antibody at 1:5,000 dilution.

Nano differential scanning fluorimetry (nanoDSF)

NanoDSF measurements were carried out using Prometheus Panta (NanoTemper Technologies).

Purified protein samples were diluted with DSF buffer containing 50 mM HEPES pH 7.4, 150 mM NaCl,

and 0.001% (w/v) LMNG to 1 μM. Buffers with different concentrations of choline, phosphocholine, or

betaine were prepared by serial dilutions of DSF buffer containing 4 mM of the compounds. The protein

samples were mixed with an equal volume of DSF buffer or the compound-containing buffer with a

final protein concentration of 0.5 μM and then incubated at room temperature for 15 min. A volume

of 10 μl mixed solution was used per Prometheus high sensitivity capillary (NanoTemper Technologies).

A temperature ramp of 1 °C min^{-1} from 25 to 95 °C was applied while the intrinsic protein fluorescence

at 330 and 350 nm was recorded. Analysis of the initial ratio of F_{350}/F_{330} was carried out using Python

libraries including pandas, numpy, scipy and seaborn in Visual Studio Code (Microsoft). Three technical

replicates were recorded for data analysis.

Proteoliposome preparation

1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoethanolamine (POPE, Avanti) and 1-Palmitoyl-2-oleoyl-*sn*-

glycero-3-phospho-(1'-rac-glycerol) (POPG, Avanti) in chloroform (25mg ml^{-1}) were mixed at the ratio

of 3:1 (w/w), followed by gentle evaporation of the chloroform and overnight desiccation under a

stream of nitrogen gas. The dried lipids were suspended in internal buffer comprised of 50 mM HEPES

pH 7.4, 150 mM NaCl to a final concentration of 12.5 mg ml^{-1} and sonicated until homogenous. The

lipid solution was subjected to three freeze-thaw cycles and extruded through a 400-nm polycarbonate

filter (Avestin) to generate unilamellar vesicles. The liposomes were destabilized by the addition of 0.13%

(w/v) Triton-X100 (Sigma) at room temperature. Purified DDM/CHS-solubilized FLVCR1 or FLVCR2

samples was added at a 1:50 protein-to-lipid mass ratio and incubated for 15 min at 4 °C. Protein-free

liposomes were prepared in parallel using the same concentration of DDM/CHS. The detergent was

removed by several additions of SM2-BioBeads (Bio-Rad). Firstly, 0.05 g ml^{-1} of BioBeads were added

to the sample and incubated with gentle agitation for 1 h at 4 °C, then 0.05 g ml^{-1} BioBeads were added

followed by incubation overnight at 4 °C, and finally 0.08 g ml^{-1} BioBeads were added with 2-h

incubation at 4 °C. BioBeads were removed by filtration and liposomes were collected by

ultracentrifugation at 264,000g for 20 min and resuspended at a concentration of 8 mg ml^{-1} lipids in

internal buffer. The reconstituted liposomes were flash frozen in liquid nitrogen and stored at −80 °C.

SSM-based electrophysiology (SSME) assays

SSM-based electrophysiology was performed on a SURFE^{2}R N1[53] (Nanion Technologies). For all

experiments, preparation of 3-mm sensors was performed as described[54]. For measurements, a single

solution exchange configuration was employed that consisted of three phases of 0.5 s duration each:

flow of non-activating solution (NA), activating solution (A), and non-activating solution (NA). Both NA

and A were prepared from the same main buffer (30 mM HEPES, 30 mM MES, and 300 mM NaCl at pH

7.4), while A was supplemented with choline chloride and NA was prepared by adding NaCl in an

equimolar manner. For the choline dose response curve, 50 mM choline chloride and 50 mM NaCl were

used to achieve A0 and NA (with the highest salt concentration), followed by dilution of A0 using NA

to generate all A solutions with choline concentrations ranging between 50 to 0.02 mM choline, while

keeping the total osmolarity constant. Proteoliposomes or liposomes (negative control) were thawed

and diluted 1:5 in NA, and sonicated using a bath sonicator for 20 s. A volume of 10 µl of the diluted

proteoliposomes were added to the sensors, and then centrifuged at 2,000g for 30 min. Assays were

performed on both sample and negative control to distinguish currents originating from FLVCRs and

artifact currents induced by the choline$^+$/Na$^+$ exchange due to interactions of the ions with the

membrane surface. Currents corresponding to different substrate concentrations were measured on

the same sensor, with a total of four different sensors. To determine transport kinetics of choline for

each FLVCR transporter, the measured currents for a given substrate concentration had the average

value of the corresponding negative control current subtracted from it. For the measurement at

different pH conditions, again the single solution exchange configuration was used. The main buffer

was titrated to different pH values that are physiologically relevant, ranging between 6 and 8, before

splitting into A supplemented with 15 mM choline chloride and the corresponding NA supplemented

with 15 mM NaCl. Before each measurement at a given pH, the pH was equilibrated via incubation for

3 minutes in NA solution at the given pH. Subsequently, peak currents were measured upon the

addition of A at the same pH. All pH values were measured sequentially using the same sensor, with a

total of four sensors. For both dose response curves and pH measurements, the error bars represent

the s.d. As the peak current signal can be a combined effect of electrogenic PSS and transport events[24],

we use the EC_{50} parameter to describe the concentration that generates the half maximum response.

Analysis of the data was performed in OriginPro (OriginLab Corporation).

Cryo-EM sample preparation

In order to collect cryo-EM data of FLVCR1 and FLVCR2 in different sample conditions, different

combinations of FLVCR proteins and putative substrate molecules were prepared. For both as-isolated

samples of FLVCRs, the protein concentration was adjusted to approximately 1.5 mg ml^{-1} and subjected

to plunge freezing. For samples supplemented with heme, heme loading was performed prior to the

SEC during protein purification with a 10-fold molar excess to the protein concentration. Peak fractions

were pooled and concentrated to 1.5 mg ml^{-1} before sample vitrification. For samples supplemented

with choline, purified proteins were adjusted to 1.5 mg ml^{-1} and choline was added at a final

concentration of 1 mM. The samples were incubated for 10 minutes at room temperature before

plunge freezing. Identical plunge freezing conditions were applied for all samples: Quantifoil R1.2/1.3

copper grids (mesh 300) were washed in chloroform and subsequently glow-discharged with a PELCO

easiGlow device at 15 mA for 90 seconds. A volume of 4 μl sample was applied to a grid and blotting

was performed for 4 seconds at 4 °C, 100% humidity with nominal blot force 20 immediately before

freezing in liquid ethane, using a Vitrobot Mark IV device (Thermo Scientific).

Cryo-EM image recording

For each cryo-EM sample, a dataset was recorded in Energy-Filtered Transmission Electron Microscopy (EF-TEM) mode using either a Titan Krios G3i or a Krios G4 microscope (Thermo Scientific), both operated at 300 kV. Electron-optical alignments were adjusted with EPU 3.0 - 3.4 (Thermo Scientific). Images were recorded using automation strategies of EPU 3.0 - 3.4 in electron counting mode with either a Gatan K3 (installed on Krios G3i) or a Falcon4 (installed on Krios G4) direct electron detector. For Gatan K3 detector, a nominal magnification of 105,000, corresponding to a calibrated pixel size of 0.837 Å was applied, and dose fractionated movies (80 frames) were recorded at an electron flux of approximately 15 e^- x $pixel^{-1}$ x s^{-1} for 4 s, corresponding to a total dose of ~80 e^-/A^2. For Falcon4 detector, a nominal magnification of 215,000, corresponding to a calibrated pixel size 0.573 Å was applied, dose fractionated movies were recorded in electron-event representation (EER) format at an electron flux of approximately 4 e^- x $pixel^{-1}$ x s^{-1} for 5 s, corresponding to a total dose of ~50 e^-/A^2. Images were recorded between -1.1 and -2.0 μm nominal defocus. Data collection quality was monitored through EPU v. 3.0-3.4 and CryoSparc Live (versions 3.0 and 4.0)[55].

Cryo-EM image processing

For each acquired dataset, the same cryo-EM image processing approach was applied: MotionCor2 was used to correct for beam-induced motion and to generate dose-weighted images[56]. Gctf was used to determine the contrast transfer function (CTF) parameters and perform correction steps[57]. Images with estimated poor resolution (>4 Å) and severe astigmatism (>400 Å) were removed at this step. Particles were picked by TOPAZ and used for all further processing steps[58]. 2D classification, initial model generation, 3D classification, CTF refinement, Bayesian polishing, 3D sorting, and final map reconstructions were performed using RELION (versions 3.1 and 4.0) or cryoSPARC (versions 3.0 and 4.0)[55,59,60]. Fourier shell correlation (FSC) curves and local-resolution estimation were generated in RELION or cryoSPARC for individual final maps. A schematic overview of our processing workflow, and a summary of map qualities are shown in Extended Data Figs. 1-3, and Supplementary Table 4.

Model building and geometry refinement

The first atomic model of FLVCR1 and FLVCR2 were built into the respective EM density maps of the as-isolated state in Coot (version 0.8) or ISOLDE within ChimeraX (version 1.5 and 1.6)[61–63], using the AlphaFold predicted structures as initial templates[21]. After manual backbone tracing and docking of side chains, real-space refinement in Phenix was performed (version 1.18)[64]. Refinement results were manually inspected and corrected if required. This model was used as a template to build all subsequent atomic models. The finalized models were validated by MolProbity implemented in Phenix[65]. Map-to-model cross-validation was performed in Phenix (version 1.18). $FSC_{0.5}$ was used as cut-off to define resolution. A summary of model parameters and the corresponding cryo-EM map statistics is found in Supplementary Table 4. The finalized models were visualized using ChimeraX. The built models of both FLVCR proteins in different states were used as starting structures for MD simulations.

Molecular dynamics simulations

All simulations were run using GROMACS 2022.4[66]. The protein structures were embedded in a lipid bilayer with 75% POPE and 25% POPG with CHARMM-GUI[67] and solvated in TIP3P water with 150 mM NaCl. The charmm36m force field[68] was used with the improved WYF parameter set for cation-pi interactions[69]. Initial systems were minimized for 5000 steepest descend steps and equilibrated for 450 ps of MD in an NVT ensemble and for 1.5 ns in an NPT ensemble. Position restraints of 4000 and 2000 kJ mol^{-1} nm^{-2} in the backbone and side chain heavy atoms, respectively, were gradually released during equilibration. The z-positions of membrane phosphates, as well as lipid dihedrals, were initially restrained with force constants of 1000 kJ mol^{-1} nm^{-2} and 1000 kJ mol^{-1} rad^{-2}, respectively, which were gradually released during equilibration. The initial time step of 1 fs was increased to 2 fs during NPT equilibration. Long range electrostatic interactions were treated with particle-mesh Ewald (PME) with a cut-off of 1.2 nm[70]. Van-der-Waals interactions were cut off beyond a distance of 1.2 nm. The LINCS algorithm was used to constrain the bonds involving hydrogen atoms[71]. During equilibration, a constant temperature of 310 K was maintained with the Berendsen thermostat, using a coupling constant of 1

ps[72]. Constant pressure of 1 bar was established with a semi-isotropic Berendsen barostat and a

coupling constant of 5 ps. In the production runs, a Nosé–Hoover thermostat and a Parrinello–Rahman

barostat were used[73,74].

We used our cryo-EM structures as initial models for simulations of apo and choline-bound inward-

facing FLVCR1, apo and choline-bound inward-facing FLVCR2, and apo outward-facing FLVCR2. An initial

structure of choline-bound outward-facing FLVCR2 was generated aligning apo outward-facing FLVCR2

to choline-bound inward-facing FLVCR2 and maintaining choline in the cavity. In choline entry

simulations, the apo structures were used with 380 mM choline in solution. Simulations of heme-

bound outward- and inward-facing FLVCR2 were prepared by manually placing heme close to the

observed binding region in the N-terminus of the apo structures, outside of the lipid bilayer.

Additionally, a simulation of FLVCR2 in the outward-facing conformation was conducted, with the heme

initially positioned in front of the cavity entrance. Three replicas with random initial velocities from the

Boltzmann distribution were run for the rest of the systems. Choline release simulations were

interrupted after choline exit from the cavity, and hence have variable duration. For all other systems,

each replica was run for 1 µs. To reproduce mitochondrial membrane potential, an additional set of

simulations was run for choline-bound inward- and outward-facing, and apo outward-facing FLVCR2

with choline in solution, in which an electric field of -200 mV was applied in the z-dimension.

Visual Molecular Dynamics (VMD) and MDAnalysis were used to visualize and analyze the trajectories,

respectively[75,76].

Interior tunnels and cavities

Tunnels and cavities were mapped with MOLE 2.5 with a bottleneck radius of 1.2 Å, bottleneck

tolerance 3 Å, origin radius 5 Å, surface radius 10 Å, probe radius 5 Å and an interior threshold of 1.1 Å[77].

We calculated the volume of the cavity using CASTp with a bottleneck radius of 1.4 Å. Residues 297-

320 and 512-516 were removed from the FLVCR1 model to avoid the misattribution of the volume

between internal loops to the cavity volume[78]. Analogously, residues 272-296 and 487-502 were not

included in the cavity volume calculation of FLVCR2.

Multiple sequence alignments

Multiple sequence alignments of FLVCR1 and FLVCR2 from *homo sapiens*, *Felis catus*, *Mus musculus*,

and *Sus scrofa* were performed using Clustal Omega[79].

Data availability

Cryo-EM maps are deposited at the Electron Microscopy Data Bank under accession numbers: EMD-18334, EMD-18335, EMD-18336, EMD-18337, EMD-18338, EMD-18339. Atomic models of human FLVCR1 and FLVCR2 have been deposited to the Protein Data Bank under accession numbers: 8QCS, 8QCT, 8QCX, 8QCY, 8QCZ, 8QD0. All other data is presented in the main text or supplementary materials. Source data are provided with this paper.

Acknowledgements

We thank Hartmut Michel for support and providing infrastructural resources. We thank ChatGPT by OpenAI for assistance with language editing and polishing of the manuscript. We thank the Central Electron Microscopy Facility at MPI of Biophysics for technical support and access to instrumentation.

Funding

This work was supported by the Max Planck Society and the Nobel Laureate Fellowship of the Max Planck Society.

Author contributions

T-H.W. purified proteins, performed biochemical assays, prepared grids, collected cryo-EM data, processed cryo-EM data, refined the structure, built the model, co-drafted the manuscript, and prepared figures. A.C.C. performed MD simulations, analyzed data, co-drafted the manuscript, and prepared figures. W.J. performed cell productions, optimized purification conditions, and purified proteins. A.B. performed SSM-based assays, analyzed data, and prepared figures. S.W. calibrated and aligned the microscope. G.G. assisted in cell culturing and protein purification. G.H., D.W., and S.S. supervised the project. D.W. implemented cell production and protein purification, prepared grids, performed initial cryo-EM screening experiments, collected cryo-EM data, analyzed data, drafted the manuscript, and funded the project. D.W. and S.S. initiated the project. S.S. designed research, evaluated data, funded the project, drafted the manuscript, and generated figures.

Competing interests

The authors declare no conflict-of-interest.

References

1. Khan, A. A. & Quigley, J. G. Heme and FLVCR-related transporter families SLC48 and SLC49. *Mol. Asp. Med.* **34**, 669–682 (2013).

2. Tailor, C. S., Willett, B. J. & Kabat, D. A putative cell surface receptor for anemia-inducing feline leukemia virus subgroup C is a member of a transporter superfamily. *J. Virol.* **73**, 6500–5 (1999).

3. Brown, J. K., Fung, C. & Tailor, C. S. Comprehensive Mapping of Receptor-Functioning Domains in Feline Leukemia Virus Subgroup C Receptor FLVCR1. *J. Virol.* **80**, 1742–1751 (2006).

4. Dutt, S., Hamza, I. & Bartnikas, T. B. Molecular Mechanisms of Iron and Heme Metabolism. *Annu. Rev. Nutr.* **42**, 311–335 (2022).

5. Rajadhyaksha, A. M. *et al.* Mutations in FLVCR1 Cause Posterior Column Ataxia and Retinitis Pigmentosa. *Am. J. Hum. Genet.* **87**, 643–654 (2010).

6. Rey, M. A. *et al.* Enhanced alternative splicing of the FLVCR1 gene in Diamond Blackfan anemia disrupts FLVCR1 expression and function that are critical for erythropoiesis. *Haematologica* **93**, 1617–1626 (2008).

7. Meyer, E. *et al.* Mutations in FLVCR2 Are Associated with Proliferative Vasculopathy and Hydranencephaly-Hydrocephaly Syndrome (Fowler Syndrome). *Am. J. Hum. Genet.* **86**, 471–478 (2010).

8. Thomas, S. *et al.* High-throughput sequencing of a 4.1 Mb linkage interval reveals FLVCR2 deletions and mutations in lethal cerebral vasculopathy. *Hum. Mutat.* **31**, 1134–1141 (2010).

9. Petrillo, S. *et al.* Endothelial cells require functional FLVCR1a during developmental and adult angiogenesis. *Angiogenesis* 1–20 (2023) doi:10.1007/s10456-023-09865-w.

10. Santander, N. *et al.* Lack of Flvcr2 impairs brain angiogenesis without affecting the blood-brain barrier. *J. Clin. Investig.* **130**, 4055–4068 (2020).

11. Peng, C. *et al.* FLVCR1 promotes the proliferation and tumorigenicity of synovial sarcoma through inhibiting apoptosis and autophagy. *Int. J. Oncol.* **52**, 1559–1568 (2018).

12. Su, X. *et al.* The prognostic marker FLVCR2 associated with tumor progression and immune infiltration for acute myeloid leukemia. *Front. Cell Dev. Biol.* **10**, 978786 (2022).

13. Yan, N. Structural Biology of the Major Facilitator Superfamily Transporters. *Annu. Rev. Biophys.*

44, 257–283 (2015).

14. Quigley, J. G. *et al.* Identification of a Human Heme Exporter that Is Essential for Erythropoiesis.

Cell **118**, 757–766 (2004).

15. Duffy, S. P. *et al.* The Fowler Syndrome-Associated Protein FLVCR2 Is an Importer of Heme. *Mol.*

Cell. Biol. **30**, 5318–5324 (2010).

16. Yuan, X., Protchenko, O., Philpott, C. C. & Hamza, I. Topologically Conserved Residues Direct Heme

Transport in HRG-1-related Proteins*. *J. Biol. Chem.* **287**, 4914–4924 (2012).

17. Ponka, P., Sheftel, A. D., English, A. M., Bohle, D. S. & Garcia-Santos, D. Do Mammalian Cells Really

Need to Export and Import Heme? *Trends Biochem. Sci.* **42**, 395–406 (2017).

18. Li, Y. *et al.* MFSD7C switches mitochondrial ATP synthesis to thermogenesis in response to heme.

Nat. Commun. **11**, 4837 (2020).

19. Bärland, N. *et al.* Mechanistic basis of choline import involved in teichoic acids and

lipopolysaccharide modification. *Sci. Adv.* **8**, eabm1122 (2022).

20. Glunde, K., Bhujwalla, Z. M. & Ronen, S. M. Choline metabolism in malignant transformation. *Nat.*

Rev. Cancer **11**, 835–848 (2011).

21. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589

(2021).

22. Perez, C. *et al.* Substrate-bound outward-open state of the betaine transporter BetP provides

insights into Na+ coupling. *Nat. Commun.* **5**, 4231 (2014).

23. Bazzone, A. *et al.* Investigation of sugar binding kinetics of the E. coli sugar/H+ symporter XylE

using solid-supported membrane-based electrophysiology. *J. Biol. Chem.* **298**, 101505 (2022).

24. Bazzone, A., Zerlotti, R., Barthmes, M. & Fertig, N. Functional characterization of SGLT1 using

SSM-based electrophysiology: Kinetics of sugar binding and translocation. *Front. Physiol.* **14**, 1058583

(2023).

25. Bazzone, A. *et al.* SSM-based electrophysiology, a label-free real-time method reveals sugar

binding & transport events in SGLT1. *Biosens. Bioelectron.* **197**, 113763 (2022).

26. Sarter, M. & Parikh, V. Choline transporters, cholinergic transmission and cognition. *Nat. Rev.*

Neurosci. **6**, 48–56 (2005).

27. Drew, D., North, R. A., Nagarathinam, K. & Tanabe, M. Structures and General Transport

Mechanisms by the Major Facilitator Superfamily (MFS). *Chem. Rev.* **121**, 5289–5335 (2021).

28. Martens, C. *et al.* Direct protein-lipid interactions shape the conformational landscape of

secondary transporters. *Nat. Commun.* **9**, 4151 (2018).

29. Corradi, V. *et al.* Emerging Diversity in Lipid–Protein Interactions. *Chem. Rev.* **119**, 5775–5848

(2019).

30. Zhang, X. C. & Han, L. Uniporter substrate binding and transport: reformulating mechanistic

questions. *Biophys. Rep.* **2**, 45–54 (2016).

31. Kalailingam, P. *et al.* Deficiency of MFSD7c results in microcephaly-associated vasculopathy in

Fowler syndrome. *J. Clin. Investig.* **130**, 4081–4093 (2020).

32. Schneider, K., Zimmer, D., Nielsen, H., Herrmann, J. M. & Mühlhaus, T. iMLP, a predictor for

internal matrix targeting-like sequences in mitochondrial proteins. *Biol. Chem.* **402**, 937–943 (2021).

33. O'Donoghue, N., Sweeney, T., Donagh, R., Clarke, K. J. & Porter, R. K. Control of choline oxidation

in rat kidney mitochondria. *Biochim. Biophys. Acta (BBA) - Bioenerg.* **1787**, 1135–1139 (2009).

34. Kenny, T. C. *et al.* Integrative genetic analysis identifies FLVCR1 as a plasma-membrane choline

transporter in mammals. *Cell Metab.* **35**, 1057-1071.e12 (2023).

35. Xie, T. *et al.* Rational exploration of fold atlas for human solute carrier proteins. *Structure* **30**,

1321-1330.e5 (2022).

36. Nigam, S. K. The SLC22 Transporter Family: A Paradigm for the Impact of Drug Transporters on

Metabolic Pathways, Signaling, and Disease. *Annu. Rev. Pharmacol. Toxicol.* **58**, 663–687 (2018).

37. Ueland, P. M. Choline and betaine in health and disease. *J. Inherit. Metab. Dis.* **34**, 3–15 (2011).

38. Zeisel, S. H. Choline: Critical Role During Fetal Development and Dietary Requirements in Adults.

Annu. Rev. Nutr. **26**, 229–250 (2006).

39. Goh, Y. Q., Cheam, G. & Wang, Y. Understanding Choline Bioavailability and Utilization: First Step

Toward Personalizing Choline Nutrition. *J. Agric. Food Chem.* **69**, 10774–10789 (2021).

40. Sanchez-Lopez, E. *et al.* Choline Uptake and Metabolism Modulate Macrophage IL-1β and IL-18

Production. *Cell Metab.* **29**, 1350-1362.e7 (2019).

41. Ghorbani, P. *et al.* Choline metabolism underpins macrophage IL-4 polarization and RELMα up-

regulation in helminth infection. (2022) doi:10.1101/2022.09.30.510305.

42. Bhat, S., El-Kasaby, A., Freissmuth, M. & Sucic, S. Functional and Biochemical Consequences of

Disease Variants in Neurotransmitter Transporters: A Special Emphasis on Folding and Trafficking

Deficits. *Pharmacol. Ther.* **222**, 107785–107785 (2021).

43. Wortmann, S. B. & Mayr, J. A. Choline-related-inherited metabolic diseases—A mini review. *J.*

Inherit. Metab. Dis. **42**, 237–242 (2019).

44. Mehedint, M. G., Craciunescu, C. N. & Zeisel, S. H. Maternal dietary choline deficiency alters

angiogenesis in fetal mouse hippocampus. *Proc. Natl. Acad. Sci.* **107**, 12834–12839 (2010).

45. King, J. H. *et al.* Maternal Choline Supplementation Modulates Placental Markers of Inflammation,

Angiogenesis, and Apoptosis in a Mouse Model of Placental Insufficiency. *Nutrients* **11**, 374 (2019).

46. Kwan, S. T. (Cecilia) *et al.* Maternal choline supplementation during murine pregnancy modulates

placental markers of inflammation, apoptosis and vascularization in a fetal sex-dependent manner.

Placenta **53**, 57–65 (2017).

47. Jin, X., Wang, R., Wang, H., Long, C. & Wang, H. Brain protection against ischemic stroke using

choline as a new molecular bypass treatment. *Acta Pharmacol. Sin.* **36**, 1416–1425 (2015).

48. Irobi, J., Jonghe, P. D. & Timmerman, V. Molecular genetics of distal hereditary motor

neuropathies. *Hum. Mol. Genet.* **13**, R195–R202 (2004).

49. Wang, H. *et al.* Choline transporter mutations in severe congenital myasthenic syndrome disrupt

transporter localization. *Brain* **140**, 2838–2850 (2017).

50. Fagerberg, C. R. *et al.* Choline transporter-like 1 deficiency causes a new type of childhood-onset

neurodegeneration. *Brain* **143**, 94–111 (2019).

51. Bauché, S. *et al.* Impaired Presynaptic High-Affinity Choline Transporter Causes a Congenital

Myasthenic Syndrome with Episodic Apnea. *Am. J. Hum. Genet.* **99**, 753–761 (2016).

52. Cruz, P. M. R. *et al.* Presynaptic congenital myasthenic syndrome due to three novel mutations in

SLC5A7 encoding the sodium-dependant high-affinity choline transporter. *Neuromuscul. Disord.* **31**,

21–28 (2021).

53. Bazzone, A., Barthmes, M. & Fendler, K. Chapter Two SSM-Based Electrophysiology for

Transporter Research. *Methods Enzym.* **594**, 31–83 (2017).

54. Bazzone, A. & Barthmes, M. Functional Characterization of SLC Transporters Using Solid

Supported Membranes. *Methods Mol. Biol. (Clifton, NJ)* **2168**, 73–103 (2020).

55. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid

unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).

56. Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved cryo-

electron microscopy. *Nat. Methods* **14**, 331–332 (2017).

57. Zhang, K. Gctf: Real-time CTF determination and correction. *J. Struct. Biol.* **193**, 1–12 (2016).

58. Bepler, T. *et al.* Positive-unlabeled convolutional neural networks for particle picking in cryo-

electron micrographs. *Nat. methods* **16**, 1153–1160 (2019).

59. Scheres, S. H. W. RELION: Implementation of a Bayesian approach to cryo-EM structure

determination. *J. Struct. Biol.* **180**, 519–530 (2012).

60. Kimanius, D., Dong, L., Sharov, G., Nakane, T. & Scheres, S. H. W. New tools for automated cryo-

EM single-particle analysis in RELION-4.0. *Biochem. J.* **478**, 4169–4185 (2021).

61. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta*

Crystallogr. Sect. D **66**, 486–501 (2010).

62. Croll, T. I. ISOLDE: a physically realistic environment for model building into low-resolution

electron-density maps. *Acta Crystallogr. Sect. D, Struct. Biol.* **74**, 519–530 (2018).

63. Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and analysis:

UCSF ChimeraX Visualization System. *Protein Sci.* **27**, 14–25 (2017).

64. Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular structure

solution. *Acta Crystallogr. Sect. D, Biol. Crystallogr.* **66**, 213–21 (2009).

65. Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular crystallography.

Acta Crystallogr. Sect. D: Biol. Crystallogr. **66**, 12–21 (2010).

66. Abraham, M. J. *et al.* GROMACS: High performance molecular simulations through multi-level

parallelism from laptops to supercomputers. *SoftwareX* **1**, 19–25 (2015).

67. Jo, S., Kim, T., Iyer, V. G. & Im, W. CHARMM-GUI: A web-based graphical user interface for

CHARMM. *J. Comput. Chem.* **29**, 1859–1865 (2008).

68. Huang, J. *et al.* CHARMM36m: an improved force field for folded and intrinsically disordered

proteins. *Nat. Methods* **14**, 71–73 (2017).

69. Khan, H. M., MacKerell, A. D. & Reuter, N. Cation-π Interactions between Methylated Ammonium

Groups and Tryptophan in the CHARMM36 Additive Force Field. *J. Chem. theory Comput.* **15**, 7–12

(2018).

70. Essmann, U. *et al.* A smooth particle mesh Ewald method. *J. Chem. Phys.* **103**, 8577–8593 (1995).

71. Hess, B., Bekker, H., Berendsen, H. J. C. & Fraaije, J. G. E. M. LINCS: A linear constraint solver for

molecular simulations. *J. Comput. Chem.* **18**, 1463–1472 (1997).

72. Berendsen, H. J. C., Postma, J. P. M., Gunsteren, W. F. van, DiNola, A. & Haak, J. R. Molecular

dynamics with coupling to an external bath. *J. Chem. Phys.* **81**, 3684–3690 (1984).

73. Evans, D. J. & Holian, B. L. The Nose–Hoover thermostat. *J. Chem. Phys.* **83**, 4069–4074 (1985).

74. Parrinello, M. & Rahman, A. Polymorphic transitions in single crystals: A new molecular dynamics

method. *J. Appl. Phys.* **52**, 7182–7190 (1981).

75. Humphrey, W., Dalke, A. & Schulten, K. VMD: Visual molecular dynamics. *J. Mol. Graph.* **14**, 33–38

(1996).

76. Michaud-Agrawal, N., Denning, E. J., Woolf, T. B. & Beckstein, O. MDAnalysis: a toolkit for the

analysis of molecular dynamics simulations. *J. Comput. Chem.* **32**, 2319–27 (2010).

77. Sehnal, D. *et al.* MOLE 2.0: advanced approach for analysis of biomacromolecular channels. *J.*

Cheminformatics **5**, 39–39 (2013).

78. Tian, W., Chen, C., Lei, X., Zhao, J. & Liang, J. CASTp 3.0: computed atlas of surface topography of

proteins. *Nucleic Acids Res.* **46**, W363–W367 (2018).

79. Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence alignments

using Clustal Omega. *Mol. Syst. Biol.* **7**, 539–539 (2011).

Fig. 1: Cryo-EM structures of FLVCR1 and FLVCR2 in their inward-facing state.

Cryo-EM density (top) and atomic model (middle and bottom) of inward-facing FLVCR1 (**a**), and inward-

facing FLVCR2 (**b**). The N- and C-domains are coloured in different shades of blue and green for FLVCR1

and FLVCR2, respectively. A transparent cryo-EM density lowpass-filtered at 6 Å is shown to visualize

the detergent belt surrounding the transmembrane region. The internal view (bottom) shows the

arrangement of TMs, with inner side helices highlighted in orange (FLVCR1) or pink (FLVCR2). The

sidechain of N265 in FLVCR1 and its N-linked glycan (grey) are shown as stick models. **c**, Schematic

diagram of FLVCR family showing the topology of the secondary structure. Motifs that are not observed

in both cryo-EM structures of FLVCR1 and FLVCR2 are shown as dashed lines.

Fig. 2: Structural heterogeneity and gating mechanism of FLVCR2.

a, Cryo-EM density (top) and atomic model (bottom) of outward-facing FLVCR2. **b**, Structural

superposition of FLVCR2-OF^{apo} (dark green) and FLVCR2-IF^{apo} (light green). **c**, Cut-away views of the

surface representation showing the cavity shape of FLVCR2-IF^{apo} (left) and FLVCR2-OF^{apo} (right). Two

central aromatic residues are shown as sticks. Cross-sections of inter-domain interactions in FLVCR2-

IF^{apo} (left) and FLVCR2-OF^{apo} (right) are shown from the external (**d**), or from the internal side (**e**). **f**,

Internal view of FLVCR2-OF^{apo} showing the inter-domain interactions between H1 and H3. Residues

participating in the inter-domain interactions are shown as sticks; hydrogen bonds and salt bridges are

labelled with dashed lines.

Fig. 3: Cryo-EM structures of FLVCR1 and FLVCR2 in complex with choline and heme.

a, Cryo-EM density (left) and ribbon model (right) of the heme-bound outward-facing FLVCR2 (FLVCR2-OF^{heme}). The heme molecule is shown in ball-and-stick representation, and the corresponding cryo-EM density is coloured in purple. **b**, Close-up views of the heme-binding site in FLVCR2-OF^{heme} (top) and in FLVCR2-OF^{apo} (bottom) with locally-filtered cryo-EM density. Residues in close proximity to heme are shown as sticks. A lipid molecule (grey ball-and-stick) is fitted into the lipid-like density at the heme-binding site of FLVCR2-OF^{apo}. **c**, Time-resolved distance plot between heme and interacting residues from MD experiments are shown. A snapshot of FLVCR2-heme interaction during the simulation is shown. For simplicity, only the phosphates of POPE/POPG are shown as red spheres. **d,e**, Cryo-EM densities and atomic models of the choline-bound inward-facing FLVCR1 (FLVCR1-IF^{choline}) structure, and the choline-bound inward-facing FLVCR2 (FLVCR2-IF^{choline}) structure, respectively. The bound choline is shown as ball-and-stick model; binding site residues are shown as sticks. **f,g**, Time-resolved

distance plots between choline and the highly-conserved tryptophan and tyrosine residues forming the

choline-binding pockets of FLVCR1 (**f**) and FLVCR2 (**g**) obtained from MD simulation runs. A cation-π

interaction is assumed for distance <4 Å (grey dashed line). **h**, Protein sequence alignment of choline-

binding pocket residues (red block) in FLVCR1 and FLVCR2 across various mammalian species. Indicated

residue numbers refer to FLVCR1 and FLVCR2 from *homo sapiens*. **i**, Choline-binding affinity of FLVCR1

(left) and FLVCR2 (right) determined by nanoDSF. Data are represented as mean ± standard deviation

(s.d.; n = 3). **j**, Representative choline-induced current traces of FLVCR1 (left) or FLVCR2 (right)

containing proteoliposomes. Representative current traces of protein-free liposomes induced by 50

mM choline are shown as magenta dashed lines. Data are represented as mean ± s.d. (n = 4).

Fig. 4: Proposed model for choline transport by FLVCR2.

Schematic illustration of FLVCR2 conformations during the choline transport cycle. Green-coloured states represent experimentally obtained conformations in this study. States coloured in grey are hypothesized based on knowledge about the commonly characterized alternative-access mechanism of MFS transporters.

Extended Data Fig. 1: Single-particle cryo-EM analysis of human FLVCR1.

a,b, Summary of the data processing procedure of the as-isolated FLVCR1 sample and FLVCR1 supplemented with choline. **c,d**, Local resolution estimation (left) and Fourier shell correlation (FSC) curves (right) of the final cryo-EM maps of FLVCR1-IF^apo and -IF^choline.

856 **Extended Data Fig. 2: Single-particle cryo-EM analysis of human FLVCR1.**

857 **a-c**, Summary of the data processing procedure of the as-isolated FLVCR2, heme-supplemented, and

858 choline-supplemented FLVCR2 samples. **d-g**, Local resolution estimation (top) and FSC curves (bottom)

859 of the final cryo-EM maps of FLVCR2-IF^{apo}, FLVCR2-OF^{apo}, FLVCR2-OF^{heme}, and FLVCR2-IF^{choline}.

Extended Data Fig. 3: Cryo-EM density and model fitting.

Cryo-EM map and model fits of the transmembrane and intracellular helices of FLVCR1-IF^{apo} (**a**), FLVCR1-IF^{choline} (**b**), FLVCR2-IF^{apo} (**c**), FLVCR2-OF^{apo} (**d**), FLVCR2-OF^{heme} (**e**), and FLVCR2-IF^{choline} (**f**). The map for choline and the surrounding residues is shown for the choline-bound states of FLVCR1 and FLVCR2, respectively.

Extended Data Fig. 4: Structural comparison between FLVCR1-IF^{apo} and FLVCR2-IF^{apo}.

a, Structural comparison of FLVCR1-IF^{apo} and FLVCR2-IF^{apo} in tube representation, viewed from the lipid

bilayer (left), extracellular side (top-right), and intracellular side (bottom-right). **b**, Cut-away view of

FLVCR1-IF^{apo} in surface representation showing the cytoplasmic cavity. Two central aromatic residues

are shown as sticks. **c,d**, Cross-sections of inter-domain interactions in FLVCR1-IF^{apo} as indicated in (**b**),

viewed from the extracellular side (**c**), or from the intracellular side (**d**). Residues corresponding to the

inter-domain interaction residues in FLVCR2 are shown.

Extended Data Fig. 5: Chemicophysical properties of FLVCR structures and conservation analyses.

Surface charge, hydrophobicity, and conservation analyses of FLVCR1-IF[apo] (**a**), FLVCR1-IF[choline] (**b**), FLVCR2-IF[apo] (**c**), FLVCR2-OF[apo] (**d**), and FLVCR2-IF[choline] (**e**). From left to right: Surface viewed from the lipid bilayer and both C and N domains viewed from the domain interface coloured by electrostatic potential, the domain interface coloured by hydrophobicity, and the domain interface coloured by

sequence conservation. The cavity in the different states of both FLVCRs is outlined with dashed lines.

Two important central pocket residues (W125 and Y349 in FLVCR1; W102 and Y325 in FLVCR2) are

indicated.

Extended Data Fig. 6: Structures of bacterial and mammalian choline transporters.

Experimental or predicted structures of bacterial choline transporter LicB (PDB 7B0K; **a**) and other

known human choline transporters SLC5A7 (AlphaFold AF-Q9GZV3-F1; **b**), SLC44A1 (PDB 7WWB; **c**),

and SLC22A5 (AlphaFold AF-O76082-F1; **d**). The lower panels show the close-up views of their

respective binding sites. Choline or putative choline-binding cavities are highlighted in pink. Choline-

binding residues or residues lining a putative choline-binding site are indicated in yellow.

Extended Data Fig. 7: Binding affinity screening of FLVCRs in the presence of choline or its metabolites

by nano differential scanning fluorimetry (nanoDSF).

a-c, Representative nanoDSF measurements (F_{350}/F_{330}) along a temperature ramp from 25°C to 95°C

(top) and the respective first derivative (bottom) of detergent-solubilized FLVCR1 and FLVCR2 in the

presence of a serial dilution of choline (**a,d**), phosphocholine (**b,e**), or betaine (**c,f**). **g**, Representative

negative-control assay using a purified human sodium transporter from the SLC9 family in the presence

of a serial dilution of choline. Magenta lines represent measurements of the samples without choline.

Red vertical dashed lines represent the melting temperatures of the samples without choline and the

samples of FLVCRs with 2 mM choline.

Extended Data Fig. 8: SSM-based electrophysiological profiles of FLVCRs in proteoliposomes.

a, Current traces for FLVCR1 (left) or FLVCR2 (right) proteoliposomes with buffers at different pH as

indicated in the figure. **b**, Normalized peak currents for FLVCR1 (left) and FLVCR2 (right) based on the

measurement in (**a**). Data are represented as mean ± s.d. (n = 4).

Extended Data Fig. 9: Structural comparison of FLVCRs in their choline-bound inward-facing states.

a, Choline-binding sites of FLVCR2-OF[apo] (top) and FLVCR2-IF[choline] (bottom) with the distance between

W102 and Y325 shown as dashed lines. **b**, Cut-away views of FLVCR1-IF[choline] (top-left) and FLVCR2-

IF[choline] (top-right) showing the inward-facing cavity. Two central aromatic residues are shown as sticks.

The surfaces shown below are the IF[apo] and IF[choline] states of FLVCR1 (left) and FLVCR2 (right) viewed

from the internal side. The dashed circles indicate the peripheral channel in the choline-bound state.

c,d, Cross-sections of inter-domain interactions in FLVCR1-IF[choline] (left) and FLVCR2-IF[choline] (right) as

indicated in (**b**), viewed from the external side (**c**), or from the internal side (**d**). **e**, Structural comparison

of FLVCR1-IF^{choline} and FLVCR1-IF^{apo}, viewed from the lipid bilayer (left), external side (top-right), and

internal side (bottom-right). **h**, Structural comparison of FLVCR2-IF^{choline} and FLVCR2-IF^{apo}, viewed from

the lipid bilayer (left), external side (top-right), and internal side (bottom-right).

Extended Data Fig. 10: Lateral gate in the inward-facing state of FLVCR2.

a, An MD-snapshot of FLVCR2 in the inward-facing state showing the entry of a lipid acyl group into the lateral gate in the cytoplasmic leaflet. The POPE molecule is shown in yellow and heteroatom colour codes. For simplicity, only the phosphates of the other POPE/POPG are shown as red spheres. **b,c**, Cryo-EM density maps of FLVCR-IF[apo] and -IF[choline] in viewing angle of the lipid-binding site. The lipid/detergent density is coloured grey. Lipid/detergent density entering the lateral gate is shown in yellow. The three residues at the lateral gate are shown as stick models.

931
932