Structural and mechanistic insights into human choline and ethanolamine transport

Keiken Ri[1-5†], Tsai-Hsuan Weng[6†], Ainara Claveras Cabezudo[7,8†], Wiebke Jösting[6], Zhang Yu[1], Andre Bazzone[9], Nancy C.P. Leong[1-5], Sonja Welsch[10], Raymond T. Doty[11], Gonca Gursu[6], Tiffany Jia Ying Lim[1-5], Sarah Luise Schmidt[6], Janis L. Abkowitz[11], Gerhard Hummer[7,12]*, Di Wu[6,15]*, Long N Nguyen[1-5]*, Schara Safarian[6,13-15]*

Affiliations

[1] Department of Biochemistry, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 119228

[2] Immunology Program, Life Sciences Institute, National University of Singapore, Singapore 117456

[3] Singapore Lipidomics Incubator (SLING), Life Sciences Institute, National University of Singapore, Singapore 117456

[4] Cardiovascular Disease Research (CVD) Programme, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117545

[5] Immunology Translational Research Program, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 117456

[6] Department of Molecular Membrane Biology, Max Planck Institute of Biophysics, D-60438 Frankfurt/Main, Germany

[7] Department of Theoretical Biophysics, Max Planck Institute of Biophysics, D-60438 Frankfurt/Main, Germany

[8] IMPRS on Cellular Biophysics, D-60438 Frankfurt/Main, Germany

[9] Nanion Technologies GmbH, Ganghoferstr. 70a, 80339, Munich, Germany

[10] Central Electron Microscopy Facility, Max Planck Institute of Biophysics, D-60438 Frankfurt am Main, Germany

[11] Division of Hematology, Department of Medicine, University of Washington, Seattle, WA, 98195, USA

[12] Institute of Biophysics, Goethe University Frankfurt, D-60438 Frankfurt/Main, Germany

[13] Institute of Clinical Pharmacology, Goethe University Frankfurt, D-60438 Frankfurt/Main, Germany

[14] Fraunhofer Cluster of Excellence for Immune Mediated Diseases CIMD, D-60438 Frankfurt/Main, Germany

[15] Fraunhofer Institute for Translational Medicine and Pharmacology ITMP Frankfurt, D-60438 Frankfurt/Main, Germany

*Correspondence and requests for materials should be addressed to Gerhard Hummer (gerhard.hummer@biophys.mpg.de), Di Wu (di.wu@biophys.mpg.de), Long Nguyen (bchnnl@nus.edu.sg), and Schara Safarian (schara.safarian@biophys.mpg.de).

†These authors contributed equally.

Abstract

Human feline leukemia virus subgroup C receptor-related proteins 1 and 2 (FLVCR1 and 2) are members of the major facilitator superfamily[1]. Their dysfunction is linked to several clinical disorders, including PCARP, HSAN, and Fowler syndrome[2–7]. Earlier studies concluded that FLVCR1 may function as a putative heme exporter[8–12], while FLVCR2 was suggested to act as a heme importer[13], yet conclusive biochemical and detailed molecular evidence remained elusive for the function of both transporters[14–17]. Here, we show that FLVCR1 and FLVCR2 facilitate the transport of choline and ethanolamine across human plasma membranes, utilizing a concentration-driven substrate translocation process. Through structural and computational analyses, we have identified distinct conformational states of FLVCRs and unraveled the coordination chemistry underlying their substrate interactions. Within the binding pocket of both transporters, we identify fully conserved tryptophan and tyrosine residues holding a central role in the formation of cation-π interactions, essential for choline and ethanolamine selectivity. Our findings not only clarify the mechanisms of choline and ethanolamine transport by FLVCR1 and FLVCR2, enhancing our comprehension of disease-associated mutations that interfere with these vital processes, but also shed light on the conformational dynamics of these MFS-type proteins during the transport cycle.

Introduction

The feline leukaemia virus subgroup C receptor (FLVCR) family, a member of the major facilitator superfamily (MFS) of secondary active transporters, consists of four paralogues encoded by the human *SLC49* gene group[1]. FLVCR1 (SLC49A1) was initially identified as the cell surface receptor for feline leukaemia virus (FeLV)[18]. FLVCR2 (SLC49A2) shares 60% sequence identity with FLVCR1 in the transmembrane domain but does not bind the feline leukaemia virus subgroup C envelope protein[19]. Both transporters exhibit ubiquitous tissue distribution in humans and have significant haemato- and neuropathological implications[1,17]. Dysfunction of FLVCR1 caused by germline mutations is associated with posterior column ataxia with retinitis pigmentosa (PCARP)[2,3], and hereditary sensory and autonomic neuropathies (HSANs)[4,5]. Similarly, truncation and missense mutations in *FLVCR2* are associated with autosomal-recessive cerebral proliferative vasculopathy (Fowler syndrome)[6,7]. Furthermore, both FLVCR variants are suggested to play a key role in cell development and differentiation, including angiogenesis and tumorigenesis[20–23].

Earlier studies concluded that FLVCR1 may function as a putative heme exporter[8–12], while FLVCR2 was suggested to act as a heme importer[13], yet their definitive roles in this capacity remain elusive[14–17]. To accurately understand their functions, experimental validation at the biochemical and molecular levels is necessary, which will connect these transporters' physiological roles and clinical relevance to their specific mechanistic actions. Recent studies indicated FLVCR1 is involved in choline transport, however, the ligands for FLVCR2 still remain elusive[24]. Here, we used an integrative approach, including cell-based radioligand transport assays, single-particle analysis (SPA) cryogenic-electron microscopy (cryo-EM), structure-guided mutagenesis, and atomistic molecular dynamics (MD) simulations to characterize the ligand specificity, molecular architecture, and the conformational landscape of FLVCR1 and FLVCR2 transporters.

Results

***Choline and ethanolamine are transport substates of* FLVCR1 and FLVCR2**

We expressed the human *FLVCR1* and *FLVCR2* genes in human embryonic kidney (HEK293)

cells to substantiate and characterize their roles in cellular choline and ethanolamine transport[25,26].

Upon overproduction, FLVCR1a (hereafter referred to as FLVCR1) and FLVCR2 localized at the plasma

membrane of HEK293 cells (Fig. 1a and Supplementary Fig. 1a-d). Radioactive [^{3}H] choline transport

assays showed a discernible increase in uptake facilitated by the FLVCR2, whereas FLVCR1 did not

exhibit such an effect under the tested condition (Extended Data Fig. 1a). Notably, co-expression of

the choline kinase A (*CHKA*) gene significantly enhanced choline uptake by both transporters in dose-

and time-dependent manners (Fig. 1b-d). This suggests that choline influx mediated by FLVCR1 and

FLVCR2 is enhanced by the overproduction of downstream choline-utilizing enzymes. In order to

ascertain whether choline is the principal physiological substrate for FLVCR1, a comprehensive

metabolomic analysis was performed on liver samples from FLVCR1 knockout mice (Extended Data Fig.

2a). This led us to discover that, in addition to choline and its metabolites, ethanolamine metabolite

profiles were also affected in FLVCR1 knockout livers (Extended Data Fig. 2b,c). Subsequent cell-based

assays revealed that both FLVCRs facilitate ethanolamine uptake into cells (Extended Data Fig. 1b).

Notably, co-expression of *ETNK-1* enhanced the ethanolamine transport rate of FLVCR1 by five-fold,

but did not substantially affect the efficiency of FLVCR2 (Fig. 1e-g and Extended Data Fig. 1b). We

determined transport kinetic parameters of FLVCR1 and FLVCR2, revealing K_m values of 47.4 ± 9.8 μM

and 64.0 ± 21.0 μM for choline, 8 ± 1.5 μM and 41.5 ± 32.0 μM for ethanolamine, respectively (Fig.

1c,f). To decipher the transport mechanism of both FLVCRs, we investigated their reliance on sodium

ions and pH levels. Our uptake studies revealed that FLVCR-mediated transport of choline and

ethanolamine is not contingent upon sodium ion involvement and operates effectively across a broad

pH range (Extended Data Fig. 3a,b). This finding underscores that neither sodium nor pH gradients are

critical for the translocation of choline and ethanolamine by FLVCRs. Next, we further assessed the

mechanistic properties of FLVCR2 by performing a choline washout assay. Upon inverting the choline

gradient across the plasma membrane, we measured a significant decrease of cellular choline levels

within 1 h indicating for a bidirectional choline transport activity mediated by FLVCR2 (Extended Data

Fig. 3c). FLVCR1 showed similar properties in an ethanolamine washout experiment (Extended Data

Fig. 3c). Our findings suggest that both FLVCR1 and FLVCR2 operate as uniporters, facilitating downhill

ligand transport independent of sodium or pH gradients but aided by the typically negative membrane

potential[27].

FLVCR architecture and conformational landscape

Next, we aimed to establish structure-function relationships for the choline and ethanolamine

transport properties of FLVCR1 and FLVCR2. Wildtype FLVCR1 and FLVCR2 proteins were purified and

subjected to SPA cryo-EM (Supplementary Fig. 1a-f). We determined inward-facing conformation

structures of FLVCR1 and FLVCR2 in their apo states at 2.9 Å resolution (Fig. 2a,b, and Supplementary

Fig. 2,3). In addition, we successfully elucidated the outward-facing conformational structure of

FLVCR2 at 3.1 Å resolution from the same sample preparation (Fig. 2c and Supplementary Fig. 3). Both

FLVCR paralogs share a common MFS-type architecture[28], with their N-domain containing

transmembrane domain (TMs) 1-6 and the C-domain containing TMs 7-12 connected by a long and

flexible loop containing two short horizontal cytoplasmic helices (H1 and H2) (Extended Data Fig. 4a,b).

In the FLVCR2 structure, we resolved a short helical segment at the C-terminus (H3), while the density

of the FLVCR1 C-terminus was less pronounced (Extended Data Fig. 4a,b and Supplementary Fig. 4).

Further, we identified an N-linked glycosylation site at N265 of FLVCR1 which locates within the

extracellular loop connecting TMs 5 and 6 (EL5-6) (Fig. 2a). In contrast, no glycosylation site was

identified in the FLVCR2 structure[19] (Fig. 2b,c and Supplementary Fig. 5).

The inward-facing conformations of FLVCR1 and FLVCR2 exhibit a close resemblance to each

other, with a root mean square deviation (Cα r.m.s.d.) of 0.993 Å (Extended Data Fig. 4b). Both

structures feature a wedge-shaped solvent-accessible cavity that is mainly created by the separation

of TMs 4 and 5 of the N-domain from TMs 10 and 11 of the C-domain (Fig. 2d and Extended Data Fig.

4b and 5a). This space extends halfway across the membrane with a similar depth of around 23 Å in

both transporters. On the level of the outer leaflet, TMs 1, 2, and 5 of the N-domain and TMs 7, 8, and

11 of the C-domain pack tightly against each other and thus shield the central cavity from the

extracellular space (Fig. 2d and Extended Data Fig. 4b and 5a). Notably, TMs 1, 4, and 7 exhibit

disordered regions at their C-terminal ends, followed by short kinked helical motifs designated TMs 1b,

4b, and 7b (Fig. 2d and Extended Data Fig. 4a and 5a).

The outward-facing conformation of FLVCR2 features a cavity accessible from the extracellular

space. This cavity is formed through a 'rocker-switch' rigid-body motion that occurs during the

transition from the inward- to outward-facing state (Fig. 2d, e and Supplementary Video 1). This

motion involves a shift of the outer halves of all transmembrane segments away from the central axis.

Concurrently, the inner halves of these TMs move inward, effectively blocking the exit pathway. (Fig.

2e). The cavities in outward- and inward-facing conformations are 7.4 Å and 8.6 Å wide at their

respective openings (Fig. 2d). Both inward- and outward-facing cavities are lined by uncharged and

hydrophobic residues which are mostly conserved (Extended Data Fig. 6). Compared to the inward-

facing cavity, the outward-facing cavity demonstrates a more restricted pathway within its central

region. In this narrowed segment of the cavity, the channel ends at W102^{FLVCR2} and Y325^{FLVCR2}

(equivalent to W125^{FLVCR1} and Y349^{FLVCR1}), two residues that are highly conserved in both FLVCR

transporters (Fig. 2d and Extended Data Fig. 5a and 6).

In line with an alternating-access model, the inward-facing conformation of FLVCR2 features a

tightly-sealed extracellular gate created by inter-domain interactions. This is mainly achieved by the

juxtaposition of TM1b (N-domain) and TM7b (C-domain) (Fig. 2d and Extended Data Fig. 4b). The inter-

domain interaction between these two motifs is stabilized by a hydrogen bonding network consisting

of two pseudo-symmetry-related asparagine residues N110^{FLVCR2} and N332^{FLVCR2}, and E343^{FLVCR2} (TM8)

and N239^{FLVCR2} (EL5-6) (Extended Data Fig. 5a and Supplementary Fig. 6a). Furthermore, we identified

a stable inter-domain salt bridge between D124^{FLVCR2} (TM2) and R333^{FLVCR2} (TM7b), reinforcing the

external occlusion (Extended Data Fig. 5a and Supplementary Fig. 6b). Our cell-based mutagenesis

studies show that alanine substitutions at N110^{FLVCR2}, E343^{FLVCR2}, D124A^{FLVCR2} and R333A^{FLVCR2}

individually result in a significant decrease of choline uptake and an almost complete perturbation of

ethanolamine transport, highlighting a critical role of inter-domain interactions on the extracellular

surface for FLVCR2 functionality (Extended Data Fig. 5b and Supplementary Fig. 7). While FLVCR1

exhibits a similar extracellular hydrogen bonding network as FLVCR2, it lacks a corresponding salt

bridge interaction (Extended Data Fig. 5a). Notably, the disruption of the hydrogen bond between

N133[FLVCR1] and N359[FLVCR1] (structurally analogous to N110[FLVCR2] and N332[FLVCR2]) through an alanine

substitution of N133[FLVCR1] markedly diminishes FLVCR1's transport activity for both choline and

ethanolamine. In contrast, the E367A[FLVCR1] mutation impacts the transport of choline to a stronger

extend as compare to ethanolamine transport (Extended Data Fig. 5c).

During the transition from inward- to outward-facing conformation of FLVCR2, inter-domain

interactions contributing to the extracellular gate become disrupted, while the formation of the

intracellular gate occludes the central cavity from the cytoplasmic side. TM4b of the N-domain moves

in proximity to the N-terminal end of TM11 of the C-domain to establish a first level of occlusion (Fig.

2d,e). An interaction network consisting of several hydrogen bonds and a salt bridge is found within

this region as well (Extended Data Fig. 5a). The E435[FLVCR2] residue in TM11 plays a pivotal and

multifaceted role, forming a hydrogen bond with S203[FLVCR2] and simultaneously establishing a salt

bridge with R200[FLVCR2] in TM4b (Extended Data Fig. 5a and Supplementary Fig. 6b). Individuals with a

homozygous S203Y mutation in FLVCR2 are nonviable, likely attributable to the complete loss of

choline and ethanolamine transport activity[29] (Fig. 1b,c). Our mutagenesis studies underscore a greater

significance of S203[FLVCR2] compared to R200[FLVCR2], particularly in choline transport (Fig. 1b and Extended

Data Fig. 5b). This suggests that the interaction between E435[FLVCR2] and S203[FLVCR2] is essential for either

facilitating conformational changes during the transport cycle or maintaining the stability of FLVCR2 in

its outward-facing state. An additional inter-domain hydrogen bond is identified between S199[FLVCR2] in

TM4b and S439[FLVCR2] in TM1 (Extended Data Fig. 5a and Supplementary Fig. 6a). In the peripheral region,

K372[FLVCR2] and R374[FLVCR2] (EL8-9) approach N209[FLVCR2] (EL4-5) and S212[FLVCR2] (TM5) to form hydrogen

bond pairs and thus block the lateral accessibility of the cavity (Extended Data Fig. 5a and

Supplementary Fig. 6a). A second level of occlusion was observed beneath the cytoplasmic ends of

TMs 4, 10 and 11, where H1 and H3 are positioned in close proximity (Extended Data Fig. 5a). Here,

the backbone carbonyl group and amide nitrogen of A283[FLVCR2] (H1) form stable hydrogen bonds with

N497[FLVCR2] (H3) and Y431[FLVCR2] (IL10-11), respectively (Extended Data Fig. 5a and Supplementary Fig.

6a). Together with the loop connecting TMs 10 and 11, the two helical motifs H1 and H3 serve as a

cytoplasmic latch to secure the closure of the two domains.

Choline binding and coordination chemistry of FLVCR1 and FLVCR2

To decipher the substrate binding and coordination chemistry of FLVCR1 and FLVCR2, we

employed choline as a ligand and determined their structures at resolutions of 2.6 Å and 2.8 Å,

respectively (Fig. 3a,b and Supplementary Fig. 2, 3 and 4d,e). Both structures were captured in the

inward-facing conformation, suggesting that choline is capable of resolving the previously captured

conformational heterogeneity of FLVCR2. The choline-bound structures exhibit a Cα r.m.s.d. of 0.688

Å (FLVCR1) and 1.002 Å (FLVCR2), respectively, compared to their apo inward-facing structures

(Extended Data Fig. 7a,b). In the choline-bound structures, the binding sites are located at analogous

positions, with the choline molecule situated between the two domains, surrounded mainly by TMs 1,

2, 4, 5, 7, and 11 (Fig. 3a,b). The binding site in both transporters comprises an identical composition

of conserved residues. We observed that W102[FLVCR1] and W125[FLVCR2] in TM1 directly interact with

choline. This central coordinating tryptophan residue is located above choline, protruding its side chain

to constrain the diffusion of the molecule towards the extracellular space (Fig. 2d and Fig 3c,d). Two

additional aromatic residues of TM7, one tyrosine (Y325[FLVCR1] and Y349[FLVCR2]) and one phenylalanine

(F348[FLVCR1] and F324[FLVCR2]), line the peripheral space of the binding site and restrict the movement of

choline within the pocket. Our MD simulations demonstrate that the quaternary ammonium group of

choline forms stable cation-π interactions with the conserved tryptophan (W125[FLVCR1] and W102[FLVCR2])

and tyrosine residues (Y325[FLVCR1] and Y349[FLVCR2]) in a simultaneous manner (Fig. 3c,d). Mutations of

W125A[FLVCR1] and W102A[FLVCR2] significantly reduced choline transport activity (Fig. 3e and

Supplementary Fig. 7). MD simulations further confirmed the key role of the tryptophan in both

transporters (Fig. 3f,g). We also found that the hydroxyl group of choline is essential for ligand

recognition by both FLVCRs. Substitution to a carboxylic group as found in betaine abolishes ligand

binding and transport activity[26] (Fig. 3h). In our choline simulations, this hydroxyl shows versatile

interactions by forming transient hydrogen bonds with two proximal asparagine residues

(Q214^{FLVCR1}/Q471^{FLVCR1}, Q191^{FLVCR2}/Q447^{FLVCR2}). Further interactions come from at least two water

molecules that were consistently present around the hydroxyl moiety of the choline (Supplementary

Fig. 8 and Supplementary Video 2). Functional analyses of Q214^{FLVCR1}, Q471^{FLVCR1}, Q191^{FLVCR2}, and

Q447^{FLVCR2} indicate that changes of the local protein environment around the hydroxyl group of choline,

caused by alanine substitutions, discernibly affect the transport activities of both FLVCRs (Fig. 3e). It is

noteworthy that all these residues are fully conserved in mammalian FLVCR homologs, suggesting a

common substrate profile of FLVCRs across species (Fig. 3i).

Molecular mechanism of ligand selectivity of FLVCR1

Guided by our functional analyses, we advanced our investigation into the molecular

mechanism of ethanolamine binding by FLVCR1. We obtained the cryo-EM structure of FLVCR1

complexed with ethanolamine at a resolution of 3.1 Å, showing an inward-facing conformation

analogous to the choline-bound FLVCR1 structure (Fig. 4a, Extended Data Fig. 7c and Supplementary

Fig. 2). The ethanolamine density was identified within the abovementioned ligand binding-pocket,

with its primary amine group engaging in cation-π interactions with the conserved W125^{FLVCR1} and

Y349^{FLVCR1} residues (Fig. 4b, and Supplementary Fig. 4f). Our cryo-EM data, corroborated by MD

simulations, indicate that ethanolamine is positioned slightly further from W125^{FLVCR1} and Y349^{FLVCR1}

than choline, attributable to differences in their molecular orientations (Fig. 3a, 4a and 4c, and

Extended Data Fig. 8). While the quaternary ammonium group of choline is situated near Q214^{FLVCR1},

ethanolamine's orientation results in its amine group being distanced, consequently bringing its

hydroxyl group into closer proximity to Q214^{FLVCR1} (Fig. 4d). These nuanced molecular differences

presumably contribute to the selective substrate profile of FLVCR1 as observed in our cell-based

transport assays where FLVCR1 showed a preferential ethanolamine transport activity (Fig. 1b, c, 3e

and 4e). This is further substantiated by the Q214A^{FLVCR1} mutant variant, which results in a complete

loss of ethanolamine transport activity while only partially affecting choline transport (Fig. 3e and 4e).

Notably, alanine substitution of the analogous Q191[FLVCR2] residue abolishes the transport of both

ethanolamine and choline in FLVCR2 (Fig. 3e and 4e).

Identification of a peripheral heme binding site of FLVCR2

Our cell-based assays, using hemin as a heme analog substrate, revealed that unlike the

choline and ethanolamine uptake data, neither FLVCR1 nor FLVCR2 demonstrated heme import

activity (Fig. 5a). This observation was confirmed by our structural studies which did not reveal any

density for bound heme molecules within the ligand binding cavities of FLVCR1 or FLVCR2, respectively

(Fig. 5b). Of note, neither the dimensions, nor the residues lining the central cavities of both FLVCRs

indicate a structural adaptation for accommodating a heme molecule as observed in other heme

transporters[30–32].

However, upon determining the structure of FLVCR2 in the presence of heme we noticed that

heme resolved the previously characterized conformational heterogeneity, fully driving the

transporter into the outward-facing conformation (Fig. 5b and Supplementary Fig. 9). Additionally, we

identified a density near the N-terminus of TM1 on the intracellular side of FLVCR2, likely

corresponding to heme. This density appears to have replaced a peripherally bound lipid molecule that

we had previously observed in the outward-facing density map of apo FLVCR2 (Fig. 5b). The protein

surface in this particular region displays a patch of positive charges, suggesting the heme molecule to

be held in place by the formation of electrostatic interactions between its two propionate groups and

the side chains of R81[FLVCR2], R82[FLVCR2], and K273[FLVCR2] (Fig. 5b and Extended Data Fig. 6). The map density

of bound heme in our structure indicates local mobility of the heme, possibly due to the absence of

axial coordination of the central iron in the macrocycle scaffold (Fig. 5b). By contrast, cryo-EM studies

on FLVCR1 in the presence of heme did not reveal any heme binding or notable conformational

changes.

To further investigate heme binding in FLVCR2, MD simulations were performed within a lipid

bilayer. Heme was found to interact with the intracellular loop region rather than accessing the central

cavity, in agreement with prior functional and structural observations (Supplementary Video 3). When

placed near the binding site at the N-terminus of TM1 as indicated by our cryo-EM data, heme rapidly

disengaged from R82[FLVCR2], yet maintained contact with R83[FLVCR2], K273[FLVCR2], and I270[FLVCR2] after

insertion into the lipid bilayer (Fig. 5c,d and Supplementary Fig. 10). Subsequent simulations with

alanine substitution variants of these residues led to either reduction or complete loss of heme-binding

events, indicating the involvement of these residues for FLVCR2 interaction with heme (Fig. 5e).

Follow-up structural studies conducted in a competitive manner by the presence of both heme

and choline revealed that choline binding occurs within the central cavity of FLVCR2 irrespective of the

presence of heme and triggers conformational shifts towards the inward-facing state. This occurs

despite heme's ability to drive FLVCR2 into an outward-facing conformation in the absence of choline

(Fig. 5f and Supplementary Fig. 9). Our results from the cell-based transport assays showed that the

introduction of mutations at the heme-binding residues marginally influenced the choline and

ethanolamine transport function of FLVCR2 (Fig. 5g). Nevertheless, co-incubation of heme with either

choline or ethanolamine did not impact the transport activity of these ligands in FLVCR2 (Fig. 5h). In

conclusion, our structural studies reveal a peripheral heme binding site in FLVCR2 that does not play a

critical role in choline and ethanolamine transport.

Translocation pathway of choline and ethanolamine in FLVCRs

MFS transporters typically cycle between inward- and outward-facing conformations,

facilitating substrate translocation in an alternating-access manner, and substrate binding plays a

pivotal role in eliciting the conformational transitions[33]. Our cryo-EM data supports this mechanism in

FLVCR2, where we see a full transition from the outward-facing to inward-facing state upon choline

binding. Complementary to our structural insights into the conformational landscape, we performed

MD simulations to map the route for choline entry along the pathway in the outward-facing

conformation of FLVCR2. After spontaneously diffusing into the translocation pathway, choline initially

interacts with several residues near the protein surface, mainly D124[FLVCR2] (Supplementary Fig. 11). It

sequentially approaches the deeper recesses of the cavity, primarily engaging with conserved aromatic

residues W102[FLVCR2] and Y325[FLVCR2]. In one of the entry events, we observed choline moving

spontaneously to a position below the W102^{FLVCR2} residue within the binding site consistent with our

structural data (Supplementary Video 4).

The substantial global conformational changes triggered by substrate further alter the local

arrangement of substrate-coordinating residues within the translocation pathway as seen in our cryo-

EM maps. The rearrangement of the binding site results in a more constricted pocket, promoted by

the inward movement of the conserved residues towards the substrate molecule (Extended Data Fig.

9a). The repositioning of these residues, especially the conserved aromatic side chains W102^{FLVCR2},

F324^{FLVCR2}, and Y325^{FLVCR2}, (equivalent to W125^{FLVCR1}, F348F^{LVCR1}, and Y349^{FLVCR1}) restricts the

accessibility of the binding pocket and promotes substrate engulfment and coordination. In their

choline-bound structures, the cavities of both FLVCRs share common characteristics, exhibiting a

neutral interior but a negatively charged surface at the exit (Extended Data Fig. 6b, e). FLVCR1 features

a slightly smaller cavity volume of 513 Å^3 compared to 579 Å^3 of FLVCR2 (Extended Data Fig. 9b). While

the cavity narrows toward the intracellular side, a peripheral solvent-accessible channel emerging

from the cytoplasmic space to the binding site reveals a semi-open translocation pathway in the

substrate-bound inward-facing conformations of both transporters, which may facilitate the release

of the bound ligand (Extended Data Fig. 9b).

Discussion

While FLVCR1 has many studies linking its activity to heme export[8–12], the study of FLVCR2-mediated heme uptake has not been confirmed[13], and furthermore, conclusive biochemical and detailed molecular evidence had remained elusive for the function of both transporters[14–17]. In this study, we have investigated the choline and ethanolamine transport properties of FLVCR1 and FLVCR2 via cell-based transport assays and determined cryo-EM structures of human FLVCR variants in distinct apo, choline-bound, and ethanolamine-bound states. Our work provides valuable insights into architecture, ligand binding-chemistries, and conformational landscapes of these two MFS transporters. Via multiple lines of evidence, we arrive at the conclusion that choline and ethanolamine represent primary transport substrates of both FLVCR1 and FLVCR2, while also revealing a peripheral heme-binding site of FLVCR2 with yet unknown functional implications[10,16]. Our findings suggest that both FLVCR1 and FLVCR2 operate as uniporters, facilitating downhill ligand transport independent of sodium or pH gradients. Notably, FLVCR1 and FLVCR2 possess similar ligand binding and coordination chemistries in which a key conserved tryptophan residue (W125A^{FLVCR1}/W102A^{FLVCR2}) forms a cation-π interaction with the ammonium and amine groups of the respective ligands. Further, we characterized the functional importance of several glutamine residues within the substrate-binding site that participate in hydrogen bonds with the hydroxyl groups of choline and ethanolamine during ligand coordination or transport. Functional assays point to a ligand preference between FLVCR1 and FLVCR2. Our structural studies suggest that ligand coordination is a crucial determinant of the transport preference between choline and ethanolamine for both FLVCRs. Given that most solute carrier transporters exhibit ligand promiscuity[34,35], the specific role of FLVCR1 as an ethanolamine transporter and FLVCR2 as a choline transporter in physiological conditions has yet to be confirmed through *in vivo* studies in animal models.

Based on our structural findings and simulation data, we suggest a rocker-switch alternating-access mechanism for the transport cycle of choline import (Fig. 6, and Supplementary Video 4 and 5). We propose a transport cycle model for both FLVCRs in which the outward-facing conformation represents the state for ligand binding from the extracellular space. Substrate-induced conformational

changes drive the transporter towards its inward-facing state. Finally, ligands are released to the

intracellular space to enter metabolic pathways. Subsequent to choline-release, FLVCRs will undergo

additional conformation changes to adopt their outward-facing state in order to re-initiate the

transport cycle. Our structural data did not reveal fully occluded conformations that are key features

of alternating access. Hence, we suspect that these occluded conformations exist only transiently and

rapidly convert towards either the inward- or outward-facing conformations[33,36,37].

In summary, our research delineates the structural framework through which FLVCR1 and

FLVCR2 mediate the cellular transport of choline and ethanolamine. Identifying these proteins as

facilitative transporters provides a crucial groundwork for future characterizations of the physiological

functions of choline and ethanolamine transport *in vivo*. Furthermore, our findings provide

mechanistic insights for understanding of the disease-mechanisms pertaining to the mutations of

these essential genes.

Online Methods

Generation of inducible HEK293 stable cell lines

The complementary DNAs of full-length wildtype FLVCR1 (human SLC49A1, NCBI Reference Sequence NM_014053) and FLVCR2 (human SLC49A2, NCBI Reference Sequence: NM_017791) were cloned into pcDNA5/FRT/TO (Invitrogen) vectors, respectively. The gene for both FLVCRs was modified by a C-terminal FLAG fusion tag. Further details are found in sequence data provided as supplementary information (Supplementary Tables 2 and 3). The recombinant Flp-In T-REx293-FLVCR1 and Flp-In T-REx293-FLVCR2 cell lines were generated by using a tetracycline-inducible and commercially available Flp-In T-REx host-cell line system from Invitrogen. Flp-In T-REx293 cells were cultured in high-glucose Dulbecco's Modified Eagle's Medium (DMEM; Sigma-Aldrich) supplemented with 10% fetal bovine serum (FBS; Gibco), 1% Pen/Strep (Gibco), 1 µg ml^{-1} Zeocin (Thermo Fisher), and 15 µg ml^{-1} blasticidin S hydrochloride (AppliChem) at 37 °C in an atmosphere of 5% CO_2. Cells were not tested for mycoplasma contamination. For stable integration, the pcDNA5/FRT-FLVCR1-FLAG and pcDNA5/FRT-FLVCR2-FLAG vectors were cotransfected with the Flp recombinase encoding expression vector pOG44 (Thermo Fisher) at a 1:13 mass ratio, respectively. All transfection procedures were performed with Lipofectamine™ 2000 reagent according to the manufacturer's instructions (Thermo Fisher). To select for stable clones, transfected cells were cultivated with growth medium containing 100 µg ml^{-1} hygromycin B (AppliChem).

Transport assays in HEK293 cells

HEK293 cells were co-transfected with pcDNA3.1 plasmid and human *FLVCR1* or *FLVCR2* and human choline kinase A (*CHKA*) for choline transport assays or ethanolamine kinase 1 (*ETNK-1*) for ethanolamine transport assays using Lipofectamine™ 2000 reagent (Thermo Fisher Scientific). After 24 hours post-transfection, cells were incubated with DMEM containing 20 µM [^{3}H] choline or 2.5 µM [^{14}C] ethanolamine. The cells were incubated at 37 °C and 5% CO_2 for 1 hour for uptake of the ligands. The cells were subsequently washed with ice-cold plain DMEM and lysed with RIPA buffer by shaking for 30 minutes. The cell lysates were also quantified by scintillation counter Tricarb. Radioactive signals from cell lysates were normalized to total protein levels. For dose curve assays, indicated

concentrations of choline and ethanolamine were incubated with the cells for 1 hour at 37 °C. For time

course assays, the cells were incubated with 20 µM [^{3}H] choline or with 2.5 µM [^{14}C] ethanolamine.

The transport assays were stopped at indicated time points by adding ice-cold plain DMEM. For testing

transport activity of FLVCR1 mutants, 20 µM [^{3}H] choline and 2.5 µM [^{14}C] ethanolamine was used. For

testing transport activity of FLVCR2 mutants, 100 µM [^{3}H] choline and 2.5 µM [^{14}C] ethanolamine was

used. For transport assays of HEK293 cells overexpressed with *FLVCR1* or *FLVCR2* without co-

expressing with *CHKA* or *ETNK-1*, 20 µM [^{3}H] choline and 2.5 µM [^{14}C] ethanolamine was used.

For transport assays with indicated pH conditions, the following buffers were used: pH8.5 buffer (140

mM NaCl, 20 mM Tris-HCl pH 8.5, 2 mM CaCl$_2$, 1 g/L D-glucose), pH6.5 buffer (140 mM NaCl, 20 mM

MES pH 6.5, 2 mM CaCl$_2$, 1 g/L D-glucose), or pH7.5 buffer (140mM NaCl, 20mM HEPES-NaOH, 2mM

CaCl2, 1 g/L D-glucose). For sodium-free buffer, buffer containing 140 mM KCl, 20 mM HEPES-KOH pH

7.5, 2 mM CaCl$_2$, 1 g/L D-glucose was used. In these assays, 20 µM [^{3}H] choline or 2.5 µM [^{14}C]

ethanolamine was used, and the assays were stopped after 15 minutes of incubation with the ligands.

Radioactive signals from cell lysates were normalized to total protein levels.

Immunofluorescent staining (IF)

HEK293 cells were seeded onto 24-well plates with coverslips and maintained in Dulbecco's Modified

Eagle's Medium (Gibco) supplemented with 10% fetal bovine serum and 1% penicillin-streptomycin.

HEK293 cells were co-transfected with *FLVCR1* or *FLVCR2* with membrane expressing GFP (Addgene:

#14757) using Lipofectamine™ 2000 reagent (Thermo Fisher Scientific). The inducible HEK293 stable

cell lines overproducing FLVCR1 or FLVCR2 were seeded onto Millicell EZ SLIDE 8-well glass slides

(Merck), respectively. The stable cell lines were induced at 80% confluence by adding a final

concentration of 2 µg ml^{-1} doxycycline hydrochloride. The protein overproduction was carried out for

24 h. For permeabilization and staining, cells were washed with PBS twice and fixed in 4% PFA for 15

minutes at room temperature, followed by washing with PBS twice, and permeabilized in PBST (PBS

with 0.5% triton-X) for 15 minutes at room temperature. For immunofluorescent staining, the HEK293

cells were subsequently washed with PBS and blocked in 5% normal goat serum for one hour before

staining with FLVCR1 and FLVCR2 polyclonal antibodies at 1:250 dilutions for 1 h and then with alexa

Fluor 555 as secondary antibody at 1:500 dilutions for 1 h. The cells were counter-stained with DAPI

and imaged with laser confocal microscope (Zeiss LSM710). The overproduction stable cells were

treated with the same protocol but stained with Monoclonal ANTI-FLAG® M2-FITC antibody against

their FLAG-tags, and MitoTracker™ Red CMXRos for mitochondria localization. The cells were mounted

using ProLong™ Diamond mounting medium with DAPI and imaged with laser confocal microscope

(Confocal Microscope Leica STELLARIS 5).

Structure-guided mutagenesis

To generate the mutant plasmids for *FLVCR1* and *FLVCR2*, an overlapping PCR approach was used. The

mutated cDNA of *FLVCR1* or *FLVCR2* was cloned into pcDNA3.1 for overexpression. The mutations were

validated by Sanger sequencing. To test the transport activity of these mutants, the mutant plasmid

was either co-transfected with *CHKA* for choline transport assay or *ETNK-1* for ethanolamine transport

assay. After 24 hours of post-transfections, cells were washed with DMEM and incubated with DMEM

containing 20 μM [^{3}H] choline or 2.5 μM [^{14}C] ethanolamine for *FLVCR1* mutants and 100 μM [^{3}H]

choline or 2.5 μM [^{14}C] ethanolamine for *FLVCR2* mutants. The assays were stopped after 1 h of

incubation at 37 °C. Radioactive signal of each mutant was normalized to the total protein levels.

Choline export assay

To examine the export function, *FLVCR1* and *FLVCR2* plasmids were expressed in HEK293 cells without

co-transfection with *CHKA* or *ETNK-1*. The cells were then incubated with 200 μM [^{3}H] choline or 100

μM [^{14}C] ethanolamine for 2 hours to prepack the cells with the ligand. Subsequently, the cells were

washed to remove the ligands left over in the medium and incubated with choline/ethanolamine-free

medium for 1 h at 37 °C for the release of pre-packed ligand. The cells were washed and collected for

quantification of radioactive signals. Samples after 2 hours of incubation with the radioactive ligand

were collected to determine the levels of radioactive levels before the release and used for control.

Hemin import assay

For hemin import activity measurement, HEK293 cells were transfected with *FLVCR1* or *FLVCR2*. After

24 hours of transfection, cells were incubated with 2.5 μM [^{3}H] hemin in DMEM containing 10% FBS

for 1 h. The cells were then washed and collected for quantification of radioactive signals. For the

transport assays where hemin was co-incubated with radioactive ligands, HEK293 cells with

overexpression of *FLVCR2* were incubated with 50 μM hemin and 10 μM [^{3}H] choline or 20 μM hemin

and 2.5 μM [^{14}C] ethanolamine for 1 hour. Radioactive signals from the cells were quantified.

Deglycosylation assay

440 PNGase assay was performed on FLVCR1 and FLVCR2 proteins purified from HEK293 cells were treated

441 with PNGase (NEB). *FLVCR1* and *FLVCR2* expressed in Expi293T GnTi- cells were used for control.

442 Treated protein samples were then subjected to Western blot analysis.

Metabolomic analysis

444 Adult livers (aged 3-6 months old) from control (*FLVCR1*$^{f/f}$) and conditional *FLVCR1* knockout (*FLVCR1*$^{f/f}$

445 Mx1-Cre) mice were used for metabolomic analysis. Briefly, the mice were perfused with PBS to

446 remove blood before organ collection. Liver samples were snap-frozen before shipped for

447 metabolomics by Metabolon. The levels of metabolites were expressed as relative amount.

Production and purification of the human FLVCR1 and FLVCR2.

For protein production, the Flp-In T-REx293-FLVCR1 and Flp-In T-REx293-FLVCR2 cell lines were

cultured in roller bottles (Greiner Bio-One) in growth media containing 100 μg ml^{-1} hygromycin B for

14 d under the above-mentioned conditions. Gene expression was induced at 100% confluence by

adding a final concentration of 2 μg ml^{-1} doxycycline hydrochloride. After 72 h, cells were harvested

with Accutase solution (Sigma-Aldrich) and stored at −80 °C until further use. Harvested cells were

suspended in cold lysis buffer containing 25 mM Tris pH 7.4, 150 mM NaCl, and 0.1 g ml^{-1} SigmaFast

ethylenediaminetetraacetic acid (EDTA)-free protease inhibitor (Sigma-Aldrich) and disrupted by

stirring under high-pressure nitrogen atmosphere (750 MPa) for 45 min at 4 °C in a cell-disruption

vessel (Parr Instrument). The cell lysate was centrifuged at 8,000*g* at 4 °C for 15 min. Subsequently,

the low-velocity supernatant was centrifuged at 220,000*g* at 4 °C for 60 min. Pelleted membranes were

resuspended and stored in a storage buffer containing 25 mM Tris pH 7.4, 150 mM NaCl, 10% glycerol

(v/v), and 0.1 g ml^{-1} SigmaFast EDTA-free protease inhibitor (Sigma-Aldrich).

All purification steps of both FLVCRs were performed at 4 °C. Isolated membranes were solubilized

with 1% (w/v) lauryl maltose neopentyl glycol (LMNG; GLYCON Biochemicals) with gentle stirring for 1

h. The insoluble membrane fraction was removed via ultracentrifugation at 220,000g for 1 h.

Subsequently, the supernatant was incubated with ANTI-FLAG® M2 Affinity Gel resin (Merck) for 1 h.

The resin was preequilibrated with a buffer containing 50 mM Tris pH 7.4, 150 mM NaCl, and 0.02%

LMNG (w/v). The washing step was performed using 20 column volumes (CVs) of wash buffer [50 mM

Tris pH 7.4, 150 mM NaCl, 5% (v/v) glycerol, and 0.02% LMNG). The protein was eluted from the M2

resin with 10 CVs of the same buffer supplemented with 4 mM FLAG® Peptide (Merck). The eluted

sample was concentrated and subjected to a Superdex 200 Increase 10/300 column (GE Healthcare)

equilibrated with size exclusion chromatography (SEC) buffer [50 mM Tris pH 7.4, 150 mM NaCl, and

0.001% (w/v) LMNG]. Peak fractions were pooled, concentrated to 1.5 mg ml^{-1} using an Amicon 50-

kDa cut-off concentrator (Merck Millipore), and stored for further analysis.

Immunoblotting

Affinity-purified proteins were subjected to SDS-PAGE and immunoblotting. FLAG-tagged FLVCR1 and

FLVCR2 were detected using anti-FLAG (F3165, Sigma-Aldrich) antibodies at 1:1,000 dilution. Anti-

mouse IgG antibody conjugated with alkaline phosphatase (A9316, Sigma-Aldrich) was used as

secondary antibody at 1:5,000 dilution. Native FLVCR1 and FLVCR2 proteins were detected by

polyclonal FLVCR1 and FLVCR2 antibodies raised in-house at 1:1000 dilution. GAPDH antibody was

purchased from Santa Cruz.

Tryptophan fluorescence measurement

Tryptophan fluorescence measurements were carried out using Prometheus Panta (NanoTemper

Technologies). Purified protein samples were diluted with dilution buffer containing 50 mM HEPES pH

7.4, 150 mM NaCl, and 0.001% (w/v) LMNG to 1 μM. Buffers with different concentrations of choline

or betaine were prepared by serial dilutions of dilution buffer containing 4 mM of the compounds. The

protein samples were mixed with an equal volume of dilution buffer or the compound-containing

buffer with a final protein concentration of 0.5 μM and then incubated at room temperature for 15

min. A volume of 10 μl mixed solution was used per Prometheus high sensitivity capillary (NanoTemper

Technologies). Recorded F_{350}/F_{330} was analyzed by using Python libraries including pandas, numpy,

scipy and seaborn in Visual Studio Code (Microsoft). Three technical replicates were recorded for data

analysis.

Cryo-EM sample preparation

In order to collect cryo-EM data of FLVCR1 and FLVCR2 in different sample conditions, different

combinations of FLVCR proteins and putative substrate molecules were prepared. For both as-isolated

samples of FLVCRs, the protein concentration was adjusted to approximately 1.5 mg ml^{-1} and

subjected to plunge freezing. For samples supplemented with heme, heme loading was performed

prior to the SEC during protein purification with a 10-fold molar excess to the protein concentration.

Peak fractions were pooled and concentrated to 1.5 mg ml^{-1} before sample vitrification. For samples

supplemented with choline, purified proteins were adjusted to 1.5 mg ml^{-1} and choline was added at

a final concentration of 1 mM. The samples were incubated for 10 minutes at room temperature

before plunge freezing. Identical plunge freezing conditions were applied for all samples: Quantifoil

R1.2/1.3 copper grids (mesh 300) were washed in chloroform and subsequently glow-discharged with

a PELCO easiGlow device at 15 mA for 90 seconds. A volume of 4 µl sample was applied to a grid and

blotting was performed for 4 seconds at 4 °C, 100% humidity with nominal blot force 20 immediately

before freezing in liquid ethane, using a Vitrobot Mark IV device (Thermo Scientific).

Cryo-EM image recording

For each cryo-EM sample, a dataset was recorded in Energy-Filtered Transmission Electron Microscopy

(EF-TEM) mode using either a Titan Krios G3i or a Krios G4 microscope (Thermo Scientific), both

operated at 300 kV. Electron-optical alignments were adjusted with EPU 3.0 - 3.4 (Thermo Scientific).

Images were recorded using automation strategies of EPU 3.0 - 3.4 in electron counting mode with

either a Gatan K3 (installed on Krios G3i) or a Falcon4 (installed on Krios G4) direct electron detector.

For Gatan K3 detector, a nominal magnification of 105,000, corresponding to a calibrated pixel size of

0.837 Å was applied, and dose fractionated movies (80 frames) were recorded at an electron flux of

approximately 15 e$^-$ x pixel^{-1} x s^{-1} for 4 s, corresponding to a total dose of ~80 e$^-$/A^2. For Falcon4 detector,

a nominal magnification of 215,000, corresponding to a calibrated pixel size 0.573 Å was applied, dose

fractionated movies were recorded in electron-event representation (EER) format at an electron flux

of approximately 4 e$^-$ x pixel^{-1} x s^{-1} for 5 s, corresponding to a total dose of ~50 e$^-$/A^2. Images were

recorded between -1.1 and -2.0 μm nominal defocus. Data collection quality was monitored through

EPU v. 3.0-3.4 and CryoSparc Live (versions 3.0 and 4.0)[38].

Cryo-EM image processing

For each acquired dataset, the same cryo-EM image processing approach was applied: MotionCor2

was used to correct for beam-induced motion and to generate dose-weighted images[39]. Gctf was used

to determine the contrast transfer function (CTF) parameters and perform correction steps[40]. Images

with estimated poor resolution (>4 Å) and severe astigmatism (>400 Å) were removed at this step.

Particles were picked by TOPAZ and used for all further processing steps[41]. 2D classification, initial

model generation, 3D classification, CTF refinement, Bayesian polishing, 3D sorting, and final map

reconstructions were performed using RELION (versions 3.1 and 4.0) or cryoSPARC (versions 3.0 and

4.0)[38,42,43]. Fourier shell correlation (FSC) curves and local-resolution estimation were generated in

RELION or cryoSPARC for individual final maps. A schematic overview of our processing workflow, and

a summary of map qualities are shown in Supplementary Figs. 2, 3 and 9, and Supplementary Table 4.

Model building and geometry refinement

The first atomic model of FLVCR1 and FLVCR2 were built into the respective EM density maps of the

as-isolated state in Coot (version 0.8) or ISOLDE within ChimeraX (version 1.5 and 1.6)[44–46], using the

AlphaFold predicted structures as initial templates[47]. After manual backbone tracing and docking of

side chains, real-space refinement in Phenix was performed (version 1.18)[48]. Refinement results were

manually inspected and corrected if required. This model was used as a template to build all

subsequent atomic models. The finalized models were validated by MolProbity implemented in

Phenix[49]. Map-to-model cross-validation was performed in Phenix (version 1.18). FSC$_{0.5}$ was used as

cut-off to define resolution. A summary of model parameters and the corresponding cryo-EM map

statistics is found in Supplementary Table 4. The finalized models were visualized using ChimeraX. The

built models of both FLVCR proteins in different states were used as starting structures for MD

simulations.

Molecular dynamics simulations

All simulations were run using GROMACS 2022.4[50]. The protein structures were embedded in a lipid

bilayer with 75% POPE and 25% POPG with CHARMM-GUI[51] and solvated in TIP3P water with 150 mM

NaCl. The charmm36m force field[52] was used with the improved WYF parameter set for cation-pi

interactions[53]. Initial systems were minimized for 5000 steepest-descent steps and equilibrated for 250

ps of MD in an NVT ensemble and for 1.625 ns in an NPT ensemble. Position restraints of 4000 and

2000 kJ mol^{-1} nm^{-2} in the backbone and side chain heavy atoms, respectively, were gradually released

during equilibration. The z-positions of membrane phosphates, as well as lipid dihedrals, were initially

restrained with force constants of 1000 kJ mol^{-1} nm^{-2}, which were gradually released during

equilibration. The initial time step of 1 fs was increased to 2 fs during NPT equilibration. Long-range

electrostatic interactions were treated with particle-mesh Ewald (PME)[54] with a real-space cut-off of

1.2 nm. Van-der-Waals interactions were cut-off beyond a distance of 1.2 nm. The LINCS algorithm[55]

was used to constrain the bonds involving hydrogen atoms. During equilibration, a constant

temperature of 310 K was maintained with the Berendsen thermostat[56], using a coupling constant of

1 ps. Constant pressure of 1 bar was established with a semi-isotropic Berendsen barostat and a

coupling constant of 5 ps. In the production runs, a Nosé–Hoover thermostat[57] and a Parrinello–

Rahman barostat were used[58].

We used our cryo-EM structures as initial models for simulations of apo and choline-bound inward-

facing FLVCR1, apo and choline-bound inward-facing FLVCR2, and apo outward-facing FLVCR2. An

initial structure of choline-bound outward-facing FLVCR2 was generated aligning apo outward-facing

FLVCR2 to choline-bound inward-facing FLVCR2 and maintaining choline in the cavity. In choline entry

simulations, the apo structures were used with 380 mM choline in solution. For simulations of

ethanolamine-bound FLVCR1, the choline within the cavity of the cryo-EM structure was replaced by

this ligand. Simulations of heme-bound outward- and inward-facing FLVCR2 were prepared by

manually placing heme close to the observed binding region in the N-terminus of the apo structures,

outside of the lipid bilayer. Additionally, simulations of FLVCR2 in the outward-facing conformation

were conducted, with the heme initially positioned in front of the cavity entrance. Three replicas with

random initial velocities from the Boltzmann distribution were run for all systems. Choline release

simulations were interrupted after choline exit from the cavity, and hence have variable duration. For

all other systems, each replica was run for 1 µs.

Alanine substitution mutations were introduced using PyMol[59] and simulated with identical

parameters as those applied in the corresponding wild-type simulations. In FLVCR1, only one mutation

was introduced in the cavity residue W125. Mutations of FLVCR2 included the cavity residue W102

and the heme binding residues R82, R83 (simultaneously) and K273A.

Visual Molecular Dynamics (VMD)[60] and MDAnalysis[61] were used to visualize and analyze the

trajectories, respectively.

Interior tunnels and cavities

Tunnels and cavities were mapped with MOLE 2.5[62] with a bottleneck radius of 1.2 Å, bottleneck

tolerance 3 Å, origin radius 5 Å, surface radius 10 Å, probe radius 5 Å and an interior threshold of 1.1 Å.

We calculated the volume of the cavity using CASTp[63] with a bottleneck radius of 1.4 Å. Residues 297-

320 and 512-516 were removed from the FLVCR1 model to avoid the misattribution of the volume

between internal loops to the cavity volume. Analogously, residues 272-296 and 487-502 were not

included in the cavity volume calculation of FLVCR2.

Multiple sequence alignments

Multiple sequence alignments of FLVCR1 and FLVCR2 from *homo sapiens*, *Felis catus*, *Mus musculus*,

and *Sus scrofa* were performed using Clustal Omega[64].

Data availability

Cryo-EM maps are deposited at the Electron Microscopy Data Bank under accession numbers:EMD-18334, EMD-18335, EMD-18336, EMD-18337, EMD-18338, EMD-18339, EMD-19009, EMD-19018. Atomic models of human FLVCR1 and FLVCR2 have been deposited to the Protein Data Bank under accession numbers: 8QCS, 8QCT, 8QCX, 8QCY, 8QCZ, 8QD0, 8R8T. All other data is presented in the main text or supplementary materials. Source data are provided with this paper.

Acknowledgements

We thank Hartmut Michel for supporting and providing infrastructural resources. We thank the Central Electron Microscopy Facility at MPI of Biophysics for technical support and access to instrumentation.

Funding

This work was supported by the Max Planck Society and the Nobel Laureate Fellowship of the Max Planck Society (to D.W. and S.S.) and Singapore Ministry of Education T2EP30221-0012, T2EP30123-0014, NUHSRO/2022/067/T1 grants (to L.N.N.).

Author contributions

T-H.W. purified proteins, performed biochemical assays, prepared grids, collected cryo-EM data, processed cryo-EM data, refined the structure, built the model, co-drafted the manuscript, and prepared figures. K.R., Z.Y., and T.J.Y. L. performed mutagenesis. K.R. performed transport assays and WB for mutants, and export assays. N.C.P.L. and Z.Y. performed dose curve, time course, and glycosylation assay, and immunostaining. A.C.C. performed MD simulations, analyzed data, co-drafted the manuscript, and prepared figures. W.J. performed cell productions, optimized purification conditions, and purified proteins. A.B. performed assays and analyzed data. R.T.D and J.L.A designed and performed animal model experiments. S.W. calibrated and aligned the microscope. G.G. performed IF assays and protein purification. S.L.S performed protein purification. G.H., D.W., L.N.N., and S.S. supervised the project. D.W. implemented cell production and protein purification, prepared grids, performed initial cryo-EM screening experiments, collected cryo-EM data, analyzed data, drafted the manuscript, generated figures, and funded the project. L.N.N. designed functional assays, interpreted, analyzed data, funded the project, and co-drafted the manuscript. D.W. and S.S. initiated the project. S.S. designed research, evaluated data, funded the project, drafted the manuscript, and generated figures.

Competing interests

The authors declare no conflict-of-interest

References

1. Khan, A. A. & Quigley, J. G. Heme and FLVCR-related transporter families SLC48 and SLC49. *Mol. Asp. Med.* **34**, 669–682 (2013).

2. Rajadhyaksha, A. M. *et al.* Mutations in FLVCR1 Cause Posterior Column Ataxia and Retinitis Pigmentosa. *Am. J. Hum. Genet.* **87**, 643–654 (2010).

3. Shaibani, A., Wong, L.-J., Zhang, V. W., Lewis, R. A. & Shinawi, M. Autosomal recessive posterior column ataxia with retinitis pigmentosa caused by novel mutations in the FLVCR1 gene. *Int. J. Neurosci.* **125**, 43–49 (2015).

4. Chiabrando, D. *et al.* Mutations in the Heme Exporter FLVCR1 Cause Sensory Neurodegeneration with Loss of Pain Perception. *PLoS Genet.* **12**, e1006461 (2016).

5. Lischka, A. *et al.* Genetic landscape of congenital insensitivity to pain and hereditary sensory and autonomic neuropathies. *Brain* **146**, 4880–4890 (2023).

6. Meyer, E. *et al.* Mutations in FLVCR2 Are Associated with Proliferative Vasculopathy and Hydranencephaly-Hydrocephaly Syndrome (Fowler Syndrome). *Am. J. Hum. Genet.* **86**, 471–478 (2010).

7. Thomas, S. *et al.* High-throughput sequencing of a 4.1 Mb linkage interval reveals FLVCR2 deletions and mutations in lethal cerebral vasculopathy. *Hum. Mutat.* **31**, 1134–1141 (2010).

8. Quigley, J. G. *et al.* Identification of a Human Heme Exporter that Is Essential for Erythropoiesis. *Cell* **118**, 757–766 (2004).

9. Keel, S. B. *et al.* A Heme Export Protein Is Required for Red Blood Cell Differentiation and Iron Homeostasis. *Science* **319**, 825–828 (2008).

10. Yang, Z. *et al.* Kinetics and Specificity of Feline Leukemia Virus Subgroup C Receptor (FLVCR) Export Function and Its Dependence on Hemopexin*. *J. Biol. Chem.* **285**, 28874–28882 (2010).

11. Doty, R. T. *et al.* Coordinate expression of heme and globin is essential for effective erythropoiesis. *J. Clin. Investig.* **125**, 4681–4691 (2015).

12. Fiorito, V. *et al.* The heme synthesis-export system regulates the tricarboxylic acid cycle flux and oxidative phosphorylation. *Cell Rep.* **35**, 109252 (2021).

13. Duffy, S. P. *et al.* The Fowler Syndrome-Associated Protein FLVCR2 Is an Importer of Heme. *Mol. Cell. Biol.* **30**, 5318–5324 (2010).

14. Yuan, X., Protchenko, O., Philpott, C. C. & Hamza, I. Topologically Conserved Residues Direct Heme Transport in HRG-1-related Proteins*. *J. Biol. Chem.* **287**, 4914–4924 (2012).

15. Ponka, P., Sheftel, A. D., English, A. M., Bohle, D. S. & Garcia-Santos, D. Do Mammalian Cells Really Need to Export and Import Heme? *Trends Biochem. Sci.* **42**, 395–406 (2017).

16. Li, Y. *et al.* MFSD7C switches mitochondrial ATP synthesis to thermogenesis in response to heme. *Nat. Commun.* **11**, 4837 (2020).

17. Dutt, S., Hamza, I. & Bartnikas, T. B. Molecular Mechanisms of Iron and Heme Metabolism. *Annu. Rev. Nutr.* **42**, 311–335 (2022).

18. Tailor, C. S., Willett, B. J. & Kabat, D. A putative cell surface receptor for anemia-inducing feline leukemia virus subgroup C is a member of a transporter superfamily. *J. Virol.* **73**, 6500–5 (1999).

19. Brown, J. K., Fung, C. & Tailor, C. S. Comprehensive Mapping of Receptor-Functioning Domains in Feline Leukemia Virus Subgroup C Receptor FLVCR1. *J. Virol.* **80**, 1742–1751 (2006).

20. Petrillo, S. *et al.* Endothelial cells require functional FLVCR1a during developmental and adult angiogenesis. *Angiogenesis* 1–20 (2023) doi:10.1007/s10456-023-09865-w.

21. Santander, N. *et al.* Lack of Flvcr2 impairs brain angiogenesis without affecting the blood-brain barrier. *J. Clin. Investig.* **130**, 4055–4068 (2020).

22. Peng, C. *et al.* FLVCR1 promotes the proliferation and tumorigenicity of synovial sarcoma through inhibiting apoptosis and autophagy. *Int. J. Oncol.* **52**, 1559–1568 (2018).

23. Su, X. *et al.* The prognostic marker FLVCR2 associated with tumor progression and immune infiltration for acute myeloid leukemia. *Front. Cell Dev. Biol.* **10**, 978786 (2022).

24. Kenny, T. C. *et al.* Integrative genetic analysis identifies FLVCR1 as a plasma-membrane choline transporter in mammals. *Cell Metab.* **35**, 1057-1071.e12 (2023).

25. Ha, H. T. T. *et al.* Mfsd7b facilitates choline transport and missense mutations affect choline
transport function. *Cell. Mol. life Sci. : CMLS* **81**, 3 (2023).
26. Nguyen, X. T. A. *et al.* MFSD7c functions as a transporter of choline at the blood-brain barrier.
bioRxiv 2023.10.03.560597 (2023) doi:10.1101/2023.10.03.560597.
27. Geisler, C. E. *et al.* Hepatocyte membrane potential regulates serum insulin and insulin sensitivity
by altering hepatic GABA release. *Cell Rep.* **35**, 109298 (2021).
28. Yan, N. Structural Biology of the Major Facilitator Superfamily Transporters. *Annu. Rev. Biophys.*
44, 257–283 (2015).
29. Kalailingam, P. *et al.* Deficiency of MFSD7c results in microcephaly-associated vasculopathy in
Fowler syndrome. *J. Clin. Investig.* **130**, 4081–4093 (2020).
30. Naoe, Y. *et al.* Crystal structure of bacterial haem importer complex in the inward-facing
conformation. *Nat Commun* **7**, 13411 (2016).
31. Woo, J.-S., Zeltina, A., Goetz, B. A. & Locher, K. P. X-ray structure of the Yersinia pestis heme
transporter HmuUV. *Nat Struct Mol Biol* **19**, 1310–1315 (2012).
32. Wu, D. *et al.* Dissecting the conformational complexity and mechanism of a bacterial heme
transporter. *Nat. Chem. Biol.* **19**, 992–1003 (2023).
33. Drew, D., North, R. A., Nagarathinam, K. & Tanabe, M. Structures and General Transport
Mechanisms by the Major Facilitator Superfamily (MFS). *Chem. Rev.* **121**, 5289–5335 (2021).
34. Hediger, M. A., Clémençon, B., Burrier, R. E. & Bruford, E. A. The ABCs of membrane transporters
in health and disease (SLC series): Introduction☆☆☆. *Mol. Asp. Med.* **34**, 95–107 (2013).
35. Lin, L., Yee, S. W., Kim, R. B. & Giacomini, K. M. SLC transporters as therapeutic targets: emerging
opportunities. *Nat. Rev. Drug Discov.* **14**, 543–560 (2015).
36. Yaffe, D., Forrest, L. R. & Schuldiner, S. The ins and outs of vesicular monoamine transporters. *J.*
Gen. Physiol. **150**, 671–682 (2018).
37. Klingenberg, M. Ligand−Protein Interaction in Biomembrane Carriers. The Induced Transition Fit
of Transport Catalysis †. *Biochemistry* **44**, 8563–8570 (2005).
38. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).
39. Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved
cryo-electron microscopy. *Nat. Methods* **14**, 331–332 (2017).
40. Zhang, K. Gctf: Real-time CTF determination and correction. *J. Struct. Biol.* **193**, 1–12 (2016).
41. Bepler, T. *et al.* Positive-unlabeled convolutional neural networks for particle picking in cryo-
electron micrographs. *Nat. methods* **16**, 1153–1160 (2019).
42. Scheres, S. H. W. RELION: Implementation of a Bayesian approach to cryo-EM structure
determination. *J. Struct. Biol.* **180**, 519–530 (2012).
43. Kimanius, D., Dong, L., Sharov, G., Nakane, T. & Scheres, S. H. W. New tools for automated cryo-
EM single-particle analysis in RELION-4.0. *Biochem. J.* **478**, 4169–4185 (2021).
44. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta*
Crystallogr. Sect. D **66**, 486–501 (2010).
45. Croll, T. I. ISOLDE: a physically realistic environment for model building into low-resolution
electron-density maps. *Acta Crystallogr. Sect. D, Struct. Biol.* **74**, 519–530 (2018).
46. Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and analysis:
UCSF ChimeraX Visualization System. *Protein Sci.* **27**, 14–25 (2017).
47. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–
589 (2021).
48. Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular structure
solution. *Acta Crystallogr. Sect. D, Biol. Crystallogr.* **66**, 213–21 (2009).
49. Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular crystallography.
Acta Crystallogr. Sect. D: Biol. Crystallogr. **66**, 12–21 (2010).
50. Abraham, M. J. *et al.* GROMACS: High performance molecular simulations through multi-level
parallelism from laptops to supercomputers. *SoftwareX* **1**, 19–25 (2015).
51. Jo, S., Kim, T., Iyer, V. G. & Im, W. CHARMM-GUI: A web-based graphical user interface for
CHARMM. *J. Comput. Chem.* **29**, 1859–1865 (2008).

52. Huang, J. *et al.* CHARMM36m: an improved force field for folded and intrinsically disordered
proteins. *Nat. Methods* **14**, 71–73 (2017).
53. Khan, H. M., MacKerell, A. D. & Reuter, N. Cation-π Interactions between Methylated Ammonium
Groups and Tryptophan in the CHARMM36 Additive Force Field. *J. Chem. theory Comput.* **15**, 7–12
(2018).
54. Essmann, U. *et al.* A smooth particle mesh Ewald method. *J. Chem. Phys.* **103**, 8577–8593 (1995).
55. Hess, B., Bekker, H., Berendsen, H. J. C. & Fraaije, J. G. E. M. LINCS: A linear constraint solver for
molecular simulations. *Journal of Computational Chemistry* (1997).
56. Berendsen, H. J. C., Postma, J. P. M., Gunsteren, W. F. van, DiNola, A. & Haak, J. R. Molecular
dynamics with coupling to an external bath. *J. Chem. Phys.* **81**, 3684–3690 (1984).
57. Evans, D. J. & Holian, B. L. The Nose–Hoover thermostat. *J. Chem. Phys.* **83**, 4069–4074 (1985).
58. Parrinello, M. & Rahman, A. Polymorphic transitions in single crystals: A new molecular dynamics
method. *J. Appl. Phys.* **52**, 7182–7190 (1981).
59. Schrödinger & LLC. The PyMOL Molecular Graphics System, Version 2.0. (2015).
60. Humphrey, W., Dalke, A. & Schulten, K. VMD: Visual molecular dynamics. *J. Mol. Graph.* **14**, 33–
38 (1996).
61. Michaud-Agrawal, N., Denning, E. J., Woolf, T. B. & Beckstein, O. MDAnalysis: a toolkit for the
analysis of molecular dynamics simulations. *J. Comput. Chem.* **32**, 2319–27 (2010).
62. Sehnal, D. *et al.* MOLE 2.0: advanced approach for analysis of biomacromolecular channels. *J.*
Cheminformatics **5**, 39–39 (2013).
63. Tian, W., Chen, C., Lei, X., Zhao, J. & Liang, J. CASTp 3.0: computed atlas of surface topography of
proteins. *Nucleic Acids Res.* **46**, W363–W367 (2018).
64. Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence alignments
using Clustal Omega. *Mol. Syst. Biol.* **7**, 539–539 (2011).
751

Fig. 1: FLVCR1 and FLVCR2 are choline and ethanolamine transporters.

a, Confocal imaging shows that FLVCR1 and FLVCR2 are localized at the plasma membrane (arrows). Plasma

membrane GFP (mGFP) was used as a marker. **b**, Choline transport activities of human FLVCR1 and FLVCR2.

Choline kinase A (*CHKA*) was co-expressed with both proteins. **b,c**, Dose curves and time courses for choline

transport activities of human FLVCR1 and FLVCR2. *CHKA* was co-expressed with both proteins. **d**, Ethanolamine

transport activities of human FLVCR1 and FLVCR2. Ethanolamine kinase 1 (*ETNK-1*) was co-expressed with both

proteins. **e,f**, Dose curves and time courses for ethanolamine transport activities of human FLVCR1 and FLVCR2.

ETNK-1 was co-expressed with both proteins. The inactive S203Y mutant of FLVCR2 was used as a control in all

experiments. Each symbol represents one replicate. Data are expressed as mean ± SD. ****$P<0.0001$,

***$P<0.001$. ns, not significant. One-way ANOVA for transport activity measurement; Two-way ANOVA for dose

curve measurements.

Fig. 2: Architecture of FLVCR1 and FLVCR2 in their inward- and outward-facing states.

Cryo-EM density (top) and atomic model (middle and bottom) of apo inward-facing FLVCR1 (**a**) as well as FLVCR2

in the inward-facing (**b**) and outward-facing (**c**) states. The N- and C-domains are coloured in different shades of

blue and green for FLVCR1 and FLVCR2, respectively. A transparent cryo-EM density lowpass-filtered at 6 Å is

shown to visualize the detergent belt surrounding the transmembrane region. The sidechain of N265 in FLVCR1

and its N-linked glycan (grey) are shown as stick models. **d**, Cut-away views of the surface representation showing

the cavity shape of FLVCR2-IF^{apo} (left) and FLVCR2-OF^{apo} (right). Two central aromatic residues are shown as sticks.

e, Structural superposition of FLVCR2-OF^{apo} (dark green) and FLVCR2-IF^{apo} (light green).

Fig. 3: Cryo-EM structures of FLVCR1 and FLVCR2 in complex with choline.

a,b, Cryo-EM densities and atomic models of the choline-bound inward-facing FLVCR1 (FLVCR1-IF^choline) structure (**a**), and the choline-bound inward-facing FLVCR2 (FLVCR2-IF^choline) structure (**b**), respectively. The bound choline is shown as ball-and-stick model; binding site residues are shown as sticks. **c,d,** Time-resolved distance plots between choline and the highly-conserved tryptophan and tyrosine residues forming the choline-binding pockets of FLVCR1 (**c**) and FLVCR2 (**d**) obtained from MD simulation runs. A cation-π interaction is assumed for distance < 4 Å (grey dashed line). **e,** Choline transport activity of indicated FLVCR1 and FLVCR2 mutants. *CHKA* was co-expressed with all proteins. 20 μM [³H] choline was used for FLVCR1 mutants and 100 μM [³H] choline was used FLVCR2 mutants, respectively. Each symbol represents one replicate. Mutant transport activities were normalized to the total protein from cells and displayed as fold change with reference to mock (empty vector). **f,g,** Distance of choline to conserved aromatics for W125A^FLVCR1 (**f**) and W102A^FLVCR2 (**g**) as function of time in MD simulations (left), and choline occupancy in binding site for WT and alanine mutants (right). Error bars represent standard errors of the mean (s.e.m.) **h,** Shifts of tryptophan fluorescence of FLVCR1 and FLVCR2 in the presence of choline or betaine. **i,** Protein sequence alignment of choline-binding pocket residues (red block) in FLVCR1 and FLVCR2 across various mammalian species. Indicated residue numbers refer to FLVCR1 and FLVCR2 from *homo*

791   *sapiens*. Data shown are mean ± SD for (**e**) and (**h**), and mean ± standard error of the mean (s.e.m) for (**f**).

792   ****P<0.0001, ***P<0.001. One-way ANOVA for (**e**), and *t*-test for (**f**) and (**g**).

Fig. 4: Cryo-EM structures of FLVCR1 and FLVCR2 in complex with ethanolamine.

a, Cryo-EM density and atomic model of the ethanolamine-bound inward-facing FLVCR1 (FLVCR1-IF^ethanolamine)

structure. The bound ethanolamine is shown as ball-and-stick model; binding site residues are shown as sticks.

b, Distance of ethanolamine to conserved aromatics of FLVCR1 as function of time in MD simulation (left) and

snapshot (right). **c**, Frequency of distance between ethanolamine or choline and W125^FLVCR1 (top) or Y349^FLVCR1

(bottom) derived from Fig. 3c and 4b, respectively. **d**, Distance between Q214^FLVCR1 and the hydroxyl group of

ethanolamine (orange) or choline (pink) as function of time in MD simulation, with corresponding frequency

(right). **e**, Ethanolamine transport activities of indicated FLVCR1 and FLVCR2 mutants. *ETNK-1* was co-expressed

with all proteins. 2.5 µM [^{14}C] ethanolamine was used. Transport activities of mutant variants were normalized

to the total protein from cells and are displayed as fold change to mock (empty vector). Each symbol

represents one replicate. Data shown are mean ± SD. ****P<0.0001. One-way ANOVA.

Fig. 5: Cryo-EM structures of FLVCR2 in complex with heme.

a, Import assay of FLVCR1 and FLVCR2 for heme. 2.5 µM [³H] hemin was used. The inactive mutant of FLVCR2^S203Y

was used as a control. **b**, Cryo-EM density (left) and ribbon model (right) of the heme-bound outward-facing

FLVCR2 (FLVCR2-OF^heme). The heme molecule is shown in ball-and-stick representation, and the corresponding

cryo-EM density is coloured in purple. Right panels are the close-up views of the heme-binding site in FLVCR2-

OF^heme (top) and in FLVCR2-OF^apo (bottom) with locally-filtered cryo-EM density. Residues in close proximity to

heme are shown as sticks. A lipid molecule (grey ball-and-stick) is fitted into the lipid-like density at the heme-

binding site of FLVCR2-OF^apo. **c**, Distance of heme and interacting residues as function of time in MD simulations

(left), with a snapshot of FLVCR2-heme interactions (right) indicating the membrane layer by the phosphates of

POPE/POPG (red spheres). **d**, Distances as in (**c**, left), but for mutants R82A/R83A (top) and K273A (bottom). **e**,

Interaction frequency of heme with four heme-interacting residues for WT and mutants of FLVCR2. **f**, Cryo-EM

density of the heme-bound, choline-bound, inward-facing FLVCR2. The cryo-EM density corresponding to heme

is coloured in purple. Right panels are the cryo-EM density of the substrate-binding site (top) and heme-binding

site (bottom) in FLVCR2-IF^heme-choline with rigid-body fitting of the FLVCR2-IF^choline model. The heme-binding site is

shown with locally-filtered cryo-EM density. **g**, Choline (left) and ethanolamine (right) transport assays for

FLVCR2 heme-binding site mutants. *CHKA* or *ETNK-1* was co-expressed with WT or *FLVCR2* mutant plasmids. 100

μM [^{3}H] choline or 2.5 μM [^{14}C] ethanolamine was used, respectively. Transport activity of the mutants was

normalized to the total protein and is displayed as fold change to mock (empty vector). **h**, Choline (left) and

ethanolamine (right) transport assay for FLVCR2 in the presence of heme. For choline + hemin experiment, 10

μM [^{3}H] choline and 50 μM hemin were used, while for ethanolamine + hemin experiment, 2.5 μM [^{14}C]

ethanolamine and 25 μM hemin were used. The experiment was repeated twice for (**a**). One dataset was shown.

Each symbol represents one replicate. Data are expressed as mean ± SD for (**a**), (**g**) and (**h**), and mean ± s.e.m for

(**e**). ****$P<0.0001$, **$P<0.01$, *$P<0.05$. ns, not significant. One-way ANOVA for (**a**) and (**g**), two-way ANOVA for

(**h**), and *t*-test for (**e**).

Fig. 6: Proposed model for choline transport by FLVCR2.

Schematic illustration of FLVCR2 conformations during the choline transport cycle. Green-colored

states represent experimentally obtained conformations of this study. States colored in grey are

hypothesized based on knowledge about the commonly characterized alternative-access mechanism

of MFS transporters.

Extended Data Fig. 1: Transport activity of FLVCRs for choline or ethanolamine without

overexpression of *CHKA* or *ETNK-1*.

Choline (**a**) or ethanolamine (**b**) transport activity of FLVCR1, FLVCR2, and FLVCR2^{S203Y} without

simultaneous overexpression of *CHKA* or *ETNK-1*, respectively. 20μM [^{3}H] choline or 2.5μM [^{14}C]

ethanolamine was used, respectively. The inactive mutant of FLVCR2^{S203Y} was used as a control in all

experiments. Each experiment was repeated twice and one dataset was shown. Each symbol

represents one replicate. Data are expressed as mean ± SD. ****P<0.0001, ***P<0.001, **P<0.01. ns,

not significant. One-way ANOVA.

Extended Data Fig. 2: Metabolomic analysis of livers from *FLVCR1*-knockout mice.

a, Illustration of experimental procedures. *FLVCR1* was deleted by polyIC injection in *FLVCR1*^f/f^-Mx1-Cre (knockout) mice. Liver samples of control (*FLVCR1*^+/+^-Mx1-Cre) and knockout mice were collected at least 4 weeks post-injection for metabolomic analysis. **b,** Levels of choline and choline metabolites from control and knockout mice. **c,** Levels of ethanolamine metabolites from control and knockout mice. Each data point represents one mouse. Data are expressed as mean ± SD. ****P<0.0001, ***P<0.001, **P<0.01, *P<0.05. ns, not significant. *t*-test.

Extended Data Fig. 3: Transport properties of FLVCR1 and FLVCR2.

a, Ethanolamine transport activity of FLVCR1 at indicated pH values, and in sodium-free buffer (K⁺). **b,**

Choline transport activity of FLVCR2 at indicated pH values, and in sodium-free buffer (K⁺). The right

panels show normalized data with reference to the respective mocks. In these experiments, 2.5 µM

[¹⁴C] ethanolamine was used for FLVCR1 or 20 µM [³H] choline for FLVCR2, respectively. The cells were

co-expressed with *ETNK-1* or *CHKA* and incubated with the ligands at 37 °C for 15 mins. **c,** Export assays

of FLVCR1 with ethanolamine (left) and FLVCR2 with choline (right). In these assays, 100 µM [¹⁴C]

ethanolamine or 200 µM [³H] choline was incubated with *FLVCR1* or *FLVCR2* overexpression cells,

respectively. After 2 hours of incubation, the buffer was washed out. Intracellular [³H] choline or [¹⁴C]

866   ethanolamine from the cells was allowed to release into choline/ethanolamine-free medium for 1 hour.

867   The radioactive signal in the cells was normalized to the total protein and expressed as fold change to

868   mock. Each symbol represents one replicate. Data are expressed as mean ± SD. ****$P<0.0001$,

869   ***$P<0.001$, *$P<0.05$. ns, not significant. Two-way ANOVA for (**a**) and (**b**), One-way ANOVA for (**c**).

Extended Data Fig. 4: Architecture and structural comparison between FLVCR1 and FLVCR2.

a, Schematic diagram of FLVCR family showing the topology of the secondary structure. Motifs that

are not observed in both cryo-EM structures of FLVCR1 and FLVCR2 are shown as dashed lines. **b**,

Structural comparison of FLVCR1-IF^{apo} and FLVCR2-IF^{apo} in tube representation, viewed from the lipid

bilayer (left), extracellular side (top-right), and intracellular side (bottom-right).

 Extended Data Fig. 5: Residues for intra- and extracellular gating in FLVCR1 and FLVCR2.

 a, Cut-away view of FLVCR1-IF[apo] in surface representation showing the cytoplasmic cavity. Two central

 aromatic residues are shown as sticks. The structures of FLVCR2-IF[apo] and FLVCR2-OF[apo] are shown for

 comparison. Cross-sections of their inter-domain interactions are shown from the extracellular side

 (top), or from the intracellular side (bottom-left). Residues in FLVCR1 corresponding to the inter-

 domain interaction residues in FLVCR2 are shown. The bottom-right panel shows the inter-domain

 interactions between H1 and H3 in FLVCR2-OF[apo] viewed from the intracellular side. Residues

participating in the inter-domain interactions are shown as sticks; hydrogen bonds and salt bridges are

labelled with dashed lines. **b**, Transport assay of FLVCR2 mutants for choline (left) and ethanolamine

(right). *CHKA* or *ETNK-1* was co-expressed with wild-type *FLVCR2* and mutant plasmids. In these assays,

100 μM [^{3}H] choline or 2.5 μM [^{14}C] ethanolamine was used, respectively. **c**, Transport assay of FLVCR1

mutants for choline (left) and ethanolamine (right). *CHKA* or *ETNK-1* was co-expressed with wild-type

FLVCR1 and mutant plasmids, and 20 μM [^{3}H] choline or 2.5 μM [^{14}C] ethanolamine was used,

respectively. Each symbol represents one replicate. Data shown are mean ± SD. ****$P<0.0001$. ns, not

significant. One-way ANOVA.

Extended Data Fig. 6: Physicochemical properties of FLVCR structures and conservation analyses.

Surface charge, hydrophobicity, and conservation analyses of FLVCR1-IF^apo (**a**), FLVCR1-IF^choline (**b**),

FLVCR2-IF^apo (**c**), FLVCR2-OF^apo (**d**), and FLVCR2-IF^choline (**e**). From left to right: Surface viewed from the

lipid bilayer and both C and N domains viewed from the domain interface coloured by electrostatic

potential, the domain interface coloured by hydrophobicity, and the domain interface coloured by

sequence conservation. The cavity in the different states of both FLVCRs is outlined with dashed lines.

902   Two important central pocket residues (W125 and Y349 in FLVCR1; W102 and Y325 in FLVCR2) are

903   indicated. The FLVCR2 binding site for heme is marked by black-and-white circles.

Extended Data Fig. 7: Conformational changes of substrate-bound FLVCRs.

Structural comparison of FLVCR2-IF^{choline} and FLVCR2-IF^{apo} (**a**), FLVCR1-IF^{choline} and FLVCR1-IF^{apo} (**b**), as

well as FLVCR1-IF^{choline} and FLVCR1-IF^{ethanolamine} (**c**), viewed from the lipid bilayer (left), extracellular side

(top-right), and intracellular side (bottom-right).

Extended Data Fig. 8: Interaction chemistries and dynamics between Q214F[LVCR1] and respective

ligands..

a, Structural comparison of the binding site of ethanolamine- and choline-bound FLVCR1. **b**, Time-

resolved distance plots between Q214 and the primary amine of ethanolamine (orange) or the tertiary

amine of choline (pink) obtained from MD simulation runs. The right panel shows the accumulative

frequency of their distances during the simulations.

Extended Data Fig. 9: Translocation pathway of choline in FLVCRs.

a, Choline-binding sites of FLVCR2-OF[apo] (top) and FLVCR2-IF[choline] (bottom) with the distance between

W102 and Y325 shown as dashed lines. **b**, Cut-away views of FLVCR1-IF[choline] (top-left), FLVCR1-

IF[ethanolamine] (top-middle) and FLVCR2-IF[choline] (top-right) showing the inward-facing cavity. Two central

aromatic residues are shown as sticks. The surfaces shown below are the respective models viewed

from the intracellular side. The surfaces of IF[apo] of FLVCR1 and FLVCR2 are also shown for comparison.

The dashed circles indicate the peripheral channel in the ligand-bound state.

928