Genomic consequences of isolation and inbreeding in an island dingo population

Ana V. Leon-Apodaca[1], Manoharan Kumar[2], Andres del Castillo[1], Gabriel C. Conroy[3,4], Robert W Lamont[3,4], Steven Ogbourne[4], Kylie M. Cairns[5,6], Liz Borburgh[3], Linda Behrendorff[7], Sankar Subramanian[2,4,*], Zachary A. Szpiech[1,8,*]

[1] Department of Biology, Pennsylvania State University, PA, USA

[2] School of Science, Technology & Engineering, University of the Sunshine Coast, 1 Moreton Parade, Petrie, Queensland, Australia

[3] School of Science, Technology & Engineering, University of the Sunshine Coast, 90 Sippy Downs Drive, Sippy Downs, Queensland, Australia

[4] Centre for Bioinnovation, University of the Sunshine Coast, 90 Sippy Downs Drive, Sippy Downs, Queensland, Australia

[5] Evolution & Ecology Research Centre, School of Biological, Earth and Environmental Sciences, UNSW Australia, Sydney NSW 2052, Australia

[6] Centre for Ecosystem Science, School of Biological, Earth and Environmental Sciences, UNSW Australia, Sydney NSW 2052, Australia

[7] Queensland Parks and Wildlife Service, Department of Environment & Science, K'gari, Australia

[8] Institute for Computational and Data Sciences, Pennsylvania State University, PA, USA

*Corresponding authors: zps5164@psu.edu, ssankara@usc.edu.au

ABSTRACT

Dingoes come from an ancient canid lineage that originated in East Asia around 8000-11,000 years BP. As Australia's largest terrestrial predator, dingoes play an important ecological role. A small, protected population exists on a world heritage listed offshore island, K'gari (formerly Fraser Island). Concern regarding the persistence of dingoes on K'gari has risen due to their low genetic diversity and elevated inbreeding levels. However, whole-genome sequencing data is lacking from this population. Here, we

include five new whole-genome sequences of K'gari dingoes. We analyze a total of 18

whole genome sequences of dingoes sampled from mainland Australia and K'gari to

assess the genomic consequences of their demographic histories. Long (>1 Mb) runs of

homozygosity (ROH) — indicators of inbreeding — are elevated in all sampled dingoes.

However, K'gari dingoes showed significantly higher levels of very long ROH (>5 Mb),

providing genomic evidence for small population size, isolation, inbreeding, and a strong

founder effect. Our results suggest that, despite current levels of inbreeding, the K'gari

population is purging strongly deleterious mutations, which, in the absence of further

reductions in population size, may facilitate the persistence of small populations despite

low genetic diversity and isolation. However, there may be little to no purging of mildly

deleterious alleles, which may have important long-term consequences, and should be

considered by conservation and management programs.

SIGNIFICANCE

A long-standing question in conservation genetics is whether long-term isolation and

elevated levels of inbreeding always leads to inevitable population extinction. Here we

conduct the first-ever whole-genome analysis of a population of dingoes living in long-

term isolation on an island off the coast of Australia (K'gari). We show that these

animals are beset by very low genetic diversity, likely the result of extensive inbreeding,

and an elevated number of deleterious homozygotes. However, our results suggest that

these dingoes are likely purging highly deleterious alleles, which may have allowed

them to persist long term despite their extremely small population (<200 individuals).

INTRODUCTION

Dingoes have a complex history in Australia, as they have been simultaneously

declared a native species and an agricultural pest dependent on jurisdiction and

context. While they are protected in parts of the Australian Capital Territory, Victoria,

Queensland and the Northern Territory as a native species, 'wild dogs', including

dingoes, have been declared a pest or restricted matter in all Australian states and

territories for posing a risk to livestock (Brink et al. 2019).

The dingo plays an important ecological role in maintaining the ecosystems of Australia,

as they have been the apex terrestrial predator since the extinction of the thylacine

(Glen et al. 2007; Johnson et al. 2007; Letnic et al. 2009; Wallach et al. 2010; Letnic et

al. 2011; Letnic et al. 2012). Research using mtDNA and Y-chromosome data has

identified a close relationship between dingoes, New Guinea singing dogs (NGSD), and

Asian village dogs, suggesting that the ancestors of dingoes originated in East Asia

(Savolainen et al. 2004; Oskarsson et al. 2012; Ardalan et al. 2012; Sacks et al. 2013;

Cairns and Wilton 2016; Cairns et al. 2017). Archeological evidence and genomic data

estimate the divergence time between wolves and early dogs to be ~15,000-36,000

years BP (Germonpré et al. 2009; Germonpré et al. 2012; Larson et al. 2012; Skoglund

et al. 2015; Frantz et al. 2016; Freedman and Wayne 2017), and ~5,000-11,000 years

BP between early dogs and dingoes (Savolainen et al. 2004; Oskarsson et al. 2012;

Sacks et al. 2013; Cairns and Wilton 2016; Bergström et al. 2020; Surbakti et al. 2020;

Cairns 2021; Field et al. 2022). Although dingoes are descended from a dog-like

ancestor, they form a distinct and separate lineage from domestic dogs, wolves and

other canids (Crowther et al. 2014; Smith et al. 2019; Cairns 2021; Field et al. 2022).

At least two distinct dingo lineages have been identified in the mainland, one in the

northwest and one in the southeast, alongside one offshore lineage on K'gari (formerly

known as Fraser Island) (Cairns and Wilton 2016; Cairns et al. 2017; Cairns et al. 2018;

Greig et al. 2018; Zhang et al. 2020). K'gari is located off the coast of Queensland,

Australia, and has been listed as a UNESCO World Heritage site since 1992. In addition

to being the world's largest sand island, it is also home to a small, protected (Nature

Conservation Act 1992) population of dingoes of ~70-170 individuals that are genetically

distinct from mainland dingoes (Brown et al. 2011; Conroy et al. 2021). In contemporary

times dingoes are known for being an infamous tourist attraction for the over 400,000

tourists that visit K'gari each year (Wardell-Johnson et al. 2015), which has led to

human-wildlife conflict (O'Neill et al. 2017; Behrendorff 2021). In response to these

incidents, management strategies have been implemented throughout the island,

including non-lethal options as well as the removal of dingoes via lethal control,

although the latter is currently a last resort action and only occurs around high-use

visitor areas (Behrendorff 2021). K'gari dingoes are of particular interest to

conservationists, as well as various stakeholders in Australia, as there are concerns

regarding their persistence due to low genetic diversity and inbreeding given it is a

small, isolated population (Cairns et al. 2018; Conroy et al. 2021).

Isolated populations often experience reduced fitness resulting from increased levels of

homozygosity and a high load of deleterious mutations across their genomes. There

have been several instances in which these factors have led to inbreeding depression

and a high risk of extinction. Such is the case of the iconic Isle Royale wolves, which

are highly inbred gray wolves located on an island in Lake Superior, in northern United

States (Hedrick et al. 2019; Robinson et al. 2019). Other examples of isolated

populations in the wild that have experienced inbreeding depression are Florida

Panthers (Roelke et al. 1993) and Scandinavian gray wolves (Liberg et al. 2005;

Räikkönen et al. 2006; Kardos et al. 2018). In other instances, small, isolated

populations have managed to persist with low genetic diversity for generations, without

experiencing inbreeding depression. Such is the case of Channel Island foxes

(Robinson et al. 2016; Robinson et al. 2018), Appenine brown bears (Benazzo et al.

2017), African mountain gorillas (Xue et al. 2015), Alpine ibex (Grossen et al. 2020),

and Isle Royale moose (Kyriazis et al. 2023). It has been suggested that the purging of

strongly deleterious, recessive mutations may facilitate the persistence of long-term

small populations (Xue et al. 2015; Robinson et al. 2018; Grossen et al. 2020; Wootton

et al. 2023).

Only a handful of studies have analyzed the genetics of K'gari dingoes. While these

studies have found low genetic diversity and high levels of inbreeding on K'gari dingoes

(Cairns and Wilton 2016; Cairns et al. 2018; Conroy et al. 2021), more research is

needed to understand the consequences of inbreeding and isolation on this population.

Moreover, there is a need for new and well-informed conservation strategies for dingoes

in Australia, given that current strategies do not take into account these genetic factors.

To our knowledge, there is only one whole-genome sequence from a K'gari dingo

individual that is publicly available (Zhang et al. 2020). In this work we include five newly

sequenced samples, and we look further into the genetics of K'gari dingoes to

understand the consequences of their demographic history on the distribution of

deleterious variation, and the implications of small population sizes and isolation, to help

inform conservation efforts to ensure their long-term persistence.

RESULTS

Canid Sampling, Relatedness and Population Structure

Previous analysis on dingoes suggest there are at least three distinct lineages of

dingoes in Australia, with one being located on the northwest of the mainland, another

in the southeast of the mainland, and the third one offshore on K'gari (Fraser Island)

(Cairns and Wilton 2016; Cairns et al. 2017; Cairns et al. 2018). To expand our

understanding of genetic diversity patterns in the K'gari population, we generated high-

quality, whole-genome sequences of five dingoes with average read depth between

12.01X and 12.56X sampled from the island between 2011 and 2013. Although there is

a dingo reference genome (ASM325472v1), the newly sequenced samples in this study

were aligned to the dog reference genome (CanFam3) to take advantage of its fully

assembled chromosomes and better gene annotations. We include previously published

whole-genome sequences from dingoes from northwest and southeast mainland

Australia, from wolves, village dogs, and New Guinea singing dogs (NGSD), which are

a close relative of dingoes. All canid reads in this study were aligned to the dog

reference genome (CanFam3).

We identified close relatives in K'gari dingoes in two ways. First, we looked at the

proportion of sites at which pairs of individuals share 0, 1 or 2 alleles that are identical-

by-state (IBS). We expect Parent/Offspring to share at least one allele at a locus

identical-by-descent, thus having low values of IBS0. We expect full-siblings to share 2

alleles identical-by-descent for approximately 25% of loci, and we expect second-
degree relatives and unrelated individuals to have higher estimates of IBS0. We also
used the KING algorithm (Manichaikul et al. 2010) to infer kinship coefficients for all
dingo individuals. We identified three pairs of Parent/Offspring relatives, four pairs of
full-siblings relatives, a pair of avuncular relatives, and seven pairs that include third-
degree relatives and unrelated individuals. Furthermore, we found that none of the
K'gari dingo individuals are closely related to dingo individuals from the mainland, and
that the mainland dingoes in this study are not closely related to one another
(supplementary fig. S1; supplementary table S1).

We computed pairwise allele sharing dissimilarities to assess population structure and
generated multidimensional scaling (MDS) plots from the dissimilarity matrix. In MDS,
individuals with small allele sharing dissimilarities tend to group together. In fig. 1A we
can observe distinct taxonomic groupings, with the close relationship between the dingo
and NGSD lineages clearly captured in this analysis. Individuals within the wolf and
village dog populations were sampled from different regions around the world (see
supplementary table S2), hence the relatively loose clustering among these individuals.
Interestingly, we find that the village dog from Vietnam seems to be genetically more
similar to mainland dingoes than the village dog from Papua New Guinea (fig. 1A).
While Papua New Guinea is geographically closer to Australia than Vietnam, similar
observations regarding shared ancestry between Vietnamese village dogs and dingoes
have been reported by Cairns et al. (Cairns et al. 2023). A possible explanation
regarding the observed genotype differences among these canids may be explained by
Vietnamese village dogs having retained the ancestral alleles that also exist in the dingo

lineage while village dogs from Papua New Guinea have not, possibly because of their

different demographic histories or genetic swamping from contemporary domestic dogs.

Although dingoes from the southeast and northwest overlap with each other (i.e., they

are genetically more similar), we observe K'gari dingoes form their own separate

subgroup (fig. 1A). To look more into detail at the allele sharing differences between the

three dingo populations, we recomputed the MDS on the matrix of allele sharing

dissimilarities containing only these samples. In fig. 1B we can observe geographic

based clustering, with K'gari dingoes maintaining a tighter and more distinct group

compared to dingoes in the southeast and northwest.

Genetic diversity between and within canid populations

To characterize genetic diversity patterns within and between canid populations we

analyzed the distribution of alleles across populations. We used ADZE (Szpiech et al.

2008) to compute estimates of allelic richness, private allelic richness, and the number

of alleles shared by pairs of populations (i.e., alleles that are found only in those

populations and nowhere else). We observed reduced levels of genetic diversity in

mainland and K'gari dingoes, compared to large and outbred populations such as

wolves and village dogs, with K'gari dingoes having the lowest genetic diversity of all

the canid groups analyzed in this study (fig. 2A; fig. 2B). Furthermore, we found

mainland dingoes share more alleles with village dogs than K'gari dingoes and village

dogs (fig. 2C). Although hybridization with domestic dogs was previously highlighted as

a main concern for dingo conservation (Newsome and Corbett 1985; Jones 1990;

Corbett 2001; Wilton 2001; Elledge et al. 2008; Stephens et al. 2015), more recent

studies have shown hybridization is uncommon (Cairns et al. 2020; Cairns 2021; Cairns

et al. 2021; Cairns et al. 2023). This suggests that the alleles shared between dingoes

and their common ancestors from Southeast Asia were likely lost in the bottleneck that

founded the K'gari dingo population.

Inbreeding and homozygosity

To characterize the levels of inbreeding we looked at long contiguous stretches of

homozygous regions in individual genomes that are shared identical-by-descent (IBD),

also known as runs of homozygosity (ROH). We used GARLIC (Szpiech et al. 2017) to

call ROH, which implements a likelihood-based approach to infer ROH. The length of

ROH can inform on how recent the parental relatedness among individuals is, with long

ROH more commonly arising in offspring descended from parents with a recent

common ancestor. On the other hand, short ROH suggests a more distant common

ancestor, as recombination has had more time to break apart the underlying IBD

haplotypes (Kirin et al. 2010; Pemberton et al. 2012). The wolf and village dog

individuals included in this study were sampled from large and outbred populations from

different regions around the world (supplementary table S2), resulting in variation in the

levels of homozygosity found in their individual genomes. Overall, we observed

relatively low levels of homozygosity in village dogs and wolves, with ~7% and ~16% of

their genomes being covered by ROH >0.5 Mbp, respectively (fig. 3). In contrast, we

observed high levels of homozygosity in northwest and southeast mainland dingo

populations, with ~30% of their genomes being covered by ROH >0.5 Mbp. In K'gari

dingoes we observed even higher levels of homozygosity, with ~65% of their genomes

being covered by ROH >0.5 Mbp, and ~28% by ROH >5 Mbp (fig. 3), with the longest

individual homozygous segment being 64.2 Mbp long (supplementary fig. S3).

We estimated inbreeding coefficients as given by ROH for each population (F_{ROH}),

which represents the proportion of the autosomal genome covered by ROH >1Mbp.

Mean F_{ROH} of 0.59, 0.23, 0.23, were estimated for K'gari, Northwest, and Southeast

dingoes, respectively, and 0.35 for NGSD. For village dogs and wolves we estimated

mean F_{ROH} of 0.04 and 0.13, respectively. These results provide clear evidence for

inbreeding, isolation, small population size, and a strong founder effect in K'gari

dingoes.

Patterns of deleterious variation and genetic load

To understand the effect of inbreeding and low genetic diversity on the distribution of

deleterious variation, we looked at the burden of deleterious alleles in K'gari dingoes

compared to mainland dingoes and other canids. Alternate alleles were predicted to be

deleterious or neutral by taking into account sequence conservation and the effect of an

amino acid substitution on protein function using Ensembl VEP (McLaren et al. 2010)

and SIFT (Vaser et al. 2016) (see Methods). We found dingoes and NGSD carry a

higher number of deleterious alternate-allele homozygotes compared to village dogs

and wolves, although K'gari dingoes carry the highest number of deleterious alternate-

allele homozygotes among all the canids included in this study (fig. 4A). Conversely, we

observed K'gari dingoes carry the lowest number of deleterious and LoF alternate-allele

heterozygotes compared to the other canids (fig. 4B; fig. 4D). Furthermore, we

observed a significant difference in the number of alternate-allele LoF homozygotes,

with K'gari dingoes carrying a higher number than northwest and southeast mainland

dingoes (fig. 4C).

To refine our analysis of the differences in the burden of deleterious alleles between

K'gari and mainland dingoes, we subset our deleterious and LoF alternate-allele calls to

include only those alternate alleles which were also inferred to be the derived allele in

dingoes. We observed patterns of deleterious derived homozygotes and heterozygotes

stayed qualitatively similar to the non-polarized patterns (supplementary fig. S4A;

(supplementary fig. S4B). For the LoF category, we did not find a significant difference

in the number of derived homozygotes in K'gari and mainland dingoes (supplementary

fig. S4C), however, the mean number of LoF derived heterozygotes in K'gari dingoes is

significantly lower than in mainland dingoes (supplementary fig. S4D). Similar patterns

have been observed in other small populations with substantial inbreeding, suggesting

purging of LoF alleles, which are more likely to be strongly recessive variants than

small-effect deleterious variants, may facilitate the persistence of small and isolated

populations (Robinson et al. 2018; Grossen et al. 2020).

To test whether K'gari dingoes showed evidence of purging of deleterious variants we

used the Rxy statistic, as implemented by Xue et al. (Xue et al. 2015), which compares

the frequency of derived alleles of a particular category (e.g., deleterious) normalized by

the frequencies of a set of putatively neutral variants, which is then compared across

two populations, X and Y. When Rxy < 1 for a given site category, we consider this

evidence of purging of those alleles in population X relative to Y, when Rxy > 1 we

consider this evidence of enrichment of those alleles in population X relative to Y. When

Rxy = 1, we consider this evidence of no difference of those alleles between X and Y.

As our K'gari samples are closely related, we implemented a modified version of the

Rxy statistic, which accounts for relatedness among K'gari dingoes (see Methods). For

this analysis we looked at several allele categories: deleterious, tolerated, LoF,

synonymous and nonsynonymous sites, and a random set of intergenic sites as a

control. All Rxy calculations used an additional, independent, set of intergenic sites for

normalization. We estimated the excess number of deleterious variants in K'gari

dingoes (population X) with respect to mainland dingoes (population Y) and found

evidence of purging of LoF alleles (i.e., putative strongly deleterious alleles), but little to

no evidence of purging of mild/slightly deleterious alleles in K'gari dingoes (fig. 5).

Although we observe a small enrichment in the rest of the categories, we hypothesize

these results may be due to the small sample size and low genetic diversity in K'gari

dingoes compared to mainland dingoes. To assess the effect of sample sizes on our

results, we conducted this analysis using the same number of samples for each dingo

population and found that results stayed very similar (supplementary fig. S5).

DISCUSSION

Inbred populations frequently experience inbreeding depression and subsequently face

a high risk of extinction. However, some populations are able to persist in small

numbers and with low genetic diversity (Xue et al. 2015; Benazzo et al. 2017; Robinson

et al. 2018; Grossen et al. 2020). Therefore, understanding the complex relationship

between genetic diversity, inbreeding, genetic load, and demographic history are key to

leveraging this information for management and conservation strategies of populations

in the wild.

In large and outbred populations, strongly deleterious recessive alleles are "hidden"

from selection in heterozygous form, making large populations more prone to

experiencing inbreeding depression after going through a harsh bottleneck (Robinson et

al. 2018; Kyriazis et al. 2021). On the other hand, small and isolated populations tend to

have a higher burden of deleterious variants as a result of inbreeding (Robinson et al.

2018; Hedrick et al. 2019; Grossen et al. 2020; Khan et al. 2021; Stoffel et al. 2021).

Accumulation of long ROH in the genome, indicators of inbreeding, is associated with

increased presence of deleterious homozygotes and potential fitness effects (Szpiech et

al. 2013; Szpiech et al. 2019; Stoffel et al. 2021; Steux and Szpiech 2023; Swinford et

al. 2023).

In this work, we have generated five whole-genome sequences of K'gari dingoes to

investigate the consequences of isolation and inbreeding on this island population. We

find K'gari dingoes have lower genetic diversity and higher levels of runs of

homozygosity compared to dingoes in the mainland and canids from large populations.

Furthermore, we find a higher burden of deleterious homozygotes in K'gari dingoes

compared to canids from large and stable populations, such as wolves and village dogs

(fig. 4A).

Although the levels of ROH and estimated inbreeding found in K'gari dingoes (F_{ROH} =

0.59 on average; fig. 3) in some cases surpass those observed in wild populations of

other species already experiencing inbreeding depression (supplementary table S3)

(Kardos et al. 2018; Robinson et al. 2019; Khan et al. 2021; Stoffel et al. 2021; Taron et

al. 2021; Colpitts et al. 2022), surveillance and necropsy records from the Queensland

Parks and Wildlife Service (QPWS) and the latest report on the genetic health of K'gari

dingoes (Allen 2023) have not recorded abnormal phenotypes in K'gari dingoes.

Nonetheless, whether a small population will experience inbreeding depression or not

depends on various factors, including the size of the ancestral population and the

magnitude and history of bottleneck events (Kyriazis et al. 2021).

It has been observed that long-term small populations (i.e., populations that

experienced early bottlenecks and have maintained small population size following the

bottlenecks) can persist despite showing high levels of inbreeding by purging strongly

recessive deleterious variants (Xue et al. 2015; Benazzo et al. 2017; Robinson et al.

2018; Grossen et al. 2020; Dussex et al. 2021), whereas populations that have gone

through multiple recent bottlenecks will most likely experience inbreeding depression

despite purging (Khan et al. 2021; Stoffel et al. 2021). At least two bottlenecks in the

form of large-scale culling events have been documented during the past 120 years

(Petrie 1995; Allen et al. 2015). Moreover, according to surveillance records from

QPWS, dingoes in K'gari have maintained a small population size (<200 individuals) for

the past 30 years, and evidence of inbreeding depression has not been observed. Our

assessment of purging in this population suggests that while strongly deleterious alleles

are likely being purged, there may be little to no purging of mildly deleterious alleles in

this population. We hypothesize purging of strongly deleterious alleles may have played

a role in the persistence of K'gari dingoes, as there is no current overt evidence of

inbreeding depression despite the population having experienced recent severe

bottlenecks. However, the long-term viability of this population is uncertain given its

small population size and genetic isolation.

Current conservation and management programs treat dingoes as a single

homogeneous population, although previous research has identified at least three

distinct dingo populations (Savolainen et al. 2004; Ardalan et al. 2012; Oskarsson et al.

2012; Sacks et al. 2013; Cairns and Wilton 2016; Cairns et al. 2017; Greig et al. 2018;

Zhang et al. 2020; Cairns et al. 2023). Adding to previous evidence, we find two distinct

populations in the mainland (northwest and southeast), and one offshore in K'gari.

Cairns et al. (Cairns et al. 2017) found dingoes in K'gari share the paternal haplogroup

from northwest mainland dingoes but carry the maternal lineage from dingoes in the

southeast. Consistent with these findings, we identify long IBD segments shared

between dingoes from K'gari and northwest Australia, and between dingoes from K'gari

and the southeast (supplementary fig. S2). In light of these findings, recognizing several

dingo subpopulations may help inform conservation strategies.

This work is the first step in broadening our knowledge and understanding of the

landscape of genetic diversity and deleterious variation in K'gari dingoes in order to

better inform conservation and public policies. Our work, although limited by a small

sample size and the sampling of several closely related samples, uses standard

genomic analysis tools and indicates that the dingo population in K'gari is challenged by

inbreeding as a result of founder effects and successive bottlenecks. We find the first

evidence that K'gari dingoes may be purging strongly deleterious mutations, whilst

accumulating a higher genetic load of mild/slightly deleterious mutations. Further

research is needed to investigate the evolutionary history and genetic load of dingoes in

K'gari using a larger sample size, which in turn, would allow us to incorporate

deleterious variation to conduct genomic simulations to learn more about the future

health of this population. In addition to this, dedicated monitoring for phenotypic or

reproductive abnormalities such as reduced fertility, increased stillbirths, and congenital

birth defects is recommended, as biological monitoring for symptoms of inbreeding

depression is currently understudied in K'gari dingoes. QPWS and Butchulla Prescribed

Body Corporates are left with the challenge of preserving dingoes in K'gari, balancing

the need to address human safety concerns, and protecting the dingo population's

genetic health and viability into the future.

MATERIALS AND METHODS

Sampling and sequencing

In this study we analyze whole-genome sequencing data from a total of 41 canids,

including 18 dingo samples from both K'gari (6) and mainland Australia (4 from the

Southeast, and 8 from the Northeast). We also include 2 New Guinea singing dog

samples, 11 gray wolf samples, 1 red wolf sample, and 9 village dog samples

(supplementary table S2). We generated whole-genome data for five of the six K'gari

dingo individuals analyzed in this study. Biological samples from these five individuals

were collected between 2011 and 2013 from regions along the east coast of K'gari,

including Cathedral Beach, Eurong, and Dundubara, and were sequenced using

Illumina Hiseq 2000. Newly sequenced raw reads from K'gari dingoes and publicly

available NCBI SRA raw reads were trimmed based on quality and adapter sequences

using Trimmomatic v0.36 (Bolger et al. 2014) using the following parameters:

ILLUMINACLIP:adapter.fa:2:30:10 SLIDINGWINDOW:4:20 MINLEN:5. Reads from all

41 canids were aligned to the CanFam3 reference genome using BWA-MEM v0.7.13 (Li

and Durbin 2009) with parameters set as follows: -M -t 8 -r

"ID:sampleID\tSM:sampleName". Default parameters were used to index the reference

genome prior to alignment. Mapped reads were converted from sequence

alignment/map (SAM) format to binary alignment/map (BAM) format using the view

command of samtools v1.3.1 (Danecek et al. 2021) and the following parameters: -@ 8

-b -F 4. The resulting BAM file was sorted based on chromosomes and alignment

position with parameters set as follows: -@ 10 -o, and subsequently, indexed using

default parameters. The sorted reads were used for marking PCR duplicates using the

MarkDuplicates tool of the Picard toolset v2.25.2 (Broad Institute 2019) with default

parameters.

Genotype calling, filtering, and phasing

All samples were jointly genotyped with the Genome Analysis Toolkit (GATK) v4.2

(McKenna et al. 2010). GATK HaplotypeCaller was used to call genomic variants with

parameters set as follows: -Xmx5g -Xms5g -XX:ParallelGCThreads=4 -ERC GVCF.

Samples were merged using GATK CombineGVCFs with the following parameters: -

Xmx5g -Xms5g -XX:ParallelGCThreads=4. Lastly, GATK GenotypeGVCFs was used

with default parameters to call the variants.

The resulting VCF file containing raw genotype calls for all 41 canids was then filtered

using BCFtools v1.18 (Danecek et al. 2021) in the following way: Only sites with biallelic

snps and only sites with a site-level quality score (QUAL > 50) were retained. Any

individual genotype with (GQ <= 19) was set to missing. All sites with >= 10% missing

data were removed. All sites with no variation among the 41 samples (i.e.,

monomorphic sites) were removed. This resulted in 3,745,051 sites. Samples were then

split into three separate VCF files: all wolves (n=12), all village dogs (n=9), and all

dingoes plus NGSD (n=20). Each group-level VCF was filtered separately, in the

following way. As a result of small sample sizes, formal statistical tests for excess

heterozygosity have no power. Therefore, to filter for excess heterozygosity, only sites

where heterozygote calls made up < 75% within each species-level VCF were retained.

This resulted in 3,729,028 sites for wolves, 3,649,859 sites for village dogs, and

3,733,743 sites for dingoes plus NGSD. Finally, all samples were merged back together

by intersecting all sites among the three data sets, resulting in 3,632,373 sites.

SHAPEIT2.v904 (Delaneau et al. 2014) was used with default parameters to phase the

genotypes of all 41 canids, using the pedigree-based recombination map from

Campbell, et al. (Campbell et al. 2016). The ancestral and derived states of each variant

for the 18 dingo samples was taken from Kumar et al. (Kumar et al. 2023).

Relatedness and Population Structure

We used BCFtools v1.18 (Danecek et al. 2021) to remove sites with 2 or fewer minor

allele counts in the 18 dingo samples, which resulted in 887,759 biallelic loci. We then

used ASD v1.1.0a (Szpiech 2014) with the flag --ibs to compute the proportion of sites

at which a pair of individuals shared 0, 1, and 2 alleles Identical-By-State (IBS) for each

pair of dingoes.

To assess relatedness among K'gari dingoes we plotted IBS0 (the proportion of 0

identical alleles at a given position being shared by two individuals) vs IBS2 (the

proportion of 2 identical alleles being shared by two individuals). Based on records from

Queensland Parks and Wildlife Service (QPWS), there are three samples in the K'gari

dingo population that look to be related to one another by first-degree relationship (i.e.,

samples Y2, B44, and Y4). To verify these relationships, we used KING v2.3.1

(Manichaikul et al. 2010) with the --kinship flag, which incorporates an algorithm that

estimates a kinship coefficient between pairs of individuals independent of the

population's allele frequencies.

To assess population structure among canids we computed pairwise allele sharing

dissimilarities (Szpiech 2014) for all 41 canids included in this study and decomposed

the dissimilarity matrix with multidimensional scaling (MDS). For MDS analysis using

only the 18 dingo samples, we subset the allele sharing dissimilarities from the 41

canids and recomputed the MDS.

Genetic diversity within and between canid populations

We estimated allelic richness, private allelic richness, and the number of alleles private

to pairs of populations (found in each population but no other) using ADZE v1.0

(Szpiech et al. 2008). ADZE uses a rarefaction approach to compare these statistics at

standardized sample sizes. For these analyses we grouped the two mainland dingo

populations (northwest and southeast) and excluded the red wolf and NGSD samples

because of small sample sizes. Parameters were set as follows: COMB 1, K_RANGE 2,

and MAX_G 24. To visualize results, we used R (R Core Team 2022) to plot the mean

of each estimate calculated for each standardized sample size.

Estimating IBD and ROH

To call IBD tracts, we used RefinedIBD v17Jan20.102 (Browning and Browning 2013) to

identify Identical-By-Descent (IBD) segments with default parameters and the

CanFam3.1 genomic map (Campbell et al. 2016). We called runs of homozygosity

(ROH) in each canid population using GARLIC v1.6.0a (Szpiech et al. 2017).

Parameters were set as follows: --winsize 100, --auto-winsize, --auto-overlap-frac, --

size-bounds 1000000 3000000 5000000, --gl-type GQ, --resample 40. This results in 4

length classes of ROH: <1Mbp, ≥1Mbp to <3Mbp, ≥3Mbp to <5Mbp, and ≥5Mbp.

As GARLIC uses allele frequency information to call ROH, we account for the close

relatedness among the K'gari dingo samples, which could result in miscalling ROH in

these individuals, by using the best linear unbiased estimator for allele frequencies

(McPeek et al. 2004) computed from pooling all dingoes together. The allele frequency

at locus i, f_i, is computed as $f_i = \sum_{k=1}^{n} \frac{\sum_{j=1}^{n}(K^{-1})_{jk}}{1^T K^{-1} 1} P_k^{(i)}$, where $P_k^{(i)}$ is the fraction of

derived alleles at a locus in individual k, K is a symmetric matrix of kinship coefficients,

K^{-1} is the inverse matrix of K, 1 is a column vector of n elements with all entries equal

to 1, and 1^T is the transpose of 1. The kinship matrix for dingoes obtained from the --

make-king option in PLINK (Purcell et al. 2007), and any negative entries were set to 0

(https://github.com/szpiech/blue_af). The allele frequencies were passed to GARLIC

with the --freq-file option for use in calling ROH for the 18 dingoes and the 2 New

Guinea singing dog samples.

Results were filtered to keep only the ROH segments longer than 0.5Mbp. We

summarized the number of ROH segments (nROH) and sum total length of ROH

(sROH) within each individual and each length class. The fraction of the genome within

ROH was calculated as the total length of ROH of 0.5 Mbp or longer within an individual

and divided by the total length of the CanFam3.1 autosomal reference genome

(2,203,764,842 nucleotides). Similarly, we estimated inbreeding coefficients as given by

ROH for each population (F_{ROH}). This coefficient represents the proportion of the

autosomal genome covered by ROH 1Mbp or longer.

Assessing burden of deleterious variants

Variants called across all 41 canids were annotated with Ensembl Variant Effect

Predictor (VEP) from release 83 (McLaren et al. 2010). We used SIFT 4G (Vaser et al.

2016) to predict whether an alternate allele at a coding mutation is tolerated or

deleterious for protein function based on the CanFam3.1 genome annotations. We sort

these mutations into three categories: Deleterious, Tolerated, and Loss-of-Function

(LoF). The "Deleterious" group includes only nonsynonymous variants that were

predicted as damaging (1,921 sites). The "Tolerated" group contains all synonymous

and nonsynonymous variants predicted as tolerated (16,380 sites). If a mutation has

more than one prediction we retain the most damaging prediction, i.e., if there is at least

one damaging prediction among multiple predictions, it is classified as "Deleterious".

The LoF category is composed of mutations that are annotated as "Stop-Gain", "Stop-

Loss", or "Start-Loss" (146 sites) and is a separate category from those predicted as

"Deleterious".

We generated a second set of SIFT predictions, this time, requiring that the derived

allele is the one with the prediction. For comparison purposes, we kept both mainland

dingo populations (northwest and southeast) together. We created the same three

categories (1,357 Deleterious, 7,713 Tolerated, and 88 LoF sites) and counted the

number of derived-allele genotypes per individual for each category by summing the

number of heterozygotes plus two times the number of homozygotes.

To assess purging of deleterious variants we compute the Rxy statistic (Xue et al.

2015), which compares the derived allele frequencies of one set of variant sites (e.g.,

putatively deleterious sites) normalized by a set of putatively neutral sites (e.g.,

intergenic sites) between two populations X and Y. We set X = K'gari dingoes and Y =

Mainland dingoes, and considered several sets of sites, including SIFT-classified

Deleterious, SIFT-classified Tolerated, LoF, and sites annotated as synonymous (n =

4708), nonsynonymous (n = 4668), or intergenic sites (n = 100,000). For each of these

sets of sites, we normalize against a second (independent) set of intergenic sites (n=

100,000). To account for the close relatedness among K'gari dingoes we implement a

kinship-weighted Rxy statistic. The modified Rxy statistic

(https://github.com/szpiech/Rxy-kin) incorporates the kinship matrix for K'gari and

mainland dingoes obtained from the --make-king option in PLINK (Purcell et al. 2007),

where any negative entries are set to 0. The statistic is defined (Xue et al. 2015) as

$R_{X,Y}(D) = \frac{L_{X,Y}(D)}{L_{Y,X}(D)}$, where $L_{A,B}(D) = \frac{\sum_{i \in D} f_i^A (1 - f_i^B)}{\sum_{j \in N} f_j^A (1 - f_j^B)}$, D is a set of test sites, N is a set of

putatively neutral sites, f_m^Z is the derived allele frequency at site m in population Z. To

account for relatedness when estimating allele frequencies, f_i are computed as

described above, using weights derived from the kinship matrix. We estimate the

variance in our kinship-weighted Rxy statistic using a jackknife sampling of 10,000

iterations with 20% reduction per iteration.

DATA AVAILABILITY

Raw sequence data of each of the K'gari dingo genomes generated in this study are
available on SRA under project PRJNA1021344.

ACKNOWLEDGEMENTS

The authors would like to acknowledge the Butchulla People as the traditional
custodians of K'gari. Thanks to the Queensland Parks and Wildlife Service for access to
the K'gari wongari (dingo) DNA sample collection, and thanks to Anna María Calderón
for helpful feedback on early versions of this work. This work was supported by the
National Institute of General Medical Sciences of the National Institutes of Health under
Award Number R35GM146926 (AVLA and ZAS), by start-up funds from the
Pennsylvania State University's Department of Biology (AVLA and ZAS), and by a grant
from the Australian Dingo Foundation (KMC). Computations for this research were
performed using the Pennsylvania State University's Institute for Computational Data
Sciences' Roar supercomputer.

SUPPLEMENTARY MATERIAL

Supplementary data are available at XXXX.

CONFLICTS OF INTEREST

KMC is a scientific advisor to the Australian Dingo Foundation, The New Guinea
Singing Dog Conservation Society and The New Guinea Highland Wild Dog
Foundation. KMC and LB are members of the IUCN Canid Specialist Group Dingo
Working Group.

REFERENCES

Allen B. 2023. Genetic health and status of K'gari wongari (Fraser Island dingoes).

Queensland Government - Department of Environment and Science [Internet].

Available from:

https://parks.des.qld.gov.au/__data/assets/pdf_file/0022/319414/genetic-health-

status-kgari-wongari-summary.pdf

Allen BL, Higginbottom K, Bracks JH, Davies N, Baxter GS. 2015. Balancing dingo

conservation with human safety on Fraser Island: the numerical and demographic

effects of humane destruction of dingoes. *Australas. J. Environ. Manag.* 22:197–

215.

Ardalan A, Oskarsson M, Natanaelsson C, Wilton AN, Ahmadian A, Savolainen P.

2012. Narrow genetic basis for the Australian dingo confirmed through analysis of

paternal ancestry. *Genetica* 140:65–73.

Auton A, Rui Li Y, Kidd J, Oliveira K, Nadel J, Holloway JK, Hayward JJ, Cohen

PE, Greally JM, Wang J, et al. 2013. Genetic recombination is targeted towards

gene promoter regions in dogs. *PLoS Genet.* 9:e1003984.

Behrendorff L. 2021. Best-practice dingo management: six lessons from K'gari

(Fraser Island). *Aust. Zool.* 41:521–533.

Benazzo A, Trucchi E, Cahill JA, Maisano Delser P, Mona S, Fumagalli M,

Bunnefeld L, Cornetti L, Ghirotto S, Girardi M, et al. 2017. Survival and divergence

in a small group: The extraordinary genomic history of the endangered Apennine

brown bear stragglers. *Proc. Natl. Acad. Sci. U. S. A.* 114:E9589–E9597.

Bergström A, Frantz L, Schmidt R, Ersmark E, Lebrasseur O, Girdland-Flink L, Lin

AT, Storå J, Sjögren K-G, Anthony D, et al. 2020. Origins and genetic legacy of

prehistoric dogs. *Science* 370:557–564.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* 30:2114–2120.

Brink H, Purcell BV, Letnic M, Webster HS, Appleby RG, Jordan NR. 2019. Pets

and pests: a review of the contrasting economics and fortunes of dingoes and

domestic dogs in Australia, and a proposed new funding scheme for non-lethal

dingo management. *Wildl. Res.* 46:365.

Broad Institute. 2019. Picard Toolkit. *Broad Institute; GitHub repository* [Internet].

Available from: https://github.com/broadinstitute/picard

Browning BL, Browning SR. 2013. Improving the accuracy and efficiency of

identity-by-descent detection in population data. *Genetics* 194:459–471.

Brown SK, Pedersen NC, Jafarishorijeh S, Bannasch DL, Ahrens KD, Wu J-T,

Okon M, Sacks BN. 2011. Phylogenetic distinctiveness of Middle Eastern and

Southeast Asian village dog Y chromosomes illuminates dog origins. *PLoS One*

6:e28496.

Cairns KM. 2021. What is a dingo – origins, hybridisation and identity. *Aust. Zool.*

41:322–337.

Cairns KM, Brown SK, Sacks BN, Ballard JWO. 2017. Conservation implications for

dingoes from the maternal and paternal genome: Multiple populations, dog

introgression, and demography. *Ecol. Evol.* 7:9787–9807.

Cairns KM, Crowther MS, Nesbitt B, Letnic M. 2021. The myth of wild dogs in

Australia: are there any out there? *Aust. Mammal.* 44:67–75.

Cairns KM, Crowther MS, Parker HG, Ostrander EA, Letnic M. 2023. Genome-wide

variant analyses reveal new patterns of admixture and population structure in

Australian dingoes. *Mol. Ecol.* 32:4133–4150.

Cairns KM, Nesbitt BJ, Laffan SW, Letnic M, Crowther MS. 2020. Geographic hot

spots of dingo genetic ancestry in southeastern Australia despite hybridisation with

domestic dogs. *Conserv. Genet.* 21:77–90.

Cairns KM, Shannon LM, Koler-Matznick J, Ballard JWO, Boyko AR. 2018.

Elucidating biogeographical patterns in Australian native canids using genome wide

SNPs. *PLoS One* 13:e0198754.

Cairns KM, Wilton AN. 2016. New insights on the history of canids in Oceania

based on mitochondrial and nuclear data. *Genetica* 144:553–565.

Campbell CL, Bhérer C, Morrow BE, Boyko AR, Auton A. 2016. A Pedigree-Based

Map of Recombination in the Domestic Dog Genome. *G3* 6:3517–3524.

Colpitts J, McLoughlin PD, Poissant J. 2022. Runs of homozygosity in Sable Island

feral horses reveal the genomic consequences of inbreeding and divergence from

domestic breeds. *BMC Genomics* 23:501.

Conroy GC, Lamont RW, Bridges L, Stephens D, Wardell-Johnson A, Ogbourne

SM. 2021. Conservation concerns associated with low genetic diversity for K'gari-

Fraser Island dingoes. *Sci. Rep.* 11:9503.

Corbett LK. 2001. The dingo in Australia and Asia. Sydney: University of NSW

Press

Crowther MS, Fillios M, Colman N, Letnic M. 2014. An updated description of the

Australian dingo (Canis dingo Meyer, 1793). *J. Zool.* 293:192–203.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A,

Keane T, McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and

BCFtools. *Gigascience* [Internet] 10. Available from:

https://academic.oup.com/gigascience/article/doi/10.1093/gigascience/giab008/613

7722

Delaneau O, Marchini J, 1000 Genomes Project Consortium, 1000 Genomes

Project Consortium. 2014. Integrating sequence and array data to create an

improved 1000 Genomes Project haplotype reference panel. *Nat. Commun.*

5:3934.

Dussex N, van der Valk T, Morales HE, Wheat CW, Díez-Del-Molino D, von Seth J,

Foster Y, Kutschera VE, Guschanski K, Rhie A, et al. 2021. Population genomics of

the critically endangered kākāpō. *Cell Genom* 1:100002.

Elledge AE, Allen LR, Carlsson B-L, Wilton AN, Leung LK-P. 2008. An evaluation of

genetic analyses, skull morphology and visual appearance for assessing dingo

purity: implications for dingo conservation. *Wildl. Res.* 35:812.

Fan Z, Silva P, Gronau I, Wang S, Armero AS, Schweizer RM, Ramirez O,

Pollinger J, Galaverni M, Ortega Del-Vecchyo D, et al. 2016. Worldwide patterns of

genomic variation and admixture in gray wolves. *Genome Res.* 26:163–173.

Field MA, Yadav S, Dudchenko O, Esvaran M, Rosen BD, Skvortsova K, Edwards

RJ, Keilwagen J, Cochran BJ, Manandhar B, et al. 2022. The Australian dingo is an

early offshoot of modern breed dogs. *Sci Adv* 8:eabm5944.

Foote AD, Alexander A, Ballance LT, Constantine R, Galletti Vernazzani Muñoz B,

Guinet C, Robertson KM, Sinding M-HS, Sironi M, Tixier P, et al. 2023. "Type D"

killer whale genomes reveal long-term small population size and low genetic

diversity. *J. Hered.* 114:94–109.

Frantz LAF, Mullin VE, Pionnier-Capitan M, Lebrasseur O, Ollivier M, Perri A,

Linderholm A, Mattiangeli V, Teasdale MD, Dimopoulos EA, et al. 2016. Genomic

and archaeological evidence suggest a dual origin of domestic dogs. *Science*

352:1228–1231.

Freedman AH, Wayne RK. 2017. Deciphering the Origin of Dogs: From Fossils to

Genomes. *Annu Rev Anim Biosci* 5:281–307.

Germonpré M, Lázničková-Galetová M, Sablin MV. 2012. Palaeolithic dog skulls at

the Gravettian Předmostí site, the Czech Republic. *J. Archaeol. Sci.* 39:184–202.

Germonpré M, Sablin MV, Stevens RE, Hedges REM, Hofreiter M, Stiller M,

Després VR. 2009. Fossil dogs and wolves from Palaeolithic sites in Belgium, the

Ukraine and Russia: osteometry, ancient DNA and stable isotopes. *J. Archaeol.*

Sci. 36:473–490.

Glen AS, Dickman CR, Soulé ME, Mackey BG. 2007. Evaluating the role of the

dingo as a trophic regulator in Australian ecosystems. *Austral Ecol.* 32:492–501.

Greig K, Gosling A, Collins CJ, Boocock J, McDonald K, Addison DJ, Allen MS,

David B, Gibbs M, Higham CFW, et al. 2018. Complex history of dog (Canis

familiaris) origins and translocations in the Pacific revealed by ancient

mitogenomes. *Sci. Rep.* 8:9130.

Grossen C, Guillaume F, Keller LF, Croll D. 2020. Purging of highly deleterious

mutations through severe bottlenecks in Alpine ibex. *Nat. Commun.* 11:1001.

Hedrick PW, Robinson JA, Peterson RO, Vucetich JA. 2019. Genetics and

extinction and the example of Isle Royale wolves. *Anim. Conserv.* 22:302–309.

Jagannathan V, Drögemüller C, Leeb T, Dog Biomedical Variant Database

Consortium (DBVDC). 2019. A comprehensive biomedical variant catalogue based

on whole genome sequences of 582 dogs and eight wolves. *Anim. Genet.* 50:695–

704.

Johnson CN, Isaac JL, Fisher DO. 2007. Rarity of a top predator triggers continent-

wide collapse of mammal prey: dingoes and marsupials in Australia. *Proc. Biol. Sci.*

274:341–346.

Jones E. 1990. Physical characteristics and taxonomic status of wild canids, Canis-

familiaris, from the Eastern highlands of Victoria. *Wildl. Res.* 17:69.

Kardos M, Åkesson M, Fountain T, Flagstad Ø, Liberg O, Olason P, Sand H,

Wabakken P, Wikenros C, Ellegren H. 2018. Genomic consequences of intensive

inbreeding in an isolated wolf population. *Nat Ecol Evol* 2:124–131.

Khan A, Patel K, Shukla H, Viswanathan A, van der Valk T, Borthakur U, Nigam P,

Zachariah A, Jhala YV, Kardos M, et al. 2021. Genomic evidence for inbreeding

depression and purging of deleterious genetic variation in Indian tigers. *Proc. Natl.*

Acad. Sci. U. S. A. [Internet] 118. Available from:

http://dx.doi.org/10.1073/pnas.2023018118

Kirin M, McQuillan R, Franklin CS, Campbell H, McKeigue PM, Wilson JF. 2010.

Genomic runs of homozygosity record population history and consanguinity. *PLoS*

One 5:e13996.

Kumar M, Conroy G, Ogbourne S, Cairns K, Borburgh L, Subramanian S. 2023.

Genomic signatures of bottleneck and founder effects in dingoes. *Ecol. Evol.*

13:e10525.

Kyriazis CC, Wayne RK, Lohmueller KE. 2021. Strongly deleterious mutations are

a primary determinant of extinction risk due to inbreeding depression. *Evol Lett*

5:33–47.

Kyriazis CC, Beichman AC, Brzeski KE, Hoy SR, Peterson RO, Vucetich JA,

Vucetich LM, Lohmueller KE, Wayne RK. 2023. Genomic Underpinnings of

Population Persistence in Isle Royale Moose. *Mol. Biol. Evol.* 40:msad021.

Larson G, Karlsson EK, Perri A, Webster MT, Ho SYW, Peters J, Stahl PW, Piper
PJ, Lingaas F, Fredholm M, et al. 2012. Rethinking dog domestication by
integrating genetics, archeology, and biogeography. *Proc. Natl. Acad. Sci. U. S. A.*
109:8878–8883.

Letnic M, Crowther MS, Dickman CR, Ritchie EG. 2011. Demonising the dingo:
How much wild dogma is enough? *Curr. Zool.* 57:668–670.

Letnic M, Koch F, Gordon C, Crowther MS, Dickman CR. 2009. Keystone effects of
an alien top-predator stem extinctions of native mammals. *Proc. Biol. Sci.*
276:3249–3256.

Letnic M, Ritchie EG, Dickman CR. 2012. Top predators as biodiversity regulators:
the dingo Canis lupus dingo as a case study. *Biol. Rev. Camb. Philos. Soc.*
87:390–413.

Liberg O, Andrén H, Pedersen H-C, Sand H, Sejberg D, Wabakken P, Kesson M,
Bensch S. 2005. Severe inbreeding depression in a wild wolf (Canis lupus)
population. *Biol. Lett.* 1:17–20.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–
Wheeler transform. *Bioinformatics* 25:1754–1760.

Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen W-M. 2010. Robust
relationship inference in genome-wide association studies. *Bioinformatics* 26:2867–

2873.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella

K, Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: a

MapReduce framework for analyzing next-generation DNA sequencing data.

Genome Res. 20:1297–1303.

McLaren W, Pritchard B, Rios D, Chen Y, Flicek P, Cunningham F. 2010. Deriving

the consequences of genomic variants with the Ensembl API and SNP Effect

Predictor. *Bioinformatics* 26:2069–2070.

McPeek MS, Wu X, Ober C. 2004. Best linear unbiased allele-frequency estimation

in complex pedigrees. *Biometrics* 60:359–367.

Newsome AE, Corbett LK. 1985. The identity of the dingo III.* the incidence of

dingoes, dogs and hybrids and their coat colours in remote and settled regions of

Australia. *Aust. J. Zool.* 33:363.

O'Neill AJ, Cairns KM, Kaplan G, Healy E. 2017. Managing dingoes on Fraser

Island: culling, conflict, and an alternative. *Pac. Conserv. Biol.* 23:4.

Oskarsson MCR, Klütsch CFC, Boonyaprakob U, Wilton A, Tanabe Y, Savolainen

P. 2012. Mitochondrial DNA data indicate an introduction through Mainland

Southeast Asia for Australian dingoes and Polynesian domestic dogs. *Proc. Biol.*

Sci. 279:967–974.

Pemberton TJ, Absher D, Feldman MW, Myers RM, Rosenberg NA, Li JZ. 2012.

Genomic patterns of homozygosity in worldwide human populations. *Am. J. Hum. Genet.* 91:275–292.

Petrie R. 1995. Early Days On Fraser Island 1913-1922 - Rollo Petrie: 9780646276809 - AbeBooks. *Go Bush Safaris* [Internet]. Available from: https://www.abebooks.com/9780646276809/Early-Days-Fraser-Island-1913-1922-0646276808/plp

Plassais J, Kim J, Davis BW, Karyadi DM, Hogan AN, Harris AC, Decker B, Parker HG, Ostrander EA. 2019. Whole genome sequencing of canids reveals genomic regions under selection and variants influencing morphology. *Nat. Commun.* 10:1489.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker PIW, Daly MJ, et al. 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* 81:559–575.

Räikkönen J, Bignert A, Mortensen P, Fernholm B. 2006. Congenital defects in a highly inbred wild wolf population (Canis lupus). *Mamm. Biol.* 71:65–73.

R Core Team. 2022. R: A language and environment for statistical computing. Available from: https://www.r-project.org/

Robinson JA, Brown C, Kim BY, Lohmueller KE, Wayne RK. 2018. Purging of Strongly Deleterious Mutations Explains Long-Term Persistence and Absence of Inbreeding Depression in Island Foxes. *Curr. Biol.* 28:3487–3494.e4.

Robinson JA, Ortega-Del Vecchyo D, Fan Z, Kim BY, vonHoldt BM, Marsden CD,

Lohmueller KE, Wayne RK. 2016. Genomic Flatlining in the Endangered Island

Fox. *Curr. Biol.* 26:1183–1189.

Robinson JA, Räikkönen J, Vucetich LM, Vucetich JA, Peterson RO, Lohmueller

KE, Wayne RK. 2019. Genomic signatures of extensive inbreeding in Isle Royale

wolves, a population on the threshold of extinction. *Sci Adv* 5:eaau0757.

Roelke ME, Martenson JS, O'Brien SJ. 1993. The consequences of demographic

reduction and genetic depletion in the endangered Florida panther. *Curr. Biol.*

3:340–350.

Sacks BN, Brown SK, Stephens D, Pedersen NC, Wu J-T, Berry O. 2013. Y

chromosome analysis of dingoes and southeast asian village dogs suggests a

neolithic continental expansion from Southeast Asia followed by multiple

Austronesian dispersals. *Mol. Biol. Evol.* 30:1103–1118.

Savolainen P, Leitner T, Wilton AN, Matisoo-Smith E, Lundeberg J. 2004. A

detailed picture of the origin of the Australian dingo, obtained from the study of

mitochondrial DNA. *Proc. Natl. Acad. Sci. U. S. A.* 101:12387–12390.

Skoglund P, Ersmark E, Palkopoulou E, Dalén L. 2015. Ancient wolf genome

reveals an early divergence of domestic dog ancestors and admixture into high-

latitude breeds. *Curr. Biol.* 25:1515–1519.

Smith BP, Cairns KM, Adams JW, Newsome TM, Fillios M, Déaux EC, Parr WCH,

Letnic M, VAN Eeden LM, Appleby RG, et al. 2019. Taxonomic status of the

Australian dingo: the case for Canis dingo Meyer, 1793. *Zootaxa* 4564:zootaxa.4564.1.6.

Stephens D, Wilton AN, Fleming PJS, Berry O. 2015. Death by sex in an Australian icon: a continent-wide survey reveals extensive hybridization between dingoes and domestic dogs. *Mol. Ecol.* 24:5643–5656.

Steux C, Szpiech ZA. 2023. The maintenance of deleterious variation in wild Chinese rhesus macaques. *bioRxiv* [Internet]. Available from: http://biorxiv.org/lookup/doi/10.1101/2023.10.04.560901

Stoffel MA, Johnston SE, Pilkington JG, Pemberton JM. 2021. Mutation load decreases with haplotype age in wild Soay sheep. *Evol Lett* 5:187–195.

Surbakti S, Parker HG, McIntyre JK, Maury HK, Cairns KM, Selvig M, Pangau-Adam M, Safonpo A, Numberi L, Runtuboi DYP, et al. 2020. New Guinea highland wild dogs are the original New Guinea singing dogs. *Proc. Natl. Acad. Sci. U. S. A.* 117:24369–24376.

Swinford NA, Prall SP, Gopalan S, Williams CM, Sheehama J, Scelza BA, Henn BM. 2023. Increased homozygosity due to endogamy results in fitness consequences in a human population. *Proc. Natl. Acad. Sci. U. S. A.* 120:e2309552120.

Szpiech Z. 2014. asd. *GitHub* [Internet]. Available from: https://github.com/szpiech/asd

Szpiech ZA, Blant A, Pemberton TJ. 2017. GARLIC: Genomic Autozygosity

Regions Likelihood-based Inference and Classification. *Bioinformatics* 33:2059–

2062.

Szpiech ZA, Jakobsson M, Rosenberg NA. 2008. ADZE: a rarefaction approach for

counting alleles private to combinations of populations. *Bioinformatics* 24:2498–

2504.

Szpiech ZA, Mak ACY, White MJ, Hu D, Eng C, Burchard EG, Hernandez RD.

2019. Ancestry-Dependent Enrichment of Deleterious Homozygotes in Runs of

Homozygosity. *Am. J. Hum. Genet.* 105:747–762.

Szpiech ZA, Xu J, Pemberton TJ, Peng W, Zöllner S, Rosenberg NA, Li JZ. 2013.

Long runs of homozygosity are enriched for deleterious variation. *Am. J. Hum.*

Genet. 93:90–102.

Taron UH, Salado I, Escobar-Rodríguez M, Westbury MV, Butschkau S, Paijmans

JLA, vonHoldt BM, Hofreiter M, Leonard JA. 2021. A sliver of the past: The

decimation of the genetic diversity of the Mexican wolf. *Mol. Ecol.* 30:6340–6354.

Vaser R, Adusumalli S, Leng SN, Sikic M, Ng PC. 2016. SIFT missense predictions

for genomes. *Nat. Protoc.* 11:1–9.

Wallach AD, Johnson CN, Ritchie EG, O'Neill AJ. 2010. Predator control promotes

invasive dominated ecological states. *Ecol. Lett.* 13:1008–1018.

Wang G-D, Zhai W, Yang H-C, Fan R-X, Cao X, Zhong L, Wang L, Liu F, Wu H,

Cheng L-G, et al. 2013. The genomics of selection in dogs and the parallel

evolution between dogs and humans. *Nat. Commun.* 4:1–9.

Wardell-Johnson G, Schoeman D, Schlacher T, Wardell-Johnson A, Weston MA,

Shimizu Y, Conroy G. 2015. Re-framing values for a World Heritage future: what

type of icon will K'gari-Fraser Island become? *Australas. J. Environ. Manag.*

22:124–148.

Wilton AN. 2001. DNA methods of assessing dingo purity. In: A Symposium on the

Dingo. P.O. Box 20, Mosman NSW 2088, Australia: Royal Zoological Society of

New South Wales. p. 49–56.

Wootton E, Robert C, Taillon J, Côté SD, Shafer ABA. 2023. Genomic health is

dependent on long-term population demographic history. *Mol. Ecol.* 32:1943–1954.

Xue Y, Prado-Martinez J, Sudmant PH, Narasimhan V, Ayub Q, Szpak M,

Frandsen P, Chen Y, Yngvadottir B, Cooper DN, et al. 2015. Mountain gorilla

genomes reveal the impact of long-term population decline and inbreeding. *Science*

348:242–245.

Zhang S-J, Wang G-D, Ma P, Zhang L-L, Yin T-T, Liu Y-H, Otecko NO, Wang M,

Ma Y-P, Wang L, et al. 2020. Genomic regions under selection in the feralization of

the dingoes. *Nat. Commun.* 11:671.

Zhang W, Fan Z, Han E, Hou R, Zhang L, Galaverni M, Huang J, Liu H, Silva P, Li

P, et al. 2014. Hypoxia adaptations in the grey wolf (Canis lupus chanco) from

Qinghai-Tibet Plateau. *PLoS Genet.* 10:e1004466.

 FIGURES

Fig. 1. Reduced dimensional representation of sample clustering based on
multidimensional scaling of the allele sharing dissimilarity matrix. First two major axes of
variation plotted for (A) all 41 canids included in this study and (B) 18 dingo individuals.

Fig. 2. Genetic diversity estimates within and between canid populations. Error bars
representing 95% confidence intervals for each line are too narrow to be visible. (A)
Mean number of alleles per locus (allelic richness). (B) Mean number of private alleles

per locus (private allelic richness). (C) Mean number of shared alleles per locus

between pairs of populations.

Fig. 3. ROH coverage across individual genomes. The percentage of the genome

covered by class A ROH (0.5-1 Mbp), class B ROH (1-3 Mbp), class C ROH (3-5 Mbp)

and class D ROH (>5 Mbp) is shown on the Y-axis for each of the 41 canids included in

this study.

Fig. 4. Total number of predicted damaging genotypes in each canid population. (A)

Alternate-allele deleterious homozygotes. (B) Alternate-allele deleterious heterozygotes.

(C) Alternate-allele LoF homozygotes. (D) Alternate-allele LoF heterozygotes. P-value

significance by Wilcoxon test with K'gari dingoes as reference population. Significance

levels of p-values: (*) p-value < 0.05, (**) p-value < 0.005, (ns) p-value > 0.05.

Fig. 5. Kinship-weighted Rxy estimates for K'gari and mainland dingoes. Estimates

represent the relative excess number of derived alleles for each category in K'gari

dingoes (population X) relative to mainland dingoes (population Y). Error bars represent

the 95% confidence interval from jackknife resampling.