Genomic consequences of isolation and inbreeding in an island dingo population

Ana V. Leon-Apodaca[1], Manoharan Kumar[2], Andres del Castillo[1], Gabriel C. Conroy[3,4], Robert W. Lamont[3,4], Steven Ogbourne[4], Kylie M. Cairns[5,6], Liz Borburgh[3], Linda Behrendorff[7], Sankar Subramanian[2,4,*], Zachary A. Szpiech[1,8,*]

[1] Department of Biology, Pennsylvania State University, PA, USA

[2] School of Science, Technology & Engineering, University of the Sunshine Coast, 1 Moreton Parade, Petrie, Queensland, Australia

[3] School of Science, Technology & Engineering, University of the Sunshine Coast, 90 Sippy Downs Drive, Sippy Downs, Queensland, Australia

[4] Centre for Bioinnovation, University of the Sunshine Coast, 90 Sippy Downs Drive, Sippy Downs, Queensland, Australia

[5] Evolution & Ecology Research Centre, School of Biological, Earth and Environmental Sciences, UNSW Australia, Sydney NSW 2052, Australia

[6] Centre for Ecosystem Science, School of Biological, Earth and Environmental Sciences, UNSW Australia, Sydney NSW 2052, Australia

[7] The University of Queensland, School of Agriculture and Food Sciences, Gatton, Queensland 4343, Australia

[8] Institute for Computational and Data Sciences, Pennsylvania State University, PA, USA

*Corresponding authors

ABSTRACT

Dingoes come from an ancient canid lineage that originated in East Asia around 8000-11,000 years BP. As Australia's largest terrestrial predator, dingoes play an important ecological role. A small population exists on a world heritage listed offshore island, K'gari (formerly Fraser Island). While dingoes on K'gari are protected, dingoes on the mainland are only protected in certain regions, and, because of its controversial status in Australia, lethal control is commonly used for management. Concern regarding the persistence of dingoes on K'gari has risen due to their low genetic diversity and elevated inbreeding levels. However, whole-genome sequencing data is lacking from this population. Here, we include five new whole-genome sequences of K'gari dingoes. We analyze a total of

18 whole genome sequences of dingoes sampled from mainland Australia and K'gari to

assess the genomic consequences of their demographic histories. Long (>1 Mb) runs of

homozygosity (ROH) — indicators of inbreeding — are elevated in all sampled dingoes.

However, K'gari dingoes showed significantly higher levels of very long ROH (>5 Mb),

providing genomic evidence for small population size, isolation, inbreeding, and a strong

founder effect. Our results suggest that, despite current levels of inbreeding, the K'gari

population is purging strongly deleterious mutations, which, in the absence of further

reductions in population size, may facilitate the persistence of small populations despite

low genetic diversity and isolation. However, there is little to no purging of mildly

deleterious alleles, which may have important long-term consequences, and should be

considered by conservation and management programs.

INTRODUCTION

The taxonomic classification of the dingo within the *Canis* genus, as well as its identity

are surrounded by controversy. Debate on the most appropriate nomenclature and

taxonomic classification centers on whether the dingo should be classified as *Canis*

familiaris (Jackson et al. 2017; Jackson et al. 2019) or *Canis dingo* (Crowther et al. 2014;

Smith et al. 2019), depending on how the species concept is defined and employed. The

different lines of thought and opinions on the dingo's evolutionary history, its relationship

with wolves and domestic dogs, its domestication status, and its arrival to Australia has

made it difficult to agree on how to manage and conserve them (Cairns 2021).

While archeological evidence and genomic data estimate the divergence time between

wolves and dogs to be ~15,000-36,000 years BP (Germonpré et al. 2009; Germonpré et

al. 2012; Larson et al. 2012; Skoglund et al. 2015; Frantz et al. 2016; Freedman and

Wayne 2017), it has been estimated dingoes diverged from early dogs ~5,000-11,000

years BP (Savolainen et al. 2004; Oskarsson et al. 2012; Sacks et al. 2013; Cairns and

Wilton 2016; Bergström et al. 2020; Surbakti et al. 2020; Cairns 2021; Field et al. 2022),

forming a distinct and separate lineage from domestic dogs, wolves and other canids

(Crowther et al. 2014; Smith et al. 2019). Research using mtDNA and Y-chromosome

data has identified a close relationship between dingoes, New Guinea singing dogs

(NGSD) and Asian village dogs, suggesting that the ancestors of dingoes originated in

East Asia (Savolainen et al. 2004; Ardalan et al. 2012; Oskarsson et al. 2012; Sacks et

al. 2013; Cairns and Wilton 2016; Cairns et al. 2017). Archeological evidence suggests

dingoes arrived in Australia more than ~3,500 years BP (Milham and Thompson 1976;

Savolainen et al. 2004), remaining isolated from other canids until 1788.

Although dingoes are present across different bioregions in Australia, at least two distinct

mainland dingo lineages have been identified, one in the northwest and one in the

southeast, alongside one offshore lineage on K'gari (formerly known as Fraser Island)

(Cairns and Wilton 2016; Cairns et al. 2017; Cairns et al. 2018; Greig et al. 2018; Zhang

et al. 2020). Genome-wide SNP data suggest there may be additional biogeographic

structure in the mainland dingo population (Cairns et al. 2023). As the apex terrestrial

predator since the extinction of the thylacine, dingoes play an important role in

maintaining the ecosystems of Australia (Glen et al. 2007; Johnson et al. 2007; Letnic et

al. 2009; Wallach et al. 2010; Letnic et al. 2011; Letnic et al. 2012). But they also have a

complex history in Australia, as they are simultaneously declared native species and an

agricultural pest dependent on jurisdiction and context. The dingo is protected in parts of

the Australian Capital Territory, Victoria, Queensland and the Northern Territory as a

native species, yet at the same time, wild dogs, including dingoes, have been declared a

pest or restricted matter in all Australian states and territories for posing a risk to livestock

(Brink et al. 2019).

Hybridization with domestic dogs was previously highlighted as a main concern for dingo

conservation, since high levels of hybridization were observed in dingoes from

southeastern Australia (Newsome and Corbett 1985; Jones 1990; Corbett 2001; Wilton

2001; Elledge et al. 2008; Stephens et al. 2015; Cairns et al. 2020; Cairns 2021; Cairns

et al. 2021). However, a recently published study using genome-wide SNP data has

shown these occurrences are less common than previously thought (Cairns et al. 2023).

Policy and legislation in Australia currently employs the term "wild dog" to refer to dingoes,

feral domestic dogs and dingo-dog hybrids, assuming that pure dingoes are virtually

extinct in the wild and that dingo-dog hybrids are common (Fleming 2001; DEPI 2013;

Bird and Bowman 2016; NWDAP 2020). Disagreement on the dingo's genetic identity,

along with the contradictory legal status of this canid throughout Australia, have allowed

for lethal control to be used on dingoes (without distinction) during pest management

programmes on public and private lands across mainland Australia (van Eeden et al.

2019; Conroy et al. 2021; van Eeden et al. 2021).

K'gari is located off the coast of Queensland, Australia, and has been listed as a UNESCO

World Heritage site since 1992. In addition to being the world's largest sand island, it is

also home to a small protected (*Nature Conservation Act* 1992) population of dingoes of

~70-170 individuals that are genetically distinct from mainland dingoes (Brown et al. 2011;

Conroy et al. 2021). The island's Traditional owners (The Butchulla) have a deep and

long standing connection with dingoes (wongari) and in contemporary times are an

infamous tourist attraction for the over 400,000 tourists visit K'gari each year (Wardell-

Johnson et al. 2015), which has led to negative human-dingo interactions arising from

human associated activities (O'Neill et al. 2017; Behrendorff 2021). In response to these

incidents, management strategies have been implemented in K'gari, including non-lethal

options as well as the removal of dingoes via lethal control (Behrendorff 2021). K'gari

dingoes are of particular interest as there are concerns regarding their persistence due

to low genetic diversity and inbreeding given it is a small, isolated population (Cairns et

al. 2018; Conroy et al. 2021).

Isolated populations often experience reduced fitness resulting from increased levels of

homozygosity and a high load of deleterious mutations across their genomes. There have

been several instances in which these factors have led to inbreeding depression and a

high risk of extinction. Such is the case of the iconic Isle Royale wolves, which are highly

inbred gray wolves located on an island in Lake Superior, in northern United States

(Hedrick et al. 2019; Robinson et al. 2019). Other examples of isolated populations in the

wild that have experienced inbreeding depression are Florida Panthers (Roelke et al.

1993) and Scandinavian gray wolves (Liberg et al. 2005; Räikkönen et al. 2006; Kardos

et al. 2018). In other instances, small isolated populations have managed to persist with

low genetic diversity for generations, without experiencing inbreeding depression. Such

is the case of island foxes (Robinson et al. 2016; Robinson et al. 2018), Appenine brown

bears (Benazzo et al. 2017), African mountain gorillas (Xue et al. 2015), and Alpine ibex

(Grossen et al. 2020). It has been suggested that the purging of strongly deleterious

recessive mutations may facilitate the persistence of long-term small populations (Xue et

al. 2015; Robinson et al. 2018; Grossen et al. 2020; Wootton et al. 2023).

Only a handful of studies have looked at the genetics of K'gari dingoes. While these

studies have found low genetic diversity and high levels of inbreeding on K'gari dingoes

(Cairns and Wilton 2016; Cairns et al. 2018; Conroy et al. 2021), more research is needed

to understand the consequences of inbreeding and isolation on this population. Moreover,

there is a need for new and well-informed conservation strategies for dingoes in Australia,

given that current strategies do not consider these genetic factors. To our knowledge,

there is only one whole-genome sequence from a K'gari dingo individual that is publicly

available (Zhang et al. 2020), although the provenance of the animal is uncertain. In this

work we include five newly sequenced samples, and we look further into the genetics of

K'gari dingoes to understand the consequences of their demographic history on the

distribution of deleterious variations, and the implications of small population sizes and

isolation, to help inform conservation efforts for their long-term persistence.

METHODS

Sampling and sequencing

In this study we include whole-genome sequencing data from a total of 41 canids,

including 18 dingo samples from both K'gari (6) and mainland Australia (4 from the

Southeast, and 8 from the Northeast). We also include 2 New Guinea singing dog

samples, 11 gray wolf samples, 1 red wolf sample, and 9 village dog samples.

Sequencing data from 3 dingo samples from mainland Australia (2 from the Northwest,

and 1 from the Southeast) were obtained from a recently published study (Kumar et al.

2023). Ten dingo samples, comprising 1 from K'gari and 9 from mainland Australia (6

from the Northwest, and 3 from the Southeast), were obtained from a previous study

(Zhang et al. 2020). Eleven gray wolf samples, 1 red wolf sample, 2 New Guinea singing

dog samples, and 9 village dog samples were obtained from a different study (Plassais

et al. 2019). Reads from all of these canids were aligned to the CanFam3 reference

genome using BWA-MEM (Li and Durbin 2009). Five of the K'gari dingo samples in this

study are from new sequencing data. Biological samples from these individuals were

collected between 2011 and 2013 from regions along the east coast of K'gari, including

Cathedral Beach, Eurong, and Dundubara. Two of the three newly sequenced dingo

samples from the mainland were collected from the Northwest, and one from the

Southeast. These samples were sequenced using Illumina Hiseq 2000, and aligned to

the CanFam3 reference genome using BWA-MEM (Li and Durbin 2009). We converted

the mapped reads in the sequencing alignment mapped (SAM) format to binary

alignment/mapped (BAM) format using samtools v1.3.1 (Danecek et al. 2021). Later, we

sorted the BAM file based on chromosomes and alignment position. These sorted reads

were used for marking PCR duplicates using the Picard tool (Broad Institute 2019).

Genotype calling, filtering, and phasing

All samples were jointly genotyped with GATK v4.2 modules HaplotypeCaller,

CombineGVCFs and GenotypeGVCFs. The VCF file containing raw genotype calls for all

41 canids were then filtered in the following way. Only sites with biallelic snps and only

sites with a site-level quality score QUAL > 50 were retained. Any individual genotype

with GQ <= 19 was set to missing. All sites with >= 10% missing data were removed. All

sites with no variation among the 41 samples (i.e., monomorphic sites) were removed.

This resulted in 3,745,051 sites.

Samples were then split into three separate VCF files: all wolves (n=12), all village dogs

(n=9), and all dingoes plus NGSD (n=20). Each group-level VCF was filtered separately,

in the following way. As a result of small sample sizes, formal statistical tests for excess

heterozygosity have no power. Therefore, to filter for excess heterozygosity, only sites

where heterozygote calls made up < 75% within each species-level VCF were retained.

This resulted in 3,729,028 sites for wolves, 3,649,859 sites for village dogs, and

3,733,743 sites for dingoes plus NGSD. Finally, all samples were merged back together

by intersecting all sites among the three data sets, resulting in 3,632,373 sites.

SHAPEIT2 (Delaneau et al. 2014) was used to phase the genotypes of all 41 canids,

using the pedigree-based recombination map from Campbell, et al. (Campbell et al.

2016). The ancestral and derived states of each variant for the 18 dingo samples was

taken from Kumar et al. (Kumar et al. 2023).

Relatedness and Population Structure

We used BCFtools (Danecek et al. 2021) to remove sites with 2 or fewer minor allele

counts in the 18 dingo samples, which resulted in 887,759 biallelic loci. We then used

ASD (Szpiech 2014) to compute the proportion of sites at which a pair of individuals

shared 0, 1, and 2 alleles Identical-By-State (IBS) for each pair of dingoes.

To assess relatedness among K'gari dingoes we plotted IBS0 (the proportion of 0

identical alleles at a given position being shared by two individuals) vs IBS2 (the

proportion of 2 identical alleles being shared by two individuals). Based on records from

Queensland Parks and Wildlife Service (QPWS), there are three samples in the K'gari

dingo population that look to be related to one another by first-degree relationship (i.e.,

samples Y2, B44, and Y4). To verify these relationships, we used the KING method

(Manichaikul et al. 2010), which incorporates an algorithm that estimates a kinship

coefficient between pairs of individuals independent of the population's allele frequencies.

To assess population structure among canids we computed pairwise allele sharing

dissimilarities (Szpiech 2014) for all 41 canids included in this study and decomposed the

dissimilarity matrix with multidimensional scaling (MDS). For MDS analysis using only the

18 dingo samples, we subsetted the allele sharing dissimilarities from the 41 canids and

recomputed the MDS.

Genetic diversity within and between canid populations

We estimated allelic richness, private allelic richness, and the number of alleles private

to pairs of populations (found in each population but no other) using ADZE (Szpiech et

al. 2008). ADZE uses a rarefaction approach to compare these statistics at standardized

sample sizes. For these analyses we grouped the two mainland dingo populations

(northwest and southeast) and excluded the red wolf and NGSD samples because of

small sample sizes. Parameters were set as follows: COMB 1, K_RANGE 2, and MAX_G

24. To visualize results, we used R (R Core Team 2022) to plot the mean of each estimate

calculated for each standardized sample size.

Estimating IBD and ROH

To call IBD tracts, we used RefinedIBD (Browning and Browning 2013) to identify

Identical-By-Descent (IBD) segments with default parameters and the CanFam3.1

genomic map (Campbell et al. 2016). We called runs of homozygosity (ROH) in each

canid population using GARLIC (Szpiech et al. 2017). Parameters were set as follows: --

winsize 100, --auto-winsize, --auto-overlap-frac, --size-bounds 1000000 2000000

3000000 4000000 5000000, --gl-type GQ, --resample 40. This results in 6 length classes

of ROH: <1Mbp, ≥1Mbp to <2Mbp, ≥2Mbp to <3Mbp, ≥3Mbp to <4Mbp, ≥4Mbp to <5Mbp,

and ≥5Mbp.

As GARLIC uses allele frequency information to call ROH, we account for the close

relatedness among the K'gari dingo samples, which could result in miscalling ROH in

these individuals, by using the best linear unbiased estimator for allele frequencies

(McPeek et al. 2004) computed from pooling all dingoes together. The allele frequency at

locus i, f_i, is computed as $f_i = \sum_{k=1}^{n} \frac{\sum_{j=1}^{n}(K^{-1})_{jk}}{1^T K^{-1} 1} P_k^{(i)}$, where $P_k^{(i)}$ is the fraction of derived

alleles at a locus in individual k, K is a symmetric matrix of kinship coefficients, K^{-1} is the

inverse matrix of K, 1 is a column vector of n elements with all entries equal to 1, and 1^T

is the transpose of 1. The kinship matrix for dingoes obtained from the --make-king option

in PLINK (Purcell et al. 2007), and any negative entries were set to 0

(https://github.com/szpiech/blue_af). The allele frequencies were passed to GARLIC with

the --freq-file option for use in calling ROH for the 18 dingoes and the 2 New Guinea

singing dog samples.

Results were filtered to keep only the ROH segments longer than 0.5Mbp. We

summarized the number of ROH segments (nROH) and sum total length of ROH (sROH)

within each individual and each length class. The fraction of the genome within ROH was

calculated as the total length of ROH of 0.5 Mbp or longer within an individual and divided

by the total length of the CanFam3.1 autosomal reference genome (2,203,764,842

nucleotides). Similarly, we estimated inbreeding coefficients as given by ROH for each

population (F_{ROH}). This coefficient represents the proportion of the autosomal genome

covered by ROH 1Mbp or longer.

Assessing burden of deleterious variants

Variants called across all 41 canids were annotated with Ensembl Variant Effect Predictor

(VEP) (McLaren et al. 2010). We used SIFT (Vaser et al. 2016) to predict whether an

alternate allele at a coding mutation is tolerated or deleterious for protein function based

on the CanFam3.1 genome annotations. We sort these mutations into three categories:

Deleterious, Tolerated, and Loss-of-Function (LoF). The "Deleterious" group includes

only nonsynonymous variants that were predicted as damaging (1,921 sites). The

"Tolerated" group contains all synonymous and nonsynonymous variants predicted as

tolerated (16,380 sites). If a mutation has more than one prediction we retain the most

damaging prediction, i.e., if there is at least one damaging prediction among multiple

predictions, it is classified as "Deleterious". LoF mutations are classified as any mutation

with a "Stop-Gain", "Stop-Loss", or "Start-Loss" classification (146 sites).

We generated a second set of SIFT predictions, this time, requiring that the derived allele

is the one with the prediction. For comparison purposes, we kept both mainland dingo

populations (northwest and southeast) together. We created the same three categories

(1,357 Deleterious, 7,713 Tolerated, and 88 LoF sites) and calculated the proportion of

derived allele counts for each category on each dingo population.

To assess purging of deleterious variants we compute the Rxy statistic (Xue et al. 2015),

which compares the derived allele frequencies of one set of variant sites (e.g., putatively

deleterious sites) normalized by a set of putatively neutral sites (e.g., intergenic sites)

between two populations X and Y. We set X = K'gari dingoes and Y = Mainland dingoes,

and considered several sets of sites, including SIFT-classified Deleterious, SIFT-

classified Tolerated, LoF, and sites annotated as synonymous (n = 4708),

nonsynonymous (n = 4668), or intergenic sites (n = 10,000). For each of these sets of

sites, we normalize against a second (independent) set of intergenic sites (n= 10,000).

To account for the close relatedness among K'gari dingoes we implement a kinship-

weighted Rxy statistic. The modified Rxy statistic (https://github.com/szpiech/Rxy-kin)

incorporates the kinship matrix for K'gari and mainland dingoes obtained from the --make-

king option in PLINK (Purcell et al. 2007), where any negative entries are set to 0. The

statistic is defined (Xue et al. 2015) as

$R_{X,Y}(D) = \frac{L_{X,Y}(D)}{L_{Y,X}(D)}$, where $L_{A,B}(D) = \frac{\sum_{i \in D} f_i^A (1 - f_i^B)}{\sum_{j \in N} f_j^A (1 - f_j^B)}$, D is a set of test sites, N is a set of

putatively neutral sites, f_m^Z is the derived allele frequency at site m in population Z. To

account for relatedness when estimating allele frequencies, f_i are computed as described

above, using weights derived from the kinship matrix. We estimate the variance in our

kinship-weighted Rxy statistic using a jackknife sampling of 2,000 iterations with 20%

reduction per iteration.

RESULTS

Canid Sampling, Relatedness and Population Structure

Previous analysis on dingoes suggest there are at least three distinct lineages of dingoes

in Australia, with one being located on the northwest of the mainland, another in the

southeast of the mainland, and the third one offshore on K'gari (Fraser Island) (Cairns

and Wilton 2016; Cairns et al. 2017; Cairns et al. 2018). To expand our understanding of

genetic diversity patterns in the K'gari population, we generated high quality whole-

genome sequences of five dingoes with average read depth between 12.01x and 12.56x

sampled from the island between 2011 and 2013. Although there is a dingo reference

genome (ASM325472v1), the newly sequenced samples in this study were aligned to the

dog reference genome (CanFam3) to take advantage of its fully assembled chromosomes

and better gene annotations. We include whole genome sequences from dingoes from

northwest and southeast mainland Australia, and we also include samples from other

canids, such as wolves and village dogs, and New Guinea singing dogs (NGSD), which

are a close relative of dingoes. All canid reads in this study were aligned to the dog

reference genome (CanFam3).

We identified close relatives in K'gari dingoes in two ways. First, we looked at the

proportion of sites at which pairs of individuals share 0, 1 or 2 alleles that are considered

identical-by-state (IBS). We expect Parent/Offspring to share at least one allele at a locus

identical-by-descent, thus having low values of IBS0. We expect full-siblings to share 2

alleles identical-by-descent for approximately 25% of loci, and we expect second-degree

relatives and unrelated individuals to have higher estimates of IBS0. We also used the

KING algorithm (Manichaikul et al. 2010) to infer kinship coefficients for all dingo

individuals. We identified three pairs of Parent/Offspring relatives, four pairs of full-siblings

relatives, a pair of avuncular relatives, and seven pairs that include third-degree relatives

and unrelated individuals. Furthermore, we found that none of the K'gari dingo individuals

are closely related to dingo individuals from the mainland, and that the mainland dingoes

in this study are not closely related to one another (Suppl. Fig. 1 and Suppl. Table 1).

We computed pairwise allele sharing dissimilarities to assess population structure and

generated multidimensional scaling (MDS) plots from the dissimilarity matrix. In MDS,

individuals with small allele sharing dissimilarities tend to group together. In fig. 1A we

can observe distinct taxonomic groupings, with the close relationship between the dingo

and NGSD lineages clearly captured in this analysis. Individuals within the wolf and village

dog populations were sampled from different regions around the world (see Suppl. Table

2), hence the relatively loose clustering among these individuals. Interestingly, we find

that the village dog from Vietnam seems to be genetically more similar to mainland

dingoes than the village dog from Papua New Guinea (fig. 1A). While Papua New Guinea

is geographically closer to Australia than Vietnam, similar observations regarding shared

ancestry between Vietnamese village dogs and dingoes have been reported by Cairns et

al. (Cairns et al. 2023). A possible explanation regarding the observed genotype

differences among these canids may be explained by Vietnamese village dogs having

retained the ancestral alleles that also exist in the dingo lineage while village dogs from

Papua New Guinea have not, possibly because of their different demographic histories

or genetic swamping from contemporary domestic dogs.

Although dingoes from the southeast and northwest overlap with each other (i.e., they are

genetically more similar), we observe K'gari dingoes form their own separate subgroup

(fig 1A). To look more into detail at the allele sharing differences between the three dingo

populations, we recomputed the MDS on the matrix of allele sharing dissimilarities

containing only these samples. In fig. 1B we can observe geographic based clustering,

with K'gari dingoes maintaining a tighter and more distinct group compared to dingoes in

the southeast and northwest.

Genetic diversity between and within canid populations

To characterize genetic diversity patterns within and between canid populations we

analyzed the distribution of alleles across populations. We used ADZE (Szpiech et al.

2008) to compute estimates of allelic richness, private allelic richness, and the number of

alleles shared by pairs of populations (i.e., alleles that are found only in those populations

and nowhere else). We observed reduced levels of genetic diversity in mainland and

K'gari dingoes, compared to large and outbred populations such as wolves and village

dogs, with K'gari dingoes having the lowest genetic diversity of all the canid groups

analyzed in this study (fig. 2A and 2B). We found mainland dingoes share more alleles

with village dogs than K'gari dingoes and village dogs (fig. 2C), which suggests that the

alleles shared between dingoes and their common ancestors from Southeast Asia were

likely lost in the bottleneck that founded the K'gari dingo population.

Inbreeding and homozygosity

To characterize the levels of inbreeding we looked at long contiguous stretches of

homozygous regions in individual genomes that are shared identical-by-descent (IBD),

also known as runs of homozygosity (ROH). We used GARLIC (Szpiech et al. 2017) to

call ROH, which implements a likelihood-based approach to infer ROH. The length of

ROH can inform on how recent the parental relatedness among individuals is, with long

ROH more commonly arising in offspring descended from parents with a recent common

ancestor. On the other hand, short ROH suggests a more distant common ancestor, as

recombination has had more time to break apart the underlying IBD haplotypes (Kirin et

al. 2010; Pemberton et al. 2012). The wolf and village dog individuals included in this

study were sampled from large and outbred populations from different regions around the

world (see suppl. Table 2 for details), resulting in variation in the levels of homozygosity

found in their individual genomes. Overall, we observed relatively low levels of

homozygosity in village dogs and wolves, with ~7% and ~16% of their genomes being

covered by ROH >0.5 Mbp, respectively (fig. 3). In contrast, we observed high levels of

homozygosity in Northwest and Southeast mainland dingo populations, with ~30% of their

genomes being covered by ROH >0.5 Mbp. In K'gari dingoes we observed even higher

levels of homozygosity, with ~65% of their genomes being covered by ROH >0.5 Mbp,

and ~28% by ROH >5 Mbp (fig. 3), with the longest individual homozygous segment being

64.2 Mbp long (suppl. Fig 3).

We estimated inbreeding coefficients as given by ROH for each population (F_{ROH}), which

represents the proportion of the autosomal genome covered by ROH >1Mbp. F_{ROH} of

0.59, 0.23, 0.23, were estimated for K'gari, Northwest, and Southeast dingoes,

respectively, and F_{ROH} of 0.35 for NGSD. For village dogs and wolves we estimated

inbreeding coefficients of 0.04 and 0.13, respectively. These results provide clear

evidence for inbreeding, isolation, small population size, and a strong founder effect in

K'gari dingoes.

Patterns of deleterious variation and genetic load

To understand the effect of inbreeding and low genetic diversity on the distribution of

deleterious variation, we looked at the burden of deleterious alleles in K'gari dingoes

compared to mainland dingoes and other canids. Alternate alleles were predicted to be

deleterious or neutral by taking into account sequence conservation and the effect of an

amino acid substitution on protein function using SIFT (Vaser et al. 2016) (see Methods).

We found dingoes and NGSD carry a higher burden of deleterious alternate-allele

homozygotes compared to village dogs and wolves, although K'gari dingoes carry the

highest proportion of deleterious alternate-allele homozygotes among all the canids

included in this study (fig. 4A). On the other hand, both mainland and K'gari dingoes seem

to carry similar proportions of alternate-allele homozygous LoF variants (fig. 4B).

To refine our analysis of the differences in the burden of deleterious alleles between K'gari

and mainland dingoes, we subset our deleterious alternate allele calls to include only

those alternate alleles which were also inferred to be the derived allele in dingoes (see

Methods). We computed proportions for deleterious and LoF derived variants in each

dingo sample. Patterns of deleterious/LoF derived homozygotes and heterozygotes

stayed qualitatively similar to the non-polarized patterns. We computed the proportion of

total deleterious/LoF derived alleles per individual by counting the number of

deleterious/LoF heterozygotes and two times the number of deleterious/LoF

homozygotes (divided by two times the total number of genotypes). Both K'gari and

mainland dingoes showed a similar proportion of predicted deleterious alleles (sample

means 290 and 278, respectively, p-value > 0.05 by Mann-Whitney U test; suppl. Fig 4A),

which suggests these alleles may not be strongly deleterious as they have not been

removed from the dingo population in K'gari through negative selection. In contrast, we

observed a significant difference in the proportion of derived allele counts for LoF variants

(sample means 14 and 19 for K'gari and mainland dingoes, respectively, p-value =

0.001101 by Mann-Whitney U test), with a decrease in K'gari dingoes compared to

mainland dingoes (suppl. Fig 4B). Moreover, the mean number of homozygous LoF

variants per individual in K'gari dingoes is significantly lower than in mainland dingoes

(6.17 and 5.25, respectively, p-value = 0.017 by Mann-Whitney U test). Similar patterns

have been observed in other small populations with substantial inbreeding, suggesting

purging of LoF alleles, which are more likely to be strongly recessive variants than small-

effect deleterious variants, may facilitate the persistence of small and isolated populations

(Robinson et al. 2018; Grossen et al. 2020).

To test whether K'gari dingoes showed evidence of purging of deleterious variants we

used the Rxy statistic, as implemented by Xue et al. (Xue et al. 2015), which compares

the frequency of derived alleles of a particular category (e.g., deleterious) normalized by

the frequencies of a set of putatively neutral variants, which is then compared across two

populations, X and Y. When Rxy < 1 for a given site category, we consider this evidence

of purging of those alleles in population X relative to Y, when Rxy > 1 we consider this

evidence of enrichment of those alleles in population X relative to Y. When Rxy = 1, we

consider this evidence of no difference of those alleles between X and Y. As our K'gari

samples are closely related, we implemented a modified version of the Rxy statistic, which

accounts for relatedness among K'gari dingoes. For this analysis we looked at several

allele categories: sites that were predicted as deleterious, tolerated, LoF, sites that were

annotated as synonymous, non-synonymous, and a random set of intergenic sites as a

control. All Rxy calculations used an additional, independent, set of intergenic sites for

normalization. We estimated the excess number of deleterious variants in K'gari dingoes

(population X) with respect to mainland dingoes (population Y) and found evidence of

purging of LoF alleles (i.e., putative strongly deleterious alleles), but little to no evidence

of purging of mild/slightly deleterious alleles in K'gari dingoes (fig. 5). Although we

observe a small enrichment in the rest of the categories, we hypothesize these results

may be due to the small sample size and low genetic diversity in K'gari dingoes compared

to mainland dingoes. To assess the effect of sample sizes on our results, we conducted

this analysis using the same number of samples for each dingo population and found that

results stayed very similar (Suppl. Fig. 5).

DISCUSSION

In this work, we have generated five whole-genome sequences of K'gari dingoes to

investigate the consequences of isolation and inbreeding on this island population. We

find K'gari dingoes have lower genetic diversity, a higher level of homozygosity, and a

higher accumulation of mildly deleterious alleles compared to dingoes in the mainland

and canids from large populations. However, contrary to the idea that inbred populations

are bound to experience inbreeding depression and subsequently face a high risk of

extinction, some populations are able to persist in small numbers and with low genetic

diversity (Xue et al. 2015; Benazzo et al. 2017; Robinson et al. 2018; Grossen et al. 2020).

Understanding the complex relationship between genetic diversity, inbreeding, mutational

load, and demographic history are key to leveraging this information for management and

conservation strategies of populations in the wild.

In large and outbred populations strongly deleterious recessive alleles are "hidden" from

selection in heterozygous form, making large populations more prone to experiencing

inbreeding depression after going through a harsh bottleneck (Robinson et al. 2018;

Kyriazis et al. 2021). On the other hand, small and isolated populations tend to have a

higher burden of deleterious variants as a result of inbreeding (Robinson et al. 2018;

Hedrick et al. 2019; Grossen et al. 2020; Khan et al. 2021; Stoffel et al. 2021). We observe

a similar trend in K'gari dingoes, finding higher levels of homozygosity in this population

compared to canids from large and stable populations, such as wolves and village dogs.

Accumulation of long ROH in the genome is associated with increased presence of

deleterious homozygotes and potential fitness effects (Szpiech et al. 2013; Szpiech et al.

2019; Stoffel et al. 2021). Although the levels of ROH and estimated inbreeding found in

K'gari dingoes (F_{ROH} = 0.59 on average; fig. 3) in some cases surpass those observed in

wild populations of other species already experiencing inbreeding depression (Khan et al.

2021; Stoffel et al. 2021; Colpitts et al. 2022), surveillance and necropsy records from the

Queensland Parks and Wildlife Service (QPWS) and the latest report on the genetic

health of K'gari dingoes (Allen 2023) have not recorded abnormal phenotypes in K'gari

dingoes. Nonetheless, whether a small population will experience inbreeding depression

or not depends on various factors, including the size of the ancestral population and the

magnitude and history of bottleneck events.

It has been observed that long-term small populations (i.e., populations that experienced

early bottlenecks and have maintained small population size following the bottlenecks)

can persist despite showing high levels of inbreeding by purging strongly recessive

deleterious variants (Xue et al. 2015; Benazzo et al. 2017; Robinson et al. 2018; Grossen

et al. 2020), whereas populations that have gone through multiple recent bottlenecks will

most likely experience inbreeding depression despite purging (Khan et al. 2021; Stoffel

et al. 2021). Previous studies have documented dingoes in K'gari have experienced at

least two bottlenecks in the form of large-scale culling events during the past 120 years

(Petrie 1995; Allen et al. 2015). Our assessment of purging in this population indicates

that while strongly deleterious alleles are likely being purged, there is little to no purging

of mildly deleterious alleles in this population. This may have important long term

consequences for the health of the K'gari dingoes that should be considered by

conservation and management programs.

Understanding the genetic factors that allow inbred populations to persist can aid in the

design and implementation of appropriate conservation and management strategies. One

of the main conservation strategies employed to assist highly inbred populations is

genetic rescue, which relies on the introduction of individuals from a large population

carrying genetic diversity into the inbred population. Although genetic rescue has been

used widely as a conservation strategy, there have been instances in which the benefits

have been short-lived, such is the case of Isle Royale wolves (Hedrick et al. 2014; Hedrick

et al. 2019). It should be noted that Isle Royale wolves are a relatively recent population,

having experienced fluctuations in population size ranging between ~15 and 30

individuals since establishing in Isle Royale around the 1940s (Peterson et al. 2014). The

introduction of individuals from a large population (with high genetic diversity) into this

inbred wolf population resulted in a higher burden of strongly deleterious variants, which

then led to inbreeding depression (Robinson et al. 2019; Kyriazis et al. 2021). On the

other hand, dingoes have inhabited Australia for at least 3,500 years (Milham and

Thompson 1976; Savolainen et al. 2004). Although the time of arrival of dingoes on K'gari

remains unknown, we find dingoes from the mainland and K'gari are genetically distinct.

Moreover, our results indicate that both dingo populations have lower genetic diversity

compared to large and outbred populations, which could make mainland dingoes a

potential source population to introduce genetic diversity in K'gari dingoes. However,

closely monitoring for signs of inbreeding depression and more research (e.g. population

genetic simulations) have been recommended under similar scenarios (Robinson et al.

2018; Kyriazis et al. 2021; Robinson et al. 2022) to explore the effectiveness of this

strategy, given that non-genetic factors (e.g. environmental stressors) may also play a

role in population persistence.

Current conservation and management programs treat dingoes as a single homogeneous

population, although studies using mtDNA (Savolainen et al. 2004; Oskarsson et al. 2012;

Cairns and Wilton 2016; Greig et al. 2018) and Y-chromosome (Ardalan et al. 2012; Sacks

et al. 2013; Cairns et al. 2017) data as well as genome-wide SNP data (Cairns et al. 2023)

and whole-genome sequencing (Zhang et al. 2020) have identified at least three distinct

dingo populations. Adding to previous evidence, we find two distinct populations in the

mainland (northwest and southeast), and one offshore in K'gari. However, we identify

long IBD segments shared between dingoes from K'gari and Northwest Australia, and

between dingoes from K'gari and the Southeast (suppl. Fig. 2). These patterns may be

due to differences in female and male dingo dispersal throughout mainland Australia, with

males having a broader dispersal range than females. Cairns et al. (Cairns et al. 2017)

found dingoes in the Northwest and K'gari carry the paternal haplogroup from Northwest

dingoes, and dingoes sampled from K'gari carry the maternal lineage from dingoes in the

Southeast. In light of this, recognizing several dingo subpopulations can help inform
conservation strategies.

Although dingoes are protected on K'gari, they are euthanized due to injury, poor health,
or as a result of high-risk behavior towards humans. Moreover, QPWS have documented
conspecific killing (Behrendorff et al. 2018), vehicle strikes and snake bite, and young
dingoes may be susceptible to predation such as goannas, snakes, and birds of prey.
There is considerable debate about how the population should be managed, both in terms
of conservation and how to reduce risk to tourists visiting the island. This work helps
broaden our knowledge and understanding about the evolutionary history and genetic
health of K'gari dingoes in order to better inform conservation and public policies. Our
work, although limited by a small sample size and the sampling of several closely related
samples, indicates that the dingo population on K'gari is challenged by inbreeding as a
result of founder effects and successive bottlenecks. We find the first evidence that K'gari
dingoes may be purging strongly deleterious mutations, whilst accumulating a higher
genetic load of mild/slightly deleterious mutations. Further research is needed to
investigate the genetic mutation load of dingoes on K'gari using a larger sample size, as
well as biological monitoring of the population for symptoms of inbreeding depression
such as reduced fertility, increased stillbirths, and congenital birth defects. QPWS and
Butchulla Prescribed Body Corporates are left with the challenge of preserving dingoes
on K'gari, balancing the need to address human safety concerns and protecting the dingo
population's genetic health and viability into the future.

ACKNOWLEDGEMENTS

The authors would like to acknowledge the Butchulla People as the traditional custodians of K'gari. Thanks to the Queensland Parks and Wildlife Service for access to the K'gari wongari (dingo) DNA sample collection, and thanks to Anna María Calderón for helpful feedback on early versions of this work. This work was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R35GM146926 (AVLA and ZAS), by start-up funds from the Pennsylvania State University's Department of Biology (AVLA and ZAS), and by a grant from the Australian Dingo Foundation (KMC). Computations for this research were performed using the Pennsylvania State University's Institute for Computational Data Sciences' Roar supercomputer.

SUPPLEMENTARY MATERIAL

Supplementary data are available at XXXX.

DATA ACCESSIBILITY

Raw sequence data of each of the K'gari dingo genomes generated in this study are available on SRA under project XXXX (TBD).

CONFLICTS OF INTEREST

KMC is a scientific advisor to the Australian Dingo Foundation, The New Guinea Singing Dog Conservation Society and The New Guinea Highland Wild Dog Foundation. KMC and LB are members of the IUCN Canid Specialist Group Dingo Working Group.

REFERENCES

Allen B. 2023. Genetic health and status of K'gari wongari (Fraser Island dingoes).

Queensland Government - Department of Environment and Science [Internet].

Available from:

https://parks.des.qld.gov.au/__data/assets/pdf_file/0022/319414/genetic-health-

status-kgari-wongari-summary.pdf

Allen BL, Higginbottom K, Bracks JH, Davies N, Baxter GS. 2015. Balancing dingo

conservation with human safety on Fraser Island: the numerical and demographic

effects of humane destruction of dingoes. *Australas. J. Environ. Manag.* 22:197–

215.

Ardalan A, Oskarsson M, Natanaelsson C, Wilton AN, Ahmadian A, Savolainen P.

2012. Narrow genetic basis for the Australian dingo confirmed through analysis of

paternal ancestry. *Genetica* 140:65–73.

Behrendorff L. 2021. Best-practice dingo management: six lessons from K'gari (Fraser

Island). *Aust. Zool.* 41:521–533.

Behrendorff L, Belonje G, Allen BL. 2018. Intraspecific killing behaviour of canids: how

dingoes kill dingoes. *Ethol. Ecol. Evol.* 30:88–98.

Benazzo A, Trucchi E, Cahill JA, Maisano Delser P, Mona S, Fumagalli M, Bunnefeld L,

Cornetti L, Ghirotto S, Girardi M, et al. 2017. Survival and divergence in a small

group: The extraordinary genomic history of the endangered Apennine brown bear

stragglers. *Proc. Natl. Acad. Sci. U. S. A.* 114:E9589–E9597.

Bergström A, Frantz L, Schmidt R, Ersmark E, Lebrasseur O, Girdland-Flink L, Lin AT,

Storå J, Sjögren K-G, Anthony D, et al. 2020. Origins and genetic legacy of

prehistoric dogs. *Science* 370:557–564.

Bird P, Bowman T. 2016. South Australian Wild Dog Strategic Plan: A plan to protect

the livestock industries and public safety whilst maintaining the integrity of the dingo

as a wildlife species. Department of Primary Industries and Regions South

Australia Available from:

https://static1.squarespace.com/static/612ee136efc1241e1fcc567e/t/633cdd4b6c57

c06c0843e819/1664933201509/SOUTH+AUSTRALIAN+WILD+DOG+STRATEGIC

+PLAN+2016+%E2%80%93+2020.pdf

Brink H, Purcell BV, Letnic M, Webster HS, Appleby RG, Jordan NR. 2019. Pets and

pests: a review of the contrasting economics and fortunes of dingoes and domestic

dogs in Australia, and a proposed new funding scheme for non-lethal dingo

management. *Wildl. Res.* 46:365.

Broad Institute. 2019. Picard Toolkit. *Broad Institute; GitHub repository* [Internet].

Available from: https://github.com/broadinstitute/picard

Browning BL, Browning SR. 2013. Improving the accuracy and efficiency of identity-by-

descent detection in population data. *Genetics* 194:459–471.

Brown SK, Pedersen NC, Jafarishorijeh S, Bannasch DL, Ahrens KD, Wu J-T, Okon M,

Sacks BN. 2011. Phylogenetic distinctiveness of Middle Eastern and Southeast

Asian village dog Y chromosomes illuminates dog origins. *PLoS One* 6:e28496.

Cairns KM. 2021. What is a dingo – origins, hybridisation and identity. *Aust. Zool.*

41:322–337.

Cairns KM, Brown SK, Sacks BN, Ballard JWO. 2017. Conservation implications for

dingoes from the maternal and paternal genome: Multiple populations, dog

introgression, and demography. *Ecol. Evol.* 7:9787–9807.

Cairns KM, Crowther MS, Nesbitt B, Letnic M. 2021. The myth of wild dogs in Australia:

are there any out there? *Aust. Mammal.* 44:67–75.

Cairns KM, Crowther MS, Parker HG, Ostrander EA, Letnic M. 2023. Genome-wide

variant analyses reveal new patterns of admixture and population structure in

Australian dingoes. *Mol. Ecol.* [Internet]. Available from:

http://dx.doi.org/10.1111/mec.16998

Cairns KM, Nesbitt BJ, Laffan SW, Letnic M, Crowther MS. 2020. Geographic hot spots

of dingo genetic ancestry in southeastern Australia despite hybridisation with

domestic dogs. *Conserv. Genet.* 21:77–90.

Cairns KM, Shannon LM, Koler-Matznick J, Ballard JWO, Boyko AR. 2018. Elucidating

biogeographical patterns in Australian native canids using genome wide SNPs.

PLoS One 13:e0198754.

Cairns KM, Wilton AN. 2016. New insights on the history of canids in Oceania based on

mitochondrial and nuclear data. *Genetica* 144:553–565.

Campbell CL, Bhérer C, Morrow BE, Boyko AR, Auton A. 2016. A Pedigree-Based Map
of Recombination in the Domestic Dog Genome. *G3* 6:3517–3524.

Colpitts J, McLoughlin PD, Poissant J. 2022. Runs of homozygosity in Sable Island feral
horses reveal the genomic consequences of inbreeding and divergence from
domestic breeds. *BMC Genomics* 23:501.

Conroy GC, Lamont RW, Bridges L, Stephens D, Wardell-Johnson A, Ogbourne SM.
2021. Conservation concerns associated with low genetic diversity for K'gari-Fraser
Island dingoes. *Sci. Rep.* 11:9503.

Corbett LK. 2001. The dingo in Australia and Asia. Sydney: University of NSW Press

Crowther MS, Fillios M, Colman N, Letnic M. 2014. An updated description of the
Australian dingo (Canis dingo Meyer, 1793). *J. Zool.* 293:192–203.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane
T, McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools.
Gigascience [Internet] 10. Available from:
https://academic.oup.com/gigascience/article/doi/10.1093/gigascience/giab008/613
7722

Delaneau O, Marchini J, 1000 Genomes Project Consortium, 1000 Genomes Project
Consortium. 2014. Integrating sequence and array data to create an improved 1000
Genomes Project haplotype reference panel. *Nat. Commun.* 5:3934.

DEPI. 2013. Action statement No. 248 dingo Canis lupus subsp. dingo. Department of

Environment and Primary Industries Available from:

https://www.wildlife.vic.gov.au/__data/assets/pdf_file/0031/27877/Dingo_Canis_lup

us-dingo-1.pdf

van Eeden LM, Crowther MS, Dickman CR, Newsome TM. 2021. Wicked "wild dogs":

Australian public awareness of and attitudes towards dingoes and dingo

management. *Aust. Zool.* 41:467–479.

van Eeden LM, Dickman CR, Newsome TM, Crowther MS. 2019. What should we do

with wild dogs? Taxonomic tangles and the management of dingo-dog

hybridisation. *Aust. Zool.* 40:92–101.

Elledge AE, Allen LR, Carlsson B-L, Wilton AN, Leung LK-P. 2008. An evaluation of

genetic analyses, skull morphology and visual appearance for assessing dingo

purity: implications for dingo conservation. *Wildl. Res.* 35:812.

Field MA, Yadav S, Dudchenko O, Esvaran M, Rosen BD, Skvortsova K, Edwards RJ,

Keilwagen J, Cochran BJ, Manandhar B, et al. 2022. The Australian dingo is an

early offshoot of modern breed dogs. *Sci Adv* 8:eabm5944.

Fleming P. 2001. Managing the Impacts of Dingoes and Other Wild Dogs. Bureau of

Rural Sciences

Frantz LAF, Mullin VE, Pionnier-Capitan M, Lebrasseur O, Ollivier M, Perri A,

Linderholm A, Mattiangeli V, Teasdale MD, Dimopoulos EA, et al. 2016. Genomic

and archaeological evidence suggest a dual origin of domestic dogs. *Science*

352:1228–1231.

Freedman AH, Wayne RK. 2017. Deciphering the Origin of Dogs: From Fossils to

Genomes. *Annu Rev Anim Biosci* 5:281–307.

Germonpré M, Lázničková-Galetová M, Sablin MV. 2012. Palaeolithic dog skulls at the

Gravettian Předmostí site, the Czech Republic. *J. Archaeol. Sci.* 39:184–202.

Germonpré M, Sablin MV, Stevens RE, Hedges REM, Hofreiter M, Stiller M, Després

VR. 2009. Fossil dogs and wolves from Palaeolithic sites in Belgium, the Ukraine

and Russia: osteometry, ancient DNA and stable isotopes. *J. Archaeol. Sci.*

36:473–490.

Glen AS, Dickman CR, Soulé ME, Mackey BG. 2007. Evaluating the role of the dingo as

a trophic regulator in Australian ecosystems. *Austral Ecol.* 32:492–501.

Greig K, Gosling A, Collins CJ, Boocock J, McDonald K, Addison DJ, Allen MS, David

B, Gibbs M, Higham CFW, et al. 2018. Complex history of dog (Canis familiaris)

origins and translocations in the Pacific revealed by ancient mitogenomes. *Sci.*

Rep. 8:9130.

Grossen C, Guillaume F, Keller LF, Croll D. 2020. Purging of highly deleterious

mutations through severe bottlenecks in Alpine ibex. *Nat. Commun.* 11:1001.

Hedrick PW, Peterson RO, Vucetich LM, Adams JR, Vucetich JA. 2014. Genetic rescue

in Isle Royale wolves: genetic analysis and the collapse of the population. *Conserv.*

Genet. 15:1111–1121.

Hedrick PW, Robinson JA, Peterson RO, Vucetich JA. 2019. Genetics and extinction

and the example of Isle Royale wolves. *Anim. Conserv.* 22:302–309.

Jackson SM, Fleming PJS, Eldridge MDB, Ingleby S, Flannery T, Johnson RN, Cooper

SJB, Mitchell KJ, Souilmi Y, Cooper A, et al. 2019. The Dogma of Dingoes-

Taxonomic status of the dingo: A reply to Smith et al. *Zootaxa*

4564:zootaxa.4564.1.7.

Jackson SM, Groves CP, Fleming PJS, Aplin KP, Eldridge MDB, Gonzalez A, Helgen

KM. 2017. The Wayward Dog: Is the Australian native dog or Dingo a distinct

species? *Zootaxa* 4317:201.

Johnson CN, Isaac JL, Fisher DO. 2007. Rarity of a top predator triggers continent-wide

collapse of mammal prey: dingoes and marsupials in Australia. *Proc. Biol. Sci.*

274:341–346.

Jones E. 1990. Physical characteristics and taxonomic status of wild canids, Canis-

familiaris, from the Eastern highlands of Victoria. *Wildl. Res.* 17:69.

Kardos M, Åkesson M, Fountain T, Flagstad Ø, Liberg O, Olason P, Sand H, Wabakken

P, Wikenros C, Ellegren H. 2018. Genomic consequences of intensive inbreeding

in an isolated wolf population. *Nat Ecol Evol* 2:124–131.

Khan A, Patel K, Shukla H, Viswanathan A, van der Valk T, Borthakur U, Nigam P,

Zachariah A, Jhala YV, Kardos M, et al. 2021. Genomic evidence for inbreeding

depression and purging of deleterious genetic variation in Indian tigers. *Proc. Natl.*

Acad. Sci. U. S. A. [Internet] 118. Available from:

http://dx.doi.org/10.1073/pnas.2023018118

Kirin M, McQuillan R, Franklin CS, Campbell H, McKeigue PM, Wilson JF. 2010.

Genomic runs of homozygosity record population history and consanguinity. *PLoS*

One 5:e13996.

Kumar M, Conroy G, Ogbourne S, Cairns K, Borburgh L, Subramanian S. 2023.

Genomic signatures of bottleneck and founder effects in dingoes. Available from:

http://biorxiv.org/lookup/doi/10.1101/2023.02.05.527211

Kyriazis CC, Wayne RK, Lohmueller KE. 2021. Strongly deleterious mutations are a

primary determinant of extinction risk due to inbreeding depression. *Evol Lett* 5:33–

47.

Larson G, Karlsson EK, Perri A, Webster MT, Ho SYW, Peters J, Stahl PW, Piper PJ,

Lingaas F, Fredholm M, et al. 2012. Rethinking dog domestication by integrating

genetics, archeology, and biogeography. *Proc. Natl. Acad. Sci. U. S. A.* 109:8878–

8883.

Letnic M, Crowther MS, Dickman CR, Ritchie EG. 2011. Demonising the dingo: How

much wild dogma is enough? *Curr. Zool.* 57:668–670.

Letnic M, Koch F, Gordon C, Crowther MS, Dickman CR. 2009. Keystone effects of an

alien top-predator stem extinctions of native mammals. *Proc. Biol. Sci.* 276:3249–

3256.

Letnic M, Ritchie EG, Dickman CR. 2012. Top predators as biodiversity regulators: the

dingo Canis lupus dingo as a case study. *Biol. Rev. Camb. Philos. Soc.* 87:390–

413.

Liberg O, Andrén H, Pedersen H-C, Sand H, Sejberg D, Wabakken P, Kesson M,

Bensch S. 2005. Severe inbreeding depression in a wild wolf (Canis lupus)

population. *Biol. Lett.* 1:17–20.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler

transform. *Bioinformatics* 25:1754–1760.

Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen W-M. 2010. Robust

relationship inference in genome-wide association studies. *Bioinformatics* 26:2867–

2873.

McLaren W, Pritchard B, Rios D, Chen Y, Flicek P, Cunningham F. 2010. Deriving the

consequences of genomic variants with the Ensembl API and SNP Effect Predictor.

Bioinformatics [Internet] 26:2069–2070. Available from:

http://dx.doi.org/10.1093/bioinformatics/btq330

McPeek MS, Wu X, Ober C. 2004. Best linear unbiased allele-frequency estimation in

complex pedigrees. *Biometrics* 60:359–367.

Milham P, Thompson P. 1976. Relative antiquity of human occupation and extinct fauna

at madura cave, southeastern western Australia. *Mankind* 10:175–180.

Newsome AE, Corbett LK. 1985. The identity of the dingo III.* the incidence of dingoes,

dogs and hybrids and their coat colours in remote and settled regions of Australia.

Aust. J. Zool. 33:363.

NWDAP. 2020. National Wild dog Action Plan 2020–2030: Promoting and supporting

community-driven action for landscape-scale wild dog management. Australian

Wool Innovation Ltd Available from: https://wilddogplan.org.au/wp-

content/themes/nwdap/docs/NWDAP2020-2030.pdf

O'Neill AJ, Cairns KM, Kaplan G, Healy E. 2017. Managing dingoes on Fraser Island:

culling, conflict, and an alternative. *Pac. Conserv. Biol.* 23:4.

Oskarsson MCR, Klütsch CFC, Boonyaprakob U, Wilton A, Tanabe Y, Savolainen P.

2012. Mitochondrial DNA data indicate an introduction through Mainland Southeast

Asia for Australian dingoes and Polynesian domestic dogs. *Proc. Biol. Sci.*

279:967–974.

Pemberton TJ, Absher D, Feldman MW, Myers RM, Rosenberg NA, Li JZ. 2012.

Genomic patterns of homozygosity in worldwide human populations. *Am. J. Hum.*

Genet. 91:275–292.

Peterson RO, Vucetich JA, Bump JM, Smith DW. 2014. Trophic cascades in a

multicausal world: Isle Royale and Yellowstone. *Annu. Rev. Ecol. Evol. Syst.*

45:325–345.

Petrie R. 1995. Early Days On Fraser Island 1913-1922 - Rollo Petrie: 9780646276809

- AbeBooks. *Go Bush Safaris* [Internet]. Available from:

https://www.abebooks.com/9780646276809/Early-Days-Fraser-Island-1913-1922-

0646276808/plp

Plassais J, Kim J, Davis BW, Karyadi DM, Hogan AN, Harris AC, Decker B, Parker HG,

Ostrander EA. 2019. Whole genome sequencing of canids reveals genomic regions

under selection and variants influencing morphology. *Nat. Commun.* 10:1489.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar

P, de Bakker PIW, Daly MJ, et al. 2007. PLINK: a tool set for whole-genome

association and population-based linkage analyses. *Am. J. Hum. Genet.* 81:559–

575.

Räikkönen J, Bignert A, Mortensen P, Fernholm B. 2006. Congenital defects in a highly

inbred wild wolf population (Canis lupus). *Mamm. Biol.* 71:65–73.

R Core Team. 2022. R: A language and environment for statistical computing. Available

from: https://www.r-project.org/

Robinson JA, Brown C, Kim BY, Lohmueller KE, Wayne RK. 2018. Purging of Strongly

Deleterious Mutations Explains Long-Term Persistence and Absence of Inbreeding

Depression in Island Foxes. *Curr. Biol.* 28:3487–3494.e4.

Robinson JA, Kyriazis CC, Nigenda-Morales SF, Beichman AC, Rojas-Bracho L,

Robertson KM, Fontaine MC, Wayne RK, Lohmueller KE, Taylor BL, et al. 2022.

The critically endangered vaquita is not doomed to extinction by inbreeding

depression. *Science* 376:635–639.

Robinson JA, Ortega-Del Vecchyo D, Fan Z, Kim BY, vonHoldt BM, Marsden CD,

Lohmueller KE, Wayne RK. 2016. Genomic Flatlining in the Endangered Island

Fox. *Curr. Biol.* 26:1183–1189.

Robinson JA, Räikkönen J, Vucetich LM, Vucetich JA, Peterson RO, Lohmueller KE,

Wayne RK. 2019. Genomic signatures of extensive inbreeding in Isle Royale

wolves, a population on the threshold of extinction. *Sci Adv* 5:eaau0757.

Roelke ME, Martenson JS, O'Brien SJ. 1993. The consequences of demographic

reduction and genetic depletion in the endangered Florida panther. *Curr. Biol.*

3:340–350.

Sacks BN, Brown SK, Stephens D, Pedersen NC, Wu J-T, Berry O. 2013. Y

chromosome analysis of dingoes and southeast asian village dogs suggests a

neolithic continental expansion from Southeast Asia followed by multiple

Austronesian dispersals. *Mol. Biol. Evol.* 30:1103–1118.

Savolainen P, Leitner T, Wilton AN, Matisoo-Smith E, Lundeberg J. 2004. A detailed

picture of the origin of the Australian dingo, obtained from the study of

mitochondrial DNA. *Proc. Natl. Acad. Sci. U. S. A.* 101:12387–12390.

Skoglund P, Ersmark E, Palkopoulou E, Dalén L. 2015. Ancient wolf genome reveals an

early divergence of domestic dog ancestors and admixture into high-latitude

breeds. *Curr. Biol.* 25:1515–1519.

Smith BP, Cairns KM, Adams JW, Newsome TM, Fillios M, Déaux EC, Parr WCH,

Letnic M, VAN Eeden LM, Appleby RG, et al. 2019. Taxonomic status of the

Australian dingo: the case for Canis dingo Meyer, 1793. *Zootaxa*

4564:zootaxa.4564.1.6.

Stephens D, Wilton AN, Fleming PJS, Berry O. 2015. Death by sex in an Australian

icon: a continent-wide survey reveals extensive hybridization between dingoes and

domestic dogs. *Mol. Ecol.* 24:5643–5656.

Stoffel MA, Johnston SE, Pilkington JG, Pemberton JM. 2021. Mutation load decreases

with haplotype age in wild Soay sheep. *Evol Lett* 5:187–195.

Surbakti S, Parker HG, McIntyre JK, Maury HK, Cairns KM, Selvig M, Pangau-Adam M,

Safonpo A, Numberi L, Runtuboi DYP, et al. 2020. New Guinea highland wild dogs

are the original New Guinea singing dogs. *Proc. Natl. Acad. Sci. U. S. A.*

117:24369–24376.

Szpiech Z. 2014. asd. *GitHub* [Internet]. Available from: https://github.com/szpiech/asd

Szpiech ZA, Blant A, Pemberton TJ. 2017. *GARLIC*: Genomic Autozygosity Regions

Likelihood-based Inference and Classification. *Bioinformatics* [Internet] 33:2059–

2062. Available from: http://dx.doi.org/10.1093/bioinformatics/btx102

Szpiech ZA, Jakobsson M, Rosenberg NA. 2008. ADZE: a rarefaction approach for

counting alleles private to combinations of populations. *Bioinformatics* 24:2498–

2504.

Szpiech ZA, Mak ACY, White MJ, Hu D, Eng C, Burchard EG, Hernandez RD. 2019.

Ancestry-Dependent Enrichment of Deleterious Homozygotes in Runs of

Homozygosity. *Am. J. Hum. Genet.* 105:747–762.

Szpiech ZA, Xu J, Pemberton TJ, Peng W, Zöllner S, Rosenberg NA, Li JZ. 2013. Long

runs of homozygosity are enriched for deleterious variation. *Am. J. Hum. Genet.*

93:90–102.

Vaser R, Adusumalli S, Leng SN, Sikic M, Ng PC. 2016. SIFT missense predictions for

genomes. *Nat. Protoc.* 11:1–9.

Wallach AD, Johnson CN, Ritchie EG, O'Neill AJ. 2010. Predator control promotes

invasive dominated ecological states. *Ecol. Lett.* 13:1008–1018.

Wardell-Johnson G, Schoeman D, Schlacher T, Wardell-Johnson A, Weston MA,

Shimizu Y, Conroy G. 2015. Re-framing values for a World Heritage future: what

type of icon will K'gari-Fraser Island become? *Australas. J. Environ. Manag.*

22:124–148.

Wilton AN. 2001. DNA methods of assessing dingo purity. In: A Symposium on the

Dingo. P.O. Box 20, Mosman NSW 2088, Australia: Royal Zoological Society of

New South Wales. p. 49–56.

Wootton E, Robert C, Taillon J, Côté SD, Shafer ABA. 2023. Genomic health is

dependent on long-term population demographic history. *Mol. Ecol.* 32:1943–1954.

Xue Y, Prado-Martinez J, Sudmant PH, Narasimhan V, Ayub Q, Szpak M, Frandsen P,

Chen Y, Yngvadottir B, Cooper DN, et al. 2015. Mountain gorilla genomes reveal

the impact of long-term population decline and inbreeding. *Science* 348:242–245.

Zhang S-J, Wang G-D, Ma P, Zhang L-L, Yin T-T, Liu Y-H, Otecko NO, Wang M, Ma Y-

P, Wang L, et al. 2020. Genomic regions under selection in the feralization of the

dingoes. *Nat. Commun.* 11:671.

 FIGURES

Figure 1. Reduced dimensional representation of sample clustering based on multidimensional scaling of the allele sharing dissimilarity matrix. First two major axes of variation plotted for (A) all 41 canids included in this study and (B) 18 dingo individuals.

Figure 2. Genetic diversity estimates within and between canid populations. Error bars

representing 95% confidence intervals for each line are too narrow to be visible. (A) Mean

number of alleles per locus (allelic richness). (B) Mean number of private alleles per locus

(private allelic richness). (C) Mean number of shared alleles per locus between pairs of

populations.

Figure 3. ROH coverage across individual genomes. The percentage of the genome

covered by class A ROH (0.5-1 Mbp), class B ROH (1-2 Mbp), class C ROH (2-3 Mbp),

class D ROH (3-4 Mbp), class E ROH (4-5 Mbp), and class F ROH (>5 Mbp) is shown on

the Y-axis for each of the 41 canids included in this study.

Figure 4. Predicted damaging alleles. (A) Predicted deleterious variants. (B) Loss-of-

Function (LoF) variants. For each category solid bars show the proportion of homozygous

sites, and striped bars show the proportion of heterozygous sites for each canid individual

of each population.

852 Figure 5. Kinship-weighted R_{xy} estimates for K'gari and mainland dingoes. Estimates

853 represent the relative excess number of derived alleles for each category in K'gari

854 dingoes (population X) relative to mainland dingoes (population Y). Error bars represent

855 the minimum and maximum values for the 95% confidence interval from jackknife.