The interferon/STAT1 signaling axis is a common feature of tumor-initiating cells in breast

cancer

Eric P. Souto[1,2,3], Ping Gong[1], John D. Landua[1], R. Rajaram Srinivasan[1], Abhinaya Ganesan[1,2,3],

Lacey E. Dobrolecki[1], Stephen C. Purdy[5], Xingxin Pan[6], Michael Zeosky[6], Anna Chung[6], S.

Stephen Yi[6], Heide L. Ford[5], Michael T. Lewis[1,2,3,4*]

[1]Lester and Sue Smith Breast Center, Baylor College of Medicine, Houston, TX, 77030, USA.

[2]Cancer and Cell Biology Graduate Program, Baylor College of Medicine, Houston, TX, 77030,

USA.

[3]Dan L Duncan Comprehensive Cancer Center, Baylor College of Medicine, Houston, TX, 77030,

USA.

[4]Departments of Molecular and Cellular Biology and Radiology, Baylor College of Medicine,

Houston, TX, 77030, USA.

[5]Department of Pharmacology, University of Colorado Anschutz Medical Campus (UC-AMC),

Aurora, CO, 80045, USA; Pharmacology Graduate Program, UC-AMC, Aurora, CO, 80045, USA;

University of Colorado Cancer Center, UC-AMC, Aurora, CO, 80045, USA.

[6]Livestrong Cancer Institutes and Department of Biomedical Engineering, The University of Texas

at Austin, Austin, TX 78712, USA; Oden Institute for Computational Engineering and Sciences

and Interdisciplinary Life Sciences Graduate Programs, The University of Texas at Austin, Austin,

TX 78712, USA

[*]Corresponding Author

Michael T. Lewis

Baylor College of Medicine

One Baylor Plaza

BCM600; Room N1210

TEL: 713-798-1538

Email: mtlewis@bcm.edu

Running Title: Interferon/STAT1 signaling in tumor-initiating cells

DECLARATION OF INTERESTS

ABSTRACT

A tumor cell subpopulation of tumor-initiating cells (TIC), or "cancer stem cells", are associated with therapeutic resistance, as well as both local and distant recurrence. Enriched populations of TIC are identified by markers including aldehyde dehydrogenase (ALDH1) activity, the cell surface marker combination CD44$^+$/CD24$^-$, or fluorescent reporters for signaling pathways that regulate TIC function. We showed previously that Signal Transducer and Activator of Transcription (STAT)-mediated transcription allows enrichment for TIC in claudin-low models of human triple-negative breast cancer using a STAT-responsive reporter. However, the molecular phenotypes of STAT TIC are not well understood, and there is no existing method to lineage-trace TIC as they undergo cell state changes. Using a new STAT-responsive lineage-tracing (LT) system in conjunction with our original reporter, we enriched for cells with enhanced mammosphere-forming potential in some, but not all, basal-like triple-negative breast cancer (TNBC) xenograft models (TNBC) indicating TIC-related and TIC-independent functions for STAT signaling. Single-cell RNA sequencing (scRNAseq) of reporter-tagged xenografts and clinical samples identified a common interferon (IFN)/STAT1-associated transcriptional state, previously linked to inflammation and macrophage differentiation, in TIC. Surprisingly, most of the genes we identified are not present in previously published TIC signatures derived using bulk RNA sequencing. Finally, we demonstrated that bone marrow stromal cell antigen 2 (BST2), is a cell surface marker of this state, and that it functionally regulates TIC frequency. These results suggest TIC may exploit the IFN/STAT1 signaling axis to promote their activity, and that targeting this pathway may help eliminate TIC.

61 **Significance**

62 TIC differentially express interferon response genes, which were not previously reported in bulk

63 RNA sequencing-derived TIC signatures, highlighting the importance of coupling single-cell

64 transcriptomics with enrichment to derive TIC signatures.

INTRODUCTION

Breast cancer is a heterogeneous disease that can be classified into several histological and molecular subtypes (1,2). Triple-negative breast cancer (TNBC) is a histological subtype defined clinically by lack of estrogen receptor (ER) and/or progesterone receptor (PR), and without amplification or overexpression of the ERBB2 gene (HER2), and is an aggressive disease with poor outcomes (3).

There exists considerable intertumoral and intratumoral heterogeneity within each histological subtype. Intertumoral heterogeneity refers to variations across tumors, whereas intratumoral heterogeneity refers to the presence of molecularly distinct tumor cell populations within the same tumor. Both present major challenges clinically, particularly in TNBC, which itself represents at least six molecularly distinct subtypes (4,5). Due to the relative lack of targeted therapeutics, TNBC is typically treated with combination chemotherapies, with or without immuno-oncology agents or radiation.

With respect to intratumoral heterogeneity, there exists a subpopulation of cells, typically rare, termed tumor-initiating cells (TIC) (a.k.a. cancer stem cells (CSC)). TIC are defined functionally by their ability to self-renew, and to recapitulate clonally-derived cellular hierarchies present in the tumor from which they were isolated upon transplantation (6). Enriched TIC populations can be identified by aldehyde dehydrogenase (ALDH1) activity (7), the cell surface marker combination of $CD44^+/CD24^{low/-}$ (8), or fluorescent signaling reporters expressing pathway-specific transcription factors that regulate TIC function, including Wnt/β-catenin, Hedgehog, and Notch signaling, as well as POU5F1 (OCT4)/SOX2, and <u>S</u>ignal <u>T</u>ransducer and <u>A</u>ctivator of <u>T</u>ranscription 3 (STAT3)-mediated signaling (9-14). However, no single marker is present in all breast cancers.

Following neoadjuvant (before surgery) taxane- or anthracyclin-based chemotherapy in TNBC, residual breast cancer cells are enriched for TIC markers, including $CD44^+/CD24^{low/-}$ and aldehyde

dehydrogenase (ALDH1) activity, and isolated cells show enhanced mammosphere-forming

efficiency (MSFE), indicating that TIC are intrinsically resistant to some systemic therapies (15-

18). Thus, standard cancer therapies can eliminate many "bulk" tumor cells to shrink tumors, but

TIC survive preferentially, leading to local and/or distant recurrence. Indeed, some metastatic

tumor cells express the TIC marker CD44, and such cells have enhanced tumor-initiating capacity

(19). Thus, selective targeting of TIC, especially at metastatic sites, may be essential for effective

treatment. Unfortunately, TIC-specific assays to evaluate drug efficacy are rarely done in clinical

trials.

Here, we exploit a STAT signaling reporter, and a new STAT-responsive lineage-tracing (LT)

system, to enrich cells with enhanced mammosphere-forming potential in some, but not all, non-

claudin-low xenograft models of TNBC. Single-cell RNA sequencing (scRNAseq) on three

xenograft models identified a distinct interferon (IFN)/STAT1-associated transcriptional state in

TIC that is also present in most clinical samples evaluated, yet contains genes that have not been

included in at least five TIC signatures derived through bulk RNA sequencing. Finally, we

validated our signature by demonstrating that one of the genes, bone marrow stromal antigen 2

(BST2), can enrich for TIC, and regulates TIC frequency, but not self-renewal. Taken together,

our data suggest that TIC exploit the IFN-mediated response pathway to regulate stem cell

frequency independent of self-renewal. Thus, this pathway may represent an attractive candidate

therapeutic target in a subset of breast cancers.

MATERIALS & METHODS

STAT lineage-tracing (LT) system construction

The STAT reporter (4M67-EGFP-P2A-CreER[T2]) was generated by LR Gateway® Cloning of a

promoter donor vector and a gene donor vector into a destination vector. To generate a flexible

promoter donor vector, the EF-1α promoter sequence from the L4/R1 EF-1α promoter donor

vector (ThermoFisher Scientific, cat. A11146) was replaced with a 194bp sequence containing

multiple restriction digestion sites to facilitate simple restriction cloning of any desired promoter.

The STAT-specific promoter donor vector was generated by inserting the STAT signaling-specific

4M67 promoter from the STAT regulatory reporter (4M67-EGFP) (11) into the flexible promoter

donor vector to generate the L4-4M67-R1 promoter donor vector. To generate the gene donor

vector, a CreER[T2] sequence with attB1/B2 recombination sites and an EcoRI cutting site between

attB1 and CreER[T2] was amplified from pCAG-CreER[T2] (20) (a gift from Connie Cepko: Addgene

plasmid #14797) by 2-step PCR, then incorporated into the pDonor233 vector (a kind gift from Dr.

Kenneth L. Scott, Baylor College of Medicine, Houston, TX) with the P1/P2 compatible

recombination sites by BP Gateway cloning. Next, GFP-P2A was amplified from pUltra (21) and

inserted in front of CreER[T2] using InFusion cloning to generate L1-EGFP-P2A-CreER[T2]-L2 gene

donor vector. Gateway® Cloning was used to incorporate the 4M67 promoter donor vector and

EGFP-P2A-CreER[T2] gene donor vector into the pLenti6.4/R4R2/V5-Dest destination vector

(ThermoFisher Scientific, cat. A11146), generating 4M67-EGFP-P2A-CreER[T2].

To generate the dual color-switching lineage-tracing reporter (EFS-loxP-dsRed-loxP-Nep), the

loxP-dsRed-loxP-GFP sequence with attB1/B2 recombination sites was amplified from pMSCV-

loxP-dsRed-loxP-eGFP-Puro-WPRE (22) (a gift from Dr. Hans Clevers: Addgene plasmid

#32702) by 2-step PCR and then incorporated into the pDonor233 vector with the P1/P2

compatible recombination sites by BP Gateway cloning. Gateway® Cloning was used to

incorporate the EFS promoter donor vector and loxP-DsRed-loxP-EGFP gene donor vector into

the pLenti6.4/R4R2/V5-Dest destination vector (ThermoFisher Scientific, cat. A11146).

mNeptune2 was amplified from the mNeptune2-C1 vector (a kind gift from Dr. Michael Davidson;

Addgene plasmid #54836) and used to replace the EGFP sequence by restriction cloning,

generating EFS-$_{loxP}$dsRed$_{loxP}$-Nep.

Lentivirus production and transduction

Lineage-tracing plasmids and envelope-expressing plasmids were prepared using a plasmid maxi

kit (QIAGEN cat. 12162). To produce lentivirus, lineage-tracing plasmids and envelope-

expressing plasmids were transfected into HEK293T cells using jetPRIME® transfection reagent

(Polyplus, cat. 114-15). Lentiviral supernatants were collected every 24 hours for three days, then

concentrated. Concentrated viruses were titered using the p24-Elisa kit (Takara Bio, cat. 632200)

following manufacturer's protocols. For lentiviral transduction, 5mg/mL polybrene (Santa Cruz

Biotechnology, cat. sc-134220) was added to cell lines prior to lentiviral spinfection at 2500 RPM,

25^{o}C for 30 minutes (MOI of 5). Transduced cells were incubated at 37^{o}C, 5% CO_2 for 24 hours,

then lentivirus media was replaced with growth media. To generate LT cell lines (HepG2/C3A

STAT LT, SUM159 STAT LT, and MDA-MB-231 STAT LT), 4M67-GFP-P2A-CreER^{T2} was

transduced first, and transduced cells were selected using 10mg/mL blasticidin (Thermo Fisher

Scientific, cat. A1113903). Following blasticidin selection, cell lines were then transduced with

EFS-$_{loxP}$dsRed$_{loxP}$-Nep. Transduced cells were selected by FACS-enrichment of dsRed$^+$ cells. To

generate BST2 knockdown cell lines, SUM159 cells were lentiviral transduced with either one of

two shRNA vectors (Horizon Discovery, sh-BST2 #1: cat. RHS3979-201824601, sh-BST2 #2: cat.

RHS3979-201829090) or one non-targeting control vector (Horizon Discovery, cat. RHS6848) as

described above. Transduced cells were selected using 1μg/mL puromycin (InvivoGen, cat. ant-

pr-1).

Xenograft tumors were dissociated (isolation of single cells from human xenograft tumors). The

resulting cells were transduced with the STAT regulatory reporter (4M67-GFP) (11) overnight in

suspension in complete Mammary Epithelial Cell Growth Media (MEGM⁺) (Lonza, cat. CC-3153,

with supplements, cat. CC-4136) containing 20ng/mL basic fibroblast growth factor (Invitrogen,

cat. 13256-029), 10ng/mL epidermal growth factor (Invitrogen, cat. PHG0311), 4µg/mL Heparin

(StemCell Technologies, cat. 07980), and 40 nL/mL B27 (Invitrogen, cat. 17504-044).

Cell culture and *in vitro* treatment

HepG2/C3A, MDA-MB-231, and HEK293T cells were purchased from ATCC and were cultured

under adherent conditions with phenol red-free DMEM (Gibco, cat. 31053-028), 10% charcoal-

stripped FBS (Gibco, cat. 12-676-029), Penicillin-Streptomycin (GenDEPOT, cat. CA005-010),

1mM sodium pyruvate (Life Tech, cat. 11360070), and 2mM L-Glutamine (Thermo Fisher

Scientific, cat. 25030081). SUM159 cells were purchased from BioIVT and were cultured under

adherent conditions with phenol red-free Ham's F12 (Caisson labs, cat. HFL05-500ML) containing

5% charcoal-stripped FBS, 5mg/mL insulin (Gemini Bioproducts, cat. 700-112P), 1mg/mL

hydrocortisone (Sigma-Aldrich, cat. H-4001), and Penicillin-Streptomycin (GenDEPOT, cat.

CA005-010). Cells were incubated in a 5% CO_2 incubator at 37°C. All cell lines were authenticated

using short tandem repeat profiling, which was performed by the Cytogenetics and Cell Line

Authentication Core at MD Anderson Cancer Center (Houston, TX). Mycoplasma detection was

routinely performed using Venor®GeM qOneStep Mycoplasma detection kit (Minerva Biolabs,

cat. 11-91100). All experiments were performed with five passages of STR profiling and

mycoplasma testing.

Human recombinant IL-6 (Sigma-Aldrich, cat. I1395) was prepared as a 50mg/mL stock solution

in PBS/1%BSA. Recombinant human IFN alpha (IFNα) (Bio-Techne, cat. NBP2-35891-100ug)

and recombinant human IFN gamma (IFNγ) (Bio-Techne, cat. 285-IF-100) were prepared as a

200mg/mL stock in PBS. Tamoxifen (TAM) (Sigma-Aldrich, cat. T5648) was prepared as a 5mM

solution in 100% ethanol. HepG2/C3A STAT LT cells were treated with either IL-6 (50ng/mL),

TAM (1μM), or the combination and the cells were visualized by confocal microscopy (Leica, TCS-

SP5) and quantified by flow cytometry (BD LSRII analyzer) after 72 hours.

Generation of cell line-derived and patient-derived xenografts

To prepare for cell injection, SUM159 STAT LT or MDA-MB-231 STAT LT cells were suspended

in a 1:1 solution of their respective growth media and phenol red-free Matrigel® matrix (Corning,

cat. 356237). Cells were transplanted into the #4 epithelium-free "cleared" fat pad of 3–4-week-

old SCID/Beige mice (Charles River Laboratories, Wilmington, MA, http://www.criver.com). When

palpable tumors formed, mice were administered TAM by intraperitoneal injection to induce

dsRed to mNeptune2 color switching in STAT signaling (EGFP[+]) cells. TAM was prepared by

dissolving in sunflower seed oil (Sigma-Aldrich, cat. S5007) and stirring overnight. All animal

protocols were reviewed and approved by the Animal Protocol Review Committee at Baylor

College of Medicine.

Isolation of single cells from human xenograft tumors

Xenograft tumors were harvested and digested using 5mL digestion media (phenol red-free

DMEM/F12 [Corning cat. 16-405-CV], 100mg/mL gentamicin [Worthington, cat. LS004176],

250U/mL collagenase II [Worthington cat. LS004174] and 2μg/mL DNase I [Stemcell, cat. 07900])

per 1g tumor at 37°C in a gentleMACS dissociator (Miltenyi). The tumor digestion was passed

through a 200mM filter and centrifuged at 300g for 1 minute. Red blood cells were lysed by adding

9mL ddH$_2$O to the cell pellet for 20 seconds and lysis was stopped by the addition of 1mL 10X

PBS. To dissociate cell clumps further, the cell suspension was centrifuged at 300 g for 1 minute,

then resuspended in TrypLE Express (Gibco, cat. 12605-010) for 5 minutes. To stop the digestion,

9mL of PBS was added, and the cell suspension was passed through a 40mm cell strainer.

Analytical flow cytometry and FACS

Cells from adherent cultures were trypsinized and pelleted by centrifuging at 300g for 3 minutes.

The cells were resuspended in HBSS[+] (HBSS [Cytiva Life Sciences cat. SH30588.01], 2% FBS).

Single cells from digested xenograft tumors were labeled using a BUV395 anti-mouse cocktail

(pooled and diluted 1:40: anti-mouse H2kD [BD Biosciences, cat. 742437], anti-mouse CD45 [BD

Biosciences, cat. 567451], anti-mouse CD31 [BD Biosciences, cat. 740231], and anti-mouse

Ter119 [BD Biosciences, cat. 563827]) to exclude mouse cells. A far-red BST2 antibody was used

to label BST2-expressing cells (diluted 1:40, BioLegend, cat. 348416). CD44 was labeled with a

PE conjugated antibody (diluted 1:20, BD Biosciences, cat. 566803), and CD24 was labeled with

a FITC conjugated antibody (diluted 1:20, BD Biosciences, cat. 655154). Cell viability was

determined by Sytox Blue (diluted 1:1000, V450, Invitrogen, cat. S34857) staining. Cells were

applied to a BD LSRII analyzer for analytical flow cytometry or a BD AriaII for fluorescence-

activated cell sorting (FACS). Quantification of each cell population was performed using FlowJo

(BD Biosciences) and statistical differences between cell populations were determined by one-

way ANOVA followed by post-hoc Tukey's test, which was calculated using GraphPad Prism

version 9 (GraphPad Software, San Diego, CA, http://www.graphpad.com).

ALDEFLUOR assay

ALDH activity was assessed using the ALDEFLUOR kit (StemCell technologies, cat. 01700)

according to the manufacturer's protocol. Briefly, cells were incubated in ALDEFLUOR assay

buffer containing ALDH substrate for 30 minutes at $37^\circ C$. In each experiment, a sample of cells

was stained under identical conditions with diethylaminobenzaldehyde (DEAB), a specific ALDH

inhibitor, as a negative control. Any antibody incubation was performed immediately after

ALDEFLUOR incubation, as described in the "analytical flow cytometry and FACS" section.

Propidium Iodide cell cycle assay

Single cell suspensions were fixed in 70% ethanol overnight at $-20^\circ C$. Then, cells were washed

twice with PBS and stained in $100\mu L$ propidium iodide solution (0.001% Triton X-100, 0.2mg/mL

DNase-free RNase A (Thermo Fisher, cat. 12091-021), $20\mu g/mL$ propidium iodide (Thermo

Fisher, cat. P3566)) for 15 minutes at 37°C. After incubations, the cells were immediately applied

to a BD LSRII analyzer for analytical flow cytometry.

Immunohistochemistry and Immunofluorescence

Immunohistochemistry (IHC) and immunofluorescence (IF) were performed by the Lester and

Sue Smith Breast Center Pathology Core. Tumors were fixed in 4% paraformaldehyde, paraffin-

embedded, sectioned at 3–4 µm, and deparaffinized. The antigen retrieval buffer for pSTAT1,

pSTAT3, and pSTAT5 staining was 0.1M Tris-HCL PH 9.0. Sections were then blocked in 3%

hydrogen peroxide solution for 5 minutes and incubated by pSTAT1 Ser727 antibody (Cell

Signaling Technology, cat. 8826, 1:200), pSTAT3 Y705 antibody (Cell Signaling Technology, cat.

9145, 1:25) or pSTAT5 antibody Tyr694 (Cell Signaling Technology, cat. 9359, 1:25), followed

by Signalstain Boost IHC Detection Reagent (HRP, Rabbit) (Cell Signaling Technology, cat.

8114) for 30 minutes, visualized by 3, 3′ - Diaminobenzidine (DAB+) solution (DakoCytomation,

Glostrup, Denmark), and counterstained by Harris hematoxylin. SUM159 tumors, known for

activation of pSTAT3 (11), were used as a positive control for pSTAT3, and a human tissue array,

which included the spleen, was used as a positive control for pSTAT1.

For IF, Five IF images were quantified for each PDX model using CellProfiler (23). 0.1M Tris-HCL

PH 9.0. was used as the antigen retrieval buffer, then sections were blocked in 3% hydrogen

peroxide solution for 5 minutes and incubated with pSTAT1 Ser727 antibody (Cell Signaling

Technology, cat. 8826, 1:200), pSTAT3 Y705 antibody (Cell Signaling Technology, cat.

9145, 1:25), or a GFP antibody (Fisher Scientific, cat. NC9777966) overnight at 4°C. The following

day, slides were incubated with secondary antibodies: anti-mouse (for pSTAT1 and pSTAT3,

Thermo Fisher, cat. A-11032) and/or anti-rabbit (for GFP, Thermo Fisher, cat. A-11008) for one

hour at room temperature. Slides were then counterstained with DAPI and imaged. For each PDX

model, five representative images were quantified using FIJI (24).

Mammosphere-forming efficiency assay

Following tumor digestion and FACS enrichment of fluorescent cell populations, equal numbers

of cells from each fraction were plated in complete Mammary Epithelial Cell Growth Media

(MEGM⁺) (Lonza, cat. CC-3153, with supplements, cat. CC-4136) containing 20ng/mL basic

fibroblast growth factor (Invitrogen, cat. 13256-029), 10ng/mL epidermal growth factor (Invitrogen,

cat. PHG0311), 4µg/mL Heparin (StemCell Technologies, cat. 07980), and 400nL/mL B27

(Invitrogen, cat. 17504-044) with 0.6% HPMC (Millipore Sigma, cat. M7027) to prevent cell

aggregation, in an ultra-low attachment plate.

To prepare secondary mammospheres, primary mammospheres were collected and dissociated

into single cells using TrypLE Express (Gibco, cat. 12605-010) for 5 minutes. To inactivate the

TrypLE Express, 9mL of PBS was added, and the cell suspension was passed through a 40mm

cell strainer. Then, an equal number of cells from each fraction were plated in MEGM⁺ media with

0.6% HPMC. Secondary mammospheres were imaged on a Yokogawa CV8000, and

CellPathfinder (Yokogawa) was used to quantify the number of mammospheres in each well.

Spheres with an area greater than 300mm^2 and a diameter greater than 50mm were used to

determine mammospheres. A BioTek Cytation 5 (Agilent, Santa Clara, CA) was used to take

representative images. Statistical differences in mammosphere forming efficiency between cell

populations were determined by one-way ANOVA followed by post-hoc Tukey's test, which was

calculated using GraphPad Prism version 9 (GraphPad Software, San Diego, CA,

http://www.graphpad.com).

Limiting-dilution transplantation assay

FACS-enriched cells were suspended in growth media at a concentration of 5,000 cells per 10µl

(SUM159) or 50,000 cells per 10µl (BCM-4272) and serially diluted using growth media. Four

dilutions were prepared (SUM159: 5,000, 1,000, 200, 40 cells per 10µl, BCM-4272: 50,000, 5,000,

500, 50 cells per 10µl), then phenol red-free Matrigel® matrix (Corning, cat. 356237) was added

for a final 1:1 solution of growth media and Matrigel® before transplantation as described in

"Generation of cell line-derived and patient-derived xenografts." The mice were monitored until

the first tumor reached 10% body weight (~7-8 weeks) at which point the experiment was

terminated and tumor outgrowth was recorded. Extreme limiting-dilution analysis was used to

calculate the TIC frequency of each population (25).

scRNAseq library prep and sequencing

SUM159, BCM-4272, and BCM-15006 were sorted based on fluorescent reporter activity

(SUM159: EGFP$^+$/mNeptune2$^+$, EGFP$^+$/dsRed$^+$, EGFP$^-$/mNeptune2$^+$, EGFP$^-$/dsRed$^+$; BCM-4272

and BCM-15006: EGFP$^+$ or EGFP$^-$), and single-cell gene expression libraries were prepared

according to Chromium Single Cell Gene Expression 3v3.1 kit (10x Genomics). In brief, single

cells, reverse transcription (RT) reagents, Gel Beads containing barcoded oligonucleotides, and

oil were loaded on a Chromium controller (10x Genomics) to generate single-cell GEMS (Gel

Beads-In-Emulsions) where full-length cDNA was synthesized and barcoded for each single cell.

Subsequently the GEMS are broken and cDNA from each single cell are pooled. Following

cleanup using Dynabeads MyOne Silane Beads, cDNA is amplified by PCR. The amplified

product is fragmented to optimal size before end-repair, A-tailing, and adaptor ligation. The final

library was generated by amplification. Finally, single-cell sequencing libraries were sequenced

using NovaSeq 6000 (read configuration: 28-10-10-90; 1% PhiX Spike-in; read depth

200M/sample).

For single nuclei sequencing, Tissue processing and nuclei RNA fixation from PDX samples were

performed using Chromium Next GEM Single Cell Fixed RNA Sample Preparation Kit (10x

Genomics). Fixation was then quenched and samples were stored at -80°C until sequencing.

Single nuclei RNAseq libraries were prepared according to Chromium Single Cell Gene

Expression 3v3.1 kit (10x Genomics), similar to the cDNA synthesis and barcoding process for

each single cell. Single-cell nuclei RNA libraries were sequenced using NovaSeq (read

configuration: Paired End 28-10-10-90; 1% PhiX Spike-in; read depth 100M/sample).

scRNAseq data preprocessing and filtering

FASTQ files were mapped to the GRCh38 reference genome using the Cell Ranger Single Cell software v.6.1.2 (SUM159) or v7.0.1 (BCM-4272 and BCM-15006) (10X Genomics). Single nuclei sequencing data were integrated with the Texas Advanced Computing Center (TACC) supercomputer resources for data analysis and interpretation. In brief, gene expression counts for individual nuclei were generated from raw FASTQ files using Cell Ranger (version 3). Reads with low mapping quality (<10) or that map to multiple genomic locations were filtered out. Using Seurat v4.1.3 in R v4.2.2, the count matrix of each sample was used to create a Seurat object. Cells expressing greater than 1000 and less than 9000 genes. Cells with greater than 10% mitochondrial transcripts were excluded as they represented low-quality or dying cells. Next, fluorescent samples from the same tumor were merged into a single Seurat object (ex. BCM-4272 EGFP+ and BCM-4272 EGFP- merged to BCM-4272). To exclude the effects of cell cycle heterogeneity in the analysis, we used the CellCycleScoring function from Seurat. This function gives each cell a quantitative score for G1, G2/M, and S phases based on the expression of cell cycle phase genes (26). Subsequently, effects of cell cycle heterogeneity were regressed out of the data during data scaling.

scRNAseq clustering and UMAP visualization

Seurat v4.1.3 was used for data normalization, dimensionality reduction, and clustering (27). The filtered gene matrices from each merged sample were normalized using the NormalizeData function. The normalized datasets were centered and scaled using ScaleData. Then, the most significant principal components were found using RunPCA, and ElbowPlot was used to determine the number of principal components to use for clustering each dataset. Different resolution parameters for unsupervised clustering were examined to find the optimal number of clusters based on clear segregation of clusters. For each dataset, the first 35 principal components were used for unsupervised clustering with a resolution = 0.1 for SUM159 (yielding

nine clusters) and a resolution of 0.5 for BCM-4272 and BCM-15006 (yielding seven clusters and

six clusters, respectively) using the FindNeighbors and FindClusters functions. To visualize the

clustering for each dataset, the dimensionality was further reduced using the Seurat function

RunUMAP. The same principal components used to determine the clustering were the same used

to calculate the UMAP embedding. Each resulting cluster was further analyzed for potential

doublets or low-quality cells using two methods: quality metrics (nCount_RNA, nFeature_RNA,

and mitochondrial content) for each cluster were calculated and any outlier values (clusters with

greater, or less than, two standard deviations than the average of all clusters) were removed.

Three PDX models from an independent, single nuclei RNA seq, dataset were processed,

clustered, and visualized as described above. UCell was used to score each cell based on

similarity the 25 genes identified by intersecting SUM159-C7, 4272-C7, and 15006-C6 (28).

scRNAseq differential gene expression analysis and intersection of gene lists

To examine the different transcriptional states of tumor cells, FindAllMarkers was used to identify

differentially expressed genes in each cluster using the "bimod" likelihood-ratio test framework.

Two additional parameters were used to include differentially expressed genes: genes that

showed at least a 0.25 log2 fold expression difference, and genes that were expressed in at least

25% of the cells in a cluster. To compare the similarities between candidate TIC transcriptional

states in different xenograft models, upregulated genes from each candidate TIC cluster were

intersected using GeneOverlap (29). To examine the similarity between the TIC genes we

identified and previously published TIC signatures derived using bulk RNA sequencing (30-34),

we intersected the TIC genes common between any two xenograft models with previously

published TIC signatures using GeneOverlap (29). To examine the extent to which the 25

common TIC genes are unique to a given transcriptional state in each xenograft model, we

performed Marker Enrichment Modeling (MEM) using the 25 common TIC genes across models

as MEM labels. Similarity scores based on MEM labels were calculated and visualized with a

heatmap using the MEM_RMSD function (35).

scRNAseq gene set enrichment analysis

To associate a biological phenotype with each cluster, gene set enrichment analysis was

performed using ClusterProfiler (36). Differentially expressed genes from each cluster, as

determine using the FindAllMarkers function described above, with a log fold change threshold of

0.25 were included in the analysis. Gene sets with a false discovery rate (FDR) < 0.01 and a

Benjamini-Hochberg (BH)-adjusted p-value < 0.05 were considered significantly enriched, and

these cutoffs were established prior to analyzing the data.

Analysis of patient scRNAseq data

Publicly available human breast cancer datasets were retrieved through GEO (GSE161529,

GSE176078) (37,38). Seurat v4.1.3 was used for cell cycle regression, data normalization,

dimensionality reduction, and clustering, as described above. To identify clusters of epithelial

cells, the expression of keratins (KRT8, KRT14, KRT18, and KRT19) was evaluated. Clusters

with no keratin expression were filtered, and the epithelial dataset was re-normalized and

clustered using Seurat v4.1.3. UCell was used to score every cell in each patient samples with

more than 1,500 non-cycling epithelial cells based on similarity using the 25 genes identified by

intersecting SUM159-C7, 4272-C7, and 15006-C6 (28). Harmony was used to integrate common

cell states across patient samples (39). The patient samples were integrated by breast cancer

subtype, pre-neoplastic, or normal breast. UCell was used to score each cell in the integrated

datasets based on similarity using the 25 genes identified by intersecting SUM159-C7, 4272-C7,

and 15006-C6 (28). A two-sided Wilcoxon test was used to determine significance of TIC scores.

To identify differentially expressed genes between clusters, FindAllMarkers from Seurat v4.1.3

was used. To compare the similarities between clusters with high TIC scores across breast cancer

subtypes and our 25-gene signature, upregulated genes from each cluster were intersected using

GeneOverlap (29).

Patient survival and chemotherapy response

384   To determine whether there is any association between our 25-gene TIC signature and overall

385   survival in breast cancer patients, we used the RNAseq datasets from KM-plotter (40). ROC

386   plotter was used to determine whether our 25-gene TIC signature is associated with

387   chemotherapy response in breast cancer patients (41). Statistical significance was set at $p < 0.05$

388   before the analysis was performed.

qRT-PCR

RNA extraction was performed using the RNeasy plus mini kit (Qiagen, cat. 74134) according to

the manufacturer's protocol. Next, we performed cDNA synthesis using SuperScript™ III first-

strand synthesis system (ThermoFisher Scientific, cat. 18080051) according to the

manufacturer's protocol. Then, qPCR was performed using iTaq qPCR supermix (BioRad, cat.

1725130). Briefly, 10μL 2X Taq, 9μL cDNA, and 1μL of a 20X TaqMan probe (ThermoFisher

Scientific, cat. Hs00171632_m1 (BST2), Hs00241666_m1 (ADAR), Hs00942643_m1 (OAS2),

Hs00967084_m1   (PARP9),   Hs00393816_m1   (PARP14),   Hs01013996_m1   (STAT1),

Hs99999901_s1 (18S)) were added per reaction. The qPCR reaction was performed in a C1000-

Touch-Thermocycler (BioRad) (Initial denaturation: 95°C 30 seconds, followed by 39 cycles of

denaturation: 95°C 5 seconds and annealing/extension 60°C 30 seconds). Samples were

normalized to the expression of a housekeeping gene, 18S. Statistical differences in BST2

expression following chemotherapy were determined for each PDX independently by one-way

ANOVA followed by post-hoc Tukey's test. Statistical differences in the expression of IFN

response genes were determined for each model independently by one-way ANOVA followed by

post-hoc Tukey's test. All statistical differences were calculated using GraphPad Prism version 9

(GraphPad Software, San Diego, CA, http://www.graphpad.com).

Jess™ Automated Western Blot

BST2 expression levels in BST2 knockdown cells relative to the non-targeting control cells was

evaluated using ProteinSimple's Jess™ fully automated western blot analysis (Bio-Techne, cat.

004-650) according to the manufacturer's protocol. A total of 0.5mg/mL protein was loaded per

sample and a BST2 antibody (1:25, Novus Biologicals, cat. NBP2-27154) was used to detect

BST2 levels in each sample. The ProteinSimple's Jess™ protein normalization was used to

normalize variation in protein loading.

RESULTS

STAT reporter activity is heterogenous in a cohort of triple-negative breast cancer PDX

models

We showed previously that STAT-mediated transcriptional activity allows enrichment for TIC in

two cell line xenograft (CDX) models of claudin-low breast cancer (SUM159 and MDA-MB-231)

(11). While our focus was on STAT3 in these two models, STAT1 and STAT3 transcription factors

bind similar DNA sequences, and both can activate the reporter (42).

To determine whether STAT signaling might also be associated with TIC function in a broader

range of TNBC, we screened a cohort of 56 TNBC PDX models for pSTAT3 expression, of which

38 (68%) had detectable pSTAT3 staining in the epithelium (Supplementary Table 1). This value

is higher than, but arguably consistent with, results using clinical samples of TNBC (~37.5% (43))

given the known engraftment bias present in PDX collections toward immunologically "cold"

tumors (44).

The 38 pSTAT3 positive TNBC models were then cross-referenced with PDX showing behaviors

of interest with respect to putative roles of TIC, specifically local recurrence after apparent

complete response to one or more chemotherapies, and/or metastasis to one or more organ sites

(44,45). Based on these properties, a subset of 11 PDX models of TNBC were chosen for genetic

tagging with the 4M67-EGFP lentiviral vector (11). Nine PDX showed varying degrees of EGFP

expression, whereas candidate negative control models BCM-0046 and WHIM12, showed no

detectable EGFP expression (Supplementary Table 2). Of nine EGFP+ PDX, five models (BCM-

3611, BCM-4013, BCM-5097, HCI-028, and HCI-002) were excluded from further consideration

because the percentage of EGFP$^+$ cells was too low (<1.2%) to be experimentally tractable,

thereby yielding four remaining PDX models for use.

The chosen PDX were then evaluated for pSTAT1, pSTAT3, and pSTAT5 by IHC, as well as

EGFP expression by immunofluorescence (Fig. 1A, Fig. S1A). Consistent with pSTAT1/3

positivity by IHC, PDX models displayed their anticipated STAT reporter activity via detection of

EGFP (Fig. 1A). Reporter-positive PDX models did not exhibit pSTAT5 staining, indicating STAT5

signaling is not responsible for reporter activity. In EGFP$^+$ PDX, analytical flow cytometry (Fig.

1B) showed a continuum of EGFP reporter activity across models ranging from 7.88% (BCM-

15006) to 31.7% (MC1) of tumor cells, ranges that should allow for discrimination of differential

TIC function/gene expression in EGFP$^+$ vs. EGFP$^-$ cells, should such differences exist. Although

EGFP percentage is comparable to pSTAT1 and pSTAT3 percentage (Fig. 1A), co-IF staining

revealed little direct correlation between the reporter and pSTAT1/3 (Fig. S1B), likely due to the

26-hour half-life of GFP (46) relative to the comparatively rapid rate of STAT dephosphorylation

(47).

To determine whether STAT reporter activity enriches for cells with increased MSFE in reporter-

positive TNBC PDX models (a surrogate for TIC function), we sorted EGFP$^+$, EGFP$^-$, and a mock

sorted control from each reporter-positive model and plated them under low attachment conditions

(Fig. 1C). STAT reporter positive cells in BCM-4272 had an 8.04% MSFE, representing a more

than 100% increase relative to the EGFP$^-$ cell population (3.74% MSFE), and the MSFE of the

EGFP$^-$ population was reduced by 59% relative to a mock sorted control (6.3% MSFE) (Fig. 1C).

In contrast, STAT reporter activity did not enrich for cells with enhanced MSFE in BCM-15006,

BCM-7821, or MC1 (Fig. 1C). Thus, while the 4M67 reporter can identify cells with enhanced

MSFE in some TNBC, STAT-mediated transcription must have TIC-independent function(s).

A STAT lineage-tracing reporter provides a long-term fluorescent label to STAT signaling

cells

Given that TIC marker expression, including STAT transcriptional activity, is dynamically

regulated (48), the inability to lineage-trace STAT signaling cells as they undergo cell state

changes, or differentiate, is a major limitation of the 4M67-EGFP reporter.

To augment the 4M67-EGFP reporter, we developed a TAM-activated, Cre recombinase-

dependent, STAT signaling-specific, lentiviral LT system (Fig. S2A–B). This system consists of

two components: the first is a lentiviral vector that labels cells with active STAT signaling (EGFP$^+$),

followed by a self-cleaving P2A peptide and a TAM-activated Cre recombinase (4M67-EGFP-

P2A-CreER^{T2}) (Fig. S2A). The second component is a constitutively-expressed dual-color

switching Cre-dependent reporter vector (EFS-$_{loxP}$dsRed$_{loxP}$-mNeptune2) (Fig. S2B). Addition of

TAM drives color switching from red (dsRed) to far-red (mNeptune2) via CreER^{T2} recombination

in STAT signaling cells within a specific temporal window where the TAM concentration remains

high enough to activate Cre recombinase (6.8-hour half-life) (49) (Fig. 2A). Use of the LT system

results in four cell populations: long-lived (label-retaining) STAT signaling cells

(EGFP$^+$/mNeptune2$^+$), transient STAT signaling cells post-TAM (or cells that did not undergo Cre

recombination) (EGFP$^+$/dsRed$^+$), lineage-tagged cells (EGFP$^-$/mNeptune2$^+$), and bulk tumor cells

(EGFP$^-$/dsRed$^+$) (Fig. 2B).

To demonstrate the sensitivity and specificity of the STAT LT reporter system, we used

HepG2/C3A cells (IL-6 responsive) in a STAT3-dependent manner (11,50). Untreated STAT LT-

transduced HepG2/C3A cells (HepG2/C3A STAT LT) had minimal EGFP and mNeptune2

expression. TAM (1µM) induced mNeptune2 expression in a small percentage of cells. Addition

of IL-6 (50ng/mL) induced the EGFP reporter activity, without mNeptune2 activation, but the

addition of both IL-6 and TAM induced EGFP and mNeptune2 expression (Fig. S2C–D). These

results demonstrate the LT reporter system is both stringent and sensitive to STAT signaling, and

TAM addition induced mNeptune2 expression in STAT-responsive cells.

To demonstrate the ability of the LT system to provide a long-term fluorescent label on STAT

signaling cells *in vivo*, we introduced the LT reporters into the two TNBC cell lines used previously

to demonstrate STAT reporter activity enriches for TIC (SUM159 STAT LT and MDA-MB-231

STAT LT) (11). To ensure effective lineage-tagging of STAT signaling cells, we performed an *in*

vivo TAM dose-response analysis (Fig. 2SE–F). Given that high levels of TAM can cause toxicity

in mice (51), we aimed to find a dose that resulted in maximal loxP recombination without any

adverse effects. At 4mg, the recombination efficiency (the percentage of EGFP$^+$/Nep$^+$ cells out of

all EGFP$^+$ cells) two days after TAM treatment was 71%. The recombination efficiency, and the

total percentage of mNeptune2$^+$ cells, decreased with each successive dose, with the lowest dose

(0.006mg) being comparable to vehicle. Given high recombination with no observable toxicity at

4mg TAM, we used this dose for the remainder of our studies.

SUM159 STAT LT or MDA-MB-231 STAT LT cell lines were engrafted into SCID/beige mice, and

tumor-bearing mice were treated with 4mg TAM when tumors reached 100mm^3. Two days after

TAM treatment, we visualized all four LT cell populations in situ using confocal microscopy and

quantified their abundance in each tumor model using flow cytometry (Fig. 2C–F). In both cell line

xenografts, TAM treatment increased the percentage of EGFP$^+$/Nep$^+$ cells (from 0.25% to 1.8%

in SUM159, and from 0.02% to 0.07% in MDA-MB-231), which was accompanied by a decrease

in EGFP$^+$/dsRed$^+$ cells (from 3.08% to 0.7% in SUM159 from 0.7% to 0.4% in MDA-MB-231).

Likewise, TAM treatment increased the percentage of mNeptune2$^+$ cells (from 1.98% to 8.88% in

SUM159 and from 0.07% to 3.15% in MDA-MB-231) with a corresponding decrease in the

percentage of dsRed$^+$ cells (from 94.7% to 88.3% in SUM159 and from 96.7% to 93.3% in MDA-

MB-231). These results indicate that TAM treatment leads to Cre-dependent mNeptune2

expression *in vivo*, demonstrating the utility of the STAT LT system to study STAT TIC biology.

Long-lived STAT signaling cells in SUM159 xenografts have enhanced mammosphere-

forming efficiency

To confirm that the STAT LT system enriches for cells with enhanced MSFE from SUM159 tumors

in a manner similar to that of the 4M67-EGFP vector shown previously (11), we dissociated

SUM159 xenografts eight days after TAM treatment, sorted the four fluorescent cell populations

(EGFP$^+$/mNeptune2$^+$, EGFP$^+$/dsRed$^+$, EGFP$^-$/mNeptune2$^+$, and EGFP$^-$/dsRed$^+$), and performed

MSFE assays (Fig. 3A–B). Long-lived, label-retaining STAT signaling cells (EGFP$^+$/mNeptune2$^+$)

had a significantly higher MSFE (7.2%) compared to all other cell populations, which were not

statistically different from each other.

The cancer stem cell hypothesis postulates that cellular heterogeneity in tumors results, in part,

from hierarchical organization of cell types present, and that these hierarchies are fueled by rare,

self-renewing TIC (52). We reasoned that long-lived STAT signaling cells give rise to progenitor

cells (EGFP$^-$/mNeptune2$^+$) that will differentiate into non-TIC bulk tumor cells over time.

To test this hypothesis, we performed a 16-day time course study. We harvested SUM159 tumors

prior to TAM treatment, and again at days 2, 4, 8, and 16 after the TAM pulse (Fig. 3C–D). A total

of 74% of the EGFP$^+$ cells recombined (EGFP$^+$/dsRed$^+$ converted to EGFP$^+$/mNeptune2$^+$ cells)

two days after the TAM pulse. Between days 0 and 16, the percentage of EGFP$^-$/Nep$^+$ cells

increased from 2% to 35% of the tumor (p < 0.001), indicating lineage-tagged cells proliferate and

displace the EGFP$^-$/dsRed$^+$ cells. Furthermore, we noted the re-emergence of EGFP$^+$/dsRed$^+$

cells on the 8th day, suggesting STAT signaling activation in cells not defined by STAT signaling

at the time of the TAM pulse.

To extend the timeframe of this experiment, we serially transplanted tumor fragments two days

after the TAM pulse for a total of three transplant generations (TG) (Fig. 3E–F). Each TG was

allowed to grow for 21 days and harvested. The percentage of lineage-tagged cells increased

from 18% to 35% from TG1 to TG2, then to 61% in TG3. Furthermore, the total percentage of

EGFP$^+$/mNeptune2$^+$ cells remained unchanged over time and following serial passaging,

indicating tight regulation of STAT signaling and TIC frequency.

Interferon response genes are differentially expressed in TIC

To determine whether STAT reporter-positive cells express a distinct set of genes that may

regulate TIC biology, we FACS-enriched all reporter populations from three TNBC xenograft

models: two models where STAT reporter activity enriched for TIC (SUM159 and BCM-4272),

and one model where STAT reporter activity did not enrich for TIC (BCM-15006), and performed

scRNAseq (Fig. 4A).

To determine whether EGFP reporter activity could be used to identify TIC transcriptional states

in each model, we overlayed sample information on Uniform Manifold Approximation and

Projection (UMAP) plots to visualize whether reporter positive cells clustered together (Fig. 4B,

Fig. S3B–D). Then, we looked at the percent representation of each cluster within each reporter

population to identify clusters that were overrepresented in EGFP positive samples. In SUM159,

though all clusters are represented in the EGFP$^+$ cells, positive cells are highly overrepresented

in cluster 7 (SUM159-C7) (9.8% of the EGFP$^+$/mNeptune2$^+$, 8.2% of the EGFP$^+$/dsRed$^+$ cells, vs.

0.1% of mNeptune2$^+$ cells, and 0.07% of dsRed$^+$ cells) (Fig. 4B, Fig S3B), suggesting it represents

a TIC transcriptional state. In BCM-4272, clusters 0 (35% of EGFP$^+$ cells, 0.5% of EGFP$^-$ cells),

2 (25% of EGFP$^+$ cells, 10% of EGFP$^-$ cells), 4 (22% of EGFP$^+$ cells, 0.7% of EGFP$^-$ cells), and 7

(0.5% of EGFP$^+$ cells, 0.1% of EGFP$^-$ cells) are highly represented in the EGFP$^+$ cells, suggesting

at least one of these clusters may represent a TIC transcriptional state (Fig. 4B, Fig. S3A).

To clarify which of these clusters most likely represents a TIC transcriptional state, we performed

differential gene expression analysis between these four clusters and found that cluster 7

differentially expressed genes associated with TIC, including *ABCD1*, *CXCL11*, *NOTCH1*, *RELA*,

and *TCF7*, suggesting cluster 7 (4272-C7) represents TIC (Fig. S3B).

Because STAT reporter activity did not enrich TIC (Fig. 1C–D) in BCM-15006, we reasoned that

a cluster representing a TIC transcriptional state would not be overrepresented in either EGFP$^+$

and EGFP$^-$ cells. Cluster 6 (15006-C6) is the only cluster that was equally represented in both

EGFP$^+$ and EGFP$^-$ cells (2.1% of EGFP$^+$, 1.9% of EGFP$^-$ cells), suggesting this cluster may

represent a TIC transcriptional state (Fig. 4B, Fig S3A).

To determine whether established TIC markers CD44$^+$/CD24$^{low/-}$ and ALDH1A1 could be used to

confirm the identification of TIC transcriptional states in our data, we generated feature plots

based on their expression (Fig. S3C). In all three xenografts, CD44 had no distinct expression

pattern and CD24 was highly expressed in all cells in BCM-4272 and BCM-15006, but not

expressed in many SUM159 cells. Furthermore, ALDH1A1 was not a distinct marker of any

clusters.

To gain insights into the biological states of each cluster, we used the differentially expressed

genes that define the cells in a given cluster relative to all other cells to perform GSEA

(Supplementary Tables 4–6). SUM159-C7, 4272-C7, and 15006-C6, which we hypothesized may

represent a TIC transcriptional state, were defined by enrichment of IFNγ response, IFNα

response, and TNF-alpha signaling via NF-kB (Fig. 4D, Supplementary Table 3). To investigate

the extent to which candidate TIC clusters share transcriptional features, we intersected the

differentially expressed genes from each cluster and performed Fisher's exact test for significance

(Fig. 4E). SUM159-C7 and 4272-C7 shared 48 features (p=1.8e^{-24}), SUM159-C7 and 15006-C6

shared 38 features (p=5.7e^{-27}), and 4272-C7 and 15006-C6 shared 79 features (p=1.1e^{-110}). In

this process, we identified only 25 common up-regulated genes across all three clusters, that may

represent a new TIC signature. A list of genes from all intersections is shown in Supplementary

Table 7.

To determine whether these 25 genes participate in shared biological processes, we performed

a functional protein-protein interaction analysis using the STRING database (53) (Fig. 4F). We

found all of our genes, except GNB4, exist within the same protein-protein interaction network

(PPI enrichment: p<1.0e^{-16}), which suggests the genes in our TIC signature may be part of a

biological process responsible their regulation. To uncover this biological process, we performed

an over-representation analysis using the 25-gene TIC signature, which confirmed these genes

are overrepresented in gene sets and pathways related to IFN signaling (Fig. 4F, Fig. S4).

To determine whether our 25-gene signature can effectively distinguish cell states within each

xenograft, we used marker enrichment modeling (35) to generate similarity scores based on the

expression of each gene in the 25-gene signature and visualized the results with a heatmap (Fig.

4G). In the PDX models, 4272-C7 and 15006-C6 clustered separately from every other cell state

in their respective xenografts, demonstrating the 25-gene signature is unique to these clusters. In

contrast, SUM159-C7 is a member of the same clade as cluster 5, indicating cluster 5 also is

defined by the expression of IFN genes. To distinguish these two clusters, we performed

differential gene expression analysis and found SUM159-C7 differentially expressed genes

associated with TIC, including *TP63*, *POU5F1*, *ABCA5*, *FOXO1*, and *TCF4*, supporting our

hypothesis that SUM159-C7 represents TIC (Fig. S3D).

To determine whether additional PDX models have a similar transcriptional state, we analyzed an

independent dataset of three PDX models sequenced using isolated nuclei rather than whole cells

and gave each cell a TIC score based on the expression of the genes in our 25-gene signature

(Fig. S5). All three PDX models (BCM-4013, BCM-15166, and WHIM30) had a single cluster with

a high TIC score relative to all other clusters. Thus, our TIC signature defines distinct cell states

in three additional PDXs, suggesting this cell state may be represented in a wider cohort of TNBC

PDXs.

To confirm that IFNs can induce EGFP in cells transduced with the STAT reporter, we treated

SUM159 STAT LT, BCM-4272 4M67-EGFP, and BCM-15006 4M67-EGFP cells with IFNα or IFNγ

and confirmed that IFNγ activated the reporter in all three models more robustly than did IFNα.

IFNα did not activate the reporter in SUM159 (Fig. S6A). To determine whether IFN treatment

activates genes that define the IFN-associated TIC transcriptional state, we input our 25 genes

into INTERFEROME (54), an online database of IFN-regulated genes, which revealed all 25

genes are regulated by IFN (Fig. S6B).

To demonstrate these genes can be regulated by IFN in our models, we compared the expression

of three upstream regulators of IFN signaling (PARP9, PARP14, STAT1) and three genes

activated in response to IFN signaling (ADAR, OAS2, and BST2) from our 25 intersected TIC

genes after IFN treatment to untreated controls by qRT-PCR (Fig. S6C). IFNα or IFNγ treatment

increased the expression of all six genes relative to the untreated controls in all three TNBC

models.

To evaluate whether any candidate TIC genes we identified have been previously reported, we

compared intersections from any two xenograft models to five previously published TIC signatures

that were derived using bulk RNA sequencing, including our own (30-34), and a previously

published panel of stem cell and epithelial-to-mesenchymal transition markers (55)

(Supplementary Table 8). None of the 25 genes we identified overlapped significantly with any of

the six previously published TIC signatures. The Pece et al. (33) signature contained one of our

TIC genes. LY6E, from the 4272-C7 and 15006-C6 intersection, whereas the Colacino et al. (55)

custom panel only contained CDH3 (SUM159-C7 and 4272-C7 intersection) and MCL1 and

ITGA6 (SUM159-C7 and 15006-C6 intersection). Furthermore, a comparison of these previously

published TIC signatures did not reveal any significant gene-level commonalities, suggesting that

single cell methods may be required for accurate identification of TIC.

Interferon response genes are differentially expressed in a subset of epithelial cells in

human breast cancer patient samples

To determine whether a similar IFN/STAT1 transcriptional state could be detected in human

clinical samples, we analyzed scRNAseq data retrieved from a published human breast cancer

cell atlas that contained TNBC, HER2$^+$, ER$^+$, pre-neoplastic (BRCA1$^{+/-}$), and normal breast

samples (37). To identify epithelial cells from breast cancer patients with a similar transcriptional

state to our IFN/STAT1 TIC, we gave each cell a TIC score based on the expression of the genes

in our 25-gene TIC signature (Fig. 5A–C). This identified cells in cluster 7 in TNBC patients

(p=3.42e^{-113}), cluster 6 in HER2$^+$ patients (p=3.16e^{-287}), and cluster 3 in ER$^+$ patients (p=4.71e^{-96})

had the highest TIC scores relative to cells in all other clusters. To determine whether established

TIC markers CD44$^+$/CD24$^{low/-}$ and ALDH1A1 could be used to identify TIC transcriptional states

in patient data, we generated feature plots based on their expression (Fig. S7A). In all three breast

cancer subtypes, established TIC markers were not markers of any particular cluster, similar to

our observations in PDX models.

To determine whether a similar transcriptional state exists in pre-neoplastic or normal breast

tissue, we applied the same TIC scoring method (Fig. 5D–E). This revealed a similar IFN/STAT1

transcriptional state in pre-neoplastic tissue (cluster 2, p=$4.76e^{-115}$), but not in normal breast.

These results suggest the IFN/STAT1 transcriptional state is only present in breast cancer and

pre-malignant tissue.

To determine the extent to which clusters with high TIC scores share transcriptional features with

our 25-gene TIC signature, we intersected the genes from each cluster with a high TIC score from

each breast cancer subtype, then intersected that gene list with our 25-genes TIC signature (Fig.

5F). This revealed the IFN/STAT1 transcriptional state from breast cancer patients shared 9

genes with our 25-gene signature (Fisher's exact test: p=$9.6e^{-23}$).

To determine whether this similarity extends to pre-neoplastic tissue, we intersected the genes

from the high TIC score cluster from pre-neoplastic samples with the high TIC score intersected

gene list from all breast cancer patients, then intersected that list with our 25-gene signature (Fig.

5G). This revealed the IFN/STAT1 transcriptional state from both breast cancer patients and pre-

neoplastic tissue shared 4 genes with our 25-gene signature (Fisher's exact test: p=$2.7e^{-11}$).

To determine whether the IFN/STAT1 transcriptional state can be detected in individual breast

cancer patients, we analyzed each breast cancer patient of each subtype with more than 1,500

non-cycling epithelial cells to increase our chances of identifying rare cell subpopulations, such

as TIC (8) (Fig. S7B). We found that the IFN/STAT1 transcriptional state in most, but not all,

patients analyzed, which is consistent with TIC heterogeneity. To establish the robustness of the

TIC signature, we examined an additional cohort of TNBC patients from an additional human

breast cancer scRNAseq dataset (38). We gave each cell a TIC score based on the expression

of the genes in our 25-gene TIC signature (Fig. S8A), which revealed cluster 5 had a significantly

higher TIC score relative to all other clusters (p=4.5e^{-113}). Furthermore, the only patient with more

than 1,500 non-cycling epithelial cells also had a cluster defined by a higher TIC score (cluster 4,

p=2.3e^{-03}) (Fig. S8B). Taken together, our findings reveal that the IFN/STAT1 signaling axis we

observe in PDX models is present in human breast cancer patients.

The IFN/STAT1 signaling axis is associated with worse overall survival and lack of

response to chemotherapy in breast cancer patients

To evaluate the clinical significance of our TIC signature, we examine the association between

high expression of our 25-gene TIC signature and overall survival using RNAseq datasets from

KM-plotter (40). In all breast cancer patients, regardless of treatment status, the 25-gene TIC

signature was associated with a worse overall survival (HR=1.46, p=0.0067) (Fig. 6A). This

association with worse overall survival was more pronounced when stratifying for untreated

patients (HR=2.33, p=0.0022), but was eliminated when stratifying for patients treated with

chemotherapy (HR=0.77, p=0.23) (Fig. 6A), suggesting the TIC signature is only prognostic in

untreated patients.

Next, we stratified patients by PAM50 subtype and examined differences in overall survival in

untreated patients compared to chemotherapy-treated patients (Fig. 6B). In basal, luminal A, and

luminal B untreated patients, there was a trend of high TIC score with worse overall survival,

which reduced in luminal A chemotherapy-treated patients or reversed in basal and luminal B

chemotherapy-treated patients (Fig. 6B). High expression of TIC gene was associated with

favorable overall survival in HER2 untreated patients, but this was reduced in chemotherapy-

treated patients (Fig. 6B). Normal-like patients had a similar overall survival regardless of

chemotherapy treatment (Fig. 6B).

We and others showed previously that TIC demonstrate intrinsic resistance to systemic

chemotherapies. Thus, we evaluated whether expression of our 25-gene TIC signature is

associated with response to chemotherapy in breast cancer patients using ROC plotter (41).

Consistent with intrinsic chemotherapy resistance, high expression of our TIC signature was

associated with non-responders, and was predictive of lack of response to chemotherapy (AUC:

0.593, p=$9.7e^{-04}$) (Fig. 6C). Furthermore, this finding was consistent for all breast cancer subtypes,

except for luminal A (Fig. 6C).

BST2 allows TIC enrichment in breast cancer xenograft models.

To demonstrate conclusively that the TIC transcriptional state identifies a subpopulation enriched

for TIC function, we leveraged the observation that BST2 was a FACS-compatible cell surface

found in the intersection between IFN/STAT1 transcriptional states from all breast cancer

subtypes and pre-neoplastic tissue and our 25-gene signature. We performed analytical flow

cytometry to examine the relationship between BST2 protein levels and TIC marker

CD44$^+$/CD24$^{low/-}$ and ALDH activity. In SUM159 xenografts, BST2$^+$ cells are ~40% of the total

tumor but represent 96% of CD44$^+$/CD24$^{low/-}$ cells and 58% of ALDH$^+$ cells. In BCM-4272

xenografts, BST2$^+$ cells represented ~20% of the tumor, but 38% of CD44$^+$/CD24$^{low/-}$ cells and

92% of ALDH$^+$ cells. These results indicate BST2 expression correlates with established TIC

markers, suggesting BST2 may also allow enrichment for TIC.

To determine whether cells that highly express BST2 are enriched for mammosphere forming

potential and/or self-renewal activity we dissociated SUM159 xenografts and FACS-enriched

BST2^{high} (top 20%), BST2$^{low/-}$ (bottom 20%), and mock sorted cells to test their primary and

secondary mammosphere forming frequencies (Fig. 7B). BST2^{high} cells formed both primary and

secondary mammospheres at a greater frequency than either the BST2$^{low/-}$ cells or the mock

sorted control, with no statistically significant difference between the mock control and BST2$^{low/-}$.

To determine whether self-renewal capacity differed between the two populations, we compared

the rate of MSFE between the primary and secondary mammospheres and found the BST2^{high},

BST2$^{low/-}$, and mock sorted control form mammospheres at the same rate over passaging (Fig.

7C) indicating no differences in self-renewal capacity. Similar results were observed for BCM-

4272 (Fig. 7D and 7E). These results indicate that cells with high BST2 expression have enhanced

MSFE, but not increased self-renewal capacity.

Finally, to test whether BST2 can be used to enrich TIC directly, we compared the tumor-initiating

potential of BST2^{high} (top 20%), BST2$^{low/-}$ (bottom 20%), and mock-sorted cells from SUM159

xenografts using a limiting-dilution transplantation assay (Fig. 7F). Notably, the BST2^{high} cells

were the only population capable of forming tumors at all doses, with a TIC frequency of 1 in

1,721 cells. The BST2$^{low/-}$ cells had a 10-fold lower TIC frequency of 1 in 17,370 cells, whereas

the mock-sorted group had a modestly lower TIC frequency of 1 in 2,501 cells, likely due to the

presence of TIC with lower BST2 expression present in the 60% of cells not assayed.

Qualitatively similar results were obtained using BCM-4272 xenografts using a limiting-dilution

transplantation assay (Fig. 7G). The BST2^{high} cells were the only population capable of forming

tumors at all doses, with a TIC frequency of 1 in 1,102 cells, a much higher TIC frequency than

the mock sorted control, which had a TIC frequency of 1 in 8,859 cells. In BST2$^{low/-}$ cells, tumor-

initiating potential was depleted relative to the mock sorted control, with a TIC frequency of 1 in

18,796 cells. Taken together, our findings indicate BST2 can enrich for IFN/STAT1 signaling TIC.

BST2 regulates TIC frequency in SUM159

To explore whether BST2 regulates TIC function, we generated two BST2 knockdown SUM159

cell lines (sh-BST2 #1 and sh-BST2 #2) (Fig. 8A) and compared their primary and secondary

MSFE relative to a non-targeting control (Fig. 8B). BST2 knockdown reduced primary MSFE from

2.5% in the non-targeting control cells to 1.5% and 1.3% in shBST2 #1 and shBST2 #2,

respectively, whereas BST2 knockdown reduced secondary MSFE from 3% in the non-targeting

control cells to 1.6% and 1.2% in shBST2 #1 and shBST2 #2, respectively. These results

demonstrate BST2 regulates mammosphere-forming frequency in SUM159 cells.

To determine whether BST2 regulates self-renewal, we compared the rate of MSFE between

primary and secondary mammospheres in BST2 knockdown cells relative to a non-targeting

control (Fig. 8C). We found both BST2 knockdown cells formed mammospheres at the same rate

over passaging compared to the non-targeting control, indicating that BST2 does not regulate

self-renewal. Taken together, our data suggest BST2 functionally regulates TIC frequency

independent of self-renewal.

Because IFN regulates the expression of genes in our TIC signature, including BST2 (Fig. S6B–

C), we grew SUM159 mammospheres in the presence of IFNα or IFNγ and compared primary

and secondary mammosphere-forming frequency to an untreated control (Fig. 8D). Both IFNα

and IFNγ increased primary mammosphere frequency to 6.2% and 7.5%, respectively, compared

to the untreated control (3.4%). Similarly, IFNα and IFNγ increased secondary mammosphere

frequency to 5% and 9.3%, respectively, compared to the untreated control (2.9%). However, only

IFNγ increased rate of MSFE over passaging (from 7.5% to 9.3%), suggesting IFNγ regulates

both TIC frequency and self-renewal in SUM159 (Fig. 8E).

To determine whether BST2 is responsible for the IFN-mediated increase in MSFE, we compared

the effect of IFN on MSFE in two BST2 knockdown SUM159 cell lines (shBST2 #1 and shBST2

#2) compared to a non-targeting control (Fig. 8F). In the non-targeting control, IFNα increased

MSFE to 4.4% and IFNγ increased MSFE to 5.1% relative to the untreated control (2.7%). In both

BST2 knockdown lines, neither IFNα nor IFNγ increased MSFE relative to the untreated control,

suggesting BST2 is a mediator of the IFN-mediated increase in MSFE.

ALDH activity is a TIC marker in SUM159 (56). To evaluate whether BST2 knockdown reduces

ALDH activity, we performed an ALDEFLUOR assay in BST2 knockdown SUM159 cell lines

relative to a non-targeting control. BST2 knockdown reduced ALDH activity from 7.0% in the non-

targeting control cells to 1.6% and 0.8% in shBST2 #1 and shBST2 #2, respectively (Fig. 8G).

To determine whether this increase in TIC is due to proliferation, we compared cell cycle changes

following BST2 knockdown. Of the non-targeting control cells, 33.7% were proliferative, whereas

only 25.4% in shBST2 #1, and 13.1% in shBST2 #2, were proliferative, indicating BST2

knockdown reduces proliferation (Fig. 8H). Taken together, our data suggest that BST2 regulates

TIC frequency in SUM159, perhaps through modulating proliferation.

DISCUSSION

Herein, we extend our previous finding that STAT reporter activity enriches for TIC in two cell line xenograft models of claudin-low TNBC (11,57), and identified PDX BCM-4272 as an additional non-claudin-low model where STAT reporter activity is associated with TIC functions. Despite this finding, the reporter did not enrich cells with enhanced MSFE in the three other TNBC PDX models. Data are consistent with previous findings that STAT-mediated transcription has function(s) not necessarily associated with TIC. For example, STAT3 can promote immune evasion, metastasis, and chemoresistance (58), whereas STAT1 is associated with both tumor-promoting functions, such as immune suppression and metastasis, and tumor-suppressor functions, such as promoting apoptosis and immunosurveillance (59).

To address shortcomings of our original STAT-EGFP reporter, we developed the LT system to provide an inducible fluorescent label to tag STAT signaling TIC and other cells permanently as they undergo cell state changes. In addition, there are key questions in the cancer stem cell field that this system can address, including whether TIC from primary tumors function as metastasis-initiating cells (52,60), and whether enriched TIC populations following chemotherapy are the result of a therapeutic selective pressure, or are activated in response to the therapy (52,61).

We leveraged a synergy between scRNAseq, which has the power to detect rare cell populations, with our reporter systems to identify more precisely the TIC-enriched subpopulation of cells in multiple models of TNBC. The established TIC markers CD44$^+$/ CD24$^{low/-}$ and ALDH1A1 did not identify any unique cell states in our scRNAseq data, suggesting the expression of established TIC markers alone may not be sufficient to identify TIC in scRNAseq datasets. However, our analyses revealed that IFN response genes are differentially expressed in TIC in all three of our reporter-tagged TNBC, as well as three PDXs from an additional cohort, indicating the IFN/STAT1 signaling axis may be a common feature of TIC in a subset of TNBC. Critically, the IFN/STAT1 transcriptional state defines distinct cell states in all three breast cancer subtypes, as well as pre-

neoplastic tissue. The absence of the IFN/STAT1 state in normal breast suggests this

transcriptional state is a unique feature of some pre-neoplastic and neoplastic tissues, and may

represent a candidate cancer-specific therapeutic target.

From an evolutionary perspective, the activation of IFN response genes in tumor cells presents

multiple possibilities. On the one hand, TIC may have co-opted the IFN-mediated inflammatory

response of the immune system to regulate TIC processes related to their frequency, such as

differentiation, or cell state decisions (e.g. cellular quiescence/activation). Alternatively, and

arguably more evolutionarily likely given that multicellular organisms arose from a single self-

renewing cell, it is also possible that the immune system may have co-opted a stem cell pathway

to mediate inflammatory responses.

Using the common genes across TIC states in all three PDX models in scRNAseq data, we

derived a 25-gene TIC signature. Almost all genes we identified were not reported in previously

published TIC signatures, all of which were derived using bulk RNA sequencing (30-34,55).

Indeed, we found little agreement among the previously published TIC signatures themselves.

This suggests that it is essential to combine TIC enrichment with scRNAseq to derive high-

resolution TIC signatures.

There are limitations to consider when comparing bulk and single-cell samples. Bulk RNA

sequencing is frequently performed at much higher coverage and depth than typical scRNAseq,

which can allow detection of low expressing genes that may be missed by single-cell approaches.

On the other hand, scRNAseq can dissect transcriptome heterogeneity, allowing discovery of rare

cell types/states that would otherwise be obscured in bulk RNAseq.

With respect to translational potential, our 25-gene TIC signature was associated with worse

overall survival in untreated breast cancer patients, but not in patients treated with chemotherapy.

However, the signature was predictive of lack of response to chemotherapy, in line with previous

findings that TIC are intrinsically resistant to systemic therapies (15). These data are also

consistent with previous studies demonstrating enhanced expression of IFN genes following

cytotoxic chemotherapy (62). Indeed, a tumor cell state defined by IFN gene expression was

recently identified in TNBC scRNAseq, and was shown to be associated with tumors resistant to

standard-of-care chemotherapies (63). This raises the possibility that TIC exploit the IFN/STAT1

axis to promote their activity in cancer, including mediating chemoresistance (64).

We identified BST2 as a candidate cell surface TIC marker, as it was differentially expressed in

candidate TIC clusters across the three TNBC models we sequenced, and was a marker of

clusters with a high TIC score in each breast cancer subtype, as well as pre-neoplastic tissue.

BST2 is a transmembrane glycoprotein, whose expression is mediated by IFN, that is well-known

for tethering newly-synthesized enveloped virions to the host cell plasma membrane (65) and

activating NFκB to induce an antiviral state (66). To our knowledge, BST2 has never been

associated with TIC in any tumor type.

Beyond utility as a TIC marker, our results demonstrate that BST2 is a regulator of TIC frequency

in SUM159 and PDX BCM-4272, yet its knockdown in SUM159 did not affect self-renewal. This

indicates BST2 must regulate TIC frequency by a different mechanism, such as quorum sensing.

Quorum sensing, which is most commonly studied in bacteria, is a process of cell-to-cell

communication that allows cells to share information about cell density and adjust their gene

expression to regulate cell number (67). Organ-level quorum sensing has also been studied in

stem cells, such as hair stem cell populations (68). To the extent that we have studied it, tumors

appear to have specific TIC frequencies that are intrinsic properties of each tumor (69). If so, it

stands to reason quorum sensing is one mechanism that regulates TIC frequency. As a cell

surface protein, BST2 may be an integral component of the quorum sensing mechanism that

regulates TIC frequency.

ACKNOWLEDGMENTS

This work was supported, in part, by an F31 predoctoral fellowship (NCI CA278274) (to E.P.S), a U54 PDXNet PDX Development and Trials Center grant (NCI U54 CA224076) (to M.T.L.), an R01 grant (CA224867 to M.T.L. and H.L.F), a P30 Cancer Center Support Grant CA125123 (To C. Kent Osborne), a Core Facility Support Grant from the Cancer Research and Prevention Initiative of Texas RP170691 (to M.T.L.), the authors also acknowledge the joint participation by Diana Henry Helis Medical Research Foundation through its direct engagement in the continuous active conduct of medical research in conjunction with Baylor College of Medicine, the Komen for the Cure Promise grant, and a generous gift from the Korell family for the study of triple-negative breast cancer. S.S.Y. was supported by NIH grant R35GM133658 and Susan G. Komen Foundation grant CCR19609287. This project was also supported by the Cytometry and Cell Sorting Core at Baylor College of Medicine with funding from the CPRIT Core Facility Support Award (CPRIT-RP180672), the NIH (P30CA125123 and RR024574), the Single Cell Genomics Core at BCM with funding from NIH shared instrument grants (S10OD023469, S10OD025240) and P30EY002520, and the Genomic and RNA Profiling Core at Baylor College of Medicine with funding from NIH NCI (P30CA125123) and CPRIT (RP200504) grants. We thank Mu Wang for cloning the STAT lineage-tracing vectors as well as Dr. Koen Venken for helpful suggestions for vector design.

REFERENCES

1. Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, *et al.* Molecular portraits of human breast tumours. Nature **2000**;406:747-52

2. Sotiriou C, Wirapati P, Loi S, Harris A, Fox S, Smeds J, *et al.* Gene expression profiling in breast cancer: understanding the molecular basis of histologic grade to improve prognosis. J Natl Cancer Inst **2006**;98:262-72

3. Dent R, Trudeau M, Pritchard KI, Hanna WM, Kahn HK, Sawka CA, *et al.* Triple-negative breast cancer: clinical features and patterns of recurrence. Clin Cancer Res **2007**;13:4429-34

4. Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y, *et al.* Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. J Clin Invest **2011**;121:2750-67

5. Cancer Genome Atlas N. Comprehensive molecular portraits of human breast tumours. Nature **2012**;490:61-70

6. Reya T, Morrison SJ, Clarke MF, Weissman IL. Stem cells, cancer, and cancer stem cells. Nature **2001**;414:105-11

7. Ginestier C, Hur MH, Charafe-Jauffret E, Monville F, Dutcher J, Brown M, *et al.* ALDH1 is a marker of normal and malignant human mammary stem cells and a predictor of poor clinical outcome. Cell Stem Cell **2007**;1:555-67

8. Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF. Prospective identification of tumorigenic breast cancer cells. Proc Natl Acad Sci U S A **2003**;100:3983-8

9. Iliopoulos D, Hirsch HA, Wang G, Struhl K. Inducible formation of breast cancer stem cells and their dynamic equilibrium with non-stem cancer cells via IL6 secretion. Proc Natl Acad Sci U S A **2011**;108:1397-402

10. Marotta LL, Almendro V, Marusyk A, Shipitsin M, Schemme J, Walker SR, *et al.* The JAK2/STAT3 signaling pathway is required for growth of CD44(+)CD24(-) stem cell-like breast cancer cells in human tumors. J Clin Invest **2011**;121:2723-35

11. Wei W, Tweardy DJ, Zhang M, Zhang X, Landua J, Petrovic I, *et al.* STAT3 signaling is activated preferentially in tumor-initiating cells in claudin-low models of human breast cancer. Stem Cells **2014**;32:2571-82

12. Zhang M, Atkinson RL, Rosen JM. Selective targeting of radiation-resistant tumor-initiating cells. Proc Natl Acad Sci U S A **2010**;107:3522-7

13. Dierks C, Beigi R, Guo GR, Zirlik K, Stegert MR, Manley P, *et al.* Expansion of Bcr-Abl-positive leukemic stem cells is dependent on Hedgehog pathway activation. Cancer Cell **2008**;14:238-49

896   14.   Harrison H, Farnie G, Howell SJ, Rock RE, Stylianou S, Brennan KR, *et al.* Regulation of
897   breast cancer stem cell activity by signaling through the Notch4 receptor. Cancer Res
898   **2010**;70:709-18

899   15.   Li X, Lewis MT, Huang J, Gutierrez C, Osborne CK, Wu MF, *et al.* Intrinsic resistance of
900   tumorigenic breast cancer cells to chemotherapy. J Natl Cancer Inst **2008**;100:672-9

901   16.   Korkaya H, Kim GI, Davis A, Malik F, Henry NL, Ithimakin S, *et al.* Activation of an IL6
902   inflammatory loop mediates trastuzumab resistance in HER2+ breast cancer by
903   expanding the cancer stem cell population. Mol Cell **2012**;47:570-84

904   17.   Alamgeer M, Ganju V, Kumar B, Fox J, Hart S, White M, *et al.* Changes in aldehyde
905   dehydrogenase-1 expression during neoadjuvant chemotherapy predict outcome in locally
906   advanced breast cancer. Breast Cancer Res **2014**;16:R44

907   18.   Yu F, Yao H, Zhu P, Zhang X, Pan Q, Gong C, *et al.* let-7 regulates self renewal and
908   tumorigenicity of breast cancer cells. Cell **2007**;131:1109-23

909   19.   Liu H, Patel MR, Prescher JA, Patsialou A, Qian D, Lin J, *et al.* Cancer stem cells from
910   human breast tumors are involved in spontaneous metastases in orthotopic mouse
911   models. Proc Natl Acad Sci U S A **2010**;107:18115-20

912   20.   Matsuda T, Cepko CL. Controlled expression of transgenes introduced by in vivo
913   electroporation. Proc Natl Acad Sci U S A **2007**;104:1027-32

914   21.   Lou E, Fujisawa S, Morozov A, Barlas A, Romin Y, Dogan Y, *et al.* Tunneling nanotubes
915   provide a unique conduit for intercellular transfer of cellular contents in human malignant
916   pleural mesothelioma. PLoS One **2012**;7:e33093

917   22.   Koo BK, Stange DE, Sato T, Karthaus W, Farin HF, Huch M, *et al.* Controlled gene
918   expression in primary Lgr5 organoid cultures. Nat Methods **2011**;9:81-3

919   23.   Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A.
920   CellProfiler 4: improvements in speed, utility and usability. BMC Bioinformatics
921   **2021**;22:433

922   24.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, *et al.* Fiji: an
923   open-source platform for biological-image analysis. Nat Methods **2012**;9:676-82

924   25.   Hu Y, Smyth GK. ELDA: extreme limiting dilution analysis for comparing depleted and
925   enriched populations in stem cell and other assays. J Immunol Methods **2009**;347:70-8

926   26.   Kowalczyk MS, Tirosh I, Heckl D, Rao TN, Dixit A, Haas BJ, *et al.* Single-cell RNA-seq
927   reveals changes in cell cycle and differentiation programs upon aging of hematopoietic
928   stem cells. Genome Res **2015**;25:1860-72

929   27.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, *et al.* Integrated
930   analysis of multimodal single-cell data. Cell **2021**;184:3573-87 e29

931   28.   Andreatta M, Carmona SJ. UCell: Robust and scalable single-cell gene signature scoring.
932   Comput Struct Biotechnol J **2021**;19:3796-8

933   29.   Shen L. GeneOverlap: Test and visualize gene overlaps. R Package **2023**;1

934   30.   Charafe-Jauffret E, Ginestier C, Iovino F, Wicinski J, Cervera N, Finetti P*, et al.* Breast
935   cancer cell lines contain functional cancer stem cells with metastatic capacity and a
936   distinct molecular signature. Cancer Res **2009**;69:1302-13

937   31.   Creighton CJ, Li X, Landis M, Dixon JM, Neumeister VM, Sjolund A*, et al.* Residual breast
938   cancers after conventional therapy display mesenchymal as well as tumor-initiating
939   features. Proc Natl Acad Sci U S A **2009**;106:13820-5

940   32.   Leth-Larsen R, Terp MG, Christensen AG, Elias D, Kuhlwein T, Jensen ON*, et al.*
941   Functional heterogeneity within the CD44 high human breast cancer stem cell-like
942   compartment reveals a gene signature predictive of distant metastasis. Mol Med
943   **2012**;18:1109-21

944   33.   Pece S, Disalvatore D, Tosoni D, Vecchi M, Confalonieri S, Bertalot G*, et al.* Identification
945   and clinical validation of a multigene assay that interrogates the biology of cancer stem
946   cells and predicts metastasis in breast cancer: A retrospective consecutive study.
947   EBioMedicine **2019**;42:352-62

948   34.   Pei J, Wang Y, Li Y. Identification of key genes controlling breast cancer stem cell
949   characteristics via stemness indices analysis. J Transl Med **2020**;18:74

950   35.   Diggins KE, Greenplate AR, Leelatian N, Wogsland CE, Irish JM. Characterizing cell
951   subsets using marker enrichment modeling. Nat Methods **2017**;14:275-8

952   36.   Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z*, et al.* clusterProfiler 4.0: A universal enrichment
953   tool for interpreting omics data. Innovation (Camb) **2021**;2:100141

954   37.   Pal B, Chen Y, Vaillant F, Capaldo BD, Joyce R, Song X*, et al.* A single-cell RNA
955   expression atlas of normal, preneoplastic and tumorigenic states in the human breast.
956   EMBO J **2021**;40:e107333

957   38.   Wu SZ, Al-Eryani G, Roden DL, Junankar S, Harvey K, Andersson A*, et al.* A single-cell
958   and spatially resolved atlas of human breast cancers. Nat Genet **2021**;53:1334-47

959   39.   Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K*, et al.* Fast, sensitive and
960   accurate integration of single-cell data with Harmony. Nat Methods **2019**;16:1289-96

961   40.   Gyorffy B. Transcriptome-level discovery of survival-associated biomarkers and therapy
962   targets in non-small-cell lung cancer. Br J Pharmacol **2024**;181:362-74

963   41.   Fekete   JT,   Gyorffy   B.   ROCplot.org: Validating   predictive   biomarkers   of
964   chemotherapy/hormonal therapy/anti-HER2 therapy using transcriptomic data of 3,104
965   breast cancer patients. Int J Cancer **2019**;145:3140-51

966   42.   Ehret GB, Reichenbach P, Schindler U, Horvath CM, Fritz S, Nabholz M*, et al.* DNA
967   binding specificity of different STAT proteins. Comparison of in vitro specificity with natural
968   target sites. J Biol Chem **2001**;276:6675-88

969   43.   Wang Y, Wang Q, Tang CH, Chen HD, Hu GN, Shao JK, *et al.* p-STAT3 expression in
970   breast cancer correlates negatively with tumor size and HER2 status. Medicine (Baltimore)
971   **2021**;100:e25124

972   44.   Petrosyan V, Dobrolecki LE, Thistlethwaite L, Lewis AN, Sallas C, Srinivasan RR*, et al.*
973   Identifying biomarkers of differential chemotherapy response in TNBC patient-derived
974   xenografts with a CTD/WGCNA approach. iScience **2023**;26:105799

975   45.   Zhang X, Claerhout S, Prat A, Dobrolecki LE, Petrovic I, Lai Q*, et al.* A renewable tissue
976   resource of phenotypically stable, biologically and ethnically diverse, patient-derived
977   human breast cancer xenograft models. Cancer Res **2013**;73:4885-97

978   46.   Corish P, Tyler-Smith C. Attenuation of green fluorescent protein half-life in mammalian
979   cells. Protein Eng **1999**;12:1035-40

980   47.   Groner B. Determinants of the extent and duration of STAT3 signaling. JAKSTAT
981   **2012**;1:211-5

982   48.   Meyer MJ, Fleming JM, Ali MA, Pesesky MW, Ginsburg E, Vonderhaar BK. Dynamic
983   regulation of CD24 and the invasive, CD44posCD24neg phenotype in breast cancer cell
984   lines. Breast Cancer Res **2009**;11:R82

985   49.   Reid JM, Goetz MP, Buhrow SA, Walden C, Safgren SL, Kuffel MJ*, et al.*
986   Pharmacokinetics of endoxifen and tamoxifen in female mice: implications for comparative
987   in vivo activity studies. Cancer Chemother Pharmacol **2014**;74:1271-8

988   50.   Horvath CM, Wen Z, Darnell JE, Jr. A STAT protein domain that determines DNA
989   sequence recognition suggests a novel DNA-binding domain. Genes Dev **1995**;9:984-94

990   51.   Guo C, Yang W, Lobe CG. A Cre recombinase transgene with mosaic, widespread
991   tamoxifen-inducible action. Genesis **2002**;32:8-18

992   52.   Batlle E, Clevers H. Cancer stem cells revisited. Nat Med **2017**;23:1124-34

993   53.   Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R*, et al.* The STRING
994   database in 2023: protein-protein association networks and functional enrichment
995   analyses for any sequenced genome of interest. Nucleic Acids Res **2023**;51:D638-D46

996   54.   Rusinova I, Forster S, Yu S, Kannan A, Masse M, Cumming H*, et al.* Interferome v2.0: an
997   updated database of annotated interferon-regulated genes. Nucleic Acids Res
998   **2013**;41:D1040-6

55.   Colacino JA, Azizi E, Brooks MD, Harouaka R, Fouladdel S, McDermott SP*, et al.*
Heterogeneity of Human Breast Stem and Progenitor Cells as Revealed by Transcriptional
Profiling. Stem Cell Reports **2018**;10:1596-609

56.   Lin L, Hutzen B, Lee HF, Peng Z, Wang W, Zhao C*, et al.* Evaluation of STAT3 signaling
in ALDH+ and ALDH+/CD44+/CD24- subpopulations of breast cancer cells. PLoS One
2013;8:e82821

57.   Neve RM, Chin K, Fridlyand J, Yeh J, Baehner FL, Fevr T, *et al.* A collection of breast
1006   cancer cell lines for the study of functionally distinct cancer subtypes. Cancer Cell
1007   **2006**;10:515-27

1008   58.   Huynh J, Chand A, Gough D, Ernst M. Therapeutically exploiting STAT3 activity in cancer
1009   - using tissue repair as a road map. Nat Rev Cancer **2019**;19:82-96

1010   59.   Meissl K, Macho-Maschler S, Muller M, Strobl B. The good and the bad faces of STAT1
1011   in solid tumours. Cytokine **2017**;89:12-20

1012   60.   Massague J, Ganesh K. Metastasis-Initiating Cells and Ecosystems. Cancer Discov
1013   **2021**;11:971-94

1014   61.   Kim C, Gao R, Sei E, Brandt R, Hartman J, Hatschek T, *et al.* Chemoresistance Evolution
1015   in Triple-Negative Breast Cancer Delineated by Single-Cell Sequencing. Cell
1016   **2018**;173:879-93 e13

1017   62.   Sistigu A, Yamazaki T, Vacchelli E, Chaba K, Enot DP, Adam J, *et al.* Cancer cell-
1018   autonomous contribution of type I interferon signaling to the efficacy of chemotherapy. Nat
1019   Med **2014**;20:1301-9

1020   63.   Jacobo Jacobo M, Donnella HJ, Sobti S, Kaushik S, Goga A, Bandyopadhyay S. An
1021   inflamed tumor cell subpopulation promotes chemotherapy resistance in triple negative
1022   breast cancer. Sci Rep **2024**;14:3694

1023   64.   Zaidi MR, Merlino G. The two faces of interferon-gamma in cancer. Clin Cancer Res
1024   **2011**;17:6118-24

1025   65.   Van Damme N, Goff D, Katsura C, Jorgenson RL, Mitchell R, Johnson MC, *et al.* The
1026   interferon-induced protein BST-2 restricts HIV-1 release and is downregulated from the
1027   cell surface by the viral Vpu protein. Cell Host Microbe **2008**;3:245-52

1028   66.   Tokarev A, Suarez M, Kwan W, Fitzpatrick K, Singh R, Guatelli J. Stimulation of NF-
1029   kappaB activity by the HIV restriction factor BST2. J Virol **2013**;87:2046-57

1030   67.   Rutherford ST, Bassler BL. Bacterial quorum sensing: its role in virulence and possibilities
1031   for its control. Cold Spring Harb Perspect Med **2012**;2

1032   68.   Chen CC, Wang L, Plikus MV, Jiang TX, Murray PJ, Ramos R, *et al.* Organ-level quorum
1033   sensing directs regeneration in hair stem cell populations. Cell **2015**;161:277-90

1034   69.   Schatton T, Murphy GF, Frank NY, Yamaura K, Waaga-Gasser AM, Gasser M, *et al.*
1035   Identification of cells initiating human melanomas. Nature **2008**;451:345-9
1036
1037

FIGURES

1039

Figure 1: STAT reporter activity in breast PDX models. A) Representative IHC images

showing levels of pSTAT1 and pSTAT3 in a panel of PDX models. 4M67-EGFP reporter activity

was visualized by immunofluorescence (IF). Scale bar = 50 μm. IHC and IF were quantified using

5 representative images per PDX model. B) Flow cytometry plot illustrating the percentage of

4M67 reporter-positive cells in PDX models with demonstrated reporter activity. C)

Representative images comparing the mammosphere forming efficiency between the EGFP$^+$,

EGFP$^-$, and mock-sorted cells in four PDX models (BCM-4272: n = 10, BCM-15006: n = 9, BCM-

7821: n = 10, MC1: n = 8) with a bar graph of the mammosphere forming efficiency. Values are

mean $\pm$ SD. One-way ANOVA and Tukey's post hoc test were used to determine statistical

significance. *$p < 0.05$, ***$p < 0.001$, ****$p < 0.0001$. MSFE: mammosphere forming efficiency.

1050

Figure 2: STAT lineage-tracing reporter provides a long-term fluorescent label to STAT signaling cells. A) Schematic illustrating how the STAT lineage-tracing system functions upon the addition of TAM. Cre is not transcribed in non-signaling (EGFP⁻) cells, thus no color-switching occurs in the presence of TAM. The Cre is transcribed in signaling (EGFP⁺) cells; therefore, the addition of TAM drives color-switching from dsRed to mNeptune2. B) Graphic illustrating the four different LT cell populations that can be identified by microscopy and flow cytometry. C) Confocal images of SUM159 STAT LT xenografts and D) flow cytometry quantification of the percent of

1059   each LT population two days after 4 mg TAM treatment (Untreated: $n = 4$, TAM-treated: $n = 5$).

1060   E) Confocal images of MDA-MB-231 STAT LT xenografts and F) flow cytometry quantification of

1061   the percent of each LT population two days after 4 mg TAM treatment ($n = 3$). One-way ANOVA

1062   and Tukey's post hoc test were used to determine statistical significance. *$p < 0.05$, **$p < 0.01$,

1063   ns = not significant; values are mean $\pm$ SD. Nep: mNeptune2.

1064

1065

Figure 3: Long-lived STAT signaling cells are enriched for TIC in SUM159 xenografts. A)

SUM159 STAT LT xenografts were dissociated, then each LT population was FACS enriched and

grown as mammospheres. B) Mammosphere forming efficiency of each LT cell population from

SUM159 xenografts (n = 6). One-way ANOVA and Tukey's post hoc test were used to determine

statistical significance. C) Confocal images of a 16-day TAM time course in SUM159 xenografts

(n = 6). D) Flow cytometry quantification of the percentage of each LT cell population in SUM159

xenografts at five different time points. E) The time course was extended using a serial

transplantation assay. Tumor fragments harvested two days post TAM (TG1), were transplanted

into new hosts two additional times (n = 6). F) Flow cytometry quantification of the percentage of

each LT cell population in SUM159 xenografts at each transplant generation (TG). ***$p < 0.001$,

****$p < 0.0001$, ####$p < 0.0001$; values are mean $\pm$ SD. Nep: mNeptune2.

1077

Figure 4: Interferon response genes are differentially expressed in TIC. SUM159 STAT3 LT,

BCM-4272 4M67-EGFP, and BCM-15006 4M67-EGFP xenografts were harvested and

dissociated, then each fluorescent population was enriched and sequenced using 10X genomics.

A) Each xenograft was clustered and visualized using a UMAP plot. B) UMAP with cell identity

overlayed on the plot. C) Hallmark gene set enrichment analysis of candidate TIC transcriptional

state from each xenograft. D) Intersection of the upregulated genes in candidate TIC clusters with

accompanying pairwise comparisons using Fisher's exact test. Intersection of candidate TIC

clusters from each model reveals 25 shared genes. E) STRING database illustration of protein-

protein interactions between the 25-gene TIC signature and overrepresentation analysis results

using the 25-gene list. F) Marker enrichment modeling heatmap illustrating cluster-to-cluster

similarity based on expression of the genes in the 25-gene TIC signature. Nep: mNeptune2, FDR:

false discovery rate, p.adjust: Benjamini-Hochberg (BH)-adjusted p-values.

1091

1092

Figure 5: Interferon response genes are differentially expressed in a subset of epithelial cells in human breast cancer patient samples. A) UMAP visualization of eight human TNBC patient samples with feature plot and violin plot showing TIC score, derived from based on the expression of our 25-gene TIC signature. B) UMAP visualization of six human HER2+ patient samples with TIC score feature and violin plots. C) UMAP visualization of 19 human ER+ patient samples with TIC score feature and violin plots. D) UMAP visualization of four human pre-neoplastic (*BRCA1*+/- tissue) patient samples with TIC score feature and violin plots. E) UMAP visualization of 11 human normal breast samples with TIC score feature and violin plots. In all TIC score violin plots, P values were calculated using a two-sided Wilcoxon test. F) The intersection

between the common differentially expressed genes from clusters that were defined by a high

TIC score across all three breast cancer subtypes and our 25-gene TIC signature. G) The

intersection between the common differentially expressed genes from clusters that were defined

by a high TIC score across all three breast cancer subtypes and pre-neoplastic tissue and our 25-

gene TIC signature. The significance of gene list comparisons was calculated using Fisher's exact

test. IFN: interferon

1108

Figure 6: The IFN/STAT1 signaling axis is associated with worse overall survival and lack

of response to chemotherapy in breast cancer patients. A) Kaplan-Meier plots showing

overall survival between breast cancer patients with high or low expression of the 25-gene TIC

signature including all patients (untreated + chemotherapy), only including untreated patients, and

1114   only including patients treated with chemotherapy. B) Kaplan-Meier plots showing overall survival

1115   between breast cancer patients with high or low expression of the 25-gene TIC signature

1116   comparing untreated patients and patients treated with chemotherapy by PAM50 subtype. C)

1117   ROC curve and box plots showing the association between the 25-gene TIC signature and

1118   chemotherapy response by breast cancer subtype.

1119

1120

Figure 7: BST2 allows TIC enrichment in breast cancer xenograft models. A) Flow cytometry

plot examining the correlation between BST2 and TIC markers CD44$^+$/CD24$^-$ and ALDEFLUOR

activity in SUM159 and BCM-4272 xenografts. A two-tailed T-test was used to determine

significance. B) SUM159 xenografts were dissociated, then the BST2 high, BST2 low, and mock-

sorted populations were FACS enriched and grown as primary mammospheres (1°), then

passaged and grown as secondary mammospheres (2°) (n = 3 biological replicates). One-way

ANOVA and Tukey's post hoc test were used to determine statistical significance. C) Differences

in MSFE between primary and secondary mammospheres for each flow-sorted fraction. A two-

tailed T-test was used to determine significance. D) BCM-4272 xenografts were dissociated, then

the BST2 high, BST2 low, and mock-sorted populations were FACS enriched and grown as

primary mammospheres, then passaged and grown as secondary mammospheres. MSFE of

each cell population for each xenograft (n = 3 biological replicates). One-way ANOVA and Tukey's

post hoc test were used to determine statistical significance. E) Differences in MSFE between

primary and secondary mammospheres for each flow-sorted fraction A two-tailed T-test was used

to determine significance. F) Limiting dilution transplantation results of BST2 high, BST2 low, and

mock-sorted populations from SUM159 xenografts. TIC frequency was calculated using extreme

limiting dilution analysis and a Fisher's exact test was used to determine statistical significance

(25). G) Limiting dilution transplantation results of BST2 high, BST2 low, and mock-sorted

populations from BCM-4272 xenografts. TIC frequency was calculated using extreme limiting

dilution analysis and a Fisher's exact test was used to determine statistical significance (25). *p <

0.05, **p < 0.01, ***p <0.001, ****p < 0.0001; values are mean $\pm$ SD.

1142

Figure 8: BST2 regulates TIC frequency in SUM159. A) BST2 was knocked down in SUM159

cells using two different shRNAs (shBST2 #1 and shBST2 #2), then BST2 knockdown was

confirmed using Jess automated western blot. A two-tailed T-test was used to evaluate

significance between each knockdown line and a non-targeting control (NT). B) SUM159 NT,

shBST2 #1, and shBST2 #2 cells were grown as primary mammospheres (1°), then passaged

and grown as secondary mammospheres (2°) (n = 3 biological replicates). One-way ANOVA and

Tukey's post hoc test were used to determine statistical significance. C) Differences in MSFE

between primary and secondary mammospheres for each cell line. A two-tailed T-test was used

to determine significance. D) SUM159 cells were grown as primary mammospheres (1°) in the

presence of IFNα (100 ng/μL), IFNγ (100 ng/μL), or left untreated as a control, then passaged and

grown as secondary mammospheres (2°) in the presence of IFNα (100 ng/μL), IFNγ (100 ng/μL),

or left untreated as a control (n = 3 biological replicates). One-way ANOVA and Tukey's post hoc

test were used to determine statistical significance. E) Differences in MSFE between primary and

secondary mammospheres for treatment. A two-tailed T-test was used to determine significance.

F) SUM159 NT, shBST2 #1, and shBST2 #2 cells were grown as primary mammospheres (1°) in

the presence of IFNα (100 ng/μL), IFNγ (100 ng/μL), or left untreated (UT) as a control (n = 3

biological replicates). One-way ANOVA and Tukey's post hoc test were used to determine

statistical significance. G) Representative flow cytometry plots illustrating ALDH activity in

SUM159 NT, shBST2 #1, and shBST2 #2 cells and quantification of three biological replicates.

H) Representative cell cycle flow cytometry plots using propidium iodide in SUM159 NT, shBST2

#1, and shBST2 #2 cells and quantification of three biological replicates. *$p < 0.05$, **$p < 0.01$,

***$p <0.001$, ****$p < 0.0001$; values are mean $\pm$ SD. 1°: primary mammospheres, 2°: secondary

mammospheres, UT: untreated, IFN: interferon.

1168

1169

1170