Tap Dancing Frogs*: Posterior Toe Tapping and Feeding Behavior in *Dendrobates*

tinctorius

Running title: Toe tapping in poison frogs

Thomas Q Parrish[1] & Eva K Fischer[1,2] [*]

1 Department of Evolution, Ecology, and Behavior, School of Integrative Biology, University of

Illinois Urbana-Champaign, Urbana IL 61801, USA

2 Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign, Urbana

IL 61801, USA

*Corresponding author:

Eva K Fischer

Department of Evolution, Ecology, and Behavior

School of Integrative Biology

University of Illinois Urbana-Champaign

505 S. Goodwin Ave.

Urbana, IL 61801, USA

Email: efischer@illinois.edu

Significance Statement

The toe tapping of some amphibians is an intriguing behavior that has attracted attention

from researchers and hobbyists, yet the functional role of toe tapping remains poorly understood.

Previous studies have noted an association between toe tapping and feeding in other species,

and our work does so quantitatively and experimentally in Dyeing poison frogs, *Dendrobates*

tinctorius. We demonstrate that frogs modulate tap rate based on the presence of conspecifics,

prey accessibility, and substrate type. We suggest tapping may act as a vibrational stimulus to

facilitate prey detection and capture alongside other functions.

Abstract

Animals have myriad adaptations to help them hunt and feed in the most efficient and

effective manner. One mysterious behavior related to hunting and feeding is the posterior toe

tapping behavior of some frogs. Biologists and hobbyists alike have long noticed this behavior,

but there is little empirical data to explain its causes and consequences. To test the hypothesis

that tapping is related to feeding and modulated by environmental context, we conducted a series

of related experiments in the Dyeing poison frog, *Dendrobates tinctorius*. We reasoned that

mechanostimulation of prey by vibrational stimuli from tapping might induce prey movement and

thereby facilitate prey detection by frogs. We first confirmed that tap rate was higher during

feeding as has been observed in other species. Interestingly, this effect was heightened in the

presence of a conspecific. We next asked whether frogs tapped less under conditions when

vibrational contact between frogs and prey was not possible. Finally, we asked whether *D.*

tinctorius tap more on more vibrationally responsive surfaces and whether prey capture success

was higher when tapping. In addition to confirming an association between tapping and feeding,

our work demonstrates modulation of toe tapping based on social context, prey accessibility, and

substrate characteristics.

Key words: anuran, Dendrobatidae, pedal luring, prey capture, feeding behavior

Introduction

The toe tapping of various species of amphibians is an intriguing behavior that has attracted attention from researchers and pet owners. Variably referred to as toe tapping, twitching, wiggling, and trembling, this curious behavior is the subject of YouTube videos, TikToks, and Twitter threads. Indeed, a recent publication took advantage of online video sharing platforms to compile a toe tapping dataset of 222 videos of 389 individuals representing 63 frog species (Claessens, Ganchev, Kukk, Schutte, & Sloggett, 2020). Yet despite being readily observed and widely documented, the functional role of toe tapping remains poorly understood.

Among amphibians, toe tapping falls under the general umbrella of pedal signaling and luring, which ranges from ostentatious movements of the entire limb to subtle tapping of fingers or toes (reviewed in (Erdmann, 2017; Hödl & Amézquita, 2001)). Under this umbrella, a related set of similar-looking toe movements have been broadly discussed in the context of visual and vibrational communication. Toe tapping has been suggested to function in prey capture (Claessens et al., 2020; Hagman & Shine, 2008; McFadden, Harlow, Kozlowski, & Purcell, 2010; Sloggett & Zeilstra, 2008), mate attraction and courtship (Landestoy & Ortíz, 2015; Starnberger, Maier, Hödl, & Preininger, 2018), aggression (Bee, Reichert, & Tumulty, 2016; Furtado, Márquez, & Hartz, 2017), and has also been suggested to be a 'displacement behavior' with no functional relevance (Furtado et al., 2017; Furtado & Nomura, 2014).

We regularly observe Dyeing poison frogs (*Dendrobates tinctorius*) in our lab tapping their posterior toes (Movie S1 & S2). Toe tapping has been documented in various species of poison frog (family Dendrobatidae), with some bias toward observations in this group likely resulting from the fact that these brightly colored, diurnal frogs are popular as pets and in zoos and aquaria (Claessens et al., 2020). Two recent studies demonstrated increased toe tapping associated with feeding in poison frog species closely related to *D. tinctorius*. Schulte and König (Schulte & König, 2023) reported more tapping in response to both small (fruit flies) and large (crickets) prey items, but not conspecific calls, in *Dendrobates auratus*; and Vergara-Herrera

and colleagues (Vergara-Herrera, Cocroft, & Rueda-Solano, 2023) report detailed tap

characterization in *Dendrobates truncatus*, including accelerated tapping right before prey

capture.

Poison frogs have dietary specializations for small prey items (Santos, Coloma, &

Cannatella, 2003), and in the wild *D. tinctorius* feed on a variety of arthropod prey (Rojas &

Pašukonis, 2019). Their prey is small and quick, and prey movement is key for detection and

capture. The essential role of prey movement for prey capture in amphibians was demonstrated

by studies showing that toads readily strike at any object of the appropriate size and moving in

the correct direction (i.e., 'worm like' lines moving horizontally) (Ewert, 1987). These studies

garnered frogs and toads the reputation of being stereotyped in their hunting strategies, with

sensory constraints limiting their ability to plastically adjust behavior. However, recent years have

seen a growing number of studies cataloging the ability of these animals to integrate complex,

multi-modal sensory information (e.g. (Amézquita & Hödl, 2004; de Lourenço, Haddad, & de Sá,

2020; Deban, O'Reilly, & Nishikawaa, 2001; Hartmann, Giasson, Hartmann, & Haddad, 2005;

James et al., 2022; Narins, Grabul, Soma, Gaucher, & Hödl, 2005; Toledo, Araújo, Guimarães,

Lingnau, & Haddad, 2007)).

Though primarily explored in the context of predator evasion and conspecific

communication (e.g. (James et al., 2022; Narins, 1990; Warkentin, 2005)), the exquisite

vibrational sensitivity of anurans may also play a role in prey capture. Using a mechanical

model, Hagman and Shine (Hagman & Shine, 2008) demonstrated that cane toads vibrate their

toes close to the frequency most effective at attracting conspecific prey. Vibrational

communication and seismic stimuli through surfaces have also been shown to be important for

amphibian's arthropod prey. For example, flies can feel vibrations through their legs and react to

them (Jarman, 2002). Taken together, this led us to predict that toe tapping could be a

vibrational stimulus associated with prey capture and feeding.

Given the importance of prey movement for prey capture, we reasoned that toe tapping

might result in vibrational stimuli that startle prey and increase hunting efficiency. Under these

assumptions, we predicted that *D. tinctorius* would adjust their tapping behavior based on

environmental context. We performed a series of experiments to test these ideas. First, we

confirmed that *D. tinctorius* do indeed tap more while feeding. Second, we asked whether frogs

tapped less under conditions with visual, but no vibrational, contact between frogs and prey. Third,

we tested whether *D. tinctorius* tap more on vibrationally responsive surfaces and asked whether

prey capture success was higher when tapping. Alongside recent studies in related species, our

findings contribute to our understanding of this visually conspicuous yet functionally mysterious

behavior.

Methods

(a) Animals

D. tinctorius were housed in our animal facility at the University of Illinois Urbana-

Champaign. Frogs were housed breeding pairs in 45x45x45 cm terraria containing soil, moss,

leaf litter, live plants, coconut shelters, and a small pond. We recorded only a single frog per

terraria (i.e., per pair) in any given trial. Terraria were maintained at 70-100% humidity and 22-

24°C on a 12L:12D light cycle. We fed frogs *Drosophila hydei* or *Drosophila melanogaster* dusted

with vitamins three times weekly. All animal care and experimental procedures were approved by

the University of Illinois Animal Care and Use Committee (IACUC protocol #20147). We used the

STRANGE framework in our experimental design and note that – while toe tapping is observed

in the wild – these are captive bred and reared animals where there is potential for lab selection.

(b) Experiment 1: Tapping in feeding versus non-feeding states

In the first experiment, we tested if *D. tinctorius* tap more while feeding versus not feeding.

For feeding trials, we dropped ½ teaspoon of *Drosophila* into the frogs' terrarium. Once frogs

began hunting, we took one minute of high-speed video (240 frames per second). For non-feeding

trials, we took one minute of high-speed video while frogs were not feeding. All trials took place

between 11:00 and 14:00. We also recorded partner proximity (frogs are housed in breeding

pairs) during filming. Videos were analyzed as described below. N=12 feeding and N=10 non-

feeding trials with N=8 total individuals (some frogs were tested twice).

(c) Experiment 2: Tapping when prey is on a different surface

In the second experiment we tested whether tapping behavior was affected by prey being

on a different surface and inaccessible to the frogs. We placed *Drosophila* inside a small, clear

petri dish so that the frogs could see the prey but not feed on it, and the prey was not on the same

surface as the frog. Petri dishes were placed in frogs home terraria during normal feeding time,

and we recorded high-speed video for one minute once frogs began striking at prey. Paired trials

with freely moving flies were conducted as described above the next day. We performed N=16

inaccessible prey and N=14 free-moving prey trials with N=16 individuals (two frogs were

excluded on the second day because they were transporting tadpoles).

(d) Experiment 3: Tapping variation between surfaces

In the third experiment, we asked whether tap rate depended on substrate type. We set

up experimental terraria with four substrate types: soil (a natural, unpliable surface), leaf litter (a

natural, pliable surface), gel (1% agar; an unnatural, pliable surface), and glass (an unnatural,

unpliable surface). We chose these surfaces to obtain a variety of pliability and familiarity. We

placed frogs singly into test terraria and gave them one-hour acclimation to the novel environment.

Following acclimation, we fed frogs and collected data as in Experiment 1. In addition, we

recorded the number of strikes frogs made in attempts to catch prey, and how many of these

strikes were successful. We performed all trials one day after last feeding to control for satiety

status. Each individual performed one trial on every surface, apart from two individuals performing

two on each surface. N=13 trials per surface with N=11 individuals. All trials were analyzed for

tap rate. For strike and success rate, three trials were excluded from analysis due to video loss.

(e) Data Analysis

We quantified tap number from slow motion video and standardized data for each frog. To

calculate the number of taps per minute, we divided the number of taps on each foot by the

number of seconds the foot was visible to obtain taps per second. We then multiplied that by 60

to calculate taps per minute for each foot and added the two feet together to get total taps per

minute. For experiment three, we calculated success rate by dividing the number of successful

strikes by the total number of strikes. To test for potential biases introduced by toe visibility across

contexts, we compared visibility rates across surfaces and found no differences (main effect of

surface: $F_{3,48}$ = 0.45, p = 0.7168).

We used R v4.4.2 (Team, 2022) in Rstudio v 3.1.446 (Posit Team, 2023) for all statistical

analyses. As in previous studies (Vergara-Herrera et al., 2023), we did not find an effect of sex

on tap rate and therefore omitted this variable from further analyses. In the first experiment, we

tested for differences in tap rate using a linear mixed model that included feeding status and

partner presence as fixed effects and frog ID as a random effect. In the second experiment, we

used the same approach (excepting the random effect of individual) to compare frogs feeding on

free flies (data from Experiment 1) to frogs feeding on flies confined to a petri dish. In the third

experiment, we included substrate type as a fixed effect and frog ID as a random effect. For all

linear models, we evaluated significance via ANOVA followed by Tukey corrected post hoc testing

with the package emmeans in R. We tested for correlations between tap rate, strike rate, and

strike success using Spearman correlations. We additionally tested for a substrate-dependent

relationship between prey capture success and tapping, where capture success rate was

predicted by surface type, tap rate, and their interaction.

Results

(a) Frogs tap more when feeding, especially in the presence of a conspecific

Frogs tapped significantly more when feeding versus non-feeding (main effect of feeding:

$F_{1,17}$ = 116.65, p <0.0001) and this effect was more pronounced in the proximity of their partner

frog (feeding*partner interaction: $F_{1,18}$ = 7.36, p = 0.0143) (Fig. 1a). While partner proximity

increased tap rate in the presence of prey (feeding alone vs feeding with partner: $t_{17.8}$ = -4.121, p

= 0.0033), tap rate was not increased by partner proximity in the absence of prey (not feeding

alone vs not feeding with partner: $t_{17.8}$ = -0.277, p = 0.9923). The average tap rate was 389

taps/min (range: 305 - 468) when feeding alone, 684 taps/min (range: 492 - 997) when feeding

near a partner, 50 taps/min (range: 4 - 149) when not feeding away from a partner, and 43

taps/min (range: 5 - 92) when not feeding but near a partner.

(b) Frogs tap more when prey is accessible

Tap rate was significantly lower when prey was inaccessible and on a different surface

from the frogs ($F_{1,14}$ = 19.85, p = 0.0005) (Fig. 1b). The average tap rate when frogs could see,

but not access, flies was 50 taps/min (range: 0 - 169); significantly lower than the 166 taps/min

(range: 11 - 409) while feeding on free moving flies (Fig. 1b). Importantly, we observed this lower

tap rate even though frogs continuously attempted to catch inaccessible prey (Movie S3).

Figure 1: Increased tapping when feeding on accessible prey. **(a)** Frogs tapped significantly more when feeding versus not feeding, and this effect was more pronounced in the presence of a partner frog (dark orange) than when alone (light orange). **(b)** Tap rate was significantly lower when prey was on a different surface and inaccessible to frogs (flies inside a petri dish; Movie S3).

(c) Tap rate depends on surface type

There was a significant difference in tap rate between surface types ($F_{1,40.6} = 6.21$, p =

0.0014) (Fig. 2a). Average tap rates were 98 taps/min for soil (range: 0 - 417), 255 taps/min for

leaf (range: 24 - 528), 118 taps/min for gel (range: 0 - 440), and 64 taps/min for glass (range: 0 -

211). Post hoc comparisons revealed that frogs tapped more on leaves than all other surfaces

(gel: $t_{38.4} = 2.87$, p = 0.0317; glass: $t_{38.4} = 4.01$, p = 0.0015; soil: $t_{38.4} = 3.30$, p = 0.0107) (Fig. 2a).

Tap rates in experimental tanks (Experiment 3) were on average lower than when frogs were in

their home terraria (Experiment 1), but still higher than when not feeding in home terraria.

While tap rate differed between surface types, there was no difference in the total number

of strikes ($F_{3,28} = 1.41$, p=0.259) or success ($F_{3,28} = 0.81$, p = 0.497) based on substrate. Strike

rate increased with tap rate (r=0.318, p=0.045); however, tap rate did not significantly predict prey

capture success overall (tap rate main effect: $F_{1,30}$ = 2.29, p = 0.142) or based on surface type

(tap rate*surface interaction: $F_{3,28}$ = 1.43, p = 0.254) (Fig. 2b).

Figure 2. Surface type affects tap rate but not prey capture success. (a) Tap rates differed based on surface. Tap rate was higher on leaves (orange) as compared to soil (yellow), gel (blue), or glass (grey). (b) Prey capture success rate did not differ based on surface type and was not predicted by tap rate.

Discussion

Taken together, our findings demonstrate that tapping is associated with feeding and tap

rate is modulated by partner proximity, prey accessibility, and surface characteristics in *D.*

tinctorius. In line with our predictions, we found that the frogs tapped more while feeding, when

on the same surface as prey, and on a vibrationally responsive surface. Unexpectedly, we also

observed an increase in feeding-related tapping in the presence of a breeding partner. Counter

to our expectations, we did not find a relationship between tap rate and prey capture success. We

discuss interpretations and implications of our findings below.

An association between feeding and posterior toe tapping has long been noted by
researchers and hobbyists, and recent empirical work substantiated this relationship in two closely
related species of poison frog (Schulte & König, 2023; Vergara-Herrera et al., 2023). Experiment
1 confirmed that tapping is associated with feeding in *D. tinctorius*, with frogs increasing their tap
rate up to 10-fold in the presence of prey. Interestingly, tap rate was further increased when
feeding in proximity of a breeding partner, while partner proximity alone did not increase tap rate.
At present, we do not know the reason for this social modulation of feeding-related toe tapping
behavior (see further discussion below). We highlight that – at a maximum of almost 500 taps per
toe per minute – this behavior is incredibly fast for any vertebrate muscular movement (Rome,
2006). In brief, these findings demonstrate an association between tapping and feeding and
provide interesting avenues for further study.

Given the vibrational sensitivity of both frogs and their arthropod prey, and the importance
of prey movement for prey detection, we hypothesized that toe tapping could function as a
vibrational stimulus that facilitates prey capture. To test this idea, we examined the relationship
between tapping and surface type. When frogs could see but not capture flies, frogs tapped
significantly less but *still hunted.* We suggest that this change in tap rate may be related to
changes in vibrational stimuli and/or feedback from prey capture success. This observation
suggests that frogs might alter their tapping behavior based on prey's responses.

To test the potential connection to vibrational stimuli further, we next asked whether *D.*
tinctorius modulated tapping based on surface type. We found that frogs tapped at a significantly
higher rate on the leaf surface. This aligned with our prediction that tapping would be enhanced
on vibrationally responsive surfaces. We were concerned that tapping behavior might be impacted
by the naturalness and familiarity of each surface, and the novel environment of the test terraria.
Tap rate was overall lower in the experimental terraria than in home terraria; however, frogs
hunted on all test surfaces, with no differences in prey capture attempts or prey capture success.
Additionally, we found that frogs tapped less on soil than leaves, despite both surfaces being

familiar to frogs. Taken together, these observations demonstrate that frogs modulate tapping

behavior based on substrate type and independent of prey capture attempts.

We expected higher tap rates to be associated with increased prey capture success;

however, this prediction was not supported, in general or based on surface type. Similarly, we

found no difference in prey capture success across surface types. We suspect this lack of a

relationship may be an artifact of our feeding method in the lab, where frogs are provided a large

number of flies and prey capture success is artificially high. While this possibility requires further

study, our demonstration of an association between tapping behavior, capture attempts, and

surface type is suggestive and opens the door for further experiments.

Additional work is also needed to clarify additional, non-mutually exclusive roles and

reasons for toe tapping. One alternative explanation for tapping is a generalized displacement

and/or excitatory response (Furtado et al., 2017; Furtado & Nomura, 2014). Under this

assumption, tapping has no functional role but is a reflex or arousal response to food or

movement, akin to a dog wagging its tail. This alternative could contribute to a lower tap rate when

prey was inaccessible, assuming arousal is lower when capture is unsuccessful. However, such

an effect cannot explain all our findings here, given that frogs also modulated tap rate based on

surface type and independent of prey presence and movement or feeding success.

Another non-mutually exclusive alternative is that tapping may be involved in social

communication between frogs, in agonistic and/or affiliative contexts (Furtado et al., 2017;

Landestoy & Ortíz, 2015; Starnberger et al., 2018). In poison frogs specifically, toe tapping has

been reported during feeding (Schulte & König, 2023) and observed during courtship (Barquero

& Arguedas, 2022) in *D. auratus*; however, presentation of advertisement calls did not increase

toe tapping in this species (Schulte & König, 2023). Tapping has similarly been noted during the

later stages of courtship in *D. tinctorius* (Rojas & Pašukonis, 2019) where it may also play a role

in vibrational signaling. We observed an increase in toe tapping while feeding in the presence of

another frog, but this social modulation occurred only when prey was present. These observations

suggest that any social function – be it cooperative or antagonistic – is in addition to, not instead

of, a feeding-related role. We emphasize that our study was not designed to explicitly test a role

for toe tapping in social communication and we did not monitor tapping during courtship or

aggressive interactions. We note that conspecifics in our study were the breeding partners of focal

individuals, and that familiarity influences aggressive and affiliative behaviors. In sum, while other

non-mutually exclusive roles for posterior toe tapping exist and are of interest for future study, our

data here support a connection between toe tapping and feeding that is socially and

environmentally modulated in *D. tinctorius*.

Conclusions

Taken together, our findings confirm an association between toe tapping and feeding and

demonstrate that *D. tinctorius* modulate tapping behavior based on social and environmental

context. The observation that frogs alter their tapping frequency – but not their strike rate – based

on substrate type, suggests that tapping has a functional, context dependent role. In line with

previous studies, we suggest that tapping may stir small arthropods via substrate vibrations and

thereby help *D. tinctorius* detect prey. By quantifying tap number, we also confirm the impressive

speed with which frogs perform the behavior. Future studies characterizing tapping biomechanics,

measuring substrate vibrations caused by tapping, and testing the vibrational sensitivity of both

frogs and prey will further our understanding of tap-dancing toes.

Acknowledgements

We would like to thank Samta Oza for help with data collection, and members of the Fischer Lab

for their support and feedback on previous version of the manuscript. Thanks to Phil Anderson

for advice on project design and assistance with high-speed filming.

Data Availability

Data analyzed in the current study are provided in the Supplemental Materials.

Funding

TQP was supported by a Dr. Kirk and Mrs. Shannon Moberg Scholarship and a Robert H. Davis

Undergraduate Research Prize. EKF was supported by start-up funding from the University of

Illinois Urbana-Champaign and the National Science Foundation (IOS 21-46058).

Conflict of interest statement

The authors declare no conflicts of interest.

References

Amézquita, A., & Hödl, W. (2004). How, when, and where to perform visual displays: The case of the Amazonian frog *Hyla parviceps*. *Herpetologica*, *60*(4), 420–429. https://doi.org/10.1655/02-51

Barquero, M. D., & Arguedas, V. (2022). Mass movement and potential vibratory toe signalling in the Green and Black Poison-Dart Frog, *Dendrobates auratus* (Amphibia: Dendrobatidae). *Herpetology Notes*, *15*(February), 79–82.

Bee, M. A., Reichert, M. S., & Tumulty, J. (2016). *Assessment and Recognition of Rivals in Anuran Contests*. *Advances in the Study of Behavior* (Vol. 48). Elsevier Ltd. https://doi.org/10.1016/bs.asb.2016.01.001

Claessens, L. S. A., Ganchev, N. O., Kukk, M. M., Schutte, C. J., & Sloggett, J. J. (2020). An investigation of toe-tapping behaviour in anurans by analysis of online video resources. *Journal of Zoology*, *312*(3), 158–162. https://doi.org/10.1111/jzo.12815

de Lourenço, A. R., Haddad, C. F. B., & de Sá, F. P. (2020). Multimodal signaling in *Boana albopunctata* (Anura: Hylidae): reading visual and acoustic cues. *Phyllomedusa*, *19*(2), 201–216. https://doi.org/10.11606/ISSN.2316-9079.V19I2P201-216

Deban, S. M., O'Reilly, J. C., & Nishikawaa, K. C. (2001). The evolution of the motor control of feeding in amphibians. *American Zoologist*, *41*(6), 1280–1298. https://doi.org/10.1093/icb/41.6.1280

Erdmann, J. A. (2017). *The function of toe movement in feeding by the gulf coast toad (Incilius nebulifer)*. Southeastern Louisiana University. https://doi.org/10.13140/RG.2.2.15286.80961

Ewert, J. P. (1987). Neuroethology of releasing mechanisms: Prey-catching in toads. *Behavioral and Brain Sciences*, *10*(3), 337–368. https://doi.org/10.1017/S0140525X00023128

Furtado, R., Márquez, R., & Hartz, S. M. (2017). In front of a mirror: visual displays may not be aggressive signals in nocturnal tree frogs. *Journal of Natural History*, *51*(7–8), 443–454.

335 https://doi.org/10.1080/00222933.2016.1262078

336 Furtado, R., & Nomura, F. (2014). Visual signals or displacement activities? The function of

337 visual displays in agonistic interactions in nocturnal tree frogs. *Acta Ethologica*, *17*(1), 9–

338 14. https://doi.org/10.1007/s10211-013-0160-6

339 Hagman, M., & Shine, R. (2008). Deceptive digits: the functional significance of toe waving by

340 cannibalistic cane toads, *Chaunus marinus*. *Animal Behaviour*, *75*(1), 123–131.

341 https://doi.org/10.1016/j.anbehav.2007.04.020

342 Hartmann, M. T., Giasson, L. O. M., Hartmann, P. A., & Haddad, C. F. B. (2005). Visual

343 communication in Brazilian species of anurans from the Atlantic forest. *Journal of Natural*

344 *History*, *39*(19), 1675–1685. https://doi.org/10.1080/00222930400008744

345 Hödl, W., & Amézquita, A. (2001). Visual Signaling in Anuran Amphibians. In M. J. Ryan (Ed.),

346 *Anuran communication* (pp. 121–141). Washington: Smithsonian Institute Press.

347 James, L. S., Baier, A. L., Page, R. A., Clements, P., Hunter, K. L., Taylor, R. C., & Ryan, M. J.

348 (2022). Cross-modal facilitation of auditory discrimination in a frog. *Biology Letters*, *18*(6).

349 https://doi.org/10.1098/rsbl.2022.0098

350 Jarman, A. P. (2002). Studies of mechanosensation using the fly. *Human Molecular Genetics*,

351 *11*(10), 1215–1218. https://doi.org/10.1093/hmg/11.10.1215

352 Landestoy, M. A., & Ortíz, R. (2015). Rediscovery of the Eastern Crested Toad. *IRCF Reptiles &*

353 *Amphibians*, *22*(2), 50–55.

354 McFadden, M., Harlow, P. S., Kozlowski, S., & Purcell, D. (2010). Toe-twitching during feeding

355 in the Australian myobatrachid frog, *Pseudophryne corroboree*. *Herpetological Review*,

356 *41*(2), 153–154.

357 Narins, P. M. (1990). Seismic in Communicati Anuran Amphibians. *BioScience*, *40*(4), 268–274.

358 Retrieved from https://doi.org/10.2307/1311263

359 Narins, P. M., Grabul, D. S., Soma, K. K., Gaucher, P., & Hödl, W. (2005). Cross-modal

360 integration in a dart-poison frog. *Proceedings of the National Academy of Sciences of the*

United States of America, *102*(7), 2425–2429. https://doi.org/10.1073/pnas.0406407102

Posit Team. (2023). RStudio: Integrated Development Environment for R.

Rojas, B., & Pašukonis, A. (2019). From habitat use to social behavior: Natural history of a

voiceless poison frog, *Dendrobates tinctorius*. *PeerJ*, *7*, e7648.

https://doi.org/10.7717/peerj.7648

Rome, L. C. (2006). Design and function of superfast muscles: New insights into the physiology

of skeletal muscle. *Annual Review of Physiology*, *68*, 193–221.

https://doi.org/10.1146/annurev.physiol.68.040104.105418

Santos, J. C., Coloma, L. A., & Cannatella, D. C. (2003). Multiple, recurring origins of

aposematism and diet specialization in poison frogs. *Proceedings of the National Academy*

of Sciences, *100*(22), 12792–12797. https://doi.org/10.1073/pnas.2133521100

Schulte, L. M., & König, Y. (2023). Experimental evidence that toe-tapping behavior in the

green-and-black poison frog (*Dendrobates auratus*) is related to prey detection. *Acta*

Ethologica, *26*(2), 139–143. https://doi.org/10.1007/s10211-023-00422-8

Sloggett, J. J., & Zeilstra, I. (2008). Waving or tapping? Vibrational stimuli and the general

function of toe twitching in frogs and toads (Amphibia: Anura). *Animal Behaviour*, *76*(5), 1–

4. https://doi.org/10.1016/j.anbehav.2008.08.005

Starnberger, I., Maier, P. M., Hödl, W., & Preininger, D. (2018). Multimodal signal testing reveals

gestural tapping behavior in spotted reed frogs. *Herpetologica*, *74*(2), 127–134.

https://doi.org/10.1655/Herpetologica-D-17-00053.1

Team, R. C. (2022). R: A language and environment for statistical computing.

Toledo, L. F., Araújo, O. G. S., Guimarães, L. D., Lingnau, R., & Haddad, C. F. B. (2007). Visual

and acoustic signaling in three species of Brazilian nocturnal tree frogs (anura, hylidae).

Phyllomedusa, *6*(1), 61–68. https://doi.org/10.11606/issn.2316-9079.v6i1p61-68

Vergara-Herrera, N., Cocroft, R., & Rueda-Solano, L. A. (2023). Eating to the beat of the drum:

vibrational parameters of toe tapping behavior in *Dendrobates truncatus* (Anura:

Dendrobatidae). *Evolutionary Ecology*. https://doi.org/10.1007/s10682-023-10277-x

Warkentin, K. M. (2005). How do embryos assess risk? Vibrational cues in predator-induced

hatching of red-eyed treefrogs. *Animal Behaviour*, *70*(1), 59–71.

https://doi.org/10.1016/j.anbehav.2004.09.019

391