Title: Enhancing KCC2 function reduces interictal activity and prevents seizures in mesial temporal lobe epilepsy

Authors: Florian Donneger[1,2,3], Adrien Zanin[1,2,3], Jeremy Besson[1,2,3], Yoness Kadiri[1,2,3], Carla Pagan[1,2,3], Nicolas David[1,2,3], Marion Russeau[1,2,3], Franck Bielle[2,4,5], Bertrand Devaux[6†], Bertrand Mathon[2,5,7], Vincent Navarro[2,5,7], Francine Chassoux[6†], Jean Christophe Poncer[1,2,3]*

Affiliations:

[1] Inserm UMR-S 1270; Paris, France

[2] Sorbonne Université; Paris, France

[3] Institut du Fer à Moulin; Paris, France

[4] AP-HP, Hôpitaux Universitaires La Pitié Salpêtrière - Charles Foix, Department of Neuropathology, F-75013; Paris, France

[5] Paris Brain Institute, INSERM, CNRS; Paris, France,

[6] Department of Neurosurgery, Sainte-Anne Hospital; Paris, France

[7] AP-HP, Epilepsy Unit, Reference Center for Rare Epilepsies, and Department of Clinical Neurophysiology, Pitié Salpêtrière Hospital; Paris, France

*Corresponding author: Jean Christophe Poncer, Institut du Fer à Moulin, 8 rue des Fossés Saint Marcel, 75005 Paris, France. Tel. +33145876118, E-mail: jean-christophe.poncer@inserm.fr

† Present address: Neurosurgery Department, Lariboisière Hospital; Paris, France.

Abstract: The neuronal K/Cl cotransporter KCC2 controls intraneuronal chloride and subsequently the efficiency of GABA signaling. In many neurological disorders, including mesial temporal lobe epilepsy (mTLE), reduced KCC2 expression or function may lead to depolarizing GABA signaling, which may then contribute to pathological activity and seizures. Compensating for the dysregulation of chloride transport in the pathology therefore appears to be a promising therapeutic strategy. Two small molecules - prochlorperazine (PCPZ) and CLP-257 - recently identified from library screening as candidate KCC2 enhancers, appear to improve symptoms in animal models of disorders associated with KCC2 extinction. However, their mode of action in cortical neurons and their therapeutic potential in epilepsy remain elusive and even controversial. Here, we show in rat hippocampal neurons that PCPZ and CLP-257 act by increasing KCC2 function and clustering while decreasing its membrane diffusion, independent of phosphorylation of canonical regulatory residues. Consistently, both compounds suppress spontaneous interictal-like discharges in postoperative tissue from patients with mTLE. Chronic administration of PCPZ also reduces seizure frequency and interictal activity in a mouse model of mTLE. These results unravel the mechanism of action of two KCC2 enhancers and validate their therapeutic potential in mesial temporal lobe epilepsy.

Introduction

Epilepsy is among the most frequent neurological disorders, affecting more than 65 million people worldwide *(1)*. Mesial temporal lobe epilepsy (mTLE), the most common form of focal epilepsy in adults, is frequently associated with hippocampal sclerosis and often refractory to antiseizure medication *(2)*. Surgical resection of the epileptic focus has been shown to offer a fairly high rate (60-75%) of seizure control at 2 years in patients with mTLE *(3)*. However, surgery cannot be proposed to all patients with mTLE. In addition, it can have negative effects on cognitive functions and fewer than 50% remain seizure-free 10 years after surgery *(4)*, highlighting the need for novel therapeutic approaches in mTLE.

Several forms of pharmaco-resistant epilepsies, including mTLE, have been associated with altered expression and function of cation-chloride cotransporters (CCCs), which control transmembrane chloride gradients *(5–10)*. In neurons specifically, the potassium-chloride cotransporter KCC2 maintains a low intracellular chloride concentration, thereby ensuring a hyperpolarizing and inhibitory current upon activation of the chloride-permeable $GABA_A$ receptors *(11)*. Reduced KCC2 expression and function, as often reported in mTLE *(7, 9, 12)* but see *(13)*, leads to elevated intraneuronal chloride and may reduce or even reverse the driving force for $GABA_A$ receptors *(7, 14, 15)*. Depolarizing and excitatory GABA signaling may then paradoxically enhance neuronal activity and is thought to promote anomalous synchronization underlying epileptiform activities *(16–18)*. Thus, pharmacologically rescuing intraneuronal chloride using bumetanide, an antagonist of the chloride importer NKCC1, suppresses interictal-like activity in resective human temporal lobe tissue *in vitro (7)*. Similarly, mutations that promote KCC2 membrane stability and function *(19)* or overexpression of recombinant KCC2

(20) reduced acute seizures in animal models. Compensating for CCC dysfunction therefore

appears as a promising therapeutic strategy in pharmacoresistant epilepsy.

Bumetanide, however, failed to prevent acute neonatal seizures in two clinical trials and induced

several side-effects due to the broad peripheral NKCC1 expression *(21, 22)*. Instead, enhancing

the function of the neuron-specific chloride extruder KCC2 is expected to rescue neuronal

chloride transport in the pathology with fewer side effects. Recently, high-throughput screening

studies of compound libraries have identified the phenothiazine-derived antipsychotic

prochlorperazine (PCPZ) and the arylmethylidine family compounds CLP-257 and CLP290, as

enhancers of KCC2 function in heterologous cell lines *(23, 24)*. These compounds were also

reported to improve symptoms in animal models of disorders associated with reduced KCC2

expression *(24–28)*. However, their mode of action remains unknown and their efficacy and

selectivity towards KCC2 were recently disputed *(29, 30)*. In addition, CLP-257 was shown to

either increase *(31)* or decrease *(32)* the duration of epileptiform discharges *in vitro*, and direct

experimental evidence of the therapeutic efficacy of KCC2 enhancers in mTLE is missing.

Here, we first investigated the efficacy and mode of action of PCPZ and CLP-257 on KCC2 in

rat hippocampal neurons. We show that both compounds enhance KCC2 function and that CLP-

257 additionally acts as a positive allosteric modulator of extrasynaptic but not synaptic $GABA_A$

receptors. Both PCPZ and CLP-257 reduced KCC2 membrane diffusion and enhanced KCC2

clustering, without affecting total or plasmalemmal KCC2 expression levels, or phosphorylation

of its canonical regulatory residues. *In vitro*, both KCC2 enhancers effectively suppressed

spontaneous, interictal-like discharges in postoperative tissue from mTLE patients. Finally,

chronic PCPZ administration reduced the occurrence of seizures and other epileptiform activities

in a lithium-pilocarpine mouse model of mTLE. Collectively, our data confirm the efficacy and

decipher the mode of action of two candidate KCC2 enhancers, and demonstrate for the first time

their antiepileptic potential in a form of pharmacoresistant epilepsy.

Results

PCPZ and CLP-257 act as KCC2 enhancers in rat hippocampal neurons

We first tested whether PCPZ and CLP-257, two candidate KCC2 enhancers *(23, 24)*, potentiate KCC2 in rat hippocampal neurons *in vitro*. KCC2 function was assessed by measuring the somato-dendritic gradient of the reversal potential of currents evoked by focal GABA$_A$ receptor activation (E$_{GABA}$) in whole-cell recordings with high-chloride internal solution (**Fig. 1**, see Supplementary Material and Methods). As shown in previous studies, this gradient is nearly abolished by a specific KCC2 antagonist *(33, 34)*, but unaffected by bumetanide *(35)*, therefore providing a reliable proxy for KCC2 function. We performed a 2-hour treatment with CLP-257 or PCPZ, a paradigm used in previous studies with CLP-257 *(23, 36)*. The mean somato-dendritic E$_{GABA}$ gradient (Δ_{EGABA}) was 6.6±0.8 mV/100 µm in control neurons (DMSO; n=19). In neurons pre-incubated with PCPZ (10 µM), this gradient was significantly increased to 11.5±1.5 mV/100 µm (n=16; p=0.006; **Fig. 1, A** to **C**). Similarly, pre-incubation with CLP-257 (100 µM) increased ΔE_{GABA} from 5.3±1.1 to 9.7±1.1 mV/100 µm (n=10 and 8, respectively; p=0.014) (**Fig. 1, D** to **F**). Neither compound had an effect on neuronal membrane resistance or series resistance (Fig. S1). These data demonstrate that both PCPZ and CLP-257 effectively potentiate KCC2 function in rat hippocampal neurons.

CLP-257 but not PCPZ directly modulates GABA$_A$ receptor function

CLP-257 has been suggested to act directly to potentiate GABA$_A$ receptors *(29)* (but see *(30)*). Because GABA$_A$ receptor-mediated chloride fluxes modulate KCC2 membrane expression and function *(37)*, we asked whether PCPZ and CLP-257 may act primarily to modulate GABA$_A$

receptor function. We therefore tested their effect on GABA$_A$ receptor-mediated currents, using

focal application of isoguvacine (100 µM) or uncaging of low concentration (30 µM) of Rubi-

GABA, respectively (**Fig. 2**). PCPZ had no significant effect on the amplitude (157.5±25.4 vs.

150.7±17.5 pA for control, p=0.48) or decay time constant (130.8±10.5 vs. 127.7±10.2 ms,

p=0.362) of GABA$_A$ receptor-mediated currents (**Fig. 2, A** to **B**). In contrast, CLP-257 rapidly

increased their decay time constant (218.2±21.1 vs. 124.4±9.1 ms, p<0.001) with no detectable

effect on their amplitude (162.0±6.6 vs 164.7±4.6 pA for control, p=0.66; **Fig. 2, C** to **D**). The

effect of CLP-257 did not appear to be dose-dependent, as previously suggested *(30)*, since

application of a 10-fold lower concentration (10 µM) similarly increased the decay time constant

of GABA$_A$ receptor-mediated currents (294.1±33.8 vs. 239.7±27.2 ms, p<0.001; Fig. S2).

We next tested the effect of PCPZ and CLP-257 specifically on synaptic GABA$_A$ receptors, by

recording miniature inhibitory postsynaptic currents (mIPSCs) from hippocampal neurons.

Neither PCPZ nor CLP-257 had detectable effects on mIPSC amplitude (42.9±2.6 vs. 42.5±2.5

pA, p=0.797 and 43.8±2.9 vs. 42.3±3.4 pA, p=0.622, respectively) or decay time constant

(10.2±0.7 vs. 10.7±0.8 ms, p=0.086 and 12.5±1.0 vs. 12.4±0.8 ms, p=0.86, respectively). Taken

together, these results demonstrate that while both PCPZ and CLP-257 potentiate KCC2

function, CLP-257 also acts to specifically potentiate high-affinity, presumably extrasynaptic but

not synaptic GABA$_A$ receptor function in a benzodiazepine-like manner in rat hippocampal

neurons.

PCPZ and CLP-257 promote KCC2 membrane confinement and clustering without

affecting its total or plasmalemmal expression levels

We then investigated the mechanisms underlying the potentiation of KCC2 by PCPZ and CLP-

257. Enhancement of KCC2 function may result from either an increase in intrinsic enzymatic

activity, an increase in membrane expression of the transporter, or both *(38)*. We therefore used

surface biotinylation to test whether PCPZ or CLP-257 influenced plasmalemmal KCC2

expression. Incubation with PCPZ or CLP-257 for 2 hours had no significant effect on total

KCC2 expression (**Fig. 3A**, p=0.137 and p=0.893, respectively) or on the ratio of surface to total

expression (**Fig. 3B**, p=0.735 and p=0.8, respectively). Therefore, enhanced KCC2 function by

PCPZ or CLP-257 does not reflect an increased expression of the transporter. PCPZ and CLP-

257 also had no effect on total NKCC1 expression (p=0.100 and p=0.700, respectively) or on the

ratio of surface to total NKCC1 expression (p=0.700 and p=0.700, respectively; Fig. S3).

KCC2 function is regulated by fine tuning of its diffusion and clustering properties at the plasma

membrane *(39)*. Thus, the potentiation of KCC2 function is often associated with decreased

membrane diffusion and increased clustering *(37, 40–42)*. Membrane lateral diffusion of

recombinant Flag-tagged KCC2 was assessed in hippocampal neurons using single particle

tracking with quantum dots. The lateral diffusion of KCC2 was reduced by both PCPZ and CLP-

257, both in the vicinity of GABAergic synapses and outside. Thus, both compounds reduced the

diffusion coefficient of recombinant KCC2 (extrasynaptic: -36.5±14.9% p<0.001; synaptic: -

26.2±14.0% p=0.011 for PCPZ and extrasynaptic: -20.7±4.9% p=0.003; synaptic: +9.4±11.2%

p=0.97 for CLP-257) as well as its explored area (extrasynaptic: -39.7±17.2%   p<0.001;

synaptic: -30.6±25.4% p<0.001 for PCPZ and extrasynaptic: -25.1±2.7%   p<0.001; synaptic:

+4.9±19.4% p=0.80 for CLP-257; **Fig. 3, C** to **D**).

This effect was associated with an increase in KCC2 clustering, as shown by

immunocytochemistry assays revealing an increase in cluster intensity (+109.0±61.4% p<0.001

and +33.9±13.6% p=0.005, respectively) and area (+45.3±18.1% p<0.001 and +36.3±20.4%

p<0.001, respectively), without significant change in cluster density (-6.6±8.4% p=0.13 and -

3.5±4.2% p=0.34; **Fig. 3, E** to **F**). Furthermore, the effect of PCPZ on KCC2 clustering did not

involve D2R antagonism, as a 2-hour treatment with eticlopride did not significantly affect the

intensity, area, or density of KCC2 clusters (Fig. S4).

Overall, our results demonstrate that PCPZ and CLP-257 act to enhance net KCC2 function and

promote its clustering, with no effect on total or plasmalemmal KCC2 expression levels.

Therefore, PCPZ and CLP-257 likely induce a redistribution of membrane-inserted KCC2,

resulting in increased clustering and function.

PCPZ and CLP-257 do not affect KCC2 phosphorylation on canonical regulatory residues

The stability, clustering, and membrane turnover of KCC2 are controlled by a variety of post-

translational mechanisms, including phosphorylation of key residues, particularly in its large

carboxy-terminal domain, that regulate its net transport function *(38, 39, 43)*. Phosphorylation of

serine 940 (S940) by protein kinase C is known to increase the membrane stability and function

of KCC2 *(40, 44)*. In contrast, phosphorylation of threonine residues 906 and 1007 (T906,

T1007) by the With-no-lysine (WNK) kinase pathway has been shown to negatively regulate

KCC2 function in cortical neurons by affecting its membrane diffusion and clustering *(37, 45)*.

We therefore asked whether PCPZ and CLP-257 could act by modulating the phosphorylation of

these residues to promote KCC2 function. Immunoprecipitation assays on hippocampo-cortical

slices from adult rats (**Fig. 4, A** to **B**, see Supplementary Material and Methods) revealed no

significant effect of PCPZ or CLP-257 on KCC2 phosphorylation at either S940 (-19.7±34.4%,

p=0.597 and +20.3±18.7%, p=0.339), T1007 (-18.5±17.1%, p=0.340 and +26.4%±17.0%

p=0.195 respectively) or T906 residues (-12.7±10.9%, p=0.308 and -10.2±10.4%, p=0.384

respectively). Consistent with these results, PCPZ and CLP-257 also had no significant effect on

the phosphorylation of the WNK effectors SPAK (S373, +33.3±24.0%, p=0.238 and

+36.9±48.4%, p=0.488 respectively) and OSR1 (S325, +11.5±18.5%, p=0.568 and

+11.8±23.5%, p=0.641, respectively) or their total expression levels (SPAK, -6.0±17.4%,

p=0.748 and -12.3±6.4%, p=0.124, respectively; OSR1, -2.2 ±8.3%, p=0.800 and +26.3±12.8%,

p=0.108, respectively; **Fig. 4, C** to **D**). In contrast, the pan-WNK inhibitor WNK463 *(46)*

reduced phosphorylation of T1007 (-78.7 ±3.3%, p<0.001) and to a lesser extent S940 (-

14.4±4.7%, p=0.037) but not T906 (-13.7 ±12.5%, p=0.333; **Fig. 4, A** to **B**). These effects of

WNK463 were associated with reduced phosphorylation of SPAK (S373, -84.1±4.6%, p<0.001)

and OSR1 (S325, -81.0±5.6%, p<0.001; **Fig. 4, C** to **D**). Thus, PCPZ and CLP-257 appear to

potentiate KCC2 clustering and function independent of detectable changes in phosphorylation

of serine 940 and threonine 906 and 1007 residues. Consistent with this conclusion, these

compounds showed no direct inhibitory effect on the activity of WNK1, WNK3, SPAK or OSR1

in *in vitro* kinase assays (Fig. S5).

Finally, phosphorylation of residues Y903 and Y1087 has been reported to affect KCC2

clustering and function *(42, 47)*. Therefore, we tested the effect of PCPZ and CLP-257 on KCC2

tyrosine phosphorylation in hippocampo-cortical slices from adult rats (**Fig. 4, E** to **F**). Whereas

sodium orthovanadate (Na3VO4), a tyrosine phosphatase inhibitor, increased tyrosine

phosphorylation of KCC2 (+89.1±45.4%, p=0.029), PCPZ and CLP-257 showed no significant

effect (+6.4±14.7% and +19.0±15.0%, p=1 and p=0.343, respectively). Together, our results

demonstrate that PCPZ and CLP-257 potentiate KCC2 function, reduce its diffusion and increase

its clustering via a mechanism independent of phosphorylation of canonical regulatory residues

in its carboxy-terminal domain.

PCPZ and CLP-257 prevent spontaneous interictal-like activity in postoperative tissue

from mTLE patients *in vitro*

Human epileptic tissue therapeutically resected from patients diagnosed with mTLE provides an

invaluable tool for evaluating antiepileptic drug candidates in the pathophysiological context of

human mTLE *(48, 49)*. We therefore tested the effect of PCPZ and CLP-257 on 13 tissue

samples resected from patients diagnosed with mTLE, 11 of which were associated with

hippocampal sclerosis (HS) (**Table 1**). As previously reported *(6, 7)*, a majority of slices (29/46)

prepared from 13 postoperative tissues and recorded using multi-electrode arrays (MEA) in

control conditions (DMSO, 2 hours) displayed spontaneous, low-frequency (<40Hz) events

identified as interictal-like discharges (IILDs, **Fig. 5**). IILDs often originated from only a small

subregion of the slice, usually in the subiculum or the dentate gyrus (**Table 1**, **Fig. 5, A** to **B**). In

tissue preincubated with DMSO alone, IILDs had a mean amplitude of 57.3±13.5 µV and

occurred at a mean frequency of 0.4±0.1 Hz. As observed previously *(7)*, the NKCC1 antagonist

bumetanide (10 µM) partially reduced the amplitude and frequency of IILDs (**Fig. 5B**).

However, these events were almost completely abolished in slices of the same resected tissue

incubated for 2 hours with either 10 µM PCPZ (**Fig. 5B**) or 100 µM CLP-257 (**Fig. 5C**). Overall,

both PCPZ and CLP-257 significantly reduced the proportion of slices displaying spontaneous

IILDs (p<0.05 for both; **Fig. 5D**). This effect did not reflect compromised neuronal activity, as

IILD and multi-unit activity could be restored in slices pre-treated with PCPZ or CLP-257 by

increasing neuronal excitability with high potassium, low magnesium ACSF (**Fig. 5, E** to **F**).

In slices from a group of 5 patients with mTLE (patients #5, 7, 8, 11, 12; group I; see **Table 1**),

PCPZ virtually abolished IILDs (-97.6±13.8% change in frequency, p<0.001; **Fig. 5G**). In one

slice of patient #3 and in the slices of patient #6 (group II), however, PCPZ failed to reduce the

frequency of IILDs (+25.5±15.9%, p=0.356; **Fig. 5G**). Interestingly, neuropathological

exploration of the resected tissue from patient #6 revealed an ILAE type 3 HS (**Table 1**), a rarer

pattern of HS often associated with dual pathology *(50)*.

Similarly, preincubation with CLP-257 virtually abolished IILDs in slices from 5 out of 6

patients (-96.5±2.3% change in frequency, p<0.001; **Fig. 5G**). In slices from the remaining

patient (#9, **Table 1**), however, CLP-257 failed to reduce the frequency of IILDs (+12.6%, **Fig.**

5G). Interestingly, again, neuropathological examination of this patient's postoperative tissue did

not reveal hippocampal sclerosis, but rather a ganglioglioma associated with a somatic

BRAF^{V600E} mutation. Overall, these data demonstrate that enhancing KCC2 function prevents

epileptiform activity recorded *in vitro* in postoperative tissue from a majority of patients with

mTLE and typical ILAE type 1 HS.

PCPZ reduces interictal activity and seizures in a mouse model of mTLE

Spontaneous ictal activity is rarely observed in postoperative mTLE tissue slices *in vitro (48,*

51). To assess the anti-seizure potential of KCC2 enhancers, we therefore tested their effect in a

lithium-pilocarpine model of mTLE *(52)* adapted to mice (**Fig. 6A**). Since PCPZ but not CLP-

257 *(23)* can be readily used for chronic *in vivo* administration, we conducted these experiments

with PCPZ. Status epilepticus (SE) was induced by intraperitoneal injection of the muscarinic

agonist pilocarpine (see Supplementary Material and Methods). Mice typically developed SE

within 10 to 20 minutes after pilocarpine injection and SE was interrupted after 1 hour by i.p.

injection of diazepam and ketamine. Telemetric ECoG recordings revealed intense interictal

discharge (IID) activity and occasional seizures for hours to a few days after SE, followed by a

seizure-free, latent period that lasted 3 to 4 weeks before regular, spontaneous seizures were

detected (**Fig. 6B, E**). Typical epileptiform activity such as interictal discharges (<20 Hz) or

high-frequency oscillations (>80 Hz, HFOs, *(53–55)*) were also observed (**Fig. 6G** to **H**). In

addition to these electrophysiological biomarkers, typical histological landmarks were detected

in the hippocampus of these mice, including hippocampal sclerosis, granule cell dispersion,

mossy fiber sprouting, and reduced KCC2 immunoreactivity in the dentate gyrus (**Fig. 6C** to **D**).

After a 5-day habituation period with twice-daily saline injections (days 35-39, control period),

we tested the effect of saline vs. PCPZ (2 mg/kg, i.p., 9AM and 8PM) on the occurrence of

seizures and epileptiform events (IIDs, HFOs) during the next 5 days (days 40-44 post-SE, test

period) (**Fig. 6A, E**).

Whereas seizure frequency was not significantly affected in animals receiving saline during the

test period (0.15±0.07 vs. 0.18±0.06 seizures/hour, n=8 animals, p=0.118), animals receiving

PCPZ showed an approximately 40% reduction in seizure frequency compared with the control

period (0.09±0.05 vs. 0.14±0.05 seizures/hour, p=0.01; **Fig. 6F**). However, PCPZ had no effect

on seizure duration (29.8±1.8 vs. 31.3±3.8 s, p=0.657) or power (7422±2736 vs. 6695±2058

mV^2, p=0.466; **Fig. 6F**), suggesting that it may be acting on mechanisms involved in seizure

onset rather than maintenance. In addition, and in agreement with our *in vitro* results on

postoperative human tissue, PCPZ significantly reduced the frequency of IIDs by about 34%

compared with control (192.0±39.3 vs. 290.9±60.3 IID/hour, p=0.01; **Fig. 6G**). The effect of

PCPZ was even stronger on the frequency of HFOs, with a 58% reduction in mice receiving

PCPZ (54.61±23.73 vs. 128.68±71.88 HFO/hour p=0.012), whereas HFO frequency increased

by 38% in the saline only group (102.5±36.6 vs. 64.7±23.7 HFO/hour, p=0.027; **Fig. 6H**).

Interestingly, although we observed a circadian rhythmicity in the occurrence of seizures and

interictal activity (Fig. S6), as previously reported *(56, 57)*, the effect of PCPZ was similar

during light (Zeitgeber time 0-12 h) and dark (Zeitgeber time 12-24 h) phases. These data

demonstrate that PCPZ reduces chronic seizures and epileptiform activities (including HFOs and

IIDs) in an animal model of mTLE.

Discussion

In this study we evaluated the effect and mode of action of two candidate KCC2 enhancers, PCPZ and CLP-257, in hippocampal neurons. Both reduce the lateral diffusion of KCC2, and increase its clustering and transport function. This effect is independent of changes in KCC2 expression and the phosphorylation level of its canonical residues and the WNK/SPAK pathway. In addition to this effect on KCC2 function, we also showed that CLP-257 directly modulates extrasynaptic but not synaptic $GABA_A$Rs. Furthermore, both compounds suppressed spontaneous, interictal-like discharges recorded in postoperative hippocampal tissue resected from patients with refractory mTLE. Finally, chronic administration of PCPZ reduced the occurrence of seizures and the frequency of interictal events in a mouse model of mTLE. Thus, our results provide a proof-of-concept of the therapeutic potential of KCC2 enhancers in a drug-resistant form of epilepsy and suggest that these compounds may also have therapeutic potential in other pathologies involving decreased KCC2 expression.

Several forms of epilepsy are associated with chloride transport dysfunction *(58)*, leading to paradoxically excitatory signaling by $GABA_A$ receptors *(7, 10)*. Although pro-GABAergic drugs such as benzodiazepines and other $GABA_A$ receptor positive allosteric modulators are often effective in epilepsy *(59)*, acting to directly potentiate GABA signaling when intraneuronal chloride levels are abnormal may fail to prevent seizures. Restoration of neuronal transmembrane chloride gradients therefore appears to be a promising first-line therapeutic strategy in drug-resistant epilepsy. In principle, this may be achieved by blocking chloride import, using NKCC1 antagonists such as bumetanide. Although this approach has been successfully used in some animal models of CNS disorders *(60–63)* it is not optimal as i)

bumetanide is largely bound to plasma proteins and very poorly crosses the blood brain barrier,

ii) undergoes active efflux transport from the brain by organic anion transporters *(64, 65)* and iii)

NKCC1 is widely expressed in many peripheral organs and tissues *(66)*, raising issues regarding

the number of undesirable side effects. Instead, KCC2 is almost exclusively expressed in

neurons, making it a prime target for neurological disorders associated with deficits in neuronal

chloride homeostasis, such as mTLE. Acting to promote KCC2 function, even when its

expression may be reduced in the pathology, may then help rescuing neuronal transmembrane

chloride gradients and inhibitory GABA signaling.

CL-058 was the first compound identified as a candidate KCC2 enhancer by a small molecule

library screening in a rodent neuroblastoma-glioma hybrid cell line *(23)*. It was then chemically

optimized into CLP-257 and CLP-290, with improved EC50 and half-life for *in vivo*

applications. While CLP drugs were initially described as specific KCC2 enhancers with little or

no activity on other cation-chloride cotransporters *(23)*, a more recent study challenged this view

and showed no effect on KCC2 function *(29)*. Instead, this study found direct binding and

potentiating action on $GABA_A$ receptors. Although still controversial *(30)*, these results have

called into question CLP drugs as pure KCC2 enhancing molecules. Our results now demonstrate

that CLP-257 acts both as a *bona fide* KCC2 enhancer (Fig. 1) and a positive allosteric

modulator of extrasynaptic, but not synaptic, $GABA_A$ receptors (Fig. 2) in rat hippocampal

neurons. These data thus reconcile apparently conflicting results obtained on tonic $GABA_A$

receptor-mediated currents in hippocampal neurons *(29)* and mIPSCs from spinal cord neurons

(30). PCPZ on the other hand is a first-generation antipsychotic and is generally known as a

dopamine D2 antagonist, although it interacts with a lower affinity with serotoninergic,

adrenergic, muscarinic and histaminergic receptors *(67–69)*. It was first suggested as a candidate

KCC2 enhancer by screening Prestwick's library on HEK cells expressing recombinant KCC2

(24), and was shown to reduce the depolarizing driving force of evoked IPSCs in immature rat

spinal motoneurons after spinal cord injury. However, its effects on cortical neurons have not

been investigated. Our results now show that PCPZ also acts as an effective KCC2 enhancer,

increasing KCC2 function with no detectable effect on NKCC1. In contrast to CLP-257,

however, PCPZ did not affect either phasic or tonic $GABA_A$ receptor-mediated IPSCs (Fig. 2).

Despite their potential therapeutic interest in rescuing chloride transport function in the

pathology, the mechanisms of action of CLP-257 and PCPZ on KCC2 have not been fully

characterized. Several studies have reported an increased total *(27, 28)* or plasmalemmal *(23, 26,*

70) KCC2 expression by CLP compounds, mostly in animal models of chronic pain or spasticity.

Similarly, PCPZ was shown to rescue KCC2 membrane immunoreactivity in lumbar

motoneurons after spinal cord injury *(24)*. However, the underlying mechanisms of these effects

were not further investigated. We did not observe such changes in KCC2 expression upon

incubation with PCPZ or CLP-257 in hippocampal neurons. Thus, both biochemical and

immunofluorescence assays revealed no change in either total or plasmalemmal KCC2

expression levels (Fig. 3 and 4). Instead, increased KCC2 function was associated with reduced

membrane lateral diffusion and increased clustering of KCC2 (Fig. 4). These observations

suggest that, at least under non-pathological conditions, PCPZ and CLP-257 may primarily

induce a redistribution of the membrane pool of KCC2 rather than a change in expression levels.

Previous studies have revealed that KCC2 lateral diffusion and clustering in the plasma

membrane are tightly correlated with function and are controlled by phosphorylation of key

residues in its carboxy-terminal domain *(11, 39, 40, 71)*, making them prime candidate targets

for KCC2 modulators. In particular, phosphorylation of serine 940 and dephosphorylation of

threonines 906 and 1007 promote KCC2 clustering and membrane stability *(37, 40, 45, 47, 72)*.

This effect usually results in increased membrane expression of the transporter, which was not

observed upon PCPZ or CLP-257 treatment, making it unlikely that a modification of their

phosphorylation underlies the effect of these compounds. Consistent with this conclusion, CLP-

257 and PCPZ produced no significant change in the phosphorylation of these residues (Fig. 4)

and did not affect the WNK/SPAK/OSR1 pathway known to phosphorylate T906 and T1007 *(37,*

72). These observations are also consistent with the lack of effect of CLP290 on the

phosphorylation of these residues in a mouse model of ischemia *(28)*. Finally, phosphorylation of

tyrosine 1087 has been suggested to specifically impact KCC2 membrane clustering and function

without affecting its membrane expression level *(47, 73, 74)* and would therefore be a good

candidate target for PCPZ and CLP-257. However, KCC2 tyrosine phosphorylation was also

unaffected in hippocampo-cortical slices upon PCPZ or CLP-257 treatment. Further

investigation will be needed to test whether compensatory phosphorylation and

dephosphorylation of distinct tyrosine residues may mask the net effect of KCC2 enhancers on

Y1087.

Taken together, our data demonstrate for the first time that both PCPZ and CLP-257 enhance

KCC2 function and promote its membrane confinement and clustering independent of apparent

changes in the phosphorylation state of its canonical regulatory sites. Interestingly, a recent study

has shown that other KCC2 enhancers with unknown mode of action display the same

potentiating effect without affecting KCC2 membrane expression or the phosphorylation of S940

or T1007 residues *(75)*. PCPZ, CLP-257 and probably other KCC2 enhancers may act on

alternative phosphorylation sites, as identified in large-scale proteomics studies *(75–77)*.

Alternatively, they may interfere with KCC2 molecular interactions, without necessarily binding

directly to KCC2. Thus, functional proteomics studies *(41, 76, 78)* have shown that KCC2

interacts with a variety of protein partners, some of which regulate its membrane expression and

clustering. In particular, KCC2 interaction with the SNARE proteins SNAP23 and syntaxin 1A is

essential for mZnR/GPR39-dependent upregulation of KCC2 activity *(79)*. Other candidates

such as CIP1 *(80)*, Neto2 *(81)*, CKB *(82)*, PAM *(83)* and gephyrin *(41)* are of particular interest

since their interaction with KCC2 was shown to increase its function. Finally, KCC2 enhancers

may act to promote KCC2 accumulation into lipid rafts *(42)*, resulting in enhanced transporter

membrane clustering and function.

Although some studies have reported effects of KCC2 enhancers on epileptiform activity *in vitro*

or acute seizures such as *status epilepticus* in animal models, our work is the first to evaluate the

therapeutic potential of KCC2 enhancers in the context of mTLE. Thus, CLP-257 was shown to

either increase *(31)* or decrease *(32)* ictal-like discharges (ILDs) in two *in vitro* models of

epileptiform activity. On the other hand, inhibition of the WNK/SPAK/OSR1 pathway with the

WNK inhibitor WNK463 prevented KCC2 T1007 phosphorylation and reduced the intensity of

kainate-induced *status epilepticus* in mice *(84)*. However, it remains to be determined whether

this effect is strictly dependent on KCC2 potentiation or other effects of WNK inhibition,

including on the NKCC1 transporter *(84)*. Finally, CLP290 was shown to reduce the sensitivity

to pentylenetetrazole-induced seizures in a neonatal ischemic mouse model *(28)* while other

KCC2 enhancers prevented the development of and terminated kainate-induced *status epilepticus*

(75). Unfortunately, this study did not investigate the effect of KCC2 enhancers on spontaneous

and recurrent seizures that characterize mTLE.

Our new data now demonstrate that both PCPZ and CLP-257 increase KCC2 function and

suppress spontaneous epileptiform activity recorded *in vitro* from human postoperative tissue

from mTLE patients as well as interictal activity and seizures in a mouse model of mTLE. It is

noteworthy that out of 13 temporal lobe resective tissues, PCPZ or CLP-257 near-completely

suppressed interictal-like activity in 10 samples, whereas it had virtually no effect in 3 of them

(Fig. 5). Neuropathological analysis of these patients revealed that they had rare forms of

temporal lobe epilepsy, one related to a ganglioglioma associated with a somatic mutation of the

BRAF proto-oncogene, and the other of ILAE type 3, often associated with a dual pathology.

Further investigation of the phenotype of these non-responding tissues will be necessary to

elucidate their pathophysiological characteristics and the origin of their resistance to KCC2

enhancers.

PCPZ is a first-generation antipsychotic with FDA approval and indications in schizophrenia,

migraine as well as nausea and vomiting *(85)*. Side effects are generally mild in adults but

adverse effects such as sedation, extrapyramidal symptoms, and in some rare cases, seizures,

have been reported in children *(86)*. Instead, we now provide evidence that PCPZ shows

significant antiepileptic properties both in postoperative temporal lobe tissue from mTLE

patients *in vitro*, and in a mouse model of mTLE. *In vitro*, the effect of PCPZ was at least

comparable to that of bumetanide as described in postoperative epileptic brain tissue from

patients with epilepsy of various etiology *(5, 7, 10)*, adding to the evidence that rescuing

neuronal chloride transport may be an effective anti-epileptic strategy. *In vivo*, PCPZ reduced

seizure occurrence by about two-fold (Fig. 6) but also strongly reduced interictal discharges and

high-frequency oscillations, two hallmarks of epileptic networks *(53–55). In vivo,* PCPZ was

used at a concentration of 2 mg/kg through i.p. injections. At this dosage in mice, PCPZ likely

acts mainly on D2 dopamine receptors, but without any side effect on motility, exploratory

behavior or motor coordination of the animals *(87).* It seems unlikely that the antiseizure effect

of PCPZ *in vivo* may be mediated by D2 receptor antagonism since reduced D2 receptor

expression or function is associated with increased, not decreased, seizure threshold *(88).* In

addition, D2 agonists, not antagonists, were shown to display antiepileptic properties *(88).*

Altogether, our results demonstrate that PCPZ, like CLP-257, is an effective KCC2 enhancer and

that KCC2 activation may be an effective strategy in mTLE in patients. The recent progress in

elucidating the molecular structure of KCC2 *(89–92)* provides a unique opportunity to explore

the molecular mechanisms underlying the modulation of its expression and function and design

more specific, structure-based KCC2 enhancers with improved efficacy and specificity.

Material and Methods

Study Design

The purpose of this study was to identify the effect and mode of action of two candidate KCC2

enhancers (CLP-257 and PCPZ) and to test their antiepileptic potential in mTLE. First, the effect

of these compounds on KCC2 and $GABA_AR$ function was investigated by performing *in vitro*

electrophysiology on rat hippocampal neurons. Secondly, the effect of CLP-257 and PCPZ on

KCC2 expression, diffusion and clustering was explored by combining biochemistry, single

particle tracking and immunocytochemistry experiments respectively, in rat hippocampal

neurons. Subsequently, changes in phosphorylation of canonical KCC2 regulatory residues after

treatment with these KCC2 enhancers were addressed by performing immunoprecipitation assays

in hippocampo-cortical slices from adult rats. Finally, the antiepileptic potential of these

compounds was investigated *in vitro* by performing multi-electrode array recordings on

postoperative tissue from mTLE patients and *in vivo* by conducting ECoG recordings in a mouse

model of mTLE.

The indicated number of observations corresponds to the number of cells for electrophysiology

and immunocytochemistry data, individual quantum dots for single particle tracking,

independent primary hippocampal cultures or animals for biochemistry, animals for *in vivo*

experiments, and slices or patients for human tissue recordings. Sample size selection for

experiments was based on published experiments, pilot studies, as well as in-house expertise.

Most of the experiment's analyses were blinded to the treatment conditions. In addition, semi-

automated analysis software was used to reduce human bias. Human hippocampal slices were

randomly assigned to either the control or treatment paradigm prior to conducting the

experiments. Finally, animals were randomly assigned to treatment groups with approximately

equal numbers in each condition.

Animal experimentation

All animal procedures were carried out according to the European Community Council

directive of 24 November 1986 (86/609/EEC) and the guidelines of the French Agriculture

and Forestry Ministry for handling animals (decree 87-848), and were approved by the

"Direction Départementale de la Protection des Populations de Paris" (authorization number

D75-05-22). Procedures had been assessed by the site's animal welfare structure and the local

ethics committee n°5 (Charles Darwin). Following the favorable opinion of the ethics committee

of our project APAFIS#24889-202008050923999 v4, we obtained the authorization from the

Ministry of Research and Higher Education to carry out the experiments on animals used for

scientific purposes.

Patients

Informed consent to participate in the study was obtained from patients or their legal guardians.

The protocol received a favorable opinion of a research ethics committee (CPP Île-de-France 1 -

14388, January 16, 2017). A small part of non-essential brain tissue for neuropathological

diagnostic purposes was attributed to research by the neurosurgeon during surgery, in agreement

with the neuropathologist and the neurologist. Brain samples were obtained between 2017 and

2021 from 13 patients (4 males, 9 females, age range: 16-63 years, Table 1) diagnosed with

mesial temporal lobe epilepsy that was associated with hippocampal sclerosis (11 patients),

glioneuronal tumor (1 patient) and hippocampal atrophy with malformation of cortical

development (polymicrogyria and subcortical heterotopia, 1 patient). Patients underwent

multimodal preoperative work-up including neurological, neuropsychological assessment video-

electroencephalographic recording, magnetic resonance imaging and fluorodeoxyglucose

positron emission tomography. In addition, stereoelectroencephalography was performed in 6

patients.

Corticectomies were performed at Sainte-Anne Hospital (9 patients) and at Pitié-Salpêtrière

Hospital (4 patients) Neuropathological diagnosis was made according to the classification of the

Diagnostic Methods Commission of the International League Against Epilepsy (ILAE) (*50*).

Statistical Analysis

Means are shown $\pm$ SEM, median values are indicated with their interquartile range (IQR, 25–

75%). Means were compared using Student's parametric t-test when statistical conditions were

verified (normal distribution and homoscedasticity). Otherwise, a non-parametric Mann-Whitney

(MW) test was performed. For paired measurements, Wilcoxon test were used unless statistical

conditions were verified for Student's paired t-test. Statistical tests were performed using

SigmaPlot 13 (Systat Software). Differences were considered significant for p-values less than 5

% (* $p \leq 0.05$, ** $p < 0.01$, *** $p < 0.001$). Error bars correspond to SEM.

List of Supplementary Materials:

Material and Methods

Fig. S1. PCPZ and CLP-257 do not affect membrane or series resistance.

Fig. S2. CLP-257 modulates GABA$_A$ receptor function at micromolar concentrations.

Fig. S3. PCPZ and CLP-257 do not affect NKCC1 total or plasmalemmal expression levels.

Fig. S4. The D2R antagonist eticlopride does not affect KCC2 clustering in hippocampal

neurons.

Fig. S5. PCPZ and CLP-257 do not inhibit kinase activity of the WNK/SPAK/OSR1 pathway.

Fig. S6. PCPZ reduces seizure frequency and interictal activity independent of their circadian

modulation.

References

References

1. A. K. Ngugi, C. Bottomley, I. Kleinschmidt, J. W. Sander, C. R. Newton, Estimation of the

burden of active and life-time epilepsy: A meta-analytic approach. *Epilepsia* **51**, 883–890

(2010).

2. I. Blumcke, R. Spreafico, G. Haaker, R. Coras, K. Kobow, C. G. Bien, M. Pfäfflin, C. Elger,

G. Widman, J. Schramm, A. Becker, K. P. Braun, F. Leijten, J. C. Baayen, E. Aronica, F.

Chassoux, H. Hamer, H. Stefan, K. Rössler, M. Thom, M. C. Walker, S. M. Sisodiya, J. S.

Duncan, A. W. McEvoy, T. Pieper, H. Holthausen, M. Kudernatsch, H. J. Meencke, P. Kahane,

A. Schulze-Bonhage, J. Zentner, D. H. Heiland, H. Urbach, B. J. Steinhoff, T. Bast, L. Tassi, G.

Lo Russo, C. Özkara, B. Oz, P. Krsek, S. Vogelgesang, U. Runge, H. Lerche, Y. Weber, M.

Honavar, J. Pimentel, A. Arzimanoglou, A. Ulate-Campos, S. Noachtar, E. Hartl, O. Schijns, R.

Guerrini, C. Barba, T. S. Jacques, J. H. Cross, M. Feucht, A. Mühlebner, T. Grunwald, E. Trinka,

P. A. Winkler, A. Gil-Nagel, R. Toledano Delgado, T. Mayer, M. Lutz, B. Zountsas, K.

Garganis, F. Rosenow, A. Hermsen, T. J. von Oertzen, T. L. Diepgen, G. Avanzini,

Histopathological Findings in Brain Tissue Obtained during Epilepsy Surgery. *New England*

Journal of Medicine **377**, 1648–1656 (2017).

3. A. J. Lowe, E. David, C. J. Kilpatrick, Z. Matkovic, M. J. Cook, A. Kaye, T. J. O'Brien,

Epilepsy Surgery for Pathologically Proven Hippocampal Sclerosis Provides Long-term Seizure

Control and Improved Quality of Life. *Epilepsia* **45**, 237–242 (2004).

4. J. de Tisi, G. S. Bell, J. L. Peacock, A. W. McEvoy, W. F. J. Harkness, J. W. Sander, J. S.

Duncan, The long-term outcome of adult epilepsy surgery, patterns of seizure remission, and

relapse: a cohort study. *Lancet* **378**, 1388–1395 (2011).

5. T. Blauwblomme, E. Dossi, C. Pellegrino, E. Goubert, B. G. Iglesias, C. Sainte-Rose, N.

Rouach, R. Nabbout, G. Huberfeld, Gamma-aminobutyric acidergic transmission underlies

interictal epileptogenicity in pediatric focal cortical dysplasia. *Annals of Neurology* **85**, 204–217

(2019).

6. I. Cohen, V. Navarro, S. Clemenceau, M. Baulac, R. Miles, On the Origin of Interictal

Activity in Human Temporal Lobe Epilepsy in Vitro. *Science* **298**, 1418–1421 (2002).

7. G. Huberfeld, L. Wittner, S. Clemenceau, M. Baulac, K. Kaila, R. Miles, C. Rivera, Perturbed

Chloride Homeostasis and GABAergic Signaling in Human Temporal Lobe Epilepsy. *J.*

Neurosci. **27**, 9866–9873 (2007).

8. M. Munakata, M. Watanabe, T. Otsuki, H. Nakama, K. Arima, M. Itoh, J. Nabekura, K.

Iinuma, S. Tsuchiya, Altered distribution of KCC2 in cortical dysplasia in patients with

intractable epilepsy. *Epilepsia* **48**, 837–844 (2007).

9. A. Muñoz, P. Méndez, J. DeFelipe, F. J. Alvarez-Leefmans, Cation-Chloride Cotransporters

and GABA-ergic Innervation in the Human Epileptic Hippocampus. *Epilepsia* **48**, 663–673

(2007).

10. J. Pallud, M. Le Van Quyen, F. Bielle, C. Pellegrino, P. Varlet, N. Cresto, M. Baulac, C.

Duyckaerts, N. Kourdougli, G. Chazal, B. Devaux, C. Rivera, R. Miles, L. Capelle, G.

Huberfeld, Cortical GABAergic excitation contributes to epileptic activities around human

glioma. *Sci Transl Med* **6**, 244ra89-244ra89 (2014).

11. P. Blaesse, M. S. Airaksinen, C. Rivera, K. Kaila, Cation-Chloride Cotransporters and

Neuronal Function. *Neuron* **61**, 820–838 (2009).

12. H. R. Pathak, F. Weissinger, M. Terunuma, G. C. Carlson, F.-C. Hsu, S. J. Moss, D. A.

Coulter, Disrupted Dentate Granule Cell Chloride Regulation Enhances Synaptic Excitability

during Development of Temporal Lobe Epilepsy. *J Neurosci* **27**, 14012–14022 (2007).

13. M. R. Karlócai, L. Wittner, K. Tóth, Z. Maglóczky, Z. Katarova, G. Rásonyi, L. Erőss, S.

Czirják, P. Halász, G. Szabó, J. A. Payne, K. Kaila, T. F. Freund, Enhanced expression of

potassium-chloride cotransporter KCC2 in human temporal lobe epilepsy. *Brain Struct Funct*

221, 3601–3615 (2016).

14. M. Goutierre, S. A. Awabdh, F. Donneger, E. François, D. Gomez-Dominguez, T.

Irinopoulou, L. M. de la Prida, J. C. Poncer, KCC2 Regulates Neuronal Excitability and

Hippocampal Activity via Interaction with Task-3 Channels. *Cell Reports* **28**, 91-103.e7 (2019).

15. M. R. Kelley, R. A. Cardarelli, J. L. Smalley, T. A. Ollerhead, P. M. Andrew, N. J. Brandon,

T. Z. Deeb, S. J. Moss, Locally Reducing KCC2 Activity in the Hippocampus is Sufficient to

Induce Temporal Lobe Epilepsy. *EBioMedicine* **32**, 62–71 (2018).

16. G. Di Cristo, P. N. Awad, S. Hamidi, M. Avoli, KCC2, epileptiform synchronization, and

epileptic disorders. *Progress in Neurobiology* **162**, 1–16 (2018).

17. T. J. Ellender, J. V. Raimondo, A. Irkle, K. P. Lamsa, C. J. Akerman, Excitatory Effects of

Parvalbumin-Expressing Interneurons Maintain Hippocampal Epileptiform Activity via

Synchronous Afterdischarges. *J Neurosci* **34**, 15208–15222 (2014).

18. R. Khazipov, G. Valeeva, I. Khalilov, Depolarizing GABA and Developmental Epilepsies.

CNS Neurosci Ther **21**, 83–91 (2014).

19. Y. E. Moore, T. Z. Deeb, H. Chadchankar, N. J. Brandon, S. J. Moss, Potentiating KCC2

activity is sufficient to limit the onset and severity of seizures. *Proceedings of the National*

Academy of Sciences **115**, 10166–10171 (2018).

20. V. Magloire, J. Cornford, A. Lieb, D. M. Kullmann, I. Pavlov, KCC2 overexpression

prevents the paradoxical seizure-promoting action of somatic inhibition. *Nat Commun* **10**, 1225–

1237 (2019).

21. R. M. Pressler, G. B. Boylan, N. Marlow, M. Blennow, C. Chiron, J. H. Cross, L. S. de Vries,

B. Hallberg, L. Hellström-Westas, V. Jullien, V. Livingstone, B. Mangum, B. Murphy, D.

Murray, G. Pons, J. Rennie, R. Swarte, M. C. Toet, S. Vanhatalo, S. Zohar, Bumetanide for the

treatment of seizures in newborn babies with hypoxic ischaemic encephalopathy (NEMO): an

open-label, dose finding, and feasibility phase 1/2 trial. *The Lancet Neurology* **14**, 469–477

(2015).

22. J. S. Soul, A. M. Bergin, C. Stopp, B. Hayes, A. Singh, C. R. Fortuno, D. O'Reilly, K.

Krishnamoorthy, F. E. Jensen, V. Rofeberg, M. Dong, A. A. Vinks, D. Wypij, K. J. Staley, A

Pilot Randomized, Controlled, Double-Blind Trial of Bumetanide to Treat Neonatal Seizures.

Ann Neurol **89**, 327–340 (2021).

23. M. Gagnon, M. J. Bergeron, G. Lavertu, A. Castonguay, S. Tripathy, R. P. Bonin, J. Perez-

Sanchez, D. Boudreau, B. Wang, L. Dumas, I. Valade, K. Bachand, M. Jacob-Wagner, C. Tardif,

I. Kianicka, P. Isenring, G. Attardo, J. A. M. Coull, Y. De Koninck, Chloride extrusion

enhancers as novel therapeutics for neurological diseases. *Nat Med* **19**, 1524–1528 (2013).

24. S. Liabeuf, L. Stuhl-Gourmand, F. Gackière, R. Mancuso, I. Sanchez Brualla, P. Marino, F.

Brocard, L. Vinay, Prochlorperazine Increases KCC2 Function and Reduces Spasticity after

Spinal Cord Injury. *J Neurotrauma* **34**, 3397–3406 (2017).

25. A. J. Barbour, K. F. Hauser, A. R. McQuiston, P. E. Knapp, HIV and opiates dysregulate

K+-Cl– cotransporter 2 (KCC2) to cause GABAergic dysfunction in primary human neurons and

Tat-transgenic mice. *Neurobiol Dis* **141**, 104878 (2020).

26. J. Bilchak, K. Yeakle, G. Caron, D. Malloy, M.-P. Côté, Enhancing KCC2 activity decreases

hyperreflexia and spasticity after chronic spinal cord injury. *Exp Neurol* **338**, 113605 (2021).

27. Y. Gao, W. Zhan, Y. Jin, X. Chen, J. Cai, X. Zhou, X. Huang, Q. Zhao, W. Wang, J. Sun,

KCC2 receptor upregulation potentiates antinociceptive effect of GABAAR agonist on

remifentanil-induced hyperalgesia. *Mol Pain* **18**, 17448069221082880 (2022).

28. B. J. Sullivan, P. A. Kipnis, B. M. Carter, L.-R. Shao, S. D. Kadam, Targeting ischemia-

induced KCC2 hypofunction rescues refractory neonatal seizures and mitigates epileptogenesis

in a mouse model. *Sci Signal* **14**, eabg2648 (2021).

29. R. A. Cardarelli, K. Jones, L. I. Pisella, H. J. Wobst, L. J. McWilliams, P. M. Sharpe, M. P.

Burnham, D. J. Baker, I. Chudotvorova, J. Guyot, L. Silayeva, D. H. Morrow, N. Dekker, S.

Zicha, P. A. Davies, J. Holenz, M. E. Duggan, J. Dunlop, R. J. Mather, Q. Wang, I. Medina, N. J.

Brandon, T. Z. Deeb, S. J. Moss, The small molecule CLP257 does not modify activity of the

K+–Cl− co-transporter KCC2 but does potentiate GABAA receptor activity. *Nat Med* **23**, 1394–

1396 (2017).

30. M. Gagnon, M. J. Bergeron, J. Perez-Sanchez, I. Plasencia-Fernández, L.-E. Lorenzo, A. G.

Godin, A. Castonguay, R. P. Bonin, Y. De Koninck, Reply to The small molecule CLP257 does

not modify activity of the K+-Cl- co-transporter KCC2 but does potentiate GABAA receptor

activity. *Nat Med* **23**, 1396–1398 (2017).

31. S. Hamidi, M. Avoli, KCC2 function modulates in vitro ictogenesis. *Neurobiol Dis* **79**, 51–

58 (2015).

32. V. I. Dzhala, K. J. Staley, KCC2 Chloride Transport Contributes to the Termination of Ictal

Epileptiform Activity. *eNeuro* **8**, ENEURO.0208-20.2020 (2021).

33. G. Gauvain, I. Chamma, Q. Chevy, C. Cabezas, T. Irinopoulou, N. Bodrug, M. Carnaud, S.

Lévi, J. C. Poncer, The neuronal K-Cl cotransporter KCC2 influences postsynaptic AMPA

receptor content and lateral diffusion in dendritic spines. *Proc Natl Acad Sci U S A* **108**, 15474–

15479 (2011).

34. Y. Otsu, F. Donneger, E. J. Schwartz, J. C. Poncer, Cation–chloride cotransporters and the

polarity of GABA signalling in mouse hippocampal parvalbumin interneurons. *The Journal of*

Physiology **598**, 1865–1880 (2020).

35. S. Khirug, J. Yamada, R. Afzalov, J. Voipio, L. Khiroug, K. Kaila, GABAergic

Depolarization of the Axon Initial Segment in Cortical Principal Neurons Is Caused by the Na–

K–2Cl Cotransporter NKCC1. *J Neurosci* **28**, 4635–4639 (2008).

36. M. Chen, J. Wang, J. Jiang, X. Zheng, N. J. Justice, K. Wang, X. Ran, Y. Li, Q. Huo, J.

Zhang, H. Li, N. Lu, Y. Wang, H. Zheng, C. Long, L. Yang, E. Kim, Ed. APP modulates KCC2

expression and function in hippocampal GABAergic inhibition. *eLife* **6**, e20142 (2017).

37. M. Heubl, J. Zhang, J. C. Pressey, S. Al Awabdh, M. Renner, F. Gomez-Castro, I. Moutkine,

E. Eugène, M. Russeau, K. T. Kahle, J. C. Poncer, S. Lévi, GABAA receptor dependent synaptic

inhibition rapidly tunes KCC2 activity via the Cl−-sensitive WNK1 kinase. *Nat Commun* **8**, 1776

(2017).

38. M. A. Virtanen, P. Uvarov, M. Mavrovic, J. C. Poncer, K. Kaila, The Multifaceted Roles of

KCC2 in Cortical Development. *Trends in Neurosciences* **44**, 378–392 (2021).

39. E. Côme, M. Heubl, E. J. Schwartz, J. C. Poncer, S. Lévi, Reciprocal Regulation of KCC2

Trafficking and Synaptic Activity. *Frontiers in Cellular Neuroscience* **13,** 48 (2019)

40. I. Chamma, M. Heubl, Q. Chevy, M. Renner, I. Moutkine, E. Eugène, J. C. Poncer, S. Lévi,

Activity-Dependent Regulation of the K/Cl Transporter KCC2 Membrane Diffusion, Clustering,

and Function in Hippocampal Neurons. *J. Neurosci.* **33**, 15488–15503 (2013).

41. S. A. Awabdh, F. Donneger, M. Goutierre, M. Séveno, O. Vigy, P. Weinzettl, M. Russeau, I.

Moutkine, S. Lévi, P. Marin, J. C. Poncer, Gephyrin Interacts with the K-Cl Cotransporter KCC2

to Regulate Its Surface Expression and Function in Cortical Neurons. *J. Neurosci.* **42**, 166–182

(2022).

42. M. Watanabe, H. Wake, A. J. Moorhouse, J. Nabekura, Clustering of Neuronal K+-Cl–

Cotransporters in Lipid Rafts by Tyrosine Phosphorylation. *J Biol Chem* **284**, 27980–27988

(2009).

43. K. T. Kahle, A. R. Khanna, S. L. Alper, N. C. Adragna, P. K. Lauf, D. Sun, E. Delpire, K-Cl

cotransporters, cell volume homeostasis, and neurological disease. *Trends Mol Med* **21**, 513–523

(2015).

44. H. H. C. Lee, T. Z. Deeb, J. A. Walker, P. A. Davies, S. J. Moss, NMDA receptor activity

downregulates KCC2 resulting in depolarizing GABAA receptor mediated currents. *Nat*

Neurosci **14**, 736–743 (2011).

45. P. delos Heros, D. R. Alessi, R. Gourlay, D. G. Campbell, M. Deak, T. J. Macartney, K. T.

Kahle, J. Zhang, The WNK-regulated SPAK/OSR1 kinases directly phosphorylate and inhibit

the K+–Cl– co-transporters. *Biochem J* **458**, 559–573 (2014).

46. K. Yamada, H.-M. Park, D. F. Rigel, K. DiPetrillo, E. J. Whalen, A. Anisowicz, M. Beil, J.

Berstler, C. E. Brocklehurst, D. A. Burdick, S. L. Caplan, M. P. Capparelli, G. Chen, W. Chen,

B. Dale, L. Deng, F. Fu, N. Hamamatsu, K. Harasaki, T. Herr, P. Hoffmann, Q.-Y. Hu, W.-J.

Huang, N. Idamakanti, H. Imase, Y. Iwaki, M. Jain, J. Jeyaseelan, M. Kato, V. K. Kaushik, D.

Kohls, V. Kunjathoor, D. LaSala, J. Lee, J. Liu, Y. Luo, F. Ma, R. Mo, S. Mowbray, M. Mogi, F. Ossola, P. Pandey, S. J. Patel, S. Raghavan, B. Salem, Y. H. Shanado, G. M. Trakshel, G. Turner, H. Wakai, C. Wang, S. Weldon, J. B. Wielicki, X. Xie, L. Xu, Y. I. Yagi, K. Yasoshima, J. Yin, D. Yowe, J.-H. Zhang, G. Zheng, L. Monovich, Small-molecule WNK inhibition regulates cardiovascular and renal function. *Nat Chem Biol* **12**, 896–898 (2016).

47. H. H. C. Lee, R. Jurd, S. J. Moss, Tyrosine phosphorylation regulates the membrane trafficking of the potassium chloride co-transporter KCC2. *Mol Cell Neurosci* **45**, 173–179 (2010).

48. G. Huberfeld, L. Menendez de la Prida, J. Pallud, I. Cohen, M. Le Van Quyen, C. Adam, S. Clemenceau, M. Baulac, R. Miles, Glutamatergic pre-ictal discharges emerge at the transition to seizure in human epilepsy. *Nat Neurosci* **14**, 627–634 (2011).

49. C. Le Duigou, E. Savary, M. Morin-Brureau, D. Gomez-Dominguez, G. Milior, F. Chali, A. Sobczyk, E. Eugène, L. Kraus, J. C. Meier, D. M. Kullmann, B. Mathon, L. M. de la Prida, G. Dorfmuller, J. Pallud, S. Clemenceau, R. Miles, Imaging pathological activities of human brain tissue in organotypic culture. *J Neurosci Methods* **298**, 33–44 (2018).

50. I. Blümcke, M. Thom, E. Aronica, D. D. Armstrong, F. Bartolomei, A. Bernasconi, N. Bernasconi, C. G. Bien, F. Cendes, R. Coras, J. H. Cross, T. S. Jacques, P. Kahane, G. W. Mathern, H. Miyata, S. L. Moshé, B. Oz, Ç. Özkara, E. Perucca, S. Sisodiya, S. Wiebe, R. Spreafico, International consensus classification of hippocampal sclerosis in temporal lobe

epilepsy: A Task Force report from the ILAE Commission on Diagnostic Methods. *Epilepsia* **54**,

1315–1329 (2013).

51. G. Huberfeld, C. Le Duigou, M. Le Van Quyen, V. Navarro, M. Baulac, R. Miles, The

Paradox of the Paroxysm: Can Seizure Precipitants Help Explain Human Ictogenesis?

Neuroscientist **19**, 523–540 (2013).

52. G. Foffani, Y. G. Uzcategui, B. Gal, L. M. de la Prida, Reduced Spike-Timing Reliability

Correlates with the Emergence of Fast Ripples in the Rat Epileptic Hippocampus. *Neuron* **55**,

930–941 (2007).

53. L. Menendez de la Prida, A. J. Trevelyan, Cellular mechanisms of high frequency

oscillations in epilepsy: On the diverse sources of pathological activities. *Epilepsy Research* **97**,

308–317 (2011).

54. M. Zijlmans, P. Jiruska, R. Zelmann, F. S. S. Leijten, J. G. R. Jefferys, J. Gotman, High-

Frequency Oscillations as a New Biomarker in Epilepsy. *Ann Neurol* **71**, 169–178 (2012).

55. M. Lévesque, P. Salami, Z. Shiri, M. Avoli, Interictal oscillations and focal epileptic

disorders. *European Journal of Neuroscience* **48**, 2915–2927 (2018).

56. P. J. Karoly, V. R. Rao, N. M. Gregg, G. A. Worrell, C. Bernard, M. J. Cook, M. O. Baud,

Cycles in epilepsy. *Nat Rev Neurol* **17**, 267–284 (2021).

57. M. Baud, A. Ghestem, J.-J. Benoliel, C. Becker, C. Bernard, Endogenous multidien rhythm

of epilepsy in rats. *Experimental Neurology* **315**, 82 (2019).

58. R. J. Burman, R. E. Rosch, J. M. Wilmshurst, A. Sen, G. Ramantani, C. J. Akerman, J. V.

Raimondo, Why won't it stop? The dynamics of benzodiazepine resistance in status epilepticus.

Nat Rev Neurol **18**, 428–441 (2022).

59. L. J. Greenfield, Molecular Mechanisms of Antiseizure Drug Activity at GABAA Receptors.

Seizure **22**, 589–600 (2013).

60. A. Banerjee, R. V. Rikhye, V. Breton-Provencher, X. Tang, C. Li, K. Li, C. A. Runyan, Z.

Fu, R. Jaenisch, M. Sur, Jointly reduced inhibition and excitation underlies circuit-wide changes

in cortical processing in Rett syndrome. *Proceedings of the National Academy of Sciences* **113**,

E7287–E7296 (2016).

61. Z. Dargaei, J. Y. Bang, V. Mahadevan, C. S. Khademullah, S. Bedard, G. M. Parfitt, J. C.

Kim, M. A. Woodin, Restoring GABAergic inhibition rescues memory deficits in a Huntington's

disease mouse model. *Proc Natl Acad Sci U S A* **115**, E1618–E1626 (2018).

62. N. Kourdougli, C. Pellegrino, J.-M. Renko, S. Khirug, G. Chazal, T.-K. Kukko-Lukjanov, S.

E. Lauri, J.-L. Gaiarsa, L. Zhou, A. Peret, E. Castrén, R. K. Tuominen, V. Crépel, C. Rivera,

Depolarizing γ-aminobutyric acid contributes to glutamatergic network rewiring in epilepsy.

Annals of Neurology **81**, 251–265 (2017).

63. N. Lozovaya, R. Nardou, R. Tyzio, M. Chiesa, A. Pons-Bennaceur, S. Eftekhari, T.-T. Bui,

M. Billon-Grand, J. Rasero, P. Bonifazi, D. Guimond, J.-L. Gaiarsa, D. C. Ferrari, Y. Ben-Ari,

Early alterations in a mouse model of Rett syndrome: the GABA developmental shift is

abolished at birth. *Sci Rep* **9**, 9276 (2019).

64. W. Löscher, K. Kaila, CNS pharmacology of NKCC1 inhibitors. *Neuropharmacology* **205**,

108910 (2022).

65. A. Savardi, M. Borgogno, M. De Vivo, L. Cancedda, Pharmacological tools to target

NKCC1 in brain disorders. *Trends Pharmacol Sci* **42**, 1009–1034 (2021).

66. E. Delpire, K. B. Gagnon, in *Comprehensive Physiology*, (John Wiley & Sons, Ltd, 2018),

pp. 871–901.

67. J. S. Silvestre, J. Prous, Research on adverse drug events. I. Muscarinic M3 receptor binding

affinity could predict the risk of antipsychotics to induce type 2 diabetes. *Methods Find Exp Clin*

Pharmacol **27**, 289–304 (2005).

68. E. Richelson, A. Nelson, Antagonism by neuroleptics of neurotransmitter receptors of normal

human brain in vitro. *Eur J Pharmacol* **103**, 197–204 (1984).

69. H. Appl, T. Holzammer, S. Dove, E. Haen, A. Strasser, R. Seifert, Interactions of

recombinant human histamine H□R, H□R, H□R, and H□R receptors with 34 antidepressants

and antipsychotics. *Naunyn Schmiedebergs Arch Pharmacol* **385**, 145–170 (2012).

70. F. Ferrini, L.-E. Lorenzo, A. G. Godin, M. L. Quang, Y. De Koninck, Enhancing KCC2

function counteracts morphine-induced hyperalgesia. *Sci Rep* **7**, 3870 (2017).

71. A.-M. Hartmann, H. G. Nothwang, NKCC1 and KCC2: Structural insights into phospho-

regulation. *Front Mol Neurosci* **15**, 964488 (2022).

72. P. Friedel, K. T. Kahle, J. Zhang, N. Hertz, L. I. Pisella, E. Buhler, F. Schaller, J. Duan, A. R.

Khanna, P. N. Bishop, K. M. Shokat, I. Medina, WNK1-regulated inhibitory phosphorylation of

the KCC2 cotransporter maintains the depolarizing action of GABA in immature neurons. *Sci*

Signal **8**, ra65 (2015).

73. K. Strange, T. D. Singer, R. Morrison, E. Delpire, Dependence of KCC2 K-Cl cotransporter

activity on a conserved carboxy terminus tyrosine residue. *American Journal of Physiology-Cell*

Physiology **279**, C860–C867 (2000).

74. M. Wantanabe, J. Zhang, M. S. Mansuri, J. Duan, J. K. Karimy, E. Delpire, S. L. Alper, R. P.

Lifton, A. Fukuda, K. T. Kahle, Developmentally regulated KCC2 phosphorylation is essential

for dynamic GABA-mediated inhibition and survival. *Sci Signal* **12**, eaaw9315 (2019).

75. R. Jarvis, S. F. Josephine Ng, A. J. Nathanson, R. A. Cardarelli, K. Abiraman, F. Wade, A.

Evans-Strong, M. P. Fernandez-Campa, T. Z. Deeb, J. L. Smalley, T. Jamier, I. K. Gurrell, L.

McWilliams, A. Kawatkar, L. C. Conway, Q. Wang, R. W. Burli, N. J. Brandon, I. P. Chessell,

A. J. Goldman, J. L. Maguire, S. J. Moss, Direct activation of KCC2 arrests benzodiazepine

refractory status epilepticus and limits the subsequent neuronal injury in mice. *Cell Rep Med* **4**,

100957 (2023).

76. A. Cordshagen, W. Busch, M. Winklhofer, H. G. Nothwang, A.-M. Hartmann,

Phosphoregulation of the intracellular termini of K+-Cl− cotransporter 2 (KCC2) enables

flexible control of its activity. *J Biol Chem* **293**, 16984–16993 (2018).

77. J. L. Smalley, G. Kontou, C. Choi, Q. Ren, D. Albrecht, K. Abiraman, M. A. R. Santos, C. E.

Bope, T. Z. Deeb, P. A. Davies, N. J. Brandon, S. J. Moss, Isolation and Characterization of

Multi-Protein Complexes Enriched in the K-Cl Co-transporter 2 From Brain Plasma Membranes.

Frontiers in Molecular Neuroscience **13**, 563091 (2020).

78. M. Weber, A.-M. Hartmann, T. Beyer, A. Ripperger, H. G. Nothwang, A Novel Regulatory

Locus of Phosphorylation in the C Terminus of the Potassium Chloride Cotransporter KCC2

That Interferes with N-Ethylmaleimide or Staurosporine-mediated Activation. *J Biol Chem* **289**,

18668–18679 (2014).

79. V. Mahadevan, C. S. Khademullah, Z. Dargaei, J. Chevrier, P. Uvarov, J. Kwan, R. D.

Bagshaw, T. Pawson, A. Emili, Y. De Koninck, V. Anggono, M. Airaksinen, M. A. Woodin, M.

B. Kennedy, Ed. Native KCC2 interactome reveals PACSIN1 as a critical regulator of synaptic

inhibition. *eLife* **6**, e28270 (2017).

80. H. Asraf, M. Bogdanovic, N. Gottesman, I. Sekler, E. Aizenman, M. Hershfinkel, SNAP23

regulates KCC2 membrane insertion and activity following mZnR/GPR39 activation in

hippocampalneurons. *iScience* **25**, 103751 (2022).

81. M. Wenz, A.-M. Hartmann, E. Friauf, H. G. Nothwang, CIP1 is an activator of the K+-Cl-

cotransporter KCC2. *Biochem Biophys Res Commun* **381**, 388–392 (2009).

82. E. A. Ivakine, B. A. Acton, V. Mahadevan, J. Ormond, M. Tang, J. C. Pressey, M. Y. Huang,

D. Ng, E. Delpire, M. W. Salter, M. A. Woodin, R. R. McInnes, Neto2 is a KCC2 interacting

protein required for neuronal Cl– regulation in hippocampal neurons. *Proceedings of the*

National Academy of Sciences **110**, 3561–3566 (2013).

83. K. Inoue, J. Yamada, S. Ueno, A. Fukuda, Brain-type creatine kinase activates neuron-

specific K+-Cl- co-transporter KCC2. *J Neurochem* **96**, 598–608 (2006).

84. N. Garbarini, E. Delpire, The RCC1 domain of Protein Associated with Myc (PAM) interacts

with and regulates KCC2. *Cell Physiol Biochem* **22**, 31–44 (2008).

85. K. T. Kahle, J. Rinehart, P. de Los Heros, A. Louvi, P. Meade, N. Vazquez, S. C. Hebert, G.

Gamba, I. Gimenez, R. P. Lifton, WNK3 modulates transport of Cl- in and out of cells:

implications for control of cell volume and neuronal excitability. *Proc Natl Acad Sci U S A* **102**,

16783–16788 (2005).

86. J. S. Furyk, R. A. Meek, D. Egerton☐Warburton, Drugs for the treatment of nausea and

vomiting in adults in the emergency department setting. *Cochrane Database Syst Rev* **2015**,

CD010106 (2015).

87. M. Lau Moon Lin, P. D. Robinson, J. Flank, L. Sung, L. L. Dupuis, The Safety of

Prochlorperazine in Children: A Systematic Review and Meta-Analysis. *Drug Saf* **39**, 509–516

(2016).

88. C. Ghelardini, N. Galeotti, C. Uslenghi, I. Grazioli, A. Bartolini, Prochlorperazine induces

central antinociception mediated by the muscarinic system. *Pharmacological Research* **50**, 351–

358 (2004).

89. Y. Bozzi, E. Borrelli, The role of dopamine signaling in epileptogenesis. *Front Cell Neurosci*

7, 157 (2013).

90. X. Chi, X. Li, Y. Chen, Y. Zhang, Q. Su, Q. Zhou, Cryo-EM structures of the full-length

human KCC2 and KCC3 cation-chloride cotransporters. *Cell Res* **31**, 482–484 (2021).

91. Y. Xie, S. Chang, C. Zhao, F. Wang, S. Liu, J. Wang, E. Delpire, S. Ye, J. Guo, Structures

and an activation mechanism of human potassium-chloride cotransporters. *Science Advances* **6**,

eabc5883 (2020).

92. T. A. Chew, J. Zhang, L. Feng, High-Resolution Views and Transport Mechanisms of the

NKCC1 and KCC Transporters. *J Mol Biol* **433**, 167056 (2021).

93. S. Zhang, J. Zhou, Y. Zhang, T. Liu, P. Friedel, W. Zhuo, S. Somasekharan, K. Roy, L.

Zhang, Y. Liu, X. Meng, H. Deng, W. Zeng, G. Li, B. Forbush, M. Yang, The structural basis of

function and regulation of neuronal cotransporters NKCC1 and KCC2. *Commun Biol* **4**, 1–15

(2021).

94. N. Ankri, P. Legendre, D. S. Faber, H. Korn, Automatic detection of spontaneous synaptic

responses in central neurons. *J Neurosci Methods* **52**, 87–100 (1994).

Acknowledgments: We thank Liset M de la Prida for helpful discussions and critical reading of

the manuscript. We also thank Gilles Huberfeld for advices and assistance with handling and

recording human postoperative brain tissues, Sabine Lévi, Etienne Côme and Marianne Renner

for training and advices with single particle tracking (SPT) recordings and data analysis, Valérie

Crépel for advices on synaptoporin immunostaining, and the Imaging Facility of the Institut du

Fer à Moulin for assistance with confocal imaging.

Funding:

Fondation pour la Recherche Médicale DEQ20140329539 (JCP)

Fondation pour la Recherche Médicale FDT202106013286 (FD)

Agence Nationale de la Recherche ERANET-Neuron ANR-16-NEU3-0006-02 (JCP)

Fondation Française pour la Recherche sur l'Epilepsie (JCP)

Federation pour la Recherche sur le Cerveau (JCP)

Sorbonne Université (FD, JB, CP)

INSERM (JCP)

The Poncer lab is affiliated with the C-BRAINS network, supported by the Conseil Régional

d'Ile de France.

Author contributions:

Conceptualization: FD, JCP

Formal Analysis: FD, AZ, JB, YK, CP, ND

Funding acquisition: JCP

Investigation: FD, AZ, JB, CP

Methodology: FD, AZ, JCP

Resources: MR, FB, BD, BM, VN, FC

Software: FD, YK, ND

Supervision: JCP

Visualization: FD, JCP

Writing – original draft: FD, JCP

Writing – review & editing: FD, JCP

Conflict of interest statement: JCP, FD and ZA are co-inventors on the patent: Methods and
Pharmaceutical Compositions for Treating Refractory Epilepsy. WO2022219021. 2022 October.
All other authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary
materials.

Abbreviations:

CCCs   cation-chloride cotransporters

CNS   central nervous system

DMSO   dimethylsulfoxide

833	D2R	dopamine receptor D2
834	ECoG	electrocorticography
835	FDA	Food and Drug Administration
836	GABA	γ-Aminobutyric acid
837	HFOs	high-frequency oscillations
838	HS	hippocampal sclerosis
839	ILAE	International League Against Epilepsy
840	KCC2	K/Cl cotransporter 2
841	IID	interictal discharge
842	IILDs	interictal-like discharges
843	MEA	multi-electrode arrays
844	mIPSCs	miniature inhibitory postsynaptic currents
845	mTLE	mesial temporal lobe epilepsy
846	NKCC1	N/K/Cl cotransporter 1
847	OSR1	oxidative stress-responsive kinase 1
848	PCPZ	prochlorperazine
849	SE	status epilepticus
850	SPAK	SPS1-related proline/alanine-rich kinase
851	WNK	with-no-lysine kinase

852   **Tables and figures:**

Patient ID	Neuropathology	Sex	Age at surgery	Age at seizure onset (years)	Seizure frequency	SEEG-video	Engel score	Medication	IILD-generating area (*in vitro*)
1	Hippocampal sclerosis, ILAE type 1	F	43	childhood (precise date unknown)	monthly	no	IC	LCS, LEV, LTG	undefined
2	Hippocampal sclerosis, ILAE type 1	F	49	11	weekly	no	IA	LCS, LEV	undefined
3	Hippocampal sclerosis, ILAE type 1	F	36	23	monthly	no	IA	LEV, OXC, PRP	DG, Sub.
4	Unconfirmed diagnosis of hippocampal sclerosis. Atrophy visible on MRI + malformation of cortical development.	F	37	18	monthly	yes	IV	LTG, ESC, CLB, ESCIT	DG, undefined
5	Hippocampal sclerosis, ILAE type 1	F	63	3	monthly	no	IA	CBZ, GBP, PB	undefined
6	Hippocampal sclerosis, ILAE type 3	M	58	40	monthly	no	ID	CBZ, LCS, CLB, R	DG, Sub.
7	Hippocampal sclerosis, ILAE type 1	F	16	12	monthly	yes	III	LEV, LTG	Sub. only
8	Hippocampal sclerosis, ILAE type 2	H	41	8	monthly	no	IC	LCS, LTG	Sub., undefined
9	No neuronal loss, ganglioglioma with BRAF mutation	F	45	20	weekly	yes	IA	TPM, PRP, CLB	DG, Sub.
10	Hippocampal sclerosis, ILAE type 1	H	37	29	weekly	yes	IA	ZNS, CBZ, LTG	undefined
11	Hippocampal sclerosis, not classifiable	F	32	12	monthly	yes	IIIA	LTG, PRP, TAC, AZT, EPZ, NVL, PRED, MEL	DG, Sub.
12	Hippocampal sclerosis, ILAE type 1	F	22	12	weekly	no	ID	CBZ, LCS, LTG, PRP	DG only
13	Hippocampal sclerosis, ILAE type 2	H	54	12-13	weekly	yes	IA	CBZ, TPM, ZNS	DG only

Table 1. Clinical, neuropathological and pharmacological features of the patient

cohort. Abbreviations: AZT: Azathioprine; CBZ: Carbamazepine; CLB: Clobazam; ESCIT:

Escitalopram; ESC: Eslicarbazepine; EPZ: Esomeprazole; GBP: Gabapentine; LCS:

Lacosamide; LTG: Lamotrigine; LEV: Levetiracetam; MEL: Melatonine; NVL: Nebivolol;

OXC: Oxcarbazepine; PRP: Perampanel; PB: Phenobarbital; PRED: Prednisone; R:

Ropinirole; TAC: Tacrolimus; TPM: Topimarate; ZNS: Zonesamide. DG: Dentate gyrus.

Sub.: subiculum. Grey background indicates patients for which no significant effect of PCPZ

or CLP257 on interictal-like activity was observed *in vitro*.

Fig. 1. Prochlorperazine (PCPZ) and CLP-257 promote chloride extrusion in hippocampal neurons. (A) Representative currents evoked at varying potentials by focal application of isoguvacine (arrow, 100 µM) onto the soma or dendrite of hippocampal neurons (21-24 DIV) previously treated for 2 hours with either vehicle only (DMSO, control) or PCPZ (10 µM). **(B)** I-V curves corresponding to the traces shown in A. **(C)** 2D chemical structure of PCPZ (top) and summary graph showing somato-dendritic E_{GABA} gradient (bottom, ΔE_{GABA}) in control and PCPZ-treated neurons (n=19 neurons and n=16 neurons respectively, from 5 independent cultures). ** t-test p<0.01. **(D-E)** same as in A-B, with currents evoked at varying potentials by focal uncaging of Rubi-GABA (arrow, 30 µM) onto the soma or dendrite of hippocampal

873 neurons, pretreated for 2 hours with either DMSO alone or 100 μM CLP-257 in DMSO. (**F**) 2D

874 chemical structure of CLP-257 (top) and summary graph showing ΔE_{GABA} (bottom) in control

875 and CLP-treated neurons (n=10 and 8 neurons, respectively, from 4 independent cultures) * t-test

876 p<0.05.

Fig. 2. CLP-257 but not PCPZ modulates GABA$_A$ receptor function. (A) Left, representative

recording of currents evoked by focal somatic application of isoguvacine before and during acute

application of 10 µM PCPZ. Evoked currents were averaged and their decay was fit with a

simple exponential (red). Right, normalized amplitude and decay time constant over time of

evoked currents. (**B**) Summary graphs showing the effect of PCPZ on the peak amplitude (left)

and decay time constant (τ) of isoguvacine-evoked currents (n=10 neurons each from 2

independent cultures). (**C-D**) Same as in A-B for currents evoked by Rubi-GABA (30 µM)

uncaging before and during application of 100 µM CLP-257 (n=11 neurons each from 2

independent cultures. *** Wilcoxon test, p<0.001). (**E**) Left, representative 5-seconds mIPSC

recording before (DMSO) and during application of PCPZ. Right, superimposed averaged

mIPSCs detected from this recording. (**F**) Summary graphs showing the effect of PCPZ

application on the amplitude (left) and decay time constant (right) of mIPSCs (n=18 neurons

each from 4 independent cultures. *** Wilcoxon test, p<0.001). (**G-H**) same as E-F for CLP-257

showing no effect of CLP-257 on mIPSC amplitude and decay time constant (n=12 neurons each

from 4 independent cultures. n.s.: not significant).

Fig. 3. Both PCPZ and CLP-257 reduce KCC2 membrane diffusion and promote its clustering in hippocampal neurons. (A) Left, representative immunoblots of protein extracts from primary hippocampal cultures treated for 2 hours with either PCPZ (10 μM), CLP-257 (100 μM) or DMSO alone. Right, quantification from 3 independent experiments. No significant change in total KCC2 expression was detected upon PCPZ or CLP-257 treatment as compared to control (t-test p=0.14 and p=0.89, respectively). **(B)** Representative immunoblots (left) and quantification (right) showing biotinylated surface KCC2 fraction (bound/total) was also unaffected upon PCPZ or CLP-257 treatment (n=3 independent cultures, t-test p=0.74 and 0.80, respectively). **(C)** Representative trajectories of recombinant KCC2 (black) outside inhibitory

synapses (gephyrin-FingR mask, grey), as detected using quantum dot-based single particle

tracking, showing reduced lateral diffusion upon treatment with PCPZ or CLP-257. Scale, 1 µm.

(D) Logarithmic distributions of median diffusion coefficients (D) and explored areas (EA) of

KCC2 trajectories, outside (gephyrin -) or in the vicinity of (gephyrin +) inhibitory synapses, in

control neurons (DMSO, D extra n=614, D syn n=162 and EA extra n=1836, EA syn n=486

quantum dots) and neurons treated for 2 hours with either PCPZ (D extra n=512, D syn=131 and

EA extra n=1536, EA syn n=393 quantum dots) or CLP-257 (D extra n=697, D syn n=179 and

EA extra n=2091, EA syn n=537 quantum dots). 4 independent cultures. Mann-Whitney test *

$p<0.05$ *** $p<0.001$. **(E)** Maximum projections of confocal images of proximal dendrites after

KCC2 immunostaining in cultured hippocampal neurons treated for 2 hours with PCPZ, CLP-

257 or DMSO only. Scale, 5 µm. **(F)** Boxplots showing the distributions of integrated intensity,

area and density of KCC2 clusters in control neurons and neurons exposed to either PCPZ or

CLP-257. n=10 neurons per condition per culture and 4 independent cultures per condition. t-test

or Mann-Whitney test * $p<0.05$ *** $p<0.001$.

Fig. 4. PCPZ and CLP-257 do not affect phosphorylation of canonical KCC2 residues. (A)

Immunoblots (IB) (left) of protein lysates or proteins immunoprecipitated (IP) with phosho-

specific KCC2 antibodies, immunoblotted with antibodies against KCC2 or pS940 KCC2 and

beta-tubulin (Tuj1) as control, prepared from hippocampo-cortical slices treated for 2 hours with

either DMSO alone, CLP-257 (100 µM in DMSO), PCPZ (10 µM in DMSO), or the pan-WNK

inhibitor WNK463 (10 µM in DMSO). IgG: control immunoglobulin G. (n=3 independent

replicates 1-3). **(B)** Quantification from the 3 independent replicates shown in A. t-test

***p<0.001 **(C)** IB (left) of protein lysates, prepared as in A, immunoblotted with antibodies

against SPAK, OSR1, or their phosphorylated forms and beta-tubulin (Tuj1) as control. **(D)**

Quantification from the 3 independent replicates in B. t-test *** p<0.001. **(E)** Representative IB

(left) of proteins immunoprecipitated with an anti-KCC2 antibody and immunoblotted with an

anti-phosphotyrosine antibody, prepared from hippocampal-cortices slices treated for 2 hours

with either DMSO, CLP-257 (100 µM in DMSO), PCPZ (10 µM in DMSO), or sodium

orthovanadate (Na$_3$VO$_4$, 200 µM). **(F)** Quantification from the 4 independent replicates shown in

E. Mann-Whitney test * p<0.05.

Fig. 5. PCPZ and CLP-257 prevent spontaneous interictal-like discharges in temporal lobe

slices from mTLE patients. (A) (Left) Photomicrograph of a recorded slice of resected brain

tissue from a mTLE patient, showing multielectrode array (MEA) layout. Hippocampal and

parahippocampal regions, including the subiculum, were delineated based on slice morphology.

Electrodes were color-coded to indicate the type of activity detected (red: interictal-like

discharges (IILDs) green: no activity). Scale: 1 mm. (Right) Representative recording from this

slice, showing synchronous IILDs mostly within the dentate gyrus. **(B)** Representative local field

potential recordings and spectrograms showing spontaneous, low-frequency (<40Hz) IILDs in a

slice maintained in control conditions (2 hours in DMSO only, top), and their reduction upon

943 >10 min application of 10 µM bumetanide (middle). These events were completely absent from

944 an adjacent slice from the same tissue incubated for 2 hours in ACSF supplemented with 10 µM

945 PCPZ (bottom). Insets show individual IILDs. (**C**) Representative recordings showing

946 spontaneous IILDs in a slice from another patient incubated in control conditions (top), and the

947 absence of these events in an adjacent slice kept for 2 hours in ACSF supplemented with 100 µM

948 CLP-257 (bottom). (**D**) Summary plot showing the proportion of slices with and without IILDs.

949 χ^2 test, * p<0.05. (**E-F**) Slices from two other patients treated with either CLP or PCPZ also did

950 not display IILDs. However, application of a pro-convulsant ACSF (6.5 mM K^+ and 0.25 Mg^{2+})

951 rescued spontaneous IILDs in these slices. (**G**) Graphs showing the mean frequency of

952 spontaneous IILDs (normalized to control) recorded in slices from 7 mTLE patients incubated in

953 control ACSF (DMSO) or in the presence of 10 µM PCPZ. Two groups were distinguished:

954 responders (Group I, 5 patients) and non-responders (Group II, 2 patients). Similar data are

955 shown for slices from 6 patients, incubated in control ACSF or treated with 100 µM CLP-257.

956 Wilcoxon signed rank test, *** p<0.001.

Fig. 6. Chronic PCPZ administration reduces the occurrence of seizures and epileptiform activities in a mTLE mouse model. (A) Experimental timeline. (**B**) Graph showing the cumulative number of seizures detected in a mouse immediately after SE and up to 37 days post SE. Recording was interrupted between days 21 and 34 post SE. Note that seizures stop 2 days after SE and chronic seizures occur at a regular rate after day 34 (inset). (**C**) Top, representative confocal fluorescent micrographs of DAPI staining of hippocampal slices from naïve (no SE) and epileptic mice. Arrowheads indicate the CA1 region, which is largely sclerotic in the epileptic mouse. Also note the dispersion of the dentate gyrus granule cells. Scale: 400 µm. Bottom, confocal fluorescent micrographs of KCC2 immunolabelling in dentate gyrus, from the regions boxed in top panels, showing reduced KCC2 immunofluorescence in the dentate

gyrus of the epileptic mouse. Scale: 60 µm (**D)** Top, representative confocal fluorescent

micrographs of synaptoporin immunolabelling of the dentate gyrus of control and epileptic mice.

Scale: 200 µm. Bottom, magnification of the regions boxed in top panels, showing ectopic

synaptoporin-immunopositive mossy fibers in the stratum granulosum and stratum moleculare

only in the epileptic mouse. Scale: 20 µm. (**E)** Left, representative ECoG recording of a seizure

with insets (a,b,c) showing typical periods of the seizure. Middle, summary graphs showing

mean seizure frequency per hour in sham (no SE, n=6) and epileptic mice (n=17). No seizures

were detected in sham animals (*** Mann-Whitney test p<0.001). Right, summary graphs

showing mean seizure frequency per hour in epileptic mice receiving saline i.p. injections during

the control period (days 35-39) and either saline or PCPZ during the test period (days 40-44).

Mice receiving PCPZ injections during the test period displayed significantly less seizures than

during the control period (n=9 mice, * paired t-test p<0.05), while no change in seizure

frequency was observed in mice receiving saline injections (n=8, p=0.118). (**F**) Summary graphs

comparing mean seizure power (left) and duration (right) during the control (saline) and test

(PCPZ) periods. While reducing seizure frequency, PCPZ had no significant effect on seizure

power (paired t-test p=0.466) or duration (p=0.657, n=8 mice). (**G-H)** Left, representative ECoG

recording of an interictal discharge (IID, in G) and a high-frequency oscillation (HFO, in H).

Middle, summary plots showing mean IID (in G) and HFO (in H) frequency in sham (n=6) and

epileptic animals (n=17). IIDs and HFOs were significantly more frequent in epileptic animals

(Mann-Whitney test * p<0.05, *** p<0.001, respectively). Right, summary plots of mean IID (in

G) and HFO (in H) frequency during the control period (days 34-39) and the test period (days

40-44). Recordings from mice receiving PCPZ injections during the test period displayed

significantly reduced IID and HFO frequency as compared to the control period (* paired t-test

$p < 0.05$, n=9 mice). Recordings from mice receiving saline injections during the test period

revealed no change in IID frequency (p=0.846) and a significant increase in HFO frequency (*

$p < 0.05$, n=8 mice).