Conformational dynamics of a nicotinic receptor neurotransmitter binding site

Mrityunjay Singh[1], Dinesh C. Indurthi[2][†], Lovika Mittal[1], Anthony Auerbach[2][†], and Shailendra Asthana[1][†].

[1]Computational Biophysics and CADD Group, Computational and Mathematical Biology Center,

Translational Health Science and Technology Institute, NCR Biotech Science Cluster, Faridabad 121001,

India.

[2]Department of Physiology and Biophysics, University at Buffalo, State University of New York, Buffalo,

United States.

[†] Corresponding Authors

Shailendra Asthana: sasthana@thsti.res.in

Anthony Auerbach: Anthony.auerbach@gmail.com

Dinesh C. Indurthi: dinesh.indurthi@gmail.com

Key words: acetylcholine, allostery, agonist, molecular dynamics, ion channel

Abstract

Agonists turn on receptors because they provide net favorable binding energy to active versus resting conformations of their target sites. We used simulations to explore conformational dynamics of the weak→strong binding transition at the *Torpedo* α–δ nicotinic acetylcholine receptor orthosteric site. Using 4 agonists, the alternative site conformations were identified in trajectories generated from a single starting structure by matching binding energies calculated *in silico* with those measured experimentally *in vitro*. The weak→strong transition starts with a rotation of the agonist about its cationic center ('flip'), followed by a downward displacement of loop C that repositions αY190 ('flop'), followed by formation of H-bonds between the ligand, a structural water and the δ subunit loop E backbone ('fix'). The result is a compact, hydrophobic and stable pocket with higher affinity for agonists. The simulations reveal a transient intermediate state in the weak→strong transition.

Introduction

Adult vertebrate skeletal muscle nicotinic acetylcholine receptors (AChRs) are allosteric proteins that mediate synaptic transmission between motor neurons and muscle fibers. These pentameric ligand-gated ion channels are composed of four different subunits (2 α1, β1, δ, and ε) arranged symmetrically around a central ion-conducting pore (Rahman et al., 2020; Unwin, 1993). Binding of the neurotransmitter acetylcholine (ACh) or other agonists to two orthosteric sites in the extracellular domain (ECD) increases the probability of a global change in protein 3D conformation that activates a distant (~60.0 Å) allosteric site (a gate) in the transmembrane domain (TMD), allowing cations to cross the membrane.

At the start of the channel-opening isomerization the neurotransmitter sites switch from having a low to a high affinity for agonists (LA→HA), and at the end the gate switches from closed to open (C→O) (Grosman et al., 2000; Purohit et al., 2013). Hence, the reaction scheme for the 'gating' allosteric transition is $C_{LA} \rightleftharpoons O_{HA}$. Without any bound agonists the probability of having the O_{HA} shape (P_O) is ~10^{-6} and baseline current is negligible (Jackson, 1986; Nayak et al., 2012; Purohit & Auerbach, 2009). With or without agonists, the receptor turns on and off spontaneously (influenced only by temperature). Agonists increase P_O (that is, preferentially stabilize O_{HA}) because they provide net favorable binding energy to their target when it is in the active, high-affinity conformation. Agonists also stabilize the transition state that separates C_{LA} from O_{HA} and therefore increase P_O almost exclusively by increasing the channel-opening rate constant.

The difference in binding free energies, HA minus LA ($\Delta G_{HA}-\Delta G_{LA}$), drives the receptor's allosteric transition and determines relative agonist efficacy (maximum P_O). Another ligand attribute we use is efficiency that is a function of the ratio of binding free energies, ($1-\Delta G_{LA}/\Delta G_{HA}$) (Nayak et al.,

2019). Efficacy (efficiency) is the amount (fraction) of binding free energy delivered to the gating

apparatus. In a dose-response curve, efficacy is manifest as the high-concentration asymptote and

efficiency as the correlation between this maximum and EC_{50}.

Adult-type muscle AChRs have 2 approximately equal and independent neurotransmitter sites

(Nayak & Auerbach, 2017). The net LA→HA binding free energy change (ΔG) for two neurotransmitters

(-10.2 kcal/mol) adds to that for the unfavorable unliganded isomerization (+8.3 kcal/mol at -100 mV)

(Nayak et al., 2012) making the free energy change of the allosteric transition with 2 bound ACh

favorable (-1.9 kcal/mol). ΔG is proportional to the logarithm of the corresponding equilibrium

constant (by -RT, where R is the gas constant and T the absolute temperature, equal to -0.59 at 23 ^{o}C).

Accordingly, after 2 neurotransmitters have bound, LA→HA rearrangements at the 2 orthosteric sites

increases P_O from $\sim 10^{-6}$ to ~0.96 (Figure 1A).

The relationship between ΔG_{LA} and ΔG_{HA} binding free energies and the agonist properties

affinity, efficacy and efficiency are considered elsewhere (Auerbach, 2024). Here, our goal was to

associate the $\Delta G_{LA} \rightarrow \Delta G_{HA}$ change in binding free energy (4 agonists) with restructuring events at a

neurotransmitter site. Molecular dynamics (MD) simulations were used to explore conformational

changes in the LA→HA transition at the $\alpha - \delta$ neurotransmitter site of the *Torpedo* AChR (6UVW;

(Rahman et al., 2020)). In brief, binding free energies calculated *in silico* matched those estimated by

using electrophysiology *in vitro* (Indurthi & Auerbach, 2023), allowing the identification of LA versus HA

structures in the simulations.   In the LA→HA transition for all agonists, three prominent

rearrangements were rotation (flip) of the ligand, downward displacement (flop) of loop C, and

compaction and stabilization of the binding cavity (fix). The initial LA and final HA conformations of the

site were connected by a brief intermediate state. The simulations suggest that the movement of a

weakly-bound agonist starts the change to strong binding and initiates the protein's allosteric

transition.

Results

Hold. Traditionally, receptor activation is divided into distinct stages called 'binding' (formation of a

ligand-protein complex) and 'gating' (receptor isomerization) (Figure 1). Here, we are concerned with

the connection between these stages as molecular movements at a neurotransmitter site.

In AChRs, agonist binding is a 2-step process. First, the agonist (A) diffuses to the target and

forms an encounter complex (A+C$\rightleftarrows$A-C), after which a local 'catch' rearrangement establishes the LA

complex (A-C$\rightleftarrows$AC$_{LA}$) (Jadey & Auerbach, 2012). Gating, too, involves multiple steps (Gupta et al., 2017;

Purohit et al., 2013). First, the site undergoes a local 'hold' rearrangement that establishes the HA

complex (AC$_{LA}\rightleftarrows$AC$_{HA}$), after which other protein domains restructure without changing agonist affinity,

eventually opening the gate (Figure 1B).

The only agonist-dependent free energy changes in activation occur in the two local site

rearrangements, ΔG_{LA} (in catch) and $\Delta G_{HA} - \Delta G_{LA}$ (in hold). The terminal state in the 'binding' stage of

activation is AC$_{LA}$ and the initial state in the 'gating' stage is AC$_{HA}$. Our interest here is hold, AC$_{LA}\rightleftarrows$AC$_{HA}$,

the local rearrangement that connects these two states. The free energy changes associated with the

other events in the activation sequence, namely the formation of the encounter complex and

downstream domain restructurings, are approximately the same for all agonists and so are not

relevant to this study.

Activation free energy changes incurred outside catch and hold, namely from the chemical

potential associated with encounter complex formation (Phillips, 2020) and from rearrangements in

other domains (and water), together represent an energy offset that is approximately agonist

independent. With a few exceptions, energy coupling between the agonist binding pocket and distant

residues (>15.0 Å) typically is weak (Gupta et al., 2017), so only the agonist's energy changes in catch

and hold determine the pharmacology attributes of affinity (ΔG_{LA} or ΔG_{HA}), efficacy ($\Delta G_{HA} - \Delta G_{LA}$) and

efficiency ($1-\Delta G_{LA}/\Delta G_{HA}$). In AChRs, ΔG_{LA} and ΔG_{HA} have been measured experimentally by using

electrophysiology for 23 agonists and 53 neurotransmitter site mutations (Indurthi & Auerbach, 2023).

Apo, resting-C with a bound toxin, and high-affinity desensitized AChR structures have been

solved by using cryo-EM (Rahman et al., 2022; Rahman et al., 2020; Zarkadas et al., 2022). Desensitized

AChRs have a similar (perhaps identical) high affinity as AO_{HA} (Auerbach, 2020; Nayak & Auerbach,

2017). Below, we describe the conformation dynamics that connect the end states of the hold

rearrangement, $AC_{LA} \rightarrow AC_{HA}$ (Figure 1B, bottom), neither of which has been captured as a structure.

However, as described below, they can be identified in MD simulations that use resting-C as a start.

Figure 1: AChR activation. *A.* Cyclic, thermalized activation scheme (vertical, gating; horizontal, binding; red, main physiological pathway). Receptors isomerize globally between closed-channel/low-affinity (C_{LA}) and open-channel/high-affinity (O_{HA}) conformations (equilibrium constant L_n; n, number of bound ligands). Agonists (A) bind weakly to C_{LA} (equilibrium association constant K_{LA}, free energy change ΔG_{LA}) and strongly to O_{HA} (K_{HA}, ΔG_{HA}). In adult AChRs the 2 orthosteric sites are approximately equivalent and independent, and there is no significant external energy: $L_2/L_0 = (K_{HA}/K_{LA})^2$ (Nayak & Auerbach, 2017). B. Top, binding and gating. Bottom, catch and hold. The agonist diffuses to the target to form an encounter complex (A-C), a local 'catch' rearrangement establishes the LA complex (AC_{LA}), a local 'hold' rearrangement establishes the HA complex (AC_{HA}), the rest of the protein isomerizes without a further change in affinity to generate a conducting pore (AO). Gray, steps that incur the same energy change for all agonists used in this study; black, agonist-dependent free energy changes in catch is ΔG_{LA} and in hold (ΔG_{HA}-ΔG_{LA}). C. α-δ subunit extracellular domains, red, after toxin removal (6UWZ.pdb) and blue, apo (7QKO.pdb). There are no major deviations (C_α RMSD = 0.3 Å). D. Closeup of the desensitized *Torpedo* α-δ subunit neurotransmitter site occupied by carbamylcholine (CCh, blue) (7QL6.pdb; (Zarkadas et al., 2022)). In this is HA conformation, 3 aromatic groups in the α subunit (149-190-198) surround the agonist's cationic center (+) together provide ~90% of the binding energy for ACh (Purohit et al., 2014), and the tail points toward the δ subunit (*trans* orientation).

Docking. We began by removing α-bungarotoxin from the high-resolution (2.69 Å) structure 6UWZ

(Rahman et al., 2020) (Figure 1C) and docking agonists into the empty pocket. The docking simulations

had strong binding scores ranging from -5.82 to -9.18 kcal/mol (Figure 2A), suggesting that the

interactions were stable. However, unlike the agonist orientation apparent in cryo-EM structures of

desensitized (HA) AChRs (Noviello et al., 2021; Rahman et al., 2022; Rahman et al., 2020; Walsh et al.,

2018; Zarkadas et al., 2022) (Figure 1D), the docked agonist's tail pointed towards (_cis_) rather than

away (t_rans_) from the α subunit (Figure 2B, top). This was the case for the top 3 docking positions for

all 4 ligands.

Because the _cis_ orientation was not observed in any cryo-EM structure, we examined the top

200 docking scores for CCh. All poses were _cis_, with the tail pointing towards the α subunit and away

from the main binding cavity. Apparently, in the apo structure the _trans_ orientation is energetically

unfavorable. Hence, to start our analysis of the hold rearrangement we hypothesized that the agonist

orientations in the end states are _cis_ in AC_{LA} versus _trans_ in AC_{HA}.

Figure 2. Agonist docking and loop dynamics. A. Top, agonists (blue, cationic center): carbamylcholine (CCh), acetylcholine (ACh), epibatidine (Ebt) and epiboxidine (Ebx). Bottom, $\alpha-\delta$ site (6UVW.pdb minus toxin, resting-C) with docked agonists (top 3 poses). docked into resting-C (red): loop C is up, agonist is *cis*. docked after 200 ns simulation and removing CCh (blue): loop C is down, agonist is *trans*. B. Bottom, docking scores (mean±SD, n=3). red, resting-C; blue, 200 ns. C. C_α RMSD (mean±SD, triplicates) (ACh, cyan; CCh, green; Ebt, orange; Ebx, purple). D. close-up of the CCh-occupied pocket. Red, resting-C; orange, equilibrated (0 ns); blue, 200 ns. After 200 ns loop C has flopped, loop F has moved inward, and the agonist has flipped *cis→trans*.

The LA→HA transition. We performed 200 ns MD simulations for each of the docked complexes, in

triplicate (with different seed values). Video 1 shows a simulation with CCh on 2 times scales,

expanded for the initial 200 ns and condensed for the full 1.0 μs extension. The 200 ns simulations for

all three replicates for all agonists showed similar root-mean-square-deviation (RMSD) patterns with

differences consistently below 1Å (Figure 2C, Figure 2-Source Data 1) suggesting that the trajectories

were sufficiently robust and reproducible. Although there were initial deviations, all systems became

stable by ~120 ns. The root-mean-square-fluctuation (RMSF) of the C_α atoms showed flexibility around

the pocket mainly in loops C and F (but not E) (Figure 2-figure supplement 1). The fluctuations at the

base of the ECD were anticipated because the TMD that offers stability here was absent. As described

below, the match between experimental binding energies (whole receptors) and those calculated in

the simulations (ECD dimers) is further evidence that the ECD-TMD interfacial region does not

influence agonist binding energy.

The ECD dimer in the absence of a bound agonist (apo) exhibited a similar RMSF and RMSD

pattern for a 200 ns simulation, to that observed with ligand present (CCh) (Figure 2-figure supplement

2A, Figure 2-figure supplement 2B). The displacement of loop C apparent with agonists (see below) was

also present but less pronounced in apo (Figure 2-figure supplement 2C).

In addition to the fluctuations in loops C and F during the simulations, the ligand orientation

inverted, *cis→trans*. For all ligands, at the start of the simulations the agonist's tail points towards α,

but by the end points towards δ (Figure 2D). The agonist re-orientation represents a nearly complete

somersault (a 'flip') about the agonist's main cationic center (N^+) that remains close to αW149

throughout (see Figure 5). The final MD configuration (with CCh) aligns well with the CCh-bound cryo-

EM desensitized structure (7QL6.PDB) (RMSD < 0.5 Å) (Figure 2-figure supplement 3). Video 1 shows

that the final *trans* orientation with CCh remains constant for at least 1 μs, indicating that structure

had settled by 200 ns, with critical rearrangements conserved.

To test whether rearrangements at the binding pocket in the simulations enforced the *trans*

configuration, we removed the bound CCh from the final 200 ns MD structure and re-docked the

agonists. The preferred poses for all 4 ligands were *trans* (Figure 2A, bottom). In addition, binding

energies from docking scores were higher for the 200 ns structures compared to the initial binding site

structures, consistent with a LA-to-HA transition (Figure 2B). As a further control we carried out MD

simulations of a pentamer docked with ACh and found similar structural changes at the binding pocket

compared to the dimer (Figure 2-figure supplement 4).

Together, these results suggest that ECD dimers are adequate to study local rearrangements of

the $\alpha-\delta$ AChR neurotransmitter binding site. Further, they provide support for the hypotheses that in

the simulations the site restructures from AC_{LA} to AC_{HA}, and that the preferred agonist pose is *cis* in LA

versus *trans* in HA structures.

Principal component analysis (PCA). To identify the most prominent conformational states of the

receptor during the simulations, PCA was carried out based on the most significant fluctuations

(principal components). Considering the first two principal components (PC-1 and PC-2) that captured

the most pronounced C_α displacements (Figure 3-Source Data 1), backbone fluctuations revealed three

energy minima (m1, m2, and m3) that reflect the stabilities of loops C and F (Figure 3). To assess

consistency across independent simulations, we calculated the inner products of the top 10 principal

components from each run. The resulting heatmaps reveal the degree of similarity between runs, with

the inner product values of PC-1 and PC-2 reflecting considerable overlap (50-90%) in the principal

dynamic modes captured in each simulation (Figure 3-figure supplement 1).

For all 4 agonists, trajectories started in m1 and ended in m3 (Figure 3-figure supplement 2).

Accordingly, we hypothesized that m1 corresponds to AC_{LA} and that m3 corresponds to AC_{HA}. For all

agonists, the m2 population had a greater variance (was less stable) compared to m1 and m3. This,

along with its temporal position, suggests that m2 represents an unstable intermediate configuration

within the hold transition, between AC_{LA} and AC_{HA}. For all agonists and trajectories, m3 had the least

variance (was the most stable).

The pronounced fluctuations in loop C are evident in residue-wise RMSF plots and align well

with the contributions of the first two principal components (PC-1 and PC-2) (Figure 2-figure

supplement 2D). The apo and CCh occupied proteins adopt similar conformational spaces during the

200 ns simulation (Figure 2-figure supplement 2E). However, presence of the agonist appears to

increase conformational diversity, generating a broader distribution in the PCA plot. The bottom of the

m1, m2 and m3 energy wells in Figure 3 represent the most stable configurations of each population

and serve as reference points.

202

Figure 3. Principal Component Analysis (PCA). For each agonist, the left panel plots PC1 vs PC2, the first two principal components that capture the maximum variance in the trajectory (Figure 3-Source Data 1). Colors represent free energy value in kcal/mol (scale, upper left-bottom). For all agonists there are 3 energy minima (dark red), m1, m2, and m3, that correspond to different conformations of the protein. The right panels are 'porcupine' plots indicating that the direction and magnitude of changes between PC1 and PC2 is in loops C and F. From energy comparisons (Figure 4, Figure 4-Source Data 1) and temporal sequences (Figure 3-figure supplement 2, Figure 4-Source Data 1): m1 is AC_{LA}, m3 is AC_{HA}, and m2 is an intermediate configuration.

Computed versus experimental binding energies. Although the well bottoms in Figure 3 represent the

most stable overall conformations, they do not directly convey information regarding agonist stability

or orientation. To incorporate this information, we performed a cluster analysis of the ligand

configuration using frames from the bottom of each PCA well as input. The top three clusters, each

having RMSD of ≤ 1.0 Å, shared a similar ligand orientation (Figure 3-figure supplement 2) and were

selected to compute binding free energies (Figure 4, Figure 4-Source Data 1, Figure 4-Source Data 2).

The remaining frames were disregarded. The fraction of frames from each minimum accepted for free

energy calculations ranged from 20% (ACh, m1) to 71% (Ebt, m3). On average, the fraction of frames

selected was ~50%, showing the dynamic nature of the ligand in the pocket (Video 1).

The above procedure defined populations of structures that we hypothesized represent the end

states of hold, AC_{LA} (m1) and AC_{HA} (m3), plus a previously unidentified intermediate state (m2). We

used PBSA (Poisson-Boltzmann Surface Area) computations of the selected structures for each agonist

in each cluster population and compared binding free energies calculated *in silico* with those measured

previously *in vitro* (Indurthi & Auerbach, 2023). There are limitations with regard to PBSA accurately

predicting absolute binding free energies (Genheden & Ryde, 2015; Hou et al., 2011) that also depend

on parameterization of the ligand (Oostenbrink et al., 2004). However, because we knew the actual

binding free energies from electrophysiology experiments we used the PBSA calculations only to test

our hypothesis that m1 is AC_{LA} and m3 is AC_{HA}.

Calculated PBSA and experimental ΔG values are compared in Figure 4A left (Figure 4-Source

Data 1). For ACh and CCh there was good agreement between ΔG_{m1} and ΔG_{LA} and between ΔG_{m3} and

ΔG_{HA}. For these agonists, *in silico* values overestimated experimental ones by ~8% and ~25%. The

agreement was not good for the other 2 agonists, with calculated values overestimating experimental

ones by ~45% (Ebt) and ~130% (Ebt). However, in all cases ΔG_{m3} was more favorable than ΔG_{m1}, and

the % overestimation was approximately the same for m1 versus m3. Despite the mismatches with the

PBSA calculations, the comparison is consistent with the hypothesis.

257   Further support for the assignments comes from comparing calculated vs experimental

efficiencies that depend on the binding free energy ratio, $\Delta G_{LA}/\Delta G_{HA}$, rather than absolute ΔGs. As

described elsewhere, efficiency is the agonist's free energy change in hold relative to its total free

energy change, $(\Delta G_{HA}-\Delta G_{LA})/\Delta G_{HA}$ or $(1 -\Delta G_{LA}/\Delta G_{HA})$ (Nayak et al., 2019). Figure 4A right (Figure 4-

Source Data 1) shows that efficiency values calculated *in silico* agree almost perfectly with those

measured experimentally. This result not only further supports the hypothesis that m1 represents AC_{LA}

and m3 represents AC_{HA}, but also indicates that efficiency can be computed from structure alone. We

conclude that MD simulations of a single pdb file successfully model the end states of the hold

transition and that the trajectories likely reproduce the conformational dynamics in $AC_{LA}{\rightarrow}AC_{HA}$.

266   All agonists show an increase in van der Waals (VdW) interaction energy in this rearrangement

(Figure 4B). The calculated increase (PBSA calculations) was most pronounced with Ebt and least for

Ebx, with CCh and ACh falling in between (Figure 4-Source Data 2). $AC_{LA}{\rightarrow}AC_{HA}$ restructuring of the

binding pocket generates a smaller, more-tightly packed pocket (Tripathy et al., 2019), in part because

of the increased VdW interactions (Video 1). This is shown in Figure 4C as a reduction in binding pocket

volume and in Figure 4D as a decrease in the number of water molecules. The VdW contacts that form

concomitant to agonist flip and loop C flop establish a more compact, hydrophobic and stable local

environment (Video 1).

Figure 4. Binding free energies and pocket properties. A. Calculated (yellow) versus experimental (blue) binding free energies for 4 agonists (Figure 2A, top) (Figure 4-Source Data 1). PBSA calculations were done on clusters selected from m1 and m3 minima of PCA plots (Figure 3; Figure 3-figure supplement 2). Left, absolute ΔG and right, efficiency. The agreement in efficiencies supports the hypothesis that m1 represents AC_{LA} and m3 represents AC_{HA}. B. In $AC_{LA} \rightarrow AC_{HA}$, VdW interactions increase (left), pocket volume decreases (center) and the number of water molecules decreases (right). Overall, the pocket stabilizes, compacts and de-wets.

Flip-Flop-Fix. In $AC_{LA}{\rightarrow}AC_{HA}$, a *cis→trans* rotation of the agonist (flip) was prominent in all simulations

(Figure 5). All agonists underwent a >130^o rotation about the N$^+$-αW149 fulcrum. Also, in all instances

the flip occurred near the start of the hold rearrangement.

A second consistent and prominent restructuring event in $AC_{LA}{\rightarrow}AC_{HA}$ was the downward

displacement (flop) of loop C toward the pocket center (C_α of αW149). This well-known 'clamshell

closure' motion is conserved in related binding proteins and thought to play a role in AChR activation

by agonists (Basak et al., 2020; Hansen et al., 2005; Hibbs et al., 2009). In the simulations, loop C flop

was measured as the displacement of its tip (C_α of αC192) in m3 relative to its apo position. The

distance traveled varied with the agonist (Figure 6-table supplement 1) but we do not discern a pattern

in the extent of loop C flop with regard to agonist affinity, efficacy or efficiency.

In hold, residues that form the pocket cavity move, presumably to accommodate the reoriented

ligand. In AC_{LA}, αY190, αY93, and δW57 are spaced apart, providing an open slot that accommodates

the agonist's tail in the AC_{LA}, *cis* orientation (Figure 5- figure supplement 1). However, in AC_{HA} loop C

flop repositions αY190 closer to αY93, effectively filling the gap to create a unified surface.

Figure 5. Agonist and loop movements in hold. In each panel *left*, superimposed cartoons of AC$_{LA}$ (m1; orange) and AC$_{HA}$ (m3; blue). Loop C is upper left and loop F is lower right. In AC$_{LA}$→AC$_{HA}$ (red→blue) there is a *cis→trans* reorientation of the agonist (a flip) and a downward movement of loop C (a flop, orange arrow); *right*, flip from m1 (red) to m2 (yellow) to m3 (blue); degree, m1→m3.

The αW149, αY190, αY198 and αY93 rings surround the N^+ (Figure 1D). In adult-type mouse

AChRs, Ala substitutions of these 4 aromatic residues have different consequences and result in ACh LA

binding free energy losses of 3.0, 2.7, 2.2 and 1.1 kcal/mol, respectively (Purohit et al., 2014). δW57A

does not change ACh binding energy (Nayak & Auerbach, 2013). αW149 (in loop B) makes a cation-π

bond with N^+ that is important for ligand stabilization (Xiu et al., 2009; Zhong et al., 1998). In the

simulations, the distance between αW149 and N^+ decreases only slightly, $AC_{LA} \rightarrow AC_{HA}$ (Figure 6, Figure

6-table supplement 1). This cation-π bond appears to act as a slippery anchor (a hinge) that maintains

stability of the N^+ position even as the tail rotates about it.

In AChBP (perhaps C_{LA}), the hydroxyl group of αY93 forms a hydrogen bond with the ligand

head group (Celie et al., 2004; Hansen et al., 2005). In the AChR AC_{LA} structure, the *cis* orientation

appears to be stabilized by this hydroxyl as it is close to a functional group on the agonist's tail.

Likewise, the tail nitrogen in CCh, Ebt, and Ebx (amide or amine) forms a hydrogen bond with the

hydroxyl group of αY198, but only in the intermediate m2 state. This interaction may guide the flip and

provide transient stability. The absence of a secondary nitrogen in ACh possibly explains its alternate

re-orientation route, as only this ligand was observed to flip counterclockwise (Figure 5). It may also

explain why the m2 basin in the PCA plots for ACh is especially broad and shallow (Figure 3).

The $AC_{LA} \rightarrow AC_{HA}$ transition compacts, dehydrates and stabilizes the binding cavity (Figure 4;

Video 1). In AC_{HA}, the tail functional moiety (carbonyl or secondary ring amine) in the *trans* orientation

is sandwiched between αC192, αC193 and αY198 (loop C), αT150 (loop B) and δC108, δN109, δL111,

and δL121 (loop E) (Figure 6- figure supplement 1). As reported elsewhere (Blum et al., 2010; Rahman

et al., 2020; Zarkadas et al., 2022), in AC_{HA} the functional group found in the tail of each agonist -- the

carbonyl group in CCh and ACh and the secondary ring amine in Ebt and Ebx - forms a hydrogen bond

with the carbonyl of δN109 and amine of δL121 in the complementary δ subunit backbone *via* a

structural water (Figure 6). The agonist's flip occurs at the start of m1→m2, and the establishment of

this water-mediated H-bond completes in m2→m3 (Figure 6) (Video 1). This water-mediated bonding

and the increase in VdW interactions appear to both be crucial for stabilizing the loops and agonist in

the HA conformation of the neurotransmitter site.

Salt-bridge. Near the pocket, the αK145-αD200 salt bridge (Figure 1D) has been suggested to play an

important role in receptor activation (Beene et al., 2002; Gay & Yakel, 2007; Mukhtasimova et al.,

2005; Padgett et al., 2007; Pless et al., 2011). The hold rearrangements of αK145 are complex and

agonist dependent. In AC_{HA}, this residue i) makes an apparent contact with αY93, ii) contacts the αY190

hydroxyl group (Figure 6-table supplement 1) and, iii) approaches loop F of the complementary

subunit.

341   Interestingly, the mutation αK145A lowers the efficiencies of ACh and CCh (from 0.51 to 0.41)

but has no effect on those of Ebt and Ebx (0.41 and 0.46) (Indurthi & Auerbach, 2023). This agonist

dependence in function has a parallel in structure. In the hold simulations, the αK145-αD200 distance

(Å) increases from only 4.4 in apo to 6.1 or 5.1 in AC_{LA} (CCh or ACh). This spreading may relate to the

agonist's tail being inserted into the αY93 slot in the *cis* orientation. In AC_{LA}→AC_{HA}, for some agonists

the αK145-αD200 separation (Å) shortens substantially to 2.8 or 2.6, making it more rigid (ACh or CCh).

However, with Ebt and Ebx the αK145-αD200 distance remains unstable, fluctuating between 3-6 Å

throughout the course of the simulation. Likewise, αK145-αY190 distance stabilizes at 5.3 Å or 6.1 Å in

AC_{HA} with ACh and CCh but continues to fluctuate with Ebt and Ebx. In contrast, the αY190-αD200

distance remains stable in AC_{HA} for all ligands. This suggests that the instability in the αK145-αD200 and

αK145-αY190 distances can primarily be attributed to αK145 (Figure 6- figure supplement 2).

Figure 6: Representative snapshots in $AC_{LA}\rightarrow AC_{HA}$ (hold). *left*, rearrangements of loop C, loop F and the ligand (red, m1; yellow, m2; blue, m3); *right*, residue and ligand orientations. m1 is AC_{LA}, m2 is an intermediate state, m3 is AC_{HA}. In m1, a functional group in the agonist tail interacts with αY93 (all agonists) and αD200 (only CCh and Ebt). The position and orientation of αW149 relative to N^+ of the agonist remains nearly unchanged m1→m2→m3 and serves as a fulcrum for the *cis→trans* flip (see Figure 5). In m2, the functional nitrogen at the agonist tail (CCh, Ebt and Ebx) interacts with the hydroxyl group of αY198. For all ligands, αY190 repositioning and loop C flop (m1→m3) are correlated. In m3, the agonist fully flips to *trans*, facilitating VdW interactions, de-wetting, and the formation of water-mediated hydrogen bonds with the reactive group at its tail with δN109/δL121 backbone (loop E) via a structural water.

Discussion

After diffusing to the target, the only agonist-dependent rearrangements (energy changes) in receptor activation are in catch and hold, $A\text{-}C \rightleftarrows AC_{LA} \rightleftarrows AC_{HA}$ (Figure 1B, bottom) (Auerbach, 2024; Jadey & Auerbach, 2012). By comparing binding energies calculated *in silico* with those measured *in vitro* we were able to identify in the MD trajectories the hold end states, AC_{LA} and AC_{HA}, plus an intermediate configuration not detected in electrophysiology experiments. Three conspicuous hold rearrangements discussed below are: i) a *cis→trans* rotation of the agonist (flip), ii) an up→down displacement of loop C (flop), and iii) reductions in pocket volume, water and fluctuations (fix).

There was reasonably good agreement between calculated and experimental binding free energies for ACh and CCh, but the match was excellent when free energy ratios (efficiencies) rather than absolute values were compared (Figure 4A). Absolute *in silico* values overestimated experimental ones by approximately the same factor (different for each ligand) that cancelled in the ratio. We do not know the origin of this factor but speculate that it is caused by errors in ligand parameterization. Regardless, the results show that agonist efficiency can be a useful metric for associating structure with function, and that efficiency can be estimated from structure alone.

Below we consider the structural elements associated with flip, flop and fix, namely the agonist (*cis→trans*), loop C (up→down) and the pocket (large→small, *et al*). MD simulations of the α–δ AChR neurotransmitter site suggest the following approximate and jittery generic sequence of rearrangements in $AC_{LA} \rightarrow AC_{HA}$ (Video 1). The stable states are bold.

1. **AC_{LA} (m1)**: agonist *cis*, loop C up, pocket wide/wet/wobbly,
2. ...agonist leaves *cis* and starts to rotate about the N^{+}-αW149 fulcrum,
3. ...loop C starts to move down (towards the fulcrum).

4. **Intermediate (m2)**: agonist half-rotated, tail stabilized by αY198,

5. ...agonist rotates to *trans* (little change at the fulcrum), tail not secured in the pocket,

6. ...loop C is fully down.

7. **AC$_{HA}$ (m3):** agonist *trans*, tail secured by VdW contacts and H-bonds, pocket small/dry/rigid.

<u>Agonist.</u>   With all agonists and in all simulations, the rotation of the agonist about the N^+-αW149

fulcrum begins at the start of the AC$_{LA}{\rightarrow}$AC$_{HA}$ transition.   Movements of bound ligands have been

identified in other proteins. In AMPA-type glutamate receptors, the neurotransmitter glutamate binds

in either of 2 poses, crystallographic or inverted (Yu et al., 2018). In one MD simulation these

interconverted, suggesting the bound ligand is not rigidly fixed.   Although the alternative poses were

not associated with binding energies, the binding cleft was more closed in the crystallographic

(presumably HA) versus inverted (possibly LA) orientation.   The re-orientation of a bound glutamate in

AMPA receptors is possibly related to the flip of the agonist in AChRs.

Multiple ligand positions in binding pockets have also been reported in Glycine receptors (Yu et

al., 2014), 5-HT3 receptors (Basak et al., 2020), δ-opioid receptors (Shang et al., 2016), HIV-1 protease

(Klei et al., 2007) and transthyretin (Klabunde et al., 2000). In enzymes, bound substrate movement is

apparent in hexokinase (Bennett & Steitz, 1978) and alcohol dehydrogenase (Plapp, 2010). It is not

unusual for movement of a bound ligand to trigger an otherwise unfavorable local rearrangement that

drives protein function (an 'induced fit'; (Richard, 2022).   Not only can an untethered ligand come and

go, it can also reorient itself dramatically to change binding energy and, hence, the local environment

that couples to the full allosteric transition.

In AChRs, such ligand reorientation cannot pertain to all agonists. Some with high potency are

perfectly symmetrical (decamethonium) and some lack a tail (tetramethylammonium). Although we

have not yet identified the conformational dynamics that determine ΔG_{HA}, factors other than the flip

certainly contribute, and in proportions that are likely to be agonist dependent (see below).

The PCA plots (Figure 3) show 3 structural populations, one of which (m2) occurs in time

between m1 (AC_{LA}) and m3 (AC_{HA}) (Figure 3-figure supplement 2). In m2, the agonist's tail (CCh, Ebt and

Ebx) interacts with αY198, possibly to guide the clockwise rotation of the ligand and reduce the

probability of a return to the *cis* orientation. The PBSA calculations indicate that ΔG_{m2} (that does not

have an experimental correlate) is intermediate between ΔG_{LA} and ΔG_{HA} for ACh and CCh, but least

stable for Ebt and Ebx (Figure 4-Source Data 1). We do not have an explanation for this difference.

Loop C. A number of observations place focus on the role of the downward displacement of loop C in

receptor activation. In ACh binding proteins, its extent correlates with agonist potency (Basak et al.,

2020; Du et al., 2015; Polovinkin et al., 2018) even if voltage-clamp fluorometry of other receptors

shows it to be similar with agonists and antagonists (Chang & Weiss, 2002; Munro et al., 2019; Pless &

Lynch, 2009). Simulations of the apo AChR indicate that in the absence of agonists loop C has flopped

partially, narrowing the aqueous connection between the bath and the pocket (Figure 2-figure

supplement 2C).

Given the flop in loop C, it is puzzling that electrophysiology measurements show the agonist

association rate constant (k_{on}) is substantially *faster* to C_{HA} (narrow entry, loop C down) than to C_{LA}

(wide entry, loop C up), as well as more correlated with agonist potency (Grosman & Auerbach, 2001;

Nayak & Auerbach, 2017). Docking results offer a clue to the apparent contradiction that narrowing an

entrance speeds association up to the diffusional limit (4 x10^9 M^{-1}s^{-1}). All 4 agonists dock in the *cis*

orientation to C$_{LA}$ (wide) versus in the *trans* orientation to C$_{HA}$ (narrow) (Figure 2B). Hence, despite the

wider aperture, *trans* is disfavored and lower affinity *cis* is preferred.   We hypothesize that binding in

the *trans* orientation is prohibited in the apo pocket, and that loop C flop (and other rearrangements)

both narrow the entrance and remove this prohibition. Accordingly, k$_{on}$ with loop C up (LA) is slower

than with loop C down (HA). We do not know the structural element(s) that prohibits *trans* with a wide

pocket. Any or all of the constellation of rearrangements in unliganded C$_{LA}$→C$_{HA}$ could contribute.

As an aside, the inward displacement of loop C toward the pocket center resembles the

'clamshell closure' apparent at other receptor binding sites (Chen et al., 2017; Hansen et al., 2005;

Hibbs & Gouaux, 2011). However, in light of the k$_{on}$ experimental results, using 'closure' (or 'capping')

to describe this motion is misleading because it implies putting a lid on the pocket that would reduce,

not increase, the agonist association rate constant. For this reason, we use functionally neutral words

'flop' and 'up/down' for loop C displacement and position.

AChRs can turn on without a loop C. Truncation of loop C eliminates activation by ACh but has

almost no effect on unliganded activation (Purohit & Auerbach, 2013). Interestingly, leaving only loop

C residues αY190 and αY198 in place saves activation by ACh. Hence, repositioning of one (or both) of

these in up→down is sufficient to maintain activation by the neurotransmitter. Other agonists were

not investigated.

αY190 is our interest. An Ala substitution here results in a large loss of ACh binding energy

with about half coming from the removal of the hydroxyl group (Purohit et al., 2014). In contrast, the

mutations αY198F and αY93F have essentially no effect on ACh binding energy. These results suggest

that repositioning αY190 consequent to loop C flop is important for activation by ACh (Figure 6). The

αY190 ring fills the slot that opens next to αY93 when the agonist exits *cis*, preventing a return (Figure

5-figure supplement 1). However, the distance between the αY190 ring and N^+ does not change

significantly so it is unlikely that the deployment of αY190 increases binding energy *via* an interaction

with the ring (Figure 6-table supplement 1).

Perturbation of the salt bridge residue αD200 by the αY190 hydroxyl has been proposed to

trigger the AChR allosteric transition (Beene et al., 2002; Gay & Yakel, 2007; Mukhtasimova et al.,

2005; Padgett et al., 2007; Pless et al., 2011). In support of this idea, the mutation αK145A reduces

ACh (and choline) binding energy substantially but has no effect when the αY190 hydroxyl is absent

(Bruhova & Auerbach, 2017). In the simulations, the downward displacement of loop C positions the

αY190 hydroxyl to within ~5.0 Å of αD200 and αK145 (Figure 6-table supplement 1). We sought, but

did not find, structural water in this gap. Although there was no correlation between these separations

and the degree of loop C displacement, the gap was smaller with high- (ACh and CCh) versus low-

efficiency agonists (Ebt and Ebx). In summary, some evidence supports the above proposal, but so far,

the simulations have not illuminated a mechanism. Substrate-induced loop displacements are known

to deploy residues to enhance catalysis, for example in chymotrypsin (Ma et al., 2005) and

triosephosphate isomerase (Brown & Kollman, 1987; Nickbarg et al., 1988). Repositioning αY190 in the

AChR $AC_{LA} \rightarrow AC_{HA}$ transition could be analogous.

Although attention on loop C has focused on the role of the flop, it is also important to consider

the functional consequences of the up position. The structural elements of the up (wide cleft)

configuration that prohibit agonist docking into the *trans* orientation have not been identified. A key

requirement for allostery is that the ligand bind more strongly to one of the alternative conformations

of the orthosteric site, so attention must also be given to the conformational dynamics of the 'catch

rearrangement, A-C$\rightleftarrows$AC$_{LA}$. In both GABA$_A$ (Jones et al., 2001; Mortensen et al., 2010) and ACh

receptors (Jadey & Auerbach, 2012) the LA agonist association rate constant is correlated with potency

rather than diffusion constant, indicating that the formation of AC$_{LA}$ requires an energy (structure)

change.

<u>Pocket.</u> The αW149-N$^+$ hinge remains relatively unchanged in the hold transition (Figure 6, Figure 6-

table supplement 1). Hence, we hypothesize the bulk of the extra favorable agonist binding energy in

AC$_{HA}$ is not generated here but somewhere else, *cis$\rightarrow$trans*. For the agonists we examined, the rotation

places the tail into the main binding cavity (Figure 6-figure supplement 1) (Video 1). The *trans*

orientation is secured by VdW interactions (possibly caused by further downward loop C displacement

and dehydration) and a hydrogen bond with a structural water bonded to the loop E backbone (in the

complementary subunit). The net result to the overall pocket is a reduction in volume (Figure 4C) and

increase in stability of loops C and F (Figure 3).

In the simulations, these two structural changes, distributed over the large surface area of the

cavity, were robust and consistent for all 4 agonists. It is possible that the increase in binding free

energy in hold derives mainly from restructuring of the pocket rather than from a change in the N$^+$-

aromatic interactions or, indeed, any one local ligand-protein interaction. Compact, hydrophobic

binding pockets are associated with functional activation in other ion channels (Furukawa et al., 2005;

Sigel & Steinmann, 2012) and observed in other proteins including β2-adrenergic receptors, opioid

receptors, and HIV-1 protease (Huang et al., 2015; Rasmussen et al., 2011; Wlodawer & Erickson,

1993).

After hold, the next step In AChR gating is the rearrangement of the extracellular domain

(Auerbach, 2024; Purohit et al., 2013). In related receptors, this domain becomes more compact and

stable (a rearrangement called 'unblooming') (Sauguet et al., 2014; Taly et al., 2009), structural

changes that echo those in hold at the AChR neurotransmitter site. It is therefore possible that the

$AC_{LA} \rightarrow AC_{HA}$ rearrangement of the neurotransmitter site nucleates the subsequent restructuring of the

extracellular domain. Extending this speculation, the structural correlates of energy change in gating

might at first spread in contact area (from N^+, to pocket, to extracellular domain) and then refocus

(from transmembrane domain, to gate region, to pore water).

In the complementary δ subunit (Figure 1D), loop E ($\beta5$-$\beta6$ strands) has been implicated in

ligand discrimination (Basak et al., 2020; Muroi et al., 2009; Pless & Lynch, 2009). The double mutation

F106L+S108C in the $\beta2$ subunit of the $\alpha4\beta2$ nicotinic AChR subtype significantly decreases the action of

Ebt without altering that of ACh (Tarvin et al., 2017). In addition, loop E residues are responsible for the

binding energy differences $\alpha-\gamma$ versus $\alpha-\delta$ (Nayak et al., 2016) and reduce site instability caused by

loop C mutations (Vij et al., 2015). Although our simulations do not shed light on these matters, the

temporal sequence flip-flop-fix suggests that perturbation of the loop E backbone could also play a role

in downstream activation of the receptor.

Agonists belong to efficiency classes in which $\Delta G_{LA}/\Delta G_{HA}$ is the same for all members. So far 5

such classes have been identified experimentally in adult-type AChRs, indicating that also are 5 pairs of

AC_{LA}/AC_{HA} structural classes (Indurthi & Auerbach, 2023). Although in the simulations loops C and F

movements, salt bridge distances were agonist dependent, with only 4 ligands we cannot associate any

these rearrangements with efficiency classes. Additional simulations using more agonists and with

binding site mutations may lead to better understanding of the dynamics that underpin molecular

recognition in catch and protein animation in hold, and perhaps reveal the structural basis of their

linkage (Auerbach, 2024).

Implications. Several aspects of this work may have general implications. The LA and HA end states of

hold emerged from simulations of a single protein structure and were identified by comparing

experimental with calculated binding energies. If general, this approach could be useful because cryo-

EM structures of ligands bound with LA are lacking. Agonist efficiency was calculated accurately from

structure alone. This, too, could be useful because efficiency allows affinity and efficacy to be

calculated from each other (Indurthi & Auerbach, 2021). In AChRs the up→down movement of loop C

flop does not impose a lid on the pocket. Rather, the up position serves to enforce catch (ligand

recognition) by blocking directly binding in the *trans* orientation, and the down position serves to

stabilize the AC_{HA} pocket and, perhaps, to deploy $\alpha Y190$. It is possible that the 'clamshell closures'

apparent in other receptors also serve to activate for reasons other than imposing a steric barrier.

Finally, it appears that the movement of the agonist, bound with low affinity, is the initial event in

gating. If confirmed, we think it both fitting and beautiful that the motion of a structural element

added to the system from the outside jumpstarts the allosteric transition.

Materials and Methods

Hardware and Software. All computational studies were carried out on a Linux Ubuntu 20.04 based workstation with 126 GB RAM, 64 CPU cores, and two RTX A5000 24 GB GPU cards, as well as an Ubuntu 20.04 based server equipped with 132 GB RAM, 96 CPU cores, and two RTX A5000 24 GB GPU cards. The software utilized includes the Maestro release 2022-2 graphical user interface (GUI) of the Schrödinger software suite (BioLuminate, Schrödinger, LLC, NY, USA), Amber22 (Case et al., 2005), VMD 1.9.3 (Humphrey et al., 1996), ChemDraw Professional 15.1 (Purushotham et al., 2022), ChimeraX (Pettersen et al., 2021), and PyMOL (PyMOL Molecular Graphics System, Version 2.0, Schrödinger, LLC).

Protein preparation. Because agonist energy changes are approximately independent of energy changes in the protein outside the binding pocket, for the simulations we removed the transmembrane domain and simulated only $\alpha-\delta$ subunit extracellular domain dimers. The starting structure for the hold transition was resting C (PDB: 6UWZ) after toxin removal and is essentially the same as the apo structure (Zarkadas et al., 2022) (Figure 1C). The 4 agonists we examined were acetylcholine (ACh), carbamylcholine (CCh), epibatidine (Ebt) and epiboxidine (Ebx) (Figure 2A). These are of similar size (MW 187, 183, 209 and 178, respectively) but have different efficiencies (0.50, 0.52, 0.42, and 0.46, respectively) (Indurthi & Auerbach, 2023).

Protein preparation was conducted using the Protein Preparation Wizard (PPW) (Purushotham et al., 2022) and OPLS4 force field. Bond orders were assigned to atoms based on the CCD database, with zero bond orders given to bonded metals. Missing hydrogen atoms were inserted, and disulfide bonds were placed appropriately. During preprocessing, structural refinement was performed with Maestro's Prime utility, optimizing and adding any missing side chains and residues. Missing residues were formulated by combining SEQRS information from the PDB files and optimized using Prime's

ePLOP module. The ionization state of all heterogeneous groups, including ligands, bound metals, and

charged amino acids, were determined using the EpiK tool at a pH range of 7.0 ± 2.0. Subsequent force

field minimization allowed for water sampling and basic restraint minimization, during which only

hydrogen atoms could be freely reduced.

The hydrogen bonding (H-bond) network underwent significant modification. The terminal chi

angle for residues Asn, Gln, and His was sampled through 180-degree flips, altering their spatial H-

bonding capabilities without affecting the fit to electron density. The neutral and protonated states of

His, Asp, and Glu, as well as two His tautomers, were sampled. Hydrogens on hydroxyls and thiols were

also analyzed to enhance the H-bond network. Finally, the ProtAssign method was employed in two

modes: a "regular" mode, which completed in seconds, and an "exhaustive" mode that considered

many more states and could take minutes or hours, depending on the H-bond network complexity.

Ligand Preparation. The two-dimensional SDF structures of carbamylcholine (CCh), acetylcholine (ACh),

epibatidine (Ebt) and epiboxidine (Ebx) were retrieved from the PUBCHEM database. These agonists

were chosen because they have a similar size and represent different efficiency classes (Indurthi &

Auerbach, 2023). LigPrep (LigPrep, Maestro 11.2.014, Schrödinger LLC) was utilized to construct

ligands, produce stereoisomers, and generate tautomers using the OPLS46 force field. Additionally, the

Epik module was employed to desalt and protonate the ligands at pH 7.0. Default values were applied

to the remaining parameters, and LigPrep was also used to prepare the ligands ACh, CCh, Ebt, and Ebx,

with the option "retain specified chiralities" engaged.

In terms of ligand state assessment, Epik can calculate a penalty to quantify the energy cost of

forming each state in solution based on the Hammett and Taft methods. This EpiK state penalty is

entirely compatible with the GlideScore used for docking, as it is computed in kcal/mol units. This

compatibility facilitates the examination of how the EpiK state penalty affects the GlideScore. The

DockingScore in Glide represents the total of the GlideScore and the Epik state penalty and served as

the basis for final ranking and enrichment evaluation. Furthermore, Epik offers a technique for

handling metal binding states, involving a change in the pH range within which the state is generated.

This approach underscores the comprehensive considerations made in preparing the ligands for

docking and subsequent analyses.

Grid Generation and Molecular Docking. The receptor grid complex of 6UWZ.pdb was generated after

removing alpha-bungarotoxin from the binding pocket using Glide (Glide, Maestro 11.2.014,

Schrödinger LLC). A virtual grid box with dimensions 20Å × 20Å × 20 Å was formed, centered within the

C and F loops in the $\alpha-\delta$ binding site of the energy-minimized resting state. This grid box was used to

generate a receptor grid, and the default values were assigned for other parameters. Docking was

performed using the ligand docking wizard built into the Glide module of Schrödinger, with the default

scaling factor and partial charge cutoff set to 0.80 and 0.15, respectively. The docking procedure

consisted of two stages, extra precision (XP) followed by post-processing with prime MM-GBSA. The

best dock poses were selected based on dock score and MM-GBSA energy for further investigation.

Molecular Dynamics Simulations. MD simulations were conducted using the Amber22 software suite

within the Amber module as detailed elsewhere (Singh et al., 2021; Srivastava et al., 2022). Ligand

topology was prepared via the Leap module in AmberTools22, employing force field ff19SB (Tian et al.,

2020) and the Generalized Amber Force Field (GAFF) (Vassetti et al., 2019). The AM1BCC strategy and

GAFF2 were used to assign atomic charges and additional parameters. Before simulation, the systems

were minimized and equilibrated following a five-step minimization process, with each step involving

10,000 energy minimization steps. Systems were heated under NVT ensembles in two consecutive runs

from 0–300 K for 1 ns each, followed by a 1 ns simulation at 300 K and 1 bar pressure under the NPT

ensemble for equilibration. An additional 5 ns equilibration was performed before the 200 ns

production MD for each system, conducted under periodic boundary conditions (PBCs) and utilizing an

NPT ensemble with 300 K and 1 bar pressure. Long-range electrostatic interactions were treated using

the particle mesh Ewald method (PME), including a direct space cutoff of 12.0 Å and van der Waals

interactions. In production MD, a time step of 2.0 fs was employed, and the SHAKE algorithm was used

to keep bond lengths at equilibrium. Isobaric (NPT) conditions were maintained using the Berendsen

barostat, with temperature monitored by a Langevin thermostat. Coordinates from the production MD

were recorded in the trajectory file every 20 ps, resulting in a minimum of 10,000 frames. Trajectories

were analyzed using amber-tools CPPTRAJ (Roe & Cheatham, 2013), and further analysis and figure

generation were completed with VMD, Pymol, Xmgrace and ChimeraX. To ensure reproducibility, MD

trajectories were generated in triplicate.

Cluster Analysis. Cluster analysis of the ligand was performed using TCL scripts within the Visual

Molecular Dynamics (VMD) software, as described elsewhere (Mittal, Tonk, et al., 2021). In brief, this

script automates the identification of frames from the most stable conformations based on PCA

minima. Using these frames as input, the algorithm employed RMSD values of ligand positions to

cluster similar orientations. Specifically, clusters with an RMSD of ≤ 1Å were considered to share a

similar ligand orientation. Frames that did not fit into these top clusters were automatically excluded

by the script.

Principal Component Analysis (PCA) and Free Energy Landscape (FEL). PCA identifies collective

motions in atomic-level simulations of macromolecules (Mittal, Srivastava, et al., 2021; Srivastava et

al., 2022). It emphasizes significant patterns by reducing noise in MD simulations, revealing essential

dynamics underlying conformational changes. All equilibrated MD simulation trajectories were

analyzed by PCA computations using the CPPTRAJ algorithm using Amber tools.

Prior to PCA, least-square fitting of the protein's Cα atoms was carried out with respect to a reference

structure (average protein coordinates) to eliminate rotational and translational degrees of freedom in

the simulation box. A positional covariance matrix C (of dimensions $3N \times 3N$) was constructed using the

Cα atom coordinates, and this matrix was diagonalized to produce eigenvectors (directions of motion)

and eigenvalues (magnitudes). The elements of the covariance matrix C were obtained using:

$C_i = \langle(X_i - \langle X_i \rangle)(X_j - \langle X_j \rangle)\rangle$▯▯▯$(i, j=1,2,3.......,3N)$,   (Eq. 1)

where X_i and X_j are the Cartesian coordinates of the Cα atoms, and N is the number of Cα atoms. The $\langle\rangle$

notation represents the ensemble average of atomic positions in Cartesian space. The resulting

principal components (eigenvectors) were sorted by the total motion they captured, generating 3N

eigenvectors. The free energy landscapes (FELs; Figure3) were created using the "g_sham" module in

GROMACS, based on the probability distribution of the top two principal components (PCs). The FEL

plot aids in visualizing possible conformations and corresponding energy levels. The free energy values

(G) were calculated as

$G_i = -kT \ln(N_i/N_{max})$ (Eq. 2)

where k is the Boltzmann constant, T is the absolute temperature of the simulated system (300 K), N_i is

the population of bin i, and N_{max} is the population of the most-populated bin.

Binding Free Energy Calculation. Experimental binding free energies were estimated by using

electrophysiology, as described elsewhere (Indurthi & Auerbach, 2023). Briefly, the constants K_{HA} and

L_2 were estimated by using electrophysiology (dose-response curves, for example), and K_{HA} was

calculated using the thermodynamic cycle with L_0 known *a priori*. The logs of K_{LA} and K_{HA} are

proportional to ΔG_{LA} and ΔG_{HA}. The binding free energy of the protein-ligand docked complex (ΔG_{bind})

was calculated in two stages, after docking (using MM-GBSA in the Prime module of Schrödinger)

(Mittal, Tonk, et al., 2021; Purushotham et al., 2022) and after MD simulation on equilibrated MD

trajectories (using MM-PBSA/-GBSA in AMBER22). MM-PBSA is particularly useful to compare binding

affinities between different states and among agonists (Kollman et al., 2000; Rastelli et al., 2010; Sun et

al., 2014). In our use, MM-PBSA computations included both enthalpic (ΔH) contributions from

molecular interactions (van der Waals, electrostatics, solvation) and entropic ($T\Delta S$) contributions

calculated using normal mode analysis. By capturing the dynamic interplay between the ligand and

receptor as well as the influence of the surrounding solvent environment this approach enables

computation of a Gibbs free energy (ΔG).

656   After docking, for the static docked pose of the protein-ligand complex the MM-GBSA binding

energy calculation was performed using the OPLS4 force field and the variable-dielectric generalized

Born (VSGB) model in the Prime MM-GBSA module in Schrödinger 2022. We used ΔG_{bind_dock} to rank

protein-ligand docked complexes, selecting those with the highest negative values and dock scores as

the best ligands for further studies. After MD Simulation, in the MD equilibrated system binding

energies were calculated through the MM-PBSA/-GBSA protocol, implemented in AMBER22. This was

performed over the most stable clustered poses extracted from the equilibrated trajectory at minima

m1, m2, and m3 on the free energy landscape (FEL) over a 200 ns time frame. AMBER utilizes the

conventional g_mmpbsa module for MM-PBSA calculations using MM-PBSA.py script. These methods

calculate the energies of electrostatic interactions, van der Waals interactions, polar solvation, and

non-polar solvation in the equilibrated trajectory using a single trajectory protocol (STP). The binding

energy of each protein-ligand complex was calculated as

$\Delta G_{bind} = \Delta G_{complex} - \Delta G_{protein} - \Delta G_{ligand}$ where $\Delta G_{complex}$, $\Delta G_{protein}$, and ΔG_{ligand} represent the absolute free

energies of the complex, protein, and ligand systems, respectively. Additional terms, such as ΔH

(enthalpy), T and ΔS (temperature and total solute entropy), ΔG_{gas} (total gas phase energy), and ΔG_{solv}

(solvation energy), contribute to the calculation.

The Poisson-Boltzmann calculations were performed using an internal PBSA solver in Sander,

while Generalized Born ESURF was calculated using 'LCPO' surface areas. The vibrational mode entropy

contributions ($T\Delta S$) for protein-ligand interactions were calculated and averaged using normal-mode

analysis in Amber. The entropy term provides insights into the disorder within the system, as well as

how this disorder changes during the binding process. This value was added to the enthalpy term

computed by the MM-PBSA method to determine the absolute binding free energy (ΔG_{bind}). These

comprehensive calculations allowed for a precise evaluation of the interactions between the protein

and ligand, providing valuable insights into the binding process and its energetic characteristics.

Pocket Volume Calculation. To investigate the variations in binding site volume across all four agonist-

bound cases, pocket volume analysis was carried out using POVME3.0 (Wagner et al., 2017). From each

of the four systems, frames were extracted at consistent intervals with a stride of 2.0, culminating in

representative sets of 5000 protein structures for every system. Before commencing the volume

calculation, the trajectory was aligned, and frames were selected from VMD, serving as the initial input

for this method. Subsequently, an inclusion region was defined, encapsulating all binding pocket

conformations within the trajectory. This region's definition involved building a sphere upon which the

inclusion grid-points were computed, using different atoms of the ligand for this calculation.

References:

Auerbach, A. (2020). Pathways for nicotinic receptor desensitization. *J Gen Physiol*, *152*(10). https://doi.org/10.1085/jgp.202012639

Auerbach, A. (2024). Dynamics of receptor activation by agonists [Perspective]. *Biophysical Journal*, *123* (15). https://doi.org/https://doi.org/10.1016/j.bpj.2024.01.003

Basak, S., Kumar, A., Ramsey, S., Gibbs, E., Kapoor, A., Filizola, M., & Chakrapani, S. (2020). High-resolution structures of multiple 5-HT(3A)R-setron complexes reveal a novel mechanism of competitive inhibition. *Elife*, *9*. https://doi.org/10.7554/eLife.57870

Beene, D. L., Brandt, G. S., Zhong, W., Zacharias, N. M., Lester, H. A., & Dougherty, D. A. (2002). Cation-pi interactions in ligand recognition by serotonergic (5-HT3A) and nicotinic acetylcholine receptors: the anomalous binding properties of nicotine. *Biochemistry*, *41*(32), 10262-10269. https://doi.org/10.1021/bi020266d

Bennett, W. S., Jr., & Steitz, T. A. (1978). Glucose-induced conformational change in yeast hexokinase. *Proc Natl Acad Sci U S A*, *75*(10), 4848-4852. https://doi.org/10.1073/pnas.75.10.4848

Blum, A. P., Lester, H. A., & Dougherty, D. A. (2010). Nicotinic pharmacophore: the pyridine N of nicotine and carbonyl of acetylcholine hydrogen bond across a subunit interface to a backbone NH. *Proc Natl Acad Sci U S A*, *107*(30), 13206-13211. https://doi.org/10.1073/pnas.1007140107

Brown, F. K., & Kollman, P. A. (1987). Molecular dynamics simulations of "loop closing" in the enzyme triose phosphate isomerase. *J Mol Biol*, *198*(3), 533-546. https://doi.org/10.1016/0022-2836(87)90298-1

Bruhova, I., & Auerbach, A. (2017). Molecular recognition at cholinergic synapses: acetylcholine versus choline. *J Physiol*, *595*(4), 1253-1261. https://doi.org/10.1113/JP273291

Case, D. A., Cheatham, T. E., 3rd, Darden, T., Gohlke, H., Luo, R., Merz, K. M., Jr., Onufriev, A., Simmerling, C., Wang, B., & Woods, R. J. (2005). The Amber biomolecular simulation programs. *J Comput Chem*, *26*(16), 1668-1688. https://doi.org/10.1002/jcc.20290

Celie, P. H., van Rossum-Fikkert, S. E., van Dijk, W. J., Brejc, K., Smit, A. B., & Sixma, T. K. (2004). Nicotine and carbamylcholine binding to nicotinic acetylcholine receptors as studied in AChBP crystal structures. *Neuron*, *41*(6), 907-914. https://doi.org/10.1016/s0896-6273(04)00115-1

Chang, Y., & Weiss, D. S. (2002). Site-specific fluorescence reveals distinct structural changes with GABA receptor activation and antagonism. *Nat Neurosci*, *5*(11), 1163-1168. https://doi.org/10.1038/nn926

Chen, S., Zhao, Y., Wang, Y., Shekhar, M., Tajkhorshid, E., & Gouaux, E. (2017). Activation and Desensitization Mechanism of AMPA Receptor-TARP Complex by Cryo-EM. *Cell*, *170*(6), 1234-1246 e1214. https://doi.org/10.1016/j.cell.2017.07.045

Du, J., Lu, W., Wu, S., Cheng, Y., & Gouaux, E. (2015). Glycine receptor mechanism elucidated by electron cryo-microscopy. *Nature*, *526*(7572), 224-229. https://doi.org/10.1038/nature14853

Furukawa, H., Singh, S. K., Mancusso, R., & Gouaux, E. (2005). Subunit arrangement and function in NMDA receptors. *Nature*, *438*(7065), 185-192. https://doi.org/10.1038/nature04089

Gay, E. A., & Yakel, J. L. (2007). Gating of nicotinic ACh receptors; new insights into structural transitions triggered by agonist binding that induce channel opening. *J Physiol*, *584*(Pt 3), 727-733. https://doi.org/10.1113/jphysiol.2007.142554

Genheden, S., & Ryde, U. (2015). The MM/PBSA and MM/GBSA methods to estimate ligand-binding
affinities. *Expert Opin Drug Discov*, *10*(5), 449-461.
https://doi.org/10.1517/17460441.2015.1032936
Grosman, C., & Auerbach, A. (2001). The dissociation of acetylcholine from open nicotinic receptor
channels. *Proc Natl Acad Sci U S A*, *98*(24), 14102-14107.
https://doi.org/10.1073/pnas.251402498
Grosman, C., Zhou, M., & Auerbach, A. (2000). Mapping the conformational wave of acetylcholine
receptor channel gating. *Nature*, *403*(6771), 773-776. https://doi.org/10.1038/35001586
Gupta, S., Chakraborty, S., Vij, R., & Auerbach, A. (2017). A mechanism for acetylcholine receptor
gating based on structure, coupling, phi, and flip. *J Gen Physiol*, *149*(1), 85-103.
https://doi.org/10.1085/jgp.201611673
Hansen, S. B., Sulzenbacher, G., Huxford, T., Marchot, P., Taylor, P., & Bourne, Y. (2005). Structures of
Aplysia AChBP complexes with nicotinic agonists and antagonists reveal distinctive binding
interfaces and conformations. *Embo Journal*, *24*(20), 3635-3646.
https://doi.org/10.1038/sj.emboj.7600828
Hibbs, R. E., & Gouaux, E. (2011). Principles of activation and permeation in an anion-selective Cys-loop
receptor. *Nature*, *474*(7349), 54-60. https://doi.org/10.1038/nature10139
Hibbs, R. E., Sulzenbacher, G., Shi, J., Talley, T. T., Conrod, S., Kem, W. R., Taylor, P., Marchot, P., &
Bourne, Y. (2009). Structural determinants for interaction of partial agonists with acetylcholine
binding protein and neuronal alpha7 nicotinic acetylcholine receptor. *Embo Journal*, *28*(19),
3040-3051. https://doi.org/10.1038/emboj.2009.227
Hou, T., Wang, J., Li, Y., & Wang, W. (2011). Assessing the performance of the MM/PBSA and
MM/GBSA methods. 1. The accuracy of binding free energy calculations based on molecular
dynamics simulations. *Journal of Chemical Information and Modeling*, *51*(1), 69-82.
https://doi.org/10.1021/ci100275a
Huang, W., Manglik, A., Venkatakrishnan, A. J., Laeremans, T., Feinberg, E. N., Sanborn, A. L., Kato, H.
E., Livingston, K. E., Thorsen, T. S., Kling, R. C., Granier, S., Gmeiner, P., Husbands, S. M.,
Traynor, J. R., Weis, W. I., Steyaert, J., Dror, R. O., & Kobilka, B. K. (2015). Structural insights into
micro-opioid receptor activation. *Nature*, *524*(7565), 315-321.
https://doi.org/10.1038/nature14886
Humphrey, W., Dalke, A., & Schulten, K. (1996). VMD: visual molecular dynamics. *J Mol Graph*, *14*(1),
33-38, 27-38. https://doi.org/10.1016/0263-7855(96)00018-5
Indurthi, D., & Auerbach, A. (2021). Agonist Efficiency Estimated from Concentration Response Curve.
Biophysical Journal, *120*(3), 57a-57a. <Go to ISI>://WOS:000629601400276
Indurthi, D. C., & Auerbach, A. (2023). Agonist efficiency links binding and gating in a nicotinic receptor.
Elife, *12*. https://doi.org/10.7554/eLife.86496
Jackson, M. B. (1986). Kinetics of unliganded acetylcholine receptor channel gating. *Biophys J*, *49*(3),
663-672. https://doi.org/10.1016/S0006-3495(86)83693-1
Jadey, S., & Auerbach, A. (2012). An integrated catch-and-hold mechanism activates nicotinic
acetylcholine receptors. *J Gen Physiol*, *140*(1), 17-28. https://doi.org/10.1085/jgp.201210801
Jones, M. V., Jonas, P., Sahara, Y., & Westbrook, G. L. (2001). Microscopic kinetics and energetics
distinguish GABA(A) receptor agonists from antagonists. *Biophys J*, *81*(5), 2660-2670.
https://doi.org/10.1016/S0006-3495(01)75909-7

Klabunde, T., Petrassi, H. M., Oza, V. B., Raman, P., Kelly, J. W., & Sacchettini, J. C. (2000). Rational
design of potent human transthyretin amyloid disease inhibitors. *Nat Struct Biol*, *7*(4), 312-321.
https://doi.org/10.1038/74082
Klei, H. E., Kish, K., Lin, P. F., Guo, Q., Friborg, J., Rose, R. E., Zhang, Y., Goldfarb, V., Langley, D. R.,
Wittekind, M., & Sheriff, S. (2007). X-ray crystal structures of human immunodeficiency virus
type 1 protease mutants complexed with atazanavir. *J Virol*, *81*(17), 9525-9535.
https://doi.org/10.1128/JVI.02503-05
Kollman, P. A., Massova, I., Reyes, C., Kuhn, B., Huo, S., Chong, L., Lee, M., Lee, T., Duan, Y., Wang, W.,
Donini, O., Cieplak, P., Srinivasan, J., Case, D. A., & Cheatham, T. E., 3rd. (2000). Calculating
structures and free energies of complex molecules: combining molecular mechanics and
continuum models. *Acc Chem Res*, *33*(12), 889-897. https://doi.org/10.1021/ar000033j
Ma, W., Tang, C., & Lai, L. (2005). Specificity of trypsin and chymotrypsin: loop-motion-controlled
dynamic correlation as a determinant. *Biophys J*, *89*(2), 1183-1193.
https://doi.org/10.1529/biophysj.104.057158
Mittal, L., Srivastava, M., Kumari, A., Tonk, R. K., Awasthi, A., & Asthana, S. (2021). Interplay among
Structural Stability, Plasticity, and Energetics Determined by Conformational Attuning of
Flexible Loops in PD-1. *Journal of Chemical Information and Modeling*, *61*(1), 358-384.
https://doi.org/10.1021/acs.jcim.0c01080
Mittal, L., Tonk, R. K., Awasthi, A., & Asthana, S. (2021). Targeting cryptic-orthosteric site of PD-L1 for
inhibitor identification using structure-guided approach. *Arch Biochem Biophys*, *713*, 109059.
https://doi.org/10.1016/j.abb.2021.109059
Mortensen, M., Ebert, B., Wafford, K., & Smart, T. G. (2010). Distinct activities of GABA agonists at
synaptic- and extrasynaptic-type GABAA receptors. *J Physiol*, *588*(Pt 8), 1251-1268.
https://doi.org/10.1113/jphysiol.2009.182444
Mukhtasimova, N., Free, C., & Sine, S. M. (2005). Initial coupling of binding to gating mediated by
conserved residues in the muscle nicotinic receptor. *J Gen Physiol*, *126*(1), 23-39.
https://doi.org/10.1085/jgp.200509283
Munro, L., Ladefoged, L. K., Padmanathan, V., Andersen, S., Schiott, B., & Kristensen, A. S. (2019).
Conformational Changes in the 5-HT(3A) Receptor Extracellular Domain Measured by Voltage-
Clamp Fluorometry. *Molecular Pharmacology*, *96*(6), 720-734.
https://doi.org/10.1124/mol.119.116657
Muroi, Y., Theusch, C. M., Czajkowski, C., & Jackson, M. B. (2009). Distinct structural changes in the
GABAA receptor elicited by pentobarbital and GABA. *Biophys J*, *96*(2), 499-509.
https://doi.org/10.1016/j.bpj.2008.09.037
Nayak, T. K., & Auerbach, A. (2013). Asymmetric transmitter binding sites of fetal muscle acetylcholine
receptors shape their synaptic response. *Proc Natl Acad Sci U S A*, *110*(33), 13654-13659.
https://doi.org/10.1073/pnas.1308247110
Nayak, T. K., & Auerbach, A. (2017). Cyclic activation of endplate acetylcholine receptors. *Proc Natl*
Acad Sci U S A, *114*(45), 11914-11919. https://doi.org/10.1073/pnas.1711228114
Nayak, T. K., Chakraborty, S., Zheng, W. J., & Auerbach, A. (2016). Structural correlates of affinity in
fetal versus adult endplate nicotinic receptors. *Nature Communications*, *7*.
https://doi.org/ARTN 11352
10.1038/ncomms11352

Nayak, T. K., Purohit, P. G., & Auerbach, A. (2012). The intrinsic energy of the gating isomerization of a
neuromuscular acetylcholine receptor channel. *J Gen Physiol, 139*(5), 349-358.
https://doi.org/10.1085/jgp.201110752
Nayak, T. K., Vij, R., Bruhova, I., Shandilya, J., & Auerbach, A. (2019). Efficiency measures the
conversion of agonist binding energy into receptor conformational change. *J Gen Physiol,*
151(4), 465-477. https://doi.org/10.1085/jgp.201812215
Nickbarg, E. B., Davenport, R. C., Petsko, G. A., & Knowles, J. R. (1988). Triosephosphate isomerase:
removal of a putatively electrophilic histidine residue results in a subtle change in catalytic
mechanism. *Biochemistry, 27*(16), 5948-5960. https://doi.org/10.1021/bi00416a019
Noviello, C. M., Gharpure, A., Mukhtasimova, N., Cabuco, R., Baxter, L., Borek, D., Sine, S. M., & Hibbs,
R. E. (2021). Structure and gating mechanism of the alpha7 nicotinic acetylcholine receptor.
Cell, 184(8), 2121-2134 e2113. https://doi.org/10.1016/j.cell.2021.02.049
Oostenbrink, C., Villa, A., Mark, A. E., & van Gunsteren, W. F. (2004). A biomolecular force field based
on the free enthalpy of hydration and solvation: the GROMOS force-field parameter sets 53A5
and 53A6. *J Comput Chem, 25*(13), 1656-1676. https://doi.org/10.1002/jcc.20090
Padgett, C. L., Hanek, A. P., Lester, H. A., Dougherty, D. A., & Lummis, S. C. (2007). Unnatural amino
acid mutagenesis of the GABA(A) receptor binding site residues reveals a novel cation-pi
interaction between GABA and beta 2Tyr97. *Journal of Neuroscience, 27*(4), 886-892.
https://doi.org/10.1523/JNEUROSCI.4791-06.2007
Pettersen, E. F., Goddard, T. D., Huang, C. C., Meng, E. C., Couch, G. S., Croll, T. I., Morris, J. H., & Ferrin,
T. E. (2021). UCSF ChimeraX: Structure visualization for researchers, educators, and developers.
Protein Sci, 30(1), 70-82. https://doi.org/10.1002/pro.3943
Phillips, R. (2020). *The Molecular Switch: Signaling and Allostery*. Prinston University Press.
Plapp, B. V. (2010). Conformational changes and catalysis by alcohol dehydrogenase. *Arch Biochem*
Biophys, 493(1), 3-12. https://doi.org/10.1016/j.abb.2009.07.001
Pless, S. A., Hanek, A. P., Price, K. L., Lynch, J. W., Lester, H. A., Dougherty, D. A., & Lummis, S. C. (2011).
A cation-pi interaction at a phenylalanine residue in the glycine receptor binding site is
conserved for different agonists. *Molecular Pharmacology, 79*(4), 742-748.
https://doi.org/10.1124/mol.110.069583
Pless, S. A., & Lynch, J. W. (2009). Ligand-specific conformational changes in the alpha1 glycine
receptor ligand-binding domain. *J Biol Chem, 284*(23), 15847-15856.
https://doi.org/10.1074/jbc.M809343200
Polovinkin, L., Hassaine, G., Perot, J., Neumann, E., Jensen, A. A., Lefebvre, S. N., Corringer, P. J.,
Neyton, J., Chipot, C., Dehez, F., Schoehn, G., & Nury, H. (2018). Conformational transitions of
the serotonin 5-HT(3) receptor. *Nature, 563*(7730), 275-279. https://doi.org/10.1038/s41586-
018-0672-3
Purohit, P., & Auerbach, A. (2009). Unliganded gating of acetylcholine receptor channels. *Proc Natl*
Acad Sci U S A, 106(1), 115-120. https://doi.org/10.1073/pnas.0809272106
Purohit, P., & Auerbach, A. (2013). Loop C and the mechanism of acetylcholine receptor-channel
gating. *J Gen Physiol, 141*(4), 467-478. https://doi.org/10.1085/jgp.201210946
Purohit, P., Bruhova, I., Gupta, S., & Auerbach, A. (2014). Catch-and-Hold Activation of Muscle
Acetylcholine Receptors Having Transmitter Binding Site Mutations. *Biophysical Journal, 107*(1),
88-99. https://doi.org/10.1016/j.bpj.2014.04.057

859 Purohit, P., Gupta, S., Jadey, S., & Auerbach, A. (2013). Functional anatomy of an allosteric protein.
860   *Nature Communications*, *4*, 2984. https://doi.org/10.1038/ncomms3984

861 Purushotham, N., Singh, M., Paramesha, B., Kumar, V., Wakode, S., Banerjee, S. K., Poojary, B., &
862   Asthana, S. (2022). Design and synthesis of amino acid derivatives of substituted
863   benzimidazoles and pyrazoles as Sirt1 inhibitors. *RSC Adv*, *12*(7), 3809-3827.
864   https://doi.org/10.1039/d1ra06149f

865 Rahman, M. M., Basta, T., Teng, J., Lee, M., Worrell, B. T., Stowell, M. H. B., & Hibbs, R. E. (2022).
866   Structural mechanism of muscle nicotinic receptor desensitization and block by curare. *Nat
867   Struct Mol Biol*, *29*(4), 386-394. https://doi.org/10.1038/s41594-022-00737-3

868 Rahman, M. M., Teng, J., Worrell, B. T., Noviello, C. M., Lee, M., Karlin, A., Stowell, M. H. B., & Hibbs, R.
869   E. (2020). Structure of the Native Muscle-type Nicotinic Receptor and Inhibition by Snake
870   Venom Toxins. *Neuron*, *106*(6), 952-962 e955. https://doi.org/10.1016/j.neuron.2020.03.012

871 Rasmussen, S. G., DeVree, B. T., Zou, Y., Kruse, A. C., Chung, K. Y., Kobilka, T. S., Thian, F. S., Chae, P. S.,
872   Pardon, E., Calinski, D., Mathiesen, J. M., Shah, S. T., Lyons, J. A., Caffrey, M., Gellman, S. H.,
873   Steyaert, J., Skiniotis, G., Weis, W. I., Sunahara, R. K., & Kobilka, B. K. (2011). Crystal structure of
874   the beta2 adrenergic receptor-Gs protein complex. *Nature*, *477*(7366), 549-555.
875   https://doi.org/10.1038/nature10361

876 Rastelli, G., Del Rio, A., Degliesposti, G., & Sgobba, M. (2010). Fast and accurate predictions of binding
877   free energies using MM-PBSA and MM-GBSA. *J Comput Chem*, *31*(4), 797-810.
878   https://doi.org/10.1002/jcc.21372

879 Richard, J. P. (2022). Enabling Role of Ligand-Driven Conformational Changes in Enzyme Evolution.
880   *Biochemistry*, *61*(15), 1533-1542. https://doi.org/10.1021/acs.biochem.2c00178

881 Roe, D. R., & Cheatham, T. E., 3rd. (2013). PTRAJ and CPPTRAJ: Software for Processing and Analysis of
882   Molecular Dynamics Trajectory Data. *J Chem Theory Comput*, *9*(7), 3084-3095.
883   https://doi.org/10.1021/ct400341p

884 Sauguet, L., Shahsavar, A., Poitevin, F., Huon, C., Menny, A., Nemecz, A., Haouz, A., Changeux, J. P.,
885   Corringer, P. J., & Delarue, M. (2014). Crystal structures of a pentameric ligand-gated ion
886   channel provide a mechanism for activation. *Proc Natl Acad Sci U S A*, *111*(3), 966-971.
887   https://doi.org/10.1073/pnas.1314997111

888 Shang, Y., Yeatman, H. R., Provasi, D., Alt, A., Christopoulos, A., Canals, M., & Filizola, M. (2016).
889   Proposed Mode of Binding and Action of Positive Allosteric Modulators at Opioid Receptors.
890   *ACS Chem Biol*, *11*(5), 1220-1229. https://doi.org/10.1021/acschembio.5b00712

891 Sigel, E., & Steinmann, M. E. (2012). Structure, function, and modulation of GABA(A) receptors. *J Biol
892   Chem*, *287*(48), 40224-40231. https://doi.org/10.1074/jbc.R112.386664

893 Singh, M., Srivastava, M., Wakode, S. R., & Asthana, S. (2021). Elucidation of Structural Determinants
894   Delineates the Residues Playing Key Roles in Differential Dynamics and Selective Inhibition of
895   Sirt1-3. *Journal of Chemical Information and Modeling*, *61*(3), 1105-1124.
896   https://doi.org/10.1021/acs.jcim.0c01193

897 Srivastava, M., Mittal, L., Kumari, A., Agrahari, A. K., Singh, M., Mathur, R., & Asthana, S. (2022).
898   Characterizing (un)binding mechanism of USP7 inhibitors to unravel the cause of enhanced
899   binding potencies at allosteric checkpoint. *Protein Sci*, *31*(9), e4398.
900   https://doi.org/10.1002/pro.4398

901 Sun, H., Li, Y., Tian, S., Xu, L., & Hou, T. (2014). Assessing the performance of MM/PBSA and MM/GBSA
902   methods. 4. Accuracies of MM/PBSA and MM/GBSA methodologies evaluated by various

simulation protocols using PDBbind data set. *Phys Chem Chem Phys*, *16*(31), 16719-16729.
https://doi.org/10.1039/c4cp01388c
Taly, A., Corringer, P. J., Guedin, D., Lestage, P., & Changeux, J. P. (2009). Nicotinic receptors: allosteric
transitions and therapeutic targets in the nervous system. *Nature Reviews Drug Discovery*, *8*(9),
733-750. https://doi.org/10.1038/nrd2927
Tarvin, R. D., Borghese, C. M., Sachs, W., Santos, J. C., Lu, Y., O'Connell, L. A., Cannatella, D. C., Harris, R.
A., & Zakon, H. H. (2017). Interacting amino acid replacements allow poison frogs to evolve
epibatidine resistance. *Science*, *357*(6357), 1261-1266.
https://doi.org/10.1126/science.aan5061
Tian, C., Kasavajhala, K., Belfon, K. A. A., Raguette, L., Huang, H., Migues, A. N., Bickel, J., Wang, Y.,
Pincay, J., Wu, Q., & Simmerling, C. (2020). ff19SB: Amino-Acid-Specific Protein Backbone
Parameters Trained against Quantum Mechanics Energy Surfaces in Solution. *J Chem Theory*
Comput, *16*(1), 528-552. https://doi.org/10.1021/acs.jctc.9b00591
Tripathy, S., Zheng, W., & Auerbach, A. (2019). A single molecular distance predicts agonist binding
energy in nicotinic receptors. *J Gen Physiol*, *151*(4), 452-464.
https://doi.org/10.1085/jgp.201812212
Unwin, N. (1993). Nicotinic acetylcholine receptor at 9 A resolution. *J Mol Biol*, *229*(4), 1101-1124.
https://doi.org/10.1006/jmbi.1993.1107
Vassetti, D., Pagliai, M., & Procacci, P. (2019). Assessment of GAFF2 and OPLS-AA General Force Fields
in Combination with the Water Models TIP3P, SPCE, and OPC3 for the Solvation Free Energy of
Druglike Organic Molecules. *J Chem Theory Comput*, *15*(3), 1983-1995.
https://doi.org/10.1021/acs.jctc.8b01039
Vij, R., Purohit, P., & Auerbach, A. (2015). Modal affinities of endplate acetylcholine receptors caused
by loop C mutations. *J Gen Physiol*, *146*(5), 375-386. https://doi.org/10.1085/jgp.201511503
Wagner, J. R., Sorensen, J., Hensley, N., Wong, C., Zhu, C., Perison, T., & Amaro, R. E. (2017). POVME
3.0: Software for Mapping Binding Pocket Flexibility. *J Chem Theory Comput*, *13*(9), 4584-4592.
https://doi.org/10.1021/acs.jctc.7b00500
Walsh, R. M., Jr., Roh, S. H., Gharpure, A., Morales-Perez, C. L., Teng, J., & Hibbs, R. E. (2018). Structural
principles of distinct assemblies of the human alpha4beta2 nicotinic receptor. *Nature*,
557(7704), 261-265. https://doi.org/10.1038/s41586-018-0081-7
Wlodawer, A., & Erickson, J. W. (1993). Structure-based inhibitors of HIV-1 protease. *Annu Rev*
Biochem, *62*, 543-585. https://doi.org/10.1146/annurev.bi.62.070193.002551
Xiu, X., Puskar, N. L., Shanata, J. A., Lester, H. A., & Dougherty, D. A. (2009). Nicotine binding to brain
receptors requires a strong cation-pi interaction. *Nature*, *458*(7237), 534-537.
https://doi.org/10.1038/nature07768
Yu, A., Salazar, H., Plested, A. J. R., & Lau, A. Y. (2018). Neurotransmitter Funneling Optimizes
Glutamate Receptor Kinetics. *Neuron*, *97*(1), 139-149 e134.
https://doi.org/10.1016/j.neuron.2017.11.024
Yu, R., Hurdiss, E., Greiner, T., Lape, R., Sivilotti, L., & Biggin, P. C. (2014). Agonist and antagonist
binding in human glycine receptors. *Biochemistry*, *53*(38), 6041-6051.
https://doi.org/10.1021/bi500815f
Zarkadas, E., Pebay-Peyroula, E., Thompson, M. J., Schoehn, G., Uchanski, T., Steyaert, J., Chipot, C.,
Dehez, F., Baenziger, J. E., & Nury, H. (2022). Conformational transitions and ligand-binding to a

muscle-type nicotinic acetylcholine receptor. *Neuron, 110*(8), 1358-1370 e1355.
https://doi.org/10.1016/j.neuron.2022.01.013
Zhong, W., Gallivan, J. P., Zhang, Y., Li, L., Lester, H. A., & Dougherty, D. A. (1998). From ab initio
quantum mechanics to molecular neurobiology: a cation-pi binding site in the nicotinic
receptor. *Proc Natl Acad Sci U S A, 95*(21), 12088-12093.
https://doi.org/10.1073/pnas.95.21.12088
952