The Atlantic Forest of South America: spatiotemporal dynamics of remaining

vegetation and implications for conservation

Maurício Humberto Vancine[1,*], Renata L. Muylaert[2], Bernardo Brandão Niebuhr[3], Júlia Emi

de Faria Oshima[4], Vinicius Tonetti[1], Rodrigo Bernardo[1], Carlos De Angelo[5], Marcos Reis

Rosa[6], Carlos Henrique Grohmann[7], Milton Cezar Ribeiro[1,8,*]

[1] Universidade Estadual Paulista (Unesp), Instituto de Biociências, Departamento de Biodiversidade,

Laboratório de Ecologia Espacial e Conservação, Rio Claro, SP, Brazil.

[2] Molecular Epidemiology and Public Health Laboratory, Hopkirk Research Institute, Massey

University, Palmerston North, New Zealand.

[3] Department of Terrestrial Biodiversity, Norwegian Institute for Nature Research (NINA), Trondheim,

Norway.

[4] Universidade de São Paulo (USP), Institute of Biosciences, Movement Ecology Laboratory, Ecology

Department, Rua do Matão, 321, 05508-090, São Paulo, SP, Brazil.

[5] Instituto de Ciencias de la Tierra, Biodiversidad y Ambiente, Río Cuarto, Córdoba, Argentina.

[6] Universidade de São Paulo, Department of Geography, São Paulo, Brazil.

[7] Universidade de São Paulo, Institute of Energy and Environment, Spatial Analysis and Modelling

Lab (SPAMLab), Prof. Luciano Gualberto Avenue, 1289, São Paulo, 05508-010, Brazil.

[8] Universidade Estadual Paulista (Unesp), Environmental Studies Center (CEA), Laboratório de

Ecologia Espacial e Conservação, Rio Claro, SP, Brazil.

* Correspondence

Maurício Humberto Vancine and Milton Cezar Ribeiro, Universidade Estadual Paulista (Unesp),

Institute of Biosciences, Department of Biodiversity, Rio Claro, SP, Brazil. Emails:

mauricio.vancine@gmail.com (MHV); milton.c.ribeiro@unesp.br (MCR).

Highlights

- There is 23% forest and 36% natural vegetation cover remaining in the Atlantic Forest.
- Between 1986-2020, native forest cover decreased by 2.4% and natural vegetation by 3.6%.
- Since 2005, there has been a 1 Mha increase in forest area by small fragments (1 ha).
- Roads and railways reduced by 56%-89% fragment size, especially on large fragments.
- Alarmingly, 97% of fragments are small (<50 ha) and 60% are under edge effect (<90 m).

Abstract

The Atlantic Forest in South America (AF) is one of the world's most diverse and threatened biodiversity hotspots. We present a comprehensive spatiotemporal analysis of 34 years of AF landscape change between 1986-2020. We analyzed landscape metrics of forest vegetation only (FV), forest plus other natural vegetation (NV), and investigated the sensitivity of metrics to linear infrastructure. Currently, remnants comprise about 23% of FV and 36% of NV, and have decreased by 2.4% and 3.6% since 1986, respectively. Linear infrastructure negatively affected large fragments (>500,000 ha) by breaking them apart. Our findings suggest that AF protection legislation adopted in mid-2005 has taken effect: between 1986-2005, there was a loss of FV and NV (3% and 3.45%) and a decrease in the number of FV and NV fragments (8.6% and 8.3%). Between 2005-2020, there was a relative recovery of FV (1 Mha; 0.6%), slight loss of NV (0.25 Mha; 0.15%) and increase in the number of FV and NV fragments (12% and 9%). Still, 97% of the vegetation fragments are small (<50 ha), with an average fragment size between 16 and 26 ha. Furthermore, 50-60% of the vegetation is <90 m from its edges, and the isolation between fragments is high (250-830 m). Alarmingly, protected areas and indigenous territories cover only 10% of the AF and are very far from any fragments (>10 km). Our work highlights the importance of legislation and landscape dynamics analysis to help monitor and keep track of AF biodiversity conservation and restoration programs in the future.

Keywords: Landscape structure; Habitat loss; Habitat fragmentation; Edge effect; Isolation; Connectivity.

1. Introduction

Habitat loss, fragmentation, and degradation caused by human-induced changes are identified as the main drivers of biodiversity loss worldwide (Chase et al., 2020). The accelerated land use conversion resulting from these changes has affected especially forest ecosystems, causing a decrease in fragment size and an increase in edge effects (Fischer et al., 2021; Hansen et al., 2020). In recent decades, tropical and subtropical regions have lost >100 million hectares (Mha) of natural forests due to anthropogenic activities (Zalles et al., 2021). Despite the large impacts, few studies presented a spatiotemporal panorama temporal long enough to describe and analyze the landscape structure dynamics, especially in the Americas, where the most diverse and threatened biodiversity hotspot in the world remains: the Atlantic Forest in South America (AF) (Sloan et al., 2014).

The AF covers almost all the coast of Brazil and portions of Paraguay and Argentina. Before European colonization, its vegetation covered over 1.6 million km^2 (Marques et al., 2021). Due to its high environmental heterogeneity, topographic variability, and pre-historic process of formation, the AF has a high species diversity and endemism (Peres et al., 2020): it hosts more than 18,000 species of plants (Flora e Funga do Brasil, 2023) and 3,500 species of vertebrates (Figueiredo et al., 2021; Reis et al., 2016). In addition, the AF provides ecosystem services for >150 million people, such as water provisioning, hydroelectric energy generation, food production, pollination, soil protection, climate regulation, carbon storage, air quality, and cultural services (Joly et al., 2014).

The intensification of degradation arises with the Portuguese colonization and degradation of agricultural processes such as large plantation systems (sugarcane and coffee), extensive cattle production, energy demand (charcoal), fires, and urban and industrial growth (Solórzano et al., 2021). These habitat transformations have affected the biodiversity in the AF for different taxonomic groups (Püttker et al., 2020) and ecological processes, such as seed dispersal (Marjakangas et al., 2020), carbon storage (de Lima et al., 2020), pollination (Varassin et al., 2021), and top-down regulation through top predators (Paviolo et al. 2016). In addition, other processes pose risks to the remaining landscapes within the AF, such as defaunation (Galetti et al., 2021) and climate change (Vale et al., 2021).

Despite the recent changes on the AF, few studies have analyzed the landscape structure in a space-time context on large time scales. In the most comprehensive study to our knowledge, Ribeiro et al. (2009) showed that only 11-16% of the forest cover remained in 2005, 83% of which was concentrated on isolated fragments smaller than 50 ha, and half of all forests were <100 m from their edges. After that, Tabarelli et al. (2010) and Ribeiro et al. (2011) showed a large proportion of forests remained in high elevations (>1600 m).

Based on finer scale satellite data (5 m-spatial resolution), Rezende et al. (2018) estimated

28% of remaining AF vegetation. In more recent studies, using data from MapBiomas

(Souza et al., 2020), Bicudo da Silva et al. (2020) showed that landscape composition did

not change between 1985-2018, and that the loss in areas of montane vegetation was

smaller than at lower elevations. Rosa et al. (2021) showed that the relative temporal

stability of AF native forest cover (28 Mha) in recent years, was in fact due to the loss of old-

growth native forests in flatter terrains, and the growth of young forests in marginal

agricultural areas, resulting in increased isolation.

Despite these studies, there is a demand for refined data to understand how

landscape structure varied over time in AF. Currently, Brazilian initiatives such as

MapBiomas have been mapping land use and land cover (LULC) change with wide thematic

coverage, high spatiotemporal resolution, and standardized classification (Souza et al.,

2020). This allows for the calculation and comparison of landscape metrics for large

territorial extensions and time periods to understand the landscape dynamics of entire

domains (Bicudo da Silva et al. 2020; Rosa et al. 2021). In addition, the AF has a high

density of linear infrastructure since it hosts a high (and increasing) human population. This

severely impacts natural vegetation and biodiversity and must be considered in landscape

structure analyses (Cassimiro et al., 2023).

Here, we analyzed the spatiotemporal dynamics of the landscape structure of

vegetation in the AF every five years from 1986-2020. To accomplish this large-scale

evaluation, we used a wide delimitation of Atlantic Forest, including Brazil, Argentina, and

Paraguay. We accounted for forest vegetation types only (FV) and forest plus other natural

vegetation types (NV) and quantified the effect of linear infrastructure on the AF landscape

metrics. To understand the spatiotemporal vegetation dynamics, we calculated the following

landscape metrics for all FV and NV fragments in the AF domain: fragment size, number of

fragments, fragment temporal dynamic, habitat amount, edge area, isolation, functional

connectivity, and distance from protected areas (PA) and indigenous territories (IT). These

metrics were generated through an approach that allows an ecological interpretation of the

influence of the landscape structure on organisms, by accounting for species mobility, gap-

crossing abilities, and sensitivity to edge effect (Riva and Nielsen, 2020).

2. Methods

2.1 Study region

AF extends from 3ºS to 33ºS, and from 35ºW to 58ºW with about 163 Mha, covering

large coastal and inland portions of Brazil, Argentina, and Paraguay (Marques et al., 2021)

(Fig. S1a). Due to this extension, the AF boundaries create important ecotones with other
vegetation domains such as Cerrado, Caatinga, Chaco and Pampa (Marques et al., 2021).
The vegetation from AF is a complex mosaic mainly composed of five vegetation types—
Dense Ombrophilous, Open Ombrophilous, Mixed Ombrophilous, Semideciduous Seasonal,
and Deciduous Seasonal (Joly et al., 2014). Additionally, the AF also includes mangroves
and coastal scrub vegetation (Marques et al., 2021). Besides, there are many marginal
habitats such as altitude grasslands (*campos rupestres* and *campos de altitude*), oceanic
islands, beaches, rocky shores, dunes, marshes, inland swamps, and mountain forest
(*brejos de altitude*) in the Northeast region (Scarano, 2002). Therefore, we used an
integrative delimitation adapted from Muylaert et al. (2018), which encompasses the main
proposed delimitations across several associated ecosystems. This delimitation was
produced by overlapping available AF delimitations (Table S1 and Fig. S1[b-e]) and
adjusting the delimitation in the Eastern coastal areas using the Brazilian territorial
delimitation from IBGE (https://www.ibge.gov.br) for 2021. This step ensures that areas of
coastal vegetation such as mangroves, dunes, and wooded sandbank/sandy coastal plain
vegetation (hereafter *restinga*) (Scarano, 2002) are better represented. The final delimitation
has a total area of 162,742,129 ha, distributed within 3653 municipalities from 18 Brazilian
states (93.1%), 70 municipalities of one province in Argentina (1.6%), and 127 municipalities
from 11 departments in Paraguay (5.3%) (Fig. S1a).
208
209   *2.2 Mapping*
210
We compiled LULC maps for Brazil, Argentina, and Paraguay from MapBiomas
Brazil collection 7 (https://mapbiomas.org/) and MapBiomas Bosque Atlántico collection 2
(https://bosqueatlantico.mapbiomas.org/) (Souza et al., 2020). These datasets reconstruct
annual LULC information at 30-m spatial resolution from 1985 to 2021, based on a pixel-
based random forest classifier of Landsat satellite images using Google Earth Engine, with
AF general accuracy of 89.8% (Souza et al., 2020). We used the interval beginning in 1986
and ending in 2020. We excluded the years 1985 and 2021, as there was no validation for
the previous and subsequent year, respectively. Furthermore, we defined two vegetation
classes for analysis: only forest vegetation types or "Forest Vegetation" (FV) and both forest
and other natural vegetation types or "Natural Vegetation" (NV) (Table S2), for every fifth
year between 1986 and 2020 (Fig. S2a-h).
We used roads and railways to trim their overlapping FV and NV (henceforth called
"trimmed" and "not trimmed" scenarios). This procedure enabled us to avoid overestimating
large fragments of vegetation and check the metrics' sensitivity to linear infrastructure, since
these structures decrease landscape connectivity and threaten multiple taxonomic groups

(Cassimiro et al., 2023). Thus, we analyzed four vegetation maps: "FV not trimmed", "FV trimmed", "NV not trimmed", and "NV trimmed". Further, we analyzed the overlap between FV and NV fragments with Protected Areas (PA) and Indigenous Territories (IT). Details of road, railway, PA, and IT maps are presented in the Data section in the Supplementary Material. All geospatial datasets were rasterized and warped to 30 m-spatial resolution (112663 × 83307 ≈ 9.4 billion cells) using the Albers Conical Equal Area Brazil (SIRGAS 2000) projection (https://spatialreference.org/ref/sr-org/albers-conical-equal-area-brazil-sirgas-2000/). International map displays were generated using Natural Earth (1:10,000,000) data and QGIS 3.22 LTR (QGIS Development Team, 2023).

2.3 Landscape metrics

All landscape metrics were processed in GRASS GIS 8.2.1 (Neteler et al., 2012) through the R 4.3.0 (R Core Team, 2023), using the *rgrass* package (Bivand, 2022). We calculated six landscape metrics: number of fragments, fragment size, edge area, isolation, functional connectivity, and distance from PA and IT (Table S3 and Figure S13). The number of fragments and fragment size allowed us to account for the number and area of remaining vegetation fragments for different size classes (Table S3). Fragments were defined using the eight-neighbor rule (Queen's case), which defines areas connected to pixels in eight directions (Turner and Gardner, 2015). We also examined the area and number of fragments that appeared and disappeared throughout time, and the areas of increase, reduction, and stability of fragments that remained in the landscape (Table S3) (Rosa et al., 2021). Edge area was calculated for different edge depths (distance from the edge of the fragment) (Table S3), allowing us to assess the amount and percentage of forest area subjected to edge effects (Harper and Macdonald, 2011).

Two metrics of functional connectivity were computed for different gap-crossing distances (species' capacities to cross the non-habitat) (Table S3). First, we calculated the sum of the areas of all fragments closer than the gap-crossing distance, which can be interpreted as the functional available area of each clump of fragments (Awade and Metzger, 2008). Second, we computed the expected cluster size as the mean fragment clump size, and then compared it with the highest cluster size in the entire study region. Isolation was calculated using an index adapted from the "Empty Space Function" (Dale and Fortin, 2014), similar to Ribeiro et al. (2009): we computed a Euclidean distance map from all the fragments, extracted its values and calculated the mean. We repeated this process by removing different-sized fragments in several steps (see Table S3 for classes of distances), and then created new Euclidean distance maps to recompute the mean distance values. These values represented the isolation of fragments while also providing insights about the

importance of the smaller fragments (*stepping stones*) (Diniz et al., 2021). We calculated the

amount of FV, NV, and vegetation classes (see Table S2) covered by PA and IT, and the

shortest Euclidean distance from each FV and NV pixel to these areas (see Table S3 for

classes of distances).

3. Results

3.1 Number of fragments and fragment size distribution

272   Roads and railways greatly impacted the large-sized fragments, depending on the

year and the scenario considered. These effects were mainly reflected in vegetation

fragments larger than 500,000 hectares, for which the maximum fragment size decreased by

56%-89% (Fig. 1, Fig. S3, and Table S4). By accounting for linear infrastructure, the >1 Mha

fragment size class ceased to exist for FV for all years and was heavily reduced for NV, and

the total area and number of fragments increased for fragments of all size classes <500,000

ha for FV and NV (Fig. 1, Fig. 2, and Fig. S4). Despite this effect for large fragments, our

results showed no difference between the scenarios "trimmed" and "not trimmed" for other

landscape metrics. Therefore, we chose to demonstrate the results with the linear

infrastructure effect (trimmed scenario) in the main text and present the additional results in

the Supplementary Material.

283   For the trimmed scenario, about 97% of the fragments have an area of less than 50

ha, with 0.3% of variation over the years. However, between 1986 and 2020 the total area

increased from 18.8% to 22.1% for FV and from 11.6% to 13.4% for NV (Fig. 1 and Fig. S3).

For fragments between 50 ha and 25,000 ha, the proportion of the total number of fragments

is low (2.5%), varying for FV from 2.44% in 1986 to 2.66% in 2020, with a maximum value of

2.76 % in 2005; and for NV with 2.34% in 1986 to 2.61% in 2020, and a maximum of 2.66%

in 2005. However, total area increased from 1986 to 2020, going from 39.8% to 45.9% for

FV, but very similar since 2005 (45.1%); and for NV, from 29.6% to 35.1% (Fig. 1 and Fig.

S3). For the last category of fragment area, above 25,000 ha, we found a very small

proportion of number of fragments (0.001%), with values falling from 0.0081% to 0.0058%

for FV, and from 0.0125% to 0.0116% for NV, between 1986 and 2020. Total area values for

FV fragments in these categories fell from 41.4% to 32% and for NV from 58.7% to 51.5%,

between 1986 and 2020 (Fig. 1 and Fig. S3).

Figure 1. Distribution of FV and NV fragment sizes across the AF (1986 and 2020), trimmed and not trimmed by linear infrastructure. %A: percentage of the total area; %NF: percentage

of the number of fragments. See Fig. 3S for other years (1990-2015). Please note the

difference scales in the x-axis between the FV and NV plots.

In 1986, the largest FV fragments were localized in the coast of Bahia (*South of*

Bahia—cabruca region), São Paulo, Paraná, and Santa Catarina states (*Serra do Mar*

region), and inland areas of Paraná, Santa Catarina and Rio Grande do Sul states in Brazil.

For the same period, there were large FV fragments in the Misiones region in Argentina and

the east portion of Paraguay (Fig. 2a). We observed the same for NV, with additions of huge

fragments in portions of Bahia, Minas Gerais, and Piauí states, mainly in the regions named

São Francisco and *Brejos Nordestinos* (see these region concepts in Ribeiro et al., 2009)

(Fig. 2c). In 2020, these same regions concentrated the largest fragments of FV and NV, but

with a decrease in the area of these fragments (Fig. 2[b-d]). The exception was Paraguay,

where there was a vast deforestation process, mainly for FV (Fig. 2[c-d]). Our results also

show a large effect of roads and railways on maximum fragment size, for the same regions

(compare Fig. 2[a-d] with Fig. S4[a-d]).

Figure 2. Fragment area for FV in 1986 (a) and 2020 (b), and for NV in 1986 (c) and 2020 (d), trimmed for the entire AF.

The spatial-temporal analysis revealed a turning point for the AF landscape structure in 2005. For the first period (1986-2005), the number of fragments decreased by 8.6% for FV and 8.3% for NV; for the second one (2005-2020), it increased to 11.9% for FV and 9% for NV (Fig. 3a). From 2010 onwards, the number of FV and NV fragments tended to become more like each other (Fig. 3a). The average fragment size in the first period for FV dropped by 3.5% (18.5 to 17.9 ha) and remained stable for NV, dropping 0.3% (28.5 to 28.4%). In the second period, the FV had a great drop of 8.2% for FV (17.9 to 16.4 ha) and 8.6% for NV (28.4 to 25.9 ha) (Fig. 3b).

Figure 3. Distribution of number of fragments (a) and average fragment sizes (b) of FV and

NV across the AF from 1986 to 2020, trimmed and not trimmed by roads and railways. The

gray lines represent when the legislation and restoration programs were created (See

Discussion for details).

The temporal dynamics of the landscape (Table S3) from 1986 to 2005 revealed a

reduction in the total area of 4.78 Mha of FV (3%) and 5.56 Mha of NV (3.4%) (Table S5).

However, between 2005 and 2020, there was an increase of 985,000 ha of FV (0.6%) and a

small decrease of 240,000 ha of NV (0.15%). Considering the balance of fragments gained

and lost, in the first period there was a sharp drop in the number of fragments for FV

(242,000) and NV (227,000), but in the second period there was an increase for FV

(380,000) and NV (310,000). Between 1986-2005, the average size of lost FV and NV

fragments (1.2 to 1.35 ha) was greater than the size of restored fragments (1.08 to 1.14 ha);

between 2005-2020, this pattern reversed, with the average size of fragments lost was

smaller (0.94 to 0.97 ha) than that of fragments gained (1.03 to 1.08 ha).

3.2 Forest and natural vegetation cover

The proportion of the Atlantic Forest domain covered by forests and natural

vegetation decreased in the past 34 years, from 25.26% (41.1 Mha) to 22.86% (37.2 Mha)

for FV and from 39.86% (64.8 Mha) to 36.27% (59 Mha) for NV (Fig. 4 and Table S4). For

the entire period, in Brazil the percentages decreased from 22.85% (34.6 Mha) to 22.27%

(33.7 Mha) for FV and from 37.34% (56.5 Mha) to 35.25% (53.4 Mha) for NV, with a stable

proportion since 2005 (Fig. S5). NV was mainly composed of savannas, grasslands, and

wetlands, besides the forest formations. In Argentina, the loss of vegetation cover was

proportionately larger, from 67.38% (1.8 Mha) to 56.9% (1.52 Mha) for FV, and 67.99% (1.82

Mha) to 57.34% (1.53 Mha) for NV, showing an increase in the rate of deforestation in the

last five years (Fig. S5). In Paraguay, the loss of vegetation cover was higher than in the other countries, dropping from 54.57% (4.7 Mha) to 22.85% (2 Mha) for FV, and 75.26% (6.5 Mha) to 47.74% (4.1 Mha) for NV (Fig. S5), but it has maintained its remnants since 2005.

Figure 4. Vegetation cover for FV and NV through the years, trimmed. Abbreviations in Table S2.

Beyond presenting the percentages for the integrative delimitation (Fig. S1a), we also present results for five other delimitations (Table S6). The results for 2020 (Fig. S1[b-f]) varied for FV from 23.15% (31.6 Mha) for the delimitation of Da Silva and Casteleti (2003) trimmed up to 26.70% (32.2 Mha) for the delimitation of Dinerstein et al. (2017) disregarding the effect of roads and railways. The same occurred for NV, ranging from 31.45% (34.8 Mha) for the IBGE (2019) trimmed to 35.98% (46.3 Mha) for the delimitation of the Atlantic Forest Law (2006) not trimmed (Table S6).

3.3 Core and edge area

The percentage of FV and NV remaining less than 90 m from the edge increased over time, going from 52% to 59% for FV and 42% to 48% for NV, as well as the percentage less than 240 m, from 76% to 82% for FV and 66% to 72% for NV (Fig. 5[a-b]). Conversely, the amount of FV and NV more than 500 m from any edge decreases, from 12% to 9% and from 20% to 15%, respectively. The maximum edge distances for FV and NV were quite different, being around 11 km for the FV, and 32 km for the latter, showing that NV creates large core areas (Fig. 5[a-b]). From 90 m onwards, there is an inversion in the edge percentage over time: <90 m, there is a gradual increase in the percentage between 1986 and 2020; >90 m, the percentage of vegetation starts to decrease, showing the conversion of fragment core areas to edge areas (Fig. 5[c-d]).

Figure 5. Cumulative (a and b) and per class (c and d) area under edge effect at different

depths for the FV and NV remaining in AF trimmed.

3.4 Functional connectivity

Considering the functional connectivity for species that cannot cross areas of non-habitat (i.e., gap-crossing equals 0 m), the average functionally connected area for FV decreased 11.7% (18.42 to 16.26 ha) (Fig. 6a), and 9.3% for NV (28.2 to 25.5 ha) between 1986 and 2020 (Fig. 6b). The same pattern occurs for 60 m of gap-crossing for both types of vegetation. However, for gap-crossing values between 120 and 180 m, functional connectivity decreased until 2005 and then increased. For values above 240 m, functional connectivity also decreased until 2005, then increased, with its value in 2020 greater than in 1986. The functional connectivity of the NV was always higher in numerical terms for the same years, but they followed the same patterns of annual trends and gap-crossing of the FV.

When we analyzed the highest functionally connected vegetation cluster, we noticed

similarities in the pattern of the curves generated for FV and NV: both curves started with a

low relation between the highest functionally connected FV and NV cluster (about 2% for FV

and 4% for NV for all years), indicating low connectivity when we do not consider a value of

gap-crossing (Fig. 6[c-d]). Despite that, around 600 m of gap-crossing, both reached their

respective horizontal asymptotes, which were about 90% for FV and 75% for NV for every

year, with a significant divergence between years between 60 and 600 m cross-level

difference values, ranging from 18% to 75%. For the not trimmed scenario, there was a

difference in the highest functionally connected initial percentages due to the different values

of the largest fragments over the years (Fig. S7[c-d]).

Figure 6. Expected cluster size (a-b) (average functional size; ha) of functionally connected

fragments of FV and NV for different functional distance values (meters) for the AF trimmed

by roads and railways. Highest functionally connected vegetation cluster (c-d) (% of total

remaining of FV and NV) estimated across varying functional distances (meters) for the AF.

 3.5 Mean isolation

Small fragments were key to reducing isolation in all analyzed scenarios. For

example, when we disregard fragments <50 ha, the isolation increases to 79-83% for FV

and 78-85% NV (Fig. 7[a-b]). Furthermore, isolation was highly reduced in 60-85% when

considering NV also for all temporal scenarios and road and rail effects analyzed (Fig. 7[a-b],

Fig. S8[a-b]). In 1986, the mean isolation for the entire AF region was 773 m for FV and 273

m for NV. The isolation reached its maximum values in 1995, with values of 949 m and 291

m for FV and NV, respectively. After that, the isolation had a slow decrease until 2015, going

to 902 m for FV and 266 m for FV; and more recently, it fell to 832 m for FV and 253 m for

NV in 2020 (Fig. 7[a-b]). When we disregarded fragments smaller than the size classes

defined in Table S3, there was a significant increase in isolation, a gradual increase for each

size that we disregarded, varying to 4-22 km for FV and 1-12 km for NV. In general, isolation

peaks for each class fragment size disregarded occurred between 1990 and 2000 for FV

and NV, which after these dates began to decrease, reaching the lowest values of the

historical series in 2020, both for FV and for NV.

Figure 7. Influence of the smallest fragment size (ha) on the isolation (m) between

fragments of FV and NV trimmed for the AF. Smallest fragments size: 0 ha (all fragments),

50 ha, 100 ha, 150 ha, 200 ha, 250 ha, 350 ha, 500 ha, and 1000 ha.

3.6 Protected areas and indigenous territories distance

Protected areas (PA) covered 4.6 Mha (2.84%) and indigenous territories (IT)

covered 1.3 Mha (0.81%) of the AF limit. These values represent 12.4% and 7.8% of the

total FV and NV area for PA, and 3.6% and 2.2% of the total FV and NV area for IT in 2020.

However, only 3.1 Mha (8.4%) of FV and 4.1 Mha (7%) of NV remaining overlaps with PA

(Fig. 8[a-b]), and only 0.56 Mha (1.5%) of FV and 0.76 Mha (1.3%) of NV overlaps with IT
(Fig. 8[c-d]), since other types of land cover occur within PA and IT. Only 2.7% of FV and
2.2% of NV are within 1 km of PA and 0.8% of FV and 0.7% of NV to IT. For vegetation
within 10 km, there are 23.4% of FV and 19.2% of NV of PA, and 9.5% of FV and 8.7% of
NV of IT (Fig. 8). On the other hand, 68.2% of the FV and 73.9% of NV are over 10 km away
from PA, and 89% of the FV and 90.2% of NV are over 10 km away from IT, demonstrating
the lack of protection for these fragments of remaining vegetation (Fig. 8).

The vegetation class with the largest area overlap is the Forest formation, with 3 Mha
(8.2%) for PA and 536,000 ha (1.5%) for IT (Fig. S10). The class with the largest cover
under protection is the *restinga* with 133,000 ha (21.6%), Mangrove with 23,000 ha (11.9%),
and Herbaceous sandbank vegetation with 31,000 ha (9.6%) (Fig. S10b). For IT, the class
with the most cover proportion is the Other non-forest formations 1,400 ha (8.6%), followed
by *restinga* with 22,000 ha (3.6%), and Mangrove with 4,600 ha (2.4%) (Fig. S10d). The
Savanna formation class is the smallest cover protection with only 507,000 ha (3.8%) in PA
(Fig. S10b) and 120,000 ha (0.9%) in IT (Fig. S10d), whose total area is second in terms of
total area, 13 Mha (22.5%).

Figure 8. Remaining vegetation of AF remnants (area and percentage) and their distance
(meters) from protected areas (PA; a – FV and b – NV) and indigenous territories (IT; c – FV
and d – NV) per class, trimmed.

4. Discussion

4.1 Main results

Our results showed great changes in the spatial-temporal dynamics of the landscape
structure for the entire AF, emphasizing the likely great importance of environmental
legislation for its conservation. These changes varied depending on the vegetation types and
roads and railways scenarios considered. For example, fragment size was highly sensitive to
linear structures, especially for large fragments of FV and NV (>500,000 ha), but other
metrics were not affected as percentage, connectivity, or isolation. In 34-year period, there
was a high loss rate for both FV and NV, with apparent differences in trends between Brazil,
Argentina, and Paraguay. Our findings revealed yet a turning point for the Atlantic Forest
after 2005. In the first period (1986-2005), the number of vegetation fragments and forest
cover decreased, following the strong trend of forest loss from the previous years. In the
second period (2005-2020), the number of vegetation fragments increased, and the forest
cover was more stable. Nevertheless, more than 97% of fragments are smaller than 50 ha,
and mean fragment size decreased being currently equal to 16 ha for FV and 26 ha for NV.
Besides, there was an increase in the amount of fragment closest to the <90 m from edge
(50-60%), although the insulation is decreasing, reaching values equal to 830 m for FV and
250 m for NV, with the connectivity increasing.
These results bring a new panorama to the remaining AF vegetation, mainly because
our analysis was more spatially and temporally comprehensive, integrated different types of
vegetation, and considered for the first time a wide distribution of AF, including Argentina
and Paraguay. Although legislation and restoration actions appear to be positively affecting
vegetation restoration processes, there is still a scenario of intense habitat fragmentation.
Allied to this, NV played a fundamental role in increasing the connectivity of the vegetation
remnants, despite having been little favored by protection areas. Our results also showed
that although there is an overlap of about 10% of the vegetation with PA and IT, a large part
of the vegetation is located far from these areas (>10 km) and given the great anthropic
concentration in the AF and recent urban expansion, a portion significant amount of
vegetation could be impacted in the recent future.

4.2 Number and fragment size distribution

Our analyses of temporal dynamics demonstrated that between 2005-2020 there was

an increase in habitat amount, number of fragments, and higher mean fragment gain for FV.

This corroborates that there was a regeneration process since new fragments appeared,

which confirms the results found by Rosa et al. (2021) and Dias et al. (2023), who

highlighted the replacement of older vegetation by younger vegetation in the AF. However,

this replacement can lead to the loss of quality of habitat fragments, altering landscape

features, and affecting vital ecological processes and ecosystem functioning, such as carbon

cycling (Piffer et al., 2022) and vegetation structure (Faria et al., 2023). The effect of roads

and railways was more pronounced in FV than in NV, due to their greater density in large

forest fragments located in Serra do Mar, southern Bahia in Brazil, and in the region of

Misiones in Argentina. Roads and railways have a huge impact on biodiversity, modifying the

movement pattern, reducing connectivity, and causing roadkill, which leads to population

declines and local extinction (Cassimiro et al., 2023; Martinez Pardo et al., 2023).

When analyzing the distribution of fragment sizes, we noticed a reduction in the

number and percentage in relation to the total area of the remnants for fragments >1 Mha of

FV and NV between 1986 and 2020 and a clear increase in fragments <50 ha. These results

showed a worrying pattern, much worse than Ribeiro et al. (2009), and can be explained by

the increase in mapping quality, with MapBiomas standardized approaches for mapping

vegetation fragments (including fragments <3 ha) that are considered secondary vegetation

in detail (Rosa et al., 2021). The increase in the proportion of smaller fragments has a direct

impact on the maintenance of species diversity and population size of multiple taxonomic

groups. Several works have estimated fragment size and habitat amount thresholds for

assemblage diversity in AF, such as terrestrial mammals (Magioli et al., 2015), bats

(Muylaert et al., 2016), birds (Barbosa et al., 2017), and multiple groups (Banks-Leite et al.,

2014). However, since 97% of the fragments are <50 ha in AF, the general scenario is

already well under the thresholds that are known to affect biodiversity composition. Then,

approaches for conservation should be comprehensive and focus on single large *and*

several small (SLASS) fragments (Szangolies et al., 2022). The SLASS approach can be

more beneficial for conserving the AF biodiversity than choosing a unique type of

conservation approach (SLOSS debate, see Fahrig et al. 2021).

4.3 Forest and natural vegetation cover

Determining how much of the AF vegetation cover is left has always been a complex

task. While we found 22.86% for FV and 36.27% for NV using an integrative AF delimitation

in 2020, for the same year, the vegetation cover values varied from 23.15% to 35.98% for different delimitations and vegetation types. Here, we highlighted the use of MapBiomas mapping version 7, with image standards and classification methods, making the comparison of annual maps possible due to the decrease in random error between them (Souza et al., 2020). Over the years, several studies have shown values ranging from 8% to 28% for different years, AF delimitation, and vegetation types (Bicudo da Silva et al., 2020; Da Silva and Casteleti, 2003; Rezende et al., 2018; Ribeiro et al., 2009). These estimates vary according to mapping resolution, size of vegetation fragments, types of vegetation (forest or non-forest), vegetation quality (primary or secondary forest), and AF delimitation (Ribeiro et al., 2009). Therefore, we emphasize that these percentage values must be used with awareness of their calculation specificities and limited comparability. Due to their simplification and high variability intrinsic to each source, their values must be presented to meet detailed and well-defined objectives, meeting specific criteria.

The vegetation cover showed a considerable decrease over time, mainly between 1986 and 2005. After 2005, the percentage of vegetation stabilized or increased, mainly in Brazil and for FV. These effects can be related to nature conservation laws, which were initiated almost in the same period in Brazil (Atlantic Forest Law in 2006, and Native Vegetation Protection Law in 2012), Argentina (Forest Law in 2007), and Paraguay (The Zero Deforestation Law in 2004) (Silva et al., 2017; Dam et al., 2019). In Brazil, specific conservation laws were established from 1998 on (Fauna Protection in 1988, and National System of Conservation Units (SNUC) in 2000), and more recently the 2012 legislation created the Rural Environmental Registry (CAR), which requires environmental information from private rural properties. CAR can be a fundamental tool to direct vegetation restoration efforts through legal reserves (LR) and permanent preservation areas (PPA) (da Silva et al., 2023). Since 2009, the Pact for the Restoration of the Atlantic Forest (https://pactomataaatlantica.org.br) has been encouraging the restoration with the goal to restore 15 Mha by 2050 (Melo et al., 2013), with about 700,000 ha forest restored between 2011 and 2015 (Crouzeilles et al., 2019). Yet, Bicudo da Silva et al. (2023) showed that between 2001-2015 there was a process called "forest transition" (declines in forest cover cease and recoveries in forest cover begin) (Rudel et al., 2005), due to the stagnation of agricultural activities, the emergence of non-agricultural rural activities, and the decrease in precipitation leading to soil abandonment and favoring regeneration.

In Argentina, the percentage of forest has been reduced linearly since the 1990s, with the combined effect of the advance of small-scale agriculture associated with population growth and road construction in some areas, and the increase of monospecific forest plantations incentivized by government subsidies and the participation of large timber companies (Izquierdo et al., 2008). The forest loss rate was lower during 2005-2015,

potentially because of the effect of the certified wood market in this region and the approval
of the National Forest Law and the implementation of the National Fund for the Enrichment
and Conservation of Native Forests (FVSA & WWF, 2017). However, forest loss increased in
the last period (2015-2020) most likely due to higher levels of economic growth and the
impact of long-term police on the expansion of agriculture and cattle raising in this province
(Mohebalian et al., 2022). Paraguay showed the highest rates of deforestation of the entire
Atlantic Forest between 1986-2005 due to the massive expansion of agriculture. However,
since the creation of the Zero Deforestation Law and the implementation of associated
mechanisms, there has been a recent stabilization of vegetation loss (Da Ponte et al., 2017;
FVSA & WWF, 2017).
594
595 *4.4 Core and edge area*
596
Our results showed that 50% of the remaining vegetation is under the effect of a 90
m edge, and about 75% is under the effect of a 240 m edge and almost 90% is under the
effect of a 500 m edge, results very similar to Haddad et al. (2015). Over time, there was an
increase in vegetation located less than 90 m from the edges, revealing a pronounced edge
effect threshold in the AF. Below this threshold, there is an intensification of edge effects,
and above it, there is a decrease in the amount of vegetation core. This threshold is probably
associated with the massive number and small average size of fragments we detected.
Importantly, small fragments are more subject to edge effects due to their size and shape
(Fahrig, 2003). The edge effect changes the AF landscape features such as microclimate
and carbon cycle (Magnago et al., 2017, 2015) depending on the fragment shape (Banks-
Leite et al., 2013) and the matrix effect (Adorno et al., 2021). In that regard, numerous
studies have demonstrated the negative effects of edge changes for epiphyte plants, small
mammalian, and birds in the AF (de la Sancha et al., 2023; Morante-Filho et al., 2018;
Parra-Sanchez and Banks-Leite, 2020). Added to that, Pivello et al. (2021) identified that AF
is highly fire-sensitive, which changes the conditions of the edges and vice versa. Some
measures such as forested or agroforestry matrices and strips of trees being planted,
forming a buffer around the remaining fragments can reduce the edge effect (Gama-
Rodrigues et al., 2021; Tavares et al., 2019).
615
616 *4.5 Functional connectivity and mean isolation*
617
Functional connectivity and isolation had similar response patterns over time, with
their worst values between 1990 and 2000, but from 2005 onwards there were clear signs of
improvement. The vegetation amount has not changed noticeably since 2005, and this

improvement was due to the appearance of new fragments that increased the connectivity of
the landscape, probably through stepping stones. In this way, small fragments (<50 ha,
which represents 97% of AF remnants) play a fundamental role in keeping large fragments
connected, even more important for species that can cross the matrix (Diniz et al., 2021).
Furthermore, NV plays a key role in decreasing the isolation of the remnants, although there
may be fewer forest-specialist species that use this type of vegetation, it can be critical to
maintaining AF connectivity (Lyra-Jorge et al., 2010).
However, practices such as agroecology and forestry can increase the connectivity
by increasing the permeability of the matrix (Tubenchlak et al., 2021). In addition, The
Atlantic Forest Restoration Pact and Rural Environment Registry (CAR) police are a great
opportunity to create and improve ecological corridors (da Silva et al., 2023; Melo et al.,
2013). Finally, although connectivity and isolation were not apparently sensitive to the roads
and railways effect, this lack of sensitivity may be due to short-distance divisions into FV or
NV fragments, as the additional cost that these linear structures cause, preventing animals
from crossing short distances (Martinez Pardo et al., 2023), were not considered. Thus, it is
essential to propose fauna passages for improving landscape permeability to maintain
wildlife gene flow and reduce roadkill (Cassimiro et al., 2023; Teixeira et al., 2022;
Zimmermann Teixeira et al., 2017).
639
640 *4.6 Protected areas and indigenous territories*
641
Alarmingly, our results showed that the proportion of PA (10% for FV and 8.3% for
NV) is far below the targets (30% land surface by 2030) of the post-2020 Global Biodiversity
Framework (Jung et al., 2021). Moreover, these values are higher but consistent with those
found in previous years, such as 9.3% by Ribeiro et al. (2009) and 9% by Rezende et al.
(2018). We highlight that IT, despite not being PA, has proven to be fundamental for forest
restoration in AF (Benzeev et al., 2023). Noticeably, 70% and 90% of vegetation is more
than 10 km distant from PA and IT. Our findings are more alarming than those found by
Ribeiro et al. (2009). Forest formation has the largest area in PA (8.2%) and smaller for IT
(1.5%). This result is expected because of its large contribution to AF composition (62.1%)
because forests have been commonly the main target for PA creation. In addition, *restinga*
and mangroves had a high overlap with PA and TI (40%), due to the high density of these
protective measures on the Brazilian coast, especially in Serra do Mar. However, despite
this high proportion of protection, these ecosystems have faced many threats in recent
decades, which can affect several functions of ecosystems and local populations (Diniz et
al., 2019). Savanna formation was critical to ensuring connectivity, however, this class has
the lowest proportion of PA and TI (4.7%) despite representing 23% of the amount of

vegetation, possibly because this vegetation formation is not guaranteed by specific protection laws. Since deforestation outside PA and IT has been lower than in private rural areas (da Silva et al., 2023), these areas are essential to ensure biodiversity conservation (Avigliano et al., 2019; Benzeev et al., 2023). Therefore, it is necessary to create new PA and IT, and strengthen the connection network between existing ones, as well as restrictions in their surroundings to promote the restoration of vegetation.

5. Conclusion

To our knowledge, this is the first work that analyzed the spatiotemporal dynamics of the entire AF landscape structure through multiple landscape metrics, considering a broad tri-national delimitation, only forest vegetation and both forest and other natural vegetation, and the effect of roads and railways. Our findings allow a detailed understanding of the habitat fragmentation process in the AF in the last three and half decades. The number of FV fragments has increased, which comes accompanied by an important increment of vegetation. Besides that, NV—fundamental to promote connectivity—is far from being under enough protection. Overall, the fragmentation scenarios in Argentina, Brazil, and Paraguay are equally worrying (97% of fragments are very small and 60% are under edge effect). We also highlight the substantial effect of roads and railways on breaking large FV fragments apart, likely disrupting the functional connectivity of several ecological processes. These results lead us to reinforce the need for conservation and restoration actions, such as investing in implementing conservation plans for large fragments, promoting the connectivity of small fragments, managing the matrix to minimize edge effects and improve connectivity, and leading restoration actions in key areas, such as large and isolated fragments and indigenous territory. Added to this, we highlight the importance of planning and building fauna passages to improve landscape connectivity and reduce wildlife roadkill. Finally, the protection legislation implemented in mid-2005, combined with the restoration initiatives started in 2009, and the implementation of the CAR in 2012 appear to be having an effect in starting a process of AF restoration. The continuity and expansion of these measures are essential to guarantee the continuity of this AF process in the future, given the new threats of climate change and the expansion of urban and agricultural areas.

Funding information

MHV was supported by grants #2013/02883-7, #2017/09676-8, and #2022/01899-6, São Paulo Research Foundation (FAPESP), and CAPES (grant numbers 88887.513979/2020-00 and 1588183). RLM was supported by Bryce Carmine and Anne Carmine (née Percival),

through the Massey University Foundation. CHG is a research fellow of the National Council for Scientific and Technological Development - CNPq (grant 311209/2021-1). JEFO was supported by FAPESP projects #2014/23132-2, and #2021/02132-2. MCR was supported by FAPESP (processes #2013/50421-2; #2020/01779-5; #2021/08322-3; #2021/08534-0; #2021/10195-0; #2021/10639-5; #2022/10760-1) and CNPq (processes #442147/2020-1; #440145/2022-8; #402765/2021-4; #313016/2021-6; #440145/2022-8), and São Paulo State University - UNESP for their financial support. This study was financed in part by CAPES Brazil – Finance Code 001. This study is also part of the *Center for Research on Biodiversity Dynamics and Climate Change*, which is financed by FAPESP.

CRediT authorship contribution statement

MHV: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Visualization, Writing— original draft, Writing—review and editing. **RLM**: Conceptualization, Writing—review and editing. **BBN**: Conceptualization, Methodology, Software, Writing—review and editing. **JEFO**: Conceptualization, Writing—review and editing. **VT**: Conceptualization, Writing—review and editing. **RB**: Writing—review and editing. **CDA**: Writing—review and editing. **MRR**: Writing—review and editing. **CHG**: Methodology, Software, Writing—review and editing. **MCR**: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Resources, Software, Supervision, Writing—review and editing. All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Code provided in GitHub (https://github.com/LEEClab/ms-atlantic-forest-spatiotemporal-dynamics). Data and code are provided in Open Science Files (OSF) (https://doi.org/10.17605/OSF.IO/RFWBZ).

Acknowledgements

We thank Luis Fernando Guedes Pinto from SOS Mata Atlântica for assistance with MapBiomas data and Atlantic Forest delimitation. We also thank Alexandre Uezu for valuable contributions to the manuscript. We thank all members from Spatial Ecology and Conservation Lab (LEEC) for their help with fruitful discussions and suggestions, especially Rafael Souza. MHV thanks João Giovanelli, Rodrigo Nobre, Rafael Fávero, and Thiago Gonçalves-Souza for their valuable proposals for data analysis and discussion of results, Alejandro R. Vargas by vegetation cover in Argentina and Paraguay, and Maurício E. Graipel by for pointing out the role of legislation in large-scale vegetation dynamics.

References

Adorno, B.F.C.B., Barros, F.M., Cezar Ribeiro, M., da Silva, V.X., Hasui, É., 2021. Landscape heterogeneity shapes bird phylogenetic responses at forest–matrix interfaces in Atlantic Forest, Brazil. Biotropica 53, 409–421.

Avigliano, E., Rosso, J.J., Lijtmaer, D., Ondarza, P., Piacentini, L., Izquierdo, M., Cirigliano, A., Romano, G., Nuñez Bustos, E., Porta, A., Mabragaña, E., Grassi, E., Palermo, J., Bukowski, B., Tubaro, P., Schenone, N., 2019. Biodiversity and threats in non-protected areas: A multidisciplinary and multi-taxa approach focused on the Atlantic Forest. Heliyon 5, e02292.

Awade, M., Metzger, J.P., 2008. Using gap-crossing capacity to evaluate functional connectivity of two Atlantic rainforest birds and their response to fragmentation. Austral Ecology 33, 863–871.

Banks-Leite, C., Ewers, R.M., Metzger, J.P., 2013. The confounded effects of habitat disturbance at the local, patch and landscape scale on understorey birds of the Atlantic Forest: Implications for the development of landscape-based indicators. Ecological Indicators, Linking landscape structure and biodiversity 31, 82–88.

Banks-Leite, C., Pardini, R., Tambosi, L.R., Pearse, W.D., Bueno, A.A., Bruscagin, R.T., Condez, T.H., Dixo, M., Igari, A.T., Martensen, A.C., Metzger, J.P., 2014. Using ecological thresholds to evaluate the costs and benefits of set-asides in a biodiversity hotspot. Science 345, 1041–1045.

Barbosa, K.V. de C., Knogge, C., Develey, P.F., Jenkins, C.N., Uezu, A., 2017. Use of small Atlantic Forest fragments by birds in Southeast Brazil. Natureza & Conservação 15, 42–46.

Benzeev, R., Zhang, S., Rauber, M.A., Vance, E.A., Newton, P., 2023. Formalizing tenure of Indigenous lands improved forest outcomes in the Atlantic Forest of Brazil. PNAS Nexus 2, pgac287.

Bicudo da Silva, R.F., Millington, J.D.A., Moran, E.F., Batistella, M., Liu, J., 2020. Three decades of land-use and land-cover change in mountain regions of the Brazilian Atlantic Forest. Landscape and Urban Planning 204, 103948.

Bicudo da Silva, R.F., Moran, E., Viña, A., Millington, J.D.A., Dou, Y., Vieira, S.A., Lopez, M.C., Liu, J., 2023. Toward a forest transition across the Brazilian Atlantic Forest biome. Frontiers in Forests and Global Change 6.

Bivand R, 2022. rgrass: Interface Between 'GRASS' Geographical Information System and 'R'. R package version 0.3-6.

Cassimiro, I.M.F., Ribeiro, M.C., Assis, J.C., 2023. How did the animal come to cross the road? Drawing insights on animal movement from existing roadkill data and expert knowledge. Landsc Ecol. https://doi.org/10.1007/s10980-023-01671-5

Chase, J.M., Blowes, S.A., Knight, T.M., Gerstner, K., May, F., 2020. Ecosystem decay exacerbates biodiversity loss with habitat loss. Nature 584, 238–243.

Crouzeilles, R., Santiami, E., Rosa, M., Pugliese, L., Brancalion, P.H.S., Rodrigues, R.R., Metzger, J.P., Calmon, M., Scaramuzza, C.A. de M., Matsumoto, M.H., Padovezi, A., Benini, R. de M., Chaves, R.B., Metzker, T., Fernandes, R.B., Scarano, F.R., Schmitt, J., Lui, G., Christ, P., Vieira, R.M., Senta, M.M.D., Malaguti, G.A., Strassburg, B.B.N., Pinto, S., 2019. There is hope for achieving ambitious Atlantic Forest restoration commitments. Perspectives in Ecology and Conservation 17, 80–83.

Da Ponte, E., Mack, B., Wohlfart, C., Rodas, O., Fleckenstein, M., Oppelt, N., Dech, S., Kuenzer, C., 2017. Assessing Forest Cover Dynamics and Forest Perception in the Atlantic Forest of Paraguay, Combining Remote Sensing and Household Level Data. Forests 8, 389.

Da Silva, J.M., Casteleti, C.H.M., 2003. Status of the Biodiversity of the Atlantic Forest of Brazil, in: Galindo-Leal, C., Câmara, I. (Eds.), Washington (DC): C. Galind-Leal, IG Câmara. The Atlantic Forest of South America: Biodiversity Status, Threats, and Outlook. Conservation International. pp. 3–11.

da Silva, R.F.B., de Castro Victoria, D., Nossack, F.Á., Viña, A., Millington, J.D.A., Vieira, S.A., Batistella, M., Moran, E., Liu, J., 2023. Slow-down of deforestation following a Brazilian forest policy was less effective on private lands than in all conservation areas. Commun Earth Environ 4, 1–12.

Dale, M.R.T., Fortin, M.-J., 2014. Spatial analysis: a guide for ecologists, Second Edition. ed. Cambridge University Press, Cambridge; New York.

Dam, J. van, Hombergh, H. van den, Hilders, M., 2019. An analysis of existing laws on forest protection in the main soy-producing countries in Latin America.

de la Sancha, N.U., González-Maya, J.F., Boyle, S.A., Pérez-Estigarribia, P.E., Urbina-Cardona, J.N., McIntyre, N.E., 2023. Bioindicators of edge effects within Atlantic Forest remnants: Conservation implications in a threatened biodiversity hotspot. Diversity and Distributions 29, 349–363.

de Lima, R.A.F., Oliveira, A.A., Pitta, G.R., de Gasper, A.L., Vibrans, A.C., Chave, J., ter Steege, H., Prado, P.I., 2020. The erosion of biodiversity and biomass in the Atlantic Forest biodiversity hotspot. Nat Commun 11, 6347.

Dias, T. da C., Silveira, L.F., Francisco, M.R., 2023. Spatiotemporal dynamics reveals forest rejuvenation, fragmentation, and edge effects in an Atlantic Forest hotspot, the Pernambuco Endemism Center, northeastern Brazil. PLOS ONE 18, e0291234.

Diniz, C., Cortinhas, L., Nerino, G., Rodrigues, J., Sadeck, L., Adami, M., Souza-Filho, P., 2019. Brazilian Mangrove Status: Three Decades of Satellite Data Analysis. Remote Sensing 11, 808.

Diniz, M.F., Coelho, M.T.P., de Sousa, F.G., Hasui, É., Loyola, R., 2021. The underestimated role of small fragments for carnivore dispersal in the Atlantic Forest. Perspectives in Ecology and Conservation 19, 81–89. https://doi.org/10.1016/j.pecon.2020.12.001

Fahrig, L., 2003. Effects of Habitat Fragmentation on Biodiversity. Annu. Rev. Ecol. Evol. Syst. 34, 487–515.

Fahrig, L., Watling, J.I., Arnillas, C.A., Arroyo-Rodríguez, V., Jörger-Hickfang, T., Müller, J., Pereira, H.M., Riva, F., Rösch, V., Seibold, S., Tscharntke, T., May, F., 2022. Resolving the SLOSS dilemma for biodiversity conservation: a research agenda. Biological Reviews 97, 99–114.

Faria, D., Morante-Filho, J.C., Baumgarten, J., Bovendorp, R.S., Cazetta, E., Gaiotto, F.A., Mariano-Neto, E., Mielke, M.S., Pessoa, M.S., Rocha-Santos, L., Santos, A.S., Soares, L.A.S.S., Talora, D.C., Vieira, E.M., Benchimol, M., 2023. The breakdown of ecosystem functionality driven by deforestation in a global biodiversity hotspot. Biological Conservation 283, 110126.

Figueiredo, M. de S.L., Weber, M.M., Brasileiro, C.A., Cerqueira, R., Grelle, C.E.V., Jenkins, C.N., Solidade, C.V., Thomé, M.T.C., Vale, M.M., Lorini, M.L., 2021. Tetrapod Diversity in the Atlantic Forest: Maps and Gaps, in: Marques, M.C.M., Grelle, C.E.V. (Eds.), The Atlantic Forest. Springer International Publishing, Cham, pp. 185–204.

Fischer, R., Taubert, F., Müller, M.S., Groeneveld, J., Lehmann, S., Wiegand, T., Huth, A., 2021. Accelerated forest fragmentation leads to critical increase in tropical forest edge area. Science Advances 7, eabg7012.

Flora e Funga do Brasil. Jardim Botânico do Rio de Janeiro. Available at: http://floradobrasil.jbrj.gov.br. Accessed on: 10 Oct 2023.

Fundación Vida Silvestre Argentina & WWF. 2017. State of the Atlantic Forest: Three Countries, 148 Million People, One of the Richest Forests on Earth. Fundación Vida Silvestre Argentina & WWF. Puerto Iguazú, Argentina. pp 146.

Galetti, M., Gonçalves, F., Villar, N., Zipparro, V.B., Paz, C., Mendes, C., Lautenschlager, L., Souza, Y., Akkawi, P., Pedrosa, F., Bulascoschi, L., Bello, C., Sevá, A.P., Sales, L., Genes, L., Abra, F., Bovendorp, R.S., 2021. Causes and Consequences of Large-Scale Defaunation in the Atlantic Forest, in: Marques, M.C.M., Grelle, C.E.V. (Eds.), The Atlantic Forest: History, Biodiversity, Threats and Opportunities of the Mega-Diverse Forest. Springer International Publishing, Cham, pp. 297–324.

Gama-Rodrigues, A.C., Müller, M.W., Gama-Rodrigues, E.F., Mendes, F.A.T., 2021. Cacao-based agroforestry systems in the Atlantic Forest and Amazon Biomes: An ecoregional analysis of land use. Agricultural Systems 194, 103270.

Haddad, N.M., Brudvig, L.A., Clobert, J., Davies, K.F., Gonzalez, A., Holt, R.D., Lovejoy, T.E., Sexton, J.O., Austin, M.P., Collins, C.D., Cook, W.M., Damschen, E.I., Ewers, R.M., Foster, B.L., Jenkins, C.N., King, A.J., Laurance, W.F., Levey, D.J., Margules, C.R., Melbourne, B.A., Nicholls, A.O., Orrock, J.L., Song, D.-X., Townshend, J.R., 2015. Habitat fragmentation and its lasting impact on Earth's ecosystems. Sci. Adv. 1, e1500052.

Hansen, M.C., Wang, L., Song, X.-P., Tyukavina, A., Turubanova, S., Potapov, P.V., Stehman, S.V., 2020. The fate of tropical forest fragments. Science Advances 6, eaax8574.

Harper, K.A., Macdonald, S.E., 2011. Quantifying distance of edge influence: a comparison of methods and a new randomization method. Ecosphere 2, art94.

IBGE — Instituto Brasileiro de Geografia e Estatística, 2008. Mapa da área de aplicação da Lei no 11.428 de 2006.

Izquierdo, A., C. De Angelo & M. Aide. 2008. Thirty years of human demography and land-use change in the Atlantic Forest of Misiones, Argentina: a test of the forest transition model. Ecology and Society 13 (2):3.

Joly, C.A., Metzger, J.P., Tabarelli, M., 2014. Experiences from the Brazilian Atlantic Forest: ecological findings and conservation initiatives. New Phytol 204, 459–473.

Jung, M., Arnell, A., de Lamo, X., García-Rangel, S., Lewis, M., Mark, J., Merow, C., Miles, L., Ondo, I., Pironon, S., Ravilious, C., Rivers, M., Schepashenko, D., Tallowin, O., van Soesbergen, A., Govaerts, R., Boyle, B.L., Enquist, B.J., Feng, X., Gallagher, R., Maitner, B., Meiri, S., Mulligan, M., Ofer, G., Roll, U., Hanson, J.O., Jetz, W., Di Marco, M., McGowan, J., Rinnan, D.S., Sachs, J.D., Lesiv, M., Adams, V.M., Andrew, S.C., Burger, J.R., Hannah, L., Marquet, P.A., McCarthy, J.K., Morueta-Holme, N., Newman, E.A., Park, D.S., Roehrdanz, P.R., Svenning, J.-C., Violle, C., Wieringa, J.J., Wynne, G., Fritz, S., Strassburg, B.B.N., Obersteiner, M., Kapos, V., Burgess, N., Schmidt-Traub, G., Visconti, P., 2021. Areas of global importance for conserving terrestrial biodiversity, carbon and water. Nat Ecol Evol 1–11.

Law 26.331. 2007. Presupuestos mínimos de protección ambiental de los bosques nativos. Senado y Cámara de Diputados de la Nación Argentina. Buenos Aires, Argentina.

Lyra-Jorge, M.C., Ribeiro, M.C., Ciocheti, G., Tambosi, L.R., Pivello, V.R., 2010. Influence of multi-scale landscape structure on the occurrence of carnivorous mammals in a human-modified savanna, Brazil. Eur J Wildl Res 56, 359–368.

Magioli, M., Ribeiro, M.C., Ferraz, K.M.P.M.B., Rodrigues, M.G., 2015. Thresholds in the relationship between functional diversity and patch size for mammals in the Brazilian Atlantic Forest. Animal Conservation 18, 499–511.

Magnago, L.F.S., Magrach, A., Barlow, J., Schaefer, C.E.G.R., Laurance, W.F., Martins, S.V., Edwards, D.P., 2017. Do fragment size and edge effects predict carbon stocks

in trees and lianas in tropical forests? Functional Ecology 31, 542–552.

Magnago, L.F.S., Rocha, M.F., Meyer, L., Martins, S.V., Meira-Neto, J.A.A., 2015. Microclimatic conditions at forest edges have significant impacts on vegetation structure in large Atlantic forest fragments. Biodivers Conserv 24, 2305–2318.

Marjakangas, E., Abrego, N., Grøtan, V., Lima, R.A.F., Bello, C., Bovendorp, R.S., Culot, L., Hasui, É., Lima, F., Muylaert, R.L., Niebuhr, B.B., Oliveira, A.A., Pereira, L.A., Prado, P.I., Stevens, R.D., Vancine, M.H., Ribeiro, M.C., Galetti, M., Ovaskainen, O., 2020. Fragmented tropical forests lose mutualistic plant–animal interactions. Divers. Distrib. 26, 154–168.

Marques, M.C.M., Trindade, W., Bohn, A., Grelle, C.E.V., 2021. The Atlantic Forest: An Introduction to the Megadiverse Forest of South America, in: Marques, M.C.M., Grelle, C.E.V. (Eds.), The Atlantic Forest: History, Biodiversity, Threats and Opportunities of the Mega-Diverse Forest. Springer International Publishing, Cham, pp. 3–23.

Martinez Pardo, J., Saura, S., Insaurralde, A., Di Bitetti, M.S., Paviolo, A., De Angelo, C., 2023. Much more than forest loss: four decades of habitat connectivity decline for Atlantic Forest jaguars. Landscape Ecology, 38, 41-57.

Melo, F.P.L., Pinto, S.R.R., Brancalion, P.H.S., Castro, P.S., Rodrigues, R.R., Aronson, J., Tabarelli, M., 2013. Priority setting for scaling-up tropical forest restoration projects: Early lessons from the Atlantic Forest Restoration Pact. Environmental Science & Policy 33, 395–404.

Mohebalian, P.M., Lopez, L.N., Tischner, A.B., Aguilar, F.X., 2022. Deforestation in South America's tri-national Paraná Atlantic Forest: Trends and associational factors. Forest Policy and Economics 137, 102697.

Morante-Filho, J.C., Arroyo-Rodríguez, V., Pessoa, M. de S., Cazetta, E., Faria, D., 2018. Direct and cascading effects of landscape structure on tropical forest and non-forest frugivorous birds. Ecological Applications 28, 2024–2032.

Muylaert, R.L., Stevens, R.D., Ribeiro, M.C., 2016. Threshold effect of habitat loss on bat richness in cerrado-forest landscapes. Ecological Applications 26, 1854–1867.

Muylaert, R.L., Vancine, M.H., Bernardo, R., Oshima, J.E.F., Sobral-Souza, T., Tonetti, V.R., Niebuhr, B.B., Ribeiro, M.C., 2018. Uma nota sobre os limites territoriais da Mata Atlântica. Oecol. Aust. 22, 302–311.

Neteler, M., Bowman, M.H., Landa, M., Metz, M., 2012. GRASS GIS: A multi-purpose open source GIS. Environmental Modelling & Software 31, 124–130.

Parra-Sanchez, E., Banks-Leite, C., 2020. The magnitude and extent of edge effects on vascular epiphytes across the Brazilian Atlantic Forest. Sci Rep 10, 18847.

Paviolo, A., C. De Angelo, K. M. P. M. B. Ferraz, R. G. Morato, J. Martínez Pardo, A. C. Srbek-Araujo, B. Beisegel, F. Lima, D. Sana, M. Xavier da Silva, M. Velázquez, L. Cullen Jr, P. G. Crawshaw Jr., M. L. S. P. Jorge, P. M. Galetti Jr, M. S. Di Bitetti, R. Cunha de Paula, E. Eizirik, T. M. Aide, M. P. Cruz, M. Perilli, A. S. M. C. Souza, V. A. Quiroga, E. Nakaro, F. Ramírez, S. Fernández, S. A. Costa, E. A. Moraes & F. C. C. Azevedo. 2016. A biodiversity hotspot losing its top predator: The challenge of jaguar conservation in the Atlantic Forest of South America. Scientific Reports 6:37147.

Peres, E.A., Pinto-da-Rocha, R., Lohmann, L.G., Michelangeli, F.A., Miyaki, C.Y., Carnaval, A.C., 2020. Patterns of Species and Lineage Diversity in the Atlantic Rainforest of Brazil, in: Rull, V., Carnaval, A.C. (Eds.), Neotropical Diversification: Patterns and Processes, Fascinating Life Sciences. Springer International Publishing, Cham, pp. 415–447.

Piffer, P.R., Calaboni, A., Rosa, M.R., Schwartz, N.B., Tambosi, L.R., Uriarte, M., 2022. Ephemeral forest regeneration limits carbon sequestration potential in the Brazilian Atlantic Forest. Global Change Biology 28, 630–643.

Pivello, V.R., Vieira, I., Christianini, A.V., Ribeiro, D.B., da Silva Menezes, L., Berlinck, C.N., Melo, F.P.L., Marengo, J.A., Tornquist, C.G., Tomas, W.M., Overbeck, G.E., 2021. Understanding Brazil's catastrophic fires: Causes, consequences and policy needed to prevent future tragedies. Perspectives in Ecology and Conservation 19, 233–255.

Püttker, T., Crouzeilles, R., Almeida-Gomes, M., Schmoeller, M., Maurenza, D., Alves-Pinto, H., Pardini, R., Vieira, M.V., Banks-Leite, C., Fonseca, C.R., Metzger, J.P., Accacio, G.M., Alexandrino, E.R., Barros, C.S., Bogoni, J.A., Boscolo, D., Brancalion, P.H.S., Bueno, A.A., Cambui, E.C.B., Canale, G.R., Cerqueira, R., Cesar, R.G., Colletta, G.D., Delciellos, A.C., Dixo, M., Estavillo, C., Esteves, C.F., Falcão, F., Farah, F.T., Faria, D., Ferraz, K.M.P.M.B., Ferraz, S.F.B., Ferreira, P.A., Graipel, M.E., Grelle, C.E.V., Hernández, M.I.M., Ivanauskas, N., Laps, R.R., Leal, I.R., Lima, M.M., Lion, M.B., Magioli, M., Magnago, L.F.S., Mangueira, J.R.A.S., Marciano-Jr, E., Mariano-Neto, E., Marques, M.C.M., Martins, S.V., Matos, M.A., Matos, F.A.R., Miachir, J.I., Morante-Filho, J.M., Olifiers, N., Oliveira-Santos, L.G.R., Paciencia, M.L.B., Paglia, A.P., Passamani, M., Peres, C.A., Pinto Leite, C.M., Porto, T.J., Querido, L.C.A., Reis, L.C., Rezende, A.A., Rigueira, D.M.G., Rocha, P.L.B., Rocha-Santos, L., Rodrigues, R.R., Santos, R.A.S., Santos, J.S., Silveira, M.S., Simonelli, M., Tabarelli, M., Vasconcelos, R.N., Viana, B.F., Vieira, E.M., Prevedello, J.A., 2020. Indirect effects of habitat loss via habitat fragmentation: A cross-taxa analysis of forest-dependent species. Biological Conservation 241, 108368.

QGIS Development Team, 2023. QGIS Geographic Information System. Open Source Geospatial Foundation Project.

R Core Team, 2023. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Reis, R.E., Albert, J.S., Di Dario, F., Mincarone, M.M., Petry, P., Rocha, L.A., 2016. Fish biodiversity and conservation in South America. Journal of Fish Biology 89, 12–47.

Rezende, C.L., Scarano, F.R., Assad, E.D., Joly, C.A., Metzger, J.P., Strassburg, B.B.N., Tabarelli, M., Fonseca, G.A., Mittermeier, R.A., 2018. From hotspot to hopespot: An opportunity for the Brazilian Atlantic Forest. Perspectives in Ecology and Conservation 16, 208–214.

Ribeiro, M.C., Metzger, J.P., Martensen, A.C., Ponzoni, F.J., Hirota, M.M., 2009. The Brazilian Atlantic Forest: How much is left, and how is the remaining forest distributed? Implications for conservation. Biological Conservation 142, 1141–1153.

Riva, F., Nielsen, S.E., 2020. Six key steps for functional landscape analyses of habitat change. Landscape Ecol 35, 1495–1504.

Rosa, M.R., Brancalion, P.H.S., Crouzeilles, R., Tambosi, L.R., Piffer, P.R., Lenti, F.E.B., Hirota, M., Santiami, E., Metzger, J.P., 2021. Hidden destruction of older forests threatens Brazil's Atlantic Forest and challenges restoration programs. Sci. Adv. 7, eabc4547.

Rudel, T.K., Coomes, O.T., Moran, E., Achard, F., Angelsen, A., Xu, J., Lambin, E., 2005. Forest transitions: towards a global understanding of land use change. Global Environmental Change 15, 23–31.

Scarano, F.R., 2002. Structure, Function and Floristic Relationships of Plant Communities in Stressful Habitats Marginal to the Brazilian Atlantic Rainforest. Ann Bot 90, 517–524.

Silva, R.F.B. da, Batistella, M., Moran, E.F., 2017. Socioeconomic changes and environmental policies as dimensions of regional land transitions in the Atlantic Forest, Brazil. Environmental Science & Policy 74, 14–22.

Sloan, S., Jenkins, C.N., Joppa, L.N., Gaveau, D.L.A., Laurance, W.F., 2014. Remaining natural vegetation in the global biodiversity hotspots. Biological Conservation 177, 12–24.

Solórzano, A., Brasil, L.S.C. de A., de Oliveira, R.R., 2021. The Atlantic Forest Ecological History: From Pre-colonial Times to the Anthropocene, in: Marques, M.C.M., Grelle, C.E.V. (Eds.), The Atlantic Forest: History, Biodiversity, Threats and Opportunities of the Mega-Diverse Forest. Springer International Publishing, Cham, pp. 25–44.

Souza, C.M., Z. Shimbo, J., Rosa, M.R., Parente, L.L., A. Alencar, A., Rudorff, B.F.T., Hasenack, H., Matsumoto, M., G. Ferreira, L., Souza-Filho, P.W.M., de Oliveira, S.W., Rocha, W.F., Fonseca, A.V., Marques, C.B., Diniz, C.G., Costa, D., Monteiro, D., Rosa, E.R., Vélez-Martin, E., Weber, E.J., Lenti, F.E.B., Paternost, F.F., Pareyn, F.G.C., Siqueira, J.V., Viera, J.L., Neto, L.C.F., Saraiva, M.M., Sales, M.H., Salgado,

M.P.G., Vasconcelos, R., Galano, S., Mesquita, V.V., Azevedo, T., 2020. Reconstructing Three Decades of Land Use and Land Cover Changes in Brazilian Biomes with Landsat Archive and Earth Engine. Remote Sensing 12, 2735.

Szangolies, L., Rohwäder, M.-S., Jeltsch, F., 2022. Single large AND several small habitat patches: A community perspective on their importance for biodiversity. Basic and Applied Ecology 65, 16–27.

Tabarelli, M., Aguiar, A.V., Ribeiro, M.C., Metzger, J.P., Peres, C.A., 2010. Prospects for biodiversity conservation in the Atlantic Forest: Lessons from aging human-modified landscapes. Biological Conservation 143, 2328–2340.

Tavares, A., Beiroz, W., Fialho, A., Frazão, F., Macedo, R., Louzada, J., Audino, L., 2019. Eucalyptus plantations as hybrid ecosystems: Implications for species conservation in the Brazilian Atlantic forest. Forest Ecology and Management 433, 131–139.

Teixeira, F.Z., Silva, L.G. da, Abra, F., Rosa, C., Buss, G., Guerreiro, M., Costa, E.R., Medeiros, A.S. de M., Gordo, M., Secco, H., 2022. A reviewof the application of canopy bridges in the conservation of primates and other arboreal animals across Brazil. Folia Primatologica 93, 479–492.

Tubenchlak, F., Badari, C.G., de Freitas Strauch, G., de Moraes, L.F.D., 2021. Changing the Agriculture Paradigm in the Brazilian Atlantic Forest: The Importance of Agroforestry, in: Marques, M.C.M., Grelle, C.E.V. (Eds.), The Atlantic Forest. Springer International Publishing, Cham, pp. 369–388.

Turner, M.G., Gardner, R.H., 2015. Landscape Ecology in Theory and Practice. Springer New York, New York, NY.

Vale, M.M., Arias, P.A., Ortega, G., Cardoso, M., Oliveira, B.F.A., Loyola, R., Scarano, F.R., 2021. Climate Change and Biodiversity in the Atlantic Forest: Best Climatic Models, Predicted Changes and Impacts, and Adaptation Options, in: Marques, M.C.M., Grelle, C.E.V. (Eds.), The Atlantic Forest: History, Biodiversity, Threats and Opportunities of the Mega-Diverse Forest. Springer International Publishing, Cham, pp. 253–267.

Varassin, I.G., Agostini, K., Wolowski, M., Freitas, L., 2021. Pollination Systems in the Atlantic Forest: Characterisation, Threats, and Opportunities, in: Marques, M.C.M., Grelle, C.E.V. (Eds.), The Atlantic Forest: History, Biodiversity, Threats and Opportunities of the Mega-Diverse Forest. Springer International Publishing, Cham, pp. 325–344.

Zalles, V., Hansen, M.C., Potapov, P.V., Parker, D., Stehman, S.V., Pickens, A.H., Parente, L.L., Ferreira, L.G., Song, X.-P., Hernandez-Serna, A., Kommareddy, I., 2021. Rapid expansion of human impact on natural land in South America since 1985. Science Advances 7, eabg1620.

Zimmermann Teixeira, F., Kindel, A., Hartz, S.M., Mitchell, S., Fahrig, L., 2017. When road-kill hotspots do not indicate the best sites for road-kill mitigation. Journal of Applied Ecology 54, 1544–1551.