Erdr1 orchestrates macrophage polarization and determines cell fate via dynamic interaction with YAP1 and Mid1

Yuhang Wang[1,2,✉]

Affiliations:

1. Department of Microbiology and Immunology, University of Iowa, Iowa City, IA 52242, USA
2. Department of Pathology, University of Iowa, Iowa City, IA 52242, USA (Current)

✉ Corresponding author:
Yuhang Wang, Email: yuhang-wang@uiowa.edu

Abstract:

Erythroid differentiation regulator 1 (Erdr1) is a stress-induced, widely distributed, extremely conserved secreted factor found in both humans and mice. Initially identified as an inducer of hemoglobin synthesis, it has emerged as a multifunctional protein, especially in immune cells. Although Erdr1 has been implicated in T cells and NK cell function, its role in macrophage remains unclear. This study aims to explore the precise function of Erdr1 in IL-1β production in macrophages and uncover the underlying mechanisms. Data manifest Erdr1 could play an inhibition role in IL-1β production, which also has been reported by previous research. What significance is we discovered Erdr1 has the capacity to promote IL-1β production which is associated with Erdr1 dose and cell density. We observed that Erdr1 has distinct expression patterns in pro-inflammatory (M1) and anti-inflammatory (M2) macrophages compared to naive macrophages. We hypothesized that Erdr1 dual modulates IL-1β production by binding with distinct adaptors via concentration change. Mechanistically, we demonstrated that Erdr1 has dynamic interaction with YAP1 and Mid1 by distinct domains. These intricate interplays not only govern the dynamic regulation of IL-1β production but also mediate macrophage functional and metabolic reprogramming and determine cell fate. This study highlights that Erdr1 orchestrates macrophage polarization by regulating YAP1 through the non-classical Hippo pathway.

Introduction

Erdr1 was initially identified as a stress-induced factor that acts as a hemoglobin synthesis-inducing factor in leukemia cells and was extremely conserved between humans and mice [1,2]. Subsequent research has demonstrated its important role in immune response [3-6]. Erdr1 has been shown to play a critical role in T cell proliferation, activation [3,4], and apoptosis [5]. Recombinant Erdr1 protein treatment strengthens TCR signal in T cells upon activation by CD3 antibody in vitro [3,4], while in vivo, it significantly promotes thymocyte proliferation [3]. However, endogenous overexpression of Erdr1 induces T cell apoptosis upon activation [5]. The conflicting functions of Erdr1 on T cells are most likely due to the Erdr1 concentration and cell context. Apart from T cells, Erdr1 protein has also been shown to boost NK cell activation by promoting F-actin accumulation [6]. Additionally, Erdr1 has been implicated in the pathogenesis of inflammatory disorders such as psoriasis [7] and rosacea [8]. Notably, Erdr1 was downregulated in these patients and has a negative correlation with the pro-inflammatory cytokine IL-18 [8,9], a readout cytokine of the inflammasome [10,11]. Animal experiments have manifested the broadly anti-inflammatory effect of Erdr1 protein in treating inflammatory disorders such as psoriasis [9], rosacea [8], colitis [12], and arthritis [13]. Erdr1 has also been reported to play a negative role in macrophage migration[14]. However, the function and mechanism of Erdr1 in macrophages are still unknown. Our preliminary data manifested that Erdr1 acts as an anti-inflammatory factor and could play an inhibition role in LPS-induced IL-1β production, which is consistent with previous reports. Interestingly, we discovered that Erdr1 also has the capacity to facilitate IL-1β production. The dual role of Erdr1 over IL-1β production is correlated with Erdr1 dose and cell density. Particularly, we found that at decent cell density, Erdr1 is significantly down-regulated in LPS-induced pro-inflammatory macrophages (M1) but dramatically upregulated in IL-4-induced anti-inflammatory macrophages (M2) compared to naive macrophages (M0). The correlation of Erdr1 level with IL-1β production in macrophages follows a bell-shaped model. We hypothesize that, compared to basal Erdr1 level in steady state (M0), the up-regulated Erdr1 inhibits IL-1β production and mediates macrophage M2 polarization, while down-regulated Erdr1 promotes IL-1β production and mediates macrophage M1 polarization, which was achieved by interacts with different adaptors via Erdr1 conformation change under different concentration. We aim to investigate the candidates' adaptors of Erdr1 and uncover the mechanism of Erdr1's dual role in IL-1β production and macrophage M1/M2 polarization.

Previous research has demonstrated that secreted Erdr1 mediates cellular adaptation, it promotes cell survival at a low concentration and low cell density but enhances cell death at high concentrations and high cell density [1], indicating Erdr1 involved in maintaining cellular homeostasis. Furthermore, the observation that the Erdr1 C-terminal domain was undetectable at cell-cell contact sites suggested that it may be inactivated at these sites [1]. Cell density-dependent inhibition, also known as contact inhibition, is a mechanism that regulates cell proliferation by modulating the Hippo-YAP1 pathway [15-17], which inspired us to link Erdr1 with the YAP1 signaling pathway. Moreover, through searching the database, we discovered the strong correlation of Erdr1 with YAP1 in various studies [5,18-21]. Firstly, Erdr1 has been found to directly regulate YAP1 signaling. For instance, in T cells, the overexpression of Erdr1 augments the expression of YAP1 and its signature gene, Amotl1, following T-cell stimulation [5]. Secondly, YAP1

knockout has also been shown to impact Erdr1 expression. In Treg cells, YAP1 knockout
prominently upregulates Erdr1 upon stimulation [19], and similarly, YAP1 knockout leads to a
remarkable increase in Erdr1 expression in Müller glia cells during degeneration induction [20].
However, it is worth noting that YAP1 knockout in mouse incisor epithelium results in a significant
downregulation of Erdr1 [18]. These intriguing findings suggest that YAP1 may regulate Erdr1 in a
cell type-specific manner. Given the robust correlation of Erdr1 with YAP1, we hypothesized YAP1
is one of the binding adaptors of Erdr1. Cell density-dependent inhibition was also observed play
role in controlling inflammation [22,23], we hypothesized that Erdr1 regulates IL-1β production by
directly modulating YAP1 in macrophages.
95
YAP1, the core of hippo pathway, has been established to mediate macrophage polarization and
facilitate pro-inflammatory cytokine production in macrophages [24-26]. The function of YAP1 is
intricately linked to its expression levels and subcellular localization. YAP1 ON (Hippo OFF)
occurs when YAP1 is translocated into the cell nuclear, where it interacts with its transcriptional
co-activators, thereby facilitating the activation of the YAP1 signaling pathway. Conversely, YAP1
OFF (Hippo ON) achieved by capturing YAP1 in the cytoplasm, leading to the inhibition of the
YAP1 signaling pathway [27]. We hypothesized that Erdr1 regulates IL-1β production achieving by
controlling YAP1 ON/OFF. Our preliminary data proved Erdr1 has a strong interaction with YAP1
in the cytoplasm in M2 but not M1 macrophages. We predicted Erdr1 can also drive YAP1 ON
and promote M1 polarization by disconnecting with YAP1 and binding to another adaptor, Mid1.
Mid1 is a highly conserved gene expressed ubiquitously in humans and mice [28]. Initially identified
for its essential role in development [28], Mid1 has also been shown to play a crucial role in
macrophage activation [29-31]. Our hypothesis is based on several lines of evidence. Firstly, Erdr1
and Mid1 are closely linked in gene mapping, as they are neighbor genes located in the PAR
region of the sex chromosomes X and Y in mice [32,33]. Secondly, Mid1 is functionally linked to
Erdr1, YAP1, and YAP1 downstream genes such as CTGF, Cry61, and IL-1β, as shown by
previous microarray data and experiments [19,34-40]. Thirdly, Erdr1 and Mid1 are known to play a
shared role in regulating YAP1 signature gene expression [36,38,41]. As an intrinsically disordered
protein (IDP) [42], Erdr1 lacks of fixed structure and has the potential to play multiple functions by
binding partners highly dynamically and cooperatively. Overall, our hypothesis is that Erdr1
interacts dynamically with YAP1 and Mid1 to orchestrate macrophage activation and to determine
the extent of inflammation.
118
Mechanistically, we evidenced that Erdr1 modulates IL-1β production through dynamically
interacting with YAP1 and Mid1 by distinct domains. Upregulated Erdr1 acts as an inhibitor of IL-
1β production by sequestering YAP1 directly in the cytoplasm (YAP1 OFF), driving macrophage
M2 polarization. Conversely, downregulated Erdr1 acts as an activator for IL-1β production via
disconnecting with YAP1 but interacting with Mid1, ultimately driving YAP1 translocate to the
nuclear (YAP1 ON), which promotes macrophage M1 polarization. More importantly, we also
manifested that Erdr1, by concentration change, decides metabolism strength and mediates
macrophage metabolic polarization towards a pro-anabolic or pro-catabolic state. Ultimately,
Erdr1 decides cell fate, determining whether cells survive or undergo programmed cell death
through either a pro-apoptotic or pro-pyroptotic pathway. This study highlights that Erdr1

dynamically drives and orchestrates macrophage polarization and determines cell fate by
regulating YAP1 through the non-classical Hippo pathway.
131
132 **Results**
133
Erdr1 is distinctly regulated in M1/M2 macrophages
Macrophages can be polarized into pro-inflammatory M1 macrophages and anti-inflammatory M2
macrophages [43,44], which are the two extremes of a spectrum of macrophage phenotypes. M1
macrophages can be induced through various stimuli, such as lipopolysaccharide (LPS), IFN-
gamma (IFNγ), while M2 macrophages can be induced by interleukin-4 (IL-4), IL-13, etc. [43,44]. We
first investigated if Erdr1 expression was regulated during the macrophage polarization process.
Here, we induced the polarization of M0 macrophages (Bone marrow-derived macrophage,
BMDM) into M1 macrophages by stimulating with 1 µg/ml of LPS for 24 hours, or into M2
macrophages by stimulating with 20 ng/ml of IL-4 for 24 hours and confirming M1 marker genes
(IL-1β, IL-6 and TNF) and M2 marker gene (IL10 and TGFβ) expression by qPCR (data not
shown). We observed Erdr1 expression was dramatically reduced in M1 macrophages at mRNA
level (Fig. 1a) and protein level (Fig. 1b), whereas it was significantly elevated in M2 macrophages
(Fig. 1a and Fig. 1b), compared to M0 macrophages. Further investigation was conducted to
assess the secretion and accumulation of Erdr1 in the supernatant of M1 and M2 macrophages,
considering it was reported to secrete upon cellular stress. Immunoblot analysis revealed a
dramatically elevated presence of Erdr1 in the supernatant of M2 macrophages when compared
to M0 macrophages (Fig. 1b). Surprisingly, there was also a significantly increased level of Erdr1
detected in the supernatant of M1 macrophages (Fig. 1b), despite the decrease in endogenous
Erdr1 expression (Fig. 1a and Fig. 1b). This increased secretion of Erdr1 in the medium of M1
macrophages might serve as a strategy to rapidly decrease the intracellular levels of Erdr1, rather
than an induced expression factor. These findings manifest Erdr1 exhibits significant but distinct
expression patterns in M1/M2 macrophage, implying Erdr1 might play roles both in M1 and M2
macrophage.
157
We conducted further investigations into the subcellular localization of Erdr1 in the polarized
macrophage, employing immunofluorescence techniques. Our findings revealed Erdr1 has
increased nuclear localization in M1 macrophage but increased cytoplasm localization in M2
macrophage compared to M0 macrophage (Fig. 1c). We further confirmed these observations by
ImageJ analysis, with significantly augmented Erdr1 Nuclear/Cytoplasm intensity ratio in M1
macrophages, while a decreased ratio in M2 macrophages, compared to M0 (Fig. 1d). These data
display Erdr1 has distinct subcellular localization pattern in M1/M2 macrophages.
165
IL-1β is one of the most important pro-inflammatory cytokines induced in M1 macrophages [43,44].
IL-1β is processed during the Inflammasome activation process, as well as IL-18 [10,11]. Erdr1 has
been shown to have a negative correlation with IL-18 [45]. Consistent with that, here we
demonstrated a negative correlation of endogenous Erdr1 level with IL-1β and IL-18 level in
macrophages (Fig. 1e). IL-10, which is the anti-inflammatory cytokines induced in M2

macrophages, has positive correlation with Erdr1 (data not shown). This indicated Erdr1 is a
signature gene for M2 macrophages.
173
174 Taken together, our observations manifest that Erdr1 exhibits distinct regulation in M1 and M2
175 macrophages. Most importantly, the significant upregulated Erdr1 acts as a marker gene for M2
176 macrophages, while if and how the downregulated Erdr1 plays role in M1 macrophage still not
177 clear.
178
Erdr1 has dose-dependent bell-shaped effect in LPS-induced IL-1β production in
macrophages
We further aimed to elucidate the function of Erdr1 in regulating IL-1β production in M1
macrophages. Because we observed Erdr1 exhibits both endogenous regulation and induced
secretion during macrophage activation, we conducted experiment to investigate how
endogenous and exogenous Erdr1 affect the production of IL-1β in response to LPS stimulation.
185
First, we assessed the endogenous Erdr1 function on LPS-induced IL-1β production. M0
macrophages were genetically modified to either knockout or overexpress Erdr1 through
transduction with Erdr1 CRISPR/Cas9 knockout or Erdr1 overexpression lentiviruses,
respectively confirmed Erdr1 expression by qPCR (Fig. 2a and Fig. 2d). We observed that Erdr1
knockdown significantly suppressed LPS-induced IL-1β production at mRNA level in cell lysate
(Fig. 2b) and protein level in the supernatant (Fig. 2c). Interestingly, we observed that moderate
overexpression of Erdr1 (Erdr1 OE1, about 4.3 folds higher) dramatically enhanced LPS-induced
IL-1β production, while a high-level overexpression of Erdr1 (Erdr1 OE2, about 28.6 folds higher)
inhibited LPS-induced IL-1β production (Fig. 2e and Fig. 2f). Notably, we discovered a bell-shaped
correlation between the endogenous Erdr1 and IL-1β production (Fig. 2g), suggesting that Erdr1
plays a dual role in regulating IL-1β production.
197
Subsequently, we investigated the impact of the secreted Erdr1 in the medium on LPS-induced
IL-1β production. To achieve this, we employed a range of concentrations of Erdr1 neutralization
antibody to block the secreted Erdr1 in the medium. Interestingly, we observed a bell-shaped
relationship between the Erdr1 antibody concentration and IL-1β production (Fig. 2h). These
results suggest that an optimal concentration of Erdr1 in the medium exerts a positive effect on
promoting IL-1β production. However, an excessive accumulation of Erdr1 in the medium appears
to trigger a negative feedback mechanism, ultimately inhibiting IL-1β production.
205
Since purified Erdr1 protein has been shown to play an inhibition role in inflammation [7,8,46]. We
next tested Erdr1 protein effect in IL-1β production in macrophages through an exogenous protein
supplementation experiment. Purified full-length Erdr1 was prepared and supplemented with a
series of concentrations as indicated to the medium. Notably, data manifested Erdr1 protein has
a dose-dependent bell-shaped effect in regulating LPS-induced IL-1β production (Fig. 2i). These
findings provide striking evidence that a low dose of Erdr1 in the medium has the ability to induce
IL-1β production, whereas a high dose of Erdr1 exerts an inhibitory role in IL-1β production. This
indicates that the function of Erdr1 in IL-1β production is highly correlated to Erdr1 dose.

In summary, our data support that Erdr1 could play an inhibition role in IL-1β production, which has been reported. But we for the first time, demonstrate that Erdr1 also has the potential to promote IL-1β production. These data suggest the influence of Erdr1 in IL-1β production following a dose-dependent bell-shaped model (Fig. 2j).

Erdr1 effect in IL-1β production also correlates with cell density in macrophages

Our observations above were obtained from samples with cell density of $0.6*10^6$ cells/ml. Given that prior research has reported Erdr1 effect in cell viability was influenced by cell density[1], we then investigated if Erdr1's role in IL-1β expression was affected by cell density, and if Erdr1 affects cell viability in macrophage.

We seeded macrophages with 3 different densities, low density ($0.1*10^6$ cells/ml), moderate density ($0.5*10^6$ cells/ml), and high density ($2.5*10^6$ cells/ml). Consistent with previous research [22,23], we confirmed that both basal and LPS-stimulated IL-1β expression exhibited an increase at low density while being suppressed at higher density compared to levels at moderate density (Fig. 3a). We also observed that high cell density protects the cell from death (Fig. 3b). Interestingly, we noted cell density affects basal endogenous Erdr1 expression levels (Fig. 3c), Specifically, heightened basal Erdr1 expression was evident in scenarios of high cell density, whereas low cell density exhibited reduced Erdr1 expression, compared to the moderate one (Fig. 3c). Furthermore, we observed Erdr1 is upregulated under conditions of low cell density upon LPS stimulation, this response is contrary to our observations in both moderate and high-density scenarios (Fig. 3c). Additionally, we also noted that cell density influences the subcellular localization of Erdr1, it tends to localize in the cytoplasm at high density, while favors nuclear localization at low density (Supplemental Fig.1a and 1b). These findings reveal that basal and LPS-induced endogenous Erdr1 expression and subcellular localization were affected by cell density. Moreover, endogenous Erdr1 demonstrates a positive correlation with IL-1β at low cell density, while manifests a negative correlation with IL-1β at high cell densities.

Subsequently, we investigated the impact of supplementing Erdr1 protein on both IL-1β expression and cell viability under conditions of low and high cell densities. The data clearly reveal that Erdr1 protein exerts a dose-dependent enhancing effect on IL-1β expression (Fig. 3d) and facilitates cell death exclusively at low cell density (Fig. 3e). In contrast, at high cell density, Erdr1 displays a dose-dependent inhibitory role in IL-1β expression (Fig. 3g), while also promotes cell survive after LPS stimulation (Fig. 3h). These observations strongly suggest that the function of Erdr1 in IL-1β modulation is profoundly influenced by cell density. Overall, Erdr1 emerges as an amplifier of pro-inflammatory signaling under conditions of low cell density yet transits into an inhibitor of pro-inflammatory signaling when cell density is high.

We proceeded to investigate the effects of Erdr1 protein on endogenous Erdr1 expression. Intriguingly, our data reveal that Erdr1 protein treatment led to an upregulation of endogenous Erdr1 in both low and high cell density settings (Fig. 3f and 3i). Notably, upon LPS stimulation, endogenous Erdr1 exhibits an upregulation in low cell density scenarios, a response further

augments by Erdr1 protein treatment (Fig. 3f), ultimately intensifies IL-1β expression (Fig. 3d) and
aggravates cell death (Fig. 3e). In contrast, under conditions of high cell density, LPS stimulation
resulted in the downregulation of Erdr1, an effect that was counteracted by Erdr1 protein
treatment (Fig. 3i), subsequently inhibits IL-1β expression (Fig. 3g) and reduces cell death (Fig.
3h). Remarkably, even in the absence of LPS stimulation, Erdr1 exhibits the ability to upregulate
endogenous Erdr1 expression (Fig. 3f and 3i) and drive cell death (Fig 3e and 3h) both at low and
high cell density. These observations suggest endogenous Erdr1 plays a crucial role in
maintaining cellular homeostasis. It becomes apparent that controlled fluctuations of endogenous
Erdr1 level facilitate cellular survival. However, when Erdr1 levels veer to extremes- either too low
or excessively high- it becomes a mediator of cell death.
267
268 In summary, these data clearly demonstrate that the effect of Erdr1 on IL-1β expression is closely
269 correlated with cell density.
270
271 **Erdr1 promotes IL-1β production through activating YAP1 and Mid1 signaling pathway**
272 We further to explore the mechanism of unique promotion role of Erdr1 in regulating IL-1β
273 production. We hypothesized YAP1 and Mid1 are involved in this process.
274
To validate this hypothesis, we initially examined if and how YAP1 and Mid1 are regulated in
M1/M2 macrophages. Notably, we observed remarkable upregulation of Mid1 in mRNA level (Fig.
4a) and protein level (Fig. 4b) in M1 macrophages compare to M0 macrophages. We also
detected increased YAP1 expression in M1 macrophages at the protein level (Fig. 4b). In contrast,
we observed a significant decreased expression of Mid1 (Fig. 4a and Fig. 4b) and YAP1 (Fig 4b)
in M2 macrophages. These findings indicate that Mid1 and YAP1 are also distinct regulated in
M1/M2 macrophages. Utilizing immunofluorescence, we investigated the subcellular localization
of YAP1 and Mid1 in M1/M2 macrophages. YAP1 has increased nuclear localization in M1
macrophages, but increased cytoplasm localization in M2 macrophages (Fig. 4c). Whereas we
observed remarkable increased cytoplasm localization of Mid1 in M1 macrophages, but increased
nuclear localization in M2 macrophages, compared to M0 macrophages (Fig. 4d). We further
quantitated YAP1 and Mid1 Nuclear to Cytoplasm intensity ratio (Nuc/Cyt ratio) by the ImageJ,
data manifest YAP1 has increased Nuc/Cyt ratio in M1 macrophages, while decreased in M2
macrophages when compared to M0 macrophages (Fig. 4e). In the case of Mid1, there was a
dramatically reduction in the Nuc/Cyt ratio within M1 macrophages, but a significant elevation
within M2 macrophages when compared to M0 macrophages (Fig. 4f). These findings reveal that
Mid1 and YAP1 also have dynamic regulation and distinct subcellular localization patterns during
macrophage polarization.
293
We further investigated if YAP1 and Mid1 are involved in Erdr1-mediated LPS-induced IL-1β
production. Data demonstrate that knockout of Mid1 or YAP1 or both led to significant blockade
of IL-1β production in macrophages induced by LPS (Fig. 4f), indicating Mid1 and YAP1 are both
essential for IL-1β production in M1 macrophages. Moderate overexpression of Erdr1 (Erdr1 OE1)
has been demonstrated promote LPS-induced IL-1β production in macrophage (Fig. 2f), however,
this promotion role is effectively blocked in Mid1 knockout, YAP1 knockout, or Mid1/YAP1 double

knockout macrophages (Fig. 4g). These suggest Erdr1 plays roles in promoting IL-1β production
by YAP1 and Mid1 signaling pathways.
302
We next tested if Erdr1 is the upstream regulator of the YAP1 and Mid1 signaling. Our data display
that Erdr1 OE1 macrophage which enhances LPS-induced IL-1β production (Fig. 2f), also
promotes LPS-induced Mid1 and YAP1 expression (Fig. 4h and Fig. 4i). Whereas, in Erdr1
deficiency macrophage, which blocks IL-1β production, Mid1 and YAP1 expression are both
dramatic suppressed (Fig. 4h and Fig. 4i). These data manifest that Erdr1 regulates IL-1β
production is achieved by direct regulating YAP1 and Mid1 expression. Furthermore, we prove
other YAP1 and Mid1 downstream genes, such as CTGF and Cyr61 [38,47,48], are also regulated by
Erdr1 (Fig. 4j), with Erdr1 overexpression (Erdr1 OE1) promotes CTGF and Cyr61 expression,
but they are inhibited in Erdr1 deficiency macrophage (Fig. 4j), providing additional evidence that
Erdr1 acts as an upstream regulator of YAP1 and Mid1.
313
314 Taken together, these findings provide compelling evidence that Erdr1 plays a crucial role in
315 promoting IL-1β production by activating YAP1 and Mid1 signaling pathways in macrophages.
316
Erdr1 displays distinct interaction with YAP1 and Mid1 by different domains and mediates
macrophage polarization
We hypothesized Erdr1 plays dual role in IL-1β production by distinct and dynamic interaction
with YAP1 and Mid1. To investigate this, we employed Co-immunoprecipitation (Co-IP) to detect
the potential interaction of Erdr1 with YAP1 and Mid1 by using 293T cells. Specifically, we
hypothesized that the C-terminal domain of Erdr1 is the region responsible for interacting with
YAP1. This hypothesis is grounded in previous research, which has shown that the Erdr1 C-
terminal 81 amino acids are involved in contact inhibition [2]. Though protein sequence and function
prediction via UniProt, we predicted the C-terminal 32 amino acids are critical for YAP1 binding.
We observed full-length Erdr1 (Erdr1-177) has direct interaction with YAP1, whereas the Erdr1
variant which lacking the C-terminal 32 amino acids (Erdr1Δ32, or Erdr1-145) fails to exhibit this
interaction (Fig. 5a). These reveal Erdr1 has direct interaction with YAP1 by its C-terminal 32
amino acid. Furthermore, our data prove Mid1 could direct interaction with Erdr1-177 and even a
stronger interaction with Erdr1-145 in Co-IP experiments (Fig. 5b), suggesting Erdr1 has the
potential to interact with Mid1, particularly when it exposes its Erdr1-145 domain.
332
According to our hypothesis, by concentration change and the concomitant alterations in
conformation, Erdr1 has different affinity with YAP1 and Mid1. Erdr1- YAP1 high affinity or, Erdr1-
Mid1 high affinity, drive and polarize macrophage to two extremely inflammation states. Erdr1-
177 (high concentration) which interacts with YAP1 can be considered to harbor anti-inflammatory
conformation. Erdr1-145, which interacts with Mid1, can be considered to display pro-
inflammatory conformation. Erdr1-177 and Erdr1-145 potentially can drive macrophage polarize
into M2 and M1 states, respectively. To further investigate this hypothesis, we employed purified
Erdr1-177 and Erdr1-145 proteins to assess their roles in mediating macrophage polarization.
Our findings revealed that 1 μg/ml Erdr1-177 could spontaneously induce the expression of M2
marker genes, such as IL-10 and TGFβ (Supplemental Fig. 2a). Conversely, 1 μg/ml Erdr1-145

induced the expression of IL-6 and TNF, which are marker genes for M1 macrophages
(Supplemental Fig. 2b). Notably, we observed that Erdr1-145 exhibits a dose-dependent effect
on priming IL-1β production and could enhance LPS-induced IL-1β production (Fig. 5c). These
data provide support for the notion that Erdr1 has the potential to act as a suppressor or an
amplifier in modulating IL-1β production through conformational changes.

To investigate the interactions between Erdr1 and YAP1, as well as Erdr1 and Mid1 at the cellular
level, we conducted experiments involving the overexpression of Erdr1-177 or Erdr1-145 along
with YAP1 or Mid1 in Hela cells, with HA-tags and Flag-tags added at the C-terminus as indicated
(Fig. 5d and Fig. 5e). Notably, Erdr1-177-HA co-localized specifically with YAP1-Flag in the
cytoplasm, suggesting a direct interaction between these two proteins. Consistent with our Co-IP
data, we did not observe a direct interaction between Erdr1-145 and YAP1 (Fig. 5d). On the other
hand, we observed that both Erdr1-177 and Erdr1-145 interact with Mid1 at the nuclear envelope.
Notably, Erdr1-145, but not Erdr1-177, demonstrated the ability to activate Mid1 and induce its
translocation to the cytoplasm (Fig. 5e). In contrast, Erdr1-177 promoted the nuclear localization
of Mid1, which represents the inhibitory state of Mid1 (Fig. 5e). These data highlight the distinct
roles of different Erdr1 conformations in regulating YAP1 and Mid1, with Erdr1-177 licensing
YAP1 OFF/Mid OFF, while Erdr1-145 forcing Mid1 ON/YAP1 ON.

We further explored the impact of Erdr1-177 and Erdr1-145 on the subcellular localization of
endogenous YAP1 in macrophages. BMDM were overexpressed with Erdr1-177 or Erdr1-145 by
lentivirus transduction. We observed that the overexpression of Erdr1-177 results in the
sequestration of YAP1 in the cytoplasm (Fig. 5f) and induces a significant upregulated of IL-10
and TGF-β mRNA expression (Supplemental Fig. 2c), suggesting Erdr1-177 overexpression
mediates macrophage polarization towards the M2 phenotype. However, overexpression of
Erdr1-145 results in the nuclear localization of YAP1 (Fig. 5f) and a significant upregulation in the
expression of IL-6 and TNF (Supplemental Fig. 2d), indicating that overexpression of Erdr1-145
drives macrophage polarization towards the M1 phenotype. These findings provide substantial
evidence that Erdr1-177 polarizes macrophages towards the M2 state by regulating YAP1 OFF,
while Erdr1-145 polarizes macrophages towards the M1 state by regulating YAP1 ON.

Furthermore, we illustrated the dynamic co-localization of endogenous Erdr1 and YAP1 in
macrophages. Through immunofluorescence analysis, we detected a robust interaction between
Erdr1 and YAP1 in the cytoplasm of M2 macrophages, whereas such interaction was not
observed in M1 macrophages (Fig. 5g). These finding exhibit a distinct interaction pattern
between endogenous Erdr1 and YAP1 in M1 and M2 macrophages.

Collectively, these observations strongly support our hypothesis that the dynamic interaction of
Erdr1 with YAP1 and Mid1 drives and coordinates macrophage polarization.

Erdr1 orchestrates macrophage functional and metabolic reprogramming and determines
cell fate

Macrophage polarization is a vital functional response and adaptation to microenvironmental
stimuli [43,44]. Activated macrophage not only exhibits distinct functions but also involves metabolic
reprogramming [49] and ultimately have different cell death patterns [50-52]. To further elucidate the
role of Erdr1 in macrophage polarization, we further performed comprehensive analysis of Erdr1
role in inflammation, metabolic and cell death phenotype.

Erdr1-177 and Erdr1-145 protein, which represent different conformation of Erdr1, were observed
to play distinct roles in regulating IL-1β production at indicated doses (1 μg/ml) with optimal cell
density ($0.6*10^6$ cell/ml) (Fig. 6a and 6b). Erdr1-177 inhibits IL-1β production upon LPS
stimulation at mRNA level (Fig. 6a) and protein level in supernatant (Fig. 6b), while Erdr1-145
amplifies basal and LPS-induced IL-1β production (Fig. 6a and Fig. 6b). We validated these
findings using immunoblotting analysis, manifesting Erdr1-177 inhibits pro-IL-1β cleaved to
matured IL-1β, whereas Erdr1-145 promotes IL-1β maturation (Fig. 6d). Furthermore, we found
Erdr1-177 inhibits Mid1 and YAP1, but Erdr1-145 promotes Mid1 and YAP1 expression (Fig. 6c
and 6d), manifesting that Mid1 and YAP1 are also distinctly regulated by different Erdr1
conformations. These findings provide strong evidence that the highly dynamic fluctuations in
conformation of Erdr1 have the capacity to dynamically regulate the levels of IL-1β, YAP1, and
Mid1.

We then investigated if Erdr1 plays a role in mediating metabolic transition. Macrophage M1
polarization is characterized by the activation of mTORC1-S6K signaling [49], which represents an
anabolic state. In contrast, macrophage M2 polarization is characterized by the activation of
AMPK signaling [49], representing a catabolic state. Our findings demonstrated that Erdr1-177
protein inhibits p-S6K activation while promotes p-AMPK activation in basal and LPS-stimulated
macrophages (Fig. 6e). Conversely, Erdr1-145 protein strikingly promotes p-S6K activation but
inhibits p-AMPK activation in basal and LPS stimulated macrophage, compared to vehicle control
(Fig. 6e). These data support that Erdr1 has the potential to polarized macrophage metabolism
to pro-anabolic state or pro-catabolic state by conformation changes. Furthermore, we observed
that Erdr1 deficiency macrophage exhibits low basal and LPS-induced p-S6K and p-AMPK
activation, whereas Erdr1 overexpression macrophage displays high basal and LPS-induced p-
S6K and p-AMPK activation (Fig. 6g), suggesting Erdr1 is essential both for anabolic and
catabolic metabolism. These observations indicate that Erdr1 decides the metabolic activity,
phenotype, and transition within macrophages via concentration and conformational changes.

Furthermore, distinct activated macrophages undergo different cell death phenotype. M2
macrophage undergo apoptotic death, which elicit anti-inflammatory response. By contrast, M1
macrophage undergo pyroptosis, which is a proinflammatory form of cell death [50-52]. Our data
show that both Erdr1-177 and Erdr1-145 protein supplementation promote unstimulated
macrophages death (Supplemental Fig. 3a). However, Erdr1-177 protects LPS-induced
macrophage from death but Erdr1-145 aggravates LPS-induced cell death (Supplemental Fig.
3a). Our data further demonstrated that Erdr1-177 promotes apoptosis by enhancing the
activation of caspase3 and caspase8, while inhibiting pyroptosis by reducing caspase1 activation
in macrophages (Fig. 6f), indicating Erdr1-177 acts as a pro-apoptotic factor. Conversely, Erdr1-

145 inhibits apoptosis but promotes pyroptosis in unstimulated and LPS-stimulated macrophages
(Fig. 6f), supporting Erdr1 has the potential to be a pro-pyroptotic factor. Furthermore, we
observed Erdr1 knockout reduces, whereas Erdr1 overexpression enhances both caspase3/8
and caspase1 activation in macrophage compared to control (Fig. 6h), implying high level of Erdr1
is essential for caspase-dependent cell death process. Even lack of caspase-dependent
programmed cell death, we still found that Erdr1 deficiency in macrophages result in reduced cell
viability (Supplemental Fig. 3b). It is not hard to understand this considering both catabolic and
anabolic metabolism are blocked in Erdr1 deficiency macrophage (Fig. 6g), indicating basal
expression of Erdr1 is essential for cell survive. In summary, these findings suggest that a narrow
range of Erdr1 mediates cell survive, otherwise dysregulated Erdr1 determines macrophage
programmed cell death through either a pro-apoptotic or pro-pyroptotic pathway.
439
440 Taken together, these data reveal dynamic change of Erdr1 not only orchestrates inflammation
441 state, but also mediates metabolic transitions, and determines cell fate. These findings uncover
442 a crucial role of Erdr1 in determining macrophage polarization.
443

Discussion

445
This study aims to elucidate the function of Erdr1 in IL-1β production and macrophage
polarization. Erdr1 was discovered to exhibit a negative correlation with IL-18 [45] and purified Erdr1
protein played the anti-inflammatory role by inhibiting the production of various cytokines such as
TNF, IL-6, and IL-1β [7,46]. Our data also manifest the anti-inflammatory role of Erdr1. What
interesting is we uncover that Erdr1 has the potential to display pro-inflammatory role which highly
associated with Erdr1 dose and cell densities. At optimal cell density, both endogenous and
exogenous Erdr1 plays role in IL-1β production following a bell-shaped model. This Erdr1 dose-
dependent threshold model has also been noted in Erdr1-mediated metastasis in murine
embryonic fibroblasts (MEF) [34,53], indicating Erdr1 has highly dynamic functions.
455
We evidence that the dual role of Erdr1 in IL-1β production is achieved by Erdr1 endogenous
concentration change which displayed differently in M1/M2 macrophages. We found Erdr1 was
significantly upregulated in M2 macrophages, and might act as a M2 marker gene, which is
consistent with previous research [54]. It is worth noting that we observed Erdr1 was dramatically
suppressed in M1 macrophages at optimal cell density. Down-regulated Erdr1 was also observed
in other immune cells upon activation and non-immune cells. Erdr1 exhibited downregulation in
CD4 T cells upon TCR stimulation [5]. Erdr1 was also downregulated in tissues of patients with
inflammation disorder disease [8,9,46], tumor cell [45], and remarkably inhibited in bleomycin-induced
fibroblast [41]. These indicate that reduced Erdr1 level might represent a way to activate specific
functions of the cell. Our data support the idea that decreased Erdr1 undergoes conformation
change and promotes IL-1β production and drives the macrophage pro-inflammatory polarization,
whereas elevated Erdr1 inhibits IL-1β production and drives the macrophage anti-inflammatory
polarization.
469

We figure out Erdr1 drives macrophage polarization by distinct interaction with YAP1 or Mid1.
Consistent with previous research [24,26,38], we observed the significantly up-regulated expression
of YAP1 and Mid1 in pro-inflammatory macrophages. Notably, YAP1 and Mid1 are both essential
for Erdr1-mediated LPS-induced IL-1β production. We reveal Erdr1 acts as an upstream regulator
of YAP1 and Mid1, regulates their expression, subcellular localization, and downstream gene
expression, which includes IL-1β. Additionally, YAP1 and Mid1 have also been identified as
promoters of fibrosis [38,47]. Down-regulated Erdr1 was also reported to involve in the pathogenesis
of bleomycin-induced fibrosis in fibroblasts [41]. CTGF, the central mediator of fibrosis [55] and a
signature gene of YAP1 and Mid1 signaling [38,48], is significantly regulated by Erdr1 (Fig. 4j). This
indicates by regulating YAP1 and Mid1, Erdr1 might not only modulate inflammation but also
regulate fibrosis. Except for the role of Erdr1 in IL-1β, we also reveal Erdr1 distinctly drive M2
maker gene (IL-10 and TNFβ) or other M1 marker gene (IL-6 and TNF) expression. Our data
demonstrate that Erdr1 acts as the driving force for macrophage M2/M1 functional polarization:
1. Erdr1-YAP1 interaction drives anti-inflammatory M2 state; 2. Erdr1-Mid1 interaction drives pro-
inflammatory M1 state (Fig. 7).
485
Metabolic reprogramming stands as a pivotal hallmark of macrophage polarization. YAP1 and
Mid1 have both been documented to enhance pro-anabolic response [18,56-59]. Notably, Erdr1 has
been implicated in YAP1 and Mid1-mediated metabolism [18,40]. YAP1 has been reported to be
essential for mTOR signaling, and its deficiency in dental epithelial stem cells resulted in
decreased Erdr1 expression, ultimately leading to the inhibition of mTOR signaling [18]. Moreover,
elevated levels of Erdr1 and Mid1 have been observed to participate in unique metabolism
(dampened oxidative phosphorylation) in β-cells harboring the SLC30A8 R138X mutation [40].
These suggest Erdr1 might enhance mTOR signaling by working with YAP1 and Mid1, which is
consistent with our observation that Erdr1 by activating YAP1 and Mid1 signaling, has the pro-
anabolic capability by activating p-S6K. However, our data also manifest Erdr1 has the pro-
catabolic potential by activating p-AMPK, when at a high concentration. This phenomenon was
also observed in previous research in T cells when Erdr1 was upregulated to a high level [60]. In
fact, Erdr1 highly expressed T cells was found to have increased both glycolysis and oxidative
phosphorylation [60], which are downstream of mTORC1 and p-AMPK respectively [49,61], indicating
Erdr1 is involved in both anabolic and catabolic metabolism. Our data for the first time prove that
Erdr1 is essential for both anabolic and catabolic metabolism by Erdr1 knockout and
overexpression experiment (Fig.6g). However, the two different conformations of Erdr1 can only
mediate the metabolic polarization to pro-catabolic or pro-anabolic state, with antagonize each
other (Fig.6e). This represents by binding with YAP1 and Mid1 with different affinities, Erdr1 might
mediate the metabolic polarized to different extreme states. Supporting this, β cell which has
increased expression of Erdr1 and Mid1 exhibited increase insulin secretion but dampened
oxidative phosphorylation [40]. This further support Erdr1-Mid1 affinity has the potential to prime
mTORC1-glycolysis but inhibit p-AMPK-oxidative phosphorylation pathway. Collectively, our data
suggest Erdr1 mediates macrophage M2/M1 metabolic polarization: 1. Erdr1-YAP1 affinity primes
pro-catabolic response; 2. Erdr1-Mid1 affinity drives pro-anabolic response (Fig. 7). As we know,
p-AMPK activation promotes macrophage M2 polarization [62], while mTOR-p-S6K activation

promotes IL-1β production in M1 macrophages[63]. Thus, these further reinforce the idea that Erdr1
mediates the macrophage polarization also by driving metabolic transition.
514
YAP1 was reported to play important roles in regulating cell death phenotypes [64,65]. Activated
YAP1 restrains apoptosis [64,65], but potentiates inflammatory activation and promotes pyroptosis
in macrophage [25]. Conversely, YAP1 ablation intensifies caspase3/8/9- dependent cell death [66].
Previous studies reported Erdr1 acts as a pro-apoptotic factor via inducing caspase3 activation
[5,67]. Our data support that and suggest Erdr1 by driving YAP1 OFF, could induce caspase3 and
caspase8 activation and promote apoptosis (Fig. 6f and 6h). Erdr1 has also been associated with
caspase1 and involved in progressive degenerative disease [68]. Our data for the first time figure
out the pro-pyroptotic role of Erdr1 (Fig. 6f and 6h) by driving YAP1 ON. In Erdr1-deficient
macrophages, both apoptosis and pyroptosis were inhibited (Fig. 6h), thereby illustrating the
involvement of Erdr1 in both pro-apoptotic and pro-pyroptotic pathways. Erdr1 was observed to
only mediate cell survival at a narrow range, otherwise cause rapid cell death [53] . We also
observed this phenomenon in Erdr1 deficiency or overexpressed macrophages. They both elicits
cell death (Supplemental Fig. 3b) but manifest distinct cell death patterns, with Erdr1 deficiency
induces caspase-independent cell death, while Erdr1 overexpression causes caspase-dependent
cell death (Fig. 6h). These support that Erdr1 determines cell fate: 1. Erdr1-YAP1 affinity induces
pro-apoptotic response. 2. Erdr1-Mid1 affinity primes pro-pyroptotic activation. Additionally,
negative feedback mechanisms exist by heightened YAP1 or elevated extracellular accumulation
of Erdr1, which foster Erdr1-YAP1 interactions and consequently facilitate cells apoptotic death,
act to alleviate the inflammation. We emphasize that the fine and dynamic interaction between
Erdr1 and YAP1 or Mid1 play pivotal roles in facilitating cellular adaptation and survive. Otherwise,
a polarized interaction with YAP1 or Mid1 triggers cellular demise and decides the pattern of cell
death.
537
YAP1 serves as a downstream effector of the Hippo pathway which is a highly conserved
signaling pathway that governs organ size through the regulation of cell proliferation, apoptosis,
and stem cell self-renewal [27]. Over the years, extensive research has revealed that the Hippo ON
(YAP1 OFF) can be achieved through a sequential cascade of kinase activations [27], which
including the formation of a complex between Mst1/2 kinases and SAV1 and subsequently
phosphorylates and activates LATS1/2 kinases [27]. However, different from LATS1/2 kinases
dependent way in regulating YAP1, a unique YAP1 regulation were observed in Tregs cells which
exhibited altered levels of Erdr1 and Mid1[19]. As we know, Hippo ON can be achieved by high cell
density [17,69]. In this study, data support that cell density regulates Hippo ON/Hippo OFF achieved
by Erdr1 concentration change. Collectively, we have successfully demonstrated that Erdr1 acts
as an upstream regulator of YAP1 through direct interaction. Most importantly, we found that the
interaction with YAP1 is highly linked to Erdr1 concentration and concomitant conformational
shifts. These findings suggest the critical role of Erdr1 in regulating YAP1 through the non-
classical Hippo pathway.
552
However, the research progress of Erdr1 was hindered by several obstacles: Firstly, Erdr1 global
knockout resulted in 100% embryonic lethality [5,70], which limited research in mice. This also

indicates Erdr1 plays a crucial role in cell physiological processes. Secondly, although Erdr1 has
been mapped to sex chromosomes in mice, due to it has a high GC content [53] and contains many
repeats [32], making identifying the entire genome sequence of Erdr1 unavailable until recently [33].
Thirdly, although human Erdr1 mRNA and protein have been detected with 100% identical to mice
Erdr1 [2,53], as of now there is still lack of accessible human Erdr1 mRNA and protein information
within the NCBI database, attributed to the incomplete sequencing of the Erdr1 localization region
in humans [71], this hampers Erdr1 being identified as a pivotal target in humans for further
research.
563
In this study, at the cellular level, we demonstrate that Erdr1 plays a crucial role in orchestrating
macrophage functional and metabolic reprogramming and ultimately determining cell fate through
dynamic interaction with YAP1 and Mid1. Our research highlights the dynamic regulation of Erdr1
as the dominant factor for the macrophage spectrum oscillating between the M1 and M2 extreme
polarized states. Understanding the molecular mechanism of macrophage polarization
orchestrated by Erdr1 stands to profoundly enhance our ability to therapeutically modulate
inflammation more precisely and efficiently. Moreover, it's crucial to highlight our significant
observation that Erdr1 regulates YAP1 signaling through the non-classical Hippo pathway. These
findings imply the critical role of Erdr1 in maintaining cellular and tissue homeostasis, extending
beyond its impact on macrophages alone.
574
575 **Materials and Methods**
576
577 **Cell culture**
578 293T and Hela cells were cultured with DMEM medium in the presence of 10% FBS, 100 IU/ml
579 penicillin, and 100 μg/ml streptomycin. RAW246.7 were cultured with RPMI-1640 medium in the
580 presence of 10% FBS, 100 IU/ml penicillin, 100 μg/ml streptomycin.
581
Mouse BMDMs
C57BL/6 mice were killed, and femurs were removed. Bone marrow was harvested by flushing
with BMDM medium (RPMI-1640 medium supplemented with 10% FBS, antibiotics, l-glutamine,
and 10 μg/ml M-CSF). The harvested cells were centrifuged at 300 × g for 10 min and were
treated with ACK lysis buffer (ThermoFisher, A1049201) for 1 min to remove RBCs. ACK lysis
buffer was neutralized by adding 10 volumes of PBS, and cells were spun down at 300 × g for 10
min. The pelleted cells were resuspended in BMDM medium and were plated at $0.6*10^6$ cells/ml
for culture. The medium was replaced on days 4, 5, and 6 of culture before cell use on day 7. In
some experiments, BMDMs were treated with indicated concentrations of LPS, IL-4, or purified
Erdr1 proteins. Some experiment seeding different cell density as indicated. All animal studies
were approved by the University of Iowa Animal Care and Use Committee and meet stipulations
of the Guide for the Care and Use of Laboratory Animals.
594
Plasmid preparation and transduction
Full length Erdr1 (Erdr1-177) and Erdr1 deficiency C-terminal 32 amino acid (Erdr1-145) with C-
terminal HA tags was synthesized (GenScript) and cloned into pcDNA3.1 plasmid using In-Fusion

cloning (Clontech). YAP1 and Mid1 with C-terminal Flag were synthesized (GenScript) and cloned
into pcDNA3.1 plasmid using In-Fusion cloning (Clontech). Plasmids were sequenced before use.
For transient expression, plasmids were transfected into 293 T cell or Hela cell using
Lipofectamine™ 3000 Transfection Reagent (Life Technologies) according to the manufacturer's
instructions. Lentiviral vectors: For gene knockout, we prepared following transfer plasmid. Erdr1,
Mid1 and YAP1 Crispr/Cas9 knockout transfer plasmid were purchased from GenScript; For Erdr1
overexpression, full length Erdr1 (Erdr1-177) C-terminal HA tags were cloned into pHAGE vector.
Lentivirus were produced by transfer lentiviral vector (transfer plasmid, psPAX2 and pMD2.G) into
293T cells following transfection protocol. Once produced, lentivirus was used for infection
macrophage with different MOI.
608
609 **Cell viability assay**
610 Cell viability assessed by MTT assay. MTT assay kit was purchased from Abcam (ab211091) and
611 were performed according to the manufacturer's protocol.
612
Coimmunoprecipitation (Co-IP) assay
Cells were harvested with ice-cold PBS. Cell pellets were then lysed in immunoprecipitation buffer
(IP lysis buffer: 25 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% NP-40 and 5% glycerol,
1 mM DTT, and 1× protease/phosphatase inhibitors) for 30 min on ice and cleared by
centrifugation at 13,000 × g for 15 min. The Erdr1-HA&YAP1-Flag or Erdr1-HA&Mid1-Flag was
immunoprecipitated by Anti-FLAG® M2 Magnetic Beads for 2h, rotating end over end at 4°C.
After the beads were pelleted, washed three times with IP lysis buffer, the immunocomplexes
containing the Flag tagged interaction protein were then prior to elution with 2× SDS-sample buffer
for 5 min at 95°C, followed by Immunoblotting analysis.
622
Recombinant Erdr1
Generation of recombinant Erdr1 was performed by following protocol. Briefly, Erdr1-177 and
Erdr1-145 with C-terminal His-tag was cloned into the pET28a vector. E. coli was transformed
with this plasmid and cultured in medium containing ampicillin. IPTG was introduced for protein
induction. Erdr1-177-His and Erdr1-145-His was purified on a nickel column and purity was >95%
as confirmed by SDS-PAGE and Immunoblot. Potential endotoxin was removed using High-
Capacity Endotoxin Removing Spin Columns (88274, ThermoFisher).
630
631 **Anti-Erdr1 polyclonal antibody**
632 Anti-Erdr1 peptide Ab (36–51 amino acids of Erdr1, N-RAPRPPRHTRHTRHTR-C) was got as a
633 gift from Dr. June Round at Utah University[5]. Erdr1 neutralizing antibody were got as a gift from
634 Dr. Timothy L. Denning [12].
635
Immunofluorescence
Cells were seeded into 24 well plates containing 12 mm coated coverslips (Platinum Line). After
indicated culture or treatment times, cells were fixed by incubation with PBS containing 4% (w/v)
paraformaldehyde (Merck, #104005100) for 15 min at room temperature. After washing three
times with PBS for 5 min, fixed cells were blocked and permeabilized by incubation with

500 µl methanol for 5 min. PBS containing 5% (w/v) horse serum (Sigma, #A7906) were used for
blocking for 1 h at room temperature. Primary antibodies (Supplemental table S1) were diluted in
5% (w/v) horse serum/PBS and applied to coverslips and incubated at 4°C overnight. After three
washes with PBS, coverslips were incubated with secondary antibody diluted in 5% (w/v) horse
serum/PBS at room temperature for 1 h. Cells were washed three times with PBS. Cells were
stained nucleic and mounted onto glass slides using Antifade Mounting Medium with DAPI
(VECTASHIELD). Slides were viewed using Zeiss LSM 710 Confocal Microscope.
648

Immunoblotting
Total cell extracts were lysed using radioimmunoprecipitation (RIPA) assay buffer, which was
supplemented with Complete Mini Protease Inhibitor Cocktail and Phosphatase Inhibitor Cocktail
from Roche. The lysates were put on ice for a period of 30 minutes and vortexed every 5 minutes.
Following centrifugation at 15,000 rpm for 15 minutes at 4°C, the supernatants were collected.
Proteins were resolved on a SDS polyacrylamide gel and transferred to a 0.45-µm PVDF
membrane. For immunoblot analysis, the indicated primary antibodies were used and listed in
Supplemental table S1 and were used at 1000-fold dilution. Horseradish peroxidase (HRP)-
conjugated secondary antibodies were used depending on the host species of the primary
antibodies.
659

660 **Cytokine Quantification**
661 IL-1β quantification was performed by harvesting cell supernatants at the indicated time points
662 and treatment and assaying for IL-1β levels using a mouse IL-1β ELISA kit (DY401-050),
663 according to the manufacturer's protocol.
664

qPCR and primers
Total RNA was extracted from macrophages at the indicated time points and treatment using
TRIzol reagent (Invitrogen). 2 µg of RNA were used for cDNA synthesis. 2 µl of cDNA were added
to 23 µl of PCR mixture containing 2xSYBR Green Master Mix (ABI) and 0.2 µM of forward and
reverse primers. Amplification was performed in an ABI Prism 7500 thermocycler. Genes and
forward and reverse primers are shown in Supplemental table S2. Cycle threshold (Ct) values
were normalized to those of the housekeeping gene β-actin by the following equation: $\Delta Ct = Ct$
(gene of interest) - Ct (β-actin). All results are shown as a ratio to β-actin calculated as $2^{-(\Delta Ct)}$.
673

674 **Statistics.**
675 A Student's t test was used to analyze differences in mean values between groups. All results are
676 expressed as means ± SD. P values less than 0.05 were considered statistically significant; $*P <$
677 0.05; $**P < 0.01$; $***P < 0.001$, $****P < 0.0001$.
678

679 **Authors' Disclosures**
680 The author declares no conflict of interest.
681

682 **Authors' Contributions**

683　Y.W. developed the conception, designed, and performed the experiments, interpreted the data,
684　and wrote the manuscript.
685

686　**Acknowledgments**
687　This work is partially supported by the National Institutes of Health (NIH) Grant P01 AI060699.
688　Sincerely thank Dr. Stanley Perlman from The University of Iowa for his significant contribution to
689　the manuscript revision. Sincere appreciation to Dr. Ning Li from Beijing Capital Agribusiness Co.,
690　Ltd., for actively participating in the discussion and for contributing to the manuscript revision.
691

Reference

1　Dormer, P., Spitzer, E. & Moller, W. EDR is a stress-related survival factor from stroma and other
tissues acting on early haematopoietic progenitors (E-Mix). *Cytokine* **27**, 47-57 (2004).
https://doi.org:10.1016/j.cyto.2004.03.014
2　Dormer, P., Spitzer, E., Frankenberger, M. & Kremmer, E. Erythroid differentiation regulator
(EDR), a novel, highly conserved factor I. Induction of haemoglobin synthesis in
erythroleukaemic cells. *Cytokine* **26**, 231-242 (2004). https://doi.org:10.1016/j.cyto.2004.02.005
3　Kim, M. S. *et al.* Erythroid differentiation regulator 1 strengthens TCR signaling in thymocytes by
modulating calcium flux. *Cell Immunol* **336**, 28-33 (2019).
https://doi.org:10.1016/j.cellimm.2018.12.004
4　Kim, M. S. *et al.* Erythroid Differentiation Regulator 1 Strengthens TCR Signaling by Enhancing
PLCgamma1 Signal Transduction Pathway. *Int J Mol Sci* **23** (2022).
https://doi.org:10.3390/ijms23020844
5　Soto, R. *et al.* Microbiota promotes systemic T-cell survival through suppression of an apoptotic
factor. *Proc Natl Acad Sci U S A* **114**, 5497-5502 (2017).
https://doi.org:10.1073/pnas.1619336114
6　Lee, H. R. *et al.* ERDR1 enhances human NK cell cytotoxicity through an actin-regulated
degranulation-dependent pathway. *Cell Immunol* **292**, 78-84 (2014).
https://doi.org:10.1016/j.cellimm.2014.10.002
7　Kim, K. E. *et al.* Therapeutic effect of erythroid differentiation regulator 1 (Erdr1) on collagen-
induced arthritis in DBA/1J mouse. *Oncotarget* **7**, 76354-76361 (2016).
https://doi.org:10.18632/oncotarget.13047
8　Kim, M. *et al.* Recombinant erythroid differentiation regulator 1 inhibits both inflammation and
angiogenesis in a mouse model of rosacea. *Exp Dermatol* **24**, 680-685 (2015).
https://doi.org:10.1111/exd.12745
9　Kim, K. E. *et al.* Downregulation of erythroid differentiation regulator 1 (Erdr1) plays a critical
role in psoriasis pathogenesis. *Exp Dermatol* **25**, 570-572 (2016).
https://doi.org:10.1111/exd.13005
10　Zheng, D., Liwinski, T. & Elinav, E. Inflammasome activation and regulation: toward a better
understanding of complex mechanisms. *Cell Discov* **6**, 36 (2020).
https://doi.org:10.1038/s41421-020-0167-x
11　Kelley, N., Jeltema, D., Duan, Y. & He, Y. The NLRP3 Inflammasome: An Overview of Mechanisms
of Activation and Regulation. *Int J Mol Sci* **20** (2019). https://doi.org:10.3390/ijms20133328
12　Abo, H. *et al.* Erythroid differentiation regulator-1 induced by microbiota in early life drives
intestinal stem cell proliferation and regeneration. *Nat Commun* **11**, 513 (2020).
https://doi.org:10.1038/s41467-019-14258-z

13 Kim, M. S. *et al.* Erythroid Differentiation Regulator 1 Ameliorates Collagen-Induced Arthritis via
Activation of Regulatory T Cells. *Int J Mol Sci* **21** (2020). https://doi.org:10.3390/ijms21249555
14 Wu, J. M. F. *et al.* Prostaglandin E(2) Receptor 2 Modulates Macrophage Activity for Cardiac
Repair. *J Am Heart Assoc* **7**, e009216 (2018). https://doi.org:10.1161/JAHA.118.009216
15 Gumbiner, B. M. & Kim, N. G. The Hippo-YAP signaling pathway and contact inhibition of growth.
J Cell Sci **127**, 709-717 (2014). https://doi.org:10.1242/jcs.140103
16 Pavel, M. *et al.* Contact inhibition controls cell survival and proliferation via YAP/TAZ-autophagy
axis. *Nat Commun* **9**, 2961 (2018). https://doi.org:10.1038/s41467-018-05388-x
17 Zhao, B. *et al.* Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact
inhibition and tissue growth control. *Genes Dev* **21**, 2747-2761 (2007).
https://doi.org:10.1101/gad.1602907
18 Hu, J. K. *et al.* An FAK-YAP-mTOR Signaling Axis Regulates Stem Cell-Based Tissue Renewal in
Mice. *Cell Stem Cell* **21**, 91-106 e106 (2017). https://doi.org:10.1016/j.stem.2017.03.023
19 Ni, X. *et al.* YAP Is Essential for Treg-Mediated Suppression of Antitumor Immunity. *Cancer
Discov* **8**, 1026-1043 (2018). https://doi.org:10.1158/2159-8290.CD-17-1124
20 Hamon, A. *et al.* Linking YAP to Muller Glia Quiescence Exit in the Degenerative Retina. *Cell Rep
* **27**, 1712-1725 e1716 (2019). https://doi.org:10.1016/j.celrep.2019.04.045
21 Gong, S. *et al.* Endothelial Conditional Knockdown of NMMHC IIA (Nonmuscle Myosin Heavy
Chain IIA) Attenuates Blood-Brain Barrier Damage During Ischemia-Reperfusion Injury. *Stroke
* **52**, 1053-1064 (2021). https://doi.org:10.1161/STROKEAHA.120.031410
22 Lee, C. M. & Hu, J. Cell density during differentiation can alter the phenotype of bone marrow-
derived macrophages. *Cell Biosci* **3**, 30 (2013). https://doi.org:10.1186/2045-3701-3-30
23 Vaughan-Jackson, A. *et al.* Density dependent regulation of inflammatory responses in
macrophages. *Front Immunol* **13**, 895488 (2022). https://doi.org:10.3389/fimmu.2022.895488
24 Zhou, X. *et al.* YAP Aggravates Inflammatory Bowel Disease by Regulating M1/M2 Macrophage
Polarization and Gut Microbial Homeostasis. *Cell Rep* **27**, 1176-1189 e1175 (2019).
https://doi.org:10.1016/j.celrep.2019.03.028
25 Meli, V. S. *et al.* YAP-mediated mechanotransduction tunes the macrophage inflammatory
response. *Sci Adv* **6** (2020). https://doi.org:10.1126/sciadv.abb8471
26 Mia, M. M. *et al.* YAP/TAZ deficiency reprograms macrophage phenotype and improves infarct
healing and cardiac function after myocardial infarction. *PLoS Biol* **18**, e3000941 (2020).
https://doi.org:10.1371/journal.pbio.3000941
27 Ma, S., Meng, Z., Chen, R. & Guan, K. L. The Hippo Pathway: Biology and Pathophysiology. *Annu
Rev Biochem* **88**, 577-604 (2019). https://doi.org:10.1146/annurev-biochem-013118-111829
28 Lu, T. *et al.* X-linked microtubule-associated protein, Mid1, regulates axon development. *Proc
Natl Acad Sci U S A* **110**, 19131-19136 (2013). https://doi.org:10.1073/pnas.1303687110
29 Collison, A. M. *et al.* TRAIL signals through the ubiquitin ligase MID1 to promote pulmonary
fibrosis. *BMC Pulm Med* **19**, 31 (2019). https://doi.org:10.1186/s12890-019-0786-x
30 Chen, X. *et al.* Irradiated lung cancer cell-derived exosomes modulate macrophage polarization
by inhibiting MID1 via miR-4655-5p. *Mol Immunol* **155**, 58-68 (2023).
https://doi.org:10.1016/j.molimm.2023.01.009
31 Fang, M. *et al.* TRIM18 is a critical regulator of viral myocarditis and organ inflammation. *J
Biomed Sci* **29**, 55 (2022). https://doi.org:10.1186/s12929-022-00840-z
32 Chu, H. P. *et al.* PAR-TERRA directs homologous sex chromosome pairing. *Nat Struct Mol Biol* **24**,
620-631 (2017). https://doi.org:10.1038/nsmb.3432
33 Kasahara, T., Mekada, K., Abe, K., Ashworth, A. & Kato, T. Complete sequencing of the mouse
pseudoautosomal region, the most rapidly evolving 'chromosome'. *bioRxiv*,
2022.2003.2026.485930 (2022). https://doi.org:10.1101/2022.03.26.485930

34 Mango, R. L. *et al.* C-C chemokine receptor 5 on pulmonary mesenchymal cells promotes
experimental metastasis via the induction of erythroid differentiation regulator 1. *Mol Cancer*
Res **12**, 274-282 (2014). https://doi.org:10.1158/1541-7786.MCR-13-0164
35 Pagliari, S. *et al.* YAP-TEAD1 control of cytoskeleton dynamics and intracellular tension guides
human pluripotent stem cell mesoderm specification. *Cell Death Differ* **28**, 1193-1207 (2021).
https://doi.org:10.1038/s41418-020-00643-5
36 Kovačić, N. L. N. Mid1 is a novel mediator of subchondral bone resorption in antigen-induced
arthritis. (4th annual meeting of the European Calcified Tissue Society, 11., Budapest, Hungary,
2019).
37 Mohseni, M. *et al.* A genetic screen identifies an LKB1-MARK signalling axis controlling the
Hippo-YAP pathway. *Nat Cell Biol* **16**, 108-117 (2014). https://doi.org:10.1038/ncb2884
38 Chen, Q., Gao, C., Wang, M., Fei, X. & Zhao, N. TRIM18-Regulated STAT3 Signaling Pathway via
PTP1B Promotes Renal Epithelial-Mesenchymal Transition, Inflammation, and Fibrosis in
Diabetic Kidney Disease. *Front Physiol* **12**, 709506 (2021).
https://doi.org:10.3389/fphys.2021.709506
39 Perez, E. C. *et al.* The axis IL-10/claudin-10 is implicated in the modulation of aggressiveness of
melanoma cells by B-1 lymphocytes. *PLoS One* **12**, e0187333 (2017).
https://doi.org:10.1371/journal.pone.0187333
40 Kleiner, S. *et al.* Mice harboring the human SLC30A8 R138X loss-of-function mutation have
increased insulin secretory capacity. *Proc Natl Acad Sci U S A* **115**, E7642-E7649 (2018).
https://doi.org:10.1073/pnas.1721418115
41 Wang, Y. *et al.* The methyl-CpG-binding domain 2 facilitates pulmonary fibrosis by orchestrating
fibroblast to myofibroblast differentiation. *Eur Respir J* **60** (2022).
https://doi.org:10.1183/13993003.03697-2020
42 Wright, P. E. & Dyson, H. J. Intrinsically disordered proteins in cellular signalling and regulation.
Nat Rev Mol Cell Biol **16**, 18-29 (2015). https://doi.org:10.1038/nrm3920
43 Orecchioni, M., Ghosheh, Y., Pramod, A. B. & Ley, K. Macrophage Polarization: Different Gene
Signatures in M1(LPS+) vs. Classically and M2(LPS-) vs. Alternatively Activated Macrophages.
Front Immunol **10**, 1084 (2019). https://doi.org:10.3389/fimmu.2019.01084
44 Murray, P. J. Macrophage Polarization. *Annu Rev Physiol* **79**, 541-566 (2017).
https://doi.org:10.1146/annurev-physiol-022516-034339
45 Jung, M. K. *et al.* Erythroid differentiation regulator 1, an interleukin 18-regulated gene, acts as a
metastasis suppressor in melanoma. *J Invest Dermatol* **131**, 2096-2104 (2011).
https://doi.org:10.1038/jid.2011.170
46 Houh, Y. K., Kim, K. E., Park, H. J. & Cho, D. Roles of Erythroid Differentiation Regulator 1 (Erdr1)
on Inflammatory Skin Diseases. *Int J Mol Sci* **17** (2016). https://doi.org:10.3390/ijms17122059
47 Liu, F. *et al.* Mechanosignaling through YAP and TAZ drives fibroblast activation and fibrosis. *Am*
J Physiol Lung Cell Mol Physiol **308**, L344-357 (2015).
https://doi.org:10.1152/ajplung.00300.2014
48 Shome, D., von Woedtke, T., Riedel, K. & Masur, K. The HIPPO Transducer YAP and Its Targets
CTGF and Cyr61 Drive a Paracrine Signalling in Cold Atmospheric Plasma-Mediated Wound
Healing. *Oxid Med Cell Longev* **2020**, 4910280 (2020). https://doi.org:10.1155/2020/4910280
49 Kelly, B. & O'Neill, L. A. Metabolic reprogramming in macrophages and dendritic cells in innate
immunity. *Cell Res* **25**, 771-784 (2015). https://doi.org:10.1038/cr.2015.68
50 Cong, Y. *et al.* Macrophages in aseptic loosening: Characteristics, functions, and mechanisms.
Front Immunol **14**, 1122057 (2023). https://doi.org:10.3389/fimmu.2023.1122057

51   Kloditz, K. & Fadeel, B. Three cell deaths and a funeral: macrophage clearance of cells
undergoing distinct modes of cell death. *Cell Death Discov* **5**, 65 (2019).
https://doi.org:10.1038/s41420-019-0146-x

52   Robinson, N. *et al.* Programmed necrotic cell death of macrophages: Focus on pyroptosis,
necroptosis, and parthanatos. *Redox Biol* **26**, 101239 (2019).
https://doi.org:10.1016/j.redox.2019.101239

53   Mango, R. L. *Stromal promotion of metastasis by erythroid differentiation regulator 1*, University
of North Carolina at Chapel Hill, (2010).

54   Lee, D. C. *et al.* Aging enhances classical activation but mitigates alternative activation in the
central nervous system. *Neurobiol Aging* **34**, 1610-1620 (2013).
https://doi.org:10.1016/j.neurobiolaging.2012.12.014

55   Lipson, K. E., Wong, C., Teng, Y. & Spong, S. CTGF is a central mediator of tissue remodeling and
fibrosis and its inhibition can reverse the process of fibrosis. *Fibrogenesis Tissue Repair* **5**, S24
(2012). https://doi.org:10.1186/1755-1536-5-S1-S24

56   Liu, E., Knutzen, C. A., Krauss, S., Schweiger, S. & Chiang, G. G. Control of mTORC1 signaling by
the Opitz syndrome protein MID1. *Proc Natl Acad Sci U S A* **108**, 8680-8685 (2011).
https://doi.org:10.1073/pnas.1100131108

57   Posey, K. L. *et al.* Novel mTORC1 Mechanism Suggests Therapeutic Targets for COMPopathies.
Am J Pathol **189**, 132-146 (2019). https://doi.org:10.1016/j.ajpath.2018.09.008

58   Boding, L. *et al.* Midline 1 directs lytic granule exocytosis and cytotoxicity of mouse killer T cells.
Eur J Immunol **44**, 3109-3118 (2014). https://doi.org:10.1002/eji.201344388

59   Matthes, F. *et al.* Inhibition of the MID1 protein complex: a novel approach targeting APP
protein synthesis. *Cell Death Discov* **4**, 4 (2018). https://doi.org:10.1038/s41420-017-0003-8

60   Muri, J. *et al.* The thioredoxin-1 system is essential for fueling DNA synthesis during T-cell
metabolic reprogramming and proliferation. *Nat Commun* **9**, 1851 (2018).
https://doi.org:10.1038/s41467-018-04274-w

61   Wculek, S. K., Dunphy, G., Heras-Murillo, I., Mastrangelo, A. & Sancho, D. Metabolism of tissue
macrophages in homeostasis and pathology. *Cell Mol Immunol* **19**, 384-408 (2022).
https://doi.org:10.1038/s41423-021-00791-9

62   Sag, D., Carling, D., Stout, R. D. & Suttles, J. Adenosine 5'-monophosphate-activated protein
kinase promotes macrophage polarization to an anti-inflammatory functional phenotype. *J*
Immunol **181**, 8633-8641 (2008). https://doi.org:10.4049/jimmunol.181.12.8633

63   Kim, B. *et al.* Cytoplasmic zinc promotes IL-1beta production by monocytes and macrophages
through mTORC1-induced glycolysis in rheumatoid arthritis. *Sci Signal* **15**, eabi7400 (2022).
https://doi.org:10.1126/scisignal.abi7400

64   Cheng, Y., Mao, M. & Lu, Y. The biology of YAP in programmed cell death. *Biomark Res* **10**, 34
(2022). https://doi.org:10.1186/s40364-022-00365-5

65   Zhang, X., Abdelrahman, A., Vollmar, B. & Zechner, D. The Ambivalent Function of YAP in
Apoptosis and Cancer. *Int J Mol Sci* **19** (2018). https://doi.org:10.3390/ijms19123770

66   LeBlanc, L. *et al.* Yap1 safeguards mouse embryonic stem cells from excessive apoptosis during
differentiation. *Elife* **7** (2018). https://doi.org:10.7554/eLife.40167

67   Kim, H. J. *et al.* Erythroid differentiation regulator 1 (Erdr1) is a proapototic factor in human
keratinocytes. *Exp Dermatol* **20**, 920-925 (2011). https://doi.org:10.1111/j.1600-
0625.2011.01354.x

68   Lee, J. H., Kim, C. H., Kim, D. G. & Ahn, Y. S. Microarray analysis of differentially expressed genes
in the brains of tubby mice. *Korean J Physiol Pharmacol* **13**, 91-97 (2009).
https://doi.org:10.4196/kjpp.2009.13.2.91

869 69 Yu, F. X., Zhao, B. & Guan, K. L. Hippo Pathway in Organ Size Control, Tissue Homeostasis, and
870   Cancer. *Cell* **163**, 811-828 (2015). https://doi.org:10.1016/j.cell.2015.10.044
871 70 Zuo, E. *et al.* One-step generation of complete gene knockout mice and monkeys by
872   CRISPR/Cas9-mediated gene editing with multiple sgRNAs. *Cell Res* **27**, 933-945 (2017).
873   https://doi.org:10.1038/cr.2017.81
874 71 Rhie, A. *et al.* The complete sequence of a human Y chromosome. *Nature* (2023).
875   https://doi.org:10.1038/s41586-023-06457-y

 Figures and Figures Legends

Fig. 1 Erdr1 is distinctly regulated in M1/M2 macrophage.
M0 macrophage used in this study was Bone marrow-derived macrophage (BMDM). Cell was seeded with density $0.6*10^6$ cells/ml in 24 well plates. M0 macrophage was induced into M1 macrophage by treating with 1 μg/ml LPS for 24 hours, or into M2 macrophage by treating with 20 ng/ml IL-4 for 24 hours. The same experiments were also conducted and got repeatable results in RAW246.7 cell. **a,** qPCR analysis of Erdr1 mRNA expression in M0, M1, and M2 macrophage, qPCR normalized to β-actin expression (n=4). **b,** Immunoblot analysis of Erdr1 level in cell lysates and supernatant of M0, M1, and M2 macrophage. β-actin as the internal control. **c,** The Immunofluorescence staining of Erdr1 in M0, M1, and M2 macrophage and counterstained by DAPI. The white arrows in the M1 macrophage display Erdr1 is mainly localized in nuclear. The white arrows in the M2 macrophage display Erdr1 is mainly localized in cytoplasm. Scale bar in represents 20 μm. **d,** Erdr1 Nuclear to Cytoplasm (Neu/Cyt) intensity ratio was quantified by ImageJ based on immunostaining. n = 5 with more than 10 macrophages were analyzed per sample. **e,** Curve Fitting correlation of Erdr1&IL-1β fold change and Erdr1&IL-18-fold change using Nonlinear Regression (n=4). Data represent mean ± SD. *$p<0.05$, **$p<0.01$, ****$p<0.001$. Data are representative of at least three independent experiments.

Fig. 2 Erdr1 has bell-shape effect in LPS induced IL-1β production in macrophages.
a,b,c, Knockout of Erdr1 in BMDM by Lentivirus transduction for 40 hours, and then treated without or with 1 µg/ml LPS for 24 hours for the following analysis. Erdr1 KO1 and Erdr1 KO2 represent two knockout vectors, the vehicle is transfected with the control Lentivirus vector (n=4). **a,** qPCR detecting Erdr1 expression after lentivirus transduction for 40 hours. **b,** qPCR detecting IL-1β expression. **c,** Detecting IL-1β production at supernatant by ELISA. **d,e,f,** Overexpression of Erdr1 in BMDM by Lentivirus transduction for 40 hours and then treated without or with 1 µg/ml LPS for 24 hours for the following analysis. Erdr1 OE1 and Erdr1 OE2 represent two transfection titers of Lentivirus, the vehicle is transfected with the control Lentivirus vector (n=4). **d,** qPCR detecting Erdr1 expression after lentivirus transduction for 40 hours. **e,** qPCR detecting IL-1β expression. **f,** Detecting IL-1β production at supernatant by ELISA. **g,** The correlation of endogenous Erdr1 level (qPCR) and IL-1β level (ELISA) summary from (**c**) and (**f**). **h,** Detecting IL-1β level at supernatant by ELISA in 1 µg/ml LPS-induced BMDM with the indicated concentration of Erdr1 antibodies treatment (n=4). **i,** Detecting IL-1β level at supernatant by ELISA in 1 µg/ml LPS-induced BMDM with the indicated concentration of purified full-length Erdr1 protein treatment (n=4). **j,** Threshold Model of Erdr1 in IL-1β Production: A Bell-Shaped Function in IL-1β Production. 1,2, low level of Erdr1 has positive correlation with IL-1β. 3, The optimal Erdr1 concentration promotes the peak expression of IL-1β. 4,5, high level of Erdr1 has negative correlation with IL-1β level. These show Erdr1 has both positive and negative roles in IL-1β production by dose change. **a-i,** Cell density is 0.6×10^6/ml and seeded in 24 well plates. All qPCR normalized to β-actin expression. Data represent mean ± SD. *p<0.05, **p<0.01, ***p<0.001 and, ****p<0.0001. Data are representative of three independent experiments.

956
957

Fig. 3 Erdr1 effect in IL-1β production correlates with cell density in macrophages.

a,b,c, BMDM were seeded with the indicated cell density: Low density (Low, 0.1*10^6 cells/ml), Moderate density (Mod, 0.5*10^6 cells/ml), or High density (High, 2.5*10^6 cells/ml) in 24 well plates for 8 hours, and then treated without or with 1 µg/ml LPS for 24 hours or 48 hours (only for MTT assay) and perform following analysis: **a,** Detecting IL-1β expression by qPCR (n=4). **b,** Detecting cell viability by MTT assay (n=6). **c.** Detecting Erdr1 expression by qPCR (n=4). **d,e,f,** BMDM were seeded with Low cell density (0.1*10^6 cells/ml) in 24 well plates for 8 hours, and then treated without or with 1 µg/ml LPS for 24 hours together with supplemented with the indicated concentration of Erdr1 (0, 0.001, 0.01, 0.1, 1, 10 µg/ml), and perform following analysis: **d,** Detecting IL-1β expression by qPCR (n=4). **e,** Detecting cell viability by MTT assay (n=6). **f.** Detecting Erdr1 expression by qPCR (n=4). **g,h,i,** BMDM were seeded with High cell density (2.5*10^6 cells/ml) in 24 well plates for 8 hours, and then treated without or with 1 µg/ml LPS for 24 hours together with supplemented with the indicated concentration of Erdr1 (0, 0.001, 0.01, 0.1, 1, 10 µg/ml), and perform following analysis: **g,** Detecting IL-1β expression by qPCR (n=4). **h,** Detecting cell viability by MTT assay (n=6). **i.** Detecting Erdr1 expression by qPCR (n=4). All qPCR normalized to β-actin expression. The same experiments were also conducted and got repeatable results in RAW246.7 cell. Data represent mean ± SD. *p<0.05, **p<0.01, ***p<0.001 and, ****p<0.0001. Data are representative of two independent experiments.

956
957
978
979

Fig. 4 Erdr1 modulates IL-1β production through regulation of YAP1 and Mid1 signaling pathways.

a, qPCR analysis of Mid1 expression in M0, M1, and M2 macrophage (n=4). **b,** Immunoblot analysis of YAP1 and Mid1 levels in the cell lysates of M0, M1, and M2 macrophage. **c,d,** The Immunofluorescence staining of YAP1 (**c**) and Mid1 (**d**) in M0, M1, and M2 macrophage and counterstained by DAPI. Scale bar in represents 20 μm. **e,f,** Nuclear to Cytoplasm (Neu/Cyt) intensity ratios YAP1 (**e**) and Mid1 (**f**) were quantified by ImageJ. n = 5 with more than 10 macrophages were analyzed per sample. **g,** Detecting IL-1β production by ELISA in indicated BMDM group before and after 1 μg/ml LPS stimulation for 24 hours (n=4). Mid1 KO, YAP1 KO, and Double KO represent knockout Mid1, YAP1, and both Mid1/YAP1 in BMDM by transfection CRISPR/cas9 Lentivirus; Erdr1 OE1 represent moderate overexpression of Erdr1 (about 4 folds) in BMDM by transfection Lentivirus; the vehicle is transfected with the control Lentivirus vector (n=4). **h,i,j,** BMDM were knockout Erdr1 (Erdr1 KO1) or moderate overexpressed Erdr1 (Erdr1 OE1), then treated without or with 1 μg/ml LPS for 24 hours for following analysis. **h,** qPCR analysis of Mid1 expression(n=4); **i,** Immunoblot analysis of YAP1 and Mid1 expression. **j,** qPCR analysis of CTGF (left) and Cyr61 (right) (n=4). All qPCR normalized to β-actin expression. (A-J) Cell density is 0.6*10^6/ml and seeded in 24 well plates. Data represent mean ± SD. *p<0.05, **p<0.01, ***p<0.001 and, ****p<0.0001. Data are representative of two or three independent experiments.

1000
1001

1002

Fig. 5 Erdr1 displays distinct interaction with YAP1 and Mid1 by different domains and mediates macrophage polarization.

a, b, Co-immunoprecipitation (Co-IP) and immunoblot analysis. Flag tag protein: YAP-Flag (**a**), Mid1-Flag (**b**), and HA-tag protein (Erdr1-177-HA and Erdr1-145-HA) were co-expressed in 293 T cells for 24 hours and cell lysates were subjected to IP using Flag beads. Both inputs and Co-IP fractions (Output) were subjected to Immunoblotting with anti-Flag or anti-HA antibodies as indicated. Erdr1-177, Full-length Erdr1; Erdr1-145, Deficiency in c-terminal 32 amino acid domain. Flag tag and HA tag were added at the C-terminal of the gene. **c,** Detecting IL-1β production at supernatant with the indicated concentration of Erdr1-145 treatment in BMDM without or with 1 µg/ml LPS stimulation for 24 hours (n=4). **d,** Co-overexpression of YAP1-Flag with Erdr1-177-HA or Erdr1-145-HA in Hela cell and fluorescence staining using anti-HA and anti-Flag antibodies. White arrows point to the direct interaction of Erdr1-177-HA with YAP1-Flag in the cytoplasm. **e,** Co-overexpression of Mid1-Flag with Erdr1-177-HA or Erdr1-145-HA in Hela cell and fluorescence staining using anti-HA and anti-Flag antibodies. Red arrows point to the direct interaction of Mid1-Flag and Erdr1-177-HA/Erdr1-145-HA at the nuclear membrane. White arrows indicated Erdr1-145 promoted Mid1 localized at cytoplasm. **f,** Immunofluorescence staining of overexpressed Erdr1-177-HA or Erdr1-145-HA by anti-HA antibody and endogenous YAP1 in BMDM. White arrows indicated the direct interaction of Erdr1-177 with YAP1 in the cytoplasm. **g,** Immunofluorescence staining of endogenous Erdr1 and YAP1 in M0, M1, and M2 macrophages. Scale bar was marked in (**d**), it in represents 20 µm. Data represent mean ± SD. *p<0.05, ***p<0.001, ****p<0.001. Data are representative of two independent experiments.

1024

1025

1026

Fig. 6 Erdr1 mediates macrophage functional and metabolic polarization and determines cell death phenotypes.

a-f, BMDM were treated as follows for 24 hours: 1. Vehicle; 2. Erdr1-177 (1µg/ml) 3. Erdr1-145 (1µg/ml); 4. Vehicle+ LPS (1µg/ml); 5. Erdr1-177 (1µg/ml) + LPS (1µg/ml); 6. Erdr1-145 (1µg/ml) + LPS (1µg/ml), Cell lysates and supernatant were collected for the following analysis. **a,** qPCR analysis of IL-1β expression in the cell lysates (n=4). **b,** Detecting IL-1β production at supernatant by ELISA (n=4). **c,** qPCR analysis of Mid1 expression in the cell lysates. (**a**) and (**c**) qPCR normalized to β-actin expression. **d,** Immunoblot analysis of IL-1β, YAP1, and Mid1 levels in cell lysates and IL-1β in the supernatant; p-IL-1β, pro-IL-1β; c-IL-1β, cleaved-IL-1β. **e,** Immunoblot analysis of key proteins involved in metabolism as indicated. **f,** Immunoblot analysis of key proteins involved in apoptosis and pyroptosis as indicated. **g, h,** BMDM were genetic knockout of Erdr1 (Erdr1 KO) or overexpressed Erdr1 (Erdr1 OE) by lentivirus transduction, the vehicle represents transduction with vehicle control lentivirus. Cell lysates were collected for immunoblot analysis with indicated antibodies without and with 1µg/ml LPS treatment for 24 hours. p-S6K, phospho-S6K; t-S6K, total S6K; p-AMPK, phospho-AMPK; t-AMPK, total AMPK; p-Caspase1, pro-caspase1; c-Caspase1, cleaved caspase1; p-Caspase3, pro-caspase3; c-caspase3, cleaved caspase3; c-caspase8, cleaved caspase8. **a-h,** Cell density is 0.6*10^6/ml and seeded in 24 well plates. Data represent mean ± SD. *p<0.05, **p<0.01, ***p<0.001 and, ****p<0.0001. Data are representative of three independent experiments.

1046

1047

Fig. 7 Working model of Erdr1 in macrophages

Erdr1 has concentration and conformation change during macrophage polarization and plays multifunction in M1 and M2 macrophage. **a,** The M0 macrophage represents the steady state of macrophage with basal levels of Erdr1, Mid1, and YAP1 expression, accompanied by a basal metabolism level. Erdr1 acts as a cell survival factor. **b,** M0 macrophages are induced to M2 macrophage with 20 ng/ml IL-4 treatment for 24 hours. In M2 macrophages, Erdr1 is endogenously upregulated, with increased cytoplasm localization and increased secretion to the extracellular matrix. The upregulated Erdr1 interacts with YAP1 through its C-terminal domain (Erdr1-CBD), which consists of 32 amino acids. This interaction sequesters YAP1 in the cytoplasm and reduces YAP1 expression (YAP OFF), consequently leading to Mid1 inactivation (Mid1 OFF). Erdr1 acts as an anti-inflammatory, pro-catabolic and pro-apoptotic factor. **c,** M1 macrophages are induced from M0 macrophages by 1 µg/ml LPS treatment for 24 hours. In M1 macrophages, Erdr1 is endogenously downregulated and increased nuclear localization but increased release to extracellular matrix. The downregulated Erdr1 undergoes conformation shift and disconnecting with YAP1 but exhibits a high affinity with Mid1 through its Erdr1-145 domain (Erdr1-145, or Erdr1Δ32) at the nuclear envelope and force YAP1 ON. Erdr1 acts as a pro-inflammatory, pro-anabolic and pro-pyroptotic factor. Additionally, negative feedback regulation can be initiated in M1 macrophage by up-regulated endogenous YAP1 and increased accumulation of secreted Erdr1. This figure illustrates the main expression level (indicated by area size) and subcellular localization of Erdr1, Mid1, and YAP1 in M0, M1 and M2 macrophages. This represents situations at optimal cell density.

1069