Multimodal subspace independent vector analysis effectively captures the latent relationships between brain structure and function

Xinhui Li,[1,2*] Peter Kochunov,[3] Tulay Adali,[4] Rogers F. Silva,[1**] Vince D. Calhoun[1,2**]

[1]Tri-institutional Center for Translational Research in Neuroimaging and Data Science,

Georgia State University, Georgia Institute of Technology, Emory University, Atlanta, GA, USA

[2]School of Electrical and Computer Engineering, Georgia Institute of Technology, Atlanta, GA, USA

[3]Maryland Psychiatric Research Center, Department of Psychiatry, School of Medicine,

University of Maryland, Baltimore, MD, USA

[4]Department of Computer Science and Electrical Engineering,

University of Maryland Baltimore County, Baltimore, MD, USA

*Correspondence: xinhuili@gatech.edu

**These authors jointly supervised and equally contributed to this work.

March 2, 2026

Abstract: A key challenge in neuroscience is to infer the relationships between brain structure and function from high-dimensional, multimodal neuroimaging data. While conventional multivariate approaches often simplify statistical assumptions and estimate one-dimensional independent sources shared across modalities, the true relationships between latent sources are likely more complex—statistical dependence may exist both within and between modalities, and possibly span more than one dimension per modality. Here we present Multimodal Subspace Independent Vector Analysis (MSIVA), a methodology to capture both joint and unique vector sources from multiple data modalities by defining both cross-modal and unimodal subspaces with variable dimensions. In particular, MSIVA enables flexible estimation of varying-size independent subspaces within modalities and their one-to-one linkage to cor-

responding subspaces across modalities. A key advantage is its ability to capture subject-level variability at the voxel level within independent subspaces, in contrast to traditional methods that share the same independent components across subjects. We evaluated three initialization workflows with five candidate subspace structures in multiple synthetic datasets and two large multimodal neuroimaging datasets, both including structural MRI (sMRI) and functional MRI (fMRI). After confirming that MSIVA successfully recovered the ground-truth subspace structures in the synthetic data, we then applied MSIVA to identify the latent subspace structure in the neuroimaging data. Subsequent subspace-specific canonical correlation analysis, brain-phenotype prediction, and voxelwise brain-age delta analysis suggest that the estimated sources from MSIVA with the optimal subspace structure are strongly associated with multiple phenotype variables, including age, sex, schizophrenia, lifestyle factors, and cognitive functions. Further, we identified modality- and group-specific brain regions related to multiple phenotype measures such as age (for example, cerebellum, precentral gyrus, and cingulate gyrus in sMRI; occipital lobe and superior frontal gyrus in fMRI), sex (for example, cerebellum in sMRI, frontal lobe in fMRI, and precuneus in both sMRI and fMRI), schizophrenia (for example, cerebellar, frontal, and insular cortices in sMRI; occipital pole, lingual gyrus, and precuneus in fMRI), shedding light on phenotypic and neuropsychiatric biomarkers of linked brain structure and function.

Keywords: multimodal fusion; latent variable models; structural and functional brain imaging; age; sex; schizophrenia

1   Introduction

Neuroimaging techniques such as magnetic resonance imaging (MRI) have been developed to understand the structural and functional properties of the brain, as well as their relationships to behavior. However, it is challenging to directly associate behavior measures with raw MRI data, which typically includes tens of thousands of voxels and subjects. Although the data in its original space appears complex, its intrinsic dimensionality can be significantly lower. Recent studies have found that neural representations in low-dimensional subspaces form the basis that supports motor functions such as reaching (Churchland et al., 2012; Pandarinath et al., 2018) and timing (Remington et al., 2018; Wang et al., 2018), and cognitive functions such as perception (Bao et al., 2020; Chang & Tsao, 2017; Semedo et al., 2019; She et al., 2024), generalization (Bernardi et al., 2020; Boyle et al., 2024; Courellis et al., 2024), and decision-making (Hajnal et al., 2024; Johnston et al., 2024). Hence, it is important to develop latent

variable models to learn low-dimensional representations and structures from high-dimensional data. In addition, each neuroimaging modality has its own strengths and weaknesses, and only captures limited information about the brain. For example, structural MRI (sMRI) provides high-resolution anatomical structure of the brain but does not capture temporal dynamics, while functional MRI (fMRI) measures blood-oxygenation-level-dependent (BOLD) signals over time at the cost of lower spatial resolution. Joint analysis of sMRI and fMRI can offer rich spatio-temporal information in the brain that is not captured by a single modality. With the increasing availability of multimodal neuroimaging datasets, it is necessary to develop multivariate approaches to effectively capture interpretable and multifaceted information about the brain and its disorders from multiple imaging modalities (Calhoun & Sui, 2016; Lahat et al., 2015; Sui et al., 2012; Zhang et al., 2020).

A variety of data-driven multivariate approaches have been developed to jointly analyze multiple neuroimaging datasets or data modalities, including joint independent component analysis (jICA) (Calhoun & Adali, 2008; Calhoun, Adali, Giuliani, et al., 2006; Calhoun, Adali, Pearlson, & Kiehl, 2006; Franco et al., 2008), linked ICA (Groves et al., 2011), multimodal canonical correlation analysis (mCCA) (Correa et al., 2008, 2010; Mohammadi-Nejad et al., 2017), jICA+mCCA (Sui et al., 2011, 2013), and independent vector analysis (IVA) (Adali et al., 2015a, 2015b). Notably, a unified framework Multidataset Independent Subspace Analysis (MISA) (Silva et al., 2020) has recently been introduced, encompassing multiple latent variable models, such as ICA (Comon, 1994), IVA (Adali et al., 2014; Kim et al., 2006), and independent subspace analysis (ISA) (Cardoso, 1998; Hyvärinen & Hoyer, 2000; Szabó et al., 2012). MISA can be applied to identify latent sources from multiple neuroimaging modalities, including sMRI and fMRI (Silva et al., 2020). More recently, a multimodal IVA (MMIVA) fusion method built upon MISA has been proposed to identify linked biomarkers related to age, sex, cognition, and psychosis in two large multimodal neuroimaging datasets (Silva et al., 2024). Many existing approaches, including MMIVA, assume that sources are one-dimensional and independent within each modality, that is, the subspace structure is an identity matrix. However, brain networks are hierarchically organized across multiple spatial and temporal scales (Betzel & Bassett, 2017; Felleman & Van Essen, 1991; Friston, 2008; Kiebel et al., 2008; Zeki & Shipp, 1988). This hierarchical organization implies that the underlying relationships between true latent sources are likely more complex—statistical dependence may exist both within and across modalities, and span more than one dimension per modality. Sources from the same modality may be linked, potentially grouped by their anatomical or functional properties. For instance, spatial dependencies between corresponding sources have been observed in task-based and resting-state fMRI studies (Ma et al., 2010, 2011; McKeown et al., 1998), which MMIVA would not optimally capture.

Aiming to better detect the statistical relationships from multimodal data, we present a novel method-

ology, Multimodal Subspace Independent Vector Analysis (MSIVA)[1], that captures linkage of vector sources by defining cross-modal and unimodal subspaces with variable dimensions. MSIVA is built upon MMIVA by defining a block diagonal matrix as the subspace structure, instead of the identity matrix used in MMIVA. We default the initialization of MSIVA to the weight matrices learned by multimodal group principal component analysis (MGPCA) and unimodal ICAs combined. MSIVA can simultaneously estimate two types of latent sources—those linked across all modalities (along with their underlying relationships) and those unique to a specific modality. Moreover, by leveraging higher-dimensional subspaces, MSIVA sources show greater representation power, which supports downstream analyses at both individual and voxel levels.

To comprehensively assess the effectiveness of MSIVA, we compared three different initialization strategies across five candidate subspace structures. We first simulated multiple synthetic datasets to evaluate whether MSIVA can successfully reconstruct the ground-truth subspace structures. Next, we applied MSIVA to two large multimodal neuroimaging datasets, the UK Biobank dataset (Miller et al., 2016) and a schizophrenia (SZ) patient dataset combined from several studies (Aine et al., 2017; Keator et al., 2016; Tamminga et al., 2014). Using canonical correlation analysis (CCA) (Hotelling, 1992), we conducted a follow-up assessment of each MSIVA cross-modal subspace separately and identified projections within the optimal subspace structure yielding the post-CCA linked sources. We then performed age regression, sex classification, and diagnosis classification to investigate the associations between these linked sources and phenotype measures. Furthermore, we proposed a voxelwise brain-age delta analysis using reconstructed data from MSIVA.

Simulation results showed that MSIVA successfully recovered the ground-truth subspace structures including high-dimensional subspaces. In two independent neuroimaging datasets, the optimal latent structure identified by MSIVA consisted of five two-dimensional cross-modal subspaces, instead of the one-dimensional independent subspaces used in MMIVA. Brain-phenotype modeling revealed the post-CCA MSIVA sources were related to age, sex, and SZ-related effects, while the brain-age gap was explained by several phenotype measures, including lifestyle factors and cognitive test scores. Overall, our findings suggest that MSIVA can effectively reveal linked multidimensional latent sources related to phenotype variables from multimodal neuroimaging data, thereby uncovering linked phenotypic and neuropsychiatric biomarkers of brain structure and function.

---

[1]A preliminary version of this work was published at the 2023 IEEE 20th International Symposium on Biomedical Imaging (ISBI) (X. Li, Adali, et al., 2023). The present study extends the conference publication and provides a more comprehensive evaluation and analysis of the proposed method by (1) including the schizophrenia patient neuroimaging data analysis, (2) incorporating an identity matrix as an additional candidate subspace structure, (3) performing a detailed multimodal initialization comparison, (4) introducing the voxelwise brain-age delta analysis, (5) adding extensive visualizations and interpretations of the multimodal patterns and their associations, and (6) performing a quantitative comparison of loss values and other proposed metrics.

2   Methods

2.1   Multimodal subspace independent vector analysis

We consider the following problem that each observed data modality is a linear mixture of latent sources:

$$\mathbf{X}^{[m]} = \mathbf{A}^{[m]}\mathbf{S}^{[m]}, \tag{1}$$

where $\mathbf{X}^{[m]} \in \mathbb{R}^{V \times N}$ is the observed data, $\mathbf{A}^{[m]} \in \mathbb{R}^{V \times C}$ is a linear mixing matrix, $\mathbf{S}^{[m]} \in \mathbb{R}^{C \times N}$ is the latent source matrix, m is the modality index, V is the input feature dimensionality, C is the number of latent sources $(C < V)$, and N is the number of samples. Sources across M modalities are either statistically dependent or independent, according to the subspace structure S defined using available *a priori* information. We aim to recover the latent sources $\hat{\mathbf{S}}^{[m]} \in \mathbb{R}^{C \times N}$ by estimating a linear unmixing matrix $\mathbf{W}^{[m]} \in \mathbb{R}^{C \times V}$:

$$\hat{\mathbf{S}}^{[m]} = \mathbf{W}^{[m]}\mathbf{X}^{[m]}. \tag{2}$$

We refer to our proposed approach as Multimodal Subspace Independent Vector Analysis (MSIVA) because it extends MMIVA, linking equal size, *higher-dimensional* subspaces across modalities. We consider five candidate subspace structure priors that define different types of multimodal relationships (Figure 1) and three initialization workflows that capture different amounts of joint cross-modal information (Figure 2). Given a candidate subspace structure, MSIVA consists of iterative combinatorial optimization of the source estimates (cross-modal subspace alignment) and numerical optimization of the MISA loss (Equation 6). This process is repeated for each of the five candidate subspace structures, followed by a best-fit determination based on the final quantitative metrics of all candidates.

2.1.1   Designing subspace structure priors

Higher-dimensional subspaces are more likely to explain complex variability in the data, such as statistical dependence. In designing subspace structures, we consider two main objectives. First, unlike MMIVA, which assumes all sources are statistically independent, our approach aims to identify groups of linked (*not* independent) sources within each modality, while assuming independence between groups. These groups of linked sources are referred to as *subspaces*. Second, we aim to detect statistical dependence between subspaces across different modalities, which involves solving an expensive combinatorial optimization problem. To simplify this process, we restrict the search space by assuming that cross-modal dependence occurs only between higher-dimensional subspaces (two-dimensional or greater) of the same size across

Figure 1: **Five plausible candidate subspace structure priors ($S_1 - S_5$) for two modalities (M_1, M_2).** Each panel depicts the idealized association between sources from two modalities (M_1, M_2), across five different plausible scenarios ($S_1 - S_5$). The size of each block represents the number of sources within a subspace (the subspace size). The colorful subspaces highlighted in blue are linked between modalities, whereas the black subspaces highlighted in green (1×1 blocks in $S_1 - S_4$) are specific to each modality (no cross-modal correlation). For each modality, sources within the same subspace are statistically dependent while sources in different subspaces are statistically independent.

modalities. Additionally, we treat all modality-specific subspaces as one-dimensional ($1D$), meaning each represents a single unlinked unimodal source.

Building on the MISA framework (Silva et al., 2020), we require user-defined candidate subspace structure priors that specify the expected linkage patterns. The goal of MSIVA is to determine which one of the candidate subspace structures best fits the observed data. We consider two questions when designing the subspace structure. First, *how many sources should be included?* We recommend estimating the model order of the real data using information-theoretic criteria (Y.-O. Li et al., 2007) and selecting the number of sources that captures sufficient information on all modalities. Second, *how should the sources be grouped?* In the functional imaging literature, two- to four-dimensional clusters have been used to hierarchically group sources (Ke et al., 2019; Ma et al., 2010, 2011). Accordingly, we define subspace structures with two- to four-dimensional cross-modal subspaces. We consider subspace structures that include as many groups of uniform dimension as possible, as well as one subspace configuration that contains groups of different dimensions. Therefore, we propose five plausible subspace structures ($S_1 - S_5$) in two modalities (M_1, M_2), all with 12 sources in each modality (Figure 1):

- S_1: One two-dimensional ($2D$) cross-modal subspace, one three-dimensional ($3D$) cross-modal subspace, one four-dimensional ($4D$) cross-modal subspace, and three $1D$ unimodal subspaces.

- S_2: Five $2D$ cross-modal subspaces and two $1D$ unimodal subspaces.

- S_3: Three $3D$ cross-modal subspaces and three $1D$ unimodal subspaces.

- S_4: Two $4D$ cross-modal subspaces and four $1D$ unimodal subspaces.

- S_5: Twelve $1D$ cross-modal subspaces (no unimodal subspaces, identical to MMIVA).

2.1.2 MSIVA default initialization workflow

Our previous research demonstrates that proper initialization of unmixing weights can pre-align latent subspaces, leading to faster convergence to optimal solutions (X. Li, Khosravinezhad, et al., 2023; Silva et al., 2024). Based on these findings, we evaluated three initialization workflows incorporating different amounts of shared cross-modal information.

MSIVA default initialization workflow first utilizes multimodal group principal component analysis (MG-PCA) to identify common principal components across all modalities and then applies unimodal ICA on the MGPCA-reduced data of each modality. Unlike principal component analysis (PCA) that identifies orthogonal directions of maximal variation for each modality separately, MGPCA identifies directions of maximal *common* variation across all modalities while ensuring equal variance contribution of each modality. Eigenvectors were computed based on the weighted average of the covariance matrices:

$$\boldsymbol{\Sigma}_{\text{avg}} = \frac{1}{M}\sum_{m=1}^{M} N\frac{\boldsymbol{\Sigma}^{[m]}}{\text{trace}\left(\boldsymbol{\Sigma}^{[m]}\right)} = \frac{1}{M}\sum_{m=1}^{M} N\frac{\mathbf{X}^{[m]^\top}\mathbf{X}^{[m]}}{||\mathbf{X}^{[m]}||_{\text{Fr}}^2}, \tag{3}$$

where $\boldsymbol{\Sigma}^{[m]} = \frac{\mathbf{X}^{[m]^\top}\mathbf{X}^{[m]}}{V-1} \approx \mathbb{E}\left[\mathbf{X}^{[m]^\top}\mathbf{X}^{[m]}\right]$, $\mathbb{E}[\cdot]$ is the expectation operator, and $||\cdot||_{\text{Fr}}$ indicates the Frobenius norm. The scaling factor $\frac{\text{trace}\left(\boldsymbol{\Sigma}^{[m]}\right)}{N}$ is the ratio of the variance in the modality to the number of samples. We define the whitening matrix $\mathbf{W}_{\text{MGPCA}}^{[m]}$ as follows:

$$\mathbf{W}_{\text{MGPCA}}^{[m]} = \sqrt{N-1}\boldsymbol{\Lambda}^{-\frac{1}{2}}\mathbf{U}^{[m]^\top}\lambda^{[m]}, \tag{4}$$

where $\boldsymbol{\Lambda}$ and $\mathbf{Q}$ are the top C eigenvalues and eigenvectors of $\boldsymbol{\Sigma}_{\text{avg}}$, respectively, $\mathbf{U}^{[m]} = \left(\lambda^{[m]}\mathbf{X}^{[m]}\right)\mathbf{Q}\boldsymbol{\Lambda}^{-\frac{1}{2}}$, $\lambda^{[m]} = \sqrt{\frac{N}{M(V-1)\text{trace}(\boldsymbol{\Sigma}^{[m]})}} = \sqrt{\frac{N}{M||\mathbf{X}^{[m]}||_{\text{Fr}}^2}}$.

Next, the MGPCA-reduced data from each modality $\mathbf{X}_{\text{r}}^{[m]} = \mathbf{W}_{\text{MGPCA}}^{[m]}\mathbf{X}^{[m]}$ underwent a separate ICA estimation using the Infomax algorithm (Bell & Sejnowski, 1995) initialized with an identity matrix to obtain C independent sources per modality $\hat{\mathbf{S}}_{\text{Infomax}}^{[m]} = \mathbf{W}_{\text{Infomax}}^{[m]}\mathbf{X}_{\text{r}}^{[m]}$. These estimates were further optimized by running MISA as a unimodal ICA model initialized with $\mathbf{W}_{\text{Infomax}}^{[m]}$, leading to the final unimodal ICA source estimates $\hat{\mathbf{S}}_{\text{ICA}}^{[m]} = \mathbf{W}_{\text{ICA}}^{[m]}\mathbf{X}_{\text{r}}^{[m]}$. Finally, MSIVA was initialized by the combined MGPCA+ICA estimates $\mathbf{W}_0^{[m]} = \mathbf{W}_{\text{ICA}}^{[m]}\mathbf{W}_{\text{MGPCA}}^{[m]}$ from both modalities. Subsequently, we compared MSIVA with a fully unimodal initialization workflow and a fully multimodal initialization workflow to comprehensively evaluate performance.

Figure 2: **Overview of three proposed initialization workflows.** From left to right, the unimodal, MSIVA default, and multimodal initialization approaches are: separate PCAs followed by separate ICAs (PCA + ICA); multimodal group PCA with separate ICAs per modality (MGPCA + ICA); multimodal group PCA with group ICA (MGPCA + GICA). MGPCA + ICA was used as the MSIVA default initialization workflow. After initialization, the greedy combinatorial optimization and the numerical optimization with the MISA loss were performed for sufficient iterations until the loss value converged.

2.1.3   Unimodal initialization workflow

The unimodal initialization workflow simply applied PCA and ICA on each modality separately. We first projected the imaging data matrix from each modality $\mathbf{X}^{[m]}$ into a reduced data matrix $\mathbf{X}_{\mathrm{r}}^{[m]}$ with C principal components and obtained the corresponding whitening matrix $\mathbf{W}_{\mathrm{PCA}}^{[m]}$. Next, we applied ICA on each reduced data matrix $\mathbf{X}_{\mathrm{r}}^{[m]}$ to obtain C independent sources and the corresponding unmixing matrix $\mathbf{W}_{\mathrm{ICA}}^{[m]}$. In this workflow, the MISA initialization matrix was defined as $\mathbf{W}_{0}^{[m]} = \mathbf{W}_{\mathrm{ICA}}^{[m]} \mathbf{W}_{\mathrm{PCA}}^{[m]}$.

2.1.4   Multimodal initialization workflow

The multimodal initialization workflow sequentially applied MGPCA (same as in Equation 4) and group ICA (GICA) across all data modalities, resulting in the weight matrices $\mathbf{W}_{\mathrm{MGPCA}}^{[m]}$ and $\mathbf{W}_{\mathrm{GICA}}^{[m]}$. GICA performed ICA on the combined MGPCA-reduced data from all M modalities, that is, $\mathbf{X}_{\mathrm{r}} = \sum_{m=1}^{M} \mathbf{X}_{\mathrm{r}}^{[m]}$. In this workflow, MISA was initialized by $\mathbf{W}_{0}^{[m]} = \mathbf{W}_{\mathrm{GICA}}^{[m]} \mathbf{W}_{\mathrm{MGPCA}}^{[m]}$.

2.1.5   Alternating combinatorial and numerical optimization

All three workflows utilize MISA's greedy combinatorial optimization and objective function to estimate latent sources. MISA uses the relative gradients and the L-BFGS algorithm (Liu & Nocedal, 1989) in a barrier-type numerical optimizer (`fmincon` from MATLAB Optimization Toolbox)[2] . The MISA loss function $\mathcal{L}_{\mathrm{MISA}}(\cdot)$ (Silva et al., 2020) is defined as the Kullback-Leibler (KL) divergence between the joint distribution of all recovered sources $p\left(\hat{\mathbf{S}}\right)$ and the product of all K subspace distributions $q\left(\hat{\mathbf{S}}\right) = \prod_{k=1}^{K} p\left(\hat{\mathbf{S}}_k\right)$, which is equivalent to mutual information among K subspaces. Thus, subspaces are assumed to be statistically independent of each other within each modality. Also, sources within a subspace are considered to be dependent (or linked) to one another. The distribution of each subspace is modeled as the joint multivariate Kotz distribution (Kotz, 1975; Nadarajah, 2003) of sources within that subspace as follows:

$$p\left(\hat{\mathbf{S}}_k\right) = \frac{\beta \lambda^{\nu} \Gamma\left(\frac{d_k}{2}\right) \left(\hat{\mathbf{S}}_k^{\top} \mathbf{D}_k^{-1} \hat{\mathbf{S}}_k\right)^{\eta-1}}{\pi^{\frac{d_k}{2}} \left(\det\left(\mathbf{D}_k\right)\right)^{\frac{1}{2}} \Gamma\left(\nu\right)} e^{-\lambda \left(\hat{\mathbf{S}}_k^{\top} \mathbf{D}_k^{-1} \hat{\mathbf{S}}_k\right)^{\beta}}, \tag{5}$$

where d_k is the dimensionality of the k-th subspace ($d_k = C_k \cdot M$ if the subspace is cross-modal—C_k being the subspace size in each modality—and $d_k = 1$ if the subspace is unimodal). $\beta > 0$, $\lambda > 0$,

---

[2]Stochastic gradient descent is a valid alternative approach. See Silva et al., 2020 for the derivation of the gradients.

and $\eta > \frac{2-d_k}{2}$ are the Kotz parameters that control the shape of the distribution, the kurtosis, and the hole size, respectively. For brevity, we define $\nu \triangleq \frac{2\eta+d_k-2}{2\beta} > 0$ and $\alpha \triangleq \frac{\Gamma(\nu+\beta^{-1})}{\lambda^{\beta^{-1}}d_k\Gamma(\nu)}$. $\Gamma(\cdot)$ denotes the gamma function. $\det(\cdot)$ denotes the determinant. The positive definite dispersion matrix $\mathbf{D}_k$ is given by $\mathbf{D}_k = \alpha^{-1}\mathbf{\Sigma}_k$, where $\mathbf{\Sigma}_k$ is the source covariance matrix in the k-th subspace. In MSIVA, $\mathbf{\Sigma}_k$ is iteratively reset as the current estimate of source *correlation* matrix for subspace k, while the Kotz parameters are set to $\beta = 0.5$, $\lambda = 0.8966$, $\eta = 1$, producing a distribution slightly less peaked than a multivariate Laplace distribution with source variance $\alpha = \frac{\pi^2}{3}$, corresponding to the variance of the successful Logistic distribution used in Infomax.

We want to minimize the loss function $\mathcal{L}_{\mathsf{MISA}}(\cdot)$ by solving the following optimization problem:

$$
\begin{aligned}
\min \mathcal{L}_{\mathsf{MISA}}\left(\hat{\mathbf{S}}\right) &= \min D_{\mathsf{KL}}\left(p\left(\hat{\mathbf{S}}\right) \middle\| q\left(\hat{\mathbf{S}}\right)\right) \\
&= \min D_{\mathsf{KL}}\left(p\left(\hat{\mathbf{S}}\right) \middle\| \prod_{k=1}^{K} p\left(\hat{\mathbf{S}}_k\right)\right) \\
&= \min \mathbb{E}\left[\ln p\left(\hat{\mathbf{S}}\right)\right] - \sum_{k=1}^{K}\mathbb{E}\left[\ln p\left(\hat{\mathbf{S}}_k\right)\right] \\
&= \min_{\substack{\hat{\mathbf{W}},\mathbf{P}_k, \\ k=1,\ldots,K}} \mathbb{E}\left[\ln p\left(\hat{\mathbf{W}}\mathbf{X}\right)\right] - \sum_{k=1}^{K}\mathbb{E}\left[\ln p\left(\mathbf{P}_k\hat{\mathbf{W}}\mathbf{X}\right)\right],
\end{aligned}
\tag{6}
$$

where $\hat{\mathbf{S}} = \left[\hat{\mathbf{S}}^{[1]}; \ldots; \hat{\mathbf{S}}^{[M]}\right] \in \mathbb{R}^{MC \times N}$ represents all estimated sources from all M modalities[3]. $\mathbf{X} = \left[\mathbf{X}^{[1]}; \ldots; \mathbf{X}^{[M]}\right] \in \mathbb{R}^{MV \times N}$ represents the concatenated data from all M modalities. $\hat{\mathbf{W}} \in \mathbb{R}^{MC \times MV}$ is the estimated block-diagonal unmixing matrix, such that $\hat{\mathbf{S}}^{[m]} = \hat{\mathbf{W}}^{[m]}\mathbf{X}^{[m]}$. $\mathbf{P}_k \in \mathbb{R}^{C_k \times MC}$ is the k-th subspace assignment matrix defined by the subspace structure S in Section 2.1.1, and C_k is the number of sources assigned to the k-th subspace.

We alternated between greedy combinatorial optimization and MISA numerical optimization until the loss value converged. Combinatorial and numerical optimizations were performed 10 times for synthetic data and 20 times for neuroimaging data. The goal of combinatorial optimization is to escape potential local minima found by the numerical optimizer. During each combinatorial optimization, the rows of the modality-specific unmixing matrices $\hat{\mathbf{W}}^{[m]}$ were permuted. The numerical optimizer then resumed for an additional 150 iterations, returning to combinatorial optimization afterwards. Finally, we selected the $\hat{\mathbf{W}}^{[m]}$ with the lowest loss value across all iterations performed.

---

[3]We use semicolon (;) to denote that matrices are stacked vertically and comma (,) to denote that matrices are stacked horizontally.

2.2 Datasets

2.2.1 Synthetic data

For each subspace structure S, we generated a synthetic dataset with two modalities $\mathbf{X} = \left[\mathbf{X}^{[1]}; \mathbf{X}^{[2]}\right] \in \mathbb{R}^{2V \times N}$, where V is the size of input features ($V = 20000$) and N is the number of samples ($N = 3000$). V is a sufficiently large number for covariance estimation in PCA or MGPCA. N is chosen to approximate the number of samples in the UK Biobank neuroimaging dataset in this study (see Section 2.2.2). Each data modality is a linear mixture of $C = 12$ sources spanning the subspaces defined in S, $\mathbf{X}^{[m]} = \mathbf{A}^{[m]}\mathbf{S}^{[m]}$, $\mathbf{A}^{[m]} \in \mathbb{R}^{V \times C}$, $\mathbf{S}^{[m]} \in \mathbb{R}^{C \times N}$, $m \in \{1, 2\}$. Each subspace $\mathbf{S}_k$ is independently sampled from a multivariate Laplace distribution and hence, the marginal distributions correspond to the individual sources. Cross-modal sources within each linked subspace are dependent with correlation coefficients uniformly sampled from 0.65 to 0.85. Unimodal sources ($1D$ subspaces in $S_1 - S_4$) also have Laplace distribution but independent from all other sources and subspaces. Thus, they do not correlate with any other sources.

2.2.2 Neuroimaging data

We utilized two large multimodal neuroimaging datasets including two imaging modalities: T1-weighted structural MRI (sMRI) and resting-state functional MRI (fMRI). The first dataset is from the UK Biobank study (Miller et al., 2016). 2907 subjects from two sites (age mean $\pm$ standard deviation: 62.09 ± 7.32 years; age median: 63 years; age range: $46 - 79$ years; 1452 males, 1455 females) were used for formal analysis after excluding subjects with more than 4% missing phenotype measures (Smith et al., 2015). The second dataset includes 999 patients and controls (age mean $\pm$ standard deviation: 38.61 ± 13.13 years; age median: 39 years; age range: $15 - 65$ years; 625 males, 374 females; 538 controls, 337 patients diagnosed with schizophrenia, 63 patients with bipolar disorder, 11 patients with schizoaffective disorder, 28 schizoaffective bipolar-type probands, and 22 schizoaffective depression-type probands) combined across several studies, including Bipolar and Schizophrenia Network for Intermediate Phenotypes (BSNIP) (Tamminga et al., 2014), Center for Biomedical Research Excellence (COBRE) (Aine et al., 2017), Function Biomedical Informatics Research Network (FBIRN) (Keator et al., 2016), and Maryland Psychiatric Research Center (MPRC).

For each dataset, we preprocessed sMRI and fMRI to obtain the gray matter (GM) and mean-scaled amplitude of low frequency fluctuations (mALFF) feature maps, respectively. We resampled each GM or mALFF feature map to $3 \times 3 \times 3\text{mm}^3$ resolution and applied a group-level GM mask on the feature

map, resulting in 44318 voxels. Data acquisition and preprocessing details are described in Appendix A.

Next, for each data modality in each dataset, we performed variance normalization (removed mean and divided by standard deviation) for each subject, and then removed the mean across all subjects for each voxel. Lastly, we regressed out site effects for each dataset as follows:

$$\mathbf{X}^{[m]} \longleftarrow \mathbf{X}^{[m]} - \mathbf{X}^{[m]}\mathbf{L}\left(\mathbf{L}^{\top}\mathbf{L}\right)^{-1}\mathbf{L}^{\top}, \tag{7}$$

where $\mathbf{L} = [\mathbf{1}, \boldsymbol{\ell}]$, with $\mathbf{1} \in \mathbb{R}^{N}$ being a column vector of ones and $\boldsymbol{\ell}$ being one-hot encoded site labels.

2.3   Quantitative evaluation metrics

2.3.1   Normalized multidataset Moreau-Amari intersymbol interference

The normalized multidataset Moreau-Amari intersymbol interference (MISI) (Amari et al., 1996; Macchi & Moreau, 1995; Silva et al., 2020) was used to evaluate the residual interference between the estimated unmixing matrix $\hat{\mathbf{W}}$ and the ground-truth mixing matrix $\mathbf{A}$:

$$\text{MISI}(\mathbf{H}) = \frac{1}{2K(K-1)}\left[\sum_{i=1}^{K}\left(-1 + \sum_{j=1}^{K}\frac{|h_{ij}|}{\max\limits_{k}|h_{ik}|}\right) + \sum_{j=1}^{K}\left(-1 + \sum_{i=1}^{K}\frac{|h_{ij}|}{\max\limits_{k}|h_{kj}|}\right)\right], \tag{8}$$

where $h_{ij} = \mathbf{1}^{\top}\left|\mathbf{P}_i\hat{\mathbf{W}}\mathbf{A}\mathbf{P}_j\right|\mathbf{1}$ is the sum of absolute values from all elements corresponding to subspaces i and j in the interference matrix $\mathbf{H} = \hat{\mathbf{W}}\mathbf{A}$.

2.3.2   MISA loss

We also reported the corresponding MISA loss value defined in Equation 6. When evaluating performance on synthetic data, we prioritized the MISI metric and interference matrix $\mathbf{H}$ as they leverage the ground-truth information. We also examined whether the loss values are consistent with the MISI values.

2.3.3   Mean correlation coefficient and minimum distance

We used two metrics to summarize cross-modal source correlations: the mean correlation coefficient (MCC) and the minimum distance (MD). Both metrics were computed in a two-stage manner. First, we constructed an aggregated correlation matrix across cross-modal subspaces. Then, we computed the

MCC or MD on the aggregated matrix. This approach ensures balanced contributions from subspaces with varying dimensionalities. We use the term *multimodal* MCC/MD to refer to correlations computed between recovered sources and ground-truth sources within each modality, after min–max subspace aggregation. We use *cross-modal* MCC/MD to denote correlations computed between recovered sources across two different modalities, without reference to the ground truth.

The proposed metrics included three distinct correlation matrices: (1) absolute correlations between recovered sources and ground-truth sources of the first modality $\mathbf{R}^{[1]} \in \mathbb{R}^{C \times C}$; (2) absolute correlations between recovered sources and ground-truth sources of the second modality $\mathbf{R}^{[2]} \in \mathbb{R}^{C \times C}$; (3) absolute cross-modal correlations between recovered sources across the two modalities $\mathbf{R}^{\mathrm{cm}} \in \mathbb{R}^{C \times C}$. The correlations of the first modality $\mathbf{R}^{[1]}$ were sorted into a block-diagonal structure consistent with the predefined subspace structure, and the same permutation was then applied to the correlations of the second modality $\mathbf{R}^{[2]}$. The cross-modal correlations $\mathbf{R}^{\mathrm{cm}}$ did not involve the ground-truth, and they were left unsorted to preserve their original linkage. In the following equations, $\delta(\texttt{condition}) = 1$ when the specified condition is satisfied.

Multimodal mean correlation coefficient.   Given a modality-specific correlation matrix $\mathbf{R}^{[m]}$, we first calculated the aggregated modality-specific correlation matrix $\bar{\mathbf{R}}^{[m]} \in \mathbb{R}^{K \times K}$ by aggregating cross-modal subspaces:

$$\bar{\mathbf{R}}^{[m]}_{kk'} = \frac{1}{C_k + C_{k'}} \left(\mathbf{1}^{\top} \cdot \mathrm{rowmax}\left(\mathbf{R}^{[m]}_{[kk']} \right) + \mathrm{columnmax}\left(\mathbf{R}^{[m]}_{[kk']} \right) \cdot \mathbf{1} \right), \tag{9}$$

where K is the number of cross-modal subspaces in a given subspace structure, and the modality index $m \in \{1, 2\}$. $\mathbf{R}^{[m]}_{[kk']} \in \mathbb{R}^{C_k \times C_{k'}}$ is the correlation block between the k-th subspace and the k'-th subspace in $\mathbf{R}^{[m]}$. $\mathbf{1}$ denotes a column vector of all ones. $\mathrm{rowmax}\left(\mathbf{R} \right)$ denotes a column vector containing the maximum value of each row in the matrix $\mathbf{R}$, and $\mathrm{columnmax}\left(\mathbf{R} \right)$ denotes a row vector containing the maximum value of each column in $\mathbf{R}$. C_k is the dimension of the k-th cross-modal subspace.

We then obtained the multimodal correlation matrix $\bar{\mathbf{R}}^{\mathrm{mm}}$ by taking the element-wise minimum for the diagonal elements and the element-wise maximum for the off-diagonal elements between the two aggregated unimodal correlation matrices ($\bar{\mathbf{R}}^{[1]}$ and $\bar{\mathbf{R}}^{[2]}$):

$$\bar{\mathbf{R}}^{\mathrm{mm}}_{kk'} = \max\left(\bar{\mathbf{R}}^{[1]}_{kk'}, \bar{\mathbf{R}}^{[2]}_{kk'} \right) \cdot \delta\left(k \neq k' \right) + \min\left(\bar{\mathbf{R}}^{[1]}_{kk'}, \bar{\mathbf{R}}^{[2]}_{kk'} \right) \cdot \delta\left(k = k' \right). \tag{10}$$

We then computed the multimodal mean correlation coefficient (MMCC) by aggregating across the

diagonal of $\bar{\mathbf{R}}^{\mathsf{mm}}$:

$$\text{MMCC}\left(\bar{\mathbf{R}}^{\mathsf{mm}}\right) = \frac{1}{K}\sum_{k=1}^{K}\bar{\mathbf{R}}_{kk}^{\mathsf{mm}}. \tag{11}$$

Cross-modal mean correlation coefficient.   Similarly, we first calculated the aggregated cross-modal correlation matrix $\bar{\mathbf{R}}^{\mathsf{cm}}$:

$$\bar{\mathbf{R}}_{kk'}^{\mathsf{cm}} = \frac{1}{C_k + C_{k'}}\left(\mathbf{1}^{\top}\cdot\text{rowmax}\left(\mathbf{R}_{[kk']}^{\mathsf{cm}}\right) + \text{columnmax}\left(\mathbf{R}_{[kk']}^{\mathsf{cm}}\right)\cdot\mathbf{1}\right). \tag{12}$$

We then computed the cross-modal mean correlation coefficient (CMCC) by aggregating across the diagonal of $\bar{\mathbf{R}}^{\mathsf{cm}}$:

$$\text{CMCC}\left(\bar{\mathbf{R}}^{\mathsf{cm}}\right) = \frac{1}{K}\sum_{k=1}^{K}\bar{\mathbf{R}}_{kk}^{\mathsf{cm}}. \tag{13}$$

Multimodal and cross-modal minimum distance.   We computed the multimodal minimum distance (MMD) and the cross-modal minimum distance (CMD) by aggregating *all* (diagonal and off-diagonal) elements of the matrices $\bar{\mathbf{R}}^{\mathsf{mm}}$ and $\bar{\mathbf{R}}^{\mathsf{cm}}$, respectively. The MMD and CMD account for not only block-diagonal alignment, but also off-diagonal separation.

$$\text{MMD}\left(\bar{\mathbf{R}}^{\mathsf{mm}}\right) = \frac{1}{K^2}\sum_{k=1}^{K}\sum_{k'=1}^{K}\left(\bar{\mathbf{R}}_{kk'}^{\mathsf{mm}}\cdot\delta\left(k\neq k'\right) + \left(1 - \bar{\mathbf{R}}_{kk'}^{\mathsf{mm}}\right)\cdot\delta\left(k = k'\right)\right), \tag{14}$$

$$\text{CMD}\left(\bar{\mathbf{R}}^{\mathsf{cm}}\right) = \frac{1}{K^2}\sum_{k=1}^{K}\sum_{k'=1}^{K}\left(\bar{\mathbf{R}}_{kk'}^{\mathsf{cm}}\cdot\delta\left(k\neq k'\right) + \left(1 - \bar{\mathbf{R}}_{kk'}^{\mathsf{cm}}\right)\cdot\delta\left(k = k'\right)\right). \tag{15}$$

2.4   Experiments

2.4.1   Synthetic data experiment

We first verified whether the proposed approaches including MSIVA can identify and distinguish the correct subspace structure (the one used to generate the data) from the incorrect ones in synthetic data. For each of the five subspace structures ($S_1 - S_5$) described in Section 2.1.1, we generated a synthetic dataset where the data distribution is defined by the corresponding subspace structure. Next, we conducted experiments on all combinations of five subspace structures (Figure 1) and three

initialization workflows (Figure 2). Finally, we visualized the interference matrices $\mathbf{H}^{[m]} = \hat{\mathbf{W}}^{[m]}\mathbf{A}^{[m]}$[4] to confirm if the subspace structures were recovered. In addition, we reported MISI and final MISA loss values.

2.4.2 Neuroimaging data experiment

We performed experiments on each of two multimodal neuroimaging datasets separately, using each of the same five candidate subspace structures $S_1 - S_5$, and identified the optimal subspace structure as the one yielding the lowest final MISA loss value. Note that the MISI is not available because the ground-truth subspace structure is unknown in the real data. Instead, we reported the cross-modal metrics (CMCC and CMD) that measure cross-modal subspace alignment. We computed cross-modal source correlations for each subspace structure using both the *linear* Pearson correlation coefficient and the *nonlinear* randomized dependence coefficient (RDC) (Lopez-Paz et al., 2013).

To further assess the cross-modal linkage strength of the estimated subspaces *within* the optimal subspace structure, separate post-hoc CCA of each cross-modal subspace was used to recover projections with the maximum correlation between the two modalities:

$$(\hat{\mathbf{p}}_k, \hat{\mathbf{q}}_k) = \underset{\mathbf{p}_k, \mathbf{q}_k}{\arg\max} \; \text{corr}\left(\mathbf{p}_k^\top \hat{\mathbf{S}}_k^{[1]}, \mathbf{q}_k^\top \hat{\mathbf{S}}_k^{[2]}\right), \tag{16}$$

where $\mathbf{p}_k \in \mathbb{R}^{C_k}$ and $\mathbf{q}_k \in \mathbb{R}^{C_k}$ are the CCA projection vectors for the kth cross-modal subspace, and $\hat{\mathbf{S}}_k^{[1]} \in \mathbb{R}^{C_k \times N}$ and $\hat{\mathbf{S}}_k^{[2]} \in \mathbb{R}^{C_k \times N}$ are the recovered sources in the kth cross-modal subspace for two modalities. After estimation, post-CCA sources in the kth cross-modal subspace are obtained as $\hat{\mathbf{p}}_k^\top \hat{\mathbf{S}}_k^{[1]}$ and $\hat{\mathbf{q}}_k^\top \hat{\mathbf{S}}_k^{[2]}$. This assessment is sensible because linear transformations of individual sources within the same subspace are considered equivalently optimal[5] (Cardoso, 1998; Szabó et al., 2012).

2.5 Brain-phenotype prediction

To evaluate the association between phenotype measures and cross-modal post-CCA sources, we performed age prediction and sex classification tasks for the UKB dataset, as well as age prediction and binary diagnosis classification tasks (controls versus patients with SZ) for the patient dataset. Specifically, we trained a ridge regression model to predict age and a support vector machine with a linear

---

[4]The absolute values of $\mathbf{H}^{[m]} = \hat{\mathbf{W}}^{[m]}\mathbf{A}^{[m]}$ entries are reported since, in general, signs are not uniquely identifiable.

[5]While a subspace is uniquely identifiable, the individual sources within each subspace are not, warranting arbitrary transformation within the subspace.

kernel to classify sex groups or diagnosis groups. For the UKB dataset, 2907 subjects were stratified into a training set of 2000 subjects and a holdout test set of 907 subjects. For the patient dataset, 999 subjects were stratified into a training set of 699 subjects and a holdout test set of 300 subjects in the age prediction task; 538 controls and 337 patients with SZ were grouped into a training set of 612 subjects and a test set of 263 subjects in the diagnosis classification task. We performed 10-fold cross-validation to choose the best hyperparameter (regularization parameter set: $\{0.1, 0.2, \ldots, 1\}$) on the training set, then trained the model using all training subjects and evaluated it on the holdout test set. Age regression performance was measured by mean absolute error (MAE) between predicted age and chronological age. Sex or diagnosis classification performance was assessed via *balanced* accuracy: $0.5 \times$(true positive rate $+$ true negative rate).

2.6   Brain-age delta analysis on UK Biobank data

A key benefit of MSIVA is that the estimated multimodal sources are more expressive by leveraging higher-dimensional ($\geq 2D$) cross-modal subspaces. To demonstrate the utility of higher-dimensional subspaces, we proposed to conduct a two-stage *voxelwise* brain-age delta analysis using the UKB estimated sources from the optimal subspace structure. For each voxel in the reconstructed subspace ($\hat{\mathbf{X}}_k^{[m]} = \hat{\mathbf{A}}_k^{[m]} \hat{\mathbf{S}}_k^{[m]}$[6]), we estimated an initial age delta at the first stage and corrected it for age dependence and other confound variables at the second stage (Smith et al., 2019, 2020):

$$\boldsymbol{\delta}_1 = \hat{\mathbf{X}}_i \boldsymbol{\beta}_1 - \mathbf{y}, \tag{17}$$

$$\boldsymbol{\delta}_2 = \boldsymbol{\delta}_1 - \mathbf{Y} \boldsymbol{\beta}_2, \tag{18}$$

where $\hat{\mathbf{X}}_i$ indicates the i-th voxel's reconstructed patterns from each subspace. Namely, they include shared patterns from each cross-modal subspace[7], reconstructed sMRI patterns from each cross-modal subspace, and reconstructed data from each unimodal subspace (see Appendix B for more details). $\mathbf{y} \in \mathbb{R}^N$ is the demeaned chronological age. $\mathbf{Y} \in \mathbb{R}^{N \times 10}$ includes the confound variables: the demeaned linear, quadratic, cubic age terms, sex, the interaction between sex and each of the three age terms, the framewise displacement variable, and the spatial normalization variables from sMRI and fMRI. An advantage of the procedure described in Smith et al., 2019, 2020 is that it yields a breakdown of $\boldsymbol{\delta}_2$ per predictor in $\hat{\mathbf{X}}_i$. Lastly, we partialized $\boldsymbol{\delta}_2$ to remove residual associations between each predictor and the

---

[6]The modality-specific mixing matrix was estimated as the least-squares solution: $\hat{\mathbf{A}}^{[m]} = \mathbf{X}^{[m]} \hat{\mathbf{S}}^{[m]\top} \left(\hat{\mathbf{S}}^{[m]} \hat{\mathbf{S}}^{[m]\top} \right)^{-1}$.

[7]For each voxel i and subspace k, we utilized singular value decomposition (SVD) across all modalities $(\hat{\mathbf{X}}_{k[i,:]}^{[1]}, \hat{\mathbf{X}}_{k[i,:]}^{[2]})$ to capture the shared *reconstructed* multimodal information of each cross-modal subspace.

Figure 3: **Performance on synthetic data: MISI (lower is better).** Each row represents the ground-truth subspace structure used to generate the data and each column represents the test subspace structure used to fit the model. If a workflow could correctly identify all ground-truth subspace structures, the lowest MISI values would occur along the main diagonal. MISI ≤ 0.1 is considered a heuristic threshold for good source separation (Silva et al., 2014). The unimodal initialization workflow (PCA+ICA) and the MSIVA default initialization workflow (MGPCA+ICA) led to the lowest MISI values (≤ 0.02) along the main diagonal, indicating that these two initialization approaches were adequate and enabled correct identification of the ground-truth subspace structures *and* differentiation from the incorrect ones. However, the multimodal initialization workflow (MGPCA+GICA) failed to detect subspace structures S_1, S_3, and S_4 with high MISI values in the main diagonal. Thus, the MSIVA default initialization and the unimodal initialization are considered better than the multimodal initialization.

other predictors, obtaining the partialized brain-age delta, $\boldsymbol{\delta}_{2p}$.

We then correlated the voxelwise brain-age delta $\boldsymbol{\delta}_{2p}$ with 25 non-imaging phenotype variables such as lifestyle factors and cognitive test scores (see Appendix C for the full list of phenotype variables) to investigate multimodal brain-phenotype relationships. This voxelwise brain-age delta analysis allows us to visualize a voxel-level spatial map showing how each phenotype variable relates to the difference between chronological and estimated brain age.

3   Results

3.1   MSIVA identifies ground-truth subspace structures in synthetic data

We verified whether the proposed approaches, including three different initialization workflows, could identify the correct subspace structures used for data generation in synthetic datasets. As discussed in Appendix D, the MSIVA default initialization method (MGPCA+ICA) achieved a well-balanced combination of unimodal separation and cross-modal alignment in most cases. Nonetheless, after iterative combinatorial and numerical optimization, both the unimodal initialization workflow (PCA+ICA) and

Table 1: **Synthetic data: Final MISA loss values (lower is better).** Each row represents the ground-truth (GT) subspace structure used to generate the data and each column represents the test subspace structure used to fit the model. The lowest loss value along the *row* is highlighted in bold, which determines the selected subspace. Approaches performing consistently well in relation to the MISI in Figure 3 would contain bold loss values *only* along the diagonal. The loss values were consistent with the MISI values. The unimodal initialization and the MSIVA default initialization correctly identified the ground-truth subspace structures. However, the multimodal initialization loss values incorrectly implied that S_3, S_1, and S_2 were better when S_1, S_3, and S_4 were the ground-truth subspace structures, respectively.

Unimodal initialization	S_1^{Test}	S_2^{Test}	S_3^{Test}	S_4^{Test}	S_5^{Test}
S_1^{GT}	**42.692**	42.879	42.766	43.054	43.231
S_2^{GT}	42.903	**42.301**	42.843	42.823	42.918
S_3^{GT}	42.710	42.864	**42.635**	43.073	43.256
S_4^{GT}	43.101	43.238	43.195	**42.975**	43.507
S_5^{GT}	43.391	42.997	43.492	43.767	**42.021**
MSIVA default initialization	S_1^{Test}	S_2^{Test}	S_3^{Test}	S_4^{Test}	S_5^{Test}
S_1^{GT}	**42.676**	42.880	42.750	43.002	43.110
S_2^{GT}	42.614	**42.288**	42.739	42.876	42.745
S_3^{GT}	42.694	42.848	**42.620**	42.868	43.070
S_4^{GT}	33.205	33.275	33.272	**33.115**	33.675
S_5^{GT}	43.398	42.946	43.383	43.969	**42.005**
Multimodal initialization	S_1^{Test}	S_2^{Test}	S_3^{Test}	S_4^{Test}	S_5^{Test}
S_1^{GT}	23.897	23.957	**23.838**	24.179	24.199
S_2^{GT}	27.786	**27.442**	27.810	28.032	28.158
S_3^{GT}	**23.864**	24.012	23.964	24.234	24.346
S_4^{GT}	17.582	**17.328**	17.85	18.404	20.275
S_5^{GT}	36.758	36.316	36.752	37.147	**35.262**

the MSIVA default initialization workflow led to the lowest MISI values (≤ 0.02) along the main diagonal in Figure 3, demonstrating that both approaches correctly recovered the ground truth when the correct subspace structure was provided. The multimodal initialization workflow (MGPCA+GICA), on the other hand, showed suboptimal performance with elevated MISI values along the main diagonal and was thus excluded from subsequent neuroimaging data experiments.

We confirmed that loss values successfully converged within each numerical optimization run (Appendix E Figure E.1). According to Table 1, the final loss values were consistent with the MISI results. In addition, we thoroughly examined the relationship between loss values and MISI values across all numerical optimization runs. Loss values were strongly correlated with MISI values (unimodal initialization: $R^2 \geq 0.95$; MSIVA default initialization: $R^2 \geq 0.58$) when the ground-truth and test subspace structures were

Figure 4: **Synthetic data: Interference matrices $\mathbf{H}^{[m]} = \hat{\mathbf{W}}^{[m]}\mathbf{A}^{[m]}$ corresponding to the diagonal MISI values in Figure 3.** Cross-modal subspaces are highlighted in blue while modality-specific subspaces are highlighted in green. The subspace permutation, applied for ease of interpretation, is identical for both modalities. The correct subspace structures were identified and aligned across both modalities by three workflows (rows II-VII), in accordance with the ground-truth simulation design (row I), except that the multimodal initialization failed to recover S_1, S_3, and S_4 (rows VI-VII).

matched, indicating that loss serves as a valid proxy for MISI under consistent initialization (Appendix E Figure E.2). The loss values obtained with the multimodal initialization workflow failed to detect the ground-truth subspace structures containing $3D$ or $4D$ subspace(s) (S_1, S_3, and S_4).

As presented in Figure 4, the recovered subspace structures from the MSIVA default initialization workflow (rows IV-V) and the unimodal initialization workflow (rows II-III) under the correct subspace structure aligned well with the proposed ground truth (row I), confirming the effectiveness of MSIVA. However, the multimodal initialization workflow (rows VI-VII) could not recover the ground-truth subspace structures for S_1, S_3, and S_4 even when given the correct subspace structure, suggesting that cross-modal alignment optimization becomes more challenging in the presence of high-dimensional subspaces.

3.2   MSIVA detects the latent subspace structure in neuroimaging data

The model order was estimated for each neuroimaging dataset using three information-theoretic criteria (Y.-O. Li et al., 2007). Twelve sources captured the majority of information in the patient dataset and a substantial amount in the UK Biobank dataset (Appendix F). Hence, twelve sources were included in each subspace structure.

We next applied the MSIVA default initialization workflow and the unimodal initialization workflow on two large multimodal neuroimaging datasets separately—the UK Biobank (UKB) dataset and the combined patient dataset—to detect their latent subspace structures. In the UKB neuroimaging dataset, within-modal self-correlation patterns (Figure 5, rows I-II and IV-V) indicated negligible residual dependence between subspaces, as expected. Dependence within subspaces is acceptable, but dependence between subspaces is not. The MSIVA default initialization workflow recovered stronger cross-modal correlations (higher CMCCs and lower CMDs) than the unimodal initialization workflow for all predefined subspace structures (Figure 5, row VI versus row III). Results from the nonlinear dependence measure also confirmed that sources in cross-modal subspaces were linked across modalities, while sources in different subspaces within each modality were independent (Appendix G Figure G.1). Among all combinations of two initialization workflows and five candidate subspace structures, the MSIVA default initialization workflow (MGPCA+ICA) with the subspace structure S_2 yielded the lowest final MISA loss value 46.775 (Table 2), suggesting that MSIVA S_2 best fits the latent structure of this dataset.

Similarly, in the patient dataset, the MSIVA default initialization workflow showed stronger cross-modal correlations (dependence) for all five subspace structures (Figures 6 and G.2, row VI versus row III). Same as the UKB dataset, MSIVA S_2 yielded the lowest final loss value 45.674 in all cases (Table 2). These results imply that MSIVA, using the default initialization and the subspace structure S_2 (including

Figure 5: **UKB neuroimaging data: Within-modal and cross-modal Pearson correlations of the recovered sources before applying post-hoc CCA.** Cross-modal subspaces are highlighted in blue while unimodal subspaces are highlighted in green. Within-modal self-correlation patterns indicated negligible residual dependence between subspaces (rows I-II and IV-V). The MSIVA default initialization workflow showed stronger cross-modal correlations (higher CMCCs and lower CMDs) than the unimodal initialization workflow for all predefined subspace structures (row VI versus row III).

Figure 6: **Patient neuroimaging data: Within-modal and cross-modal Pearson correlations of the recovered sources before applying post-hoc CCA.** Cross-modal subspaces are highlighted in blue while unimodal subspaces are highlighted in green. Within-modal self-correlation patterns indicated negligible residual dependence between subspaces (rows I-II and IV-V). The MSIVA default initialization workflow showed stronger cross-modal correlations (higher CMCCs and lower CMDs) than the unimodal initialization workflow for all predefined subspace structures (row VI versus row III).

Table 2: **Neuroimaging data: Final MISA loss values (lower is better).** MSIVA with the sub-space structure S_2 consistently yielded the lowest loss values in both multimodal neuroimaging datasets, thus it was considered as the optimal approach to capture the latent subspace structure in these two neuroimaging datasets.

Subspace structure	S_1	S_2	S_3	S_4	S_5
UK Biobank dataset					
Unimodal initialization	47.735	47.811	47.768	47.778	47.999
MSIVA default initialization	46.794	**46.775**	46.798	46.892	46.924
Patient dataset					
Unimodal initialization	47.361	47.350	47.336	47.404	47.527
MSIVA default initialization	45.775	**45.674**	45.788	45.924	45.696

five linked $2D$ subspaces), provides a better fit to the statistical relationships in these datasets.

3.3   MSIVA reveals linked phenotypic and neuropsychiatric biomarkers

After identifying the neuroimaging sources, we asked whether the linked sources were biologically mean-ingful. To answer this question, we evaluated the brain-phenotype relationships between phenotype variables and neuroimaging sources estimated by MSIVA (with the optimal subspace structure S_2 se-lected based on Table 2). In the UKB dataset, the cross-modal CCA projections in each linked subspace showed that subspaces 1, 3, 4, and 5 were significantly associated with age groups, particularly cross-modal source 9 in subspace 5 (Figure 7 rows I and II). Subspaces 1, 2, 3, and 4 showed significant sex differences, especially cross-modal source 7 in subspace 4 (Figure 7 rows III and IV). Furthermore, we used the post-CCA sources from each linked subspace to predict age and sex. The age regression and sex classification performance also confirmed that subspace 5 was strongly associated with age while subspace 4 was strongly associated with sex. More specifically, the age prediction MAE in subspace 5 was the lowest (5.378 years), and the sex classification balanced accuracy was the highest in subspace 4 (79.933%). In the patient dataset, the cross-modal CCA projections in each linked subspace revealed significant age effects in the cross-modal sources 1, 3, 8, 9, and 10, as well as SZ-related effects in both sources 9 and 10 from subspace 5 (Figure 8). These associations were verified by the age regression and diagnosis classification results. The sex effect in the patient dataset was not significant (Appendix H).

Next, we utilized a dual-coded visualization (Allen et al., 2012) for the modality- and group-specific geometric median spatial maps of the reconstructed data $\hat{\mathbf{X}}_k^{[m]} = \hat{\mathbf{A}}_k^{[m]} \hat{\mathbf{S}}_k^{[m]}$ from each representative subspace k (Figures 9 and 10). Voxel intensity was mapped to both color hue and opacity. Contours

Figure 7: **UKB neuroimaging data: Post-CCA sources from MSIVA S_2 cross-modal subspaces, color coded by age or sex.** Rows I and II show the age effect; rows III and IV show the sex effect. The Pearson correlation coefficient (r_c, c is the source index) indicates the correlation of post-CCA sources between two modalities (high r_c means similar expression profiles, not similar spatial organization). We performed age regression and sex classification, measured by mean absolute error (MAE; lower is better) and balanced accuracy (Acc.; higher is better), to assess the associations between MSIVA linked sources from each cross-modal subspace and phenotype measures (age and sex). Sources from subspace 5 yielded the best performance in age regression (MAE $= 5.378$ years), while sources from subspace 4 achieved the best performance in sex classification (Acc. $= 79.933\%$). The p-value indicates statistical difference between two age or sex groups ($p^{[1]}$: sMRI; $p^{[2]}$: fMRI; two-sample t-test, FDR correction for all subjects). Cross-modal sources 1, 5, 8, and 9 were primarily associated with aging ($p^{[1]} < 0.001, p^{[2]} < 0.001$). Cross-modal sources 1, 3, 5, and 7 were more linked to sex differences ($p^{[1]} < 0.001, p^{[2]} < 0.001$).

Figure 8: **Patient neuroimaging data: Post-CCA sources from MSIVA S_2 cross-modal subspaces, color coded by age or diagnosis labels.** Rows I and II show the age effect; rows III and IV show the SZ effect. The Pearson correlation coefficient (r_c, c is the source index) shows the correlation of post-CCA sources between two modalities (high r_c means similar expression profiles, not similar spatial organization). We performed age regression and diagnosis classification, measured by mean absolute error (MAE; lower is better) and balanced accuracy (Acc.; higher is better), to assess the associations between MSIVA linked sources from each cross-modal subspace and phenotype measures. Sources from subspace 5 yielded the best age regression and diagnosis classification performance (MAE = 10.307 years; Acc. = 61.404%), whereas subspace 2 performed similarly (MAE = 10.470 years; Acc. = 57.624%). The p-value indicates statistical difference between two age (younger versus older) or diagnosis (HC versus SZ) groups ($p^{[1]}$: sMRI; $p^{[2]}$: fMRI; two-sample t-test, FDR correction for all subjects). Cross-modal sources 1, 3, 8, 9 and 10 were primarily associated with aging ($p^{[1]} < 0.001, p^{[2]} < 0.001$). Both sources 9 and 10 in subspace 5 exhibited notable associations with SZ-related effects ($p^{[1]} < 0.05, p^{[2]} < 0.001$).

highlighted brain regions with top 15% of voxelwise cross-modal correlations for each group (negative correlations: black; positive correlations: magenta), after eliminating small clusters of voxels using morphological dilation and erosion.

In the UKB dataset, source 9 from subspace 5 showed the strongest age effect, while source 7 from subspace 4 showed the strongest sex effect (Figure 9). *Subspace 5*: age effects were observed in the cerebellum, precentral gyrus, cingulate gyrus, and paracingulate gyrus in sMRI; the occipital pole, lateral occipital cortex, superior frontal gyrus, and precuneus in fMRI. In particular, younger subjects ($<$ median age in the UKB dataset; $46-62$ years) showed higher positive voxel intensities in these areas, while older subjects ($\geq$ median age in the UKB dataset; $63-79$ years) showed negative intensities in the same areas. *Subspace 4*: Sex effects were observed in the frontal lobe, occipital lobe, and precuneus in both sMRI and fMRI. Female participants exhibited strong positive intensities in the cerebellum (sMRI), lateral occipital cortex (fMRI), subcallosal area (fMRI), and precuneus cortex (sMRI and fMRI), and negative intensities in the frontal pole and postcentral gyrus (fMRI). Male participants showed the opposite patterns. Spatial maps for the other MSIVA S_2 cross-modal subspaces in the UKB dataset are presented in Appendix I Figure I.1.

In the patient dataset, sources from subspace 5 were significantly associated with different age and diagnosis groups (Figure 10). The younger control participants showed high positive intensities in the cerebellum, temporal pole, and frontal operculum cortex in sMRI; the lingual gyrus, occipital pole, and precuneus cortex in fMRI. They also exhibited negative intensities in the middle temporal gyrus, inferior temporal gyrus, and occipital fusiform gyrus in sMRI. Additionally, both strong positive and negative voxel intensities were observed in the frontal lobe of sMRI. The older patient group ($\geq$ median age in the patient dataset; 39–65 years) showed decreased sMRI intensities in the cerebellum, paracingulate gyrus, insular cortex, frontal pole, middle frontal gyrus, and inferior frontal gyrus, increased sMRI intensities in the middle temporal gyrus and lateral occipital cortex, as well as reduced fMRI intensities in the lingual gyrus, precuneus cortex, and occipital pole, compared to age-matched controls. Spatial maps for the other MSIVA S_2 linked subspaces in the patient dataset are shown in Appendix I Figure I.2.

In addition, the number of voxels with significant cross-modal correlations ($p < 0.01$, Bonferroni correction for 44318 voxels) in older patients diagnosed with SZ (25623) was 18.6% less than their age-matched control subjects (31482) in subspace 5. Particularly, brain areas with reduced structure-function agreement included the insular cortex, lingual gyrus, occipital pole, inferior frontal gyrus, and paracingulate gyrus. Apart from subspace 5, consistent reductions were observed in the other three linked subspaces (Appendix I Figure I.3 subspaces 1-3), suggesting decreased coupling between brain structure and function for older patients.

Figure 9: **UKB neuroimaging data: Spatial maps of group-specific reconstructed data from MSIVA S_2 sources related to age and sex effects.** Axial slices show the geometric median of the reconstructed data $(\hat{\mathbf{X}}_k^{[m]})$ for each modality (sMRI or fMRI) and each group (younger: $46-62$ years, older: $63-79$ years; male or female). Voxel intensity is mapped to both color hue and opacity. Contours highlight brain areas with top 15% of voxelwise cross-modal correlations for each group (negative correlations: black; positive correlations: magenta). Histograms show voxelwise cross-modal correlations for each group (colored dashed lines: top 15% of negative or positive correlations for each group; black dotted lines: $p = \frac{0.01}{44318}$, Bonferroni correction for 44318 voxels). The reported R^2 indicates the proportion of variance captured by the subspace in each modality.

Figure 10: **Patient neuroimaging data: Spatial maps of group-specific reconstructed data from MSIVA S_2 sources related to age and SZ interaction effects.** Axial slices show the geometric median of the reconstructed data $(\hat{\mathbf{X}}_k^{[m]})$ for each modality (sMRI or fMRI) and each group (younger: $15-38$ years, older: $39-65$ years; control or patient). Voxel intensity is mapped to both color hue and opacity. Contours highlight brain areas with top 15% of voxelwise cross-modal correlations for each group (negative correlations: black; positive correlations: magenta). Histograms show voxelwise cross-modal correlations for each group (colored dashed lines: top 15% of negative or positive correlations for each group; black dotted lines: $p = \frac{0.01}{44318}$, Bonferroni correction for 44318 voxels). The reported R^2 indicates the proportion of variance captured by the subspace in each modality.

Figure 11: **Number of voxels with significant Pearson correlation between corrected brain-age delta δ_{2p} and phenotype variables.** Brain-age gap shows significant positive (left) and negative (right) associations with phenotype variables including physical exercise, time spent watching TV, sleep duration, and fluid intelligence ($p < 0.05$, false discovery rate correction for 44318 voxels, 25 phenotype variables, and 14 predictors).

3.4   Brain-age gap is associated with lifestyle factors and cognitive functions

We performed a two-stage voxelwise brain-age delta analysis using the UKB sources estimated by MSIVA using the optimal subspace structure S_2 (see Appendix B for details). We investigated whether the brain-age gap was associated with other phenotype variables by computing the Pearson correlation between δ_{2p} and each phenotype variable for each voxel. To examine effects specific to shared multimodal variability, we applied voxelwise singular value decomposition (SVD) to the combined reconstructed data from both modalities ($\hat{\mathbf{X}}_k^{[1]}$ and $\hat{\mathbf{X}}_k^{[2]}$) for each of the five cross-modal subspaces. The brain-age deltas corresponding to the top SVD-shared voxel-level features of the cross-modal subspaces 2, 4, 5 were significantly associated with various phenotype variables, including time spent watching TV, sleep duration, fluid intelligence, and physical exercise (Figure 11). In particular, predictor 5 (SVD-shared feature from cross-modal subspace 5), which showed the strongest age association (Figure 7 and Appendix B Figure B.1), positively correlated with time spent watching TV and mean time to correctly identify matches (cognitive performance), and negatively correlated with the first principal component of physical exercise variables.

Figure 12 shows the relevant spatial maps of predictor 5 (SVD 5). According to Figure 7, subspace 5 showed the strongest association with the chronological age, which aligned with the strong $\boldsymbol{\beta}_1$ coefficients and $\sigma(\boldsymbol{\delta}_{2p})$ spatial maps from the first step of brain-age delta analysis (Figure 12, panel A, rows I and II). The geometric median of brain-age delta $\boldsymbol{\delta}_{2p}$ was slightly negative (Figure 12, panel A, row III),

Figure 12: **Spatial maps of predictor 5 (SVD 5) from brain-age delta analysis.** (A) Spatial maps of β_1, standard deviation of δ_{2p}, and geometric median of δ_{2p}. Voxel value is mapped to both color hue and opacity. (B) Voxelwise correlations between δ_{2p} and phenotype variables time spent watching TV, mean time to correctly identify matches, the first principal component of physical exercise variables. The voxelwise correlation is mapped to both color hue and opacity. The contours outline the brain regions where the correlations are significant ($p < 0.05$, false discovery rate correction for 44318 voxels, 25 phenotype variables, and 14 predictors). 14431 voxels overlap within the contours in these three spatial maps.

indicating that biological age is slightly lower than chronological age (the brain appears younger). We also present spatial maps for three phenotype variables that exhibited strong associations with $\boldsymbol{\delta}_{2p}$: time spent watching TV, mean time to correctly identify matches, and the first principal component of physical exercise variables (Figure 12, panel B). Particularly, significant effects were observed in the cerebellum, postcentral gyrus, cingulate gyrus, precuneus cortex, occipital lobe, and caudate nucleus for time to watch TV; the frontal pole, precentral gyrus, and insular cortex for time to identify matches; the cerebellum, occipital fusiform gyrus, and caudate nucleus for physical exercise measure. If the correlation on the spatial map is negative (as in the first principal component of physical exercise), $\boldsymbol{\delta}_{2p}$ decreases as the phenotype score increases and the brain appears younger. If it is positive (as in time to watch TV or identify matches), $\boldsymbol{\delta}_{2p}$ increases as the phenotype score increases and the brain appears older. Therefore, the more physical exercise, the younger the brain looks; the more time spent watching TV or identifying correct matches, the older the brain looks. These findings indicate that increased physical activity and reduced TV time can potentially improve brain health.

4   Discussion

We present a novel multivariate methodology, Multimodal Subspace Independent Vector Analysis (MSIVA), to capture both cross-modal and unimodal sources. We first showed that MSIVA successfully identified the ground truth when given the correct subspace structure, according to the MISI and interference matrix results, and verified that the correct subspace structures led to the lowest loss values for all synthetic data experiments. We next applied MSIVA to two large multimodal neuroimaging datasets and identified the best-fitted latent subspace structure from five candidates. Among all combinations of different initialization workflows and subspace structures, the MSIVA default initialization method with the subspace structure S_2 output the lowest loss value, thus being considered as the best fit to the latent structure in both neuroimaging datasets. The CCA projections within each cross-modal subspace were strongly associated with age, sex and SZ-related effects, as verified through the phenotype prediction tasks. Moreover, the voxelwise brain-age delta analysis on the UKB dataset identified key non-imaging phenotype variables, including lifestyle factors and cognitive performance, that were significantly correlated with voxel-level brain-age gap.

Recommendations for model order and subspace structure priors. The success of MSIVA depends on properly identifying the model order and the candidate subspace structures. We propose estimating the number of sources using common information derived from MGPCA, guided by information-theoretic criteria (Y.-O. Li et al., 2007). Since functional imaging sources can form hierarchical clusters ranging

from two to four dimensions (Ke et al., 2019; Ma et al., 2010, 2011), we recommend evaluating subspaces with sizes within that range. Future research is encouraged to explore a broader set of subspace structures.

Evaluation of initialization workflows. We evaluated three initialization workflows that captured different amounts of joint information. Interestingly, MSIVA default initialization and unimodal initialization achieved comparative performance and outperformed a multimodal initialization workflow. One possible explanation is that multimodal initialization may overfit the cross-modal information. Conversely, unimodal initialization does not leverage any cross-modal information, which risks yielding potentially unrecoverable misalignment. MSIVA default initialization, which captures an intermediate level of cross-modal information, appears to strike the best balance among the three initialization workflows. Analyses of within- and cross-modal correlations support this hypothesis (Appendix D). Regardless of the initialization method—MSIVA default initialization or unimodal initialization—MISA effectively aligns the sources during the optimization process.

Comparison of MSIVA and MMIVA. MSIVA can be viewed as an extension of MMIVA which (1) uses a different default initialization method (MSIVA: MGPCA+ICA initialization; MMIVA: MGPCA+GICA initialization) and (2) allows for arbitrary subspace structures (MSIVA: flexible subspace structures like $S_1 - S_5$ and more; MMIVA: rigid subspace structures like an identity matrix S_5). To further investigate the relationships between the estimated sources from MSIVA and MMIVA, we compared MSIVA (with the subspace structure S_2) and MMIVA by using MSIVA S_2 sources to predict MMIVA sources, as well as using matched MMIVA sources to predict MSIVA S_2 sources. We found that a pair of MSIVA S_2 sources from each subspace could predict variability in more than two MMIVA sources, while pairs of matched MMIVA sources could also explain variability in more than two MSIVA S_2 sources (Appendix J Figures J.1-J.4). Hence, there was no perfect one-to-one mapping between MSIVA S_2 sources and MMIVA sources. We conclude that MSIVA and MMIVA apportion variability to their sources in different ways. We also note that the mismatch appears to be more pronounced in the patient dataset than in the UKB dataset, which may be related to inherent characteristics of the patient data, such as higher population heterogeneity and smaller sample size. Furthermore, we compared phenotype prediction performance using the MSIVA S_2 post-CCA sources from each cross-modal subspace and the matched pairs of MMIVA sources. MSIVA linked sources from a cross-modal subspace more accurately predicted phenotype measures (age and sex in the UKB dataset and diagnosis in the patient dataset) than pairs of independent MMIVA sources, indicating that higher-dimensional subspaces in MSIVA better capture phenotype-related variability than pairs of linked one-dimensional sources in MMIVA (Appendix J Table J.1).

Brain regions associated with phenotypes and disorders. MSIVA learns complex statistical rela-

tionships across two neuroimaging modalities, gray matter (GM) maps from T1-weighted structural MRI (sMRI) and mean-scaled amplitude of low frequency fluctuations (mALFF) maps from resting-state functional MRI (fMRI), by identifying cross-modal dependence in high-dimensional subspaces. This approach reveals structurally and functionally relevant brain networks and identifies source groups associated with specific phenotypes and disorders.

In the UKB dataset, four cross-modal sources (1, 5, 8, and 9) were significantly associated with age groups, with source 9 from subspace 5 showing the strongest association (Figure 7). Spatial maps reconstructed from subspace 5 revealed that older participants exhibited reduced sMRI intensities in the cerebellum, precentral gyrus, cingulate gyrus, and paracingulate gyrus, as well as reduced fMRI intensities in the occipital pole, precuneus cortex, superior frontal gyrus, and lateral occipital cortex (Figure 9). Several age-related brain regions identified in this study align with findings reported in the previous literature. For example, cerebellar volume has been reported to be associated with age-related decline (Jernigan et al., 2001; Luft et al., 1999; Romero et al., 2021). Hogstrom et al., 2013 has observed strong age effects in the precentral gyrus and weak age effects in the cingulate gyrus using structural brain imaging. In addition, the reductions observed in the visual areas from the mALFF feature maps are consistent with our previous findings using MMIVA (Silva et al., 2024). Prior research on functional networks has also identified significant associations with aging in the occipital lobe (Scheinost et al., 2015).

Regarding sex effects, four sources (1, 3, 5, and 7) showed significant associations with sex, with source 7 from subspace 4 showing the strongest effect (Figure 7). Spatial maps from subspace 4 indicated that males had lower sMRI intensities in the cerebellum and precuneus, lower fMRI intensities in the lateral occipital cortex and precuneus, and higher fMRI intensities in the frontal pole and postcentral gyrus (Figure 10). Previous studies have also reported sex differences in the gray matter volume of the cerebellum (Fan et al., 2010) and the precuneus cortex (Ruigrok et al., 2014), as well as in the frontal and occipital areas using functional measures (Tian et al., 2011). In particular, MMIVA has revealed that females exhibit higher cerebellar sMRI intensities than males (Silva et al., 2024).

In the patient dataset, both sources in subspace 5 showed significant associations with aging and schizophrenia (SZ) (Figure 8). Using reconstructed data from subspace 5 sources, we found that SZ patients, compared with age-matched controls, showed decreased gray matter intensities in the cerebellar, frontal, and insular cortices, and increased intensities in the middle temporal gyrus and lateral occipital cortex. They also exhibited distinct functional reductions in the precuneus, lingual gyrus, and occipital pole (Figure 10). These results align with previous findings of significantly reduced cerebellar gray matter volume in SZ patients (Moberget et al., 2018; Picard et al., 2008), as well as SZ-related alterations in

anatomical networks involving the frontal, medial temporal, and insular cortices (Bassett et al., 2008). According to cross-modal correlations, the older patient group shows reduced structure-function coupling in the insular cortex, lingual gyrus, occipital pole, inferior frontal gyrus, and paracingulate gyrus (Appendix I Figure I.3). Similarly, Cocchi et al., 2014 used diffusion MRI and resting-state fMRI to examine structure–function coupling in SZ, reporting reduced white matter integrity and functional connectivity in the frontal, temporal, thalamic, and striatal regions, as well as disrupted functional networks in the parietal, occipital, and temporal cortices where coupling remained intact. Notably, both studies identified frontal and occipital regions, suggesting consistent structure–function alterations in these cortical areas across imaging modalities.

Apart from resting-state data, MSIVA can be applied to task-related or naturalistic data to uncover structural–functional relationships under task conditions or naturalistic stimuli, as well as to other high-dimensional modalities such as genomics, thereby advancing understanding of the molecular mechanisms underlying brain development and disorders.

Limitations and future directions. A limitation of our current work is the subspace structure used in MSIVA. MSIVA selects the best-fitting subspace structure for the data from a predefined set.However, it is not computationally efficient to exhaustively evaluate the merits of other potential subspace structures. Additionally, we make two assumptions on the subspace structure: the cross-modal subspaces share the same dimensionality per modality, and the unimodal subspaces are all one-dimensional. Yet, it is possible that these assumptions might not represent the true underlying structure of the data. In future work, we plan to apply data-driven subspace structures such as the NeuroMark template (Du et al., 2020; Fu et al., 2024), or learn the underlying subspace structure from the data directly in an unsupervised manner. In this study, we favored 12 latent sources to approximate each data modality in part for computational efficiency during the combinatorial optimization, although 12 sources may be insufficient to recover all multimodal links (Song et al., 2016). Further workflow optimization and *joint* model order estimation methods are needed to effectively determine the sizes and efficiently estimate the alignment of subspaces with higher dimensionality.

We used the loss value to select the optimal subspace structure in the neuroimaging data in the absence of ground-truth information. However, its baseline can vary across initialization methods, thus it may not always serve as a gold standard of goodness of fit. However, when the same initialization workflow is used, loss values demonstrate consistency across test subspace structures, showing significant correlation with MISI values. Therefore, we recommend comparing loss values only under the same initialization. Also, we suggest evaluating performance using multiple metrics in addition to the loss value, such as the cross-modal mean correlation coefficient (CMCC) and the cross-modal minimum distance (CMD),

which measure the cross-modal subspace alignment without groud-truth information, exercising caution to not over interpret these as indicators of correct estimation but rather as diagnostics to grossly verify that the results are consistent with the tested subspace structure.

While this study focused on two modalities, MSIVA can be applied to two or more modalities by design. We encourage future work to evaluate its performance on additional data modalities. Another promising future direction is to evaluate alternative initialization methods. For instance, the geometric mean or Log-Euclidean framework—commonly regarded as more principled in applications such as diffusion tensor imaging (DTI) (Arsigny et al., 2006) and electroencephalography (EEG) (Tibermacine et al., 2024)— could serve as valid alternatives to adjust covariance matrices before aggregation in MGPCA. In addition, a key limitation is the linear mixing assumption in MSIVA. MSIVA, like ICA, assumes that each data modality is a linear mixture of latent sources, but the true mixing process in neuroimaging data may be nonlinear, especially considering the multiple physiological effects and nonlinear transformations during fMRI modeling and data preprocessing stages. To address this limitation, we plan to develop *nonlinear* latent variable models that estimate multimodal sources which are nonlinearly mixed.

5   Conclusions

Our proposed multivariate methodology MSIVA effectively captures both within- and cross-modal sources, as well as their underlying subspace structure, from multiple synthetic and neuroimaging datasets. According to brain-phenotype modeling, the estimated sources from the MSIVA cross-modal subspaces are strongly associated with phenotype variables including age, sex, and psychosis. Subsequent brain-age delta analysis shows that voxel-wise brain-age gap in the recovered cross-modal subspaces is strongly related to lifestyle and cognitive function measures. Our results support that MSIVA can be applied to uncover linked phenotypic and neuropsychiatric biomarkers of brain structure and function at the voxel level from multimodal neuroimaging data.

Ethics statement

The authors have no conflicts of interest to declare. This study used the UK Biobank Resource under Application Number 34175. Ethical approval was not required, as confirmed by the license attached to the open access data. Large language models such as Claude and ChatGPT were used to correct grammar mistakes at the sentence level.

Data and code availability

The UK Biobank dataset can be accessed at https://www.ukbiobank.ac.uk/. The BSNIP and MPRC datasets are available through the NIMH Data Archive (NDA) https://nda.nih.gov/. The COBRE dataset is available from the Collaborative Informatics and Neuroimaging Suite (COINS) https://coins.trendscenter.org/. The FBIRN phase III dataset cannot be shared directly due to the Institutional Review Board (IRB) restrictions. Individuals interested in requesting access can contact Vince D. Calhoun, vcalhoun@gsu.edu.

Analysis and visualization code for this study is publicly available at https://github.com/trendscenter/MSIVA.git. Code for brain-age delta analysis is adapted from https://www.fmrib.ox.ac.uk/datasets/BrainAgeDelta/. Code for dual-coded images is adapted from https://trendscenter.org/x/datavis/.

Author contributions

Xinhui Li: Conceptualization; formal analysis; investigation; methodology; software; validation; visualization; writing - original draft; writing - review and editing. **Peter Kochunov**: Data curation; writing - reviewing and editing. **Tulay Adali**: Funding acquisition; investigation; writing - reviewing and editing. **Rogers F. Silva**: Conceptualization; formal analysis; investigation; methodology; project administration; software; supervision; writing - review and editing. **Vince D. Calhoun**: Conceptualization; funding acquisition; investigation; project administration; resources; supervision; writing - review & editing.

Funding

This work was supported by the National Science Foundation (NSF) grants (NSF2112455 and NSF2316420) and the National Institutes of Health (NIH) grant (R01MH123610). Additionally, X.L. was supported by the Georgia Tech/Emory NIH/NIBIB Training Program in Computational Neural-engineering (T32EB025816).

Declaration of competing interests

The authors have no competing interests to declare.

Acknowledgments

We acknowledge the FBIRN team who coordinated and performed the data acquisition, including Adrian Preda, Aysenil Belger, Bryon A. Mueller, Daniel H. Mathalon, Daniel S. O'Leary, Jessica A. Turner, Juan R. Bustillo, Judith M. Ford, Kelvin O. Lim, Steven G. Potkin, and Theo G.M. van Erp.

References

Adali, T., Anderson, M., & Fu, G.-S. (2014). Diversity in independent component and vector analyses: Identifiability, algorithms, and applications in medical imaging. *IEEE Signal Processing Magazine*, *31*(3), 18–33.

Adali, T., Levin-Schwartz, Y., & Calhoun, V. D. (2015a). Multimodal data fusion using source separation: Application to medical imaging. *Proceedings of the IEEE*, *103*(9), 1494–1506.

Adali, T., Levin-Schwartz, Y., & Calhoun, V. D. (2015b). Multimodal data fusion using source separation: Two effective models based on ica and iva and their properties. *Proceedings of the IEEE*, *103*(9), 1478–1493.

Aine, C., Bockholt, H. J., Bustillo, J. R., Cañive, J. M., Caprihan, A., Gasparovic, C., Hanlon, F. M., Houck, J. M., Jung, R. E., Lauriello, J., et al. (2017). Multimodal neuroimaging in schizophrenia: Description and dissemination. *Neuroinformatics*, *15*(4), 343–364.

Akaike, H. (1998). Information theory and an extension of the maximum likelihood principle. In *Selected papers of hirotugu akaike* (pp. 199–213). Springer.

Allen, E. A., Erhardt, E. B., & Calhoun, V. D. (2012). Data visualization in the neurosciences: Overcoming the curse of dimensionality. *Neuron*, *74*(4), 603–608.

Amari, S.-I., Cichocki, A., & Yang, H. H. (1996). A New Learning Algorithm for Blind Signal Separation. *Proc NIPS 1996*, *8*, 757–763.

Arsigny, V., Fillard, P., Pennec, X., & Ayache, N. (2006). Log-euclidean metrics for fast and simple calculus on diffusion tensors. *Magnetic Resonance in Medicine: An Official Journal of the International Society for Magnetic Resonance in Medicine*, *56*(2), 411–421.

Ashburner, J., Barnes, G., Chen, C.-C., Daunizeau, J., Flandin, G., Friston, K., Kiebel, S., Kilner, J., Litvak, V., Moran, R., et al. (2014). Spm12 manual. *Wellcome Trust Centre for Neuroimaging, London, UK*, *2464*(4).

Bao, P., She, L., McGill, M., & Tsao, D. Y. (2020). A map of object space in primate inferotemporal cortex. *Nature*, *583*(7814), 103–108.

Bassett, D. S., Bullmore, E., Verchinski, B. A., Mattay, V. S., Weinberger, D. R., & Meyer-Lindenberg, A. (2008). Hierarchical organization of human cortical networks in health and schizophrenia. *Journal of Neuroscience*, *28*(37), 9239–9248. https://doi.org/10.1523/JNEUROSCI.1929-08.2008

Bell, A. J., & Sejnowski, T. J. (1995). An information-maximization approach to blind separation and blind deconvolution. *Neural computation*, *7*(6), 1129–1159.

Bernardi, S., Benna, M. K., Rigotti, M., Munuera, J., Fusi, S., & Salzman, C. D. (2020). The geometry of abstraction in the hippocampus and prefrontal cortex. *Cell*, *183*(4), 954–967.

Betzel, R. F., & Bassett, D. S. (2017). Multi-scale brain networks. *Neuroimage*, *160*, 73–83.

Boyle, L. M., Posani, L., Irfan, S., Siegelbaum, S. A., & Fusi, S. (2024). Tuned geometries of hippocampal representations meet the computational demands of social memory. *Neuron*, *112*(8), 1358–1371.

Calhoun, V. D., & Adali, T. (2008). Feature-based fusion of medical imaging data. *IEEE Transactions on Information Technology in Biomedicine*, *13*(5), 711–720.

Calhoun, V. D., Adali, T., Giuliani, N., Pekar, J., Kiehl, K., & Pearlson, G. (2006). Method for multimodal analysis of independent source differences in schizophrenia: Combining gray matter structural and auditory oddball functional data. *Human brain mapping*, *27*(1), 47–62.

Calhoun, V. D., Adali, T., Pearlson, G. D., & Kiehl, K. A. (2006). Neuronal chronometry of target detection: Fusion of hemodynamic and event-related potential data. *Neuroimage*, *30*(2), 544–553.

Calhoun, V. D., & Sui, J. (2016). Multimodal fusion of brain imaging data: A key to finding the missing link (s) in complex mental illness. *Biological psychiatry: cognitive neuroscience and neuroimaging*, *1*(3), 230–244.

Cardoso, J.-F. (1998). Multidimensional independent component analysis. *Proceedings of the 1998 IEEE International Conference on Acoustics, Speech and Signal Processing, ICASSP'98 (Cat. No. 98CH36181)*, *4*, 1941–1944.

Cavanaugh, J. E. (1999). A large-sample model selection criterion based on kullback's symmetric divergence. *Statistics & Probability Letters*, *42*(4), 333–343.

Chang, L., & Tsao, D. Y. (2017). The code for facial identity in the primate brain. *Cell*, *169*(6), 1013–1028.

Churchland, M. M., Cunningham, J. P., Kaufman, M. T., Foster, J. D., Nuyujukian, P., Ryu, S. I., & Shenoy, K. V. (2012). Neural population dynamics during reaching. *Nature*, *487*(7405), 51–56.

Cocchi, L., Harding, I. H., Lord, A., Pantelis, C., Yucel, M., & Zalesky, A. (2014). Disruption of structure–function coupling in the schizophrenia connectome. *NeuroImage: Clinical*, *4*, 779–787. https://doi.org/https://doi.org/10.1016/j.nicl.2014.05.004

Comon, P. (1994). Independent component analysis, a new concept? *Signal processing*, *36*(3), 287–314.

Correa, N. M., Adali, T., Li, Y.-O., & Calhoun, V. D. (2010). Canonical correlation analysis for data fusion and group inferences. *IEEE signal processing magazine*, *27*(4), 39–50.

Correa, N. M., Li, Y.-O., Adali, T., & Calhoun, V. D. (2008). Canonical correlation analysis for feature-based fusion of biomedical imaging modalities and its application to detection of associative networks in schizophrenia. *IEEE journal of selected topics in signal processing*, *2*(6), 998–1007.

Courellis, H. S., Minxha, J., Cardenas, A. R., Kimmel, D. L., Reed, C. M., Valiante, T. A., Salzman, C. D., Mamelak, A. N., Fusi, S., & Rutishauser, U. (2024). Abstract representations emerge in human hippocampal neurons during inference. *Nature*, 1–9.

Du, Y., Fu, Z., Sui, J., Gao, S., Xing, Y., Lin, D., Salman, M., Abrol, A., Rahaman, M. A., Chen, J., et al. (2020). Neuromark: An automated and adaptive ica based pipeline to identify reproducible fmri markers of brain disorders. *NeuroImage: Clinical*, *28*, 102375.

Fan, L., Tang, Y., Sun, B., Gong, G., Chen, Z. J., Lin, X., Yu, T., Li, Z., Evans, A. C., & Liu, S. (2010). Sexual dimorphism and asymmetry in human cerebellum: An mri-based morphometric study. *Brain research*, *1353*, 60–73.

Felleman, D. J., & Van Essen, D. C. (1991). Distributed hierarchical processing in the primate cerebral cortex. *Cerebral cortex (New York, NY: 1991)*, *1*(1), 1–47.

Franco, A. R., Ling, J., Caprihan, A., Calhoun, V. D., Jung, R. E., Heileman, G. L., & Mayer, A. R. (2008). Multimodal and multi-tissue measures of connectivity revealed by joint independent component analysis. *IEEE journal of selected topics in signal processing*, *2*(6), 986–997.

Friston, K. (2008). Hierarchical models in the brain. *PLoS computational biology*, *4*(11), e1000211.

Fu, Z., Batta, I., Wu, L., Abrol, A., Agcaoglu, O., Salman, M. S., Du, Y., Iraji, A., Shultz, S., Sui, J., et al. (2024). Searching reproducible brain features using neuromark: Templates for different age populations and imaging modalities. *NeuroImage*, *292*, 120617.

Giakoumatos, C., Nanda, P., Mathew, I., Tandon, N., Shah, J., Bishop, J., Clementz, B., Pearlson, G., Sweeney, J., Tamminga, C., et al. (2015). Effects of lithium on cortical thickness and hippocampal subfield volumes in psychotic bipolar disorder. *Journal of psychiatric research*, *61*, 180–187.

Griffanti, L., Salimi-Khorshidi, G., Beckmann, C. F., Auerbach, E. J., Douaud, G., Sexton, C. E., Zsoldos, E., Ebmeier, K. P., Filippini, N., Mackay, C. E., et al. (2014). Ica-based artefact removal and accelerated fmri acquisition for improved resting state network imaging. *Neuroimage*, *95*, 232–247.

Groves, A. R., Beckmann, C. F., Smith, S. M., & Woolrich, M. W. (2011). Linked independent component analysis for multimodal data fusion. *Neuroimage*, *54*(3), 2198–2217.

Hajnal, M. A., Tran, D., Szabó, Z., Albert, A., Safaryan, K., Einstein, M., Vallejo Martelo, M., Polack, P.-O., Golshani, P., & Orbán, G. (2024). Shifts in attention drive context-dependent subspace

encoding in anterior cingulate cortex in mice during decision making. *Nature communications*, *15*(1), 5559.

Hogstrom, L. J., Westlye, L. T., Walhovd, K. B., & Fjell, A. M. (2013). The structure of the cerebral cortex across adult life: Age-related patterns of surface area, thickness, and gyrification. *Cerebral cortex*, *23*(11), 2521–2530.

Hotelling, H. (1992). Relations between two sets of variates. In *Breakthroughs in statistics* (pp. 162–190). Springer.

Hyvärinen, A., & Hoyer, P. (2000). Emergence of phase-and shift-invariant features by decomposition of natural images into independent feature subspaces. *Neural computation*, *12*(7), 1705–1720.

Jernigan, T. L., Archibald, S. L., Fennema-Notestine, C., Gamst, A. C., Stout, J. C., Bonner, J., & Hesselink, J. R. (2001). Effects of age on tissues and regions of the cerebrum and cerebellum. *Neurobiology of aging*, *22*(4), 581–594.

Johnston, W. J., Fine, J. M., Yoo, S. B. M., Ebitz, R. B., & Hayden, B. Y. (2024). Semi-orthogonal subspaces for value mediate a binding and generalization trade-off. *Nature Neuroscience*, 1–13.

Ke, Q., Zhang, J., Wei, W., Damaševičius, R., & Woźniak, M. (2019). Adaptive independent subspace analysis of brain magnetic resonance imaging data. *IEEE Access*, *7*, 12252–12261. https://doi.org/10.1109/ACCESS.2019.2893496

Keator, D. B., van Erp, T. G., Turner, J. A., Glover, G. H., Mueller, B. A., Liu, T. T., Voyvodic, J. T., Rasmussen, J., Calhoun, V. D., Lee, H. J., et al. (2016). The function biomedical informatics research network data repository. *Neuroimage*, *124*, 1074–1079.

Kiebel, S. J., Daunizeau, J., & Friston, K. J. (2008). A hierarchy of time-scales and the brain. *PLoS computational biology*, *4*(11), e1000209.

Kim, T., Eltoft, T., & Lee, T.-W. (2006). Independent vector analysis: An extension of ica to multivariate components. *International conference on independent component analysis and signal separation*, 165–172.

Kotz, S. (1975). Multivariate distributions at a cross road. In *A modern course on statistical distributions in scientific work* (pp. 247–270). Springer.

Lahat, D., Adali, T., & Jutten, C. (2015). Multimodal data fusion: An overview of methods, challenges, and prospects. *Proceedings of the IEEE*, *103*(9), 1449–1477.

Li, X., Adali, T., Silva, R. F., & Calhoun, V. D. (2023). Multimodal subspace independent vector analysis better captures hidden relationships in multimodal neuroimaging data. *2023 IEEE 20th International Symposium on Biomedical Imaging (ISBI)*, 1–5.

Li, X., Khosravinezhad, D., Calhoun, V. D., & Silva, R. F. (2023). Evaluating trade-offs in iva of multimodal neuroimaging using cross-platform multidataset independent subspace analysis. *2023 IEEE*

20th International Symposium on Biomedical Imaging (ISBI), 1–5. https://doi.org/10.1109/ISBI53787.2023.10230492

Li, Y.-O., Adalı, T., & Calhoun, V. D. (2007). Estimating the number of independent components for functional magnetic resonance imaging data. *Human brain mapping*, *28*(11), 1251–1266.

Liu, D. C., & Nocedal, J. (1989). On the limited memory bfgs method for large scale optimization. *Mathematical programming*, *45*(1-3), 503–528.

Lopez-Paz, D., Hennig, P., & Schölkopf, B. (2013). The randomized dependence coefficient. *Advances in neural information processing systems*, *26*.

Luft, A. R., Skalej, M., Schulz, J. B., Welte, D., Kolb, R., Bürk, K., Klockgether, T., & Voigt, K. (1999). Patterns of age-related shrinkage in cerebellum and brainstem observed in vivo using three-dimensional mri volumetry. *Cerebral Cortex*, *9*(7), 712–721.

Ma, S., Correa, N. M., Li, X.-L., Eichele, T., Calhoun, V. D., & Adali, T. (2011). Automatic identification of functional clusters in fmri data using spatial dependence. *IEEE Transactions on Biomedical Engineering*, *58*(12), 3406–3417.

Ma, S., Li, X.-L., Correa, N. M., Adali, T., & Calhoun, V. D. (2010). Independent subspace analysis with prior information for fmri data. *2010 IEEE International Conference on Acoustics, Speech and Signal Processing*, 1922–1925.

Macchi, O., & Moreau, E. (1995). Self-adaptive source separation by direct or recursive networks. *Proc ICDSP 1995*, 122–129.

McKeown, M. J., Jung, T.-P., Makeig, S., Brown, G., Kindermann, S. S., Lee, T.-W., & Sejnowski, T. J. (1998). Spatially independent activity patterns in functional mri data during the stroop color-naming task. *Proceedings of the National Academy of Sciences*, *95*(3), 803–810.

Miller, K. L., Alfaro-Almagro, F., Bangerter, N. K., Thomas, D. L., Yacoub, E., Xu, J., Bartsch, A. J., Jbabdi, S., Sotiropoulos, S. N., Andersson, J. L., et al. (2016). Multimodal population brain imaging in the uk biobank prospective epidemiological study. *Nature neuroscience*, *19*(11), 1523–1536.

Moberget, T., Doan, N., Alnæs, D., Kaufmann, T., Córdova-Palomera, A., Lagerberg, T., Diedrichsen, J., Schwarz, E., Zink, M., Eisenacher, S., et al. (2018). Cerebellar volume and cerebellocerebral structural covariance in schizophrenia: A multisite mega-analysis of 983 patients and 1349 healthy controls. *Molecular psychiatry*, *23*(6), 1512–1520.

Mohammadi-Nejad, A.-R., Hossein-Zadeh, G.-A., & Soltanian-Zadeh, H. (2017). Structured and sparse canonical correlation analysis as a brain-wide multi-modal data fusion approach. *IEEE transactions on medical imaging*, *36*(7), 1438–1448.

Nadarajah, S. (2003). The kotz-type distribution with applications. *Statistics: A Journal of Theoretical and Applied Statistics*, *37*(4), 341–358.

Pandarinath, C., O'Shea, D. J., Collins, J., Jozefowicz, R., Stavisky, S. D., Kao, J. C., Trautmann, E. M., Kaufman, M. T., Ryu, S. I., Hochberg, L. R., et al. (2018). Inferring single-trial neural population dynamics using sequential auto-encoders. *Nature methods*, *15*(10), 805–815.

Picard, H., Amado, I., Mouchet-Mages, S., Olié, J.-P., & Krebs, M.-O. (2008). The role of the cerebellum in schizophrenia: An update of clinical, cognitive, and functional evidences. *Schizophrenia bulletin*, *34*(1), 155–172.

Qi, S., Sui, J., Pearlson, G., Bustillo, J., Perrone-Bizzozero, N. I., Kochunov, P., Turner, J. A., Fu, Z., Shao, W., Jiang, R., Yang, X., Liu, J., Du, Y., Chen, J., Zhang, D., & Calhoun, V. D. (2022). Derivation and utility of schizophrenia polygenic risk associated multimodal MRI frontotemporal network. *Nat Commun*, *13*(1), 4929. https://doi.org/10.1038/s41467-022-32513-8

Remington, E. D., Narain, D., Hosseini, E. A., & Jazayeri, M. (2018). Flexible sensorimotor computations through rapid reconfiguration of cortical dynamics. *Neuron*, *98*(5), 1005–1019.

Rissanen, J. (1978). Modeling by shortest data description. *Automatica*, *14*(5), 465–471.

Romero, J. E., Coupe, P., Lanuza, E., Catheline, G., Manjón, J. V., & Initiative, A. D. N. (2021). Toward a unified analysis of cerebellum maturation and aging across the entire lifespan: A mri analysis. *Human Brain Mapping*, *42*(5), 1287–1303.

Ruigrok, A. N., Salimi-Khorshidi, G., Lai, M.-C., Baron-Cohen, S., Lombardo, M. V., Tait, R. J., & Suckling, J. (2014). A meta-analysis of sex differences in human brain structure. *Neuroscience & Biobehavioral Reviews*, *39*, 34–50.

Scheinost, D., Finn, E. S., Tokoglu, F., Shen, X., Papademetris, X., Hampson, M., & Constable, R. T. (2015). Sex differences in normal age trajectories of functional brain networks. *Human brain mapping*, *36*(4), 1524–1535.

Schijven, D., Postema, M. C., Fukunaga, M., Matsumoto, J., Miura, K., de Zwarte, S. M., Van Haren, N. E., Cahn, W., Hulshoff Pol, H. E., Kahn, R. S., et al. (2023). Large-scale analysis of structural brain asymmetries in schizophrenia via the enigma consortium. *Proceedings of the National Academy of Sciences*, *120*(14), e2213880120.

Semedo, J. D., Zandvakili, A., Machens, C. K., Byron, M. Y., & Kohn, A. (2019). Cortical areas interact through a communication subspace. *Neuron*, *102*(1), 249–259.

She, L., Benna, M. K., Shi, Y., Fusi, S., & Tsao, D. Y. (2024). Temporal multiplexing of perception and memory codes in it cortex. *Nature*, 1–8.

Silva, R. F., Plis, S. M., Adalı, T., & Calhoun, V. D. (2014). A statistically motivated framework for simulation of stochastic data fusion models applied to multimodal neuroimaging. *NeuroImage*, *102, Part 1*, 92–117. https://doi.org/10.1016/j.neuroimage.2014.04.035

Silva, R. F., Damaraju, E., Li, X., Kochunov, P., Ford, J. M., Mathalon, D. H., Turner, J. A., van Erp, T. G., Adali, T., & Calhoun, V. D. (2024). A method for multimodal iva fusion within a misa

unified model reveals markers of age, sex, cognition, and schizophrenia in large neuroimaging studies. *Human Brain Mapping*, *45*(17), e70037.

Silva, R. F., Plis, S. M., Adalı, T., Pattichis, M. S., & Calhoun, V. D. (2020). Multidataset independent subspace analysis with application to multimodal fusion. *IEEE Transactions on Image Processing*, *30*, 588–602.

Smith, S. M., Elliott, L. T., Alfaro-Almagro, F., McCarthy, P., Nichols, T. E., Douaud, G., & Miller, K. L. (2020). Brain aging comprises many modes of structural and functional change with distinct genetic and biophysical associations. *elife*, *9*, e52677.

Smith, S. M., Nichols, T. E., Vidaurre, D., Winkler, A. M., Behrens, T. E., Glasser, M. F., Ugurbil, K., Barch, D. M., Van Essen, D. C., & Miller, K. L. (2015). A positive-negative mode of population covariation links brain connectivity, demographics and behavior. *Nature neuroscience*, *18*(11), 1565–1567.

Smith, S. M., Vidaurre, D., Alfaro-Almagro, F., Nichols, T. E., & Miller, K. L. (2019). Estimation of brain age delta from brain imaging. *Neuroimage*, *200*, 528–539.

Song, Y., Schreier, P. J., Ramírez, D., & Hasija, T. (2016). Canonical correlation analysis of high-dimensional data with very small sample support. *Signal Processing*, *128*, 449–458.

Sui, J., Adali, T., Yu, Q., Chen, J., & Calhoun, V. D. (2012). A review of multivariate methods for multimodal fusion of brain imaging data. *Journal of neuroscience methods*, *204*(1), 68–81.

Sui, J., He, H., Pearlson, G. D., Adali, T., Kiehl, K. A., Yu, Q., Clark, V. P., Castro, E., White, T., Mueller, B. A., et al. (2013). Three-way (n-way) fusion of brain imaging data based on mcca+jica and its application to discriminating schizophrenia. *NeuroImage*, *66*, 119–132.

Sui, J., Pearlson, G., Caprihan, A., Adali, T., Kiehl, K. A., Liu, J., Yamamoto, J., & Calhoun, V. D. (2011). Discriminating schizophrenia and bipolar disorder by fusing fmri and dti in a multimodal cca+ joint ica model. *Neuroimage*, *57*(3), 839–855.

Szabó, Z., Póczos, B., & L\Horincz, A. (2012). Separation theorem for independent subspace analysis and its consequences. *Pattern Recognit*, *45*(4), 1782–1791. https://doi.org/10.1016/j.patcog.2011.09.007

Tamminga, C. A., Pearlson, G., Keshavan, M., Sweeney, J., Clementz, B., & Thaker, G. (2014). Bipolar and schizophrenia network for intermediate phenotypes: Outcomes across the psychosis continuum. *Schizophrenia bulletin*, *40*(Suppl_2), S131–S137.

Tian, L., Wang, J., Yan, C., & He, Y. (2011). Hemisphere-and gender-related differences in small-world brain networks: A resting-state functional mri study. *Neuroimage*, *54*(1), 191–202.

Tibermacine, I. E., Russo, S., Tibermacine, A., Rabehi, A., Nail, B., Kadri, K., & Napoli, C. (2024). Riemannian geometry-based eeg approaches: A literature review. *arXiv preprint arXiv:2407.20250*.

Wang, J., Narain, D., Hosseini, E. A., & Jazayeri, M. (2018). Flexible timing by temporal scaling of cortical responses. *Nature neuroscience*, *21*(1), 102–110.

Zeki, S., & Shipp, S. (1988). The functional logic of cortical connections. *Nature*, *335*(6188), 311–317.

Zhang, Y.-D., Dong, Z., Wang, S.-H., Yu, X., Yao, X., Zhou, Q., Hu, H., Li, M., Jiménez-Mesa, C., Ramirez, J., et al. (2020). Advances in multimodal data fusion in neuroimaging: Overview, challenges, and novel orientation. *Information Fusion*, *64*, 149–187.

Zhao, N., Yuan, L.-X., Jia, X.-Z., Zhou, X.-F., Deng, X.-P., He, H.-J., Zhong, J., Wang, J., & Zang, Y.-F. (2018). Intra-and inter-scanner reliability of voxel-wise whole-brain analytic metrics for resting state fmri. *Frontiers in neuroinformatics*, *12*, 54.

Appendix A   Data acquisition and preprocessing

A.1   UK Biobank dataset

A.1.1   Acquisition parameters

T1-weighted structural MRI (sMRI) images were acquired using a 3D MPRAGE sequence with the following parameters: repetition time $(\text{TR}) = 2000$ms, inversion time $(\text{TI}) = 880$ms, in-plane acceleration factor $= 2$, voxel size $= 1 \times 1 \times 1$mm^3, acquisition matrix $= 208 \times 256 \times 256$. Resting-state functional MRI (fMRI) were acquired with the following parameters: $\text{TR} = 735$ms, echo time $(\text{TE}) = 39$ms, multiband factor $= 8$, in-plane acceleration factor $= 1$, flip angle $= 52°$, voxel size $= 2.4 \times 2.4 \times 2.4$mm^3, acquisition matrix $= 88 \times 88 \times 64$.

A.1.2   Preprocessing steps

For sMRI preprocessing, we performed tissue segmentation and normalization to the Montreal Neurological Institute (MNI) template using the statistical parametric mapping toolbox (SPM12, http://www.fil.ion.ucl.ac.uk/spm/) (Ashburner et al., 2014), leading to gray matter (GM), white matter (WM), and cerebrospinal fluid (CSF) tissue probability maps. Next, the normalized GM tissue probability maps were spatially smoothed using a Gaussian kernel with a full width at half maximum (FWHM) $= 10$mm. The smoothed images were then resampled to $3 \times 3 \times 3$mm^3. We next defined a group mask for GM voxels. Specifically, an average GM tissue probability map from all subjects was obtained from the normalized GM tissue probability maps at $1 \times 1 \times 1$mm^3 resolution. This group-average GM map was binarized at a threshold of 0.2 and resampled to $3 \times 3 \times 3$mm^3 resolution, resulting in 44318 voxels.

For fMRI preprocessing, we utilized the distortion corrected, FIX-denoised (Griffanti et al., 2014), normalized fMRI data from the UK Biobank data resource to compute subject-specific amplitude of low frequency fluctuations (ALFF) maps, defined as the area under the low frequency band $[0.01 - 0.08$ Hz] power spectrum of each voxel time course in each scan. We then calculated a mean-scaled ALFF (mALFF) map for each subject, which is the subject-specific ALFF map divided by its global mean ALFF value for greater test-retest reliability (Zhao et al., 2018). The mALFF maps were smoothed using a 6mm FWHM Gaussian filter and resampled to $3 \times 3 \times 3$mm^3 isotropic voxels. We applied the same group-average GM mask for the mALFF maps, resulting in 44318 voxels.

A.2 Patient datasets

A.2.1 Acquisition parameters

BSNIP. We used the BSNIP dataset collected at two sites: (1) Baltimore with a 3-Tesla Siemens Trio Tim scanner and (2) Hartford with a 3-Tesla Siemens Allegra scanner. Isotropic T1-weighted MPRAGE scans were acquired using the following parameters: TR $= 6.7$ms, TE $= 3.1$ms, flip angle $= 8°$, matrix size $= 256 \times 240$, total scan time $= 10 : 52.6$min, 170 sagittal slices, slice thickness $= 1$mm, voxel size $= 1 \times 1 \times 1.2$mm^3 (Giakoumatos et al., 2015). Resting-state fMRI scans were obtained with the following parameters: (1) Baltimore, TR $= 2210$ms, TE $= 30$ms, flip angle $= 70°$, number of slices $= 36$, voxel size $= 3.4 \times 3.4 \times 4$mm^3, and 140 time points; (2) Hartford, TR $= 1500$ms, TE $= 27$ms, flip angle $= 70°$, number of slices $= 29$, voxel size $= 3.4 \times 3.4 \times 5$mm^3, and 210 time points.

COBRE. The COBRE dataset was collected at a single site using a 3-Tesla Siemens Tim Trio scanner. A high-resolution T1-weighted multi-echo MPRAGE sequence was used with the following parameters: TR $= 2530$ms, TE $= [1.64, 3.5, 5.36, 7.22, 9.08]$ms, TI $= 900$ms, flip angle $= 7°$, acquisition matrix $= 256 \times 256 \times 176$, voxel size $= 1 \times 1 \times 1$mm^3, number of echos $= 5$, pixel bandwidth $= 650$Hz, total scan time $= 6$min. Resting-state fMRI scans were collected with a standard single-shot full k-space echo-planar imaging (EPI) sequence: TR $= 2000$ms, TE $= 29$ms, voxel size $= 3.75 \times 3.75 \times 4.55$mm^3, slice gap $= 1.05$mm, flip angle $= 75°$, number of slices $= 32$, field of view (FOV) $= 240 \times 240$mm^2, matrix size $= 64 \times 64$, and 149 volumes. See https://fcon_1000.projects.nitrc.org/indi/retro/cobre.html for more details.

FBIRN. The FBIRN phase III dataset was collected from seven sites. Out of seven sites, six sites used 3-Tesla Siemens Tim Trio scanners and one site used a 3-Tesla General Electric (GE) Discovery MR750 scanner. A high-resolution Siemens MPRAGE sequence was acquired with the following parameters: TR/TE/TI $= 2300/2.94/1100$ms, flip angle $= 9°$, acquisition matrix $= 256 \times 256 \times 160$. Likewise, a GE IR-SPGR sequence was acquired with the following parameters: TR/TE/TI $= 5.95/1.99/45$ms, flip angle $= 12°$, acquisition matrix $= 256 \times 256 \times 166$, FOV $= 220 \times 220$mm^2, voxel size $= 0.86 \times 0.86 \times 1.2$ mm^3, collected in the sagittal plane with GRAPPA/ASSET acceleration factor $= 2$, and NEX $= 1$ (Qi et al., 2022). The same resting-state fMRI parameters were used across all seven sites: a standard gradient EPI sequence, TR/TE $= 2000/30$ms, voxel size $= 3.4375 \times 3.4375 \times 4$mm^3, slice gap $= 1$mm, flip angle $= 77°$, FOV $= 220 \times 220$mm^2, and 162 volumes (Qi et al., 2022).

MPRC. The MPRC dataset was collected at three sites, each using a different 3-Tesla Siemens scanner, with a standard EPI sequence. T1-weighted 3D MPRAGE sequence was collected in the saggital plane with voxel size $= 1 \times 1 \times 1$mm^3 using a Siemens Allegra scanner (TE/TR/TI $= 4.3/2500/1000$ms, flip

angle $= 8°$) or a Siemens Trio scanner (TE/TR/TI $= 2.9/2300/900$ms, flip angle $= 9°$) (Schijven et al., 2023). Resting-state fMRI scans were collected using the following scanners and parameters: 3-Tesla Siemens Allegra scanner (TR/TE $= 2000/27$ms, voxel size $= 3.44 \times 3.44 \times 4$mm^3, FOV $= 220 \times 220$mm^2, and 150 volumes); 3-Tesla Siemens Trio scanner (TR/TE $= 2210/30$ms, voxel size $= 3.44 \times 3.44 \times 4$mm^3, FOV $= 220 \times 220$mm^2, and 140 volumes); and 3-Tesla Siemens Tim Trio scanner (TR/TE $= 2000/30$ms, voxel size $= 1.72 \times 1.72 \times 4$mm^3, FOV $= 220 \times 220$mm^2, and 444 volumes) (Qi et al., 2022).

A.2.2 Preprocessing steps

All sMRI datasets were preprocessed using SPM12, following the steps described in Qi et al., 2022. Specifically, the data were normalized to the MNI template using unified segmentation, resampled to $3 \times 3 \times 3$mm^3, and segmented into GM, WM, and CSF using modulated normalization, leading to GM volume maps. These GM volume maps were then smoothed using a 6mm FWHM Gaussian kernel. To ensure proper segmentation for all subjects, outlier detection was performed using spatial Pearson correlation with the template image.

All fMRI datasets underwent the preprocessing steps as outlined in Qi et al., 2022: removal of the initial five scans to eliminate T1 equilibration effects, slice timing correction, realignment, normalization to the EPI template with $3 \times 3 \times 3$mm^3 resolution, spatial smoothing using a 6mm FWHM Gaussian kernel, regression of nuisance covariates (including six head motion parameters, CSF, WM) and global signal from the voxelwise time course using a general linear model, and computation of the mALFF maps.

Appendix B   Voxelwise brain-age delta analysis on UK Biobank data

We performed a voxelwise brain-age delta analysis using the estimated sources $\hat{\mathbf{S}}$ from MSIVA subspace structure S_2 in the UK Biobank dataset. We describe the steps to construct imaging-derived predictors as follows.

1. Reconstruction. We first reconstructed the modality- and subspace-specific imaging feature $\hat{\mathbf{X}}_k^{[m]} = \hat{\mathbf{A}}_k^{[m]}\hat{\mathbf{S}}_k^{[m]}$ for each of five multimodal subspaces ($\hat{\mathbf{A}}_k^{[m]} \in \mathbb{R}^{V \times 2}$, $\hat{\mathbf{S}}_k^{[m]} \in \mathbb{R}^{2 \times N}$, $k \in \{1, ..., 5\}$) and each of four unimodal subspaces ($\hat{\mathbf{A}}_k^{[m]} \in \mathbb{R}^{V \times 1}$, $\hat{\mathbf{S}}_k^{[m]} \in \mathbb{R}^{1 \times N}$, $k \in \{6, ..., 9\}$), where k is the subspace index. Here, subspaces 6 and 7 are used exclusively for sMRI and subspaces 8 and 9 are used exclusively for fMRI.

2. Singular value decomposition (SVD). SVD was applied on cross-modal features to capture the shared information between two modalities. For each voxel i in the reconstructed imaging data $\hat{\mathbf{X}}_{k[i,:]}^{[m]} \in \mathbb{R}^N$ from each of five cross-modal subspaces, we concatenated the two modalities $\hat{\mathbf{X}}_{k[i,:]} = \left[\hat{\mathbf{X}}_{k[i,:]}^{[1]}, \hat{\mathbf{X}}_{k[i,:]}^{[2]}\right]$, $\hat{\mathbf{X}}_{k[i,:]} \in \mathbb{R}^{N \times 2}$, normalized $\hat{\mathbf{X}}_{k[i,:]}$ along the rows[8], and then performed SVD on $\hat{\mathbf{X}}_{k[i,:]}$, that is, $\hat{\mathbf{X}}_{k[i,:]} = \mathbf{U}\boldsymbol{\Sigma}\mathbf{V}^\top$. Next, we multiplied $\hat{\mathbf{X}}_{k[i,:]}$ by the first left singular vector $\mathbf{V}_{[:,1]} \in \mathbb{R}^{2 \times 1}$ corresponding to the largest singular value $\lambda_{\max}$, leading to $\hat{\mathbf{X}}_{k[i,:]}^{\mathsf{SVD}} = \hat{\mathbf{X}}_{k[i,:]}\mathbf{V}_{[:,1]}$, $\hat{\mathbf{X}}_{k[i,:]}^{\mathsf{SVD}} \in \mathbb{R}^{N \times 1}$. We then normalized $\hat{\mathbf{X}}_{k[i,:]}^{\mathsf{SVD}}$ to obtain the normalized $\hat{\mathbf{X}}_{k[i,:]}^{\mathsf{SVD}'}$.

3. Partialization and normalization. To assess each predictor's unique variance, we partialized and normalized $\hat{\mathbf{X}}_{k[i,:]}^{\mathsf{SVD}'}$ (from five cross-modal subspaces in sMRI) and $\hat{\mathbf{X}}_{k[i,:]}^{[m]}$ (from four unimodal subspaces in both modalities) to remove SVD-related confounds from $\hat{\mathbf{X}}_{k[i,:]}^{[m]}$, leading to $\hat{\mathbf{X}}_{k[i,:]}^{[m]'}$. This partialization and normalization procedure follows Smith et al., 2020 to remove confounds from other predictors.

4. Concatenation. We concatenated the SVD results from Step 2 (without partialization or extra normalization) $\hat{\mathbf{X}}_{k[i,:]}^{\mathsf{SVD}'}$ from five cross-modal subspaces, and modality-specific partialized and normalized $\hat{\mathbf{X}}_{k[i,:]}^{[m]'}$ from five cross-modal subspaces and four unimodal subspaces, resulting in 14 predictors in total, $\hat{\mathbf{X}}_i = \left[\hat{\mathbf{X}}_{1[i,:]}^{\mathsf{SVD}'}, \ldots, \hat{\mathbf{X}}_{5[i,:]}^{\mathsf{SVD}'}, \hat{\mathbf{X}}_{1[i,:]}^{[1]'}, \ldots, \hat{\mathbf{X}}_{7[i,:]}^{[1]'}, \hat{\mathbf{X}}_{8[i,:]}^{[2]'}, \hat{\mathbf{X}}_{9[i,:]}^{[2]'}\right]$, $\hat{\mathbf{X}}_i \in \mathbb{R}^{N \times 14}$.

For each voxel i, we performed a two-stage brain age prediction where the first stage estimates the initial delta and the second stage further removes age dependence and other confound factors from the delta

---

[8]We removed mean and divided by standard deviation along the rows (subject dimension) of $\hat{\mathbf{X}}_{k[i,:]}$.

(Smith et al., 2019, 2020):

$$\boldsymbol{\delta}_1 = \hat{\mathbf{X}}_i \boldsymbol{\beta}_1 - \mathbf{y}, \tag{19}$$

$$\boldsymbol{\delta}_2 = \boldsymbol{\delta}_1 - \mathbf{Y} \boldsymbol{\beta}_2, \tag{20}$$

where $\mathbf{y} \in \mathbb{R}^N$ is the chronological age after removing the mean age across subjects. $\mathbf{Y} \in \mathbb{R}^{N \times 10}$ includes the confound variables:

1. the demeaned linear age term,

2. the demeaned quadratic age term after regressing out the linear age effects and normalizing to have the same standard deviation as the linear age term,

3. the demeaned cubic age term after regressing out the linear and quadratic age effects and normalizing to have the same standard deviation as the linear age term,

4. sex,

5. the interaction between sex and each of the three age terms,

6. the framewise displacement variable,

7. the spatial normalization variables from sMRI and fMRI.

Finally, we partialized $\boldsymbol{\delta}_2$ to remove residual associations, obtaining the partialized brain-age delta, $\boldsymbol{\delta}_{2p}$.

Figure B.1: **Spatial maps of β_1 in voxelwise brain-age delta analysis.** Voxel intensity is mapped to both color hue and opacity. SVD 5 showed the strongest age association among all predictors.

Appendix C   UK Biobank phenotype variables

We used 25 phenotype variables, including lifestyle measures and cognitive test scores, to investigate their associations with brain-age delta. We describe the process of selecting the phenotype variables as follows.

We first excluded variables with extreme values from original 64 non-imaging variables using a two-step approach:

1. We calculated the sum of squared absolute median deviations (d) for each variable.

2. We excluded any variable where $\max(d) > 100 \times \text{mean}(d)$, as these extreme outliers could skew statistical analysis.

This initial screening resulted in 54 variables, including age, sex, fluid intelligence, physical activity measures, alcohol intake frequency, cognitive test scores, time spent watching TV, and sleep duration. We further reduced or excluded phenotype variables from these 54 variables as described below:

1. We applied PCA to decompose 28 physical exercise variables into 8 principal components.

2. We removed five age-related variables due to high correlation with other age variables. These variables were "age when attended assessment center", "age when first sexual intercourse", "age started wearing glasses", "years since first sexual intercourse", and "years since started wearing glasses".

3. We excluded two variables related to a cognitive test ("time to answer" and "log time to answer") due to distinct population distributions resulting from two different cognitive tests used during data collection.

4. Finally, we removed the sex variable and another log variable ("log pm score").

This selection process ultimately yielded 25 variables in total for our analysis, as listed in Table C.1.

Table C.1: **54 UK Biobank phenotype variables.** 28 physical exercise variables in purple were reduced to 8 principal components by PCA. Variables in red were excluded in brain-age delta analysis. Variables without IDs were created by R.F.S. based on the original variables and not included in the original UK Biobank dataset.

Variable ID	Variable Name
f399 2 2	number of incorrect matches in round
f400 2 2	time to complete round
f699 2 0	length of time at current address
f864 2 0	number of daysweek walked 10 minutes
f874 2 0	duration of walks
f884 2 0	number of daysweek of moderate physical activity 10 minutes
f894 2 0	duration of moderate activity
f904 2 0	number of daysweek of vigorous physical activity 10 minutes
f914 2 0	duration of vigorous activity
f943 2 0	frequency of stair climbing in last 4 weeks
f971 2 0	frequency of walking for pleasure in last 4 weeks
f981 2 0	duration walking for pleasure
f991 2 0	frequency of strenuous sports in last 4 weeks
f1001 2 0	duration of strenuous sports
f1011 2 0	frequency of light diy in last 4 weeks
f1021 2 0	duration of light diy
f1050 2 0	time spend outdoors in summer
f1060 2 0	time spent outdoors in winter
f1070 2 0	time spent watching television tv
f1080 2 0	time spent using computer
f1160 2 0	sleep duration
f1438 2 0	bread intake
f1488 2 0	tea intake
f1498 2 0	coffee intake
f1558 2 0	alcohol intake frequency
f2139 2 0	age first had sexual intercourse
f2217 2 0	age started wearing glasses or contact lenses
f2624 2 0	frequency of heavy diy in last 4 weeks
f2634 2 0	duration of heavy diy
f3637 2 0	frequency of other exercises in last 4 weeks
f3647 2 0	duration of other exercises
f4288 2 0	time to answer
f4609 2 0	longest period of depression
f20016 2 0	fluid intelligence score
f20023 2 0	mean time to correctly identify matches
f20128 2 0	number of fluid intelligence questions attempted within time limit
f21003 2 0	age when attended assessment centre
f31 0 0	sex
	total hours walked 10 minutes
	total hours moderate physical activity 10 minutes
	total hours vigorous physical activity 10 minutes
	total hours of walking for pleasure in last 4 weeks
	total hours of strenuous sports in last 4 weeks
	total hours of other exercises in last 4 weeks
	total hours of light diy in last 4 weeks
	total hours of heavy diy in last 4 weeks
	number of physical activities wrt walking for pleasure
	years since first sexual intercourse
	years since started wearing glasses
	log time to answer
	inverse log duration screen displayed
	inverse log number of attempts
	log pm score
	fluid intelligence interaction

Appendix D   Comparison of initialization workflows

The choice of weight matrix initialization may affect whether the model can converge to the optimal solution. To better understand the impact of different initialization workflows, we evaluated the initial source estimates obtained after each initialization. Figure D.1 displays the within-modal source correlations (diagonal blocks) and cross-modal source correlations (off-diagonal blocks) following different initialization approaches.

Unimodal initialization applies PCA and ICA to each data modality separately. Since independent sources are estimated without accessing information from other datasets, this method can successfully separate sources within each modality but cannot align sources across modalities, nor group them by subspace within modality (Figure D.1, top row).

In contrast, MSIVA default initialization first applies MGPCA to both datasets simultaneously, followed by ICA on each dimensionally reduced dataset separately. MGPCA identifies orthogonal components using the average covariance matrix across modalities, which preserves cross-modal information during dimensionality reduction. ICA then further separates these MGPCA-derived components into independent sources. This "hybrid" approach effectively separates unimodal sources and identifies substantial cross-modal linkage (evidenced by diagonal structure in cross-modal correlations) for smaller subspace structures (S_1, S_2, and S_5). However, it fails to capture the cross-modal linkage for more complex subspace structures such as S_3 and S_4 (Figure D.1, middle row).

Lastly, multimodal initialization sequentially applies MGPCA and GICA on both datasets. The method demonstrates reasonable results for S_5, which contains one-dimensional sources, but exhibits limitations for other subspace structures containing higher-dimensional subspaces. Specifically, it neither properly separates sources within each modality (high off-diagonal values in within-modal correlations), nor aligns sources between the two modalities, as indicated by the absence of diagonal structure in cross-modal correlations (Figure D.1, bottom row).

While perfect separation and alignment are not expected immediately after initialization—since subsequent combinatorial and numerical optimization steps will further refine these results—achieving a well-balanced combination of unimodal separation and cross-modal alignment during initialization generally increases the likelihood of convergence to the optimal solution.

Figure D.1: **Within-modal and cross-modal source correlations across initialization workflows.** From top to bottom, each row shows within- and cross-modal source correlations after unimodal, MSIVA default, and multimodal initialization, respectively. Diagonal blocks represent within-modal correlations; off-diagonal blocks represent cross-modal correlations. MSIVA default initialization achieved the lowest MISI values in four of the five evaluated subspace structures.

Appendix E   Loss curves during numerical optimization

Figure 3 summarizes experiments conducted under 25 conditions and 3 initialization workflows (75 experiments in total). In each experiment, we alternated between combinatorial optimization and numerical optimization 10 times on synthetic data. Each numerical optimization run was capped at 150 iterations. As shown in Figure E.1, the normalized loss curves indicate that most numerical optimization runs converged within 150 iterations (i.e., the loss plateaued), except for multimodal initialization, likely due to suboptimal initialization (Figure D.1). All experiments were executed on an HPC node with two CPUs and 100 GB of memory. Runtime varied substantially across test subspace structures, ranging from 1 minute for S_5^{Test} to over 250 minutes for S_4^{Test}. Specifically, S_5^{Test} completed in 1 minute, S_1^{Test} in approximately 20 minutes, S_3^{Test} in around 35 minutes, and S_2^{Test} in about 120 minutes. For S_4^{Test}, the combinatorial optimization phase was substantially more time-consuming, requiring 253–489 minutes.

Figure E.2 shows the relationships between MISI and other metrics across five subspace structures when the ground-truth and test subspace structures are correctly matched. The loss values were significantly correlated with the MISI values (unimodal initialization: $R^2 \geq 0.95$; MSIVA default initialization: $R^2 \geq 0.58$), suggesting that loss serves as a valid proxy for MISI in both initialization workflows.

The multimodal mean correlation coefficient (MMCC), which leveraged ground-truth information, showed strong agreement with MISI under both unimodal and MSIVA default initialization workflows (unimodal: $R^2 \geq 0.82$; MSIVA: $R^2 \geq 0.79$). Similarly, the multimodal minimum distance (MMD) was significantly correlated with MISI for both initialization strategies (unimodal: $R^2 \geq 0.69$; MSIVA: $R^2 \geq 0.77$). Notably, the minimum MMD value was closer to zero than the minimum value of $1-$MMCC, implying that MMD may serve as a more reliable metric than MMCC.

The cross-modal metrics—the cross-modal mean correlation coefficient (CMCC) and the cross-modal minimum distance (CMD)—do not leverage ground-truth information, and were not particularly reliable, as they did not exhibit consistently significant correlations with MISI across the five evaluated subspace structures. Specifically, significant associations were observed only for subspace structures S_2 and S_4 in the unimodal initialization workflow, and for S_2 in the MSIVA default initialization workflow. Among all measures that did not use ground-truth information, only the loss was meaningfully associated with MISI.

Figure E.1: **Loss curves during numerical optimization.** Panels a, g, m, s, and y ($S^{\text{GT}} = S^{\text{Test}}$) confirm that only the unimodal and MSIVA workflows consistently converge to the lowest loss values and detect the correct subspace structures. Vertical colored stripes mark the start and end of numerical optimization runs (≤ 150 iterations each), with combinatorial optimization performed between successive numerical runs. The *left* y-axis shows the normalized loss values after subtracting the minimum loss across the five different test subspace structures *under the same initialization workflow*. The *right* y-axis shows other performance metrics at the end of each numerical optimization run: MISI, $1-$MMCC, $1-$CMCC, MMD, and CMD. The lowest (best) MISI within a workflow is highlighted in a circle ($\bigcirc/\bigcirc$), while the lowest MISI across all three workflows is highlighted in a square ($\square/\square$).

Figure E.2: **Scatter plots of MISI versus other metrics.** Each row presents scatter plots of MISI versus another metric (normalized loss, $1-$MMCC, $1-$CMCC, MMD, or CMD) across the five evaluated subspace structures, under the condition that the test subspace structure matches the ground truth. Each point corresponds to the result from the final iteration of a combinatorial optimization run. In each plot, the blue or orange vertical dotted line indicates the minimum MISI value, and the horizontal dotted line indicates the minimum value of the corresponding comparison metric. The gray vertical line marks the MISI threshold of 0.1, which is commonly used as a heuristic criterion for good source separation (MISI ≤ 0.1). For points with MISI ≤ 0.1, we fit a linear regression line and reported R^2 and p-values. The number of asterisks (1, 2, 3, 4) denotes the level of statistical significance in the linear regression analysis ($p < 0.05$, $p < 0.01$, $p < 0.001$, $p < 0.0001$), while "ns" indicates a non-significant result. Normalized loss values were computed by subtracting the minimum loss across the five test subspace structures within the same initialization workflow. Loss values were significantly correlated with MISI, suggesting that loss serves as a valid proxy for MISI. The multimodal metrics, $1-$MMCC and MMD, which leveraged ground-truth information, also showed significant correlations with MISI. In contrast, the cross-modal metrics, $1-$CMCC and CMD, which did not use ground-truth information, were generally not correlated with MISI. Among all measures that did not use ground-truth information, only the loss was meaningfully associated with MISI.

Appendix F   Order selection

In order to select the model order (the number of sources), we evaluated three information-theoretic criteria (Y.-O. Li et al., 2007) based on the singular values of the MGPCA covariance matrix derived from each multimodal neuroimaging dataset: the Akaike Information Criterion (AIC) (Akaike, 1998), the Kullback-Leibler Information Criterion (KIC) (Cavanaugh, 1999), and the Minimum Description Length (MDL) (Rissanen, 1978).

As shown in Figure F.1, the selected orders using AIC, KIC, and MDL are 25, 18, and 10 for the UKB neuroimaging dataset, and 12, 8, and 4 for the patient dataset, respectively. In the patient dataset, the maximum selected order is 12. In the UKB dataset, although AIC and KIC suggest higher orders, we note that the curves are nearly flat as the number of sources is 12 or higher, suggesting that 12 sources already capture a large amount of the information. Considering the increasing computational cost with more sources, we chose 12 sources for both datasets in this study.

Figure F.1: **Order selection for multimodal neuroimaging data.** The selected orders using AIC, KIC, and MDL are 25, 18, and 10 for the UKB neuroimaging dataset, and 12, 8, and 4 for the patient dataset, respectively. Twelve sources captured the majority of information in the patient dataset and a substantial amount in the UK Biobank dataset. Therefore, we used twelve sources for both datasets in this study.

Appendix G Nonlinear source dependence in neuroimaging data

Apart from Pearson correlation, we calculated the randomized correlation coefficents (RDCs) (Lopez-Paz et al., 2013) to measure source correlations of random nonlinear copula projections in neuroimaging data. The RDC results (Figures G.1, G.2) were largely consistent with those measured by the linear Pearson correlation coefficient (Figures 5, 6). The low RDC values outside the block subspace structure indicated very weak residual dependence between subspaces, suggesting that different subspaces are nearly independent. The MSIVA default initialization workflow exhibited stronger cross-modal correlations (higher CMCCs and lower CMDs) than the unimodal initialization workflow for all predefined subspace structures (row VI versus row III) in both neuroimaging datasets.

Figure G.1: **UKB neuroimaging data: Within-modal and cross-modal correlations of random nonlinear copula projections of the recovered sources before applying post-hoc CCA.** Cross-modal subspaces are highlighted in blue while unimodal subspaces are highlighted in green. Within-modal self-correlation patterns showed very weak residual dependence between subspaces (rows I-II and IV-V). The MSIVA default initialization workflow exhibited stronger cross-modal correlations (higher CMCCs and lower CMDs) than the unimodal initialization workflow for all predefined subspace structures (row VI versus row III).

Figure G.2: **Patient neuroimaging data: Within-modal and cross-modal correlations of random nonlinear copula projections of the recovered sources before applying post-hoc CCA.** Cross-modal subspaces are highlighted in blue while unimodal subspaces are highlighted in green. Within-modal self-correlation patterns showed weak residual dependence between subspaces (rows I-II and IV-V). The MSIVA default initialization workflow showed stronger cross-modal correlations (higher CMCCs and lower CMDs) than the unimodal initialization workflow for all predefined subspace structures (row VI versus row III).

Appendix H   Sex effects in patient neuroimaging data

For the patient neuroimaging data, we evaluated sex effects in the post-CCA sources from MSIVA S_2 cross-modal subspaces (Figure H.1). The balanced accuracy for sex prediction was approximately 50% for each source. Moreover, no significant sex effects were observed for any source ($p^{[1]} > 0.05, p^{[2]} > 0.05$ for all sources; two-sample t-test with FDR correction for all subjects).

Figure H.1: **Patient neuroimaging data: Post-CCA sources from MSIVA S_2 cross-modal subspaces, color coded by sex.** The Pearson correlation coefficient (r) shows the correlation of post-CCA sources between two modalities. The p-value shows statistical differences between two sex groups for sMRI and fMRI, respectively ($p^{[1]}$: sMRI; $p^{[2]}$: fMRI; two-sample t-test, FDR correction for all subjects). Sex effects were not significant in the patient neuroimaging dataset.

Appendix I   MSIVA S_2 reconstructed neuroimaging data

Figure I.1 shows spatial maps of group-specific reconstructed data from each of five MSIVA S_2 linked subspaces in the UKB dataset. Similarly, Figure I.2 shows results related to the age and SZ interaction effects in the patient dataset. In each panel, axial slices show the geometric median of the reconstructed subspace data $(\hat{\mathbf{X}}_k^{[m]})$ for each modality and each group. Voxel intensity is mapped to both color hue and opacity. Contours highlight brain areas with top 15% of voxelwise cross-modal correlations for each group (negative correlations: black; positive correlations: magenta). Histograms show voxelwise cross-modal correlations for each group (colored dashed lines: top 15% of negative or positive correlations for each group; black dotted lines: $p = \frac{0.01}{44318}$, Bonferroni correction for 44318 voxels). The reported R^2 indicates the proportion of variance captured by the subspace in each modality.

Figure I.3 illustrates the number of voxels that show significant cross-modal correlations for age and sex groups in the UKB dataset (rows I and II), and for age and diagnosis groups in the patient dataset (rows III and IV). The number of voxels for older patients diagnosed with SZ is consistently less than that for their age-matched control subjects in four of five subspaces, implying reduced brain structure-function coupling in the older patient group.

Subspace 1 ($\hat{S}_1^{[1]}$ vs $\hat{S}_1^{[2]}$)

Subspace 2 ($\hat{S}_2^{[1]}$ vs $\hat{S}_2^{[2]}$)

Subspace 3 ($\hat{S}_3^{[1]}$ vs $\hat{S}_3^{[2]}$)

(a) Subspaces 1-3.

(b) Subspaces 4-5.

Figure I.1: **UKB neuroimaging data: Spatial maps of group-specific reconstructed data from MSIVA S_2 sources related to age and sex effects.** Axial slices show the geometric median of the reconstructed data $(\hat{\mathbf{X}}_k^{[m]})$ for each modality (sMRI or fMRI) and each group (younger: $46-62$ years, older: $63-79$ years; male or female). Voxel intensity is mapped to both color hue and opacity. Contours highlight brain areas with top 15% of voxelwise cross-modal correlations for each group (negative correlations: black; positive correlations: magenta). Histograms show voxelwise cross-modal correlations for each group (colored dashed lines: top 15% of negative or positive correlations for each group; black dotted lines: $p = \frac{0.01}{44318}$, Bonferroni correction for 44318 voxels). The reported R^2 indicates the proportion of variance captured by the subspace in each modality.

Subspace 1 ($\hat{S}_1^{[1]}$ vs $\hat{S}_1^{[2]}$)

Subspace 2 ($\hat{S}_2^{[1]}$ vs $\hat{S}_2^{[2]}$)

Subspace 3 ($\hat{S}_3^{[1]}$ vs $\hat{S}_3^{[2]}$)

(a) Subspaces 1-3.

(b) Subspaces 4-5.

Figure I.2: **Patient neuroimaging data: Spatial maps of group-specific reconstructed data from MSIVA S_2 sources related to age and SZ interaction effects.** Axial slices show the geometric median of the reconstructed data $(\hat{\mathbf{X}}_k^{[m]})$ for each modality (sMRI or fMRI) and each group (younger: $15 - 38$ years, older: $39 - 65$ years; control or patient). Voxel intensity is mapped to both color hue and opacity. Contours highlight brain areas with top 15% of voxelwise cross-modal correlations for each group (negative correlations: black; positive correlations: magenta). Histograms show voxelwise cross-modal correlations for each group (colored dashed lines: top 15% of negative or positive correlations for each group; black dotted lines: $p = \frac{0.01}{44318}$, Bonferroni correction for 44318 voxels). The reported R^2 indicates the proportion of variance captured by the subspace in each modality.

Figure I.3: **Number of voxels that show significant cross-modal correlations for age and sex groups in the UKB dataset (rows I and II), and for age and diagnosis groups in the patient dataset (rows III and IV).** Rows I and III display the number of voxels with both positive and negative correlations $(+/-)$, while rows II and IV display the number of voxels with only positive correlations $(+)$. The number of voxels for older patients diagnosed with SZ is consistently less than that for their age-matched controls in four of five subspaces, implying reduced brain structure-function coupling in older patients.

Appendix J   Comparison of MSIVA S_2 and MMIVA sources

J.1   Experiments and evaluation metrics

MSIVA can be viewed as an extension of MMIVA with two main differences. First, MSIVA uses a flexible *block* diagonal subspace structure, while MMIVA uses a rigid identity matrix as the subspace structure. Second, MSIVA uses MGPCA and separate ICAs initialization, while MMIVA uses MGPCA and group ICA initialization (multimodal initialization). To further investigate similarities and differences of the recovered sources from MSIVA and MMIVA, we compared MSIVA with the subspace structure S_2 and MMIVA with the subspace structure S_5 through the following experiments:

1. We performed multiple linear regression (MLR) for each modality using MSIVA S_2 post-CCA sources from each cross-modal subspace $\mathbf{X}_i^{[m]}$ to predict each MMIVA source $\mathbf{y}_j^{[m]}$:

$$\mathbf{y}_j^{[m]} = \mathbf{X}_i^{[m]}\boldsymbol{\beta}, \tag{21}$$

   where $i \in \{1, \ldots, 5\}$ is the cross-modal subspace index in MSIVA S_2, and $j \in \{1, \ldots, 12\}$ is the subspace index in MMIVA.

2. We performed multivariate analysis of variance (MANOVA) for each modality using a pair of matched MMIVA sources $\left[\mathbf{y}_j^{[m]}, \mathbf{y}_k^{[m]}\right]$ from Step 1 to predict MSIVA S_2 post-CCA sources from each cross-modal subspace $\mathbf{X}_i^{[m]}$:

$$\mathbf{X}_i^{[m]} = \left[\mathbf{y}_j^{[m]}, \mathbf{y}_k^{[m]}\right]\boldsymbol{\beta}. \tag{22}$$

   Here, (j, k) are a pair of matched subspace indices in MMIVA, and i is the cross-modal subspace index in MSIVA S_2.

3. We performed brain-phenotype prediction using MSIVA S_2 post-CCA sources from each cross-modal subspace, as well as each pair of matched linked 1D MMIVA sources identified in Step 1. As described in Section 2.5, we performed age prediction and sex classification tasks for the UKB dataset, as well as age prediction and binary diagnosis classification tasks (controls versus patients with SZ) for the patient dataset. We measured mean absolute error (MAE) for age regression and balanced accuracy for binary classification.

We measured the adjusted R^2 (R^2_{adj}) from MLR as shown below:

$$R^2 = 1 - \frac{\sum_{i=1}^{N}(y_i - \hat{y}_i)^2}{\sum_{i=1}^{N}(y_i - \bar{y})^2}, \quad \bar{y} = \frac{1}{N}\sum_{i=1}^{N}y_i, \tag{23}$$

$$R^2_{adj} = 1 - (1 - R^2)\frac{N-1}{N - N_P - 1}, \tag{24}$$

where N is the number of samples (here subjects) and N_P is the number of predictors.

J.2   Results

Figures J.1 and J.3 show the adjusted R^2 when using pairs of MSIVA S_2 sources from each cross-modal subspace to predict each of the 12 MMIVA sources for the UKB dataset and the patient dataset, respectively. We reordered MMIVA sources to identify the most likely correspondence between MSIVA S_2 sources and MMIVA sources. We notice that there exists some correspondence between MSIVA S_2 sources and MMIVA sources. For example, for UKB sMRI data, MSIVA S_2 subspace 3 sources match MMIVA source 3 ($R^2_{adj} = 0.96$), MSIVA S_2 subspace 5 sources match MMIVA source 5 ($R^2_{adj} = 0.87$), and MSIVA S_2 subspace 4 sources match MMIVA source 2 ($R^2_{adj} = 0.73$). We also observe that more than two columns show high R^2_{adj} (> 0.2) for each row, indicating that every two MSIVA S_2 sources from each subspace can predict variability for more than two MMIVA sources. Note that the prediction results for fMRI are very consistent with those for sMRI.

Next, we performed MANOVA using every two matched MMIVA sources to predict the pair of MSIVA S_2 sources from each cross-modal subspace. Figures J.2 and J.4 illustrate the Pillai's trace divided by the number of modalities ($M = 2$). Note that the maximum possible diagonal Pillai's trace value is 2 for two modalities, and dividing the Pillai's trace by 2 shows the variance explained for each modality. We observe large off-diagonal values per column, indicating that every pair of matched MMIVA sources predicts variability of more than two MSIVA S_2 sources. Note that the prediction results for fMRI are very consistent with those for sMRI.

Finally, we performed phenotype prediction using MSIVA S_2 post-CCA sources from each cross-modal subspace, as well as each pair of matched MMIVA sources. Table J.1 lists the detailed prediction performance. For the UKB dataset, MSIVA sources from subspaces 5 and 4 achieved the best age regression MAE and sex classification balanced accuracy, respectively. For the patient dataset, MMIVA matched sources from subspace 2 yielded the best age regression performance, while MSIVA sources from subspaces 5 showed the best diagnosis classification performance. Overall, the linked sources from

MSIVA S_2 showed stronger associations with age, sex, and SZ-related effects than the paired independent sources from MMIVA.

Therefore, we conclude that MSIVA and MMIVA apportion variability to their sources in different ways. There is no perfect one-to-one mapping between MSIVA S_2 sources and MMIVA sources. We also note that the mismatch appears to be more pronounced in the patient dataset than in the UKB dataset, which may be related to inherent characteristics of the patient data, such as higher population heterogeneity and smaller sample size. MSIVA linked sources from a cross-modal subspace more accurately predicted phenotype measures (age and sex in the UKB dataset and diagnosis in the patient dataset) than independent pairs of matched linked 1D MMIVA sources, indicating that higher-dimensional subspaces in MSIVA better capture phenotype-related variability than one-dimensional sources in MMIVA.

Figure J.1: **UKB neuroimaging data: Adjusted R^2 using MSIVA sources to predict MMIVA sources.**

Figure J.2: **UKB neuroimaging data: Pillai's trace value using matched MMIVA sources to predict MSIVA sources.**

Figure J.3: **Patient neuroimaging data: Adjusted** R^2 **using MSIVA sources to predict MMIVA sources.**

Figure J.4: **Patient neuroimaging data: Pillai's trace value using matched MMIVA sources to predict MSIVA sources.**

Table J.1: **Phenotype prediction performance using post-CCA sources from MSIVA subspace structure S_2 and matched sources from MMIVA.** For the UKB dataset, MSIVA sources from subspaces 5 and 4 achieved the best age regression and sex classification performance, respectively. For the patient dataset, MMIVA matched sources from subspace 2 yielded the best age regression performance, while MSIVA sources from subspaces 5 showed the best diagnosis classification performance. Overall, the linked sources obtained by MSIVA S_2 showed stronger associations with age, sex, and SZ-related effects than the paired independent sources from MMIVA. Note that we estimated sources for the UKB data and the patient data separately, thus subspaces in the UKB dataset do not correspond to those in the patient dataset.

Subspace	1	2	3	4	5
UK Biobank dataset					
MSIVA age mean absolute error (years)	5.674	6.163	5.892	5.847	**5.378**
MMIVA age mean absolute error (years)	6.020	5.541	6.100	6.054	5.470
MSIVA sex balanced accuracy (%)	59.542	64.496	59.206	**79.933**	52.699
MMIVA sex balanced accuracy (%)	58.874	63.729	66.483	60.085	75.415
Patient dataset					
MSIVA age mean absolute error (years)	10.720	10.470	11.226	11.445	10.307
MMIVA age mean absolute error (years)	10.959	**9.822**	10.903	11.679	10.835
MSIVA diagnosis balanced accuracy (%)	50.565	57.624	50.000	49.691	**61.404**
MMIVA diagnosis balanced accuracy (%)	50.752	57.747	59.919	50.000	57.695