Young Eurasian Scops Owls make individually distinctive vocalizations

Fabrizio Grieco

Wageningen, The Netherlands; email fabriziogrieco1@gmail.com

Abstract. Vocal distinctiveness is expected to occur more often in colonial-breeding species as parents need to recognize their offspring in a large group of conspecifics. In this study, I examined the calls of the young of a territorial nocturnal raptor, the Scops Owl both before and after fledging. Contrary to what is expected for a territorial species, the calls of young owlets showed a high degree of repeatability in many acoustic features, particularly call duration, spectral centre of gravity, and spectral skewness. Discriminant analysis assigned 52.5% of the calls to the correct individual, and 77.5% when restricting analysis within broods. Inter-individual differences appear before fledging and were stable for multiple days. The results suggest that the vocalizations of young Scops Owls could potentially act as identity signals. However, the functional significance of acoustic signatures remains unclear; a few hypotheses are discussed. Acoustic distinctiveness could merely reflect the development of vocal individuality that is typical of the adults, with no net benefit to parents and offspring. However, it could also play a role in parent-offspring communication, preventing misdirected parental care to unrelated young, or helping the parents feed preferentially specific young, as was suggested earlier for this species. Acoustic signatures could also help brood mates maintain contact after fledging or mediate competition for food between them. Further studies are needed to determine whether parents and offspring do assess identity signals in vocalizations.

Keywords: Acoustic signature, food-begging, fledglings, individuality, nestlings, Otus scops, parent-offspring communication, repeatability.

Statements and Declarations

Funding and competing interests. The author did not receive support from any organization for the submitted work. The author has no relevant financial or non-financial interests to disclose.

Data availability. Data supporting the results are available from the author upon request.

Ethical approval. The observations reported in this work comply with the current laws of the country where they were carried out. No ethical approval was required.

INTRODUCTION

The ability to discriminate among conspecifics is an important characteristic in many animals and is often required for mate and kin recognition, and parent-offspring communication (Tibbetts & Dale, 2007). Studies of bird vocalizations have found a plethora of apparent adaptations to increase the distinctiveness of individuals (Falls, 1982; Lambrechts & Dhondt, 1995; Carlson et al., 2020) in contexts such as mate and kin recognition (e.g. Sharp et al., 2005; Vignal et al., 2008), neighbour-stranger discrimination (Stoddard, 1996), and parent-offspring communication ((Beer, 1971; Tibbetts & Dale, 2007). The latter case occurs often in bird species that breed in dense and noisy colonies, like penguins and seabirds (Buckley & Buckley, 1972; Lefevre et al., 1998; Jouventin & Aubin, 2002). In colonial swallow species, nestlings make individually distinctive calls that parents recognize (Beecher et al., 1981; Stoddard & Beecher, 1983; Loesche et al., 1991; Strickler, 2013). On the contrary, calls of nestlings of non-colonial species show weak or absent individuality (Medvin & Beecher, 1986; Medvin et al., 1992, 1993; Leonard et al., 1997 but see Barg & Mumme, 1994). The observation that parent-offspring recognition is better developed in colonial than in noncolonial species led to the hypothesis that coloniality is a strong selection pressure for the evolution and maintenance of the distinctiveness of calls (Beecher, 1990). In colonially-breeding species, the parents must identify their own young among many unrelated ones, particularly after their young leave the nest and become mobile, so that their location is an unreliable indicator of identity. In those cases, the risk of misdirected parental care is high.

Individual variation in vocalizations is a prerequisite for the ability to discriminate between conspecifics (Falls, 1982). Features of the signal that is sent out, like its dominant frequency, or duration, must show weak intra-individual variability combined with high inter-individual variability. However, vocal traits often vary along with physical or behavioural conditions of the individual. To be a reliable indicator of identity, a signal must also be temporally stable, that is, it must remain unaltered along significant time scales (Gilbert & McGregor, 1994; Peake et al., 1998; Lengagne, 2001; Grava et al., 2008; Klenova et al.,

2009, 2012). Among *Strigiformes*, vocal signatures have often been found in adult birds (e.g. Eagle owls:

Lengagne, 2001; Great horned owls, Odom et al., 2013; Barred owls, Freeman, 2000; Tawny owls,

Galeotti & Pavan, 1991; Pigmy owl: Galeotti et al., 1993; Western screech-owls, Tripp & Otter, 2006;

Ural owl, Zhou et al., 2020; Northern spotted owl, Fitton, 1991; Eastern screech owl Cavanagh &

Ritchison, 1987), but less often in nestlings and fledglings (Dreiss et al. 2014).

The Eurasian Scops Owl (*Otus scops*, hereafter 'Scops Owl') is a small (body length 19-20 cm),

nocturnal, primarily insectivorous owl, that breeds in open habitats in the Mediterranean zone and

Central-Eastern Europe (Mikkola, 1983; Cramp, 1985). The species is mainly territorial, although it

sometimes forms "loose colonies" of breeding pairs with inter-nest distances as short as 15 m (Grieco

2018; for an overview see Cramp 1985). The advertising call of the adult is highly repeatable (Koenig,

1973; Dragonetti, 2007) and can be used to track individuals unobtrusively (Galeotti & Sacchi, 2001;

Denac & Trilar, 2006; Grieco, 2018). The young owlets fledge at about 21 days from hatching and stay

together while they are fed by their parents up to independence. The young owlets produce short (0.15 s)

"tschp" calls at intervals of 0.5-1 second, interpreted as food solicitation calls, and thought to gradually

evolve into the adult advertising call after independence (Koenig 1973). However, there is no quantitative

data about their inter-individual variability and stability over time.

Because the species is predominantly territorial, and the fledged brood mates do not mix with other,

unrelated fledglings (unpubl. data), the parents likely do not need to recognize their offspring

individually. Therefore, the expectation is that the degree of individuality in the vocalizations of the

young during the period of dependence on the parents would be low to moderate.

This observational study had two aims: (1) To provide the first detailed description of the food-begging

calls of the young Scops Owls and their inter-individual variation based on spectrographic analysis; (2)

To determine whether individuals produce calls that are consistently similar in their structure. For this

purpose, I recorded the vocalizations of young owlets both before and after nest departure. Especially

information about post-fledging behavior is important, since most of the studies of vocalizations and

begging behaviour of young birds have focused on nestlings, while the post-fledging phase is

underrepresented (Draganoiu et al., 2006; Middleton et al., 2007), particularly for nocturnal bird species

(Kouba et al., 2014).

METHODS

Study area and species

The study was carried out at the State Reserve Metaponto, Italy (40°22'24"N, 16°50'19"E), a reforested

area dominated by Aleppo pine *Pinus halepensis* along the sea coast. Forest plots are mixed with human

settlements, especially recreational resorts and camping parks. The Scops Owl is a primarily cavity-

nesting species but because tree cavities are virtually absent in this area, breeding pairs occupy old twig

nests, particularly those of Magpies (*Pica pica*). The young leave their nest about 21 days after hatching

(Koenig, 1973), and on the following nights, they move further and further away from the nest. The

parents keep feeding the young regularly up to 40 days after hatching, then increasingly less often. At 45

days the young are fully able to catch prey (Koenig 1973), and become gradually independent and

disperse. Up to that time, the owlets are inseparable but they do not come in contact with other broods

(pers. obs.).

Sound recording and call selection

I recorded vocalizations of 20 Scops Owl nestlings and fledged young from six broods between July and

August 2021 and July 2022. Recording sessions were carried out between 19:30 and 21:30 and between

00:00 to 04:30 and were made opportunistically. The study area was actively searched for nests and

fledged broods. To find active nests, Magpie twig nests were watched in the evening and in the morning

for at least half an hour to detect adult owls bringing food (see also Grieco 2018). Fledged young were

found by listening to their calls at a distance. To record sound, I used a Telinga Pro 8 MK2 stereo

microphone mounted on a parabolic dish with a diameter of 52 cm, and a Marantz PMD 661 MK2 digital

sound recorder. Each recording lasted at least one minute and was stored in a WAV file at a sampling rate

of 44.1 kHz, and with 16-bit depth. In most cases, recordings were made a few meters from the nest tree

or the fledged young.

Among all the recordings available, I selected those that contained the calls of all the individuals of the

brood that could be measured: brood 1-3 (3 individuals, 1 recording each); brood 4 (4 individuals, 2

recordings); brood 5 (4 individuals, 5 recordings), and brood 6 (3 individuals of which one disappeared;

13 recordings in total). An additional recording was selected for broods 5 and 6 to illustrate the changes in

the calls' structure during the visits by the adults. The total length of the audio analyzed was 30996 s (8 h

36 min 36 s). Recordings lasted on average 1291 ± 812 (SD) seconds (n = 24).

Because hatching was not observed directly, the exact age of the nestlings was unknown. For four of the

six broods, the fledging date was known, and the first recording was made 7 days before fledging (brood

6) to 10 days after fledging (brood 2). For the other two broods with an unknown fledging date, their

behavior suggests that they were recorded at least one week after fledging.

For each of the 20 young owlets and each recording, I selected the 10 loudest calls that were free of

background noise and did not overlap calls of their mates or adults. This resulted in the main data set of

490 calls. It is important to state that calls were assigned based on the visual appearance of their

spectrogram, that is, two calls with similar spectrograms were assumed to be from the same individual.

The entire work is based on this assumption. Because the owlets were not individually marked, the

assumption could have produced overestimates of distinctiveness. However, when the young owls were

clearly separated from their brood mates – and consequently the position of the caller could not be

mistaken – the spectrograms of the calls given by a specific individual were of the same type, and that

consistency could be verified for as long as half an hour of continuous calling (e.g. Figure 4 in Results).

In a second analysis, I tested whether vocal differences between brood mates were stable over days. For

this purpose, I selected recordings of brood 6 that covered the longest period of time, i.e., 10 days (July 20

to July 29, 2022). This second data set only includes the calls of two young out of three, because one

disappeared shortly after leaving the nest (July 22).

Sound analysis

Sound analysis was carried out in Audacity 2.1 (https://www.audacityteam.org), Praat 6.0

(https://www.fon.hum.uva.nl/praat/), and Sound Analysis Pro (http://soundanalysispro.com/) depending

on the acoustic features examined (see below). Recordings were first filtered with a 1000 high-pass filter

in Praat 6.0 to remove low-frequency background noise. The following FFT settings were used: FFT

window size 2048 samples, Hanning window, temporal resolution 46 ms, frequency resolution 21.5 Hz.

Two temporal features and eight spectral features were measured:

1. Call duration (s). The time from the onset to the end of the call, measured from the waveform in

Praat.

2. Time from the previous call (s). The interval between the onset time of the previous call and that

of the current call, measured from the waveform in Praat.

3. Peak frequency(Hz). The frequency at maximum amplitude. To measure it, the entire call

duration was selected in Audacity and the function Plot Spectrum was used with the following

settings: algorithm Spectrum, function Hanning window, window size 2048.

4. Peak frequency 1/3 (Hz). The frequency at maximum amplitude within the first third of the call

duration, measured the same way as Peak frequency.

5. Peak frequency 3/3 (Hz). The frequency at maximum amplitude within the last third of the call

duration, measured the same way as Peak frequency. Peak frequency 1/3 ad Peak Frequency 3/3

were chosen because the initial and the ending parts of the calls were often very different -

whereas the initial part is generally noisy, the ending part is more structured.

6. Centre of gravity (Hz). The average frequency over the entire frequency domain, weighted by the

sound amplitude. It was measured in Praat by selecting the call waveform and calling the function

Spectrum > View spectral slice, then Query > Get centre of gravity, with power setting 2.

7. Pitch (Hz). The average pitch across the call duration. It was measured in Praat by selecting the

waveform and calling the function Pitch > Get Pitch, with the following settings: frequency range

1000-10000, Analysis method Autocorrelation, Silence threshold 0.03, Voicing threshold 0.1.

The intended use of Pitch was to describe the perceived tone of a sound.

8. Average entropy. The average Wiener entropy across the call duration, measured with Sound

Analysis Pro. Entropy measures the extent to which the sound contains a mixture of frequencies,

as opposed to a single frequency or tone. The noisier a sound is, the more energy is smeared

within a wide range. Entropy is a pure number and varies between 0 to 1, but in this work it is

reported on a logarithmic scale, ranging between minus infinite (pure tones, complete order) and

zero (white noise, complete disorder).

9. Minimum entropy. The minimum Wiener entropy within the call duration (see Average Entropy

above for details). Minimum Entropy aims at capturing the more structured sound, usually at the

end of the call (see an example in Online Resource S1).

10. Spectrum skewness. A measure of the symmetry of the spectrum, that is. It was measured in Praat

by selecting the waveform and calling the function Spectrum > View spectral slice, then Query >

Get skewness, with power setting 2. Skewness is a pure number; it is negative when frequencies

are concentrated on the right of the spectrum, and positive when the frequencies are concentrated

on the left of the spectrum. If skewness is zero, the spectrum is perfectly symmetrical.

Statistical analysis

For each of the acoustic features defined above, I tested whether the 10 measurements taken for each

individual were more similar to each other than to those of the other brood mates. Data were first tested

for normality. For each recording, I considered the residuals of the measurements after fitting the

individual means (i.e., each measurement diminished by the mean within the recording). Kolmogorov-

Smirnov tests were applied to the sets of measures in each recording (10 acoustic features and 23

recordings). After Bonferroni correction for multiple testing, where alpha was reduced by dividing 0.05

by the number of recordings, a few features showed signs of non-normality. The null hypothesis of

normality was rejected in five out of 23 cases for Peak frequency and Peak Frequency 1/3 and in two out

of 23 for Peak Frequency 3/3. Overall, the effects of non-normality should not be fatal for the ANOVA

but the results of Peak Frequency and Peak Frequency 1/3 should be taken with caution.

To assess the repeatability of acoustic features, I performed a one-factor ANOVA for each recording and

calculated the Repeatability (R) using the formula by Lessells & Boag (1987). In the main data set, each

brood resulted in one to five values of R for each acoustic feature, depending on how many recordings

were available for that brood. Furthermore, 95% Confidence intervals were calculated from the individual

R values from the six broods using the method described by Nakagawa & Schielzeth (2010). Similarly,

the second data set provided multiple values of R for each acoustic feature. For this data set, which

should be considered illustrative as it was based on one brood (brood 6), the confidence intervals were not

calculated.

Another way to look at individual differences in vocalizations is to use linear discriminant analysis (LDA)

to test whether a specific call could be attributed to an individual based on its acoustic features. To keep a

balanced design with an equal number of calls per individual, I selected one recording (10 calls) for each

of the 20 owlets. Among the 10 acoustic features, seven were entered as explanatory variables. The other

three features (Time from the previous call, Peak frequency, and Average entropy) were excluded because

the former showed low inter-individual variation according to the Repeatability analysis (see Results) and

the other two were highly correlated with other features (Peak frequency correlated with Peak frequency

1/3 and Average entropy correlated with Minimum entropy). LDA was performed in XLSTAT for Excel

(Lumivero, 2023). Classification accuracy was obtained with leave-one-out cross-validation, that is, each

of the 200 calls was classified using a discriminant function computed from the remaining 199 calls.

RESULTS

The calls of young Scops Owls

Food calls of young Scops Owls showed marked variation in all the temporal and spectral features

measured (Table 1). First, some calls exhibited a wide spectrum, with peak frequency around 4000-5000

kHz, while in others the spectrum was more asymmetric, with the loudest sound around 2000 Hz (Figure

1A). Second, the calls of the same form were also consistent in their duration and occurred sometimes

with a consistent rhythm (Figure 1A, B). Third, fine-scale spectral structure, in the form of spots or

scribbles in the spectrograms, could be easily detected through visual inspection (Figure 1C). Finally,

calls that looked similar also did sound different to the human ear (Online Resource S6). Figure 2 shows a

summary of the spectrogram types for each of the six broods examined.

Covariation between vocal features

Young owls that gave high-frequency calls also produced less skewed calls, that is, with a broader

spectrum and less clear frequency peaks (Table 2 and Online Resource 2). The negative relationship

between frequency and spectrum skewness was also apparent when considering individuals within broods

(Online Resource 3), so it was not the result of fundamental differences in the spectral structure between

broods. Frequency measurements co-varied in general with each other, as expected. However, Peak

Frequency 3/3, which measured the dominant frequency at the end of the call, did not correlate

significantly with any other measure. Call duration and time from the previous call did not co-vary with

spectral measures, although longer calls tended to be low-frequency and more skewed (Online Resource

2).

Distinctiveness of calls

Repeatability estimates within broods were significantly greater than zero for all the measured features

(Table 3). Call duration showed the highest repeatability with the lowest uncertainty of the estimate,

suggesting that it was the best indicator of individual distinctiveness. Frequency features and spectral

skewness showed moderate to high repeatability, while average and minimum entropy were somewhat

less repeatable. On the contrary, the time from the previous call, a measure of the calling rhythm, showed

the lowest repeatability.

Discriminant analysis confirmed a substantial level of individuality in the calls. On average, 52.5% of the

200 calls were correctly assigned to an individual owl. Calls of some individuals were more distinctive

than those of others, with individual-based classification accuracy ranging from 30% to 90% ($n = 20$).

When LDA was repeated within broods, classification accuracy increased to 77.5%, with little difference

between individuals in broods of three siblings and those in broods of four siblings (79.2% vs. 75.0%;

range 50% to 100% in both cases).

Effect of parental visits on the call structure

Young owls in the post-fledging phase vocalized continually for a long time, also in the absence of the

parents. Temporal changes in the acoustic structure occurred around the visits of the parents (Figure 3).

At the arrival of the parent, the young generally tended to get closer to it, while their calls became more

frequent and shorter. During the actual feeding event, which occurred a few seconds later, the calls

changed most profoundly: frequency measures and entropy tended to increase, while the spectral

skewness decreased especially for the call types with a highly-skewed spectrum (Figure 3, bottom-right

chart). In short, the young gave shorter, noisier, and less distinctive calls (see for example call duration in

Figure 3 and S4). However, the spectral centre of gravity still differed between individuals in two broods

(Figure 3 and S4).

Stability of between-siblings differences

Analysis of a separate dataset of vocalizations of brood mates recorded over multiple days revealed that

individual differences in acoustic features emerged well before fledging. Peculiarities in the spectrograms

appeared seven days before fledging (Online Resource 5), but a proper quantitative comparison between

calls could not be done because of the low sound amplitude. Two days before fledging, and in the

following 10 days, almost all call features exhibited significant inter-individual differences (Figure 4).

The results were generally in line with the patterns found in the large data set with multiple broods. Call

duration was the acoustic feature that showed consistently the highest repeatability (Figure 4, top-right).

Inter-individual differences were also found in most of the recordings, particularly for minimum entropy,

spectral skewness, centre of gravity, and peak frequency 3/3 (Figure 4). Therefore, inter-individual

variability could be potentially used to discriminate between brood mates for several days, also after

fledging.

DISCUSSION

The results of this study may be summarized as follows: (1) The main call of young Scops Owls is more complex than previously described (Koenig, 1973) as it exhibited larger variability (for example, in its duration) and marked spectral complexity. (2) Significant interindividual differences were found in almost all temporal and spectral features measured. (3) Individuality of vocal characteristics appeared before fledging and could be traced for ten days. (4) Calls were less distinctive when the young interacted with the provisioning parents, but the latter could have enough time to perceive differences immediately before the actual feeding event. Taken together, these findings suggest that vocalizations of young Scops allow not only discrimination, that is, telling whether two calls are from different individuals, but also identification, that is, labeling individuals, at least within broods.

Acoustic distinctiveness in parent-offspring communication

The results of this study are in contrast with what one would expect if vocal signatures develop primarily to make the offspring recognizable in a large group of fledglings, as it occurs in colonial-breeding species. The Scops Owl is to a large extent territorial (Cramp, 1985), and fledged brood mates tend to stick together and do not intermingle with other, unrelated ones. Therefore, it is unlikely that vocal signatures in fledglings serve the function to prevent misdirected parental care. In the study population, however, breeding density is unusually high: nests can be as close as 12 m from one another and males defend partially overlapping territories (Grieco, 2018, 2022). Fledgled broods have sometimes been observed moving outside their natal territory, even entering other males' territories (pers. obs.), which suggests that unrelated fledglings may sometimes be at a short distance from each other. A semi-colonial organization of breeding pairs is poorly known but probably occurs frequently in the South-European distribution

range of the species where local breeding density can be high. Some degree of intermingling of young

could provide some selection pressure for vocal distinctiveness in this species. A similar mechanism has

been suggested by Barg and Mumme (1994) to explain the apparent contradictory finding that in the

Florida Scrub Jays, a highly territorial species with some limited intermingling of young after fledging,

the parents can discriminate between offspring and nonoffspring (Barg and Mumme, 1994).

There are a few possible alternative explanations for the occurrence of acoustic signatures in young Scops

Owls. First, the acoustic structure in young birds' calls could merely reflect the development of

individuality at later stages. In several species, including the Scops Owl (Koenig 1973), begging calls

gradually transition to adult calls (Redondo & Exposito, 1990; Brittan-Powell et al., 1997; Marques et al.,

2011). Young owlets might not "broadcast" their individuality, and their voice would simply reflect their

individuality in adulthood, with no net advantage early in life for parents and offspring. Second, parents

could use acoustic signatures to discriminate between their own nestlings and alien (parasitic) nestlings,

as found in other species (Colombelli-Négrel et al., 2012; Yang et al., 2015). However, Scops Owls are

not subject to any known brood parasites (Mikkola, 1983; Cramp, 1985), and laying eggs by multiple

females in the same nest is a rare event (Bavoux et al., 1991). Therefore, the acoustic distinctiveness of

nestlings is unlikely to be of adaptive value in the context of the rejection of unrelated nestlings.

There is another, intriguing scenario where parents may be using acoustic signatures of their offspring to

feed preferentially some individuals over others, as suggested earlier for this species (Parejo et al., 2010).

Theory (Lessells, 2002) and an increasing number of empirical studies suggest that in some circumstances

the parents do not distribute food equally among nestlings, based on nestling size, body condition, sex, or

age (Krebs, 2001; Smiseth et al., 2003; de Ayala et al., 2007; Dickens & Hartley, 2007; Dugas, 2009;

Avilés et al., 2011; Ersoy et al., 2021). For example, females and male parents differ in the amount of

care given to the smallest nestlings (Slagsvold, 1997), although it is not always clear whether nestling size

is the primary cue used in this discrimination. Condition-dependent parental favouritism mediated by

visual cues has been suggested to occur in several species (Jourdie et al., 2004; Bize et al., 2006; Galván

et al., 2008; Avilés et al., 2011), including nocturnal raptors (Avilés & Parejo, 2013). In an experimental

study on Scops Owls, Parejo et al. (2010) found that when the ultraviolet reflectance of the cere (i.e., the

skin above the beak) of the nestlings was experimentally reduced, thereby simulating lower body

condition, those nestlings gained more weight than their control siblings. The authors concluded that the

parents somehow perceived the lower brightness of the cere and fed preferentially those nestlings over the

untreated siblings. The study suggested that parent Scops Owls do attend to cues of quality of nestlings.

Discriminating between offspring by sight only is efficient in the nestling-rearing phase, where both

parent and offspring are confined in the small environment of the nest. However, when the offspring leave

the nest and become mobile, a recognition system based on visual cues of limited size like the beak cere

could could be aided by acoustic distinctiveness (Draganoiu et al., 2006). The parents could learn to

associate cues of quality in the offspring, like their body size, sex, or hatching order, with the vocal

signatures learnt during the nestling-rearing phase so that they could adjust their food-provisioning

decisions. With an important limitation though - this mechanism could only work if the cues of quality

remain stable in the short/medium term (i.e. over days to weeks) so that they can be associated with an

individual's voice.

Why vocalize in the absence of parents?

In contrast with what occurs in many bird species, the young Scops Owls spent significant time calling

when the parents were absent (Figure 3, Figure S4). This behavior has often been described as "parent-

absent begging" (Budden & Wright, 2001) or "secondary begging" (Leonard & Horn, 2001).

Vocalizations in the absence of parents are known in other owl species, both during the nestling (Roulin

et al., 2000, 2009) and in the post-fledging stage (Olsen et al., 2020). Prolonged vocalizing is presumably

costly, in terms of energy requirements (Kilner & Johnstone, 1997), predation risk (Magrath et al., 2010),

and vigilance costs (Roulin, 2001). Because any begging signal that does not elicit any parental responses

is wasteful, one may wonder what other benefits of this behavior could be. First, because the fledglings

are mobile, by continually vocalizing they could inform the parents about their location and need. A

second possibility is that siblings may need to recognize each other to keep contact after they leave the

nest (Ligout et al., 2016). In this scenario, vocal distinctiveness would play a role in sib-sib

communication. Furthermore, repeated calling may be a "quick on-trigger" strategy to attract the attention

of the parents. This behavior presumably results in many false alarms (that is, soliciting a parent that is

absent at that time) but at the same time reduces costly misses if the parent is in the vicinity (Dor et al.,

2007). The high cost of misses could be particularly important after fledging when each individual does

not know whether the parent will arrive next to it or its siblings. Finally, vocal individuality could be

advantageous in another, interesting scenario of sibling competition. Through repeated calling, the

siblings could be displaying their willingness to compete for the next food item delivered by the parents -

a form of negotiation known in another nocturnal raptor, the Barn Owl (Roulin et al., 2000; Roulin,

2002). Each individual could recognize which mates are more motivated to compete, and thereby adjust

its competitive behavior accordingly (Roulin et al., 2009; Dreiss et al., 2014).

Concluding, this work has revealed that vocalizations of juvenile Scops Owls potentially carry individual

information, at least within broods. This should encourage researchers to address the many unanswered

questions outlined above and gain insights into the relationships between nesting ecology,

social/territorial organization, and communication in this still poorly known species.

ACKNOWLEDGMENTS

REFERENCES

Avilés, J. M., & Parejo, D. (2013). Colour also matters for nocturnal birds: Owlet bill coloration advertises quality and influences parental feeding behaviour in little owls. Oecologia 173 399–408. https://doi.org/10.1007/s00442-013-2625-8

Avilés, J. M., Parejo, D., & Rodríguez, J. (2011). Parental favouritism strategies in the asynchronously hatching European Roller (Coracias garrulus). Behav. Ecol. Sociobiol. 65 1549–1557. https://doi.org/10.1007/s00265-011-1164-8

Barg, J. J., & Mumme, R. L. (1994). Parental recognition of juvenile begging calls in the Florida scrub jay. Auk 111 459–464. https://doi.org/10.2307/4088610

Bavoux, C., Burneleau, G., & Nicolau-Guillaumet, P. (1991). Aspects de la biologie de reproduction du hibou petit-duc Otus scops. Alauda 59 65–71.

Beecher, M. D. (1990). The evolution of parent-offspring recognition in swallows. In D. A. Dewsbury (Ed.), *Comparative Issues in Comparative Psychology* (pp. 360–380). Sinauer Associates.

Beecher, M. D., Beecher, I. M., & Hahn, S. (1981). Parent-offspring recognition in bank swallows (Riparia riparia): II. Development and acoustic basis. Anim. Behav. 29 95–101.

Beer, C. G. (1971). Individual Recognition of Voice in the Social Behavior of Birds (D. S. Lehrman, R. A. Hinde, & E. B. T.-A. in the S. of B. Shaw (eds.); Vol. 3, pp. 27–74). Academic Press. https://doi.org/https://doi.org/10.1016/S0065-3454(08)60154-0

Bize, P., Piault, R., Moureau, B., & Heeb, P. (2006). A UV signal of offspring condition mediates context-dependent parental favouritism. Proc. R. Soc. B Biol. Sci. 273 2063–2068. https://doi.org/10.1098/rspb.2006.3546

Brittan-Powell, E. F., Dooling, R. J., & Farabaugh, S. M. (1997). Vocal Development in Budgerigars (Melopsittacus Undulatus): Contact Calls. J. Comp. Psychol. 111 226–241. https://doi.org/10.1037/0735-7036.111.3.226

Buckley, P. A., & Buckley, F. G. (1972). Individual egg and chick recognition by adult royal terns (Sterna maxima maxima). Anim. Behav. 20 457-IN1. https://doi.org/https://doi.org/10.1016/S0003-3472(72)80009-5

Budden, A. E., & Wright, J. (2001). Falling on deaf ears: The adaptive significance of begging in the absence of a parent. Behav. Ecol. Sociobiol. 49 474–481. https://doi.org/10.1007/s002650100323

Carlson, N. V, Kelly, E. M., & Couzin, I. (2020). Individual vocal recognition across taxa: a review of the
literature and a look into the future. Philos. Trans. R. Soc. B Biol. Sci. 375 20190479.
https://doi.org/10.1098/rstb.2019.0479

Cavanagh, P. M., & Ritchison, G. (1987). Variation in the Bounce and Whinny Songs of the Eastern
Screech-Owl. Wilson Bull. 99 620–627. http://www.jstor.org/stable/4162469

Colombelli-Négrel, D., Hauber, M. E., Robertson, J., Sulloway, F. J., Hoi, H., Griggio, M., &
Kleindorfer, S. (2012). Embryonic learning of vocal passwords in superb fairy-wrens reveals
intruder cuckoo nestlings. Curr. Biol. 22 2155–2160. https://doi.org/10.1016/j.cub.2012.09.025

Cramp, S. (1985). Eurasian Scops Owl. In *The Birds of the Western Palearctic, Vol. 4.* Oxford University
Press; Oxford.

de Ayala, R. M., Saino, N., Møller, A. P., & Anselmi, C. (2007). Mouth coloration of nestlings covaries
with offspring quality and influences parental feeding behavior. Behav. Ecol. 18 526–534.
https://doi.org/10.1093/beheco/arm006

Denac, K., & Trilar, T. (2006). Individual recognition of Scops Owls (Otus scops) by spectrographic
analysis of their calls: a preliminary study. Razpr. IV. Razreda SAZU 47 87–97.

Dickens, M., & Hartley, I. R. (2007). Differences in parental food allocation rules: Evidence for sexual
conflict in the blue tit? Behav. Ecol. 18 674–679. https://doi.org/10.1093/beheco/arm029

Dor, R., Kedar, H., Winkler, D. W., & Lotem, A. (2007). Begging in the absence of parents: A "quick on
the trigger" strategy to minimize costly misses. Behav. Ecol. 18 97–102.
https://doi.org/10.1093/beheco/arl056

Draganoiu, T. I., Nagle, L., Musseau, R., & Kreutzer, M. (2006). In a songbird, the black redstart, parents
use acoustic cues to discriminate between their different fledglings. Anim. Behav. 71 1039–1046.
https://doi.org/10.1016/j.anbehav.2005.06.022

Dragonetti, M. (2007). Individuality in Scops Owl Otus scops vocalisations. Bioacoustics 16 147–172.
https://doi.org/10.1080/09524622.2007.9753573

Dreiss, A. N., Ruppli, C. A., & Roulin, A. (2014). Individual vocal signatures in barn owl nestlings: Does
individual recognition have an adaptive role in sibling vocal competition? J. Evol. Biol. 27 63–75.
https://doi.org/10.1111/jeb.12277

Dugas, M. B. (2009). House sparrow, Passer domesticus, parents preferentially feed nestlings with mouth

colours that appear carotenoid-rich. Anim. Behav. 78 767–772.

Ersoy, S., Maag, N., Boehly, T., Boucherie, P. H., & Bugnyar, T. (2021). Sex-specific parental care
during postfledging in common ravens. Anim. Behav. 181 95–103.
https://doi.org/10.1016/j.anbehav.2021.09.004

Falls, J. B. (1982). Individual recognition by sounds in birds. In D. E. Kroodsma & E. H. Miller (Eds.),
Acoustic communication in birds (pp. 237–278). Academic Press; New York.

Fitton, S. D. (1991). Vocal learning and call structure of male northern spotted owls in northwestern
California. Humboldt State University.

Freeman, P. L. (2000). Identification of individual Barred owls using spectrogram analysis and auditory
cues. J. Raptor Res. 34 85–92.

Galeotti, P., Paladin, M., & Pavan, G. (1993). Individually distinct hooting in male Pygmy Owls
Glaucidium passerinum: A multivariate approach. Ornis Scand. 24 15–20.
https://doi.org/10.2307/3676404

Galeotti, P., & Pavan, G. (1991). Individual recognition of male Tawny owls (Strix aluco) using
spectrograms of their territorial calls. Ethol. Ecol. Evol. 3 113–126.

Galeotti, P., & Sacchi, R. (2001). Turnover of territorial Scops Owls Otus scops as estimated by
spectrographic analyses of male hoots. J. Avian Biol. 32 256–262. https://doi.org/10.1111/j.0908-
8857.2001.320308.x

Galván, I., Amo, L., & Sanz, J. J. (2008). Ultraviolet-blue reflectance of some nestling plumage patches
mediates parental favouritism in great tits Parus major. J. Avian Biol. 39 277–282.
https://doi.org/https://doi.org/10.1111/j.0908-8857.2008.04273.x

Gilbert, G., & McGregor, P. K. (1994). Vocal individuality as a census tool: Practical consideration
illustrated by a study of two rare species. J. F. Ornithol. 65 335–348.

Grava, T., Mathevon, N., Place, E., & Balluet, P. (2008). Individual acoustic monitoring of the European
Eagle Owl Bubo bubo. Ibis (Lond. 1859). 150 279–287. https://doi.org/10.1111/j.1474-
919X.2007.00776.x

Grieco, F. (2018). Aggregation of Eurasian Scops Owls Otus scops Breeding in Magpie Pica pica Nests.
Ardea 106 177–191. https://doi.org/10.5253/arde.v106i2.a9

Grieco, F. (2022). Vocal behaviour reveals asymmetries in neighbour relationships in a semi-colonial

raptor, the Eurasian Scops Owl Otus scops. Behaviour 160 27–60.
https://doi.org/10.1163/1568539X-bja10188

Jourdie, V., Moureau, B., Bennett, A. T. D., & Heeb, P. (2004). Ultraviolet reflectance by the skin of nestlings. Nature 431 262. https://doi.org/10.1038/431262a

Jouventin, P., & Aubin, T. (2002). Acoustic systems are adapted to breeding ecologies : individual recognition in nesting penguins. Anim. Behav. 64 747–757. https://doi.org/10.1006/anbe.2002.4002

Kilner, R., & Johnstone, R. A. (1997). Begging the question: are offspring solicitation behaviours signals of need? Trends Ecol. Evol. 12 11–15.

Klenova, A. V., Volodin, I. A., & Volodina, E. V. (2009). Examination of pair-duet stability to promote long-term monitoring of the endangered red-crowned crane (Grus japonensis). J. Ethol. 27 401–406. https://doi.org/10.1007/s10164-008-0133-9

Klenova, A. V., Zubakin, V. A., & Zubakina, E. V. (2012). Inter- and intra-season stability of vocal individual signatures in a social seabird, the crested auklet. Acta Ethol. 15 141–152. https://doi.org/10.1007/s10211-011-0120-y

Koenig, L. (1973). Das Aktionssystem der Zwergohreule Otus scops scops (Linné 1758). Fortschritte Der Verhal. Adv. Ethol. 13 1–124.

Kouba, M., Bartoš, L., & Št'Astný, K. (2014). Factors affecting vocalization in Tengmalm's owl (Aegolius funereus) fledglings during post-fledging dependence period: Scramble competition or honest signalling of need? PLoS One 9. https://doi.org/10.1371/journal.pone.0095594

Krebs, E. A. (2001). Begging and food distribution in crimson rosella (Platycercus elegans) broods: why don't hungry chicks beg more? Behav. Ecol. Sociobiol. 50 20–30.

Lambrechts, M. M., & Dhondt, A. A. (1995). Individual voice discrimination in birds. In D. M. Power (Ed.), *Current Ornithology 12* (pp. 115–139). Plenum Press; New York. https://doi.org/10.1007/978-1-4615-1835-8_4

Lefevre, K., Montgomerie, R., & Gaston, A. J. (1998). Parent-offspring recognition in thick-billed murres (Aves: Alcidae). Anim. Behav. 55 925–938.

Lengagne, T. (2001). Temporal stability in the individual features in the calls of Eagle owls (Bubo bubo). Behaviour 138 1407–1419.

Leonard, M. L., & Horn, A. G. (2001). Begging in the absence of parents by nestling tree swallows.

Behav. Ecol. 12 501–505. https://doi.org/10.1093/beheco/12.4.501

Leonard, M. L., Horn, A. G., Brown, C. R., & Fernandez, N. J. (1997). Parent-offspring recognition in

tree swallows, Tachycineta bicolor. Anim. Behav. 54 1107–1116.

https://doi.org/https://doi.org/10.1006/anbe.1997.0559

Lessells, C. M. (2002). Parentally biased favouritism: Why should parents specialize in caring for

different offspring? Philos. Trans. R. Soc. B Biol. Sci. 357 381–403.

https://doi.org/10.1098/rstb.2001.0928

Lessells, C. M., & Boag, P. T. (1987). Unrepeatable Repeatabilities: A Common Mistake. Auk 104 116–

121. https://doi.org/10.2307/4087240

Ligout, S., Dentressangle, F., Mathevon, N., & Vignal, C. (2016). Not for Parents Only: Begging Calls

Allow Nest-Mate Discrimination in Juvenile Zebra Finches. Ethology 122 193–206.

https://doi.org/https://doi.org/10.1111/eth.12450

Loesche, P., Stoddard, P. K., Higgins, B. J., & Beecher, M. D. (1991). Signature Versus Perceptual

Adaptations for Individual Vocal Recognition in Swallows. Behaviour 118 15–25.

https://doi.org/https://doi.org/10.1163/156853991X00175

Lumivero. (2023). XLSTAT statistical and data analysis solution. New York, USA.

https://www.xlstat.com/en

Magrath, R. D., Haff, T. M., Horn, A. G., & Leonard, M. L. (2010). Calling in the face of danger:

predation risk and acoustic communication by parent birds and their offspring. Adv. Study Behav.

41 187–253. https://doi.org/https://doi.org/10.1016/S0065-3454(10)41006-2

Marques, P. A. M., De Araújo, C. B., & Vicentel, L. (2011). Nestling call modification during early

development in a colonial passerine. Bioacoustics 20 45–58.

https://doi.org/10.1080/09524622.2011.9753632

Medvin, M. B., & Beecher, M. D. (1986). Parent-offspring recognition in the barn swallow (Hirundo

rustica). Anim. Behav. 34 1627–1639. https://doi.org/https://doi.org/10.1016/S0003-3472(86)80251-

2

Medvin, M. B., Stoddard, P. K., & Beecher, M. D. (1992). Strong sib-sib call similarity in cliff swallows

not barn swallows. Ethology 90 17–28.

Medvin, M. B., Stoddard, P. K., & Beecher, M. D. (1993). Signals for parent-offspring recognition: a

comparative analysis of the begging calls of cliff swallows and barn swallows. Anim. Behav. 45

841–850. https://doi.org/https://doi.org/10.1006/anbe.1993.1105

Middleton, H. A., Green, D. J., & Krebs, E. A. (2007). Fledgling begging and parental responsiveness in

American dippers (Cinclus mexicanus). Behaviour 144 485–501.

https://doi.org/10.1163/156853907780713064

Mikkola, H. (1983). Owls of Europe. In *Calton, London* (1st ed.).

Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: A practical

guide for biologists. Biol. Rev. 85 935–956. https://doi.org/10.1111/j.1469-185X.2010.00141.x

Odom, K. J., Slaght, J. C., & Gutiérrez, R. J. (2013). Distinctiveness in the territorial calls of great horned

owls within and among years. J. Raptor Res. 47 21–30. https://doi.org/10.3356/JRR-12-11.1

Olsen, J., Trost, S., Fuentes, E., & Debus, S. (2020). *Behaviour and family association during the post-*

fledging period in Southern Boobooks Ninox boobook. 61–70.

Parejo, D., Avilés, J. M., & Rodríguez, J. (2010). Visual cues and parental favouritism in a nocturnal bird.

Biol. Lett. 6 171–173. https://doi.org/10.1098/rsbl.2009.0769

Peake, T. M., McGregor, P. K., Smith, K. W., Tyler, G., Gilbert, G., & GREEN, R. E. (1998).

Individuality in Corncrake Crex crex vocalizations. Ibis (Lond. 1859). 140 120–127.

https://doi.org/https://doi.org/10.1111/j.1474-919X.1998.tb04548.x

Redondo, T., & Exposito, F. (1990). Structural variations in the begging calls of nestling magpies Pica

pica and their role in the development of adult voice. Ethology 84 307–318.

Roulin, A. (2001). On the cost of begging vocalization: Implications of vigilance. Behav. Ecol. 12 506–

511. https://doi.org/10.1093/beheco/12.4.506

Roulin, A. (2002). The sibling negotiation hypothesis. In J. Wright & M. L. Leonard (Eds.), *The evolution*

of begging: competition, cooperation and communication (pp. 107–126). Springer; Dordrecht.

Roulin, A., Dreiss, A., Fioravanti, C., & Bize, P. (2009). Vocal sib-sib interactions: how siblings adjust

signalling level to each other. Anim. Behav. 77 717–725.

https://doi.org/10.1016/j.anbehav.2008.12.004

Roulin, A., Kölliker, M., & Richner, H. (2000). Barn owl (Tyto alba) siblings vocally negotiate resources.

Proc. R. Soc. B Biol. Sci. 267 459–463. https://doi.org/10.1098/rspb.2000.1022

Sharp, S. P., McGowan, A., Wood, M. J., & Hatchwell, B. J. (2005). Learned kin recognition cues in a

social bird. Nature 434 1127–1130. https://doi.org/10.1038/nature03522

Slagsvold, T. (1997). Brood division in birds in relation to offspring size: sibling rivalry and parental
control. Anim. Behav. 54 1357–1368. https://doi.org/https://doi.org/10.1006/anbe.1997.0530

Smiseth, P. T., Bu, R. J., Eikenæs, A. K., & Amundsen, T. (2003). Food limitation in asynchronous
bluethroat broods: Effects on food distribution, nestling begging, and parental provisioning rules.
Behav. Ecol. 14 793–801. https://doi.org/10.1093/beheco/arg083

Stoddard, P. K. (1996). Vocal recognition of neighbors in territorial passerines. In D. E. Kroodsma & E.
H. Miller (Eds.), *Ecology and evolution of acoustic communication in birds* (pp. 356–374). Cornell
University Press; Ithaca, NY.

Stoddard, P. K., & Beecher, M. D. (1983). Parental recognition of offspring in the cliff swallow. Auk 100
795–799.

Strickler, S. A. (2013). Recognition of young in a colonially nesting bird. Ethology 119 130–137.

Tibbetts, E. A., & Dale, J. (2007). Individual recognition: it is good to be different. Trends Ecol. Evol. 22
529–537. https://doi.org/10.1016/j.tree.2007.09.001

Tripp, T. M., & Otter, K. A. (2006). Vocal individuality as a potential long-term monitoring tool for
Western Screech-owls, Megascops kennicottii. Can. J. Zool. 84 744–753.
https://doi.org/10.1139/Z06-055

Vignal, C., Mathevon, N., & Mottin, S. (2008). Mate recognition by female zebra finch: Analysis of
individuality in male call and first investigations on female decoding process. Behav. Processes 77
191–198. https://doi.org/10.1016/j.beproc.2007.09.003

Yang, C., Wang, L., Chen, M., Liang, W., & Møller, A. P. (2015). Nestling recognition in red-rumped
and barn swallows. Behav. Ecol. Sociobiol. 69 1821–1826. https://doi.org/10.1007/s00265-015-
1994-x

Zhou, B., Xia, C.-W., Chen, Z.-R., & Deng, W.-H. (2020). Individual Identification of Male Ural Owls
Based On Territorial Calls. J. Raptor Res. 54 57–65. https://doi.org/10.3356/0892-1016-54.1.57

TABLES

Table 1

Descriptive statistics of ten acoustic variables of 670 vocalizations of 20 young Scops Owls after they left

the nest. The standard deviation (SD) represents the inter-individual variation. (*) Minimum and

maximum of individual averages. (**) Minimum and maximum of the single measurements (n=670).

	Average	SD	Range (individuals)*	Range (calls)**
Call duration (s)	0.182	0.045	0.117 – 0.279	0.094 – 0.423
Time from previous call (s)	1.775	0.244	1.410 – 2.302	0.770 – 6.490
Peak frequency (Hz)	1924.9	701.7	1316.4 – 4082.1	1112 – 5963
Peak frequency 1/3 (Hz)	1984.6	690.1	1321.8 – 3937.6	1113 – 5393
Peak frequency 3/3 (Hz)	1577.6	235.0	1221.5 – 2093.6	1074 – 5459
Centre of gravity (Hz)	2296.6	560.2	1722.7 – 3717.9	1468 – 4924
Pitch (Hz)	1820.3	484.2	1334.8 – 3551.0	1202 – 5046
Average entropy	-2.73	0.50	-3.92 – -1.80	-4.80 – -1.18
Minimum entropy	-3.89	0.73	-5.12 – -2.39	-7.25 – -1.45
Spectrum skewness	2.24	1.13	-0.04 – 4.11	-1.04 – 9.36

	Time from previous call	Peak frequency	Peak frequency 1/3	Peak frequency 3/3	Centre of gravity	Pitch	Average entropy	Minimum entropy	Spectrum skewness
Call Duration	0.27 ns	-0.43 ns	-0.43 ns	0.06 ns	-0.52 *	-0.37 ns	-0.24 ns	-0.41 ns	0.47 *
Time from previous call		-0.002 ns	-0.08 ns	0.42 ns	-0.09 ns	-0.14 ns	-0.07 ns	-0.18 ns	0.34 ns
Peak frequency			**0.99** ***	0.52 *	**0.93** ***	**0.84** **	0.38 ns	0.43 ns	**-0.77** **
Peak frequency 1/3				0.50 *	**0.92** ***	**0.85** **	0.38 ns	0.42 ns	**-0.77** **
Peak frequency 3/3					0.48 *	0.56 *	-0.05 ns	-0.17 ns	-0.22 ns
Centre of gravity						**0.84** **	0.49 *	0.55 *	**-0.90** ***
Pitch							0.14 ns	0.17 ns	-0.66 *
Average entropy								**0.92** ***	-0.50 *
Minimum entropy									-0.61 *

584   **Table 3.** Estimates of repeatability of acoustic features within six Scops Owl broods. Significance refers

585   to the F tests for each of the 14 recordings. Some broods are represented more often than others (see

586   Methods). *P* values are adjusted after Bonferroni correction ($P < 0.0036$).

	Average	CI	Significant	
Call duration (s)	0.91	0.87 – 0.96	14 out of 14	
Time from previous call (s)	0.16	0.06 – 0.26	2 out of 14	
Peak frequency (Hz)	0.60	0.41 – 0.79	11 out of 14	
Peak frequency 1/3 (Hz)	0.54	0.48 – 0.61	12 out of 14	
Peak frequency 3/3 (Hz)	0.55	0.36-0.74	13 out of 14	
Centre of gravity (Hz)	0.56	0.22 – 0.91	11 out of 14	
Pitch (Hz)	0.59	0.37 – 0.81	12 out of 14	

Average entropy	0.36	0.05 – 0.68	7 out of 14	
Minimum entropy	0.39	0.08 – 0.71	10 out of 14	
Spectrum skewness	0.62	0.41 – 0.83	11 out of 14	

Fig. 1

Example of spectral and temporal variation in vocalizations of young Scops Owls. **A**. Spectrogram of the

calls of two young one week after leaving the nest. (a) Short call with many frequency components (see

the spectrum on the left). (b) Longer call with a narrower bandwidth, and a main component around 2

kHz (see the spectrum on the right). **B**. Spectrogram of a train of calls by the same two young owls as in

A. The symbols below indicate the intervals between subsequent calls, which may show some regularity

as in this example. **C**. Spectrogram of the calls of two owlets and b in another brood. The arrows indicate

individually-repeatable frequency peaks, usually captured with Peak frequency 1/3 and Peak frequency

3/3; see Methods. The time bar represents one second. The vertical axis scale is in kHz

Fig. 2

Representative spectrograms of the calls of young Scops Owls of six broods (1-2 from 2021, 3-6 from

2022). The letters indicate features that help recognition of individuals. (a) Differences in call duration.

(b) S-shape patterns, usually at the lower end of the frequency range. (c) Wider frequency band in the

first part of the call, resulting in a higher centre of gravity and lower spectral skewness. (d) Mass of

frequency concentrated at specific points, especially at the end of the call (the "dots" in the spectrograms

indicated with arrows). (e) The dominant frequency is generally lower than in calls of other brood mates.

(f) More structured sound, with high spectral skewness

Fig. 3

Temporal variation of the calls' acoustic features in two fledglings (brood 5, compare with Figure 3, last

two spectrograms) during a 31-minute recording, which included two visits of one of their parents

(indicated by the owl symbols). During this period the owlets were always separated, so it was possible to

accurately tell where the caller was. Time is expressed as the time remaining to the first visit (0 s). Each

point represents the loudest call selected in a 30-second interval. The thin lines represent the running

average calculated with three values per data point. The five figures on the right show representative

spectrograms at five time points spread over the recording time; the numbers indicate the time points on

the top-left chart. In each figure, the first spectrogram refers to the individual represented with open

circles in the charts. Note the effect of the parental visits on call duration, time from the previous call,

spectral centre of gravity and spectral skewness of both individuals

Fig. 4

Temporal stability of the ten acoustic variables of the calls of two young scops owls (triangles and circles)

in 13 sets of measurements over 10 nights. The young left the nest on July 22. The squares indicate the

resulting repeatability R. Filled squares indicate statistically significant R after Bonferroni correction for

multiple tests ($P < 0.00385$)

Figure 1

636   **Figure 2**

 Figure 3

Figure 4

ELECTRONIC SUPPLEMENTARY MATERIAL

Online Resource 1 – ESM_1.tiff

Spectrogram of the calls of two young Scops Owls: a low-frequency, longer call (a) and a high-frequency,

shorter call (b). The line indicates the temporal changes of the Wiener entropy.

Online Resource 2 – ESM_2.tiff

Correlation matrix for the ten acoustic variables measured in 20 young Scops Owls. A line indicates a

significant correlation after Bonferroni correction (n= 45 correlation tests; $P < 0.0011$)

Online Resource 3 – ESM_3.tiff

Spectrum skewness plotted against the spectrum centre of gravity of young Scops Owl calls. Points

represent individuals' averages; symbols indicate the broods those individuals belong to. The insets show

examples of spectrograms of calls at the two extremes of the range. Top: low-frequency, highly skewed

calls. Bottom: calls with less skewed spectrum and frequencies distributed on a wider range. The width of

the spectrograms represents a one-second interval. Pearson correlation, $r = -0.896$, $n = 20$, $P < 0.00001$

Online Resource 4 – ESM_4.tiff

Temporal variation acoustic features of the calls of two brood mates (Figure 3, brood 6, last two

spectrograms) over two minutes preceding the visit by an adult. The arrival of the adult is indicated by the

owl symbol; Time is expressed as the time to the feeding event (0 s). The thin lines represent the running

average calculated with three values per data point. After one fledgling was fed, the other keeps calling

with longer, lower-frequency, lower-entropy calls

Online Resource 5 – ESM_5.tiff

Call spectrograms of young Scops Owls seven days before fledging. The first fragment includes three

calls, the others two calls each. The arrows indicate the call with a more structured sound, similar to that

detected later in the same brood (compare Figure 2, brood 6), suggesting that it could have been from the

same individual.

Online Resource 6 – ESM_6.m4v

Fragment of recording of alternated vocalizations of two brood mates (brood 6 in Figure 2). Although the

call spectrograms are similar, the two young make distinctive sounds. The sequence is as follows (a and b

indicate the two individuals, b slightly low-pitched than a): 15:07 a-b-a-b-a, 15:11 b-a-b-a-b-a, 15:16 b-a-

b-a-b-a. Note the "dots" in the spectrograms at specific frequencies. In the background, calls of adult

Scops Owls and crickets. This fragment was not included in the analysis because of the pervasive

background noise

681