A novel MARV glycoprotein-specific antibody with potentials of broad-spectrum

neutralization to filovirus

Running title: An anti-filovirus antibody fused with NPC2

Yuting Zhang[1,2†], Min Zhang[1†], Haiyan Wu[1†], Xinwei Wang[1,2†], Hang Zheng[1,2],

Junjuan Feng[1,2], Jing Wang[1], Longlong Luo[1], He Xiao[1], Chunxia Qiao[1], Xinying Li[1],

Yuanqiang Zheng[2], Weijin Huang[3], Youchun Wang[3], Yi Wang[4*], Yanchun Shi[2*],

Jiannan Feng[1*], Guojiang Chen[1*]

[1]State Key Laboratory of Toxicology and Medical Countermeasures, Institute of

Pharmacology and Toxicology, Beijing, China

[2]Inner Mongolia Key Lab of Molecular Biology, School of Basic Medical Sciences,

Inner Mongolia Medical University, Hohhot, China

[3]Division of HIV/AIDS and Sex-transmitted Virus Vaccines, National Institutes for

Food and Drug Control, Beijing, China

[4]Department of Hematology, Fifth Medical Center of Chinese PLA General Hospital,

Beijing, China

*Address correspondence to Yi Wang, yk091023@163.com; Yanchun Shi,

ycshi5388@163.com; Jiannan Feng, fengjiannan1970@qq.com or Guojiang Chen,

jyk62033@163.com.

†These authors contributed equally to this work

Abstract

Marburg virus (MARV) is one of the filovirus species that causes a deadly hemorrhagic fever in humans, with mortality rates up to 90%. Neutralizing antibodies represent ideal candidates to prevent or treat virus disease. However, no antibody has been approved for MARV treatment to date. In this study, we identified a novel human antibody named AF-03 that targeted MARV glycoprotein (GP). AF-03 possessed a high binding affinity to MARV GP and showed neutralizing and protective activities against the pseudotyped MARV in vitro and in vivo. Epitope identification, including molecular docking and experiment-based analysis of mutated species, revealed that AF-03 recognized the Niemann-Pick C1 (NPC1) binding domain within GP1. Interestingly, we found the neutralizing activity of AF-03 to pseudotyped Ebola viruses (EBOV, SUDV, and BDBV) harboring cleaved GP instead of full-length GP. Furthermore, NPC2-fused AF-03 exhibited neutralizing activity to several filovirus species and EBOV mutants via binding to CI-MPR. In conclusion, this work demonstrates that AF-03 represents a promising therapeutic cargo for filovirus-caused disease.

Keywords Marburg, glycoprotein, neutralizing antibody, NPC1, filovirus, NPC2

Introduction

Filoviruses are a nonsegmented negative-sense RNA viruses, comprised of six genera, Ebolavirus, Marburgvirus, Cuevavirus, Striavirus, Thamnovirus and a recently discovered sixth genus, Dianlovirus[1-3]. The Marburgvirus genus consists of Marburg virus (MARV) and Ravn virus (RAVN)[1,4,5]. The former includes three strains-- Uganda, Angola and Musoke. The Ebola virus genus includes six distinct species Zaire Ebola virus (EBOV), Bundibugyo virus (BDBV), Sudan virus (SUDV), Reston virus (RESTV), Taii Forest virus (TAFV) and Bombali virus (BOMV), the first three of which cause severe hemorrhagic fevers[6,7]. The genus Cuevavirus (Lloviu virus, LLOV) was isolated from Miniopterus schreibersii bats in Spain and Hungary and potently infected monkey and human cells[8,9]. The genus Měnglà virus (MLAV) was discovered in the liver of a bat from Mengla, Yunnan, China in 2019. So far, only an almost complete RNA sequence of the viral genome is available, there are no viable MLAVs isolated.[10] MARV and EBOV infect humans and non-human primates, causing Marburg virus disease (MVD) and EBOV virus disease (EVD) with an incubation period of 2-21 days[11]. The symptoms of MVD include severe headache and high fever rapidly within 5 days of the onset of symptoms, followed by diarrhea and vomiting, leading to up to 90% fatality rate[11]. MARV and EBOV have a high potential to cause a public health emergency.

Glycoprotein (GP) on the surface of filoviruses is a type I transmembrane protein and consists of GP1 and GP2 subunits[12,13]. It is inserted into the virus envelope in the form of homotrimeric spikes[14] and is responsible for virus attachment and entry. The

furin cleaves Marburg GP at the amino acid 435 into two subunits, GP1 and GP2,

which remains linked by a disulfide bond[15]. GP1 contains a receptor binding domain

(RBD), a glycan cap, and a heavily glycosylated mucin-like domain (MLD), which

mediates binding to entry factors and receptors[16]. GP2 has a partial MLD, a

transmembrane domain for viral anchoring to the envelop surface, and a fusion

peptide required for the fusion of virus and cell membranes[17-19]. In the Ebola virus,

the furin cleavage site is located at residue 501 and the entire MLD is attached to the

GP1 subunit[20]. Marburg virus contains 66 amino acids on GP2 that are absent from

the Ebola virus MLD, and are called "wings" due to their outward projection and

flexibility[17].

Currently, GP is a major target for antibodies validated in filovirus-infected

animals and clinical trials because it is exposed on the surface of the virus and plays a

key role in viral entry[21]. Filoviruses initially enter cells by endocytosis[22,23]. Once

inside the endosome, GP is cleaved by host cathepsins and glycan cap and MLD are

removed, enabling GP to bind to the Niemann-Pick C1 (NPC1) receptor[24,25].

Interestingly, Ebola virus entry requires cathepsin B cleavage[26], which is redundant

for Marburg virus entry[27,28]. Hashiguchi et al. propose that the receptor binding

domain is masked by glycan cap and MLD in the Ebola virus, whereas it is partially

exposed in the Marburg virus[16].

To date, there is no licensed treatment or vaccine for Marburg infection. Herein,

we utilized phage display technology to screen an antibody in a well-established

antibody library[29,30] and obtained a novel human antibody with prominent neutralizing

activity. Furthermore, NPC2 fusion at the N terminus of the light chain of this

antibody renders broad-spectrum inhibition of cell entry of filovirus species and

mutants.

Materials and methods

Cell lines and plasmids

Human embryonic kidney cells HEK293T and human hepatoma cells Huh7 were purchased from ATCC. These cell lines were cultured in Dulbecco's modified Eagle's medium (DMEM) (Gibco, 11965e092) supplemented with 100 units/ml penicillin, 100 units/ml streptomycin (Gibco, 15140) and 10% fetal bovine serum (Gibco, 10099) in a humidified atmosphere (5% CO_2, 95% air) at 37°C. ExpiCHO-S cells were purchased from Gibco and cultured in ExpiCHO™ Expression Medium (Gibco, A29100) in a humidified atmosphere (8% CO_2, 92% air) on an orbital shaker platform.

MARV (AFV31370.1), Angola (Q1PD50.1), Musoke (YP_001531156.1), Ravn (YP_009055225.1), TAFV (Q66810), RESTV (Q66799), BOMV (YP_009513277.1) MLAV (YP_010087186.1), LLOV (JF828358) GPs plasmid synthesized by GENEWIZ and then cloned into the expression vector pcDNA3.1. EBOV (A0A068J419), BDBV (AYI50382), SUDV (Q7T9D9) GPs and HIV-based vector pSG3. Δenv. cmvFluc plasmids were kindly gifted by China Institute for Food and Drug control.

Preparation of full-length antibody and antigen

AF-03 was selected from a human phage antibody library, which displays on the surface of M13 bacteriophage particles. Screening procedures were as described in detail previously[34,35]. Phage antibodies that bound to MARV GP protein were obtained to express full-length IgGs using a standard protocol. In brief, the VH and

VL region of AF-03 were constructed into a mammalian full-length

immunoglobulin expression vector pFRT-KIgG1 (Thermo, V601020), to generate

plasmid AF-03. The human NPC2 gene (aa20-151) was linked to the VL of AF-03 by

a short linker "TVAAP" and then constructed into pFRT-KIgG1 (designated as

AF03-NL). The AF-03 and AF03-NL plasimid were transfected into ExpiCHO-S cells

using the ExpiFectamine™ CHO Transfection Kit (Gibco, A29129) following the

manufacturer's instructions. Purification was performed using ÄKTA prime plus

system (GE Healthcare) with protein A column (GE Healthcare). MARV GP (Uganda

strain) (aa 20-648, Δ277-455), CI-MPR1-3 (aa36-466) and NPC2 (aa20-151) gene

with six-histidine-tagged at C-terminus was cloned into mammalian expression vector

pcDNA3.1 and then transfected into HEK293T cells. MARV GP was purified using

nickel column (GE Healthcare, 11003399). The concentration of proteins and

antibodies were quantified by bicinchoninic acid (BCA) method.

ELISA

The 96-well enzyme-labelled array plates were coated with 2 μg/ml MARV-GP and

mutated MARV GP ($Q^{128}S$-$N^{129}S$) respectively and incubated overnight at 4°C. Wells

were washed for three times and blocked for 1 h at 37°C. A series of 12

concentrations of AF-03 and MR78 were added and incubated for 1 h at 37 °C. Bound

antibodies were detected with horseradish peroxidase (HRP)-labeled goat anti-human

IgG secondary antibody (Invitrogen, A18817) at room temperature for 30 min.

Binding signals were visualized using a TMB substrate (CWBIO, CW0050S) and the

reaction was stopped by adding 2 N H_2SO_4. The light absorbance at 450 nm was

measured by microplate reader (Thermo Fisher Scientific). The concentration for 50%

of maximal effect was defined as EC_{50}.

For competitive ELISA, biotinylated AF-03 (1 μg/ml) was coated. MR78 and

control mAb (Herceptin) in 3-fold serial dilution (ranging from 0.27 to 200 ug/ml)

and added to the plates. After 1 h incubation at 37°C, the plates were washed and the

bound Biotin-AF-03 was detected by adding horseradish peroxidase (HRP)-labeled

Streptavidin (Thermo, S911). After a further 30 min incubation at room temperature,

the plates were washed and TMB was added. The reaction was stopped by adding 2 N

H_2SO_4. Absorbance was measured at 450 nm using a plate reader.

BLI analysis of antibody affinity

The binding affinity of AF-03 to MARV GP was measured by the Fortebio biofilm

interferometry technique using anti-huamn IgG Fc capture (AHC) biosensors. In brief,

8 μg/ml antibody was diluted to PBS containing 0.02% Tween 20 and immobilized in

biosensors. The biosensors were then immersed with a serial dilutions of MARV GP

to determine the association constant. Programs setting were following: baseline 60 s,

loading 60 s, baseline 60 s, association 180 s, dissociation 220 s, regeneration 5 s.

Baseline was performed with PBST buffer and regeneration was performed with 10

mM glycine-HCl (pH 1.7) thrice. The data were analyzed using FortéBio Data

Analysis 9.0 (Sartorius, FortéBio®).

Computer□guided homology modeling and molecular docking

The three-dimensional theoretical structure of fragment variable (FV) was constructed

using computer-guided homology modeling approach based on the amino acid

sequences of the variable structural domains of the heavy and light chains of AF-03,

and the conserved regions and loop structural domains were identified. The 3-D

structure of the AF-03 Fv fragment was optimized under the consistent valence force

field (CVFF) using the steepest descent and conjugate gradient minimization methods.

The final minimized 3-D structure was evaluated by means of Ramachandran

diagrams. In addition, the 3-D theoretical structure of the MARV GP protein was

obtained and optimised using the CVFF force field. Under molecular docking method,

the 3-D complex structures AF-03 Fv fragment and GP were obtained and optimized.

With the determined 3-D structure of the AF-03 Fv fragment and GP, 50-ns molecular

dynamics were performed with the Discovery_3 module. All calculations were

performed using Insight II 2000 software (MSI Co., San Diego) with IBM

workstation.

Pseudovirus preparation

HIVΔenvor (pSG3.Δenv.cmvFluc) bearing MARV, mutated MARV ($Q^{128}S$-$N^{129}S$,

$T^{204}A$-$Q^{205}A$-$T^{206}A$, $Y^{218}A$, $K^{222}A$ and $C^{226}Y$), EBOV (parental and 17 mutants

indicated), SUDV, BDBV, TAFV, RESTV, BOMV, RAVN, MLAV and LLOV GPs

were prepared by liposome-mediated transfection of HEK293T cells using JetPRIME

(Polyplus Transfection, 25Y1801N5). Cells were seeded in 6-well plates at a density

of 7×10^{5}cells/well and transfected with 2 μg plasmids (0.4 μg GP and 1.6 μg

HIVΔenvor) when cells reached 60-80% confluence. Supernatants were collected 48

h after transfection, centrifuged to remove cell debris at 3,000 rpm for 10 min, filtered

through a 0.45 μm-pore filter (Millipore,SLHUR33RB) and stored at -80°C.

Pseudovirus entry and antibody neutralization assay

Huh7 and HEK293T cells (3×10^4 cells/100 µl/well) were infected with 100 µl

pseudovirus, which contained a luciferase reporter gene respectively. The luciferase

activity were measured in a fluorescence microplate reader (Promega). The operation

steps were following: after 36 h incubation at 37°C, 100 µl of culture medium were

discarded and addition with 100 µl of Bright-Glo luciferase reagent (Promega, E6120)

in each well. Mixtures were transfered to 96-well whiteboards after 2-minutes

reaction to detect the relative luciferase intensity.

For AF-03 neutralization of MARV assays, 50 µl mAb was 3-fold serially

diluted and separately mixed with MARV pseudovirus at the same volume. The

mixture were incubated at 37°C for 1 h, followed by the addition of 100 µl cells

(3×10^4 cells/well). 50% of maximal inhibitory concentration was defined as IC_{50}. IC_{50}

values were determined by non-linear regression with least-squares fit in GraphPad

Prism 8 (GraphPad Software).

In terms of AF-03 neutralization of ebola virus assays, pseudovirus (EBOV,

SUDV and BDBV) were processed with thermolysin as previously described [36].

Briefly, Pseudotyped ebola virus were incubated at 37°C for 1 h with 200 µg/ml

thermolysin (Sigma, T7902). The reaction was stopped by addition of 400 µM

Phosphoramidon (Sigma, R7385) on ice for 20 min. The remaining steps followed

AF-03 neutralization of MARV assays.

For AF03-NL neutralization assays, 50 µl/well of serial diluted AF03-NL and

AF-03 were incubated with cells at 37°C for 2 h to enable internalization of the

antibodies. 50 μl/well of diluted pseudovirus was added to a 96-well plate and

incubated at 37°C for 36 h. Bright-Glo luciferase reagent was added to detect the

relative luciferase intensity.

Bioluminescent imaging in vivo

Four-week-old female BALB/c mice were purchased from Beijing Vital River

Laboratory Animal Technology Co. Ltd. Mice were intraperitoneally injected with

MARV pseudovirus (0.2 ml/mouse). AF-03 (10, 3 or 1 mg/kg) or control antibody

Herceptin (10 mg/kg) were injected via intravenous route 24 h and 4 h before the

pseudovirus injection respectively. Bioluminescent signals were monitored at Day 5.

Briefly, D-luciferin (150 mg/kg body weight) (PerkinElmer, 122799) was

intraperitoneally injected into the mice, and then exposed to Isoflurane alkyl for

anesthesia. Bioluminescence was measured by the IVIS Lumina Series III Imaging

System (Xenogen, Baltimore, MD, USA) with the living Image software. The signals

emitted from different regions of the body were measured and presented as total

fluxes. All data are presented as mean values ± SEM.

Evaluation of internalization

HEK293T cells were counted 3×10^5 cells, cold PBS washed twice and discarded the

supernatant. 20 μg/ml AF-03/AF03-NL was added and incubated at 4°C for 30 min.

Then internalization group was transferred to 37°C for internalization for 30 min,

while the control group continued to be incubated at 4°C for 30 min to adhere to the

cell surface. The cells were washed with cold PBS, PE-anti-human IgG Fc secondary

antibody (Biolegend, 41070) was added and incubated for 30 min at 4°C. The cells

were collected for analysis.

pHrodo Red labelling and immuno-localization

AF03-NL and AF-03 were covalently labeled with pH-sensitive pHrodo red

succinimidyl ester (Thermo, P36600) according to the manufacturer's instructions.

Antibodies were incubated with 10-fold molar excess of pHrodo red succinimidyl

ester for 1 h at room temperature. Excess unconjugated dye was removed using PD-10

desalting columns (GE Healthcare). pHrodo Red-labeled antibodies were exchanged

into HEPES buffer and concentrated in an Amicon Ultra centrifugal filter unit with a

nominal molecular weight cutoff of 30 kDa. Antibody concentration and degree of

labeling was determined according to the manufacturer's instructions.

HEK293T cells ($1\sim2\times10^5$ cells per dish) were cultured overnight in confocal dish

pre-treated with polylysine (Beyotime, ST508) and then incubated with pHrodo

Red-labeled AF-03 and AF03-NL (20 μg/ml) at 37°C for 30 min. Unbound antibodies

were removed by washing with cold PBS. As well, cells were stained with cell

membrane dye (DiD) (Thermo, V22887) and Hoechst33342 (Thermo, H1398) at

37°C for 15 min. Single cells were analyzed for pHrodo Red fluorescence

on.Confocal Microscopy (dragonfly 200).

CI-MPR knockin and knockdown

Huh7 Cells were seeded in 6-well plates at 3×10^5 cells/well and transfected with 2 μg

CI-MPR expression plasmids (JetPRIME) when cells reached 40-60% confluence and

cultured for 24 h. For CI-MPR silencing, HEK293T cells were seeded in 6-well plates

and transfected with siRNA-CI-MPR (Ribbio) and cultured for 48-72 h. CI-MPR

expression was detected by flow cytometry.

Flow cytometry

The cells were harvested and stained with FITC-conjugated anti-CI-MPR antibody

(Biolegend, 364207) on ice for 30 min. Cells were washed twice and detected on

FACSAria II (BD Biosciences). Data analysis was performed using the FlowJo

software.

Statistical analyses

Data were analyzed, and the graphs were plotted using Prism software (GraphPad

Prism 8, San Diego, USA). The data are presented as the mean ± standard error of the

mean. Intergroup differences were compared using unpaired t-tests or ANOVA. The *P*

<0.05 was considered statistically significant.

Results

Characteristics of AF-03

AF-03 was selected from a well-established phage surface display antibodies

library with immense diversity in the selected complementarity determining region

(CDR) loops[34,35]. We further subcloned VH and VL sequences of the antibody into a

mammalian full-length immunoglobulin expression vector for full‑length IgG

expression. As shown in Fig. 1A, AF-03 was eukaryotically expressed with the purity

over 95%. To determine the binding affinity, recombinant MARV GP without MLD

was prepared (Fig. 1A). ELISA analysis showed that AF-03 bound to MARV GP with

EC_{50} of 0.16 µg/ml (Fig. 1B). Furthermore, BLI assay was done to determine the

binding kinetics and showed that AF-03 bound to MARV GP with high affinity (K_D

value was less than 1×10^{-12}M) (Fig. 1C). To identify determinants of MARV GP

binding to AF-03, we utilized computer-guided homology modeling and molecular

docking to generate computer models of MARV GP in complex with AF-03. Firstly,

we obtained the theoretical 3-D structure of AF-03 Fv (Fig. 1D). Based on the 3-D

structure of AF-03 and MARV GP separately, the 3-D complex structure of AF-03 and

MARV GP achieved utilizing the molecular docking method, as shown in Fig. 1E.

Overall, these data suggest the potency of AF-03 binding to MARV GP.

Epitope mapping of MARV GP bound to AF-03

Under CVFF forcefield, chosen steepest descent and conjugate gradient

minimization methods, after 30,000 steps of minimization with the convergence

criterion 0.02 kCal/mol, the optimized structure of the AF-03 Fv was evaluated. Using

a Ramachandran plot, the assignment of the whole heavy atoms of the AF-03 Fv was

in the credible range. Through analyzing the van der Waals interactions,

inter-molecular hydrogen bonds, polar interactions and electrostatic interactions

between AF-03 and MARV GP, the key amino acid residues of MARV GP were

selected for amino acid point mutations (Fig. 1F). T^{204}-Q^{205}-T^{206}, Y^{218} and K^{222} were

mutated to alanine, and Q^{128}-N^{129} were mutated to serine as well as C^{226} was mutated

to tyrosine. Firstly, We investigated if this mutated MARV species was still sensitive

to AF-03 treatment. The inhibition assay revealed the impairment of neutralizing

activity of AF-03 to mutated MARV pseudovirus, which indicates that Q^{128}/N^{129}/C^{226}

functions as key amino acids responsible for AF-03 neutralization (Fig. 2A left panel).

To this end, we constructed the mutated MARV GP. ELISA assay showed that

mutation of Q^{128}S-N^{129}S or C^{226}Y significantly disrupted binding of GP to AF-03,

while the binding and neutralizing capacity of MR78 to mutant GP   and pseudovirus

(C^{226}Y), a mAb reported to be isolated from Marburg virus survivors[16], was not

almost affected (Fig. 2A right panel and B). Furthermore, we analyzed the secondary

structure of the MARV GP and its mutants. By circular dichroism, the structure of

both mutants was not obviously different from that of the parent GP (Fig. 3C, Tab. s1).

Therefore, the weakened binding to the antibody was not due to the conformational

change of the protein caused by the mutation. Competitive ELISA showed that AF-03

and MR78 could compete with each other to bind to MARV GP (Fig. 3D). These

results indicate the epitopes of these two mAbs overlapped partially.

In vitro neutralizing activity of AF-03

Given the high binding affinity of AF-03 to MARV GP, we sought to determine

whether AF-03 could impede MARV pseudotyped viral entry. An in vitro

neutralization assay was developed based on full-length MARV GP-pseudotyped

virus using a HIVΔenvor (pSG3.Δenv.cmvFluc). Liver and adrenal glands have been

reported to be the early targets of MARV infection [14,37]. Therefore, we first tested the

entry of MARV to hepatocyte cell line (Huh7) and HEK293T cells (renal cell line) by

measuring the relative luciferase intensity. These two cell lines were susceptible to

MARV cell entry. We used MR78 and cetuximab as positive and negative controls

respectively. As expected, cetuximab had no effects on pseutotyped MARV entry. In

contrast, AF-03 actively inhibited viral entry to HEK293T cells, with IC_{50} value of

0.13 μg/ml. As well, IC_{50} of MR78 was 0.44 μg/ml (Fig. 3A left panel). In Huh7 cells,

IC_{50} of AF-03 and MR78 was 0.4 and 1.03 μg/ml, respectively (Fig. 3A right panel).

These results suggest that AF-03 had more potency of neutralization than MR78. We

also conducted AF-03 neutralization experiments on pseudotyped Angola, Musoke

and Ravn strains and showed strong and comparable neutralizing ability to all these

strains (IC_{50} was 0.32, 0.12 and 0.15 μg/ml respectively) (Fig. 3B). Taken together,

these data suggest that AF-03 harbors prominent neutralizing activity to MARV

infection.

In vivo preventive efficacy of AF-03

To verify the in vivo preventive efficacy, AF-03 was intravenously injected into

mice before pseudotyped MARV exposure (-24 h and -4 h) respectively. The

bioluminescence intensity was measured on day 5 after pseudovirus injection. As

shown in Fig. 3C and D, AF-03-treated group displayed lower bioluminescence activity compared with the control group, while the treatment with control antibodies had no effects. Administration of 1 mg/kg AF-03 prior to the injection of MARV could decrease viral infection to approximately 50% level and increasing doses of AF-03 led to higher preventive efficacy. This indicates clearly that AF-03 is capable of preventing against MARV infection in a dose-dependent manner and represents a potential candidate for MARV prophylaxis.

AF-03 impedes cell entry of EBOV, SUDV and BDBV harboring GPcl

Given the close structural similarity of Marburg virus to ebola virus, to determine whether AF-03 was also available to the treatment of EBOV infection, we conducted neutralization of pseudotyped EBOV, SUDV and BDBV with AF-03. In addition, considering that glycan cap or mucin-like domain are known to mask the putative receptor-binding domain on EBOV, SUDV and BDBV GP, leading to blockade of engagement between AF-03 and GP[16]. Furthermore, in vitro cleavage of particles with thermolysin functionally mimics the cleavage due to the combination of cathepsin L and cathepsin B. Accordingly, glycan cap and mucin-like domain were enzymatically cleaved by digesting GP with 250 μg/ml thermolysin at 37°C. Consequently, the inhibitory function of AF-03 on cell entry of all three species of ebola virus bearing cleaved GP was much stronger than those bearing uncleaved GP (Fig. 4). Overall, these data suggest that AF-03 has therapeutic potentials for EBOV, SUDV and BDBV infection.

The potency of NPC2-fused AF-03 to be delivered into the endosome

Given the inability of AF-03 to transport into endosomal compartment where intact

GP is cleaved by cathepsin B/L, we engaged NPC2 to the N-terminus of light chain of

AF-03 (Fig. s1A), according to a protocol described previously[38]. As well, we

produced the 1-3 domain of CI-MPR (Fig. s1A), which is a ligand for NPC2 and

expressed on the cellular and endosomal membrane[32]. The results showed that

NPC2-fused AF-03 (termed AF03-NL), rather than AF-03, bound to CI-MPR1-3 (Fig.

s1B). Next we investigated the internalization of AF03-NL. AF03-NL or AF-03 was

incubated with HEK293T cells, which expressed CI-MPR (Fig. s2A), at 4□ for

attachment. As expected, AF03-NL instead of AF-03 adhered to the cell surface,

detected by fluorescence-labelled secondary IgG. Upon endocytosis, the fluorescence

on the cell surface decreased dramatically, implying the occurrence of AF03-NL

internalization (Fig. 5A). To further address this issue, AF-03 and AF03-NL were

labelled by pHrodo Red dye that is sensitive to acidic niche. Accord with Fig. 5A,

pHrodo Red-conjugated AF03-NL was observed in the acidic endosomal

compartment by flow cytometry and fluorescence microscopy respectively (Fig.   5B

and C). In contrast, AF-03 was not seen in the endosome.

Pan-filovirus inhibition of cell entry by AF03-NL via engagement between NPC2

and CI-MPR

We first compared the binding of AF03-NL/AF-03 to MARV GP. ELISA showed

relatively weak binding activity of AF03-NL compared with AF-03 (Fig. s3A). We

thereafter evaluated the neutralizing activity of AF03-NL to MARV pseudovirus.

Intriguingly, AF03-NL showed stronger neutralizing activity than AF-03 (The IC_{50}

was 0.057 and 0.284 μg/ml, respectively) (Fig. s3B), which may be attributed to

sustained tethering of AF03-NL to pseudovirus at extracellular space as well as

endosomal compartment. Next we compared the neutralizing activity of AF03-NL and

AF-03 to a series of filovirus species. AF03-NL displayed superior neutralizing

activity to other nine filovirus species. While, no or weak inhibition of entry by AF-03

was found (Fig. 6A). Furthermore, AF03-NL, instead of AF-03, also actively inhibited

cell entry of 17 EBOV mutants that were detected in natural hosts (Fig. 6B).

To investigate the mechanisms mediating the potency of AF03-NL, we produced

NPC2 protein (Fig.7A) and then examined the inhibition of EBOV entry by AF03-NL

or the mixture of AF-03 and NPC2. AF-03, NPC2 alone or in combination did not

inhibit EBOV entry. Conversely, AF03-NL actively impeded this process (Fig. 7B).

To clarify whether this effect is CI-MPR-dependent, CI-MPR in HEK293T cells was

silenced (Fig. 7C). The result showed that CI-MPR knockdown rendered significant

abrogation of the inhibitory ability of AF03-NL (Fig. 7C). We also introduced

CI-MPR into Huh7 cell line that is null for this receptor (Fig. s2B). The inhibitory

effects of AF03-NL were augmented in CI-MPR-overexpressed cells compared with

empty vector-introduced counterparts (Fig. 7D). Taken together, these data indicate

that the inhibitory potency of AF03-NL is dependent on the interaction between NPC2

and CI-MPR.

Discussion

The Marburg virus was initially identified after simultaneous outbreaks in Marburg and Frankfurt in Germany in 1967[16,39]. To date, there have been a dozen outbreaks of Marburg virus infection in humans[40]. Giving the recurrence of Marburg virus outbreaks and its high virulence and lethality, there is an urgent need to develop prophylactic and therapeutic interventions for Marburg infections. MARV GP is a surface viral protein, which is responsible for host receptor binding and cell entry thus provides an attractive target for the development of antagonists. Flyak et al. screened several MARV GP-specific neutralizing antibodies from the plasma of a MARV-infected survivor, which achieved 100% protection in mice subjected to mouse-adapted MARV challenge[41]. The MARV GP-specific antibody cocktail was also developed, 3 mAbs cocktail could protect hamster from lethal hamster-adapted MARV infection, while treatment with either one or two antibodies failed[42].

In this study, we selected an antibody from a human antibody phage library and the affinity constant reached the 1×10^{-12}M level. The neutralizing activities of the antibody were demonstrated by utilizing pseudotyped MARV Uganda strain. The results showed that AF-03 effectively inhibits HIVΔenvor (pSG3.Δenv.cmvFluc) pseudotyped MARV viral entry at IC_{50} of 0.13 and 0.4 μg/ml in HEK293T and Huh7 respectively. Furthermore, compared with control antibody, AF-03 exhibited a protective property against pseudovirus infection in mice. Epitope mapping results showed that GP Q^{128}-N^{129} and C^{226} was the binding and functional epitope that interacted with AF-03, which means AF-03 targeting the interface of GP-NPC1

interaction, considering that N129 is known to be located in the NPC1 binding

domain. RBDs are highly conserved among filovirus species, so they are an attractive

target for broadly effective anti-filovirus drug development[49]. We found that AF-03

also bound to EBOV GPcl and can neutralized ebola viruses bearing in vitro cleaved

GP, suggesting that AF-03 represents a good candidate for endosome-delivering

strategy by ligation to another mAb against a surface-exposed EBOV GP epitope or a

ligand peptide for host cation-independent mannose-6-phosphate receptor[38], which

will ultimately affords cross-reactivity against multiple filovirus species. Accordingly,

we designed NPC2-fused AF-03 and demonstrated its broad-spectrum inhibitory

capacity to filovirus species and EBOV mutants. Future investigations on the

inhibition of AF03-NL to authentic virus infection in vitro and in vivo are warranted.

Overall, our study identified a high-affinity anti-MARV antibody AF-03

targeting a conserved and hidden site at the filovirus GPcl-NPC1 interface, which was

capable of neutralizing MARV infection both in vitro and in vivo. Furthermore,

AF-03 may be a potential candidate for the effective protection against pan-filovirus

species infection. Investigations on AF-03 treatment of mice challenged by authentic

virus are undergoing.

Abbreviations

MARV: Marburg virus; EBOV: Ebola virus; SUDV: Sudan virus; BDBV: Bundibugy virus; mAb: monoclonal antibody; PBS: phosphate-buffered saline; GP: glycoprotein; RAVN: Ravn virus; RESTV: Reston virus; TAFV: Tai forest virus; MVD: Marburg virus disease; EVD: EBOV virus; RBD: receptor binding domain; MLD: mucin like domain; NPC1: Niemann-Pick C1; CDR: complementarity determining region; CVFF: consistent valence force field; FV: fragment variable; CI-MPR: cation-independent mannose-6-phosphate receptor; NPC2: Niemann-Pick C2

Compliance with ethics guidelines

Yuting Zhang, Min Zhang, Haiyan Wu, Xinwei Wang, Hang Zheng, Junjuan Feng,

Jing Wang, Longlong Luo, He Xiao, Chunxia Qiao, Xinying Li, Yuanqiang Zheng,

Weijin Huang, Youchun Wang, Yi Wang, Yanchun Shi, Jiannan Feng, Guojiang Chen

declare that they have no competing interests.

Author contributions

Guojiang Chen, Jiannan Feng, Yanchun Shi and Yi Wang conceived and designed

this study. Guojiang Chen, Jiannan Feng and Yuanqiang Zheng provided funding

support. Yuting Zhang, Min Zhang, Haiyan Wu and Xinwei Wang performed of the

experiments and prepared the manuscript. Hang Zheng and Junjuan Feng were

involved in optimization of the experimental protocols. Jing Wang, Longlong Luo, He

Xiao, Chunxia Qiao, Xinying Li, Yuanqiang Zheng, Weijin Huang, Youchun Wang

provided methodological support. All authors contributed to the article and approved

the submitted version.

Acknowledgments

This work is granted from the National Natural Science Foundation of China (81672803, 31771010, 81871252).

460 **Ethics approval**

461 The animal study was reviewed and approved by the Institutional Animal Care and

462 Use Committee of Academy of Military Medical Sciences.

References

1.Amarasinghe, G.K., Ayllón, M.A., Bào, Y., Basler, C.F., Bavari, S., Blasdell, K.R., Briese, T., Brown, P.A., Bukreyev, A., Balkema-Buschmann, A., et al. (2019). Taxonomy of the order Mononegavirales: update 2019. Arch Virol *164*, 1967-1980. 10.1007/s00705-019-04247-4.

2.Baseler, L., Chertow, D.S., Johnson, K.M., Feldmann, H., and Morens, D.M. (2017). The Pathogenesis of Ebola Virus Disease. Annu Rev Pathol *12*, 387-418. 10.1146/annurev-pathol-052016-100506.

3.Kuhn, J.H., Adachi, T., Adhikari, N.K.J., Arribas, J.R., Bah, I.E., Bausch, D.G., Bhadelia, N., Borchert, M., Brantsæter, A.B., Brett-Major, D.M., et al. (2019). New filovirus disease classification and nomenclature. Nat Rev Microbiol *17*, 261-263. 10.1038/s41579-019-0187-4.

4.Bào, Y., Amarasinghe, G.K., Basler, C.F., Bavari, S., Bukreyev, A., Chandran, K., Dolnik, O., Dye, J.M., Ebihara, H., Formenty, P., et al. (2017). Implementation of Objective PASC-Derived Taxon Demarcation Criteria for Official Classification of Filoviruses. Viruses *9*. 10.3390/v9050106.

5.Ristanović, E.S., Kokoškov, N.S., Crozier, I., Kuhn, J.H., and Gligić, A.S. (2020). A Forgotten Episode of Marburg Virus Disease: Belgrade, Yugoslavia, 1967. Microbiol Mol Biol Rev *84*. 10.1128/MMBR.00095-19.

6.Fan, P., Chi, X., Liu, G., Zhang, G., Chen, Z., Liu, Y., Fang, T., Li, J., Banadyga, L., He, S., et al. (2020). Potent neutralizing monoclonal antibodies against Ebola virus isolated from vaccinated donors. mAbs *12*, 1742457. 10.1080/19420862.2020.1742457.

7.Goldstein, T., Anthony, S.J., Gbakima, A., Bird, B.H., Bangura, J., Tremeau-Bravard, A., Belaganahalli, M.N., Wells, H.L., Dhanota, J.K., Liang, E., et al. (2018). The discovery of Bombali virus adds further support for bats as hosts of ebolaviruses. Nat Microbiol *3*, 1084-1089. 10.1038/s41564-018-0227-2.

8.Negredo, A., Palacios, G., Vázquez-Morón, S., González, F., Dopazo, H., Molero, F., Juste, J., Quetglas, J., Savji, N., de la Cruz Martínez, M., et al. (2011). Discovery of an ebolavirus-like filovirus in europe. PLoS Pathog *7*, e1002304. 10.1371/journal.ppat.1002304.

9.Kemenesi, G., Tóth, G.E., Mayora-Neto, M., Scott, S., Temperton, N., Wright, E., Mühlberger, E., Hume, A.J., Suder, E.L., Zana, B., et al. (2022). Isolation of infectious Lloviu virus from Schreiber's bats in Hungary. Nat Commun *13*, 1706. 10.1038/s41467-022-29298-1.

10.Yang, X.L., Tan, C.W., Anderson, D.E., Jiang, R.D., Li, B., Zhang, W., Zhu, Y., Lim, X.F., Zhou, P., Liu, X.L., et al. (2019). Characterization of a filovirus (Měnglà virus) from Rousettus bats in China. Nat Microbiol *4*, 390-395. 10.1038/s41564-018-0328-y.

11.Mehedi, M., Groseth, A., Feldmann, H., and Ebihara, H. (2011). Clinical aspects of Marburg hemorrhagic fever. Future Virol *6*, 1091-1106. 10.2217/fvl.11.79.

12.Beniac, D.R., and Booth, T.F. (2017). Structure of the Ebola virus glycoprotein spike within the virion envelope at 11 A resolution. Sci Rep *7*, 46374. 10.1038/srep46374.

13.Lee, J.E., and Saphire, E.O. (2009). Neutralizing ebolavirus: structural insights into the envelope glycoprotein and antibodies targeted against it. Curr Opin Struct Biol *19*, 408-417. 10.1016/j.sbi.2009.05.004.

14.Brauburger, K., Hume, A.J., Mühlberger, E., and Olejnik, J. (2012). Forty-five years of Marburg virus research. Viruses *4*, 1878-1927. 10.3390/v4101878.

15.Schafer, A., Xiong, R., Cooper, L., Nowar, R., Lee, H., Li, Y., Ramirez, B.E., Peet, N.P., Caffrey, M., Thatcher, G.R.J., et al. (2021). Evidence for distinct mechanisms of small molecule inhibitors of filovirus entry. PLoS Pathog *17*, e1009312. 10.1371/journal.ppat.1009312.

16.Hashiguchi, T., Fusco, M.L., Bornholdt, Z.A., Lee, J.E., Flyak, A.I., Matsuoka, R., Kohda, D., Yanagi, Y.,

Hammel, M., Jr, J.E.C., and Saphire, E.O. (2015). Structural basis for Marburg virus neutralization by a
cross-reactive human antibody. Cell *160*, 904-912. 10.1016/j.cell.2015.01.041.
17.Fusco, M.L., Hashiguchi, T., Cassan, R., Biggins, J.E., Murin, C.D., Warfield, K.L., Li, S., Holtsberg, F.W.,
Shulenin, S., Vu, H., et al. (2015). Protective mAbs and Cross-Reactive mAbs Raised by Immunization
with Engineered Marburg Virus GPs. PLoS Pathog *11*, e1005016. 10.1371/journal.ppat.1005016.
18.Gregory, S.M., Harada, E., Liang, B., Delos, S.E., White, J.M., and Tamm, L.K. (2011). Structure and
function of the complete internal fusion loop from Ebolavirus glycoprotein 2. Proc Natl Acad Sci U S A
108, 11211-11216. 10.1073/pnas.1104760108.
19.Lee, J., Nyenhuis, D.A., Nelson, E.A., Cafiso, D.S., White, J.M., and Tamm, L.K. (2017). Structure of
the Ebola virus envelope protein MPER/TM domain and its interaction with the fusion loop explains
their fusion activity. Proc Natl Acad Sci U S A *114*, E7987-E7996. 10.1073/pnas.1708052114.
20.Volchkov, V.E., Volchkova, V.A., Ströher, U., Becker, S., Dolnik, O., Cieplik, M., Garten, W., Klenk, H.D.,
and Feldmann, H. (2000). Proteolytic processing of Marburg virus glycoprotein. Virology *268*, 1-6.
10.1006/viro.1999.0110.
21.Dye, J.M., Herbert, A.S., Kuehne, A.I., Barth, J.F., Muhammad, M.A., Zak, S.E., Ortiz, R.A., Prugar, L.I.,
and Pratt, W.D. (2012). Postexposure antibody prophylaxis protects nonhuman primates from filovirus
disease. Proc Natl Acad Sci U S A *109*, 5034-5039. 10.1073/pnas.1200409109.
22.Aleksandrowicz, P., Marzi, A., Biedenkopf, N., Beimforde, N., Becker, S., Hoenen, T., Feldmann, H.,
and Schnittler, H.-J. (2011). Ebola virus enters host cells by macropinocytosis and clathrin-mediated
endocytosis. J Infect Dis *204 Suppl 3*, S957-967. 10.1093/infdis/jir326.
23.Saeed, M.F., Kolokoltsov, A.A., Albrecht, T., and Davey, R.A. (2010). Cellular entry of ebola virus
involves uptake by a macropinocytosis-like mechanism and subsequent trafficking through early and
late endosomes. PLoS Pathog *6*, e1001110. 10.1371/journal.ppat.1001110.
24.Carette, J.E., Raaben, M., Wong, A.C., Herbert, A.S., Obernosterer, G., Mulherkar, N., Kuehne, A.I.,
Kranzusch, P.J., Griffin, A.M., Ruthel, G., et al. (2011). Ebola virus entry requires the cholesterol
transporter Niemann-Pick C1. Nature *477*, 340-343. 10.1038/nature10348.
25.Brecher, M., Schornberg, K.L., Delos, S.E., Fusco, M.L., Saphire, E.O., and White, J.M. (2012).
Cathepsin cleavage potentiates the Ebola virus glycoprotein to undergo a subsequent fusion-relevant
conformational change. J Virol *86*, 364-372. 10.1128/JVI.05708-11.
26.Martinez, O., Johnson, J., Manicassamy, B., Rong, L., Olinger, G.G., Hensley, L.E., and Basler, C.F.
(2010). Zaire Ebola virus entry into human dendritic cells is insensitive to cathepsin L inhibition.
Cellular Microbiology *12*, 148-157. 10.1111/j.1462-5822.2009.01385.x.
27.Gnirss, K., Kühl, A., Karsten, C., Glowacka, I., Bertram, S., Kaup, F., Hofmann, H., and Pöhlmann, S.
(2012). Cathepsins B and L activate Ebola but not Marburg virus glycoproteins for efficient entry into
cell lines and macrophages independent of TMPRSS2 expression. Virology *424*, 3-10.
10.1016/j.virol.2011.11.031.
28.Misasi, J., Chandran, K., Yang, J.-Y., Considine, B., Filone, C.M., Côté, M., Sullivan, N., Fabozzi, G.,
Hensley, L., and Cunningham, J. (2012). Filoviruses require endosomal cysteine proteases for entry but
exhibit distinct protease preferences. J Virol *86*, 3284-3292. 10.1128/JVI.06346-11.
29.Hu, N., Qiao, C., Wang, J., Wang, Z., Li, X., Zhou, L., Wu, J., Zhang, D., Feng, J., Shen, B., et al. (2021).
Identification of a novel protective human monoclonal antibody, LXY8, that targets the key
neutralizing epitopes of staphylococcal enterotoxin B. Biochem Biophys Res Commun *549*, 120-127.
10.1016/j.bbrc.2021.02.057.
30.Wang, Z., Hu, N., Zhou, Y., Shi, N., Shen, B., Luo, L., and Feng, J. (2022). Structure-guided affinity

maturation of a novel human antibody targeting the SARS-CoV-2 nucleocapsid protein. Scientific Reports *12*, 8469. 10.1038/s41598-022-12242-0.

31.Dahms, N.M., Olson, L.J., and Kim, J.J. (2008). Strategies for carbohydrate recognition by the mannose 6-phosphate receptors. Glycobiology *18*, 664-678. 10.1093/glycob/cwn061.

32.Bohnsack, R.N., Song, X., Olson, L.J., Kudo, M., Gotschall, R.R., Canfield, W.M., Cummings, R.D., Smith, D.F., and Dahms, N.M. (2009). Cation-independent mannose 6-phosphate receptor: a composite of distinct phosphomannosyl binding sites. J Biol Chem *284*, 35215-35226. 10.1074/jbc.M109.056184.

33.Grubb, J.H., Vogler, C., and Sly, W.S. (2010). New strategies for enzyme replacement therapy for lysosomal storage diseases. Rejuvenation Res *13*, 229-236. 10.1089/rej.2009.0920.

34.Duan, Y., Luo, L., Qiao, C., Li, X., Wang, J., Liu, H., Zhou, T., Shen, B., Lv, M., and Feng, J. (2019). A novel human anti-AXL monoclonal antibody attenuates tumour cell migration. Scandinavian journal of immunology. 10.1111/sji.1277.

35.Liu, F., Guan, Z., Liu, Y., Li, J., Liu, C., Gao, Y., Ma, Y., Feng, J., Shen, B., and Yang, G. (2021). Identification of a Human Anti-Alpha-Toxin Monoclonal Antibody Against Staphylococcus aureus Infection. Frontiers in Microbiology *12*, 1875-. 10.3389/fmicb.2021.692279.

36.Markosyan, R.M., Miao, C., Zheng, Y.-M., Melikyan, G.B., Liu, S.-L., and Cohen, F.S. (2016). Induction of Cell-Cell Fusion by Ebola Virus Glycoprotein: Low pH Is Not a Trigger. PLoS Pathog *12*, e1005373. 10.1371/journal.ppat.1005373.

37.Shifflett, K., and Marzi, A. (2019). Marburg virus pathogenesis - differences and similarities in humans and animal models. Virol J *16*, 165. 10.1186/s12985-019-1272-z.

38.Wirchnianski, A.S., Wec, A.Z., Nyakatura, E.K., Herbert, A.S., Slough, M.M., Kuehne, A.I., Mittler, E., Jangra, R.K., Teruya, J., Dye, J.M., et al. (2021). Two Distinct Lysosomal Targeting Strategies Afford Trojan Horse Antibodies With Pan-Filovirus Activity. Frontiers in immunology *12*, 729851. 10.3389/fimmu.2021.729851.

39.Abir, M.H., Rahman, T., Das, A., Etu, S.N., Nafiz, I.H., Rakib, A., Mitra, S., Emran, T.B., Dhama, K., Islam, A., et al. (2022). Pathogenicity and virulence of Marburg virus. Virulence *13*, 609-633. 10.1080/21505594.2022.2054760.

40.Araf, Y., Maliha, S.T., Zhai, J., and Zheng, C. (2023). Marburg virus outbreak in 2022: a public health concern. Lancet Microbe *4*, e9. 10.1016/s2666-5247(22)00258-0.

41.Flyak, A.I., Ilinykh, P.A., Murin, C.D., Garron, T., Shen, X., Fusco, M.L., Hashiguchi, T., Bornholdt, Z.A., Slaughter, J.C., Sapparapu, G., et al. (2015). Mechanism of human antibody-mediated neutralization of Marburg virus. Cell *160*, 893-903. 10.1016/j.cell.2015.01.031.

42.Marzi, A., Haddock, E., Kajihara, M., Feldmann, H., and Takada, A. (2018). Monoclonal Antibody Cocktail Protects Hamsters From Lethal Marburg Virus Infection. J Infect Dis *218*, S662-S665. 10.1093/infdis/jiy235.

43.Kan, S.H., Troitskaya, L.A., Sinow, C.S., Haitz, K., Todd, A.K., Di Stefano, A., Le, S.Q., Dickson, P.I., and Tippin, B.L. (2014). Insulin-like growth factor II peptide fusion enables uptake and lysosomal delivery of α-N-acetylglucosaminidase to mucopolysaccharidosis type IIIB fibroblasts. Biochem J *458*, 281-289. 10.1042/bj20130845.

44.Aoyagi-Scharber, M., Crippen-Harmon, D., Lawrence, R., Vincelette, J., Yogalingam, G., Prill, H., Yip, B.K., Baridon, B., Vitelli, C., Lee, A., et al. (2017). Clearance of Heparan Sulfate and Attenuation of CNS Pathology by Intracerebroventricular BMN 250 in Sanfilippo Type B Mice. Mol Ther Methods Clin Dev

6, 43-53. 10.1016/j.omtm.2017.05.009.

45.Kan, S.H., Aoyagi-Scharber, M., Le, S.Q., Vincelette, J., Ohmi, K., Bullens, S., Wendt, D.J.,

Christianson, T.M., Tiger, P.M., Brown, J.R., et al. (2014). Delivery of an enzyme-IGFII fusion protein to

the mouse brain is therapeutic for mucopolysaccharidosis type IIIB. Proc Natl Acad Sci U S A *111*,

14870-14875. 10.1073/pnas.1416660111.

46.Berthe, M.L., Esslimani Sahla, M., Roger, P., Gleizes, M., Lemamy, G.J., Brouillet, J.P., and Rochefort,

H. (2003). Mannose-6-phosphate/insulin-like growth factor-II receptor expression levels during the

progression from normal human mammary tissue to invasive breast carcinomas. Eur J Cancer *39*,

635-642. 10.1016/s0959-8049(02)00627-5.

47.Jonson, T., Albrechtsson, E., Axelson, J., Heidenblad, M., Gorunova, L., Johansson, B., and Höglund,

M. (2001). Altered expression of TGFB receptors and mitogenic effects of TGFB in pancreatic

carcinomas. Int J Oncol *19*, 71-81.

48.Pavelić, K., Kolak, T., Kapitanović, S., Radosević, S., Spaventi, S., Kruslin, B., and Pavelić, J. (2003).

Gastric cancer: the role of insulin-like growth factor 2 (IGF 2) and its receptors (IGF 1R and M6-P/IGF

2R). J Pathol *201*, 430-438. 10.1002/path.1465.

49.Densumite, J., Phanthong, S., Seesuay, W., Sookrung, N., Chaisri, U., and Chaicumpa, W. (2021).

Engineered Human Monoclonal scFv to Receptor Binding Domain of Ebolavirus. Vaccines *9*.

10.3390/vaccines9050457.

 Figure legends

 Figure 1. Binding activity of mAb AF-03 to MARV GP and its epitopes.

(A) AF-03 and MARV GP proteins are examined by SDS-PAGE. NR, non-reducing;

R, reducing. (B) The binding activity of AF-03 to MARV GP is determined by ELISA.

The absorbance is detected at 450 nm. (C) The binding kinetics of AF-03 to MARV

GP is detected by BLI with a 180-s association step followed by a 220-s dissociation

step. Experiments are independently repeated at least three times, and the data from

one representative experiment is shown. (D) The 3-D ribbon structures of the AF-03

Fv fragment. The red ribbon denotes H-CDR1, the light blue denotes H-CDR2, the

pink denotes H-CDR3, the orange denotes L-CDR1, the deep blue denotes L-CDR2,

the purple denotes L-CDR3. (E) AF-03 and MARV GP complex derived from

theoretical modeling. The green ribbon denotes the orientation of the MARV GP

fragment, the yellow denotes AF-03 VLCDR, the pink denotes AF-03 VHCDR, the

deep blue denotes AF-03 VL and the red ribbon denotes AF-03 VH. (F) By molecular

docking analysis of van der Waals interaction, intermolecular hydrogen bonding,

polarity interaction and electrostatic interaction, the key amino acid residues of

MARV GP are screened.

Figure 2. AF-03 Epitope Identification.

(A) The neutralization activity of AF-03 or MR78 to mutated pseudovirus

($Q^{128}S$-$N^{129}S$, $Q^{204}A$-$T^{205}A$-$Q^{206}A$, $Y^{218}A$, $K^{222}A$, $C^{226}Y$) is evaluated in HEK293T

cells. The inhibition rate is analyzed. (B) The binding of AF-03 and MR78 to mutant

GP ($Q^{128}S$-$N^{129}S$ or $C^{226}Y$) is examined by ELISA respectively. (C) Secondary

structure of mutants and MARV GP is detected by CD. (D) The epitope overlapping

between AF-03 and MR78 is examined by the competition ELISA.

Figure 3. In vitro and in vivo neutralization of MARV pseudovirus infection by

AF-03.

(A) Pseudotypic MARV-Uganda is incubated with AF-03, MR78 or control mAb at

37°C for 1 h before infecting HEK293T cells (left) and Huh7 cells (right) respectively.

Luciferase is assayed and inhibition rates are calculated. (B) Pseudotypic

MARV-Angola, Musoke and Ravn infect HEK293T cells respectively and

neutralization activity of AF-03 to these species is determined. (C) AF-03 (10, 3,

1mg/kg) is administrated at 24 and 4 h before intraperitoneal injection of pseudovirus.

On day 5, bioluminescence signals are detected by an IVIS Lumina Series III imaging

system. (D) The average radiance value based on the luminescence of (C),*$p<0.05$,*

$*p<0.01$, $***p<0.001$.

Figure 4. The neutralization activity of AF-03 to EBOV, SUDV and BDBV

harboring cleaved GP.

Pseudotypic EBOV, SUDV, and BDBV are processed with thermolysin at 37°C.

Inhibition of these ebola virus infection harboring GP or GPcl by AF-03 is examined

by luciferase assay. *$p<0.05$, $**p<0.01$, $***p<0.001$*

Figure 5. Cellular internalization of AF03-NL.

(A) AF-03 or AF03-NL is incubated with cells at 4□ for 1h to prevent internalization

and then at 37□ for another 2 h to allow internalization. PE-conjugated secondary

antibody is added prior to analysis by flow cytometry. (B,C) pHrodo Red-labelled

AF-03 or AF03-NL is incubated with cells at 37□ for 1h and analyzed by flow

cytometry (B) and fluorescence microscopy (C) respectively. The red arrow denotes

internalized AF03-NL. Experiments are independently repeated at least three times,

and the data from one representative experiment is shown.

Figure 6. Pan-filovirus entry inhibition by AF03-NL.

(A,B) AF-03 or AF03-NL is incubated with HEK293T cells at 37°C for 2 h prior to

exposure to pseudotypic filovirus species (A) and EBOV mutants (B). Luciferase is

assayed and inhibition rates are calculated.

Figure 7. The requirement of CI-MPR for the neutralization activity of

AF03-NL.

(A) NPC2 protein is examined by SDS-PAGE. NR, non-reducing; R, reducing. (B)

AF03-NL, AF-03, NPC2 alone or equimolar combination of AF-03 and NPC2 is

incubated with HEK293T cells at 37°C for 2 h prior to exposure to pseudotypic

EBOV. Luciferase is assayed and inhibition rates are calculated. (C) HEK293T cells

are treated with siRNA-CI-MPR or negative control vector (NC) respectively and

CI-MPR expression is detected by flow cytometry. AF03-NL is incubated with

siCI-MPR or NC-treated HEK293T cells at 37°C for 2 h respectively prior to

exposure to pseudotypic EBOV. (D) CI-MPR is introduced into Huh7 cells and its

expression is detected by flow cytometry. AF03-NL is incubated with CI-MPR or

NC-knockin Huh7 cells at 37°C for 2 h respectively prior to exposure to pseudotypic

EBOV. Luciferase is assayed and inhibition rates are calculated.

A

kDa   M   AF-03   GP
           NR   R   NR   R

B

EC$_{50}$: 0.16μg/ml

OD450 (nm) vs AF-03(μg/ml)

C

K$_D$<1pM

nm vs Time (sec)

D

AF-03_VH   AF-03_VL
H_CDR2   L_CDR1   L_CDR2
H_CDR1   L_CDR3
H_CDR3   AF-03_FV

E

AF-03_VH
AF-03_VH_CDR
AF-03_VL
AF-03_VL_CDR
GP Modeling

F

AF-03_VH_CDR
T204
K222
T204
AF-03_VL_CDR

A

MOCK   AF-03   AF03-NL

0.66%   8.44%   41.3%

rate(%) 0.66   rate(%) 8.44   rate(%) 41.2

4°C

7.86%   18.8%

rate(%) 7.86   rate(%) 18.8

37°C

PE

B

37°C

MOCK   AF-03   AF03-NL

3.15%   2.46%   37.1%

rate(%) 3.15   rate(%) 2.46   rate(%) 37.1

pHrodo Red

C

37°C

MOCK   AF-03   AF03-NL

20 μm   20 μm   20 μm

A

B

C

D