Elevated temperature fatally disrupts nuclear divisions in the early *Drosophila* embryo.

Girish Kale[1,*], Pratika Agarwal[1], J Jaime Diaz-Larrosa[1], Steffen Lemke[1,2,*]

[1] Centre for Organismal Studies, University of Heidelberg, Heidelberg. Germany
[2] Institute of Biology, University of Hohenheim, Stuttgart, Germany
* Correspondance: G.K. (girish.kale@cos.uni-heidelberg.de), S.L. (steffen.lemke@uni-hohenheim.de)

Abstract:
Temperature fluctuations can challenge animal survival, especially in embryos of animals that can't
actively regulate their body temperature. Previous work in insects suggests that spatial gradients
and temporal variations in temperature can affect embryonic development in a region-specific
manner. Yet, it is unclear if elevated temperature produces developmental defects, and if so, how
the disruption may manifest. Here we use early embryonic development of the fruit fly *Drosophila*
melanogaster as a model to address consequences of elevated temperature on fly embryonic
development.

We identified the syncytial blastoderm stage as a critical time window as being particularly sensitive
to elevated temperature. During the syncytial blastoderm stage involves 4 rounds of meta-
synchronous nuclear divisions at the embryo cortex, followed by cellularization to form the cellular
blastoderm. Live imaging these processes at elevated temperature revealed an increase in mitotic
failures and a loss of cortical nuclei, both, during syncytial- and cellular-blastoderm stage.
Additionally, embryos developing at elevated temperature show a local crowding of nuclei and an
increase in asynchrony between the nuclear cycles in the middle vs anterior/posterior poles.
Interestingly, nuclear crowding and nuclear cycle asynchrony cooperatively amplify the frequency of
mitotic failures, leading to holes in the blastoderm epithelium. Strikingly, the exposure to elevated
temperature during this developmental stage also increased embryonic lethality. We propose that
the loss of nuclei results in a loss of patterning information and the abrogation of proper embryonic
development. We identified genes that regulate mitosis, especially the ones mediating the
interaction between microtubules and F-actin, and functionally tested their capacity to rescue the
embryonic lethality.

Our results expose a vulnerability at elevated temperatures in the interaction between cortical actin
and microtubules, which leads to mitotic failures, and can be potentially rescued by modulating the
expression of just a few factors. We propose that the expression levels of the corresponding genes
could be used as indicators to predict fitness effects on insect populations that are exposed to
increasing temperature variations.

Introduction:

Temperature is a key factor determining the fidelity of embryonic development. Studies have shown occurrence of birth defects due to hyperthermia during pregnancy (Edwards 2006; Miller and Church 2013; Graham 2020), in spite of active regulation of core temperature. Here, the cell biological basis for the developmental defects due to an exposure to high temperature, remains elusive. It is currently unclear whether this phenomenon could be observed and studied also in invertebrate species, for which the developmental process is well described, allowing for a quantitative characterization of the potential developmental defects arising due to high temperature. Here we explore whether the exposure to high temperature is detrimental to the development in the embryos of *Drosophila melanogaster* (henceforth, *Drosophila*), addressing the cell biological basis for possible defects, and asking what are the consequences in terms of population level parameters, such as embryo survival.

Previous studies have shown that the early embryonic development of *Drosophila* embryos (first ~6 hours) is especially sensitive to acute heat shock (Welte et al. 1993), and have indicated that acute heat shock causes a loss of coordination in developmental processes to induce developmental defects (Crews, McCleery, and Hutson 2016). Given that the survival after a heat shock relies on transcription of heat-shock proteins (Graziosi et al. 1980), and that zygotic transcription is progressively activated during maternal-to-zygotic transition (Pritchard and Schubiger 1996), it is likely that the developmental processes prior to the completion of this transition are especially susceptible to heat shock. In line with these expectations, previous studies have demonstrated a disruption of the normal dynamics of cytoskeletal elements due to an exposure to high temperature (Walter, Petersen, and Biessmann 1990; Zheng et al. 2013; Figard et al. 2019). Even more elusive are the effects of prolonged exposure to elevated temperature below the heat shock limits. Here we focus on such temperature sensitivity of early embryonic development in *Drosophila*, and ask what is the cell biological basis for developmental defects due to an exposure to elevated temperature, and whether there is a link to the population level consequences on embryo survival.

Post fertilization, the *Drosophila* embryos undergo 9 rounds of syncytial nuclear divisions deep in the yolk. Following these divisions, most nuclei migrate to embryo periphery, where 4 more rounds of nuclear division take place during a stage called, syncytial blastoderm (Foe and Alberts 1983). These nuclear divisions are facilitated by pseudo-cleavage furrows, i.e., temporary membrane invaginations resembling a half-cell that slowly form around the nuclei during interphase-metaphase, and are then rapidly reabsorbed into the embryo cortex during anaphase-telophase (Xie and Blankenship 2018). Following these 13 rounds of nuclear divisions, the resulting ~6000 nuclei simultaneously segregate into individual cells, in the process of cellularization, producing the blastoderm epithelium (Mazumdar and Mazumdar 2002). At this point gastrulation starts, which involves cell shape changes (Martin 2020), cell contact rearrangements (Lecuit and Yap 2015), and further rounds of cell divisions in isolated mitotic domains (Foe 1989): all facilitating the early morphogenesis. Cell biological basis for these early developmental processes is well described, and would allow us to identify the defects at elevated temperature, if any.

In our experiments, we exposed *Drosophila* embryos to the elevated temperature, and compared them with normal temperature controls. Following previous studies, we chose 29°C as elevated temperature (Kuntz and Eisen 2014; Chong, Amourda, and Saunders 2018) and asked whether increased temperature up to 29°C resulted in developmental defects. Notably, this temperature is well below the 32°C and higher range required to elicit a robust heat shock response (Vazquez, Pauli,

and Tissières 1993), and no major developmental defects have been reported so far at 29°C. We
found that exposing embryos to elevated temperature during early embryonic development leads to
a significant increase in the population level embryonic lethality. When we traced defects from the
organism to the cellular level, we observed a loss of nuclei during syncytial- and cellular-blastoderm
stage, which was caused by mitotic failures. Additionally, in embryos developing at elevated
temperature, we observed and quantified the nuclear crowding in the central region of the embryo,
as well as greater asynchrony in the nuclear division cycles between central regions vs
anterior/posterior poles of the embryo. What's more, we could show that the nuclear crowding and
nuclear cycle asynchrony combinatorially cooperate to locally increase the propensity of mitotic
failures. Such local increase in mitotic failures results in holes in the blastoderm, i.e., region devoid
of nuclei/cells, to disrupt the continuity of this primary epithelial tissue. We overexpressed a few
candidate genes to boost the robustness of nuclear divisions to rescue the mitotic failures defect,
focusing on the regulators of cell polarity, microtubule- F-actin interactions, and mitotic spindle
orientation, to test their ability to rescue the described defects, and ameliorate the embryo lethality
at elevated temperature. Our results suggest a vulnerability in microtubule- F-actin interaction
during the well conserved process of nuclear division, when occurring at elevated temperature. Our
work establishes a novel link between cell biological defects due to an exposure to elevated
temperature, and a reduction in organismal fitness at population level at elevated temperature.

Results:

Early exposure to elevated temperature reduces embryo survival.
To test whether elevated temperature had an effect on developmental robustness during the first
hours of embryonic development, we exposed young embryos to 29°C and asked whether this
would alter the fraction of embryos that hatch into larvae. To associate possible fitness effects with
defined developmental windows (see above), embryos were exposed to elevated temperature
either from deposition till before the onset of gastrulation (~3 hours) or for a duration of ~3 hours
from the onset of gastrulation (Figure 1A), to keep the temperature exposures comparable. In both
cases we observed a significant increase in embryonic lethality (Figure 1B). Our findings reveal a
particular vulnerability during the syncytial phase of embryonic fly development and suggest that
elevated temperature leads to defects in the setup of the insect blastoderm. This is surprising
because previous studies indicated that genetic patterning and setting up of positional information
within the blastoderm was overall robust against temperature fluctuations (Lucchetta et al. 2005;
Niemuth and Wolf 1995).

In addition to genetic patterning, nuclear dynamics could be challenged during the first hours of
embryonic development. In this developmental window, nuclei divide deep in the yolk, then
translocate to the embryo cortex, where they need to be anchored robustly while undergoing
further rounds of nuclear division. A normal distribution of nuclei ultimately determines a normal
morphogenesis of the blastoderm epithelium. Thus, to address whether defects in the blastoderm
could stem from defects in nuclear dynamics, we analyzed behavior of syncytial nuclei as they
translocated to the egg cortex and divided. To follow nuclear dynamics during blastoderm
formation, we used Histone-RFP to visualize nuclei, and Jupiter-GFP to visualize microtubules,
keeping track of nuclear division cycles. Starting with nuclear migration from the yolk to periphery
(occurring during the interphase of the 10th nuclear cycle), we observed nuclei through their
peripheral divisions during syncytial blastoderm (mitoses 10 through 13), and then through
cellularization (interphase 14), until the onset of gastrulation could be inferred from the movement

and division of nuclei (beyond interphase 14). During this time window, in embryos developing at
elevated temperature, we did not observe any significant defects in nuclear distribution, as they
translocated normally from deep inside the yolk to embryo cortex. However, embryos that
developed at 29°C much more frequently "lost" nuclei from their cortical layer than if they
developed at 25°C, i.e., nuclei that were initially positioned within the cortical layer of the egg,
appeared to get expelled, descended into the yolk, and did not become part of the embryonic
blastoderm (Figure 1C-E), with a variability in the extent of the nuclear loss, producing holes in the
blastoderm of a fraction of embryos (Figure S1). Further, in the embryos fixed and stained with
DRAQ5, we imaged the nuclear distribution in the entire embryo volume, to observe that the loss of
nuclei wasn't symmetric in the two halves of the embryos, and albeit in the central region of the
embryos, the location of the nuclear loss varied across embryos (Figure S2). The embryo-to-embryo
variability, both in the distribution and intensity of nuclear loss argued for the possibility that these
defects were self-organized rather than dictated by the genetic make-up.

Taken together, we identified the syncytial blastoderm stage of embryo development as the most
sensitive window, based on the embryo lethality analysis. Additionally, the loss of nuclei during this
stage indicated a defect in nuclear anchorage at the embryo cortex, rather than nuclear divisions
deep in the yolk, or the translocation of nuclei from deep inside the yolk to the embryo cortex.

A closer observation of live imaged embryos revealed three striking phenomena during blastoderm
formation in embryos developing at elevated temperature compared to normal temperature
controls: an extensive loss of nuclei from the egg cortex (as above), a time shift and slowdown of
nuclear divisions cycles in the central region of the embryo compared to its poles, and nuclear
crowding in the central region of the embryo. Thus, we decided to quantitatively describe these
defects, in an attempt to describe the necessary conditions to produce a blastoderm hole, and link
these defects to the embryo lethality phenotype.

At elevated temperature, embryos lose nuclei from the cortex after defective divisions.
A closer look at the loss of cortical nuclei indicated that the nuclei were lost either individually or as
pairs of two adjacent nuclei. We could also observe periods of drastic displacement of nuclei at
elevated temperature (see below). Based on these observations, we imagined various mechanisms
for the observed loss of nuclei. For instance, individual nuclei might be lost due to out-of-plane
nuclear divisions, while a pair of nuclei might be lost due to a defective in-plane nuclear division. In
case of defects in nuclear divisions, it is possible that the cell cycle checkpoints are triggered. Nuclei
being dislodged from the embryo cortex during the drastic displacements could also explain both
individual and pairwise loss, though in such scenario, whether the pairs of nuclei need to be
somehow associated with each other, influencing each other's loss. Here we considered these
possibilities, and asked to what extent do they account for the observed loss of nuclei.

To address whether cortical nuclei appeared as getting lost because of an increased frequency of
out-of-plane divisions, we visually tracked nuclei from their last division to the time they were no
longer associated with the cortex. We could detect out-of-plane divisions in embryos at
temperatures ranging from 22 to 29°C, only during the mitoses of the 10th nuclear cycle. These out-
of-plane divisions could be easily identified, as they formed a characteristic mitotic spindle that was
oriented perpendicular to the embryo surface (Figure S3A, B), and after the division only one of the
daughter nuclei remained attached to the cortex. However, the overall frequency of out-of-plane
divisions was very low (<5%): neither did it differ significantly between any of the observed

temperatures (Figure S3C), nor could it explain the loss of nuclei during later nuclear cycles.
Similarly, losses of individual nuclei after an in-plane division were rather infrequent, and did not
differ significantly within the range of observed temperatures (Figure S3D, E). This suggested that
the increased loss of nuclei observed in embryos developing at 29°C could neither be accounted for
with individual nuclei dropping into the yolk, irrespective of whether the loss was due to them being
mechanically dislodged.

For the observed pairwise loss of nuclei, our tracking revealed that these were sister nuclei born
from the same mitosis. This suggested that specific characteristics of cortical nuclear divisions might
determine the future fate of the newborn nuclei and trigger their release from the egg cortex,
possibly in response to failed checkpoints during mitosis. In all of the pairwise loss of nuclei, we
observed chromosome segregation defects (Figure 1F) that could lead to aneuploidy, suggesting that
the subsequent expulsion of daughter nuclei from the egg cortex was caused by a defective mitosis.
Following defective mitoses either at normal or elevated temperature, we could not detect a clear
time course of nuclear expulsion as the corresponding nuclei could be lost in the immediately
following or in later interphases (Figure S4), suggesting that nuclear expulsion after defective mitosis
was a passive process and not triggered by molecular signaling at failed checkpoints.

To confirm that the observed loss of nuclei at 29°C could be attributed to an overall increase in
defective mitoses, we asked if the frequency of defective mitoses was changing depending on the
temperature. To address this question, we focused on interphase 14, as that's when we observed
the most nuclear losses, and counted the pairwise loss of nuclei after mitotic failures at 22, 25, 28,
and 29°C (see methods). While the frequency of mitotic failures was overall low at 22 to 28°C, it
increased significantly at 29°C (Figure 1G). Taken together, our findings identify nuclear divisions in
the blastoderm as a thermolabile cellular process (Figure 1H) that shows a disproportionate increase
in failures at the transition from 28°C to 29°C, resulting in the loss of defectively born sister nuclei
from the egg cortex. Interestingly, in a fraction of embryos the mitotic failures occurred in great
numbers and in close vicinity, leading to the holes in the blastoderm epithelium (Figure 1G, hollow
markers). Thus, the formation of blastoderm holes (Figure S1) could be directly attributed to
clustered mitotic failures, though it remained unclear why some embryos have the aggregates of
mitotic failures while others didn't.

To explore the potential cause of mitotic failures at elevated temperatures, we tracked nuclear
clusters starting from individual mother nucleus (end of interphase 10) to the 8 great-granddaughter
nuclei (end of interphase 13), in order to quantitatively describe their movement (Figure S5A-C). This
allowed us to assess mechanical stresses that would be experienced by nuclei dividing at the
blastoderm cortex, as we distinguished the movement of nuclear clusters during interphase-
metaphase vs. anaphase-telophase. Our results showed that the nuclear clusters had 2 components
to their movement. First, the clusters exhibited a constant drift-like movement, which we captured
using a linear fit to the displacement of the center of the nuclear clusters over time (Figure S5B
dotted lines). The slope of the linear fit indicated that the clusters drifted towards posterior (Figure
S5D, E). The second component was composed of deviations in the centers of nuclear cluster from
the linear fit, i.e., the displacements relative to the first component. These deviations were
significantly greater at elevated temperature, both during the interphase-metaphase period (Figure
S5F, G) and during the anaphase-telophase period (Figure S5H, I), though the displacements along
AP-axis were more drastic than those along DV-axis, and those during anaphase-telophase were 2-3-
fold greater than those during interphase-metaphase. For context, the length scale of the

displacements during anaphase-telophase could be up to 4-times the length of the spindle itself,
indicating an extensive propensity for displacement in x-y plane, especially along the AP axis. We
also observed that the spindles were getting displaced in z-axis, again, during anaphase-telophase
period. In embryos developing at 29°C, we could see two waves of nuclear displacement, originating
from the anterior and the posterior pole, meeting at the central region of the embryo (Figure S6).
Overall, these quantifications and observations indicated that the mitotic spindles are displaced
drastically while nuclei are dividing, and argued for the possibility of mechanical defects during
nuclear divisions.

Taken together, we observed the effects of exposure to elevated temperature, at organismal level as
embryo lethality (Figure 1B), at tissue level as blastoderm holes (Figure 1E, Figure S1, S2), and at
cellular level as mitotic failures (Figure 1F, G). Embryos with most severe form of mitotic failure
phenotype had blastoderm hole, while the fraction of embryos with large blastoderm holes
correlated well with increase in embryo lethality, indicting a link from cellular till organismal
phenotype. Given that the loss of nuclei occurs after mitosis, and that the nuclei move drastically
during anaphase-telophase period, we further argue that nuclear anchorage is specifically
challenged during mitosis.

Nuclei divide with a time shift and are crowded in embryos at elevated temperature.
The most severe loss of nuclei was often observed in the central domain of the embryo (Figure 1C-E,
Figure S1 and S2). Notably, these areas coincided with the "collision" of two waves of nuclear
displacements along z-axis (Figure S6), i.e., the future apical-basal axis in the blastoderm cells (A.
Schmidt and Grosshans 2018). We could further observe that, not only anaphase, but all other
phases of nuclear division were delayed in the central region of the embryos developing at 29°C. This
resulted in regions of embryo being segregated into domains of nuclear division phases (Figure 2A).
Conversely, the same phase of nuclear division cycle could be observed at different timepoints in
different regions of the embryo (Figure S7). Thus, we decided to quantify this asynchrony of nuclear
division cycles as a function of temperature (see methods, also Figure S7). Our quantifications show
that the asynchrony significantly increases in embryos developing at 29°C as compared to other
temperatures (Figure 2B). Thus, we quantitatively describe the defect of asynchrony in nuclear
cycles at elevated temperature (Figure 2C).

To quantify nuclear crowding, we focused on the last nuclear division cycle and its preceding
interphase (interphase 13, see methods). To quantify the space available for cortical blastoderm
nuclei, we used Voronoi tessellation and measured distances between nuclei using pseudo-cell area.
The distances between nuclei overall were rather homogeneous in embryos developing at 22°C
(Figure 2D, H, L). Already at 25°C (Figure 2E, I, M) and 28°C (Figure 2F, J, n), the available space for
cortical nuclei was no longer uniform across the embryo, and it showed signs of local crowding. At
29°C, the space available for nuclei along the anterior-posterior axis varied substantially and
revealed strong nuclear crowding in the central region of the embryo (Figure 2G, K, O). We
quantified the extent of crowding (see methods) as a function of temperature, and found a
progressive increase in the nuclear crowding (Figure 2P). Thus, we also quantitatively describe the
defect of nuclear crowding at elevated temperature (Figure 2Q).

Taken together, we have two additional defects at elevated temperature, nuclear crowding and
nuclear cycle asynchrony, besides mitotic failures. We also note that the severity of these defects
(based on the quantifications) can vary between embryos, e.g., the embryos with most extensive

failure in mitosis (blastoderm holes) don't have the highest levels of nuclear crowding (Figure 2P), or
strongest asynchrony (Figure 2B), etc. Though, based on the quantifications and statistical
significances we note that; one, severe mitotic failures (blastoderm holes) appear in a threshold-
effect-like manner, appearing only in a fraction of embryos at 29°C; two, the nuclear crowding and
asynchrony in nuclear cycles appears progressively as function of temperature, although the
progression is not linear; and three, the fraction of embryos with large blastoderm holes best
explains the increase in embryo lethality. Thus, we hypothesize a hierarchy of the defects, purely
based on the fraction of embryos in which they appear as a function of temperature: exposure to
elevated temperature will increase the frequency of mitotic failures as primary defect; nuclear
crowding and nuclear cycle asynchrony will secondarily dictate the distribution of mitotic failures,
producing extensive mitotic failures (blastoderm holes) as the ultimate effect in a fraction of
embryo; eventually leading to a reduction in embryo fitness and an increase in embryo lethality.

Regions of embryo with greater nuclear crowding and with delayed nuclear cycles are associated
with greater frequency of mitotic failures.
To test whether nuclear crowding and nuclear cycle asynchrony could influence the propensity of
mitotic failures, we decided to compute the fraction of mitotic failures occurring in the regions of
the embryo that had nuclear crowding and/or delayed nuclear division cycles. We constructed
regions of the embryo with maximum nuclear crowding (crowded regions), maximum delay in
nuclear division cycles (delayed regions), and a combination of crowded and delayed regions
(combined region) (see methods).

We defined the crowded regions (Figure 3A), and delayed regions (Figure 3B) based of the
distribution and dynamics of nuclei in live imaged embryos. Then, we overlayed the map of mitotic
failures on the crowded regions (Figure 3C), delayed regions (Figure 3D), and the combined regions
(Figure 3E). The crowded and delayed regions had rather comparable areas across temperatures
(Figure 3F, G), indicating that any difference in the fraction of mitotic defects would be solely due to
the nuclear crowding or due to the delays in nuclear cycles. Interestingly, the areas of the combined
regions were smaller at elevated temperature (Figure 3H), indicating that there was an increase in
the overlap between the crowded and delayed regions, and that we should expect a similar
reduction in the fraction of mitotic failures, if the mitotic failures were distributed randomly.

The fraction of mitotic failures in crowded region increased progressively as function of
temperature, reaching statistical significance for comparison between 22 and 29°C (Figure 3I), while
the fraction of mitotic failures in delayed regions did not increase progressively, showing an abrupt
increase from 28 to 29°C with only marginal statistical significance (Figure 3J). Interestingly, we see
that the smaller area fractions in the combined regions correspond to even greater fraction of
mitotic failures (Figure 3K). Additionally, the distribution of the fraction of mitotic failures in the
combined region was unimodal, in contrast to the bimodal distribution of mitotic failures themselves
(Figure 1G). Thus, we conclude that the combinatorial effect of nuclear crowding and asynchrony
significantly increases the propensity of mitotic failures. We would further argue that nuclear
crowding has a stronger impact on the propensity of mitotic failures than that due to nuclear cycle
asynchrony.

Taken together, we find large clusters of defective mitosis and subsequent loss of nuclei in the
center of about 10-20% of embryos developing at 29°C (Figure 1G). Nuclear crowding and nuclear
cycle asynchrony cooperate to produce "hot spots" of mitotic failures (Figure 3). The aggregated

losses of nuclei in the center of the embryo persist until the end of blastoderm formation and leads
to visible holes in the newly formed embryonic epithelium (Figure S1). These holes indicate a severe
disruption in epithelial integrity and approximately match the increased rate of embryonic lethality
for development at 29°C (Figure 1B). Going further, we attempted to rescue the embryo lethality
defect using over-expression of various candidate genes, and if a rescue was possible, further
decided to quantify whether the rescue could be explained through fraction of mitotic failures in the
crowded region and the combined region.

Identifying temperature-sensitive vulnerabilities in a deeply conserved cellular process
The analysis of early fly development at moderately elevated temperatures, in our experiments, has
uncovered the first three hours of fly development as sensitive to environmental perturbations,
which can affect embryo fitness and increase embryonic lethality at elevated temperatures. This
increase in embryonic lethality correlates with specific perturbations in fundamental nuclear
dynamics, i.e., the timing and fidelity of nuclear divisions within the spatial constraints of a cortex
densely packed with nuclei. To address how a deeply conserved and presumably robust process like
mitosis could be affected so dramatically at a temperature lower than those eliciting a heat-shock
response, we functionally probed for possible vulnerabilities in core molecular interactions that
control nuclear dynamics in the early fly embryo.

In stark deviation from canonical cytokinesis, cortical nuclear divisions in the syncytial fly embryo
rely on cell polarity cues that are set up at shallow metaphase furrows of the egg cortex, where they
orient mitotic spindles parallel to the cell surface (A. Schmidt and Grosshans 2018; Xie and
Blankenship 2018). What's more, these furrows are transient, and are reestablished during each
nuclear cycle: including new growth of membrane between nuclei, reestablishment of cell polarity,
along with reorganization of cytoskeletal elements. The dependence on transient membrane
invaginations to establish cell polarity information represents the least conserved and potentially
most labile element of cortical nuclear divisions in insect embryos. To test whether the fidelity of
cortical mitoses at elevated temperatures hinges upon insufficiently robust cell polarity signaling, we
overexpressed two known prominent polarity proteins, Bazooka (baz, a Par-3 homolog) and Disks-
large (dlg) (Harris and Peifer 2004), and asked whether we rescue embryo lethality defect, and the
increase in the fraction of mitotic failures in crowded regions and in combined regions with
combinatorial effect of nuclear crowding and asynchrony.

When we over-expressed either *baz* or *dlg* we did not see a significant change in embryo lethality at
control temperature (25°C) as compared to the control (Figure 4A). The lethality was greater in
outcross controls at elevated temperature (29°C) similar to results in previous experiments, while
the embryos over-expressing either *baz* or *dlg* faired significantly better at the elevated temperature
(Figure 4A). We note however that the rescue of lethality at elevated temperature in *baz* over-
expressing embryos was only marginally significant, with a lot of variability between samples.
Additionally, *dlg* over-expressing embryos at normal temperature showed higher variability in
embryo lethality than their outcross control counterparts. Consistent with these results, *baz* over-
expression did not rescue the increase in the fraction of mitotic failures, neither in the crowded
regions, nor in the combined regions at elevated temperature (Figure 4B, C), though the large
blastoderm holes were absent in these embryos. Then, *dlg* over-expression rescued the increase in
fraction of mitotic failures only in the crowded region at elevated temperature (Figure 4B, C), though
the large blastoderm holes were absent in these embryos. Additionally, *dlg* over-expression

increased the fraction of mitotic defects at normal temperature in both, the crowded region and in
the combined region, occasionally producing small blastoderm holes (Figure 4B, C).

Above results indicated a temperature-dependent vulnerability more specific to the molecular core
of the mitotic spindle rather than its setup in the context of a syncytial embryo. To faithfully pull
sister chromatids towards the actin cortex on either side of the division plane, the execution of
mitosis relies on tightly regulated interactions between the actin cortex and the microtubules of the
mitotic spindle (Cao et al. 2010; Rizzelli et al. 2020). In Drosophila, these critical interactions of actin
and microtubules have been associated with a complex of three proteins: α-Catenin (αCat), which
binds to the actin cortex; β-Catenin (βCat, or armadillo in Drosophila), which binds to α-Catenin; and
Adenomatous polyposis coli 2 (Apc2), which links to β-Catenin and binds the microtubules of the
mitotic spindle (McCartney et al. 2001). Within this tripartite complex, the interaction of αCat and
βCat is direct and presumably non-regulated, whereas the interaction of βCat and Apc2 is mediated
by a specific kinase, the Drosophila homolog of the Glycogen Synthase Kinase 3β, Shaggy (sgg) (Rim,
Clevers, and Nusse 2022). Additional indication for the activity of the complex came when we fixed
and stained embryos developing at elevated temperature, to visualize nuclei and cortical F-actin (see
methods). We could observe F-actin structures in embryos at elevated temperature (Figure S9, see
panel F), which were reminiscent of those in embryos mutant for *Apc2*, β*Cat*, or *sgg*, as previously
described (McCartney et al. 2001). Thus, we hypothesized that boosting the activity of the complex
through over-expression of one or more of its components should meaningfully rescue the embryo
lethality defect, and the increase in the fraction of mitotic failures in crowded regions and in
combined regions.

To test whether the fidelity of cortical mitoses at elevated temperatures was affected by
insufficiently tight or poorly controlled coupling of actin and microtubules, we overexpressed either
αCat or *sgg*, and again asked whether the increase in embryo lethality at elevated temperature was
rescued. When we over-expressed either *αCat* or *sgg* we did not see a significant change in embryo
lethality at control temperature (25°C) as compared to the control (Figure 4D). The lethality was
greater in outcross controls at elevated temperature (29°C) similar to results in previous
experiments, while the embryos over-expressing either *αCat* or *sgg* faired significantly better at the
elevated temperature (Figure 4D). Consistent with these results, both *αCat* and *sgg* over-expressions
rescued the increase in the fraction of mitotic failures, in the crowded regions, as well as in the
combined regions at elevated temperature (Figure 4E, F), with the large blastoderm holes absent in
these embryos. These results suggested that an experimental strengthening of actin binding was
able to attenuate temperature-dependent destabilizations of the mitotic spindle. These findings are
consistent with the idea that the scaffolding activity of αCat, βCat, and Apc2, mediated by sgg,
constitutes a thermosensitive controller of nuclear division fidelity in the early blastoderm embryo.

Given the reduction in mitotic fidelity at elevated temperature, we were wondering whether any
regulator of mitosis could rescue temperature defects upon overexpression. To test this possibility,
we tested other regulators of mitosis and focused on the activity of mud and insc. In epithelia, the
activity of 'Mushroom body defect' (mud, NuMA homolog) and that of Inscuteable (insc) is known to
dictate the orientation of mitotic spindle during cell division (Mapelli and Gonzalez 2012). Similar
activity might be present for nuclear divisions during syncytial blastoderm stage, raising the
possibility that over-expression of *mud* or *insc* could rescue embryo lethality, presumably through
reducing mitotic failures. However, neither of the over-expressions ameliorated the increased
embryo lethality at elevated temperature, with mud over-expression additionally increasing the

lethality at normal temperature as well (Figure S11). This failure to rescue embryo lethality further
allowed us to argue that, the elevated temperature critically challenges the core machinery of
mitotic spindle, more so than the regulation of spindle orientation.

The natural distribution of a few regulators of spindle dynamics varies along thermoclines.
Our analyses show how elevated temperatures during the first hours of fly development affect early
embryonic cell biology; it may seem suggestive, but it was not shown that naturally occurring
changes similar to our overexpression of cytoskeletal regulators could shape the evolution of fly or
insect development in general. To address this possibility more directly, we analyzed the distribution
of standing genetic variations associated with the regulation of the spindle dynamics (combination
of GO terms, cell polarity, spindle organization, and spindle localization), and tested the hypothesis
that such variation was randomly distributed with respect to described thermoclines, i.e., continent-
scale gradients of annual mean temperature. We found that indeed, most standing genetic variation
associated with putative spindle dynamics regulators was distributed without a clear thermoclinal
pattern, although the fraction of genes showing clinal distribution was greater than those associated
with a more generic GO term like 'metabolic process' (Figure 4G, Table S1, S2). A clear and notable
outlier in this analysis was *sgg*, which was marked with a north/south clinal distribution of natural
variations independently in populations in the US, in Europe, and in Australia (see also, Table S3).
These findings strongly suggest that the thermosensitivity that *sgg* exhibits in lab conditions,
constitutes a biological property that can be exposed and selected for also in natural populations
and during the course of evolution. Similarly, we found that 'scribble', a cell polarity protein which
interacts with dlg (Khoury and Bilder 2020; 2022), was also clinal in all 3 populations. Further
experiments are required to formally test the possibility of a rescue of embryo lethality with over-
expression of *scribble*, which will be addressed in future experiments.

Discussion:

Cellular basis of the sensitivity against elevated temperature during early embryonic development
With our work, we demonstrate how the first hours of embryonic development are vulnerable to an
exposure to a non-heat-shock temperature. We identify the mechanistic basis for this vulnerability:
it is the interaction between microtubules and F-actin during mitosis is thermolabile. Given that
these early processes till blastoderm formation are conserved across insects (Lynch, El-Sherif, and
Brown 2012; Lemke, Kale, and Urbansky 2020), our results implicate a vulnerability in a plethora of
insect species. It would be interesting to investigate whether mechanisms exist in nature to
accommodate for this vulnerability. In the backdrop of this sensitivity, it is the challenge for the
embryo to establish an egg cortex that is uniformly populated with cortically anchored nuclei that
carry positional information for subsequent steps of epithelial morphogenesis and cell
differentiation. A defect in these early morphogenetic processes and cell differentiation could lead
to defects in subsequent processes, ultimately leading to developmental failure and embryo
lethality. Thus, with our data, a mechanism emerges to establish the link between the proximate
effects of elevated temperature on cytoskeletal regulation, and its ultimate impact on the survival of
embryos.

What is the best predictor of lethality in embryos exposed to elevated temperatures? In our analysis,
we found 3 defects in embryos at elevated temperature: mitotic failures, nuclear crowding, and
nuclear cycle asynchrony. Mitotic failures, especially during the last mitosis prior to cellularization,
can lead to loss of nuclei, and consequently a loss of local positional information that is advected

and lost with the nuclei. This can lead to missing cell fates in the coordinate system of the embryo,
leading to missing embryonic structures, which can pose challenges for developmental fidelity.
Comparing the fraction of dead embryos at elevated temperature vs. the severity of mitotic failures,
we would argue that the embryos with the most severe mitotic failures, i.e., those that end up with
holes in blastoderm epithelium, are the ones that don't hatch into larvae. In our over-expression
conditions, it is clear that the severe mitotic failures (blastoderm holes) are absent in the genotypes
that rescue the embryo lethality at elevated temperature. It remains to be seen if an extensive, but
scattered, loss of nuclei poses any developmental disadvantage to the embryo or not. It is possible
that such disadvantages might be apparent only during larval stage, and might directly affect larval
fitness, but our current analysis wouldn't capture that effect. Additionally, developmental
robustness can vary between species, which raises the possibility that not all fly species will be able
to accommodate such developmental disadvantages. Further, here we have presented 28 embryos
of control genotype at elevated temperature, of which 6 have large- and 5 have small blastoderm
holes. In all of these cases, the blastoderm holes appear in the region where we see an overlap
between nuclear crowding and the delay of nuclear divisions. Thus, we further argue that the severe
version of the defect at elevated temperature (blastoderm holes) occur where the three phenomena
(mitotic failures, nuclear cycle asynchrony, and nuclear crowding) spatially overlap. Thus, even
though all of the 3 defects can independently reduce embryo fitness, we demonstrate that the
defects cooperate, to induce lethality in a fraction of embryos.

Genetically increasing robustness of early embryos against environmental perturbations
In theory, the robustness of embryonic development can be higher depending on, say, the quality of
the food available to the mother. Thus, one can hypothesize a rescue of embryonic lethality at
elevated temperature through, e.g., improvement of the quality of the diet available to the adult
females. A previous study compared the effect of temperature and the atmospheric O_2 levels on
embryo survival, where they report consistently greater embryo survival at 25% Oxygen
concentration, than the normal 21% (Kuntz and Eisen 2015). These results indicate that it might also
be possible to increase the robustness of embryo development through increasing the atmospheric
O_2 concentration. However, such solution might not be heritable, and might not confer an increase
in future fitness. Previous studies have increased embryo survival in face of exposure to elevated
temperature through over-expression of heat-chock proteins (Lockwood, Julick, and Montooth
2017). Presumably, this increases the capacitance of the system to tolerate environmental insults,
akin to how heat-shock proteins are known to act as capacitors for genetic variation (Rutherford and
Lindquist 1998; Sangster, Lindquist, and Queitsch 2004). However, such a rescue would not shed
light on underlying biology of temperature defects, as the effect of such perturbations on the mitotic
failures would remain nebulous. Thus, to rescue the lethality at elevated temperature, we used our
cellular level understanding of the defects to guide our choice of candidate genes for genetic
manipulations. This approach additionally allowed us to challenge our cellular level understanding of
the defects at elevated temperature, as we construct and test hypotheses about the over-expression
of various genes towards a genetic rescue.

Various genotypes in our experiments, specifically the overexpression of *αCat*, *sgg*, *baz*, or *dlg*, act to
reduce one or more of the defects (embryo lethality, presence of blastoderm holes, fraction of
mitotic failure in crowded region, or that in the combined region) that occur due to exposure to
elevated temperature. Here, we would argue that *αCat* and *sgg* overexpression are the superior
experimental condition among these as far as potential for a rescue is concerned, as they produce

virtually no side-effects (unlike *dlg* overexpression at normal temperature), while robustly reducing
the lethality at elevated temperature (unlike *baz* overexpression). Additional support for this notion
comes from our analysis of the presence of natural variation along thermoclines, where we
identified *sgg* as an interesting candidate gene, along with *scribble*. The list of genes associated with
spindle dynamics (combination of GO terms spindle organization, spindle localization, and cell
polarity) included our candidate genes *baz*, *dlg*, *sgg*, *mud*, and *insc*. While *αCat* is not presently
associated with GO terms related to spindle dynamics, we also did not find it to be clinal in our
extensive literature search. It is possible that natural fly populations don't possess sufficient standing
genetic variation at that locus that could provide a selective advantage. In any case, an overlap
between our candidates and potential natural candidate genes increases our confidence in our
approach to mitigate the detrimental effects of an exposure to elevated temperature during early
embryonic development.

Given that both *αCat* and *sgg* overexpression show a robust rescue of embryo lethality, we would
further argue that the interaction between microtubules and cortical F-actin, as supported by the
tripartite protein complex between αCat, βCat, and Apc2, mediate by sgg, is sensitive to
temperature fluctuations. It is likely that the association/dissociation constants for one or more
binding interactions (F-actin with αCat, αCat with βCat, and βCat with Apc2) decrease/increase with
an increase in temperature. It is also likely that the kinase activity of sgg reduces with an increase in
temperature, hampering the association between βCat and Apc2. Further experiments are required
to test these ideas. However, here we also note that many nuclei manage to divide in embryos
mutant for βCat, Apc2, and sgg (McCartney et al. 2001), forming a somewhat defective blastoderm.
This would indicate presence of alternative mechanisms that can also be reinforced to increase the
robustness of mitosis, and identifying and testing such mechanisms would be of great interest to
rescue embryo lethality at elevated temperature.

How fast can nature increase the robustness against the adverse effects of elevated temperature?
A recent experiment (Rudman et al. 2022) indicates that nature might already be exploiting the
allelic plasticity in our candidate genes to increase robustness against elevated temperatures. In this
study, the researchers allowed fly populations to naturally evolve from spring through summer till
the fall season, and found that specific single nucleotide polymorphisms were selected for in the
regulatory regions of *sgg* and β*Cat*, along with many other genes. It is interesting to see that these
genes show selectable variation at the population level, and that an over-expression of *sgg* in our
analysis is capable of rescuing the embryo lethality at elevated temperature. To be clear, the study
does not comment on whether the accumulated mutations lead to an over-expression or
hyperactivation of *sgg*, and at the same time, the analysis won't pick up candidates that don't have
sufficient variation in the seed population. In any case, our functional analysis can be effectively
used to test for the function of any gene that rescues (or worsens) the effect of elevated
temperature on early embryonic development, in any fly embryo that's amenable to experimental
manipulations. We believe that our experimental pipeline holds the promise to uncover exciting
biology, while understanding how physiological processes are affected by the exposure to elevated
temperature.

Sensitivity of early embryonic development to the global warming and climate change
Climate change is a global problem that threatens the existence of life on earth as we know it (IPBES
2019). Given the gigantic scope of the problem, it is overwhelming to address the question in its
entirety. Instead, we can breakdown the problem into manageable pieces, in accordance with our

area of expertise, and address them to the best of our abilities. In this context, we decided to focus
on effect of elevated temperature on the physiology of flies, given their vital roles for the
functioning of ecosystems (Raguso 2020). In our efforts to understand the effects of global warming,
we focused on the early embryonic development in *Drosophila*, which we found to be especially
sensitive to exposure to elevated temperature. Here, we identified the syncytial blastoderm stage
during embryonic development as the earliest possible time window that's sensitive to elevated
temperature. We could also observe cascading defects in the cellular blastoderm formed thereafter.
An increase in embryonic lethality, due to the early exposure to elevated temperatures, further
indicated that, presumably, the subsequent developmental processes are affected as well. We also
see a small increase in embryonic lethality when we expose post-cellularization embryos to elevated
temperature, indicating additionally that the temperature sensitive time window might actually be
longer than just the pre-gastrulation developmental phase. It is easy to imagine that, in wild,
embryos would be exposed to elevated temperature for first few hours, starting form egg lay till well
beyond gastrulation. Given that these early processes during the embryonic development are
conserved across fly species (Lemke, Kale, and Urbansky 2020), our results raise the possibility that
many other key species, would exhibit similar defects during early embryonic development, will have
an increase in lethality, and will be overall threatened due to global warming. It is rather impressive
that an exposure to a non-heat-shock temperature during first few hours of embryonic development
is capable of reducing the fitness of the entire embryonic development to increase embryonic
lethality: an effect which is presumably conserved across the animal kingdom, and would be worth
exploring at cell biological level to improve embryo survival. Perhaps, the temperature sensitive
developmental window is so crucial as it includes gastrulation, very much in agreement with the
famous quote from Lewis Wolpert, "It is not birth, marriage, or death, but gastrulation which is truly
the most important time in your life".

Materials and methods:

<u>Fly lines and genetics</u>:

OregonR line was used as WT *Drosophila*.

Histone-RFP expresses RFP tagged version of His2Av. Insertion on 3rd chromosome. Bloomington #23650.

Jupiter-GFP expresses GFP tagged version of microtubule binding protein Jupiter. Insertion on 3rd chromosome. Always used in combination with *Histone-RFP*. Gift from Jan Felix Evers.

67-Gal4 (mat αTub-GAL4-VP16) is a maternal specific Gal4 driver. Insertion on 2nd chromosome. Always used in combination with *Histone-RFP* and *Jupiter-GFP* in the over-expression experiments (see below). Gift from Thomas Lecuit.

The following lines were used to drive the over-expression of various genes of interest in an attempt to rescue the effects of elevated temperature. *baz*, Bloomington #17613; *dlg*, Bloomington #15779; *αCat*, Bloomington #65590; *sgg*, Bloomington #15886; *mud*, Bloomington #22371; *insc*, Bloomington #39676. An outcross to *w1118* was used as control.

For various over-expression experiments, we first crossed virgins from the Gal4 line '+; 67-Gal4; Jupiter-GFP, Histone-RFP' to the males of over-expression line 'UAS-X; +; +' (X= *baz*, *dlg*, *sgg*, or *mud*) or '+; +; UAS-Z' (Z= *αCat*, or *insc*), or to males of *w1118* as control. F1 virgin females from these crosses were back crossed to males of the same paternal genotype.

Genotypes associated with various figure panels are listed in Table S3.

<u>Embryo collection</u>: For any experiment, we put males and females of appropriate genotypes into transparent cylindrical embryo collection cages, with inner diameter of ~5cm. Fine wire mesh is glued to top side of the cage to allow air exchange while keeping the flies trapped, while the bottom of the cage is closed with apple-juice agar plates, with a thin layer of yeast paste, which is where flies deposited the embryos. To prepare the agar plates, we typically prepare 1l of apple-juice agar solution. For this, we boiled 24gm of agar in 750ml of water, to which we added 250ml of fresh store-bought apple juice, while stirring on a hot plate. We make the final amount of sucrose to 25gm: the amount of additional sucrose depends on the sugar content of the apple juice. This solution is sufficient for ~80 petri-dishes of 6cm diameter. After agar is solidified, the plates can be stored for long time at 4°C. Before the embryo collection, the plates were covered with a thin layer of yeast paste (to stimulate oviposition), and were pre-incubated to the embryo collection temperature depending on the temperature regimen, for at least half an hour. *Drosophila* females are known to hold onto the embryo long after fertilization, such that a large part of the early embryo development can occur inside the abdomen. To avoid these embryos, we perform a sham embryo collection for about an hour, from which all embryos are discarded. Following this sham collection, we collect the embryos that are actually used in the experiment.

<u>Embryo lethality experiment</u>: There were 3 temperature regimes in the embryo lethality experiments, as depicted in Figure 1A. Depending on the regimen, we collected embryos for 1 hour either at normal temperature (24-25°C) or elevated temperature (29-30°C). The 24-25°C collections (corresponding to normal temperature) were done on the bench top. The 29-30°C collections (corresponding to elevated temperature) were done in a water bath, where the bottom surface of the embryo collection agar plate is in thermal contact with the heated water. For NN or EN, we

collected embryos for 1 hour either at normal temperature or elevated temperature, and then
incubated them at the same temperature for another 2 hours. Afterwards, the embryos are washed
with bleach for 10-15 seconds, then washed thoroughly with tap water, aligned on a 24x60mm
rectangular coverslip, and covered with halocarbon oil. For NE, we collected embryos for 1 hour at
normal temperature, and then incubated them at the same temperature for another 1.5 hours.
Within the next half an hour, the embryos are washed with bleach for 10-15 seconds, then washed
thoroughly with tap water, aligned on a 24x60mm rectangular coverslip, and covered with
halocarbon oil. Then, we incubated them at elevated temperature for 3 hours. Then, for all the
temperature regimen, the coverslips with embryos were incubated at normal temperature for about
a day, following which we counted the number of eclosed larvae, and the number of dead embryos.
The embryonic lethality was estimated as the percentage of dead embryos, i.e., 100*(number of
dead embryos)/ (number of dead embryos + number of larvae). We excluded the unfertilized
embryos from these calculations.

Embryo fixation and Imaging: We used 2 different temperature regimens for embryo collection for
fixation and staining. We collected embryos for 3-4 hours either at normal temperature (24-25°C) or
elevated temperature (29-30°C). The 24-25°C collections (corresponding to 25°C regimen) were
done on the bench top. The 29-30°C collections (corresponding to 29°C regimen) were done in a
water bath, where the bottom surface of the embryo collection agar plate is in thermal contact with
the heated water. Afterwards, the embryos are washed with bleach for 40-45 seconds, then washed
thoroughly with tap water, and then fixed immediately. Collecting embryos for longer duration is not
a problem here, as we only image and analyze the embryos that are before gastrulation stage,
through post hoc staging and sorting. Here, we did 2 kinds of experiments; one, where the aim was
visualizing the nuclei from the entire embryo (data in Figure S2); and two, where the aim was
visualizing the nuclei along with F-actin distributions from half embryos (data in Figure S9).
Accordingly, we used 2 different protocols for embryo fixation, staining, mounting, and imaging.

In the experiment where we wanted to *visualize all nuclei from the entire embryo*, we used heat
fixation, and stained the embryos with DRAQ5. Briefly, we transferred the processed embryos to
50ml tubes with a brush, and drained the excess water. To this tube, we added the boiled-to-
turbidity heat-fix solution (28% 500µl NaCl + 5% 200µl TritonX100 + 19.3ml water), incubated it for
~20sec with thorough mixing, and then immediately added water to cool down the mixture. Then,
we transferred the fixed embryos to 2ml microfuge tubes, sequentially added n-Heptane and
methanol in equal proportions, vigorously shook the tube for about half a minute to break-open and
remove the vitelline membrane. This was followed by 3x washes of methanol. Here, we started the
staining immediately after, although the embryos can be stored in methanol for long time. The
embryos were rehydrated with PBT (0.01% solution of Tween20 in PBS) over 3 washes, following this
we stained the embryos with 1:1000 DRAQ5 for 1 hour. Afterwards, we washed the embryos 6x with
PBT, re-suspended them in 3:1 glycerol:PBS solution, and mounted them between 2 rectangular
coverslips. The spacing between the coverslips was optimized to keep the embryos in place w/o
deforming them. Then we imaged the embryos through both coverslips, and stitched the two half
embryos during image analysis to get the full view (see below). Images were acquired on Leica SP8
DMi8 inverted confocal microscope using 20x glycerol objective (0.75NA), with 1.15x digital zoom, at
1µm step-size in z, and a final x-y resolution of 0.247 µm/pixel.

In the experiment where we wanted to *visualize the nuclei and F-actin*, we used formaldehyde
fixation, and stained the embryos with DAPI and Phallacidin. Briefly, we transferred the processed

embryos to 20ml vial with a brush, which contained the fixation mix (3500µl PBS + 37% 1000µl

formaldehyde + 4500µl n-Heptane). The vial was shaken on a platform shaker for 25 minutes

@200rpm. Then, we transferred the fixed embryos to 2ml microfuge tubes, sequentially added n-

Heptane and 90% ethanol in equal proportions, vigorously shook the tube for about half a minute to

break-open and remove the vitelline membrane. This was followed by 3x washes of 90% ethanol.

Here, we started the staining immediately after, although the embryos can be stored in 90% ethanol

for a week or so. The embryos were rehydrated with PBT (0.01% solution of Tween20 in PBS) over 3

washes, following this we stained the embryos with 1:1000 DAPI for 1 hour, followed by 1:500

Phallacidin for 1 hour. Afterwards, we washed the embryos 4x with PBT, re-suspended them in 3:1

glycerol:PBS solution, and mounted them between 2 rectangular coverslips, followed by imaging.

Images were acquired on Leica SP8 DMi8 inverted confocal microscope using 20x glycerol objective

(0.75NA), with 1.2x digital zoom, at 1µm step-size in z, and a final x-y resolution of 0.473 µm/pixel.

<u>Embryo Live imaging</u>: We used 4 experimental temperature regimes for embryo live imaging. We

collected embryos for 1 hour either at normal temperatures (22-23°C, 24-25°C, or 27-28°C) or

elevated temperature (29-30°C). The 22-23°C collections (corresponding to 22°C imaging) were done

in the fly incubator. The 24-25°C collections (corresponding to 25°C imaging) were done on the

bench top. The 27-28°C collections (corresponding to 28°C imaging) or the 29-30°C collections

(corresponding to 29°C imaging) were done in a water bath, where the bottom surface of the

embryo collection agar plate is in thermal contact with the heated water. Afterwards, the embryos

are washed with bleach for 35-40 seconds, then washed thoroughly with tap water at the same

temperature as embryo collection. We then aligned the embryos on a 25mm diameter circular

coverslip, covered them with halocarbon oil, and immediately transferred them to the temperature

controller setup on the microscope that's pre-equilibrated to the experimental temperatures for

normal temperature regime, (22, 25, or 28°C) or the elevated temperature regime (29°C). We used

the temperature controller setup from Warner instruments (locally supplied by

MultiChannelSystems). The setup consists of; QR-40LP, which holds the 25mm coverslip with

embryos; QE-1HC, which heats/cools QR-40LP along with the embryos mounted in it; LCS-1, the

liquid cooling system that isolates the QE-1HC from the effect of ambient temperature; and CL-100,

the bipolar temperature controller that can impart the temperature of interest against the baseline

set by LCS-1. We mounted QE-1HC on the microscope stage, with LCS-1 and CL-100 as the

peripherals controlling the temperature. Images were acquired on Leica SP5 DMI6000CS inverted

confocal microscope using 40x water objective (1.1NA), with 1x digital zoom, at 2µm step-size in z

with 20 z-steps overall, and a final x-y resolution of 0.379 µm/pixel. As mentioned in the main text,

we acquired images starting from interphase 10 or before, till the beginning of gastrulation

movements. We recorded the timelaspe at a frame rate of 1 frame every 30sec, 1min, or 2min

depending on whether we recorded at a time 1, 2, or 3-4 embryos in parallel.

<u>Image analysis and statistics</u>: All image analysis was done using FIJI freeware. There are 7 different

kinds of image analyses presented in the manuscript. These are; one, whole embryo reconstruction

to see distribution of nuclei in the entire embryo (fixed embryos); two, the analysis of mitotic

failures (live embryos); three, the analysis of nuclear movements (live embryos); four, analysis of

nuclear cycle asynchrony (live embryos); five, analysis of nuclear crowding (live embryos); six,

analysis of the fraction of mitotic failures (live embryos); and seven, comparing the distribution of

nuclei alongside that of F-actin (fixed embryos). All the statistical analysis was done using Prism 9

(GraphPad). This includes, the curve fitting (see below, the analysis of nuclear crowding), estimating

the non-parametric statistical significances, and estimating the 95% confidence intervals (CI) on the
medians. The straight-line fits for analysis of nuclear movements were done using ImageJ macro
language.

In the analysis to *reconstruct the nuclear distribution in the whole embryo*, we used a custom-built
image analysis pipeline to fuse the halves of the same embryo, followed by the computing the
distance of each point inside the embryo from the embryo surface. Here we refer to the two views
of the embryo as "half embryo", even though in reality the volume in either "half" spans about 70-
75% of the embryo width, providing a significant overlap between the two halves. In the pipeline, we
process the raw images, stitch the two half embryos, render the fused embryo in lateral view,
segment the embryo volume, and then segregate the nuclei as a function of their distance from the
surface of segmented embryo volume. First, we processed the raw images: to improve visualization
of the nuclei, we used 'Minimum 3D' filter with radii x=y=1, and z=3, and to reduce noise, we used
'Median 3D' filter with radii x=y=z=1. Then, we homogenized the brightness/contrast between the
two half embryos. Following these steps, we manually aligned the half embryos along their AP axis,
and then used the BigWarp (Bogovic et al. 2016) package in the BigDataViewer ecosystem (Pietzsch
et al. 2015) to align the half embryos using manually selected interest points. Essentially, we
selected either subcortical- or yolk-nuclei that could be identified in both half embryos. We further
cropped the output images to keep only a small margin around the embryo with "aligned halves", in
order to reduce the computational load in the steps to follow. After the manual alignment step, we
used BigStitcher (Preibisch, Saalfeld, and Tomancak 2009) to further align embryo, using phase
correlation method, to produce the "fused embryo". In case of many embryos, additional manual
transforms were necessary to produce a fused embryo that passes visual inspection. The fused
embryos were then rotated as necessary, and resliced to produce lateral views. Following this, we
processed the fused embryo to separate the regions of embryo that contain cortical-, subcortical,
and yolk-nuclei. First, we used the nuclear signal to segment the embryo volume, using simple
thresholding. This generated a binary image that corresponds to the embryo volume. Then using '3D
Distance Map' (a plugin in the 3DSuite), we compute the Euclidian distances in 3D, i.e., for each pixel
inside the embryo gets the grayscale value corresponding to its distance from the embryo surface.
All pixels with Euclidian distance <10µm would be considered as the cortical layer, those within
range of 10-30µm would be consider subcortical, and those farther than 30µm from embryo surface
would be considered as part of the yolk region. These distances are determined empirically, using
approximate measurements from various fused embryos. Typically, nuclei have the diameter of 5-
10µm depending on the phase of nuclear division cycle, while the cell height at the end of
cellularization is about 30µm. The pixels in these regions were then color coded and presented as
either as maximum projections or cross-sectional views in Figure S2.

In the *analysis of mitotic failures*, we tracked expelled nuclei back in time to identify the mitosis
when they were born, and to categorize the expulsion as pairwise or isolated (see examples
respectively in Figure 1F and Figure S3D). We considered a nucleus to be expelled, if its apical-most
tip was at least 12µm away from the embryo surface. To simplify the identification of expelled
nuclei, we color-coded the Histone-RFP signal as a function of distance from the embryo surface.
This is similar to above analyses, except we use the 'LocalZPojector' plugin in FIJI (Herbert et al.
2021) to define the distance from the embryo surface, while to predict the embryo surface itself we
use the Jupiter-GFP signal, which was co-imaged with Histone-RFP. Thus, first we color-code the

nuclei, and then manually identify and count the number of pairwise or isolated nuclear expulsions
during interphase 14.

In the *analysis of nuclear movements*, we manually tracked the entire progeny of a few nuclei,
starting from the mother nucleus in late interphase 10 till the 8 great-granddaughter nuclei till late
interphase 13. Having assigned all of these nuclei to a single cluster, we can then calculate the center
of the cluster at each time point. We chose nuclei along one continuous row at interphase 10, and
tracked their progeny in their individual clusters. For sake of simplicity and consistency, we only
chose those nuclear clusters whose entire progeny can be tracked, i.e., we can follow the cluster and
see the formation of all 8 great-granddaughter nuclei in interphase 13. This way, we obtained ≥ 13
clusters in any embryo that we tracked. For each of these nuclear cluster trajectories, we fitted a
straight-line to coordinates for the center of the cluster, over time (dashed lines in Figure S5B). Then,
we calculated the distance of the center of the nuclear cluster to this line at every timepoint. An
average of these distances over a specific time period was used as an estimate for deviations in the
center of nuclear clusters over that time period. We observed that the clusters move significantly
starting from anaphase till a few minutes into the subsequent interphase, as the clusters return to
their normal position. Thus, we decided to separately analyze these periods of extensive movement
vs. the periods when the cluster movement is low. To identify the periods of extensive movements
consistently and objectively, we noted that these movements start at least 1 timepoint (30 sec)
before we see the first telophase in any of the tracked clusters, while the movements mostly subside
within 3 minutes after all of the nuclear clusters have finished telophase. Thus, we defined the
window of extensive movement as starting from 30 sec before appearance of first telophase nuclei,
till 3 min after the appearance of last telophase nuclei. Once we have an objective definition of
various time periods, it is straightforward to calculate the average deviation in the center of nuclear
clusters.

In the *analysis of asynchrony in nuclear cycles*, we marked various timepoints in the timelapse
recordings to follow the progression of nuclear cycles in the anterior-, middle-, and posterior-5th of
the field of view. For sake of simplicity, we marked the transitions from a telophase 13 to interphase
14, i.e., the first timepoint when the daughter nuclei in any region go from telophase 13 to
interphase 14. If the timepoint for this occurance in the anterior is 'T_a', that in posterior is 'T_p', while
the same in the middle is 'T_m', then the 'asynchrony' can be calculated as,

$$asynchrony = T_m - \left(\frac{T_a + T_p}{2}\right)$$

Essentially, we start with the expectation that T_m should be the average of T_a and T_p (maximum
synchrony), and a deviation from this expected value would be the asynchrony. Figure S7
diagramatically demonstrates this analysis.

In the *analysis of nuclear crowding*, we analyzed the distribution of nuclei at the end of interphases,
as that's the time when the distribution is as stable as it can be. To assess nuclear crowding, we
started with segmentation of nuclei in the field of view. Here, we used the 'Find Maxima' command
to identify local intensity maxima, and imprinted those on an image. Many of these maxima marked
a nucleus. We did manual corrections to, remove the maximum that didn't correspond to a nucleus,
add a maximum for nucleus that didn't have a maximum associated with it, or correct the maxima
that were too off-center from corresponding nuclei. Then we used Voronoi tessellation to mark
territories around these maxima to generate pseudo-cells. We excluded territories that, touched the

edge of the image, that had aberrant shapes (aspect ratio>2), or those with aberrant pixel intensities
(pseudo cell containing pixels with <5 units of intensity, typically the case with ill-defined pseudo-
cells at the edge of segmentation). Thus, each nucleus is associated with a pseudo-cell area value
('Area' in μm^2), and an x-coordinate value for the center of the pseudo-cell ('X' in μm) (x-axis of the
image aligned with anterior-posterior axis of the embryo). We fitted these with the equation of a
parabola such that 'Area' can be expressed as a function of 'X',

$$Area = aX^2 + bX + c$$

Here, the curvature of the parabola will be higher in case of greater nuclear crowding (compare
Figure 2L-O). Thus, the fit parameter 'a', which dictates the curvature of the parabola, was used to
estimate the extent of crowding.

In the *analysis of the fraction of mitotic failures*, we tracked expelled nuclei back in time same as the
analysis for mitotic failures, except now we also recorded where the nuclei were lost from, to get a
map of the distribution of mitotic failures. In parallel, we generated regions in the embryo that
correspond to maximum nuclear crowding, maximum delay in nuclear divisions, and a combination
of the two regions. To have an objective definition of the area of maximum crodwing we looked to a
previous study on the effect of nuclear density on nuclear division cycles (Lu et al. 2009). This study
indicates that the nuclear division cycles seize when the embryo reaches ~70% of the normal
cellularization nuclear density. Another previous study reported the nuclear domain area (i.e., the
pseudo-cell area) to be 28.8 μm^2 at cellularization (Telley et al. 2012). We corroborate this number
with our own quantifications (data not shown). Thus, for simplicity of calculations, we assume that
Drosophila embryos have 6000 nuclei at interphase 14 each with a area of 29 μm^2. The a 70%
density of nuclei would thus correspond to 4200 nuclei at interphase 14, each with an area of 41.43
μm^2, which we approximate to 42 μm^2. Thus the back of the envelop calculation indicated that
nuclei will seize to divide when their pseudo-cell area drops below ~42μm^2. We observed that in
embryos developing at 29°C, the central regions of the embryo contained many nuclei that had
pseudo-cell areas less than 42μm^2, even at interphase 13. Thus, we used this criterion to define
areas of embryos that have likely reached the 70% cellularization nuclear density, at one nuclear
division cycle earlier, albeit only locally. Since the embryos at normal temperatures (22, 25, and
28°C) rarely have nuclei with pseudo-cell areas smaller than 42μm^2, we further decided to include all
nuclei with pseudo-cell areas in the lower quartile to construct the region with maximum nuclear
crowding. Then, to construct the region with maximum delay in nuclear divisions, we simply marked
the areas of the embryo that contain the last ~20% of the nuclei that finished nuclear cycle 13. This
percentage was decided empirically: it corresponds to a region of similar size as that corresponding
to the crowded region above. Here the rationale is drawn from previous studies which indicate that
the regulation of nuclear division cycles is coordinated at the scale of the entire embryo (Deneke et
al. 2016), and given that all of the nuclei share the same cytoplasm, a local delay in nuclear cycles
would create conflicts between the biochemical status of the cytoplasm and the cell cycle stage of
the delayed nuclei. Further, we combined the crowded and delayed regions using the logical OR
operator, to test combinatorial effects of the nuclear crowding and nuclear cycle asynchrony. Finally,
we computed the fraction of mitotic failures in areas thus constructed.

In the analysis to *compare the distribution of nuclei alongside that of F-actin*, was rather similar to
the pipeline to *reconstruct the nuclear distribution in the whole embryo*, except that we didn't have 2
half-embryos to fuse, essentially starting from segmentation of embryo volume. Here, instead of

using nuclear staining, we segmented embryo volume using the F-actin staining. Then the 3D
distance map similarly allowed us to classify pixels into cortical-, subcortical-, or yolk-regions based
on their distance from embryo surface. The pixels in these regions were then color coded and
presented as sum projections in Figure S9.

Figure and table legends:

Figure 1: Embryonic lethality and the loss of nuclear due to mitotic failures in embryos exposed to
elevated temperature.
(A, B) To assess developmental robustness, we exposed embryos to various temperature regimen
and estimated the embryo lethality. (A) Schematic showing the temperature regimen in the embryo
survival experiment, starting from the beginning of embryo collection. Here we have three
conditions depending on exposure to normal or elevated temperature regimen in the 3-hour time
windows as shown, normal-normal [NN], elevated-normal [EN], and normal-elevated [NE]. In all
cases, the first time-window also includes 1 hour of embryo collection. The embryonic development
beyond 6 hours continued at normal temperature. (B) The scatter plots show the distribution of
percent lethality in embryos from various conditions, calculated based on the fraction of embryos
that don't hatch into larvae. Each dot represents data from a sample with ≥26 embryos.

(C-E) Montages of snapshots from live embryos, expressing Histone-RFP, from late interphase 11 till
late interphase 14, in embryos developing either at normal temperatures (C), or at elevated
temperature showing mild (D), or severe (E) defect. The nuclei were labelled as a function of their
distance from the embryo surface. Here, green indicates cortical, and magenta indicates sub-cortical
nuclei that are lost. At elevated temperature, a lot more nuclei are expelled, and as a result, we can
see large gaps between cortical (green) nuclei, often coinciding with expelled (magenta) nuclei,
especially during early interphase-14.

(F-H) To assess the pairwise loss of nuclei, we counted mitotic failures that lead to such nuclear
expulsions. (F) Montage of snapshots from live embryo, from late interphase 12 to late interphase
13, where Histone-RFP marks the distribution of nuclei, cortical nuclei are in green and subcortical
(expelled) nuclei are in magenta. The pairwise loss of nuclei due to mitotic failure is exemplified in an
embryo developing at elevated temperature. The arrowheads track the segregating daughter nuclei
that will be expelled eventually, while arrows exemplify normal nuclei. Of note, this type of nuclear
loss occurs at all temperatures, though at different a frequency. (G) Scatter plot showing the
distribution of events of nuclear losses due to mitotic failure during interphase 14. (H) The schematic
to illustrate the phenomenon of mitotic failures at elevated temperature.

Scale bars in C-E are 25 µm, in F is 5 µm. B and G show scatters with median, with whiskers showing
95% CI. In G, the hollow markers indicate embryos that have a blastoderm hole, with large hollow
markers indicating a single large hole, and small hollow markers indicating either a single or multiple
small holes. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$, non-significant comparisons are
not shown.

Figure 2: Nuclear cycle asynchrony and nuclear crowding increases as a function of temperature
during embryo development.
(A-C) To assess the asynchrony in nuclear cycles, we identified the phases of nuclear cycle along the
AP axis. (A) Image shows the entire anterior-posterior extent of the field of view, from a live embryo
developing at 29°C. Histone-RFP marks the nuclei, cortical nuclei are in green and subcortical

(expelled) nuclei are in magenta. Different regions of the embryo have nuclei in different stages of
nuclear division cycle, as indicated above the image. Arrows show the direction in which the wave of
nuclear division cycle will progress. Here, we have an anterior- and a posterior-wave which will meet
in the central region of the embryo. The anterior, central, and posterior 1/5th of the field of view are
indicated below the image, with the anterior 1/5th of the embryo containing nuclei that are more
advanced in division cycle, as compared to the posterior 1/5th, which are yet more advanced than
the central 1/5th. As a result, these regions will enter the subsequent interphase in an asynchronous
manner (see also Figure S7). (B) Scatter plots showing the distribution of asynchrony for embryos at
various temperatures. (C) The schematic illustrates the progressive increase in asynchrony at
elevated temperature.

(D-Q) To assess the crowding of nuclei, we segmented the nuclei and generated pseudo-cells. The
figure shows data from of live embryos that were developing either at 22°C (D, H, L), 25°C (E, I, M),
28°C (F, J, N), or at 29°C (G, K, O). (D-G) Snapshots from live embryos at the end of interphase 13,
where Histone-RFP marks the nuclei. The outline shows the region of the image that is considered
for quantifications. (H-K) The pseudo-cells for the corresponding embryos in D-G. A pseudo cell is
centred on a nucleus, while the region of a pseudo-cell is bound by line segments that are
equidistant from neighbouring nuclei. Various pseudo-cells are colour-coded according to their area
in μm^2 according to the LUT at the bottom. (L-O) Plots showing the distribution of pseudo-cell areas
as a function of their anterior-posterior position, along with a quadratic fit through the distribution,
for embryos in D-G. (P) Scatter plots showing distribution of the coefficient of the quadratic term for
embryos at various temperatures. This essentially captures the extent of nuclear crowding, with
larger quadratic coefficient indicating greater crowding. (Q) The schematic illustrates the crowding
effect at elevated temperature. The size of the black arrows indicates the gradation in nuclear
density, from poles to the central region.

Scale bars, 20 μm. B and P shows scatters with median, with whiskers showing 95% CI. In B and P,
hollow markers indicate data from embryos that have blastoderm hole, with large hollow markers
indicating a single large hole, and small hollow markers indicating either a single or multiple small
holes. *, p<0.05; **, p<0.01; ***, p<0.001; non-significant comparisons are not shown.

Figure 3: Nuclear crowding and nuclear cycle asynchrony combinatorially enhance mitotic failure
defect.
(A, B) Snapshots from live embryo, developing at 29°C, at the end of interphase 13 (A), and 14 min
afterwards (B), where Histone-RFP marks the nuclei. The outline in A shows the region of the image
that is has nuclear crowding, while that in B bounds the region containing most delayed nuclei, i.e.,
those that are most lagging in nuclear division cycles. (C-E) The locations of lost pairs of nuclei during
interphase 14 are marked as white dots to indicate their distribution in the field of view. Then,
various regions are defined in the embryo to illustrate the fraction of mitotic failures occurring in
that region. These regions are composed based on, the crowded region only (C), the delayed region
only (D), and the combination of the crowded and delayed regions (E). (F-H) Scatter plots showing
the distribution of the area fractions in crowded (F), delayed (G), and crowded OR delayed (H)
regions of the embryos. In case of both crowded (F) and delayed (G) regions, the medians are rather
comparable across temperature, and the low variability within a given temperature is exaggerating
the differences across temperatures. In case of the area fraction of the 'crowded OR delayed'
regions (H), the increase in the overlap of crowded and delayed region reduces the overall size of the
combined region. (I-K) Plots showing the distribution of fraction of mitotic failures occurring in the

crowded region only (I), in the delayed region only (J), and in the combination of the crowded and delayed regions (K).

Scale bars, 20 μm. F-K show scatters with median, with whiskers showing 95% CI. Hollow markers indicate data from embryos that have blastoderm hole, with large hollow markers indicating a single large hole, and small hollow markers indicating either a single or multiple small holes. *, p<0.05; **, p<0.01; ***, p<0.001; non-significant comparisons are not shown.

Figure 4: Rescue of embryonic lethality at elevated temperature through over-expression of the regulators of cell polarity signalling or those of F-actin -microtubules interaction during mitosis, and genes varying along thermoclines as "natural" candidates.

(A) Scatter plot showing the distribution of percent lethality in embryos from various genotypes, at normal (25°C) vs elevated temperature (29°C), calculated based on the fraction of embryos that don't hatch into larvae. Each dot represents data from a sample with ≥12 embryos. (B, C) Scatter plots showing the distribution of the fraction of mitotic failures in crowded regions (B), and in crowded OR delayed regions (C), in the embryos at normal (22°C) vs elevated temperature (29°C). Each dot represents data from one embryo. Additional quantifications related to this experiment are presented in Figure S8.

(D) Scatter plots showing the distribution of percent lethality in embryos from various genotypes, at normal (25°C) vs elevated temperature (29°C), calculated based on the fraction of embryos that don't hatch into larvae. Each dot represents data from a sample with ≥18 embryos. (E, F) Scatter plots showing the distribution of the fraction of mitotic failures in crowded regions (E), and in crowded OR delayed regions (F), in the embryos at normal (22°C) vs elevated temperature (29°C). Each dot represents data from one embryo. Additional quantifications related to this experiment are presented in Figure S10.

(G) Pie charts showing the clinal distribution score for various genes associated with 'spindle dynamics' (combination of GO terms, cell polarity, spindle organisation, and spindle localisation, n=320 genes), or with GO term 'metabolic process' (n=6612 genes). ~40% of genes associated with spindle dynamics, while ~30% of the genes associated with metabolic process, are clinal in at least one continent.

A-F shows scatters with median, with whiskers showing 95% CI. In B, C, E, and F, hollow markers indicate data from embryos that have blastoderm hole, with large hollow markers indicating a single large hole, and small hollow markers indicating either a single or multiple small holes. *, p<0.05; **, p<0.01; ***, p<0.001; ****, p<0.0001; non-significant comparisons are not shown.

Figure S1: Blastoderm holes appear in a fraction of embryos at elevated temperature.

The figure shows the snapshots from live embryos, expressing Histone-RFP, from early interphase 14, in embryos developed at elevated temperature. The nuclei are labelled as a function of their distance from the embryo surface. Here, green indicates cortical, and magenta indicates sub-cortical nuclei that are expelled, while arrows/arrowheads indicate large/small blastoderm holes. For embryos with large blastoderm holes (embryo 05, and 08), we see many nuclei in magenta, while in case of embryos with small blastoderm holes (embryos 04, and 09), we can see only a few nuclei in magenta per hole. We would argue that development in *Drosophila* is robust enough to buffer the defects due to the small holes, but fails at 29°C when large holes appear.

Scale bar, 25 μm.

Figure S2: Nuclear loss shows left-right asymmetry in embryos developing at elevated temperature.

Distribution of nuclei in embryos during interphase 13, in fixed embryos that were developing either at normal temperature (A, D, G, H), or at elevated temperature that show either a mild defect (B, E, I, J) or a severe defect (C, F, K, L). (A-F) Maximum projections of the right and left halves of embryos, where the nuclei are labelled as a function of their distance from the embryo surface. Here, green indicates cortical nuclei, i.e., nuclei close to embryo surface, while magenta indicates sub-cortical nuclei, i.e., nuclei that are deeper in the embryo, farther from embryo surface. The yolk nuclei are not shown for improved visibility. As compared to embryos at normal temperature (A), those at elevated temperature show mildly (B) or severely (C) numerous gaps in the distribution of cortical nuclei. These gaps often also correspond to the locations of subcortical nuclei (A-C). Of note, the embryo surface prediction gets distorted at the posterior pole, due to the pole cell nuclei that are outside the embryo cortex, thus mislabelling cortical nuclei in magenta. These mis-coloured nuclei were manually erased to exclusively highlight the subcortical nuclei (D-F). (G-L) Cross-sectional view along the indicated regions in panels A-C. Here, yolk nuclei are additionally marked in yellow. The arrow indicates the direction along which we can follow the surface of the embryo along the cross-section. The cross-sections are chosen to specifically highlight the locations of subcortical nuclei.

Scale bars in A-C are 50 μm, while in G, I, and K are 25 μm.

Figure S3: Out-of-plane nuclear divisions during mitosis 10, and the low frequency of isolated nuclear expulsion during interphase 14, does not vary significantly with temperature, and cannot explain the observed loss of nuclei.

(A, B) Montages exemplifying out-of-plane division during mitosis 10 in an embryo developing at 25°C. Histone-RFP marks the nuclei (green), while Jupiter-GFP marks the microtubules (magenta). Arrowheads exemplify in-plane divisions, while arrows exemplify out-of-plane division. Each subsequent frame is 1min apart. In the x-y view of the divisions (A), only one of the daughter nuclei is clearly visible after the out-of-plane division, while the other daughter nucleus is better seen in the y-z view of the division (B), where we can see it translocating to the embryo interior. (C) Plots show the number of out-of-plane divisions, that are visible in the field of view, in embryos developing at various temperatures. For context, an embryo contains on the order of 70 mitotic spindles the field of view at interphase 10. So, even though the p-value is <0.05 for the comparison between 22°C and 29°C, and given that there is no correlation between the number of out-of-plane divisions and the presence of blastoderm holes, we would argue that the difference is not biologically significant.

(D) Montage exemplifying isolated nuclear expulsion in an embryo developing at 25°C, from telophase 12 to late interphase 13. Histone-RFP marks the distribution of nuclei, cortical nuclei are in green and subcortical (expelled) nuclei are in magenta. The arrowheads track the daughter nucleus that will be expelled, while arrows track the daughter nucleus that will be retained cortically. (E) Scatter plots showing the distribution of isolated loss of nuclei during interphase 14.

Scale bar in A is 10 μm, and in D is 5 μm. C and E show scatters with median, with whiskers showing 95% CI. The hollow markers indicate embryos that have a blastoderm hole, with large hollow markers indicating a single large hole, and small hollow markers indicating either a single or multiple small holes. *, p<0.05; non-significant comparisons are not shown.

Figure S4: Defective nuclei take different amount of time to be expelled from embryo cortex.

(A, B) The figure shows the montages of snapshots from live embryos, with a few representative

nuclei tracked from late interphase 12 to interphase 13. Top panels show Histone-RFP signal, while

bottom panels show Jupiter-GFP signal. Nuclei tagged in green/ magenta are the ones that arise

from mitotic defects during mitosis 10/ 11. All of these are expelled during interphase 13. Notice the

lack of spindle formation around the tagged nuclei. Also note that the nuclei expelled by the 5th

timepoint are not shown in the 6th timepoint. (A) An embryo developing at normal temperature,

where 1 magenta and 1 green nuclei are followed. (B) An embryo developing at elevated

temperature, where 6 magenta and 3 green nuclei are followed.

Scale bars are 20 μm. The numbers in inset indicate time in min:sec.

Figure S5: Nuclear movement parallel to the embryo surface is amplified at elevated temperature.

(A-C) An example depicting how nuclear movements, and the deviations thereof, are calculated.

Snapshots from live embryo developing at elevated temperature, where Histone-RFP marks the

distribution of nuclei, showing the mother nuclei at the end of interphase 10 (A) and the

corresponding 8 great-granddaughter nuclei, each, at the end of interphase 13 (C). The neighbouring

nuclear clusters are marked in either cyan or yellow. The trajectories of the center of nuclear

clusters (B) also indicate the increase in the number of nuclei due to divisions, as the broadening of

the tracks. The dotted lines indicate straight-lines fitted to individual trajectories, which are used as

reference for computing deviations for each time point of that trajectory. The shaded region

captures periods of greatest movement that starts with anaphase (see methods). (D, E) To quantify

the drift in the center of nuclear clusters, we compute the slope of the straight-line fits (dotted lines

in B). The median of slopes for all nuclear cluster from one embryo provides a robust estimate of the

drift in the centers of the nuclear clusters. Plots showing the distribution of these medians (one dot

per embryo), for the drift along AP axis (D), and for that along DV axis (E). (F-I) To quantify the

movements in the center of nuclear clusters during periods of maximum movement (shaded region

in B) and normal movement (unshaded regions in B), we compute the average of the deviations from

the straight-line fits (dotted lines in B). The median of such average deviations for all nuclear cluster

from one embryo provides a robust estimate of the nuclear cluster deviation. Plots showing the

distribution of these medians (one dot per embryo), for the fluctuations along AP axis (F, H), and for

those along DV axis (G, I), during interphase-metaphase (F, G) or during anaphase-telophase (H, I).

Scale bar is 25 μm. D to I show scatters with median, with whiskers showing 95% CI. Hollow markers

indicate data from embryos that have blastoderm hole, with large hollow markers indicating a single

large hole, and small hollow markers indicating either a single or multiple small holes. *, p<0.05; **,

p<0.01; ***, p<0.001; non-significant comparisons are not shown.

Figure S6: Nuclear movement along surface-yolk axis is amplified at elevated temperature.

Cross-sections from live embryos, expressing Histone-RFP, from late interphase 13 till early

interphase 14, in embryos developing either at normal temperature (A), or at elevated temperature

(B). Here, the coloured lines mark the basal tip of nuclei in the first frames (t=-7:30 min) of the

respective montage. Timepoint 00:00 corresponds to initiation of anaphase at the left/right extreme

of the images, which is also the first time we see the shift from baseline.

Arrowheads point out the local peak of the nuclear shift from the baseline, while arrow points

expelled nuclei. As we can see, at elevated temperature, the nuclei shift significantly from the

baseline, before returning to normal localisation by the time the nuclei enter the subsequent

interphase. Also, the shift from baseline occurs progressively from anterior/posterior poles towards

the central region.

Scale bars are 25 µm. Cross-sections are in the lateral region of the embryos, anterior to the left.

Figure S7: Nuclear cycle asynchrony increases as a function of the temperature during embryo
development.
(A, B) Montages of snapshots from metaphase 13 to early interphase 14, from live embryos
developing either at 25°C (A), or at 29°C (B). Histone-RFP marks the nuclei, cortical nuclei are in
green and subcortical (expelled) nuclei are in magenta. The dotted lines divide the field of view in 5
equal parts. Numbers in the inset represent time in min:sec, while 00:00 is when anterior 1/5th of the
embryo enters interphase 14. Boxes highlight entry into interphase 14 in that region. The black
arrows indicate the time point of maximum synchrony, i.e., halfway between the anterior and
posterior 1/5th of the embryo, when the nuclei in the central region should have entered interphase
14. This point serves as the reference to estimate asynchrony (see methods).

Scale bars, 25 µm.

Figure S8: Additional quantifications related to *baz* or *dlg* over-expression.
(A-F) The distribution of the pseudo-cell areas in embryos at the end of interphase 13, in live
embryos developing either at normal temperatures (A, C, and E), or at elevated temperature (B, D,
and F). The pseudo-cell areas in embryos of control genotype (A, B), in embryos over-expressing *baz*
(C, D), or in embryos over-expressing *dlg* (E, F), are shown. Anterior is to the left, dorsal up. The LUT
bar at the bottom indicates the colour code corresponding to the area of pseudo-cells. (G-K) Scatter
plots showing the distribution of the area fractions in crowded (G), and crowded OR delayed (H)
regions of the embryos, the distribution of the extent of nuclear crowding (I), the distribution of the
nuclear cycle asynchrony (J), and the number of pairwise nuclear expulsions (i.e., the number of
mitotic failures) during interphase 14 (K) for embryos from various genotypes, at normal (22°C) vs
elevated temperature (29°C). Each dot represents data from one embryo.

Scale bar, 25 µm. G-K show scatters with median, with whiskers showing 95% CI. Hollow markers
indicate data from embryos that have blastoderm hole, with large hollow markers indicating a single
large hole, and small hollow markers indicating either a single or multiple small holes. *, $p<0.05$; **,
$p<0.01$; ***, $p<0.001$; non-significant differences are omitted.

Figure S9: Elevated temperature disrupts the association between nuclei and F-actin caps.
Distribution of nuclei and F-actin in embryos during interphase 13, in fixed embryos that were
developing either at normal temperatures (A, C, D), or at elevated temperature (B, E, F). (A, B) The
images show SUM projections of the DAPI stained nuclei (top image) and Phallacidin stained F-actin
(bottom image) of representative embryos. In the top images, the nuclei are labelled as a function of
their distance from the embryo surface, where green indicates cortical, and magenta indicates sub-
cortical nuclei. The yolk nuclei are not shown. Anterior is on the left, dorsal up. (C-F) The top and
bottom images show a zoomed-in view of the indicated regions in panels A and B. Here, arrowheads
indicate the expelled nuclei (C, E top panels) and the structures of F-actin corresponding to them (C,
E bottom panels), which should otherwise be localised around cortical nuclei. Then, the arrows
indicate gaps between neighbouring nuclei (D, F top panels), which should normally be devoid of any
F-actin structures (D, bottom panel), but contain small "actin caps" in embryos at elevated
temperature (F, bottom panel).

Scale bars, 25µm. The boxed regions are 60x60 µm.

Figure S10: Additional quantifications related to *αCat* or *sgg* over-expression.

(A-F) The distribution of the pseudo-cell areas in embryos at the end of interphase 13, in live
embryos developing either at normal temperatures (A, C, and E), or at elevated temperature (B, D,
and F). The pseudo-cell areas in embryos of control genotype (A, B), in embryos over-expressing
αCat (C, D), or in embryos over-expressing *sgg* (E, F), are shown. Anterior is to the left, dorsal up. The
LUT bar at the bottom indicates the colour code corresponding to the area of pseudo-cells. (G-K)
Scatter plots showing the distribution of the area fractions in crowded (G), and crowded OR delayed
(H) regions of the embryos, the distribution of the extent of nuclear crowding (I), the distribution of
the nuclear cycle asynchrony (J), and the number of pairwise nuclear expulsions (i.e., the number of
mitotic failures) during interphase 14 (K) for embryos from various genotypes, at normal (22°C) vs
elevated temperature (29°C). Each dot represents data from one embryo.

Scale bar, 25 μm. G-K show scatters with median, with whiskers showing 95% CI. Hollow markers
indicate data from embryos that have blastoderm hole, with large hollow markers indicating a single
large hole, and small hollow markers indicating either a single or multiple small holes. *, $p<0.05$; **,
$p<0.01$; ***, $p<0.001$; non-significant differences are omitted.

**Figure S11: *mud* or *insc* over-expression does not ameliorate embryo survival at elevated
temperature.**
Scatter plot shows the distribution of percent lethality in embryos from various genotypes, at normal
(25°C) vs elevated temperature (29°C), calculated based on the fraction of embryos that don't hatch
into larvae. Each dot represents data from a sample with ≥21 embryos. As we can see in the plot,
neither *mud* nor *insc* over-expression rescues the embryo lethality at elevated temperature, with
mud over-expression increasing embryo lethality even at normal temperature (25°C).

The plot shows scatters with median, while whiskers showing 95% CI. *, $p<0.05$; **, $p<0.01$; ***,
$p<0.001$; non-significant differences are omitted.

Table S1: This table contains genes that are identified as being clinal in different continents. Columns
A to C contains lists of studies, and under the studies is a list of genes that are identified in these
studies as being clinal. Column A, B, and C respectively list studies that have data from the
continents of Australia and/or Australasia (AUS), North and/or Central America (USA), and Europe
and/or North Africa (EU).

We collect all the genes associated with the GO terms 'cell polarity', 'spindle organization', and
'spindle localization', to compile a non-redundant master list of candidate genes associated with
'spindle dynamics' in Column F. For each gene in column F, we count the number of times the gene
appears in column A to C, counting a maximum of once per column, to respectively populate
columns G to I. For each gene in column F, column J contains the sum of values in G, H, and I, to
indicate the number of continents in which a gene is clinal. Each gene gets a score from 1 to 3, and
we consider the gene to be a relevant candidate if it is found in at least 2 columns.

We collect all the genes associated with the GO term 'metabolic process', to compile a control list of
candidate genes in Column L. For each gene in column L, we count the number of times the gene
appears in column A to C, counting a maximum of once per column, to respectively populate
columns M to O. For each gene in column L, column P contains the sum of values in M, N, and O, to
indicate the number of continents in which a gene is clinal. Thus, each gene gets a score from 1 to 3.

Following articles were used to source the list of genes in column A (Oakeshott, McKechnie, and Chambers 1984; McColl and McKechnie 1999; Bettencourt et al. 2002; Frydenberg, Hoffmann, and Loeschcke 2003; Umina et al. 2005; Sawyer et al. 2006; Weeks, Mckechnie, and Hoffmann 2006; Rand et al. 2010; Kolaczkowski et al. 2011; Paaby et al. 2014), column B (Vigue and Johnson 1973; Oakeshott, McKechnie, and Chambers 1984; Berry and Kreitman 1993; Duvernell and Eanes 2000; P. S. Schmidt, Duvernell, and Eanes 2000; Verrelli and Eanes 2001; Duvernell, Schmidt, and Eanes 2003; Sezgin et al. 2004; P. S. Schmidt and Paaby 2008; Rand et al. 2010; Fabian et al. 2012; Paaby et al. 2014; Machado et al. 2016), and column C (Oakeshott, McKechnie, and Chambers 1984; Costa et al. 1992; Weeks, Mckechnie, and Hoffmann 2006; Kapun et al. 2020).

Table S2: Column A lists candidates from Table S1 column F that have a score of 2 or higher, i.e., those that show selectable variation in natural populations. For these genes, we query FlyBase (http://flybase.org/batchdownload) for additional details like, gene name (column B), gene symbol (column C), and the level of expression for these genes during the first 2 hours of embryo development (column D). The level of expression is based on published transcriptomics data (Graveley et al. 2011), which would correspond to maternal pool of mRNAs, and the most relevant developmental window for this study. The candidate genes clinal in all 3 continents are highlighted in green.

Columns F-J present a table with counts of genes that are associated with various categories of genes in Table S1 column F and L, and a further breakdown of how many of these genes are clinal in how many continents. The breackdown is also plotted as pie-charts in Figure 4G.

Table S3: This table lists various figure panels (columns A and B), and the related parental genotypes, i.e., the genotypes of females (column C) and males (column D) present in the embryo collection cages for the experiment related to a figure panel.

Additional information

Data and Code Availability:
Data and codes developed for data analysis are available upon request from corresponding authors.

Author contributions:
G.K. and S.L. planned the project. G.K. did the experiments with help from P.A. for baz and dlg overexpressions, and JJ.D.-L. for embryo fixation, and nuclei and F-actin staining. G.K. did the image analyses and statistical analyses. G.K. and S.L. wrote the manuscript, and all authors gave comments on the manuscript.

Acknowledgments:
We would like to thank Lazero Centanin, Nicolas Gompel, and Marianthi Karageorgi for insightful comments on various drafts of the manuscript. We are grateful to Jochen Wittbrodt for his enthusiasm, and for his continuous and generous support. We would also like to thank Ingrid Lohmann for *Drosophila* culture related support, and Thomas Lecuit for sharing fly lines. This work wass funded by the DFG (LE 2787/3-1) to S.L., while G.K. was supported through an HFSP Long-term postdoctoral fellowship (LT000597/2019-L).

Competing interests:
The authors declare no competing financial interest.

References:

Berry, A., and M. Kreitman. 1993. "Molecular Analysis of an Allozyme Cline: Alcohol Dehydrogenase
in Drosophila Melanogaster on the East Coast of North America." *Genetics* 134 (3): 869–93.
https://doi.org/10.1093/genetics/134.3.869.

Bettencourt, Brian R., Inyoung Kim, A. A. Hoffmann, and M. E. Feder. 2002. "Response to Natural and
Laboratory Selection at the Drosophila Hsp70 Genes." *Evolution* 56 (9): 1796–1801.
https://doi.org/10.1111/j.0014-3820.2002.tb00193.x.

Bogovic, John A., Philipp Hanslovsky, Allan Wong, and Stephan Saalfeld. 2016. "Robust Registration
of Calcium Images by Learned Contrast Synthesis." In *2016 IEEE 13th International Symposium
on Biomedical Imaging (ISBI)*, 2016-June:1123–26. IEEE.
https://doi.org/10.1109/ISBI.2016.7493463.

Cao, Jian, Justin Crest, Barbara Fasulo, and William Sullivan. 2010. "Cortical Actin Dynamics Facilitate
Early-Stage Centrosome Separation." *Current Biology : CB* 20 (8): 770–76.
https://doi.org/10.1016/j.cub.2010.02.060.

Chong, Jeronica, Christopher Amourda, and Timothy E. Saunders. 2018. "Temporal Development of
Drosophila Embryos Is Highly Robust across a Wide Temperature Range." *Interface* 15 (144).
https://doi.org/10.1098/rsif.2018.0304.

Costa, Rodolfo, Alexandre A. Peixoto, Guido Barbujani, and Charalambos P. Kyriacou. 1992. "A
Latitudinal Cline in a Drosophila Clock Gene." *Proceedings of Royal Society of London*, 43–49.

Crews, Sarah M., W Tyler McCleery, and M. Shane Hutson. 2016. "Pathway to a Phenocopy: Heat
Stress Effects in Early Embryogenesis." *Developmental Dynamics : An Official Publication of the
American Association of Anatomists* 245 (3): 402–13. https://doi.org/10.1002/dvdy.24360.

Deneke, Victoria E., Anna Melbinger, Massimo Vergassola, and Stefano Di Talia. 2016. "Waves of
Cdk1 Activity in S Phase Synchronize the Cell Cycle in Drosophila Embryos." *Developmental Cell*
38 (4): 399–412. https://doi.org/10.1016/j.devcel.2016.07.023.

Duvernell, David D., and W. F. Eanes. 2000. "Contrasting Molecular Population Genetics of Four
Hexokinases in Drosophila Melanogaster, D. Simulans and D. Yakuba." *Genetics* 156 (3): 1191–
1201. https://doi.org/10.1093/genetics/156.3.1191.

Duvernell, David D., Paul S. Schmidt, and Walter F. Eanes. 2003. "Clines and Adaptive Evolution in
the Methuselah Gene Region in Drosophila Melanogaster." *Molecular Ecology* 12 (5): 1277–85.
https://doi.org/10.1046/j.1365-294X.2003.01841.x.

Edwards, Marshall J. 2006. "Review: Hyperthermia and Fever during Pregnancy." *Birth Defects
Research. Part A, Clinical and Molecular Teratology* 76 (7): 507–16.
https://doi.org/10.1002/bdra.20277.

Fabian, Daniel K., Martin Kapun, Viola Nolte, Robert Kofler, Paul S. Schmidt, Christian Schlötterer,
and Thomas Flatt. 2012. "Genome-Wide Patterns of Latitudinal Differentiation among
Populations of Drosophila Melanogaster from North America." *Molecular Ecology* 21 (19):
4748–69. https://doi.org/10.1111/j.1365-294X.2012.05731.x.

Figard, Lauren, Liuliu Zheng, Natalie Biel, Zenghui Xue, Hasan Seede, Seth Coleman, Ido Golding, and
Anna Marie Sokac. 2019. "Cofilin-Mediated Actin Stress Response Is Maladaptive in Heat-
Stressed Embryos." *Cell Reports* 26 (13): 3493-3501.e4.
https://doi.org/10.1016/j.celrep.2019.02.092.

Foe, V. E., and B. M. Alberts. 1983. "Studies of Nuclear and Cytoplasmic Behaviour during the Five
Mitotic Cycles That Precede Gastrulation in Drosophila Embryogenesis." *Journal of Cell Science*
61 (May): 31–70. https://doi.org/10.1242/jcs.61.1.31.

Foe, V.E. 1989. "Mitotic Domains Reveal Early Commitment of Cells in Drosophila Embryos." *Trends*

1178    *in Genetics* 5 (C): 322. https://doi.org/10.1016/0168-9525(89)90120-0.

1179 Frydenberg, Jane, A. A. Hoffmann, and Volker Loeschcke. 2003. "DNA Sequence Variation and
1180    Latitudinal Associations in Hsp23, Hsp26 and Hsp27 from Natural Populations of Drosophila
1181    Melanogaster." *Molecular Ecology* 12 (8): 2025–32. https://doi.org/10.1046/j.1365-
1182    294X.2002.01882.x.

1183 Graham, John M. 2020. "Update on the Gestational Effects of Maternal Hyperthermia." *Birth Defects*
1184    *Research* 112 (12): 943–52. https://doi.org/10.1002/bdr2.1696.

1185 Graveley, Brenton R., Angela N. Brooks, Joseph W. Carlson, Michael O. Duff, Jane M. Landolin, Li
1186    Yang, Carlo G. Artieri, et al. 2011. "The Developmental Transcriptome of Drosophila
1187    Melanogaster." *Nature* 471 (7339): 473–79. https://doi.org/10.1038/nature09715.

1188 Graziosi, G., F. Micali, R. Marzari, F. de Cristini, and A. Savoini. 1980. "Variability of Response of
1189    EarlyDrosophila Embryos to Heat Shock." *Journal of Experimental Zoology* 214 (2): 141–45.
1190    https://doi.org/10.1002/jez.1402140203.

1191 Harris, Tony J C, and Mark Peifer. 2004. "Adherens Junction-Dependent and -Independent Steps in
1192    the Establishment of Epithelial Cell Polarity in Drosophila." *The Journal of Cell Biology* 167 (1):
1193    135–47. https://doi.org/10.1083/jcb.200406024.

1194 Herbert, Sébastien, Léo Valon, Laure Mancini, Nicolas Dray, Paolo Caldarelli, Jérôme Gros, Elric
1195    Esposito, et al. 2021. "LocalZProjector and DeProj: A Toolbox for Local 2D Projection and
1196    Accurate Morphometrics of Large 3D Microscopy Images." *BMC Biology* 19 (1): 136.
1197    https://doi.org/10.1186/s12915-021-01037-w.

1198 IPBES. 2019. *The Global Assessment Report of the Intergovernmental Science-Policy Platform on*
1199    *Biodiversity and Ecosystem Services*. https://doi.org/10.5281/zenodo.3831673.

1200 Kapun, Martin, Maite G Barrón, Fabian Staubach, Darren J. Obbard, R Axel W Wiberg, Jorge Vieira,
1201    Clément Goubert, et al. 2020. "Genomic Analysis of European Drosophila Melanogaster
1202    Populations Reveals Longitudinal Structure, Continent-Wide Selection, and Previously
1203    Unknown DNA Viruses." *Molecular Biology and Evolution* 37 (9): 2661–78.
1204    https://doi.org/10.1093/molbev/msaa120.

1205 Khoury, Mark J., and David Bilder. 2020. "Distinct Activities of Scrib Module Proteins Organize
1206    Epithelial Polarity." *Proceedings of the National Academy of Sciences of the United States of*
1207    *America* 117 (21): 11531–40. https://doi.org/10.1073/pnas.1918462117.

1208 ———. 2022. "Minimal Functional Domains of the Core Polarity Regulator Dlg." *Biology Open* 11 (7).
1209    https://doi.org/10.1242/bio.059408.

1210 Kolaczkowski, Bryan, Andrew D. Kern, Alisha K. Holloway, and David J. Begun. 2011. "Genomic
1211    Differentiation between Temperate and Tropical Australian Populations of Drosophila
1212    Melanogaster." *Genetics* 187 (1): 245–60. https://doi.org/10.1534/genetics.110.123059.

1213 Kuntz, Steven G., and Michael B. Eisen. 2014. "Drosophila Embryogenesis Scales Uniformly across
1214    Temperature in Developmentally Diverse Species." *PLoS Genetics* 10 (4): e1004293.
1215    https://doi.org/10.1371/journal.pgen.1004293.

1216 ———. 2015. "Oxygen Changes Drive Non-Uniform Scaling in Drosophila Melanogaster
1217    Embryogenesis." *F1000Research* 4 (May): 1102.
1218    https://doi.org/10.12688/f1000research.7221.1.

1219 Lecuit, Thomas, and Alpha S. Yap. 2015. "E-Cadherin Junctions as Active Mechanical Integrators in
1220    Tissue Dynamics." *Nature Cell Biology* 17 (5): 533–39. https://doi.org/10.1038/ncb3136.

1221 Lemke, Steffen, Girish Kale, and Silvia Urbansky. 2020. "Comparing Gastrulation in Flies: Links
1222    between Cell Biology and the Evolution of Embryonic Morphogenesis." *Mechanisms of*

Development 164 (October): 103648. https://doi.org/10.1016/j.mod.2020.103648.

Lockwood, Brent L., Cole R. Julick, and Kristi L. Montooth. 2017. "Maternal Loading of a Small Heat
Shock Protein Increases Embryo Thermal Tolerance in Drosophila Melanogaster." *The Journal*
of Experimental Biology 220 (Pt 23): 4492–4501. https://doi.org/10.1242/jeb.164848.

Lu, Xuemin, Jennifer M. Li, Olivier Elemento, Saeed Tavazoie, and Eric F. Wieschaus. 2009. "Coupling
of Zygotic Transcription to Mitotic Control at the Drosophila Mid-Blastula Transition."
Development (Cambridge, England) 136 (12): 2101–10. https://doi.org/10.1242/dev.034421.

Lynch, Jeremy A., Ezzat El-Sherif, and Susan J. Brown. 2012. "Comparisons of the Embryonic
Development of Drosophila , Nasonia , and Tribolium." *Wiley Interdisciplinary Reviews:*
Developmental Biology 1 (1): 16–39. https://doi.org/10.1002/wdev.3.

Machado, Heather E., Alan O. Bergland, Katherine R. O'Brien, Emily L. Behrman, Paul S. Schmidt, and
Dmitri A. Petrov. 2016. "Comparative Population Genomics of Latitudinal Variation in
Drosophila Simulans and Drosophila Melanogaster." *Molecular Ecology* 25 (3): 723–40.
https://doi.org/10.1111/mec.13446.

Mapelli, Marina, and Cayetano Gonzalez. 2012. "On the Inscrutable Role of Inscuteable: Structural
Basis and Functional Implications for the Competitive Binding of NuMA and Inscuteable to
LGN." *Open Biology* 2 (8): 120102. https://doi.org/10.1098/rsob.120102.

Martin, Adam C. 2020. "The Physical Mechanisms of Drosophila Gastrulation: Mesoderm and
Endoderm Invagination." *Genetics* 214 (3): 543–60.
https://doi.org/10.1534/genetics.119.301292.

Mazumdar, Aveek, and Manjari Mazumdar. 2002. "How One Becomes Many: Blastoderm
Cellularization in Drosophila Melanogaster." *BioEssays : News and Reviews in Molecular,*
Cellular and Developmental Biology 24 (11): 1012–22. https://doi.org/10.1002/bies.10184.

McCartney, Brooke M., Donald G. McEwen, Elizabeth Grevengoed, Paul Maddox, Amy Bejsovec, and
Mark Peifer. 2001. "Drosophila APC2 and Armadillo Participate in Tethering Mitotic Spindles to
Cortical Actin." *Nature Cell Biology* 3 (10): 933–38. https://doi.org/10.1038/ncb1001-933.

McColl, Gawain, and Stephen W. McKechnie. 1999. "The Drosophila Heat Shock Hsr-Omega Gene:
An Allele Frequency Cline Detected by Quantitative PCR." *Molecular Biology and Evolution* 16
(11): 1568–74. https://doi.org/10.1093/oxfordjournals.molbev.a026069.

Miller, Morton W., and Charles C. Church. 2013. "Arrhenius Thermodynamics and Birth Defects:
Chemical Teratogen Synergy. Untested, Testable, and Projected Relevance." *Birth Defects*
Research. Part C, Embryo Today : Reviews 99 (1): 50–60. https://doi.org/10.1002/bdrc.21025.

Oakeshott, J. G., S. W. McKechnie, and G. K. Chambers. 1984. "Population Genetics of the
Metabolically Related Adh, Gpdh and Tpi Polymorphisms in Drosophila Melanogaster I.
Geographic Variation in Gpdh and Tpi Allele Frequencies in Different Continents." *Genetica* 63
(1): 21–29. https://doi.org/10.1007/BF00137461.

Paaby, Annalise B., Alan O. Bergland, Emily L. Behrman, and Paul S. Schmidt. 2014. "A Highly
Pleiotropic Amino Acid Polymorphism in the Drosophila Insulin Receptor Contributes to Life-
History Adaptation." *Evolution; International Journal of Organic Evolution* 68 (12): 3395–3409.
https://doi.org/10.1111/evo.12546.

Pietzsch, Tobias, Stephan Saalfeld, Stephan Preibisch, and Pavel Tomancak. 2015. "BigDataViewer:
Visualization and Processing for Large Image Data Sets." *Nature Methods* 12 (6): 481–83.
https://doi.org/10.1038/nmeth.3392.

Preibisch, Stephan, Stephan Saalfeld, and Pavel Tomancak. 2009. "Globally Optimal Stitching of Tiled
3D Microscopic Image Acquisitions." *Bioinformatics (Oxford, England)* 25 (11): 1463–65.

https://doi.org/10.1093/bioinformatics/btp184.

Pritchard, David K., and Gerold Schubiger. 1996. "Activation of Transcription in Drosophila Embryos
Is a Gradual Process Mediated by the Nucleocytoplasmic Ratio." *Genes & Development* 10 (9):
1131–42. https://doi.org/10.1101/gad.10.9.1131.

Raguso, Robert A. 2020. "Don't Forget the Flies: Dipteran Diversity and Its Consequences for Floral
Ecology and Evolution." *Applied Entomology and Zoology* 55 (1): 1–7.
https://doi.org/10.1007/s13355-020-00668-9.

Rand, David M., Daniel M. Weinreich, Daniel Lerman, Donna Folk, and George W. Gilchrist. 2010.
"Three Selections Are Better than One: Clinal Variation of Thermal QTL from Independent
Selection Experiments in Drosophila." *Evolution* 64 (10): 2921–34.
https://doi.org/10.1111/j.1558-5646.2010.01039.x.

Rim, Ellen Youngsoo, Hans Clevers, and Roel Nusse. 2022. "The Wnt Pathway: From Signaling
Mechanisms to Synthetic Modulators." *Annual Review of Biochemistry* 91 (June): 571–98.
https://doi.org/10.1146/annurev-biochem-040320-103615.

Rizzelli, Francesca, Maria Grazia Malabarba, Sara Sigismund, and Marina Mapelli. 2020. "The
Crosstalk between Microtubules, Actin and Membranes Shapes Cell Division." *Open Biology* 10
(3): 190314. https://doi.org/10.1098/rsob.190314.

Rudman, Seth M., Sharon I. Greenblum, Subhash Rajpurohit, Nicolas J. Betancourt, Jinjoo Hanna,
Susanne Tilk, Tuya Yokoyama, Dmitri A. Petrov, and Paul S. Schmidt. 2022. "Direct Observation
of Adaptive Tracking on Ecological Time Scales in Drosophila." *Science (New York, N.Y.)* 375
(6586): eabj7484. https://doi.org/10.1126/science.abj7484.

Rutherford, Suzanne L., and Susan Lindquist. 1998. "Hsp90 as a Capacitor for Morphological
Evolution." *Nature* 396 (6709): 336–42. https://doi.org/10.1038/24550.

Sangster, Todd A., Susan Lindquist, and Christine Queitsch. 2004. "Under Cover: Causes, Effects and
Implications of Hsp90-Mediated Genetic Capacitance." *BioEssays : News and Reviews in
Molecular, Cellular and Developmental Biology* 26 (4): 348–62.
https://doi.org/10.1002/bies.20020.

Sawyer, Lesley A., Federica Sandrelli, Carlo Pasetto, Alexandre A. Peixoto, Ezio Rosato, Rodolfo
Costa, and Charalambos P. Kyriacou. 2006. "The Period Gene Thr-Gly Polymorphism in
Australian and African Drosophila Melanogaster Populations: Implications for Selection."
Genetics 174 (1): 465–80. https://doi.org/10.1534/genetics.106.058792.

Schmidt, Anja, and Jörg Grosshans. 2018. "Dynamics of Cortical Domains in Early Drosophila
Development." *Journal of Cell Science* 131 (7). https://doi.org/10.1242/jcs.212795.

Schmidt, Paul S., David D. Duvernell, and Walter F. Eanes. 2000. "Adaptive Evolution of a Candidate
Gene for Aging in Drosophila." *Proceedings of the National Academy of Sciences of the United
States of America* 97 (20): 10861–65. https://doi.org/10.1073/pnas.190338897.

Schmidt, Paul S., and Annalise B. Paaby. 2008. "Reproductive Diapause and Life-History Clines in
North American Populations of Drosophila Melanogaster." *Evolution* 62 (5): 1204–15.
https://doi.org/10.1111/j.1558-5646.2008.00351.x.

Sezgin, Efe, David D. Duvernell, Luciano M. Matzkin, Yihao Duan, Chen Tseh Zhu, Brian C. Verrelli,
and Walter F. Eanes. 2004. "Single-Locus Latitudinal Clines and Their Relationship to Temperate
Adaptation in Metabolic Genes and Derived Alleles in Drosophila Melanogaster." *Genetics* 168
(2): 923–31. https://doi.org/10.1534/genetics.104.027649.

Telley, Ivo A., Imre Gáspár, Anne Ephrussi, and Thomas Surrey. 2012. "Aster Migration Determines
the Length Scale of Nuclear Separation in the Drosophila Syncytial Embryo." *The Journal of Cell*

Biology 197 (7): 887–95. https://doi.org/10.1083/jcb.201204019.

Umina, P. A., A. R. Weeks, M. R. Kearney, S. W. McKechnie, and A. A. Hoffmann. 2005. "Evolution: A
Rapid Shift in a Classic Clinal Pattern in Drosophila Reflecting Climate Change." *Science* 308
(5722): 691–93. https://doi.org/10.1126/science.1109523.

Vazquez, Julio, Daniel Pauli, and Alfred Tissières. 1993. "Transcriptional Regulation in Drosophila
during Heat Shock: A Nuclear Run-on Analysis." *Chromosoma* 102 (4): 233–48.
https://doi.org/10.1007/BF00352397.

Verrelli, B. C., and W. F. Eanes. 2001. "Clinal Variation for Amino Acid Polymorphisms at the Pgm
Locus in Drosophila Melanogaster." *Genetics* 157 (4): 1649–63.
https://doi.org/10.1093/genetics/157.4.1649.

Vigue, C. L., and F. M. Johnson. 1973. "Isozyme Variability in Species of the Genus Drosophila. VI.
Frequency-Property-Environment Relationships of Allelic Alcohol Dehydrogenases in D.
Melanogaster." *Biochemical Genetics* 9 (3): 213–27.

Walter, Marika F., Nancy S. Petersen, and Harald Biessmann. 1990. "Heat Shock Causes the Collapse
of the Intermediate Filament Cytoskeleton in Drosophila Embryos." *Developmental Genetics* 11
(4): 270–79. https://doi.org/10.1002/dvg.1020110405.

Weeks, A. R., S. W. Mckechnie, and A. A. Hoffmann. 2006. "In Search of Clinal Variation in the Period
and Clock Timing Genes in Australian Drosophila Melanogaster Populations." *Journal of
Evolutionary Biology* 19 (2): 551–57. https://doi.org/10.1111/j.1420-9101.2005.01013.x.

Welte, Michael A., Joan M. Tetrault, Robert P. Dellavalle, and Susan L. Lindquist. 1993. "A New
Method for Manipulating Transgenes: Engineering Heat Tolerance in a Complex, Multicellular
Organism." *Current Biology* 3 (12): 842–53. https://doi.org/10.1016/0960-9822(93)90218-D.

Xie, Yi, and J. Todd Blankenship. 2018. "Differentially-Dimensioned Furrow Formation by Zygotic
Gene Expression and the MBT." *PLoS Genetics* 14 (1): e1007174.
https://doi.org/10.1371/journal.pgen.1007174.

Zheng, Liuliu, Leonardo A. Sepúlveda, Rhonald C. Lua, Olivier Lichtarge, Ido Golding, and Anna Marie
Sokac. 2013. "The Maternal-to-Zygotic Transition Targets Actin to Promote Robustness during
Morphogenesis." *PLoS Genetics* 9 (11): e1003901.
https://doi.org/10.1371/journal.pgen.1003901.

1342

Figure 1

Increased embryonic lethality at elevated temperature

A Temperature regimes during embryonic development

(N) Normal temperature = 25°C   (E) Elevated temperature = 29°C

B Lethality at the end of embryo development

Nuclear loss phenotype at elevated temperature

25°C   29°C (mild defect)   29°C (severe defect)

Mitotic failure and pairwise nuclear expulsion
(Example from embryo at 29°C)

F Histone-RFP

−5 min   0 min   5 min   10 min   15 min

G number of mitotic failures causing nuclear expulsions during interphase 14

H

Normal temperature   Elevated temperature

• • Normal mitosis   ⚊⚊ Mitotic failure and pairwise loss of nuclei

Figure 2

Asynchrony in the phases of nuclear division cycles
(Example from embryo at 29°C)

Telophase | Anaphase | Metaphase | Anaphase

Histone-RFP

A

Anterior 1/5th | Central 1/5th | Posterior 1/5th

B Asynchrony at interphase 14

***
***
*

Asynchrony (min)

22°C 25°C 28°C 29°C

C Nuclear cycles **11**, **12**, and **13** in different regions of the embryo

Normal temperature

Elevated temperature

Asynchrony at the beginning of interphase 14

Increased nuclear crowding at elevated temperature

Histone-RFP | Pseudo-cells | Distribution of pseudo-cell areas

D 22°C

E 25°C

F 28°C

G 29°C

H

I

J

K

L pseudo-cell area (μm²)

M pseudo-cell area (μm²)

N pseudo-cell area (μm²)

O pseudo-cell area (μm²)

Anterior-posterior position (μm)

25 μm² 100 μm²

P crowding at the end of interphase 13

***
*

Quadratic coefficient

0.002

0.001

0.000

22°C 25°C 28°C 29°C

Q Normal temperature | Elevated temperature

Moderate nuclear crowding in the central region

Nuclear crowding intensifies in the central region

Figure 3

Figure 4

Over-expressing regulators of 'cell polarity signalling during mitosis'
to rescue embryonic lethality at elevated temperature

Over-expressing regulators of 'F-actin -microtubule interaction during mitosis'
to rescue embryonic lethality at elevated temperature

"Natural" candidate genes varying along thermoclines

Figure S1

Figure S2

Nuclear loss phenotype due to elevated temperature

Figure S3

Histone-RFP Jupiter-GFP (Example from embryo at 25°C)

A — X-Y view

B — Y-Z view

C — out-of-plane divisions during mitosis-10

D — isolated nuclear expulsion (Example from embryo at 25°C)

E — number of isolated nuclear expulsions during interphase 14

Figure S4

A — Normal temperature (22°C)

B — Elevated temperature (29°C)

Interphase-12 | mitosis-12 | Interphase-13

● Cortical nuclei that failed to divide in mitosis 10   ● Cortical nuclei that failed to divide in mitosis 11

Figure S5

Figure S6

Figure S7

Histone-RFP

A

25°C

−03:00   mitosis 13

00:00   interphase 14…
… in the anterior

00:30   ←

01:00   … in the central region
… in the posterior

anterior 1/5th   central 1/5th   posterior 1/5th

B

29°C

−03:00   mitosis 13

00:00   interphase 14…
… in the anterior

00:30   ←
… in the posterior

02:00   … in the central region

Figure S8

Figure S9

Figure S10

Figure S11