Elevated temperature fatally disrupts nuclear divisions in the early *Drosophila* embryo.

Girish Kale[1],*, Pratika Agarwal[1], J Jaime Diaz-Larrosa[1], Steffen Lemke[1,2],*

[1] Centre for Organismal Studies, University of Heidelberg, Heidelberg. Germany
[2] Institute of Biology, University of Hohenheim, Stuttgart, Germany
* Correspondance: G.K. (girish.kale@cos.uni-heidelberg.de), S.L. (steffen.lemke@uni-hohenheim.de)

Abstract:
Temperature fluctuations can challenge animal survival, especially in embryos of animals that can't actively
regulate their body temperature. Previous work in insects suggests that spatial gradients and temporal
variations in temperature can affect embryonic development in a region-specific manner. Yet, it is unclear if
elevated temperature produces developmental defects, and if so, how the disruption may manifest. Here we
use early embryonic development of the fruit fly *Drosophila melanogaster* as a model to address
consequences of elevated temperature on fly embryonic development.

We identified the syncytial blastoderm stage as a critical time window, being particularly sensitive to elevated
temperature. The syncytial blastoderm stage involves 4 rounds of meta-synchronous nuclear divisions at the
embryo cortex, followed by cellularization to form the cellular blastoderm. Live imaging these processes at
elevated temperature revealed an increase in mitotic failures and a loss of cortical nuclei, both, during
syncytial- and cellular-blastoderm stage. Additionally, embryos developing at elevated temperature show a
local crowding of nuclei and an increase in asynchrony between the nuclear cycles in the middle vs.
anterior/posterior poles. Interestingly, nuclear crowding and nuclear cycle asynchrony cooperatively amplify
the frequency of mitotic failures, leading to holes in the blastoderm epithelium. Strikingly, the same
treatment also increased embryonic lethality. We propose that the loss of nuclei results in a loss of patterning
information and the abrogation of proper embryonic development. We identified genes that regulate
mitosis, especially the ones mediating the interaction between microtubules and F-actin, and functionally
tested their capacity to rescue the embryonic lethality.

Our results expose a vulnerability at elevated temperatures in the interaction between cortical actin and
microtubules, which leads to mitotic failures, and can be potentially rescued by modulating the expression
of just a few factors. We propose that the expression levels of the corresponding genes could be used as
indicators to predict fitness effects on insect populations that are exposed to increasing temperature
variations.

Keywords:
Syncytial blastoderm, elevated temperature, microtubule - F-actin interaction, nuclear division defects,
embryo lethality, genetic rescue, *Drosophila*

Introduction:

Temperature is a key factor determining the fidelity of embryonic development. Studies have shown occurrence of birth defects due to hyperthermia during pregnancy (Edwards 2006; Miller and Church 2013; Graham 2020), in spite of active regulation of core temperature. Here, the cell biological basis for the developmental defects due to an exposure to high temperature, remains elusive. It is currently unclear whether this phenomenon could be observed and studied also in invertebrate species, for which the developmental process is well described, allowing for a quantitative characterization of the potential developmental defects arising due to high temperature. In this context, we explore whether the exposure to elevated temperature is detrimental to embryonic development in *Drosophila melanogaster* (henceforth, *Drosophila*), addressing the cell biological basis for possible defects, and asking what are the consequences in terms of population level parameters, such as embryo survival.

Previous studies have shown that the early embryonic development of *Drosophila* embryos (first ~6 hours) is especially sensitive to acute heat-shock (Welte et al. 1993), and have indicated that acute heat-shock causes a loss of coordination in developmental processes, leading to developmental defects (Crews, McCleery, and Hutson 2016). Given that the survival after a heat-shock relies on transcription of heat-shock proteins (Graziosi et al. 1980), and that zygotic transcription is progressively activated during maternal-to-zygotic transition (Pritchard and Schubiger 1996), it is likely that the developmental processes prior to the completion of this transition are especially susceptible to heat-shock. In line with these expectations, previous studies have demonstrated a disruption of the normal dynamics of cytoskeletal elements due to an exposure to high temperature (Walter, Petersen, and Biessmann 1990; Zheng et al. 2013; Figard et al. 2019). Even more elusive are the effects of prolonged exposure to elevated temperature below the heat-shock limits. Here we focus on such temperature sensitivity of early embryonic development in *Drosophila*, and ask what is the cell biological basis for developmental defects due to an exposure to elevated temperature, and whether there is a link to the population level consequences on embryo survival.

Post fertilization, the *Drosophila* embryos undergo 9 rounds of syncytial nuclear divisions deep in the yolk. Following these divisions, most nuclei migrate to embryo periphery, where 4 more rounds of nuclear division take place during a stage called, syncytial blastoderm (Foe and Alberts 1983). These nuclear divisions are facilitated by pseudo-cleavage furrows, i.e., temporary membrane invaginations resembling a half-cell that slowly form around the nuclei during interphase-metaphase, and are then rapidly reabsorbed into the embryo cortex during anaphase-telophase (Xie and Blankenship 2018). Following these 13 rounds of nuclear divisions, the resulting ~6000 nuclei simultaneously segregate into individual cells, in the process of cellularization, producing the blastoderm epithelium (Mazumdar and Mazumdar 2002). At this point gastrulation starts, which involves cell shape changes (Martin 2020), cell contact rearrangements (Lecuit and Yap 2015), and further rounds of cell divisions in isolated mitotic domains (Foe 1989): all facilitating the early morphogenesis. Cell biological basis for these early developmental processes is well described, and would allow us to identify the defects at elevated temperature, if any.

In our experiments, we exposed *Drosophila* embryos to the elevated temperature, and compared them with normal temperature controls. Following previous studies, we chose 29°C as elevated temperature (Kuntz and Eisen 2014; Chong, Amourda, and Saunders 2018) and asked whether increased temperature up to 29°C resulted in developmental defects. Notably, this temperature is well below the 32°C and higher range required to elicit a robust heat-shock response (Vazquez, Pauli, and Tissières 1993), and no major developmental defects have been reported so far at 29°C. We found that exposing embryos to elevated temperature during early embryonic development leads to a significant increase in the population level embryonic lethality. When we traced defects from the organism to the cellular level, we observed a loss of nuclei during syncytial- and cellular-blastoderm stage, which was caused by mitotic failures. Additionally, in

embryos developing at elevated temperature, we reported and quantified the nuclear crowding in the central region of the embryo, as well as greater asynchrony in the nuclear division cycles between central regions vs. anterior/posterior poles of the embryo. Moreover, we could show that the nuclear crowding and nuclear cycle asynchrony combinatorially cooperate to locally increase the propensity of mitotic failures. Such local increase in mitotic failures results in holes in the blastoderm, i.e., region devoid of nuclei/cells, to disrupt the continuity of this primary epithelial tissue. We overexpressed a few candidate genes to boost the robustness of nuclear divisions to rescue the mitotic failures defect, focusing on the regulators of cell polarity, microtubule- F-actin interactions, and mitotic spindle orientation, to test their ability to rescue the described defects, and ameliorate the embryo lethality at elevated temperature. Our results suggest a vulnerability in microtubule- F-actin interaction during the well conserved process of nuclear division, when occurring at elevated temperature. Our work establishes a novel link between cell biological defects due to an exposure to elevated temperature, and a reduction in organismal fitness at population level at elevated temperature.

Results:

Early exposure to elevated temperature reduces embryo survival.
To test whether elevated temperature had an effect on developmental robustness during the first hours of embryonic development, we exposed young embryos to 29°C and asked whether this would alter the fraction of embryos that hatch into larvae. To associate possible fitness effects with defined developmental windows (see above), embryos were exposed to elevated temperature either from deposition till before the onset of gastrulation (~3 hours) or for a duration of ~3 hours from the onset of gastrulation (Figure 1A), to keep the temperature exposures comparable. In both cases we observed a significant increase in embryonic lethality (Figure 1B). Our findings reveal a particular vulnerability during the syncytial phase of the embryonic development and suggest that elevated temperature leads to defects in the setup of the insect blastoderm. This result is surprising, especially because previous studies indicated that genetic patterning and setting up of positional information within the blastoderm was overall robust against temperature fluctuations (Lucchetta et al. 2005; Niemuth and Wolf 1995).

In addition to genetic patterning, nuclear dynamics could be challenged during the first hours of embryonic development. In this developmental window, nuclei divide deep in the yolk, then translocate to the embryo cortex, where they need to be anchored robustly while undergoing further rounds of nuclear division. A normal distribution of nuclei ultimately determines a normal morphogenesis of the blastoderm epithelium, setting the foundation for the developmental processes to follow. Thus, to address whether defects in the blastoderm could stem from defects in nuclear dynamics, we followed nuclear dynamics till the end of blastoderm formation using Histone-RFP to visualize nuclei, and Jupiter-GFP to visualize microtubules, keeping track of nuclear division cycles. Starting with nuclear migration from the yolk to periphery (occurring during the interphase of the 10th nuclear cycle), we observed nuclei through their peripheral divisions during syncytial blastoderm (through mitoses 10 to 13), and then through cellularization (interphase 14), until the onset of gastrulation could be inferred from the movement and division of nuclei (beyond interphase 14). During this time window, we did not observe any significant defects, in the nuclear translocation from deep inside the yolk to embryo cortex, in embryos developing at elevated temperature. However during the syncytial blastoderm stage, embryos that developed at 29°C much more frequently "lost" nuclei from their cortical layer than if they developed at 25°C, i.e., nuclei that were initially positioned within the cortical layer of the egg, appeared to get expelled, descended into the yolk, and did not become part of the embryonic blastoderm (Figure 1C-E), with a variability in the extent of the nuclear loss, producing holes in the blastoderm of a fraction of embryos (Figure S1). Further, in the embryos fixed and stained with DRAQ5, we imaged the nuclear distribution in the entire embryo volume, to observe that the loss of nuclei wasn't symmetric in the two halves of the embryos, and albeit in the central region of the embryos, the location of

the nuclear loss varied across embryos (Figure S2). The embryo-to-embryo variability, both in the distribution
and intensity of nuclear loss argued for the possibility that these defects were self-organized rather than
dictated by the genetic make-up.

Taken together, we identified the syncytial blastoderm stage of embryo development as the most sensitive
window, based on the embryo lethality analysis. Additionally, the loss of nuclei during this stage indicated a
defect in nuclear anchorage at the embryo cortex, rather than nuclear divisions deep in the yolk, or the
translocation of nuclei from deep inside the yolk to the embryo cortex.

A closer observation of live imaged embryos revealed three striking phenomena during blastoderm formation
in embryos developing at elevated temperature compared to normal temperature controls: an extensive loss
of nuclei from the egg cortex (as above), a time shift and slowdown of nuclear divisions cycles in the central
region of the embryo compared to its poles, and nuclear crowding in the central region of the embryo. Thus,
we decided to quantitatively describe these defects, in an attempt to describe the necessary conditions to
produce a blastoderm hole, and link these defects to the embryo lethality phenotype.

At elevated temperature, embryos lose nuclei from the cortex after defective divisions.
A closer look at the loss of cortical nuclei indicated that the nuclei were lost either individually or as pairs of
two adjacent nuclei. We could also observe periods of drastic displacement of nuclei at elevated temperature
(see below). Based on these observations, we imagined various mechanisms for the observed loss of nuclei.
For instance, individual nuclei might be lost due to out-of-plane nuclear divisions, while a pair of nuclei might
be lost due to a defective in-plane nuclear division. In case of defects in nuclear divisions, it is possible that
the cell cycle checkpoints are triggered. Nuclei being dislodged from the embryo cortex during the drastic
displacements could also explain both individual and pairwise loss, though in such scenario, the pairs of nuclei
need to be somehow associated with each other, influencing each other's loss. Here we considered these
possibilities, and asked to what extent they account for the observed loss of nuclei.

To address whether cortical nuclei appeared as getting lost because of an increased frequency of out-of-
plane divisions, we visually tracked nuclei from their last division to the time they were no longer associated
with the cortex. We could detect out-of-plane divisions in embryos at temperatures ranging from 22 to 29°C,
only during the mitoses of the 10th nuclear cycle. These out-of-plane divisions could be easily identified, as
they formed a characteristic mitotic spindle that was oriented perpendicular to the embryo surface (Figure
S3A, B), and after the division only one of the daughter nuclei remained attached to the cortex. However, the
overall frequency of out-of-plane divisions was very low (<5%): neither did it differ significantly between any
of the observed temperatures (Figure S3C), nor could it explain the loss of nuclei during later nuclear cycles.
Similarly, losses of individual nuclei after an in-plane division were rather infrequent, and did not differ
significantly within the range of observed temperatures (Figure S3D, E). This suggested that the increased
loss of nuclei observed in embryos developing at 29°C could neither be accounted for with individual nuclei
dropping into the yolk, irrespective of whether the loss was due to them being mechanically dislodged.

For the observed pairwise loss of nuclei, our tracking revealed that these were sister nuclei born from the
same mitosis. This suggested that specific characteristics of cortical nuclear divisions might determine the
future fate of the newborn nuclei and trigger their release from the egg cortex, possibly in response to failed
checkpoints during mitosis. In the pairwise loss of nuclei, we observed chromosome segregation defects
(Figure 1F) that could lead to aneuploidy, suggesting that the subsequent expulsion of daughter nuclei from
the egg cortex was due to a prior defective mitosis. Following defective mitoses at normal or elevated
temperature, we could not detect a clear time course of nuclear expulsion: these nuclei could be lost in the
immediately following or in later interphases (Figure S4), suggesting that nuclear expulsion after defective
mitosis was a passive process and not triggered by molecular signaling at failed checkpoints.

To confirm that the observed loss of nuclei at 29°C could be attributed to an overall increase in defective
mitoses, we asked if the frequency of defective mitoses was changing depending on the temperature. To
address this question, we focused on interphase 14, as that's when we observed the most nuclear losses,
and counted the pairwise loss of nuclei after mitotic failures at 22, 25, 28, and 29°C (see methods). While the
frequency of mitotic failures was overall low at 22 to 28°C, it increased significantly at 29°C (Figure 1G). Taken
together, our findings identify nuclear divisions in the blastoderm as a thermolabile cellular process (Figure
1H) that shows a disproportionate increase in failures at the transition from 28°C to 29°C, resulting in the loss
of sister nuclei from the egg cortex, when born in defective mitoses. Interestingly, in a fraction of embryos
the mitotic failures occurred in great numbers and in close vicinity, leading to the holes in the blastoderm
epithelium (Figure 1G, hollow markers). Thus, the formation of blastoderm holes (Figure S1) could be directly
attributed to clustered mitotic failures, though it remained unclear why some embryos have the of mitotic
failures in local clusters, while others didn't.

To explore the potential cause of mitotic failures at elevated temperatures, we tracked nuclear clusters
starting from individual mother nucleus (end of interphase 10) to the 8 great-granddaughter nuclei (end of
interphase 13), in order to quantitatively describe their movement (Figure S5A-C, also see methods). This
allowed us to assess mechanical stresses that would be experienced by nuclei dividing at the blastoderm
cortex, as we distinguished the movement of nuclear clusters during interphase-metaphase vs. anaphase-
telophase. Our results showed that the nuclear clusters had 2 components to their movement. First, the
clusters exhibited a constant drift-like movement, which we captured using a linear fit to the displacement
of the center of the nuclear clusters over time (Figure S5B dotted lines). The slope of the linear fit indicated
that the clusters drifted towards the posterior end (Figure S5D, E). The second component was composed of
deviations in the centers of nuclear cluster from the linear fit, i.e., the displacements relative to the first
component. These deviations were significantly greater at elevated temperature, both during the interphase-
metaphase period (Figure S5F, G) and during the anaphase-telophase period (Figure S5H, I), though the
displacements along AP-axis were more drastic than those along DV-axis, and those during anaphase-
telophase were 2-3-fold greater than those during interphase-metaphase. For context, the length scale of
the displacements during anaphase-telophase could be up to 4-times the length of the spindle itself,
indicating an extensive propensity for displacement in x-y plane, especially along the AP axis. We also
observed that the spindles were getting displaced in z-axis, again, during anaphase-telophase period. In
embryos developing at 29°C, we could see two waves of nuclear displacement, originating from the anterior
and the posterior pole, meeting at the central region of the embryo (Figure S6). Overall, these quantifications
and observations indicated that the mitotic spindles are displaced drastically while nuclei are dividing, and
argued for the possibility of mechanical defects during nuclear divisions.

Taken together, we observed the effects of exposure to elevated temperature, at organismal level as embryo
lethality (Figure 1B), at tissue level as blastoderm holes (Figure 1E, Figure S1, S2), and at cellular level as
mitotic failures (Figure 1F, G). Embryos with most severe form of mitotic failure phenotype had blastoderm
hole, while the fraction of embryos with large blastoderm holes correlated well with increase in embryo
lethality, indicting a link from cellular till organismal phenotype. Given that the loss of nuclei occurs after
mitosis, and that the nuclei move drastically during anaphase-telophase period, we further argue that nuclear
anchorage is specifically challenged during mitosis.

Nuclei divide with a time shift and are crowded in embryos at elevated temperature.
The most severe loss of nuclei was often observed in the central domain of the embryo (Figure 1C-E, Figure
S1 and S2). Notably, these areas coincided with the "collision" of two waves of nuclear displacements along
z-axis (Figure S6), i.e., the future apical-basal axis in the blastoderm cells (A. Schmidt and Grosshans 2018).
We could further observe that, not only anaphase, but all other phases of nuclear division were delayed in

the central region of the embryos developing at 29°C. This resulted in regions of embryo being segregated
into domains of nuclear division phases (Figure 2A). Conversely, the same phase of nuclear division cycle
could be observed at different timepoints in different regions of the embryo (Figure S7). Thus, we decided to
quantify this asynchrony of nuclear division cycles as a function of temperature (see methods). Our
quantifications show that the asynchrony significantly increases in embryos developing at 29°C as compared
to other temperatures (Figure 2B). Thus, we quantitatively describe the defect of asynchrony in nuclear cycles
at elevated temperature (Figure 2C).

The central region of the embryo, which corresponded to the delayed nuclear division cycles, also had more
tightly packed nuclei. To quantify nuclear crowding, we focused on the last nuclear division cycle and its
preceding interphase, i.e., interphase 13 (see methods), as that's the time we observed maximum nuclear
crowding. To quantify the space available for cortical blastoderm nuclei, we segmented the nuclei, and used
Voronoi tessellation to define a region around each nucleus, which we call pseudo-cells. The areas of pseudo-
cells were rather homogeneous in embryos developing at 22°C (Figure 2D, H, L). Already at 25°C (Figure 2E,
I, M) and 28°C (Figure 2F, J, n), the available space for cortical nuclei was no longer uniform across the
embryo, and it showed signs of local crowding. At 29°C, the space available for nuclei along the anterior-
posterior axis varied substantially and revealed strong nuclear crowding in the central region of the embryo
(Figure 2G, K, O). We quantified the extent of crowding (see methods) as a function of temperature, and
found a progressive increase in the nuclear crowding (Figure 2P). Thus, we also quantitatively describe the
defect of nuclear crowding at elevated temperature (Figure 2Q).

Taken together, besides mitotic failure, we have two additional defects at elevated temperature, nuclear
crowding and nuclear cycle asynchrony. We also note that the severity of these defects (based on the
quantifications) can vary between embryos, e.g., the embryos with most extensive failure in mitosis
(blastoderm holes) don't have the highest levels of nuclear crowding (Figure 2P), or strongest asynchrony
(Figure 2B), etc. Though, based on the quantifications and statistical significances we note that; one, severe
mitotic failures (blastoderm holes) appear in a threshold-effect-like manner, appearing only in a fraction of
embryos at 29°C; two, the nuclear crowding and asynchrony in nuclear cycles appears progressively as
function of temperature, although the progression is not linear; and three, the fraction of embryos with large
blastoderm holes best explains the increase in embryo lethality. Thus, we hypothesize a hierarchy of the
defects, purely based on the fraction of embryos in which they appear as a function of temperature: exposure
to elevated temperature will increase the frequency of mitotic failures as primary defect; nuclear crowding
and nuclear cycle asynchrony will secondarily dictate the distribution of mitotic failures, producing extensive
mitotic failures (blastoderm holes) as the ultimate effect in a fraction of embryos; eventually leading to a
reduction in embryo fitness and an increase in embryo lethality.

Regions of embryo with greater nuclear crowding and with delayed nuclear cycles are associated with
greater frequency of mitotic failures.
To test whether nuclear crowding and nuclear cycle asynchrony could influence the propensity of mitotic
failures, we decided to compute the fraction of mitotic failures occurring in the regions of the embryo that
had nuclear crowding and/or delayed nuclear division cycles. We constructed regions of the embryo with
maximum nuclear crowding (crowded regions), maximum delay in nuclear division cycles (delayed regions),
and using an overlap of crowded and delayed regions (combined region) (see methods).

We defined the crowded regions (Figure 3A), and delayed regions (Figure 3B) based of the distribution and
dynamics of nuclei in live imaged embryos. Then, we overlayed the map of mitotic failures on the crowded
regions (Figure 3C), delayed regions (Figure 3D), and the combined regions (Figure 3E). The crowded and
delayed regions had rather comparable areas across temperatures (Figure 3F, G), indicating that any

difference in the fraction of mitotic defects would be solely due to the nuclear crowding or due to the delays
in nuclear cycles. Interestingly, the areas of the combined regions were smaller at elevated temperature
(Figure 3H), indicating that there was an increase in the overlap between the crowded and delayed regions,
and that we should expect a similar reduction in the fraction of mitotic failures, if the mitotic failures were
distributed randomly.

The fraction of mitotic failures in crowded region increased progressively as function of temperature,
reaching statistical significance for comparison between 22 and 29°C (Figure 3I), while the fraction of mitotic
failures in delayed regions did not increase progressively, showing an abrupt increase from 28 to 29°C with
only marginal statistical significance (Figure 3J). Interestingly, we see that the smaller area fractions in the
combined regions correspond to even greater fraction of mitotic failures (Figure 3K). Additionally, the
distribution of the fraction of mitotic failures in the combined region was unimodal, in contrast to the bimodal
distribution of mitotic failures themselves (Figure 1G). Thus, we conclude that the combinatorial effect of
nuclear crowding and asynchrony significantly increases the propensity of mitotic failures. We would further
argue that nuclear crowding has a stronger impact on the propensity of mitotic failures than that due to
nuclear cycle asynchrony.

Taken together, we find large clusters of defective mitoses and subsequent loss of nuclei in the center of
about 10-20% of embryos developing at 29°C (Figure 1G). Nuclear crowding and nuclear cycle asynchrony
cooperate to produce "hot spots" of mitotic failures (Figure 3). The aggregated losses of nuclei in the center
of the embryo persist until the end of blastoderm formation and leads to visible holes in the newly formed
embryonic epithelium (Figure S1). These holes indicate a severe disruption in epithelial integrity and
approximately match the increased rate of embryonic lethality for development at 29°C (Figure 1B). Going
further, we attempted to rescue the embryo lethality defect using over-expression of various candidate
genes, and if a rescue was possible, further decided to quantify whether the rescue could be explained
through fraction of mitotic failures in the crowded region and the combined region.

Identifying temperature-sensitive vulnerabilities in a deeply conserved cellular process
The analysis of early fly development at moderately elevated temperatures, in our experiments, has
uncovered the first three hours of fly development as sensitive to environmental perturbations, which can
affect embryo fitness and increase embryonic lethality at elevated temperatures. This increase in embryonic
lethality correlates with specific perturbations in fundamental nuclear dynamics, i.e., the timing and fidelity
of nuclear divisions within the spatial constraints of a cortex densely packed with nuclei. To address how a
deeply conserved and presumably robust process like mitosis could be affected so dramatically at a
temperature lower than those eliciting a heat-shock response, we functionally probed for possible
vulnerabilities in core molecular interactions that control nuclear dynamics in the early fly embryo.

In stark deviation from canonical cytokinesis, cortical nuclear divisions in the syncytial fly embryo rely on cell
polarity cues that are set up at shallow metaphase furrows of the egg cortex, where they orient mitotic
spindles parallel to the cell surface (A. Schmidt and Grosshans 2018; Xie and Blankenship 2018). What's more,
these furrows are transient, and are reestablished during each nuclear cycle: including new growth of
membrane between nuclei, reestablishment of cell polarity, and reorganization of cytoskeletal elements. The
dependence on transient membrane invaginations to establish cell polarity information represents the least
conserved and potentially most labile element of cortical nuclear divisions in insect embryos. To test whether
the fidelity of cortical mitoses at elevated temperatures hinges upon insufficiently robust cell polarity
signaling, we overexpressed two known prominent polarity proteins, Bazooka (baz, a Par-3 homolog) and
Disks-large (dlg) (Harris and Peifer 2004). We asked whether these over-expressions rescue embryo lethality

defect, and if so, whether the increase in the fraction of mitotic failures in crowded regions and in combined
regions is rescued as well.

When we over-expressed either *baz* or *dlg* we did not see a significant change in embryo lethality at control
temperature (25°C) as compared to the control (Figure 4A). The lethality was greater in outcross controls at
elevated temperature (29°C) similar to results in previous experiments (Figure 1B), while the embryos over-
expressing either *baz* or *dlg* faired significantly better at the elevated temperature (Figure 4A). We note
however that the rescue of lethality at elevated temperature in *baz* over-expressing embryos was only
marginally significant, with a lot of variability between samples. Additionally, *dlg* over-expressing embryos at
normal temperature showed higher variability in embryo lethality than their outcross control counterparts.
Consistent with these results, *baz* over-expression did not rescue the increase in the fraction of mitotic
failures, either in the crowded regions, or in the combined regions at elevated temperature (Figure 4B, C),
though the large blastoderm holes were absent in these embryos. Then, *dlg* over-expression rescued the
increase in fraction of mitotic failures only in the crowded region at elevated temperature (Figure 4B, C), and
while the large blastoderm holes were absent in these embryos, *dlg* over-expression increased the fraction
of mitotic defects at normal temperature in both, the crowded region and in the combined region,
occasionally producing small blastoderm holes (Figure 4B, C).

Above results indicated a temperature-dependent vulnerability more specific to the molecular core of the
mitotic spindle rather than its setup in the context of a syncytial embryo. To faithfully pull sister chromatids
towards the actin cortex on either side of the division plane, the execution of mitosis relies on tightly
regulated interactions between the actin cortex and the microtubules of the mitotic spindle (Cao et al. 2010;
Rizzelli et al. 2020). In *Drosophila*, these critical interactions of actin and microtubules have been associated
with a complex of three proteins: α-Catenin (αCat), which binds to the actin cortex; β-Catenin (βCat, or
armadillo in Drosophila), which binds to α-Catenin; and Adenomatous polyposis coli 2 (Apc2), which links to
βCat and binds to the microtubules of the mitotic spindle (McCartney et al. 2001). Within this tripartite
complex, the interaction of αCat and βCat is direct, and is presumably not regulated, whereas the interaction
of βCat and Apc2 is mediated by a specific kinase, the Drosophila homolog of the Glycogen Synthase Kinase
3β, Shaggy (sgg) (Rim, Clevers, and Nusse 2022). Additional indication for the activity of the complex came
when we fixed and stained embryos developing at elevated temperature, to visualize nuclei and cortical F-
actin (see methods). We could observe F-actin structures in embryos at elevated temperature (Figure S9, see
panel F), which were reminiscent of those in embryos mutant for *Apc2*, β*Cat*, or *sgg*, as previously described
(McCartney et al. 2001). Thus, we hypothesized that boosting the activity of the complex through over-
expression of one or more of its components should meaningfully rescue the embryo lethality defect, and
the increase in the fraction of mitotic failures in crowded regions and in combined regions.

To test whether the fidelity of cortical mitoses at elevated temperatures was affected by insufficiently tight
or poorly controlled coupling of actin and microtubules, we overexpressed either *αCat* or *sgg*, and again
asked whether the increase in embryo lethality at elevated temperature was rescued. When we over-
expressed either *αCat* or *sgg* we did not see a significant change in embryo lethality at control temperature
(25°C) as compared to the control (Figure 4D). The lethality was greater in outcross controls at elevated
temperature (29°C) similar to results in previous experiments (Figure 1B, Figure 4A), while the embryos over-
expressing either *αCat* or *sgg* faired significantly better at the elevated temperature (Figure 4D). Consistent
with these results, both *αCat* and *sgg* over-expressions rescued the increase in the fraction of mitotic failures,
in the crowded regions, as well as in the combined regions at elevated temperature (Figure 4E, F), with the
large blastoderm holes absent in these embryos. These results suggested that an experimental strengthening
of actin binding was able to attenuate temperature-dependent destabilizations of the mitotic spindle. These

findings are consistent with the idea that the scaffolding activity of αCat, βCat, and Apc2, mediated by sgg, constitutes a thermosensitive controller of nuclear division fidelity in the early blastoderm embryo.

Given the reduction in mitotic fidelity at elevated temperature, we were wondering whether any regulator of mitosis could rescue temperature defects upon overexpression. To test this possibility, we tested other regulators of mitosis and focused on the activity of mud and insc. In epithelia, the activity of 'Mushroom body defect' (mud, NuMA homolog) and that of Inscuteable (insc) is known to dictate the orientation of mitotic spindle during cell division (Mapelli and Gonzalez 2012). Similar activity might be present for nuclear divisions during syncytial blastoderm stage, raising the possibility that over-expression of *mud* or *insc* could rescue embryo lethality, presumably through reducing mitotic failures. However, neither of the over-expressions ameliorated the increased embryo lethality at elevated temperature, with mud over-expression additionally increasing the lethality at normal temperature as well (Figure S11). This failure to rescue embryo lethality further allowed us to argue that, the elevated temperature critically challenges the core machinery of mitotic spindle, more so than the regulation of spindle orientation.

The natural distribution of a few regulators of spindle dynamics varies along thermoclines.
Our analyses showed how elevated temperatures during the first hours of fly development affect early embryonic cell biology; it may seem suggestive, but it was not shown that naturally occurring changes similar to our overexpression of cytoskeletal regulators could shape the evolution of fly or insect development in general. To address this possibility more directly, we analyzed the distribution of standing genetic variations associated with the regulation of the spindle dynamics (combination of GO terms, cell polarity, spindle organization, and spindle localization), and tested the hypothesis that such variation was randomly distributed with respect to described thermoclines (continent-scale gradients of annual mean temperature). We found that indeed, most standing genetic variation associated with putative spindle dynamics regulators was distributed without a clear thermoclinal pattern, although the fraction of genes showing clinal distribution was greater than those associated with a more generic GO term like 'metabolic process' (Figure 4G, Table S1). A clear and notable outlier in this analysis was *sgg*, which was marked with a north/south clinal distribution of natural variations independently in populations in the US, in Europe, and in Australia (see also, Table S2). These findings strongly suggest that the thermosensitivity that *sgg* exhibits in lab conditions, constitutes a biological property that can be exposed and selected for also in natural populations and during the course of evolution. Similarly, we found that 'scribble', a cell polarity protein which interacts with dlg (Khoury and Bilder 2020; 2022), was also clinal in all 3 populations. Further experiments are required to formally test the possibility of a rescue of embryo lethality with over-expression of *scribble*, which will be addressed in future experiments.

Discussion:

Cellular basis of the sensitivity against elevated temperature during early embryonic development
With our work, we demonstrate how the first hours of embryonic development are vulnerable to an exposure to temperatures below the heat-shock range. We identify the mechanistic basis for this vulnerability: the thermolability of the interaction between microtubules and F-actin during mitosis. Given that these early processes till blastoderm formation are conserved across insects (Lynch, El-Sherif, and Brown 2012; Lemke, Kale, and Urbansky 2020), our results implicate a vulnerability in a plethora of insect species. It would be interesting to investigate whether mechanisms exist in nature to accommodate for this vulnerability. In the backdrop of this sensitivity, it is the challenge for the embryo to establish an egg cortex that is uniformly populated with cortically anchored nuclei that carry positional information for subsequent steps of epithelial morphogenesis and cell differentiation. A defect in these early morphogenetic processes and cell differentiation could lead to defects in subsequent processes, ultimately leading to developmental failure

and embryo lethality. Thus, with our data, a mechanism emerges to establish the link between the proximate
effects of elevated temperature on cytoskeletal regulation during mitosis, and its ultimate impact on the
survival of embryos.

What is the best predictor of lethality in embryos exposed to elevated temperatures? In our analysis, we
found three defects in embryos at elevated temperature: mitotic failures, nuclear crowding, and nuclear cycle
asynchrony. Mitotic failures, especially during the last mitosis prior to cellularization, can lead to loss of
nuclei, and consequently a loss of local positional information that is advected and lost with the nuclei. This
can lead to missing cell fates in the coordinate system of the embryo, leading to missing embryonic
structures, which can pose challenges for developmental fidelity. Comparing the fraction of dead embryos at
elevated temperature vs. the severity of mitotic failures, we would argue that the embryos with the most
severe mitotic failures, i.e., those that end up with holes in blastoderm epithelium, are the ones that don't
hatch into larvae. In our over-expression conditions, it is clear that the severe mitotic failures (blastoderm
holes) are absent in the genotypes that rescue the embryo lethality at elevated temperature. It remains to
be seen whether an extensive, but scattered, loss of nuclei poses any developmental disadvantage to the
embryo. It is possible that such disadvantages might be apparent only during larval stage, and might directly
affect larval fitness, but our current analysis wouldn't capture that effect. Additionally, developmental
robustness can vary between species, which raises the possibility that not all fly species will be able to
accommodate such developmental disadvantages. Further, here we have presented 28 embryos of control
genotype at elevated temperature, of which 6 have large- and 5 have small blastoderm holes. In all of these
cases, the blastoderm holes appear in the region where we see an overlap between nuclear crowding and
the delay of nuclear divisions. We further argue that the severe version of the defect at elevated temperature
(blastoderm holes) occur where the three phenomena (mitotic failures, nuclear cycle asynchrony, and
nuclear crowding) spatially overlap, further supported with our quantifications of fraction of mitotic failures
in combined regions. Thus, even though all of the 3 defects can independently reduce embryo fitness, we
demonstrate that the defects cooperate, to induce lethality in a fraction of embryos.

Genetically increasing robustness of early embryos against environmental perturbations
In theory, the robustness of embryonic development can be higher depending on, say, the quality of the food
available to the mother. Thus, one can hypothesize a rescue of embryonic lethality at elevated temperature
through, e.g., improvement of the quality of the diet available to the adult females. A previous study
compared the effect of temperature and the atmospheric O_2 levels on embryo survival, where they report
consistently greater embryo survival at 25% Oxygen concentration, than the normal 21% (Kuntz and Eisen
2015). These results indicate that it might also be possible to increase the robustness of embryo development
through increasing the atmospheric O_2 concentration. However, such solution might not be heritable, and
might not confer an increase in future fitness. Previous studies have increased embryo survival in face of
exposure to elevated temperature through over-expression of heat-shock proteins (Lockwood, Julick, and
Montooth 2017). Presumably, this increases the capacitance of the system to tolerate environmental insults,
akin to how heat-shock proteins are known to act as capacitors for genetic variation (Rutherford and
Lindquist 1998; Sangster, Lindquist, and Queitsch 2004). However, such a rescue would not shed light on
underlying biology of temperature defects, as the effect of such perturbations on the mitotic failures would
remain nebulous. Thus, to rescue the lethality at elevated temperature, we used our cellular level
understanding of the defects to guide our choice of candidate genes for genetic manipulations. This approach
additionally allowed us to challenge our cellular level understanding of the defects at elevated temperature,
as we construct and test hypotheses about the over-expression of various genes towards a genetic rescue.

Various genotypes in our experiments, specifically the overexpression of *αCat*, *sgg*, *baz*, or *dlg*, either rescue
or reduce the severity of one or more defects (embryo lethality, presence of blastoderm holes, fraction of

mitotic failure in crowded region, or that in the combined region) that occur due to an exposure to elevated
temperature. Here, we would argue that *αCat* and *sgg* overexpression are the superior experimental
condition among these as far as potential for a rescue is concerned, as they produce virtually no side-effects
(unlike *dlg* overexpression at normal temperature), while robustly reducing the lethality at elevated
temperature (unlike *baz* overexpression). Additional support for this notion comes from our analysis of the
presence of natural variation along thermoclines, where we identified *sgg* as an interesting candidate gene,
along with *scribble*. The list of genes associated with spindle dynamics (combination of GO terms spindle
organization, spindle localization, and cell polarity) included our candidate genes *baz*, *dlg*, *sgg*, *mud*, and *insc*.
While *αCat* is not presently associated with GO terms related to spindle dynamics, we also did not find it to
be clinal in our extensive literature search. It is possible that natural fly populations don't possess sufficient
standing genetic variation at that locus that could provide a selective advantage. In any case, an overlap
between our candidates and potential natural candidate genes increases our confidence in our approach to
mitigate the detrimental effects of an exposure to elevated temperature during early embryonic
development.

Given that both *αCat* and *sgg* overexpression show a robust rescue of embryo lethality, we would further
argue that the interaction between microtubules and cortical F-actin, as supported by the tripartite protein
complex between αCat, βCat, and Apc2, mediate by sgg, is sensitive to temperature fluctuations. It is likely
that the association/dissociation constants decrease/increase for one or more binding interactions (F-actin
with αCat, αCat with βCat, and βCat with Apc2) as the temperature increases. It is also likely that the kinase
activity of sgg reduces with an increase in temperature, hampering the association between βCat and Apc2.
Further experiments are required to test these ideas. However, here we also note that many nuclei manage
to divide in embryos mutant for βCat, Apc2, and sgg (McCartney et al. 2001), forming a somewhat defective
blastoderm. This would indicate presence of alternative mechanisms that can also be reinforced to increase
the robustness of mitosis, and identifying and testing such mechanisms would be of great interest to rescue
embryo lethality at elevated temperature.

How fast can nature increase the robustness against the adverse effects of elevated temperature?
In a recent study (Rudman et al. 2022), the researchers allowed fly populations to naturally evolve from spring
through summer till the fall season, and found that specific single nucleotide polymorphisms were selected
for in the regulatory regions of *sgg* and β*Cat*, along with many other genes. This further indicates that nature
might be exploiting the allelic plasticity in our candidate genes at rather rapid timescales of just a few
generations, to increase robustness against elevated temperatures. It is interesting to see that these genes
show selectable variation at the population level, and that an over-expression of *sgg* in our analysis is capable
of rescuing the embryo lethality at elevated temperature. To be clear, the study does not comment on
whether the accumulated mutations lead to an over-expression or hyperactivation of *sgg*, and at which stage
during the life cycle. At the same time, the analysis won't pick up candidates that don't have sufficient
variation in the seed population. In any case, our functional analysis can be effectively used to test for the
function of any gene that rescues (or worsens) the effect of elevated temperature on early embryonic
development, in any fly embryo that's amenable to experimental manipulations. We believe that our
experimental pipeline holds the promise to uncover exciting biology, while understanding how physiological
processes are affected by the exposure to elevated temperature.

Sensitivity of early embryonic development to the global warming and climate change
Climate change is a global problem that threatens the existence of life on earth as we know it (IPBES 2019).
In this context, we decided to focus on effect of elevated temperature on the physiology of flies, given their
vital roles for the functioning of ecosystems (Raguso 2020). We found the early embryonic development in
Drosophila to be especially sensitive to an exposure to elevated temperature. Here, we identified the

syncytial blastoderm stage during embryonic development as the earliest possible time window that's
sensitive to elevated temperature. We could also observe cascading defects in the cellular blastoderm
formed thereafter. An increase in embryonic lethality, due to the early exposure to elevated temperatures,
further indicated that, presumably, the subsequent developmental processes are affected as well. We also
see a small increase in embryonic lethality when we expose post-cellularization embryos to elevated
temperature, indicating additionally that the temperature sensitive time window might actually be longer
than just the pre-gastrulation developmental phase. It is easy to imagine that, in wild, embryos would be
exposed to elevated temperature for first few hours, starting from egg lay till well beyond gastrulation. Given
that these early processes during the embryonic development are conserved across fly species (Lemke, Kale,
and Urbansky 2020), our results raise the possibility that many other key species, would exhibit similar
defects during early embryonic development, will have an increase in lethality, and will be overall threatened
due to global warming. It is rather impressive that an exposure to a sub-heat-shock elevated temperature
during first few hours of embryonic development is capable of reducing the fitness of the entire embryonic
development to increase embryonic lethality: an effect which is presumably conserved across the animal
kingdom, and would be worth exploring at the cellular level to improve embryo survival. Perhaps, the
temperature sensitive developmental window is so crucial as it includes gastrulation, very much in agreement
with the famous quote from Lewis Wolpert, "It is not birth, marriage, or death, but gastrulation which is truly
the most important time in your life".

Materials and methods:

<u>Fly lines and genetics</u>:

OregonR line was used as WT *Drosophila*.

Histone-RFP expresses RFP tagged version of His2Av. Insertion on 3rd chromosome. Bloomington Drosophila Stock Center (BDSC) #23650.

Jupiter-GFP expresses GFP tagged version of microtubule binding protein Jupiter. Insertion on 3rd chromosome. Always used in combination with *Histone-RFP*. Gift from Jan Felix Evers.

67-Gal4 (mat αTub-GAL4-VP16) is a maternal specific Gal4 driver. Insertion on 2nd chromosome. Always used in combination with *Histone-RFP* and *Jupiter-GFP* in the over-expression experiments (see below). Gift from Thomas Lecuit.

The following lines were used to drive the over-expression of various genes of interest in an attempt to rescue the effects of elevated temperature. *baz*, BDSC #17613; *dlg*, BDSC #15779; *αCat*, BDSC #65590; *sgg*, BDSC #15886; *mud*, BDSC #22371; *insc*, BDSC #39676. An outcross to *w1118* was used as control.

For various over-expression experiments, we first crossed virgins from the Gal4 line '+; 67-Gal4; Jupiter-GFP, Histone-RFP' to the males of over-expression line 'UAS-X; +; +' (X= *baz*, *dlg*, *sgg*, or *mud*) or '+; +; UAS-Z' (Z= *αCat*, or *insc*), or to males of *w1118* as control. F1 virgin females from these crosses were back crossed to males of the same paternal genotype.

Genotypes associated with various figure panels are listed in Table S3.

<u>Embryo collection</u>: For any experiment, we put males and females of appropriate genotypes (Table S3) into transparent cylindrical embryo collection cages (inner diameter ~5 cm). Fine wire mesh is glued to top side of the cage to allow air exchange while keeping the flies trapped, and the bottom of the cage is closed with apple-juice agar plates bearing a thin layer of yeast paste, where flies deposited the embryos. To prepare the agar plates, we typically prepare 1 l of apple-juice agar solution. For this, we boiled 24 gm of agar in 750 ml of water, to which we added 250 ml of fresh store-bought apple juice, while stirring on a hot plate. We make the final amount of sucrose to 25 gm: the amount of additional sucrose depends on the sugar content of the apple juice. This solution is sufficient for ~80 petri-dishes of 6 cm diameter. After agar is solidified, the plates can be stored for long time at 4°C. Before the embryo collection, the plates were covered with a thin layer of yeast paste (to stimulate oviposition), and were pre-incubated to the embryo collection temperature depending on the temperature regimen, for at least half an hour. *Drosophila* females are known to hold onto the embryo long after fertilization, such that a large part of the early embryo development can occur inside the abdomen. To avoid these embryos, we perform a sham embryo collection for about an hour at room temperature (24-25°C), from which all embryos are discarded. Following this sham collection, we collect the embryos that are actually used in the experiment at the specific experimental temperature.

<u>Embryo lethality experiment</u>: There were 3 temperature regimes in the embryo lethality experiments, as depicted in Figure 1A. Depending on the regimen, we collected embryos for 1 hour either at normal temperature (24-25°C) or elevated temperature (29-30°C). The 24-25°C collections (corresponding to normal temperature) were done on the bench top. The 29-30°C collections (corresponding to elevated temperature) were done in a water bath, where the bottom surface of the embryo collection agar plate is in thermal contact with the heated water. For NN or EN, we collected embryos for 1 hour either at normal temperature or elevated temperature, and then incubated them at the same temperature for another 2 hours. Afterwards, the embryos are washed with bleach for 10-15 seconds, then washed thoroughly with tap water, aligned on a 24x60 mm rectangular coverslip, and covered with halocarbon oil. For NE, we collected embryos for 1 hour

at normal temperature, and then incubated them at the same temperature for another 1.5 hours. Within the
next half an hour, the embryos are washed with bleach for 10-15 seconds, then washed thoroughly with tap
water, aligned on a 24x60 mm rectangular coverslip, and covered with halocarbon oil. Then, we incubated
them at elevated temperature for 3 hours. Then, for all the temperature regimen, the coverslips with
embryos were incubated at normal temperature for about a day, following which we counted the number of
eclosed larvae, and the number of dead embryos. The embryonic lethality was estimated as,

546
$$embryo\ lethality\ = \frac{100 * number\ of\ dead\ embryos}{number\ of\ dead\ embryos\ +\ number\ of\ larvae}$$

We excluded the unfertilized embryos from these calculations.

<u>Imaging fixed embryos</u>: We used 2 different temperature regimens for embryo collection for fixation and
staining. We collected embryos for 3-4 hours either at normal temperature (24-25°C) or elevated
temperature (29-30°C). The 24-25°C collections (corresponding to 25°C regimen) were done on the bench
top. The 29-30°C collections (corresponding to 29°C regimen) were done in a water bath, where the bottom
surface of the embryo collection agar plate is in thermal contact with the heated water. Afterwards, the
embryos are washed with bleach for 40-45 seconds, washed thoroughly with tap water, and fixed
immediately. Collecting embryos for longer duration is not a problem here, as we only image and analyze the
embryos that are before gastrulation stage, through *post hoc* identification of developmental stage. Here,
we did 2 kinds of experiments; one, where the aim was visualizing the nuclei from the entire embryo (data
in Figure S2); and two, where the aim was visualizing the nuclei along with F-actin distributions from half
embryos (data in Figure S9). Accordingly, we used 2 different protocols for embryo fixation, staining,
mounting, and imaging.

To **visualize all nuclei from the entire embryo**, we used heat fixation, and stained the embryos with DRAQ5.
Briefly, we transferred the processed embryos to 50 ml tubes with a brush, and drained the excess water. To
this tube, we added the boiled-to-turbidity heat-fix solution (28% 500 µl NaCl + 5% 200 µl TritonX100 + 19.3
ml water), incubated it for ~20 sec with thorough mixing, and then immediately added water to cool down
the mixture. Then, we transferred the fixed embryos to 2 ml microfuge tubes, sequentially added n-Heptane
and methanol in equal proportions, vigorously shook the tube for about half a minute to break-open and
remove the vitelline membrane. This was followed by 3x washes of methanol. Here, we started the staining
immediately after, although the embryos can be stored in methanol for long time. The embryos were
rehydrated with PBT (0.01% solution of Tween20 in PBS) over 3 washes. Following this we stained the
embryos with 1:1000 DRAQ5 for 1 hour. Afterwards, we washed the embryos 6x with PBT, re-suspended
them in 3:1 glycerol:PBS solution, and mounted them between 2 rectangular coverslips. The spacing between
the coverslips was optimized to keep the embryos in place w/o deforming them. Then we imaged the
embryos through both coverslips, and stitched the two half embryos during image analysis to get the full
view (see below). Images were acquired on Leica SP8 DMi8 inverted confocal microscope using 20x glycerol
objective (0.75NA), with 1.15x digital zoom, at 1 µm step-size in z, and a final x-y resolution of 0.247 µm/pixel.

To **visualize the nuclei and F-actin**, we used formaldehyde fixation, and stained the embryos with DAPI and
Phallacidin. Briefly, we transferred the processed embryos to 20 ml vial with a brush, which contained the
fixation mix (3500 µl PBS + 37% 1000 µl formaldehyde + 4500 µl n-Heptane). The vial was shaken on a
platform shaker for 25 minutes @200 rpm. Then, we transferred the fixed embryos to 2 ml microfuge tubes,
sequentially added n-Heptane and 90% ethanol in equal proportions, vigorously shook the tube for about
half a minute to break-open and remove the vitelline membrane. This was followed by 3x washes of 90%
ethanol. Here, we started the staining immediately after, although the embryos can be stored in 90% ethanol
for a week or so. The embryos were rehydrated with PBT (0.01% solution of Tween20 in PBS) over 3 washes,

following this we stained the embryos with 1:1000 DAPI for 1 hour, followed by 1:500 Phallacidin for 1 hour.
Afterwards, we washed the embryos 4x with PBT, re-suspended them in 3:1 glycerol:PBS solution, and
mounted them between 2 rectangular coverslips, followed by imaging. Images were acquired on Leica SP8
DMi8 inverted confocal microscope using 20x glycerol objective (0.75NA), with 1.2x digital zoom, at 1 μm
step-size in z, and a final x-y resolution of 0.473 μm/pixel.

Imaging live embryos: We used 4 experimental temperature regimes for embryo live imaging. We collected
embryos for 1 hour either at normal temperatures (22-23°C, 24-25°C, or 27-28°C) or elevated temperature
(29-30°C). The 22-23°C collections (corresponding to 22°C imaging) were done in the fly incubator. The 24-
25°C collections (corresponding to 25°C imaging) were done on the bench top. The 27-28°C collections
(corresponding to 28°C imaging) or the 29-30°C collections (corresponding to 29°C imaging) were done in a
water bath, where the bottom surface of the embryo collection agar plate is in thermal contact with the
heated water. Afterwards, the embryos are washed with bleach for 35-40 seconds, then washed thoroughly
with tap water at the same temperature as embryo collection. We then aligned the embryos on a 25 mm
diameter circular coverslip, covered them with halocarbon oil, and immediately transferred them to the
temperature controller setup on the microscope that's pre-equilibrated to the experimental temperatures
for normal temperature regime, (22, 25, or 28°C) or the elevated temperature regime (29°C). We used the
temperature controller setup from Warner instruments (locally supplied by MultiChannelSystems). The setup
consists of; QR-40LP, which holds the 25mm coverslip with embryos; QE-1HC, which heats/cools QR-40LP
along with the embryos mounted in it; LCS-1, the liquid cooling system that isolates the QE-1HC from the
effect of ambient temperature; and CL-100, the bipolar temperature controller that can impart the
temperature of interest against the baseline set by LCS-1. We mounted QE-1HC on the microscope stage,
with LCS-1 and CL-100 as the peripherals controlling the temperature. Images were acquired on Leica SP5
DMI6000CS inverted confocal microscope using 40x water objective (1.1NA), with 1x digital zoom, at 2 μm
step-size in z with 20 z-steps overall, and a final x-y resolution of 0.379 μm/pixel. As mentioned in the main
text, we acquired images starting from interphase 10 or before, till the beginning of gastrulation movements.
We recorded the timelaspe at a frame rate of 1 frame every 30 sec, 1 min, or 2 min depending on whether
we recorded at a time 1, 2, or 3-4 embryos in parallel.

Image analysis and statistics: All image analysis was done using FIJI freeware. There are 7 different kinds of
image analyses presented in the manuscript. These are; one, whole embryo reconstruction to see distribution
of nuclei in the entire embryo (fixed embryos); two, the analysis of mitotic failures (live embryos); three, the
analysis of nuclear movements (live embryos); four, analysis of nuclear cycle asynchrony (live embryos); five,
analysis of nuclear crowding (live embryos); six, analysis of the fraction of mitotic failures (live embryos); and
seven, comparing the distribution of nuclei alongside that of F-actin (fixed embryos). All the statistical analysis
was done using Prism (GraphPad). This includes, the curve fitting (see below, the analysis of nuclear
crowding), estimating the non-parametric statistical significances, and estimating the 95% confidence
intervals (CI) on the medians. The straight-line fits for analysis of nuclear movements were done using ImageJ
macro language.

In the **analysis to reconstruct the nuclear distribution in the whole embryo**, we used a custom-built image
analysis pipeline to fuse the halves of the same embryo, followed by the computing the distance of each
point inside the embryo from the embryo surface. Here we refer to the two views of the embryo as "half
embryo", even though in reality the volume in either "half" spans about 70-75% of the embryo width,
providing a significant overlap between the two halves. In the pipeline, we process the raw images, stitch the
two half embryos, render the fused embryo in lateral view, segment the embryo volume, and then segregate
the nuclei as a function of their distance from the surface of segmented embryo volume. First, we processed
the raw images: to improve visualization of the nuclei, we used 'Minimum 3D' filter with radii x=y=1, and z=3,

and to reduce noise, we used 'Median 3D' filter with radii x=y=z=1. Then, we homogenized the
brightness/contrast between the two half embryos. Following these steps, we manually aligned the half
embryos along their AP axis, and then used the BigWarp (Bogovic et al. 2016) package in the BigDataViewer
ecosystem (Pietzsch et al. 2015) to align the half embryos using manually selected interest points. Essentially,
we selected either subcortical- or yolk-nuclei that could be identified in both half embryos. We further
cropped the output images to keep only a small margin around the embryo with "aligned halves", in order
to reduce the computational load in the steps to follow. After the manual alignment step, we used BigStitcher
(Preibisch, Saalfeld, and Tomancak 2009) to further align embryo, using phase correlation method, to
produce the "fused embryo". In case of many embryos, additional manual transforms were necessary to
produce a fused embryo that passes visual inspection. The fused embryos were then rotated as necessary,
and resliced to produce lateral views. Following this, we processed the fused embryo to separate the regions
of embryo that contain cortical-, subcortical, and yolk-nuclei. First, we used the nuclear signal to segment the
embryo volume, using simple thresholding. This generated a binary image that corresponds to the embryo
volume. Then using '3D Distance Map' (plugin in the 3DSuite), we compute the Euclidian distances in 3D, i.e.,
each pixel inside the embryo gets the grayscale value corresponding to its distance from the embryo surface.
All pixels with Euclidian distance <10 μm would be considered as the cortical layer, those within range of 10-
30 μm would be consider subcortical, and those farther than 30μm from embryo surface would be considered
as part of the yolk region. These distances are determined empirically, using approximate measurements
from various fused embryos. Typically, nuclei have the diameter of 5-10 μm depending on the phase of
nuclear division cycle, while the cell height at the end of cellularization is about 30 μm. The pixels in these
regions were then color coded and presented as either as maximum projections or cross-sectional views in
Figure S2.

In the **analysis of mitotic failures**, we tracked expelled nuclei back in time to identify the mitosis when they
were born, and to categorize the expulsion as pairwise or isolated (see examples respectively in Figure 1F
and Figure S3D). We considered a nucleus to be expelled, if its apical-most tip was at least 12 μm away from
the embryo surface. To simplify the identification of expelled nuclei, we color-coded the Histone-RFP signal
as a function of distance from the embryo surface. This is similar to above analyses, except we use the
'LocalZPojector' plugin in FIJI (Herbert et al. 2021) to define the distance from the embryo surface, while to
predict the embryo surface itself we use the Jupiter-GFP signal, which was co-imaged with Histone-RFP. Thus,
first we color-code the nuclei, and then manually identify and count the number of pairwise or isolated
nuclear expulsions during interphase 14.

In the **analysis of nuclear movements**, for a select few nuclei we manually tracked the entire progeny,
starting from the mother nucleus in late interphase 10 till the 8 great-granddaughter nuclei at late interphase
13. Assigned all of these nuclei to a single cluster, we then calculated the center of the cluster at each time
point. We chose nuclei along one continuous row at interphase 10, and tracked their progeny in their
individual clusters. For sake of simplicity and consistency, we only chose those nuclear clusters whose entire
progeny could be tracked, i.e., we could follow the cluster and see the formation of all 8 great-granddaughter
nuclei in interphase 13. This way, we obtained ≥ 13 clusters in any embryo that we tracked. For each of these
nuclear cluster trajectories, we fitted a straight-line to coordinates for the center of the cluster, over time
(dashed lines in Figure S5B). Then, we calculated the distance of the center of the nuclear cluster to this line
at every timepoint. An average of these distances over a specific time period was used as an estimate for
deviations in the center of nuclear clusters over that time period. We observed that the clusters move
significantly starting from anaphase till a few minutes into the subsequent interphase, as the clusters return
to their normal position. Thus, we decided to separately analyze these periods of extensive movement vs.
the periods when the cluster movement was low. To identify the periods of extensive movements

consistently and objectively, we noted that these movements start at least 1 timepoint (30 sec) before we
could see the first telophase in any of the tracked clusters, while the movements mostly subside within 3
minutes after all of the nuclear clusters finished telophase. Thus, we defined the window of extensive
movement as starting from 30 sec before appearance of first telophase nuclei, till 3 min after the appearance
of last telophase nuclei. Once we had an objective definition of various time periods, it was straightforward
to calculate the average deviation in the center of nuclear clusters.

In the **analysis of asynchrony in nuclear cycles**, we marked various timepoints in the timelapse recordings to
follow the progression of nuclear cycles in the anterior-, middle-, and posterior-5th of the field of view. For
sake of simplicity, we marked the transitions from a telophase 13 to interphase 14, i.e., the first timepoint
when the daughter nuclei in any region go from telophase 13 to interphase 14. If the timepoint for this
occurance in the anterior is 'T$_a$', that in posterior is 'T$_p$', while the same in the middle is 'T$_m$', then the
'asynchrony' can be calculated as,

685
$$asynchrony = T_m - \left(\frac{T_a + T_p}{2}\right)$$

Essentially, we started with the expectation that T$_m$ should be the average of T$_a$ and T$_p$ (maximum synchrony),
and a deviation from this expected value would be the asynchrony. Figure S7 diagramatically demonstrates
this analysis.

In the **analysis of nuclear crowding**, we analyzed the distribution of nuclei at the end of interphases, as that's
the time when the distribution is as stable as it can be, with maximum crowding, if any. To assess nuclear
crowding, we started with segmentation of nuclei in the field of view. Here, we used the 'Find Maxima'
command to identify local intensity maxima, and imprinted those on an image. Many of these maxima
marked a nucleus. We did manual corrections to, remove the maxima that didn't correspond to a nucleus,
add a maximum for nuclei that didn't have a maximum associated with them, or correct the maxima that
were too off-center from the corresponding nucleus. Then we used Voronoi tessellation to mark territories
around these maxima to generate pseudo-cells. We excluded territories that, touched the edge of the image,
that had aberrant shapes (aspect ratio>2), or those with aberrant pixel intensities (pseudo cell containing
pixels with <5 units of intensity, typically the case with ill-defined pseudo-cells at the edge of segmentation).
Thus, each nucleus was associated with a pseudo-cell area value ('Area' in μm^2), and an x-coordinate value
for the center of the pseudo-cell ('X' in μm) (x-axis of the image aligned with anterior-posterior axis of the
embryo). We fitted these with the equation of a parabola such that 'Area' can be expressed as a function of
'X',

703
$$Area = aX^2 + bX + c$$

Here, the curvature of the parabola will be higher in case of greater nuclear crowding (compare Figure 2L-O).
Thus, the fit parameter 'a', which dictates the curvature of the parabola, was used to estimate the extent of
crowding.

In the **analysis of the fraction of mitotic failures**, we tracked expelled nuclei back in time same as the analysis
for mitotic failures, except now we also recorded where the nuclei were lost from, to get a map of the
locations of mitotic failure in the field-of-view. In parallel, we generated regions in the embryo that
correspond to maximum nuclear crowding, maximum delay in nuclear divisions, and a combination of the
two regions. To construct the *crowded regions* (the region corresponding to maximum nuclear crowding) we
referred to previous studies, which have argued that a certain threshold of nuclear content will halt the
syncytial nuclear division cycles (Edgar, Kiehle, and Schubiger 1986), while this threshold was determined to
be ~70% of the normal cellularization nuclear density (Lu et al. 2009). Here, we argue that such threshold

nuclear content is reached in regions with nuclear crowding, albeit only locally, and should locally trigger an
arrest of nuclear divisions, in conflict with rest of the embryo. Another previous study reported the nuclear
domain area (i.e., the pseudo-cell area) to be 28.8 μm^2 at cellularization (Telley et al. 2012). We corroborate
this number with our own quantifications (data not shown). For simplicity of calculations, we assume that
Drosophila embryos have 6000 nuclei at interphase 14 each with an area of 29 μm^2. The 70% density of nuclei
would thus correspond to 4200 nuclei at interphase 14, each with an area of 41.43 μm^2, which we
approximate to 42 μm^2. Thus, the back of the envelop calculation indicated that nuclei will seize to divide
when their pseudo-cell area drops below ~42 μm^2. We observed that in embryos developing at 29°C, the
central regions of the embryo contained many nuclei that had pseudo-cell areas less than 42 μm^2, already at
interphase 13. Thus, we used this criterion to define areas of embryos that have likely reached the 70%
cellularization nuclear density, at one nuclear division cycle earlier, albeit only locally. Since the embryos at
normal temperatures (22, 25, and 28°C) rarely have nuclei with pseudo-cell areas smaller than 42 μm^2, we
further decided to include all nuclei with pseudo-cell areas in the lower quartile to construct the region with
maximum nuclear crowding. Here we also note that this description is only a rationalization of the approach,
and that it is not a comment on how the effect of nuclear crowding might actually manifest. For instance, it
is also likely that mitotic spindles are not properly established in the crowded regions, thus causing the
mitotic failure. However, this wouldn't allow us to define equivalent regions in the embryos at normal
temperatures, as it is not feasible to accurately define the size of the mitotic spindle, in our experiments.
Then, to construct the *delayed region* (the region with maximum delay in nuclear divisions), we simply
marked the areas of the embryo that contain the last ~20% of the nuclei that finished nuclear cycle 13. This
percentage was decided empirically: it corresponds to a region of similar size as that corresponding to the
crowded regions above. Here the rationale is drawn from previous studies which indicate that the regulation
of nuclear division cycles is coordinated at the scale of the entire embryo (Deneke et al. 2016), and given that
all of the nuclei share the same cytoplasm, a local delay in nuclear cycles would create conflicts between the
biochemical status of the cytoplasm and the cell cycle stage of the delayed nuclei. Finally, to construct the
combined region (combination of crowded and delayed region), we used the logical OR operator to combine
the crowded and delayed regions, to test combinatorial effects of the nuclear crowding and nuclear cycle
asynchrony. In the end, we computed the fraction of mitotic failures in the crowded, delayed, or combined
regions using,

744
$$fraction\ of\ mitotic\ failures = \frac{number\ of\ mitotic\ failures\ in\ a\ specific\ region}{total\ number\ of\ mitotic\ failures}$$

In the **analysis to compare the distribution of nuclei alongside that of F-actin**, was rather similar to the
pipeline to *reconstruct the nuclear distribution in the whole embryo*, except that we didn't have 2 half-
embryos to fuse, essentially starting from segmentation of embryo volume. Here, instead of using nuclear
staining, we segmented embryo volume using the F-actin staining. Then the 3D distance map similarly
allowed us to classify pixels into cortical-, subcortical-, or yolk-regions based on their distance from embryo
surface. The pixels in these regions were then color coded and presented as sum projections in Figure S9.

Figure and table legends:

Figure 1: Embryonic lethality and the loss of nuclei due to mitotic failures in embryos exposed to elevated
temperature.
(A, B) To assess developmental robustness, we exposed embryos to various temperature regimen and
estimated the embryo lethality. (A) Schematic showing the temperature regimen in the embryo survival
experiment, starting from the beginning of embryo collection. Here we have three conditions depending on
exposure to normal or elevated temperature regimen in the 3-hour time windows as shown, normal-normal

[NN], elevated-normal [EN], and normal-elevated [NE]. In all cases, the first time-window also includes 1 hour of embryo collection. The embryonic development beyond 6 hours continued at normal temperature. (B) The scatter plots show the distribution of percent lethality in embryos from various conditions, calculated based on the fraction of embryos that don't hatch into larvae. Each dot represents data from a sample with ≥26 embryos; n= 12 (NN), 10 (EN), 10 (NE) samples.

(C-E) Montages of snapshots from live embryos, expressing Histone-RFP, from late interphase 11 till late interphase 14, in embryos developing either at normal temperatures (C), or at elevated temperature showing mild (D), or severe (E) defect. The nuclei were labelled as a function of their distance from the embryo surface. Here, green indicates cortical, and magenta indicates sub-cortical nuclei that are lost. At elevated temperature, a lot more nuclei are expelled, and as a result, we can see large gaps between cortical (green) nuclei, often coinciding with expelled (magenta) nuclei, especially during early interphase-14.

(F-H) To assess the pairwise loss of nuclei, we counted mitotic failures that lead to such nuclear expulsions. (F) Montage of snapshots from live embryo, from late interphase 12 to late interphase 13, where Histone-RFP marks the distribution of nuclei, cortical nuclei are in green and subcortical (expelled) nuclei are in magenta. The pairwise loss of nuclei due to mitotic failure is exemplified in an embryo developing at elevated temperature. The arrowheads track the segregating daughter nuclei that will be expelled eventually, while arrows exemplify normal nuclei. Of note, this type of nuclear loss occurs at all temperatures, though at different a frequency. (G) Scatter plot showing the distribution of events of nuclear losses due to mitotic failure during interphase 14. n= 10 (22°C), 7 (25°C), 9 (28°C), and 10 (29°C) embryos. (H) The schematic to illustrate the phenomenon of mitotic failures at elevated temperature.

Scale bars; C-E, 25 μm; F, 5 μm. B and G show scatters with median, with whiskers showing 95% CI. In G, the hollow markers indicate embryos that have a blastoderm hole, with large hollow markers indicating a single large hole, and small hollow markers indicating either a single or multiple small holes. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$, non-significant comparisons are not shown.

Figure 2: Nuclear cycle asynchrony and nuclear crowding increases as a function of temperature during embryo development.

(A-C) To assess the asynchrony in nuclear cycles, we identified the phases of nuclear cycle along the AP axis. (A) Image shows the entire anterior-posterior extent of the field of view, from a live embryo developing at 29°C. Histone-RFP marks the nuclei, cortical nuclei are in green and subcortical (expelled) nuclei are in magenta. Different regions of the embryo have nuclei in different stages of nuclear division cycle, as indicated above the image. Arrows show the direction in which the wave of nuclear division cycle will progress. Here, we have an anterior- and a posterior-wave which will meet in the central region of the embryo. The anterior, central, and posterior $1/5^{th}$ of the field of view are indicated below the image, with the anterior $1/5^{th}$ of the embryo containing nuclei that are more advanced in division cycle, as compared to the posterior $1/5^{th}$, which are yet more advanced than the central $1/5^{th}$. As a result, these regions will enter the subsequent interphase in an asynchronous manner (see also Figure S7). (B) Scatter plots showing the distribution of asynchrony for embryos at various temperatures. n= 10 (22°C), 7 (25°C), 9 (28°C), and 10 (29°C) embryos. (C) The schematic illustrates the progressive increase in asynchrony at elevated temperature.

(D-Q) To assess the crowding of nuclei, we segmented the nuclei and generated pseudo-cells. The figure shows data from of live embryos that were developing either at 22°C (D, H, L), 25°C (E, I, M), 28°C (F, J, N), or at 29°C (G, K, O). (D-G) Snapshots from live embryos at the end of interphase 13, where Histone-RFP marks the nuclei. The outline shows the region of the image that is considered for quantifications. (H-K) The pseudo-cells for the corresponding embryos in D-G. A pseudo cell is centred on a nucleus, while the region of a pseudo-cell is bound by line segments that are equidistant from neighbouring nuclei. Various pseudo-cells

are colour-coded according to their area in μm^2 according to the LUT at the bottom. (L-O) Plots showing the distribution of pseudo-cell areas as a function of their anterior-posterior position, along with a quadratic fit through the distribution, for embryos in D-G. (P) Scatter plots showing distribution of the coefficient of the quadratic term for embryos at various temperatures. This essentially captures the extent of nuclear crowding, with larger quadratic coefficient indicating greater crowding. n= 10 (22°C), 7 (25°C), 9 (28°C), and 10 (29°C) embryos. (Q) The schematic illustrates the crowding effect at elevated temperature. The size of the black arrows indicates the gradation in nuclear density, from poles to the central region.

Scale bars, 20 μm. B and P shows scatters with median, with whiskers showing 95% CI. In B and P, hollow markers indicate data from embryos that have blastoderm hole, with large hollow markers indicating a single large hole, and small hollow markers indicating either a single or multiple small holes. *, p<0.05; **, p<0.01; ***, p<0.001; non-significant comparisons are not shown.

Figure 3: Nuclear crowding and nuclear cycle asynchrony combinatorially enhance mitotic failure defect.
(A, B) Snapshots from live embryo, developing at 29°C, at the end of interphase 13 (A), and 14 min afterwards (B), where Histone-RFP marks the nuclei. The outline in A shows the region of the image that is has nuclear crowding, while that in B bounds the region containing most delayed nuclei, i.e., those that are most lagging in nuclear division cycles.

(C-E) The locations of lost pairs of nuclei during interphase 14 are marked as white dots to indicate their distribution in the field of view. Then, various regions are defined in the embryo to illustrate the fraction of mitotic failures occurring in that region. These regions are composed based on, the crowded region (C), the delayed region (D), and the combined region (E).

(F-K) Scatter plots showing the distribution of the area fractions in crowded (F), delayed (G), and combined (H) regions of the embryos. In case of both crowded (F) and delayed (G) regions, the medians are rather comparable across temperature, and the low variability within a given temperature is exaggerating the differences across temperatures. In case of the area fraction of the combined regions (H), the increase in the overlap of crowded and delayed region reduces the overall size of the combined regions. Then, scatter plots showing the distribution of fraction of mitotic failures occurring in the crowded region (I), in the delayed region (J), and in the combined regions (K). n= 10 (22°C), 7 (25°C), 9 (28°C), and 10 (29°C) embryos.

Scale bars, 20 μm. F-K show scatters with median, with whiskers showing 95% CI. Hollow markers indicate data from embryos that have blastoderm hole, with large hollow markers indicating a single large hole, and small hollow markers indicating either a single or multiple small holes. *, p<0.05; **, p<0.01; ***, p<0.001; non-significant comparisons are not shown.

Figure 4: Rescue of embryonic lethality at elevated temperature through over-expression of the regulators of cell polarity signalling or those of F-actin -microtubules interaction during mitosis, and genes varying along thermoclines as "natural" candidates.
(A-C) Testing the rescue of embryonic lethality and fraction of mitotic failures with over-expression of *baz* and *dlg*. (A) Scatter plot showing the distribution of percent lethality in embryos from various genotypes, at normal (25°C) vs elevated temperature (29°C), calculated based on the fraction of embryos that don't hatch into larvae. Each dot represents data from a sample with ≥12 embryos; control, n= 9 (22°C), 9 (29°C) samples; baz-OE, n= 11 (22°C), 10 (29°C) samples; dlg-OE, n= 12 (22°C), 10 (29°C) samples. Scatter plots showing the distribution of the fraction of mitotic failures in crowded regions (B), and in crowded OR delayed regions (C), in the embryos at normal (22°C) vs elevated temperature (29°C). Each dot represents data from one embryo: control, n= 7 (22°C), 9 (29°C) embryos; baz-OE, n= 7 (22°C), 7 (29°C) embryos; dlg-OE, n= 6 (22°C), 7 (29°C) embryos. Additional quantifications related to this experiment are presented in Figure S8.

(D-F) Testing the rescue of embryonic lethality and fraction of mitotic failures with over-expression of *αCat*
and *sgg*. (D) Scatter plots showing the distribution of percent lethality in embryos from various genotypes,
at normal (25°C) vs elevated temperature (29°C), calculated based on the fraction of embryos that don't
hatch into larvae. Each dot represents data from a sample with ≥18 embryos; control, n= 10 (22°C), 11 (29°C)
samples; αCat-OE, n= 10 (22°C), 12 (29°C) samples; sgg-OE, n= 8 (22°C), 7 (29°C) samples. (E, F) Scatter plots
showing the distribution of the fraction of mitotic failures in crowded regions (E), and in crowded OR delayed
regions (F), in the embryos at normal (22°C) vs elevated temperature (29°C). Each dot represents data from
one embryo: control, n= 8 (22°C), 9 (29°C) embryos; αCat-OE, n= 8 (22°C), 8 (29°C) embryos; sgg-OE, n= 7
(22°C), 7 (29°C) embryos. Additional quantifications related to this experiment are presented in Figure S10.

(G) Pie charts showing the clinal distribution score for various genes associated with 'spindle dynamics'
(combination of GO terms, cell polarity, spindle organisation, and spindle localisation, n=320 genes), or with
GO term 'metabolic process' (n=6612 genes). ~40% of genes associated with spindle dynamics, while ~30%
of the genes associated with metabolic process, are clinal in at least one continent.

A-F shows scatters with median, with whiskers showing 95% CI. In B, C, E, and F, hollow markers indicate data
from embryos that have blastoderm hole, with large hollow markers indicating a single large hole, and small
hollow markers indicating either a single or multiple small holes. *, p<0.05; **, p<0.01; ***, p<0.001; ****,
p<0.0001; non-significant comparisons are not shown.

Figure S1: Blastoderm holes appear in a fraction of embryos at elevated temperature.
The figure shows the snapshots from live embryos, expressing Histone-RFP, from early interphase 14, in
embryos developed at elevated temperature. The nuclei are labelled as a function of their distance from the
embryo surface. Here, green indicates cortical, and magenta indicates sub-cortical nuclei that are expelled,
while arrows/arrowheads indicate large/small blastoderm holes. For embryos with large blastoderm holes
(embryo 05, and 08), we see many nuclei in magenta, while in case of embryos with small blastoderm holes
(embryos 04, and 09), we can see only a few nuclei in magenta per hole. We would argue that development
in *Drosophila* is robust enough to buffer the defects due to the small holes, but fails at 29°C when large holes
appear.

Scale bar, 25 μm.

Figure S2: Nuclear loss shows left-right asymmetry in embryos developing at elevated temperature.
Distribution of nuclei in embryos during interphase 13, in fixed embryos that were developing either at
normal temperature (A, D, G, H), or at elevated temperature that show either a mild defect (B, E, I, J) or a
severe defect (C, F, K, L). (A-F) Maximum projections of the right and left halves of embryos, where the nuclei
are labelled as a function of their distance from the embryo surface. Here, green indicates cortical nuclei, i.e.,
nuclei close to embryo surface, while magenta indicates sub-cortical nuclei, i.e., nuclei that are deeper in the
embryo, farther from embryo surface. The yolk nuclei are not shown for improved visibility. As compared to
embryos at normal temperature (A), those at elevated temperature show mildly (B) or severely (C) numerous
gaps in the distribution of cortical nuclei. These gaps often also correspond to the locations of subcortical
nuclei (A-C). Of note, the embryo surface prediction gets distorted at the posterior pole, due to the pole cell
nuclei that are outside the embryo cortex, thus mislabelling cortical nuclei in magenta. These mis-coloured
nuclei were manually erased to exclusively highlight the subcortical nuclei (D-F). (G-L) Cross-sectional view
along the indicated regions in panels A-C. Here, yolk nuclei are additionally marked in yellow. The arrow
indicates the direction along which we can follow the surface of the embryo along the cross-section. The
cross-sections are chosen to specifically highlight the locations of subcortical nuclei.

Scale bars; A-C, 50 μm; G, I, and K, 25 μm.

Figure S3: Out-of-plane nuclear divisions during mitosis 10, and the low frequency of isolated nuclear expulsion during interphase 14, does not vary significantly with temperature, and cannot explain the observed loss of nuclei.

(A, B) Montages exemplifying out-of-plane division during mitosis 10 in an embryo developing at 25°C. Histone-RFP marks the nuclei (green), while Jupiter-GFP marks the microtubules (magenta). Arrowheads exemplify in-plane divisions, while arrows exemplify out-of-plane division. Each subsequent frame is 1 min apart. In the x-y view of the divisions (A), only one of the daughter nuclei is clearly visible after the out-of-plane division, while the other daughter nucleus is better seen in the y-z view of the division (B), where we can see it translocating to the embryo interior. (C) Plots show the number of out-of-plane divisions, that are visible in the field of view, in embryos developing at various temperatures. For context, an embryo contains on the order of 70 mitotic spindles the field of view at interphase 10. So, even though the p-value is <0.05 for the comparison between 22°C and 29°C, we would argue that the difference is not biologically significant, given that there is no correlation between the number of out-of-plane divisions and the presence of blastoderm holes. n= 10 (22°C), 7 (25°C), 9 (28°C), and 10 (29°C) embryos.

(D) Montage exemplifying isolated nuclear expulsion in an embryo developing at 25°C, from telophase 12 to late interphase 13. Histone-RFP marks the distribution of nuclei, cortical nuclei are in green and subcortical (expelled) nuclei are in magenta. The arrowheads track the daughter nucleus that will be expelled, while arrows track the daughter nucleus that will be retained cortically. (E) Scatter plots showing the distribution of isolated loss of nuclei during interphase 14. n= 10 (22°C), 7 (25°C), 9 (28°C), and 10 (29°C) embryos.

Scale bar; A, 10 µm; D, 5 µm. C and E show scatters with median, with whiskers showing 95% CI. The hollow markers indicate embryos that have a blastoderm hole, with large hollow markers indicating a single large hole, and small hollow markers indicating either a single or multiple small holes. *, p<0.05; non-significant comparisons are not shown.

Figure S4: Defective nuclei take different amount of time to be expelled from embryo cortex.

(A, B) The figure shows the montages of snapshots from live embryos, with a few representative nuclei tracked from late interphase 12 to interphase 13. Top panels show Histone-RFP signal, while bottom panels show Jupiter-GFP signal. Nuclei tagged in green/magenta are the ones that arise from mitotic defects during mitosis 10/11. All of these are expelled during interphase 13. Notice the lack of spindle formation around the tagged nuclei. Also note that the nuclei expelled by the 5th timepoint are not shown in the 6th timepoint. (A) An embryo developing at normal temperature, where 1 magenta and 1 green nuclei are followed. (B) An embryo developing at elevated temperature, where 6 magenta and 3 green nuclei are followed.

Scale bars, 20 µm. The numbers in inset indicate time in min:sec.

Figure S5: Nuclear movement parallel to the embryo surface is amplified at elevated temperature.

(A-C) An example depicting how the nuclear movements are tracked, and the deviations thereof are calculated. Snapshots from live embryo developing at elevated temperature, where Histone-RFP marks the distribution of nuclei, showing the mother nuclei at the end of interphase 10 (A) and the corresponding 8 great-granddaughter nuclei, each, at the end of interphase 13 (C). The neighbouring nuclear clusters are marked in either cyan or yellow. The trajectories of the center of nuclear clusters (B) also indicate the increase in the number of nuclei due to divisions, as the broadening of the tracks. The dotted lines indicate straight-lines fitted to individual trajectories, which are used as reference for computing deviations for each time point of that trajectory. The shaded region captures periods of greatest movement that starts with anaphase (see methods).

(D-I) To quantify the drift in the center of nuclear clusters, we compute the slope of the straight-line fits (dotted lines in B). The median of slopes for all nuclear cluster from one embryo provides a robust estimate of the drift in the centers of the nuclear clusters. Plots showing the distribution of these medians (one dot per embryo), for the drift along AP axis (D), and for that along DV axis (E). To quantify the movements in the center of nuclear clusters during periods of maximum movement (shaded region in B) and normal movement (unshaded regions in B), we compute the average of the deviations from the straight-line fits (dotted lines in B). The median of such average deviations for all nuclear cluster from one embryo provides a robust estimate of the nuclear cluster deviation. Plots showing the distribution of these medians (one dot per embryo), for the fluctuations along AP axis (F, H), and for those along DV axis (G, I), during interphase-metaphase (F, G) or during anaphase-telophase (H, I). n= 10 (22°C), 7 (25°C), 9 (28°C), and 10 (29°C) embryos.

Scale bar, 25 µm. D to I show scatters with median, with whiskers showing 95% CI. Hollow markers indicate data from embryos that have blastoderm hole, with large hollow markers indicating a single large hole, and small hollow markers indicating either a single or multiple small holes. *, p<0.05; **, p<0.01; ***, p<0.001; non-significant comparisons are not shown.

Figure S6: Nuclear movement along surface-yolk axis is amplified at elevated temperature.
Cross-sections from live embryos, expressing Histone-RFP, from late interphase 13 till early interphase 14, in embryos developing either at normal temperature (A), or at elevated temperature (B). Here, the coloured lines mark the basal tip of nuclei in the first frames (t=-7:30 min) of the respective montage. Timepoint 00:00 corresponds to initiation of anaphase at the left/right extreme of the images, which is also the first time we see the shift from baseline.

Arrowheads point out the local peak of the nuclear shift from the baseline, while arrow points expelled nuclei. As we can see, at elevated temperature, the nuclei shift significantly from the baseline, before returning to normal localisation by the time the nuclei enter the subsequent interphase. Also, the shift from baseline occurs progressively from anterior/posterior poles towards the central region.

Scale bars, 25 µm. Cross-sections are in the lateral region of the embryos, anterior to the left.

Figure S7: Nuclear cycle asynchrony increases as a function of the temperature during embryo development.
(A, B) Montages of snapshots from metaphase 13 to early interphase 14, from live embryos developing either at 25°C (A), or at 29°C (B). Histone-RFP marks the nuclei, cortical nuclei are in green and subcortical (expelled) nuclei are in magenta. The dotted lines divide the field of view in 5 equal parts. Numbers in the inset represent time in min:sec, while 00:00 is when anterior $1/5^{th}$ of the embryo enters interphase 14. Boxes highlight entry into interphase 14 in that region. The black arrows indicate the time point of maximum synchrony, i.e., halfway between the anterior and posterior $1/5^{th}$ of the embryo, when the nuclei in the central region should have entered interphase 14. This point serves as the reference to estimate asynchrony (see methods).

Scale bars, 25 µm.

Figure S8: Additional quantifications related to *baz* or *dlg* over-expression.
(A-F) The distribution of the pseudo-cell areas in embryos at the end of interphase 13, in live embryos developing either at normal temperatures (A, C, and E), or at elevated temperature (B, D, and F). The pseudo-cell areas in embryos of control genotype (A, B), in embryos over-expressing *baz* (C, D), or in embryos over-expressing *dlg* (E, F), are shown. Anterior is to the left, dorsal up. The LUT bar at the bottom indicates the colour code corresponding to the area of pseudo-cells.

(G-K) Scatter plots showing the distribution of the area fractions in crowded (G), and combined (H) regions
of the embryos, the distribution of the extent of nuclear crowding (I), the distribution of the nuclear cycle
asynchrony (J), and the number of pairwise nuclear expulsions (i.e., the number of mitotic failures) (K) for
embryos from various genotypes, at normal (22°C) vs elevated temperature (29°C). Each dot represents data
from one embryo: control, n= 7 (22°C), 9 (29°C) embryos; baz-OE, n= 7 (22°C), 7 (29°C) embryos; dlg-OE, n=
6 (22°C), 7 (29°C) embryos.

Scale bar, 25 µm. G-K show scatters with median, with whiskers showing 95% CI. Hollow markers indicate
data from embryos that have blastoderm hole, with large hollow markers indicating a single large hole, and
small hollow markers indicating either a single or multiple small holes. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$;
non-significant differences are omitted.

Figure S9: Elevated temperature disrupts the association between nuclei and F-actin caps.
Distribution of nuclei and F-actin in embryos during interphase 13, in fixed embryos that were developing
either at normal temperatures (A, C, D), or at elevated temperature (B, E, F). (A, B) The images show SUM
projections of the DAPI stained nuclei (top image) and Phallacidin stained F-actin (bottom image) of
representative embryos. In the top images, the nuclei are labelled as a function of their distance from the
embryo surface, where green indicates cortical, and magenta indicates sub-cortical nuclei. The yolk nuclei
are not shown. Anterior is on the left, dorsal up. (C-F) The top and bottom images show a zoomed-in view of
the indicated regions in panels A and B. Here, arrowheads indicate the expelled nuclei (C, E top panels) and
the structures of F-actin corresponding to them (C, E bottom panels), which should otherwise be localised
around cortical nuclei. Then, the arrows indicate gaps between neighbouring nuclei (D, F top panels), which
should normally be devoid of any F-actin structures (D, bottom panel), but contain small "actin caps" in
embryos at elevated temperature (F, bottom panel).

Scale bars, 25µm. The boxed regions are 60x60 µm.

Figure S10: Additional quantifications related to *αCat* or *sgg* over-expression.
(A-F) The distribution of the pseudo-cell areas in embryos at the end of interphase 13, in live embryos
developing either at normal temperatures (A, C, and E), or at elevated temperature (B, D, and F). The pseudo-
cell areas in embryos of control genotype (A, B), in embryos over-expressing *αCat* (C, D), or in embryos over-
expressing *sgg* (E, F), are shown. Anterior is to the left, dorsal up. The LUT bar at the bottom indicates the
colour code corresponding to the area of pseudo-cells. (G-K) Scatter plots showing the distribution of the
area fractions in crowded (G), and combined (H) regions of the embryos, the distribution of the extent of
nuclear crowding (I), the distribution of the nuclear cycle asynchrony (J), and the number of pairwise nuclear
expulsions (i.e., the number of mitotic failures) (K) for embryos from various genotypes, at normal (22°C) vs
elevated temperature (29°C). Each dot represents data from one embryo: control, n= 8 (22°C), 9 (29°C)
embryos; αCat-OE, n= 8 (22°C), 8 (29°C) embryos; sgg-OE, n= 7 (22°C), 7 (29°C) embryos.

Scale bar, 25 µm. G-K show scatters with median, with whiskers showing 95% CI. Hollow markers indicate
data from embryos that have blastoderm hole, with large hollow markers indicating a single large hole, and
small hollow markers indicating either a single or multiple small holes. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$;
non-significant differences are omitted.

Figure S11: *mud* or *insc* over-expression does not ameliorate embryo survival at elevated temperature.
Scatter plot shows the distribution of percent lethality in embryos from various genotypes, at normal (25°C)
vs elevated temperature (29°C), calculated based on the fraction of embryos that don't hatch into larvae.
Each dot represents data from a sample with ≥21 embryos; control, n= 10 (22°C), 11 (29°C) samples; mud-
OE, n= 11 (22°C), 12 (29°C) samples; insc-OE, n= 7 (22°C), 10 (29°C) samples. As we can see in the plot, neither

mud nor *insc* over-expression rescues the embryo lethality at elevated temperature, with *mud* over-
expression increasing embryo lethality even at normal temperature (25°C).

The plot shows scatters with median, while whiskers showing 95% CI. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$;
non-significant differences are omitted.

Table S1: This table contains genes that are identified as being clinal in different continents. Columns A to C
contains lists of studies, and under the studies is a list of genes that are identified in these studies as being
clinal. Column A, B, and C respectively list studies that have data from the continents of Australia and/or
Australasia (AUS), North and/or Central America (USA), and Europe and/or North Africa (EU).

We collect all the genes associated with the GO terms 'cell polarity', 'spindle organization', and 'spindle
localization', to compile a non-redundant master list of candidate genes associated with 'spindle dynamics'
in Column F. For each gene in column F, we count the number of times that gene appears in column A to C,
counting a maximum of once per column, to respectively populate columns G to I. For each gene in column
F, column J contains the sum of values in G, H, and I, to indicate the number of continents in which a gene is
clinal. Each gene gets a score from 1 to 3, and we consider the gene to be a relevant candidate if it is found
in at least 2 columns, i.e., with a score of 2 or 3.

We collect all the genes associated with the GO term 'metabolic process', to compile a control list of
candidate genes in Column L. For each gene in column L, we count the number of times that gene appears in
column A to C, counting a maximum of once per column, to respectively populate columns M to O. For each
gene in column L, column P contains the sum of values in M, N, and O, to indicate the number of continents
in which a gene is clinal. Thus, each gene gets a score from 1 to 3.

Following articles were used to source the list of genes in column A (Oakeshott, McKechnie, and Chambers
1984; McColl and McKechnie 1999; Bettencourt et al. 2002; Frydenberg, Hoffmann, and Loeschcke 2003;
Umina et al. 2005; Sawyer et al. 2006; Weeks, Mckechnie, and Hoffmann 2006; Rand et al. 2010; Kolaczkowski
et al. 2011; Paaby et al. 2014), column B (Vigue and Johnson 1973; Oakeshott, McKechnie, and Chambers
1984; Berry and Kreitman 1993; Duvernell and Eanes 2000; P. S. Schmidt, Duvernell, and Eanes 2000; Verrelli
and Eanes 2001; Duvernell, Schmidt, and Eanes 2003; Sezgin et al. 2004; P. S. Schmidt and Paaby 2008; Rand
et al. 2010; Fabian et al. 2012; Paaby et al. 2014; Machado et al. 2016), and column C (Oakeshott, McKechnie,
and Chambers 1984; Costa et al. 1992; Weeks, Mckechnie, and Hoffmann 2006; Kapun et al. 2020).

Table S2: Column A lists candidates from Table S1 column F that have a score of 2 or higher, i.e., those that
show selectable variation in natural populations. For these genes, we query FlyBase
(http://flybase.org/batchdownload) for additional details like, gene name (column B), gene symbol (column
C), and the level of expression for these genes during the first 2 hours of embryo development (column D).
The level of expression is based on published transcriptomics data (Graveley et al. 2011), which would
correspond to maternal pool of mRNAs, the most relevant developmental window for this study. The
candidate genes clinal in all 3 continents are highlighted in green.

Columns F-J present a table with counts of genes that are associated with various categories of genes in Table
S1 column F and L, and a further breakdown of how many of these genes are clinal in how many continents.
The breackdown is also plotted as pie-charts in Figure 4G.

Table S3: This table lists various figure panels (columns A and B), and the related parental genotypes, i.e., the
genotypes of females (column C) and males (column D) present in the embryo collection cages for the
experiment related to a figure panel.

Additional information

Data and Code Availability:

Data and codes developed for data analysis are available upon request from corresponding authors.

Author contributions:

G.K. and S.L. planned the project. G.K. did the experiments with help from P.A. for baz and dlg overexpressions, and JJ.D.-L. for embryo fixation, and nuclei and F-actin staining. G.K. did the image analyses and statistical analyses. G.K. and S.L. wrote the manuscript, and all authors gave comments on the manuscript.

Acknowledgments:

We would like to thank Lázaro Centanin, Nicolas Gompel, and Marianthi Karageorgi for insightful comments on various drafts of the manuscript, and Lemke lab members for discussions throughout the project. We are grateful to Jochen Wittbrodt for his enthusiasm, and for his continuous and generous support. We would also like to thank Ingrid Lohmann for *Drosophila* culture related support, and Thomas Lecuit for sharing fly lines. We sincerely appreciate the efforts taken by FlyBase for maintaining curated database and Bloomington Drosophila Stock Center for providing transgenic fly lines. This work was funded by the DFG (LE 2787/3-1) to S.L., while G.K. was supported through an HFSP Long-term postdoctoral fellowship (LT000597/2019-L).

Competing interests:

The authors declare no competing financial interest.

References:

Berry, A., and M. Kreitman. 1993. "Molecular Analysis of an Allozyme Cline: Alcohol Dehydrogenase in Drosophila Melanogaster on the East Coast of North America." *Genetics* 134 (3): 869–93. https://doi.org/10.1093/genetics/134.3.869.

Bettencourt, Brian R., Inyoung Kim, A. A. Hoffmann, and M. E. Feder. 2002. "Response to Natural and Laboratory Selection at the Drosophila Hsp70 Genes." *Evolution* 56 (9): 1796–1801. https://doi.org/10.1111/j.0014-3820.2002.tb00193.x.

Bogovic, John A., Philipp Hanslovsky, Allan Wong, and Stephan Saalfeld. 2016. "Robust Registration of Calcium Images by Learned Contrast Synthesis." In *2016 IEEE 13th International Symposium on Biomedical Imaging (ISBI)*, 2016-June:1123–26. IEEE. https://doi.org/10.1109/ISBI.2016.7493463.

Cao, Jian, Justin Crest, Barbara Fasulo, and William Sullivan. 2010. "Cortical Actin Dynamics Facilitate Early-Stage Centrosome Separation." *Current Biology : CB* 20 (8): 770–76. https://doi.org/10.1016/j.cub.2010.02.060.

Chong, Jeronica, Christopher Amourda, and Timothy E. Saunders. 2018. "Temporal Development of Drosophila Embryos Is Highly Robust across a Wide Temperature Range." *Interface* 15 (144). https://doi.org/10.1098/rsif.2018.0304.

Costa, Rodolfo, Alexandre A. Peixoto, Guido Barbujani, and Charalambos P. Kyriacou. 1992. "A Latitudinal Cline in a Drosophila Clock Gene." *Proceedings of Royal Society of London*, 43–49.

Crews, Sarah M., W Tyler McCleery, and M. Shane Hutson. 2016. "Pathway to a Phenocopy: Heat Stress Effects in Early Embryogenesis." *Developmental Dynamics : An Official Publication of the American Association of Anatomists* 245 (3): 402–13. https://doi.org/10.1002/dvdy.24360.

Deneke, Victoria E., Anna Melbinger, Massimo Vergassola, and Stefano Di Talia. 2016. "Waves of Cdk1 Activity in S Phase Synchronize the Cell Cycle in Drosophila Embryos." *Developmental Cell* 38 (4): 399–412. https://doi.org/10.1016/j.devcel.2016.07.023.

Duvernell, David D., and W. F. Eanes. 2000. "Contrasting Molecular Population Genetics of Four Hexokinases in Drosophila Melanogaster, D. Simulans and D. Yakuba." *Genetics* 156 (3): 1191–1201. https://doi.org/10.1093/genetics/156.3.1191.

Duvernell, David D., Paul S. Schmidt, and Walter F. Eanes. 2003. "Clines and Adaptive Evolution in the Methuselah Gene Region in Drosophila Melanogaster." *Molecular Ecology* 12 (5): 1277–85. https://doi.org/10.1046/j.1365-294X.2003.01841.x.

Edgar, Bruce A., Caroline P. Kiehle, and Gerold Schubiger. 1986. "Cell Cycle Control by the Nucleo-
Cytoplasmic Ratio in Early Drosophila Development." *Cell* 44 (2): 365–72.
https://doi.org/10.1016/0092-8674(86)90771-3.
Edwards, Marshall J. 2006. "Review: Hyperthermia and Fever during Pregnancy." *Birth Defects Research.*
Part A, Clinical and Molecular Teratology 76 (7): 507–16. https://doi.org/10.1002/bdra.20277.
Fabian, Daniel K., Martin Kapun, Viola Nolte, Robert Kofler, Paul S. Schmidt, Christian Schlötterer, and
Thomas Flatt. 2012. "Genome-Wide Patterns of Latitudinal Differentiation among Populations of
Drosophila Melanogaster from North America." *Molecular Ecology* 21 (19): 4748–69.
https://doi.org/10.1111/j.1365-294X.2012.05731.x.
Figard, Lauren, Liuliu Zheng, Natalie Biel, Zenghui Xue, Hasan Seede, Seth Coleman, Ido Golding, and Anna
Marie Sokac. 2019. "Cofilin-Mediated Actin Stress Response Is Maladaptive in Heat-Stressed
Embryos." *Cell Reports* 26 (13): 3493-3501.e4. https://doi.org/10.1016/j.celrep.2019.02.092.
Foe, V. E., and B. M. Alberts. 1983. "Studies of Nuclear and Cytoplasmic Behaviour during the Five Mitotic
Cycles That Precede Gastrulation in Drosophila Embryogenesis." *Journal of Cell Science* 61 (May): 31–
70. https://doi.org/10.1242/jcs.61.1.31.
Foe, V.E. 1989. "Mitotic Domains Reveal Early Commitment of Cells in Drosophila Embryos." *Trends in*
Genetics 5 (C): 322. https://doi.org/10.1016/0168-9525(89)90120-0.
Frydenberg, Jane, A. A. Hoffmann, and Volker Loeschcke. 2003. "DNA Sequence Variation and Latitudinal
Associations in Hsp23, Hsp26 and Hsp27 from Natural Populations of Drosophila Melanogaster."
Molecular Ecology 12 (8): 2025–32. https://doi.org/10.1046/j.1365-294X.2002.01882.x.
Graham, John M. 2020. "Update on the Gestational Effects of Maternal Hyperthermia." *Birth Defects*
Research 112 (12): 943–52. https://doi.org/10.1002/bdr2.1696.
Graveley, Brenton R., Angela N. Brooks, Joseph W. Carlson, Michael O. Duff, Jane M. Landolin, Li Yang, Carlo
G. Artieri, et al. 2011. "The Developmental Transcriptome of Drosophila Melanogaster." *Nature* 471
(7339): 473–79. https://doi.org/10.1038/nature09715.
Graziosi, G., F. Micali, R. Marzari, F. de Cristini, and A. Savoini. 1980. "Variability of Response of
EarlyDrosophila Embryos to Heat Shock." *Journal of Experimental Zoology* 214 (2): 141–45.
https://doi.org/10.1002/jez.1402140203.
Harris, Tony J C, and Mark Peifer. 2004. "Adherens Junction-Dependent and -Independent Steps in the
Establishment of Epithelial Cell Polarity in Drosophila." *The Journal of Cell Biology* 167 (1): 135–47.
https://doi.org/10.1083/jcb.200406024.
Herbert, Sébastien, Léo Valon, Laure Mancini, Nicolas Dray, Paolo Caldarelli, Jérôme Gros, Elric Esposito, et
al. 2021. "LocalZProjector and DeProj: A Toolbox for Local 2D Projection and Accurate Morphometrics
of Large 3D Microscopy Images." *BMC Biology* 19 (1): 136. https://doi.org/10.1186/s12915-021-
01037-w.
IPBES. 2019. *The Global Assessment Report of the Intergovernmental Science-Policy Platform on Biodiversity*
and Ecosystem Services. https://doi.org/10.5281/zenodo.3831673.
Kapun, Martin, Maite G Barrón, Fabian Staubach, Darren J. Obbard, R Axel W Wiberg, Jorge Vieira, Clément
Goubert, et al. 2020. "Genomic Analysis of European Drosophila Melanogaster Populations Reveals
Longitudinal Structure, Continent-Wide Selection, and Previously Unknown DNA Viruses." *Molecular*
Biology and Evolution 37 (9): 2661–78. https://doi.org/10.1093/molbev/msaa120.
Khoury, Mark J., and David Bilder. 2020. "Distinct Activities of Scrib Module Proteins Organize Epithelial
Polarity." *Proceedings of the National Academy of Sciences of the United States of America* 117 (21):
11531–40. https://doi.org/10.1073/pnas.1918462117.
———. 2022. "Minimal Functional Domains of the Core Polarity Regulator Dlg." *Biology Open* 11 (7).
https://doi.org/10.1242/bio.059408.
Kolaczkowski, Bryan, Andrew D. Kern, Alisha K. Holloway, and David J. Begun. 2011. "Genomic
Differentiation between Temperate and Tropical Australian Populations of Drosophila Melanogaster."
Genetics 187 (1): 245–60. https://doi.org/10.1534/genetics.110.123059.
Kuntz, Steven G., and Michael B. Eisen. 2014. "Drosophila Embryogenesis Scales Uniformly across
Temperature in Developmentally Diverse Species." *PLoS Genetics* 10 (4): e1004293.
https://doi.org/10.1371/journal.pgen.1004293.
———. 2015. "Oxygen Changes Drive Non-Uniform Scaling in Drosophila Melanogaster Embryogenesis."

F1000Research 4 (May): 1102. https://doi.org/10.12688/f1000research.7221.1.

Lecuit, Thomas, and Alpha S. Yap. 2015. "E-Cadherin Junctions as Active Mechanical Integrators in Tissue Dynamics." *Nature Cell Biology* 17 (5): 533–39. https://doi.org/10.1038/ncb3136.

Lemke, Steffen, Girish Kale, and Silvia Urbansky. 2020. "Comparing Gastrulation in Flies: Links between Cell Biology and the Evolution of Embryonic Morphogenesis." *Mechanisms of Development* 164 (October): 103648. https://doi.org/10.1016/j.mod.2020.103648.

Lockwood, Brent L., Cole R. Julick, and Kristi L. Montooth. 2017. "Maternal Loading of a Small Heat Shock Protein Increases Embryo Thermal Tolerance in Drosophila Melanogaster." *The Journal of Experimental Biology* 220 (Pt 23): 4492–4501. https://doi.org/10.1242/jeb.164848.

Lu, Xuemin, Jennifer M. Li, Olivier Elemento, Saeed Tavazoie, and Eric F. Wieschaus. 2009. "Coupling of Zygotic Transcription to Mitotic Control at the Drosophila Mid-Blastula Transition." *Development (Cambridge, England)* 136 (12): 2101–10. https://doi.org/10.1242/dev.034421.

Lynch, Jeremy A., Ezzat El-Sherif, and Susan J. Brown. 2012. "Comparisons of the Embryonic Development of Drosophila , Nasonia , and Tribolium." *Wiley Interdisciplinary Reviews: Developmental Biology* 1 (1): 16–39. https://doi.org/10.1002/wdev.3.

Machado, Heather E., Alan O. Bergland, Katherine R. O'Brien, Emily L. Behrman, Paul S. Schmidt, and Dmitri A. Petrov. 2016. "Comparative Population Genomics of Latitudinal Variation in Drosophila Simulans and Drosophila Melanogaster." *Molecular Ecology* 25 (3): 723–40. https://doi.org/10.1111/mec.13446.

Mapelli, Marina, and Cayetano Gonzalez. 2012. "On the Inscrutable Role of Inscuteable: Structural Basis and Functional Implications for the Competitive Binding of NuMA and Inscuteable to LGN." *Open Biology* 2 (8): 120102. https://doi.org/10.1098/rsob.120102.

Martin, Adam C. 2020. "The Physical Mechanisms of Drosophila Gastrulation: Mesoderm and Endoderm Invagination." *Genetics* 214 (3): 543–60. https://doi.org/10.1534/genetics.119.301292.

Mazumdar, Aveek, and Manjari Mazumdar. 2002. "How One Becomes Many: Blastoderm Cellularization in Drosophila Melanogaster." *BioEssays : News and Reviews in Molecular, Cellular and Developmental Biology* 24 (11): 1012–22. https://doi.org/10.1002/bies.10184.

McCartney, Brooke M., Donald G. McEwen, Elizabeth Grevengoed, Paul Maddox, Amy Bejsovec, and Mark Peifer. 2001. "Drosophila APC2 and Armadillo Participate in Tethering Mitotic Spindles to Cortical Actin." *Nature Cell Biology* 3 (10): 933–38. https://doi.org/10.1038/ncb1001-933.

McColl, Gawain, and Stephen W. McKechnie. 1999. "The Drosophila Heat Shock Hsr-Omega Gene: An Allele Frequency Cline Detected by Quantitative PCR." *Molecular Biology and Evolution* 16 (11): 1568–74. https://doi.org/10.1093/oxfordjournals.molbev.a026069.

Miller, Morton W., and Charles C. Church. 2013. "Arrhenius Thermodynamics and Birth Defects: Chemical Teratogen Synergy. Untested, Testable, and Projected Relevance." *Birth Defects Research. Part C, Embryo Today : Reviews* 99 (1): 50–60. https://doi.org/10.1002/bdrc.21025.

Oakeshott, J. G., S. W. McKechnie, and G. K. Chambers. 1984. "Population Genetics of the Metabolically Related Adh, Gpdh and Tpi Polymorphisms in Drosophila Melanogaster I. Geographic Variation in Gpdh and Tpi Allele Frequencies in Different Continents." *Genetica* 63 (1): 21–29. https://doi.org/10.1007/BF00137461.

Paaby, Annalise B., Alan O. Bergland, Emily L. Behrman, and Paul S. Schmidt. 2014. "A Highly Pleiotropic Amino Acid Polymorphism in the Drosophila Insulin Receptor Contributes to Life-History Adaptation." *Evolution; International Journal of Organic Evolution* 68 (12): 3395–3409. https://doi.org/10.1111/evo.12546.

Pietzsch, Tobias, Stephan Saalfeld, Stephan Preibisch, and Pavel Tomancak. 2015. "BigDataViewer: Visualization and Processing for Large Image Data Sets." *Nature Methods* 12 (6): 481–83. https://doi.org/10.1038/nmeth.3392.

Preibisch, Stephan, Stephan Saalfeld, and Pavel Tomancak. 2009. "Globally Optimal Stitching of Tiled 3D Microscopic Image Acquisitions." *Bioinformatics (Oxford, England)* 25 (11): 1463–65. https://doi.org/10.1093/bioinformatics/btp184.

Pritchard, David K., and Gerold Schubiger. 1996. "Activation of Transcription in Drosophila Embryos Is a Gradual Process Mediated by the Nucleocytoplasmic Ratio." *Genes & Development* 10 (9): 1131–42. https://doi.org/10.1101/gad.10.9.1131.

Raguso, Robert A. 2020. "Don't Forget the Flies: Dipteran Diversity and Its Consequences for Floral Ecology
and Evolution." *Applied Entomology and Zoology* 55 (1): 1–7. https://doi.org/10.1007/s13355-020-
00668-9.
Rand, David M., Daniel M. Weinreich, Daniel Lerman, Donna Folk, and George W. Gilchrist. 2010. "Three
Selections Are Better than One: Clinal Variation of Thermal QTL from Independent Selection
Experiments in Drosophila." *Evolution* 64 (10): 2921–34. https://doi.org/10.1111/j.1558-
5646.2010.01039.x.
Rim, Ellen Youngsoo, Hans Clevers, and Roel Nusse. 2022. "The Wnt Pathway: From Signaling Mechanisms
to Synthetic Modulators." *Annual Review of Biochemistry* 91 (June): 571–98.
https://doi.org/10.1146/annurev-biochem-040320-103615.
Rizzelli, Francesca, Maria Grazia Malabarba, Sara Sigismund, and Marina Mapelli. 2020. "The Crosstalk
between Microtubules, Actin and Membranes Shapes Cell Division." *Open Biology* 10 (3): 190314.
https://doi.org/10.1098/rsob.190314.
Rudman, Seth M., Sharon I. Greenblum, Subhash Rajpurohit, Nicolas J. Betancourt, Jinjoo Hanna, Susanne
Tilk, Tuya Yokoyama, Dmitri A. Petrov, and Paul S. Schmidt. 2022. "Direct Observation of Adaptive
Tracking on Ecological Time Scales in Drosophila." *Science (New York, N.Y.)* 375 (6586): eabj7484.
https://doi.org/10.1126/science.abj7484.
Rutherford, Suzanne L., and Susan Lindquist. 1998. "Hsp90 as a Capacitor for Morphological Evolution."
Nature 396 (6709): 336–42. https://doi.org/10.1038/24550.
Sangster, Todd A., Susan Lindquist, and Christine Queitsch. 2004. "Under Cover: Causes, Effects and
Implications of Hsp90-Mediated Genetic Capacitance." *BioEssays : News and Reviews in Molecular,
Cellular and Developmental Biology* 26 (4): 348–62. https://doi.org/10.1002/bies.20020.
Sawyer, Lesley A., Federica Sandrelli, Carlo Pasetto, Alexandre A. Peixoto, Ezio Rosato, Rodolfo Costa, and
Charalambos P. Kyriacou. 2006. "The Period Gene Thr-Gly Polymorphism in Australian and African
Drosophila Melanogaster Populations: Implications for Selection." *Genetics* 174 (1): 465–80.
https://doi.org/10.1534/genetics.106.058792.
Schmidt, Anja, and Jörg Grosshans. 2018. "Dynamics of Cortical Domains in Early Drosophila Development."
Journal of Cell Science 131 (7). https://doi.org/10.1242/jcs.212795.
Schmidt, Paul S., David D. Duvernell, and Walter F. Eanes. 2000. "Adaptive Evolution of a Candidate Gene
for Aging in Drosophila." *Proceedings of the National Academy of Sciences of the United States of
America* 97 (20): 10861–65. https://doi.org/10.1073/pnas.190338897.
Schmidt, Paul S., and Annalise B. Paaby. 2008. "Reproductive Diapause and Life-History Clines in North
American Populations of Drosophila Melanogaster." *Evolution* 62 (5): 1204–15.
https://doi.org/10.1111/j.1558-5646.2008.00351.x.
Sezgin, Efe, David D. Duvernell, Luciano M. Matzkin, Yihao Duan, Chen Tseh Zhu, Brian C. Verrelli, and
Walter F. Eanes. 2004. "Single-Locus Latitudinal Clines and Their Relationship to Temperate
Adaptation in Metabolic Genes and Derived Alleles in Drosophila Melanogaster." *Genetics* 168 (2):
923–31. https://doi.org/10.1534/genetics.104.027649.
Telley, Ivo A., Imre Gáspár, Anne Ephrussi, and Thomas Surrey. 2012. "Aster Migration Determines the
Length Scale of Nuclear Separation in the Drosophila Syncytial Embryo." *The Journal of Cell Biology*
197 (7): 887–95. https://doi.org/10.1083/jcb.201204019.
Umina, P. A., A. R. Weeks, M. R. Kearney, S. W. McKechnie, and A. A. Hoffmann. 2005. "Evolution: A Rapid
Shift in a Classic Clinal Pattern in Drosophila Reflecting Climate Change." *Science* 308 (5722): 691–93.
https://doi.org/10.1126/science.1109523.
Vazquez, Julio, Daniel Pauli, and Alfred Tissières. 1993. "Transcriptional Regulation in Drosophila during
Heat Shock: A Nuclear Run-on Analysis." *Chromosoma* 102 (4): 233–48.
https://doi.org/10.1007/BF00352397.
Verrelli, B. C., and W. F. Eanes. 2001. "Clinal Variation for Amino Acid Polymorphisms at the Pgm Locus in
Drosophila Melanogaster." *Genetics* 157 (4): 1649–63. https://doi.org/10.1093/genetics/157.4.1649.
Vigue, C. L., and F. M. Johnson. 1973. "Isozyme Variability in Species of the Genus Drosophila. VI.
Frequency-Property-Environment Relationships of Allelic Alcohol Dehydrogenases in D.
Melanogaster." *Biochemical Genetics* 9 (3): 213–27.
Walter, Marika F., Nancy S. Petersen, and Harald Biessmann. 1990. "Heat Shock Causes the Collapse of the

Intermediate Filament Cytoskeleton in Drosophila Embryos." *Developmental Genetics* 11 (4): 270–79. https://doi.org/10.1002/dvg.1020110405.

Weeks, A. R., S. W. Mckechnie, and A. A. Hoffmann. 2006. "In Search of Clinal Variation in the Period and Clock Timing Genes in Australian Drosophila Melanogaster Populations." *Journal of Evolutionary Biology* 19 (2): 551–57. https://doi.org/10.1111/j.1420-9101.2005.01013.x.

Welte, Michael A., Joan M. Tetrault, Robert P. Dellavalle, and Susan L. Lindquist. 1993. "A New Method for Manipulating Transgenes: Engineering Heat Tolerance in a Complex, Multicellular Organism." *Current Biology* 3 (12): 842–53. https://doi.org/10.1016/0960-9822(93)90218-D.

Xie, Yi, and J. Todd Blankenship. 2018. "Differentially-Dimensioned Furrow Formation by Zygotic Gene Expression and the MBT." *PLoS Genetics* 14 (1): e1007174. https://doi.org/10.1371/journal.pgen.1007174.

Zheng, Liuliu, Leonardo A. Sepúlveda, Rhonald C. Lua, Olivier Lichtarge, Ido Golding, and Anna Marie Sokac. 2013. "The Maternal-to-Zygotic Transition Targets Actin to Promote Robustness during Morphogenesis." *PLoS Genetics* 9 (11): e1003901. https://doi.org/10.1371/journal.pgen.1003901.

Figure 1

Increased embryonic lethality at elevated temperature

A Temperature regimes during embryonic development

(N) Normal temperature = 25°C   (E) Elevated temperature = 29°C

NN

EN

NE

0hr   3hr   6hr   30hr

Time since egg fertilisation

B   Lethality at the end of embryo development

NN

EN

NE

5   10   15   20

% dead embryos

Nuclear loss phenotype at elevated temperature

25°C   29°C (mild defect)   29°C (severe defect)

C   D   E

inter-11 late

inter-12 late

inter-13 late

inter-14 early

inter-14 late

Mitotic failure and pairwise nuclear expulsion
(Example from embryo at 29°C)

F   Histone-RFP

−5 min   0 min   5 min   10 min   15 min

G   number of mitotic failures causing nuclear expulsions during interphase 14

number of lost pairs of nuclei

22°C   25°C   28°C   29°C

H

Normal temperature   Elevated temperature

•• Normal mitosis   ▬▬ Mitotic failure and pairwise loss of nuclei

1280

Figure 2

Asynchrony in the phases of nuclear division cycles
(Example from embryo at 29°C)

Increased nuclear crowding at elevated temperature

Figure 3

Figure 4

Over-expressing regulators of 'cell polarity signalling during mitosis'
to rescue embryonic lethality at elevated temperature

Over-expressing regulators of 'F-actin -microtubule interaction during mitosis'
to rescue embryonic lethality at elevated temperature

"Natural" candidate genes varying along thermoclines

1287   **Figure S1**

1288

Figure S2

Nuclear loss phenotype due to elevated temperature

Figure S3

Histone-RFP Jupiter-GFP (Example from embryo at 25°C)

A   X-Y view

B   Y-Z view

C   out-of-plane divisions during mitosis-10

number of divisions

22°C   25°C   28°C   29°C

D   isolated nuclear expulsion
(Example from embryo at 25°C)

Histone-RFP

0 min   4 min

7 min   12 min

E   number of isolated nuclear expulsions
during interphase 14

number of lost nuclei

22°C   25°C   28°C   29°C

1293 **Figure S4**

1294

Figure S5

1297   **Figure S6**

1298

1299   **Figure S7**

Histone-RFP

1300

1301 **Figure S8**

1302

Figure S9

1305   **Figure S10**

1306

1307 **Figure S11**

1308