Elevated temperature fatally disrupts nuclear divisions in the early *Drosophila* embryo.

Girish Kale[1,2,*], Pratika Agarwal[1], J Jaime Diaz-Larrosa[1], Steffen Lemke[1,2,*]

[1] Centre for Organismal Studies, University of Heidelberg, Heidelberg. Germany

[2] Institute of Biology, University of Hohenheim, Stuttgart, Germany

* Correspondence: G.K. (girish.kale@uni-hohenheim.de), S.L. (steffen.lemke@uni-hohenheim.de)

Abstract:

Temperature variations can challenge animal survival, with elevated temperatures presenting distinct
vulnerabilities at different stages of the animal life cycle. Embryonic development is known to be especially
vulnerable, though the molecular mechanisms that underlie this vulnerability remain unclear. Here we
investigate how elevated temperature affects embryogenesis in *Drosophila melanogaster*, an insect model
known for extensively characterized genetic makeup and developmental processes.

We identify the first three hours as a critical window in which *Drosophila* embryos are particularly vulnerable
to elevated temperature. This period includes the formation of a syncytial blastoderm, which involves 4
rounds of meta-synchronous nuclear divisions at the embryonic cortex, and cellularization that leads to the
formation of the cellular blastoderm. Embryos exposed to elevated temperature during this period
subsequently exhibited developmental defects at the gastrulation stage, leading to increased lethality. At
elevated temperature, we observed an increase in mitotic failures causing a loss of cortical nuclei during
cellularization. There is also a local crowding of nuclei and an increase in asynchrony between the nuclear
cycles in the center vs. the poles. Interestingly, these features cooperatively amplify the frequency of mitotic
failures, leading to holes in the blastoderm epithelium.

Further mechanistic analysis of mitotic failures revealed that they trigger DNA damage response. Focusing
on cell biological causes of mitotic failures, we found that the interaction between F-actin and microtubules
is weakened at elevated temperature. We performed functional tests to determine whether known
regulators of mitosis could rescue mitotic failures, blastoderm hole formation, and improve embryo survival.
We further analyzed genomic datasets from wild populations to determine whether these regulators also
carry signatures of adaptation in wild populations.

Our genetic rescue experiments show that in *Drosophila* embryos, the interaction between cortical F-actin
and astral microtubules of mitotic spindle is vulnerable to disruption at elevated temperatures, leading to
mitotic failures that can be potentially rescued by modulating the expression of just a few factors. We
propose that levels of expression of corresponding genes could be used as indicators to predict the effects
of increasing temperature variations on insect populations.

Keywords:
Blastoderm formation, elevated temperature, microtubule - F-actin interaction, nuclear division defects,
embryo lethality, genetic rescue, *Drosophila*

Introduction:

It has been suggested that the current rise in average global temperatures and the amplitude of temperature fluctuations can affect species fitness and result in a loss of biodiversity[1]. It is unclear whether effects on species fitness are direct consequences of the temperature rise, particularly since many species are likely experiencing a rather gradual departure from optimal conditions of temperature, which have previously evolved in a species-specific manner. Previous studies suggest that, under some circumstances, even such gradual departures could affect animal survival. For instance, studies have shown that a gradual departure from normal temperatures, e.g., experienced by fevers during pregnancy produce birth defects in humans[2–4], indicating that temperature conditions tolerable by an adult may be more detrimental to the embryo. This raises the possibility that specific aspects of embryo development are more vulnerable to temperature increase. Consistent with this notion, previous studies in model organisms have shown that temperature is a key factor that determines the fidelity of embryonic development; rat and zebrafish embryos show an increase in developmental defects at elevated temperatures[5–8]. Yet the mechanisms underlying vulnerability in embryonic development at elevated temperatures remain unclear. Here we explore whether exposure to elevated temperature is detrimental to embryonic development in the insect model *Drosophila melanogaster* (henceforth, *Drosophila*), and examine the consequences in terms of population-level parameters such as embryo survival.

In nature, animals rarely experience highly steady temperatures, but they do experience a range of temperatures to which they are generally well adapted. Increasing temperatures accelerate diverse physiological processes, including development, growth, and regeneration, making the timing of events temperature-dependent[9]. Developmental processes can be viewed as networks of biochemical reactions occurring simultaneously or in rapid succession. The Arrhenius equation, long used to describe the temperature dependence of biochemical reactions, has more recently been applied to capture the effects of temperature on the rate of physiological processes, particularly during embryonic development[10,11]. However, in contrast to chemical reactions *in vitro*, where temperatures often exceed physiological ranges, complex biological processes *in vivo* – such as embryonic development – follow the Arrhenius equation, provided that temperatures remain within the relatively narrow range typically experienced by the organism. With a transition from 28°C to 30°C, *Drosophila* no longer exhibit an acceleration of embryonic development[10–12], suggesting that 29°C marks a threshold at which elevated temperatures challenge development and exceeds the physiological range over which the Arrhenius equation accurately describes developmental rates, while lower temperatures remain effectively normal. To test this hypothesis, we analyzed the fidelity of development at an elevated temperature of 29°C, compared with controls at the normal culture temperature of 25°C.

Previous studies have found that *Drosophila* embryo development beyond the first 4-6 hours is robust to even acute exposure to high temperatures[13], suggesting that potential embryonic vulnerabilities are likely to occur and be observed during the first few hours of development. *Drosophila* embryo development starts with fertilization and egg deposition. Following fertilization, the embryo undergoes nine rounds of syncytial nuclear divisions within the yolk. Following these divisions, most nuclei are anchored to the periphery of the embryo, where four more rounds of nuclear division take place during a stage called syncytial blastoderm[14]. These divisions are facilitated by pseudo-cleavage furrows, i.e., temporary membrane invaginations that resemble a half-cell and slowly form around the nuclei during interphase-metaphase. They are subsequently reabsorbed into the embryo cortex rapidly during anaphase-telophase[15]. Following a total of 13 nuclear divisions, the resulting ~6000 nuclei simultaneously segregate into individual cells and produce the blastoderm epithelium in a process called cellularization[16]. This occurs at about 3 hrs from fertilization. It marks the beginning of gastrulation, which involves changes in cell shape[17], rearrangements of cell contacts[18],

and further rounds of cell divisions in isolated mitotic domains[19]. All of these events facilitate early
morphogenesis.

In our experiments, we found that exposing embryos to 29°C during the first 3 hours of embryonic
development leads to a significant increase in population-level embryonic lethality during the gastrulation
stage. Examinations at the cell and tissue level revealed mitotic failures, which lead to the loss of nuclei during
the syncytial blastoderm and especially the cellularization stage. Additionally, embryos developing at 29°C
showed nuclear crowding in the central region of the embryo and greater asynchrony of nuclear division
cycles between the central regions and at the poles. We show that nuclear crowding and nuclear cycle
asynchrony combine to increase the local propensity of mitotic failures. Local clusters of mitotic failures lead
to regions devoid of nuclei/cells ('holes'), which disrupts the continuity of the blastoderm as primary
epithelial tissue. The molecular dissection of mitotic failures in the blastoderm revealed that they trigger a
DNA damage response, which mitotically arrests the defective daughter nuclei, resulting in their expulsion
from the embryo cortex. We show that cytoskeletal elements, F-actin and microtubules, show incongruent
dynamics at 29°C, suggesting a weakening of their interaction during mitosis, implicating mitotic failures at
the core of developmental vulnerabilities at elevated temperature. In an attempt to enhance the resilience
of nuclear divisions at 29°C and rescue temperature-induced developmental defects, we overexpressed
specific genes that regulate cell polarity, microtubule-F-actin interactions, and mitotic spindle orientation.
Our results suggest a genetically rescuable vulnerability in microtubule-F-actin interactions during nuclear
divisions at elevated temperatures. Notably, the genes we identified as capable of providing a genetic rescue
to temperature-induced developmental defects could be recovered in a genome-wide analyses of data on
genetic variation in natural populations along temperature clines. In summary, our work identifies cell
biological defects caused by elevated temperatures and links them to population-level lethality and fitness.
This offers insights into the broader implications of vulnerability to elevated temperatures during a defined
time span in the life history of flies and provides an initial handle to quantitatively and mechanistically predict
embryo fitness at elevated temperature.

Results:

Early exposure to elevated temperature reduces embryo survival due to gastrulation defects

To explore how elevated temperature affects the early development of *Drosophila* embryos, we exposed
young embryos to 29°C and asked how many of them survived and how they developed. To investigate
alterations in the rate of embryos hatching into larvae, we first differentiated the effects of temperature
exposure by timing: from the moment of deposition until just before gastrulation begins (~3 hours), and
during a similar time starting from gastrulation onset (refer to Figure 1A for experimental temperature
regimen). In both scenarios, we observed a notable increase in embryonic lethality (Figure 1B), pinpointing a
critical vulnerability especially during the pre-gastrulation phase of development. This finding was
unexpected, as prior research suggested that the genetic mechanisms governing embryonic patterning
during this phase were resilient to temperature fluctuation[20,21], indicating a more systemic effect, possibly
rooted in cell biology.

To address when and how embryos died after early exposure to 29°C, we maintained embryos until late
cellularization stage at 29°C and then recorded their development using confocal microscopy time-lapse
imaging at 25°C. Embryonic development in these recordings was analyzed qualitatively as well as
quantitatively. Qualitatively, many embryos displayed moderate anomalies during gastrulation, including
twisting along the embryo's anterior-posterior axis (Figure 1C-D), exemplifying transient defects in the
bilateral symmetry of embryonic development. Notably, these twisted embryos seemed to recover from their
defects and continued to develop. By contrast, a minority of embryos exhibited severe gastrulation defects,

culminating in developmental cessation and lethality (Figure 1E). Quantitatively, surviving embryos after exposure to 29°C did not differ statistically significantly from those after control exposure to 25°C, neither in the rate of germband extension (GBE, Figure 1F) nor in the timing of other key developmental milestones (Figure 1G). Overall, this suggests that if embryos surpass the gastrulation stage, they proceed without detectable developmental delays.

Taken together, our findings underline increased embryonic lethality during gastrulation due to prior exposure to 29°C, highlighting the detrimental effects of elevated temperature. Previous work has demonstrated that genetic patterning is robust to temperature perturbations[20,21], suggesting that the observed defects are unlikely to be directly caused by a failure of embryonic patterning per se. To explain the observed fatal defects during gastrulation stage, we instead hypothesized that nuclei, who ultimately interpret embryonic patterning information into cell and tissue movements, might have exhibited fatal defects in their dynamic behavior during blastoderm formation. To address this possibility, we investigated nuclear dynamics during pre-gastrulation development.

A clustered loss of nuclei at elevated temperature likely causes gastrulation defects and embryo lethality

Pre-gastrulation development in flies and other insects involves nuclei dividing deep within the yolk before their arrival at embryo's cortex, where they must be securely anchored while undergoing further rounds of division. Ultimately, the tight and even cortical distribution of those syncytial nuclei is crucial for the generation of blastoderm epithelium and its subsequent morphological changes during gastrulation.

To address whether defects during gastrulation could stem from defective nuclear distribution rather than defective blastoderm patterning, we focused our analyses on the distribution of cortical nuclei during cellularization, i.e., just before gastrulation, using fixed embryos stained for DNA to visualize nuclei in the embryo volume (see methods). In embryos developing at 29°C, we observed a subcortical cluster of nuclei in a small fraction of embryos, corresponding to a hole in the cortical layer of nuclei (Figure 2A). To better visualize and differentiate cortical and subcortical nuclei, we segmented the embryo volume and color-coded the image voxels based on their distance from the embryo surface (Figure 2B). We noticed a few subcortical nuclei in embryos developing at normal temperature (Figure 2C, F, I, J), while a large fraction of embryos developing at 29°C showed an increase in subcortical nuclei, albeit not in large clusters (Figure 2D, G, K , L). Notably, however, in a small fraction of embryos developing at 29°C, we observed a notable void in the cortical layer of nuclei (henceforth, blastoderm hole), coinciding with a subcortical cluster of at least 10 nuclei (Figure 2E, H, M, N). Within the coordinate system of the embryo, the distribution of neither isolated subcortical nuclei nor blastoderm holes was left-right symmetric. Additionally, even if blastoderm holes were predominantly observed in the central region of the embryo, their location varied in their position across embryos (Figure 2O, P). These results further support the idea that loss of nuclei and blastoderm holes are unlikely to be directly caused by defective patterning: embryonic patterning is left-right symmetrical and accordingly, observed defects in patterning would likewise show left-right symmetry.

To investigate the underlying cause for the appearance of subcortical nuclei, we followed nuclear dynamics using time-lapse recordings: Histone-RFP was used to visualize chromatin, and Jupiter-GFP to visualize microtubules for keeping track of nuclear division cycles. Starting with the first appearance of nuclei at the embryo cortex (occurring during the interphase of the 10th nuclear cycle), we observed nuclei through their cortical divisions during syncytial blastoderm (through mitoses 10 to 13), then through cellularization (interphase 14), and until the onset of gastrulation (beyond interphase 14). To determine whether the nuclear loss at 29°C reflects a threshold response associated with a transition to non-Arrhenius behavior, we examined temperature dependencies additionally at 22°C, 25°C, and 28°C, temperatures within the normal physiological range.

The cortical arrival of nuclei during interphase 10, from the yolk to the embryo cortex, was comparable between embryos developing at 22°C, 25°C, 28°C, and 29°C (Figure 3A-E, top panels), indicating that the preblastoderm stage spreading of nuclei along the AP axis[22] was comparable between these temperatures. During the syncytial blastoderm stage and cellularization, however, embryos that developed at 29°C much more frequently "lost" nuclei from their cortical layer than those developing at 22, 25, and 28°C, i.e., nuclei that were initially positioned within the cortical layer of the egg, appeared to get expelled, descended into the yolk, and did not become part of the embryonic blastoderm (Figure 3, also see below), with variability in the extent of the nuclear loss, producing blastoderm holes in a fraction of embryos (Figure S1).

Taken together, in pre-gastrulation embryonic development we narrowed down the developmental vulnerability at 29°C to the syncytial blastoderm stage and cellularization, where we see nuclear loss from the embryo cortex. Notably, the fraction of embryos showing blastoderm holes (Figure 2A, Figure S1) was similar to the fraction of embryos with gastrulation defects (Figure 1F) and was consistent with embryo lethality at 29°C (Figure 1B). The embryo-to-embryo variability, both in the distribution and intensity of nuclear loss, further supports the argument that these defects were not dictated by defective embryonic patterning, and were likely caused by defects in non-deterministic self-organised processes. While consistent with the idea that the presence of blastoderm holes was causing the observed increase in embryonic lethality, the temperature-dependent mechanism leading to blastoderm holes was still unclear. To further address the mechanistic link between elevated temperature and embryo lethality, we decided to quantify the loss of nuclei and possible patterns that could be predictive of blastoderm holes.

At elevated temperature, embryos lose nuclei from the cortex after defective divisions.

Closer analyses of lost cortical nuclei indicated that these nuclei were lost either individually or as pairs of two adjacent nuclei. To identify their relative contribution to the early developmental vulnerability at elevated temperatures, we decided to quantify these two types of nuclear loss from the embryo cortex.

Our analyses first focused on the isolated loss of nuclei. An isolated nucleus could be lost by a division perpendicular to the embryo surface (out-of-plane division), where the nucleus facing the inside of the embryo may not get properly anchored to the cortex and is consequently lost into the yolk. Alternatively, an isolated nucleus may be lost after an in-plane division, e.g. if steric hindrance and insufficient space would not allow for simultaneous anchoring of two adjacent nuclei to the embryo cortex. To account for these possibilities, we visually tracked every individually lost nucleus back in time to its last division.

Out-of-plane divisions could be detected in embryos at all of our experimental temperatures. These out-of-plane divisions could be easily identified, as they formed a characteristic mitotic spindle that was oriented perpendicular to the embryo surface (Figure 4A), and after the division only one of the daughter nuclei remained attached to the cortex. However, out-of-plane divisions occurred only during the mitoses of the 10th nuclear cycle, thus couldn't explain the loss of nuclei from the 11th nuclear cycle onwards. Loss of nuclei following an in-plane division could also be detected. These losses occurred at a low frequency throughout the syncytial blastoderm stage and cellularization, at all tested temperatures (Figure 4B). When we quantified the isolated loss of nuclei, we noted that (i) the overall frequency of out-of-plane divisions was very low (<5%) (Figure 4C), (ii) loss of individual nuclei after an in-plane division was rather infrequent, and, even if slightly more frequent during interphase 14 (cellularization), (iii) the loss of individual nuclei did not differ significantly within the range of experimental temperatures (Figure 4D). The overall extremely low frequency of isolated nuclear loss – even in embryos developing at 29°C – strongly suggests that isolated nuclear loss was not the main driver of temperature-dependent nuclear loss in the fly blastoderm.

Next, we focused on the pairwise losses of nuclei. As above, we tracked lost pairs of nuclei to their last division
cycle. This tracking revealed that every pairwise loss of nuclei was, in fact, a loss of sister nuclei. This
observation suggested that specific characteristics of cortical nuclear divisions might determine the future
fate of the newborn nuclei and trigger their release from the egg cortex, possibly in response to failed
checkpoints. Consistent with this idea, we observed chromosome segregation defects, and a delay in
telophase, preceding the pairwise loss of nuclei (Figure 4E), at all of our experimental temperatures. Because
chromosome segregation defects can lead to aneuploidy, our observations suggested that the subsequent
expulsion of daughter nuclei from the egg cortex was due to prior defective mitosis. To address whether the
observed loss of nuclei at 29°C could be attributed to an overall increase in defective mitoses, we asked if
their frequency was changing depending on the temperature (see methods). For this, we focused on
interphase 14, which is when we observed the most nuclear losses. While the frequency of mitotic failures
was overall low at 22 to 28°C, it increased significantly at 29°C (Figure 4F). Of note, defective mitoses, both
at normal or at elevated temperature, were not associated with a stereotypical time course of nuclear
expulsion. Defective nuclei could be lost in the interphase immediately following defective divisions, or later
(Figure S2), suggesting that nuclear expulsion after defective mitosis was a passive process and not triggered
by molecular signaling at failed checkpoints.

Taken together, our findings identify nuclear divisions in the blastoderm as a thermolabile cellular process
that shows a disproportionate increase in failures with the transition from 28°C to 29°C, resulting in the loss
of sister nuclei from the egg cortex, when born in defective mitoses. However, which characteristics of
nuclear divisions render them thermolabile was unclear.

At elevated temperature, nuclei displace drastically, especially during divisions.

For embryos at 25°C, we noted that the nuclei were rather dynamic in their localization, showing local
movements, throughout the syncytial blastoderm stage. For embryos at 29°C, this dynamicity increased, and
we could see nuclei being displaced rather drastically. We hypothesized that such displacement poses a
challenge to nuclear anchorage at elevated temperatures, especially during mitosis, causing mitotic defects
and the paired loss of nuclei. To that end, we tracked nuclear groups starting from individual mother nuclei
at end of interphase 10 to the 8 great-granddaughter nuclei at end of interphase 13. At each time point, we
computed the center of the nuclear group, using the positions of the nuclei in that group, yielding one track
per group (Figure 5A-C, also see methods). Our tracking showed that the centers of nuclear groups had two
components to their movement, a global drift and a local displacement.

First, the nuclear groups exhibited a global consistent drift-like movement, which represented a concerted
global movement of all nuclei. We captured this component using a linear fit to the tracks of the center of
the nuclear groups over time (Figure 5B dotted lines). The slope of the linear fit indicated that the groups
drifted towards the posterior end at all of our experimental temperatures (Figure 5D, E). This component of
nuclear movements was reminiscent of the previously reported dynamic reordering of nuclei in the early
syncytial blastoderm[23]. Notably, however, the extent of drift in nuclear groups did not correlate with the
occurrence of blastoderm holes (Figure 5D, hollow markers).

Second, independent of this drift, we observed local deviations in the position of the nuclear groups over
time. We decided to measure these local deviations relative to the global drift. We reasoned these local
deviations could be a proxy for mechanical stresses experienced by the nuclear division machinery, and
quantified these deviations during interphase-metaphase and anaphase-telophase. The deviations relative
to the constant drift were more drastic along the anterior-posterior (AP) axis (Figure 5F, H) than those along
the dorsal-ventral (DV) axis (Figure 5G, I), and deviations during the anaphase-telophase period (Figure 5H)
were more drastic than those during the interphase-metaphase period (Figure 5F). In the context of length

scales typical for nuclear divisions, this meant that nuclear group deviations in embryos developing at 25°C
and during anaphase-telophase were comparable to that of the length of the spindle itself, while for embryos
at 29°C, nuclear group deviations could be up to 4-fold higher. This increase in nuclear group deviations
suggested increased mechanical stresses especially during the critical anaphase-telophase period, indicating
an extensive propensity for displacement relative to embryo cortex at a time when the dividing nuclei need
to be robustly anchored to the embryo cortex. Consistent with this interpretation, embryos with blastoderm
holes showed the highest nuclear group deviations especially during anaphase-telophase (Figure 5H, large
hollow markers).

To further test the link between mitotic failures and the short-range local nuclear displacements, i.e., nuclear
group deviations, we followed the division of each of the 8 great-granddaughter nuclei in every tracked
nuclear group at 29°C from mitosis 13 to interphase 14, expecting to see mitotic failures in nuclear groups
with greater displacements. Here, however, we failed to see a correlation between extent of nuclear group
deviations and a subsequent mitotic failure (Figure 5J). In that, the nuclear groups with one or more
subsequent mitotic failures (Figure 5J, hollow markers) do not show particularly higher deviations compared
to their normally dividing counterparts. In fact, when we count the number of nuclear groups with
subsequent mitotic failure defects, we see that 15 of these groups were below their respective embryo
medians, while only 7 are above, arguing against a correlation between increased nuclear displacements and
subsequent mitotic failures during interphase 14.

In addition to the analysis of nuclear movements parallel to the embryo cortex we wondered about the
movements orthogonal to cortex, i.e., the future apical-basal (AB) axis in the blastoderm cells[24]. In embryos
developing at 25°C, we observed that nuclei maintained a constant distance from the cortex (Figure S3A).
However, in embryos developing at 29°C, we could see two waves of nuclear displacement towards the
cortex, which originate from the anterior and the posterior, and meet at the central region of the embryo
(Figure S3B). These waves of nuclear displacement were tightly correlated with waves of nuclear divisions,
with peaks of the waves corresponding to anaphase-telophase transition. This suggests that as nuclei divide
the spindle moves upwards, presumably to gain more space. This need for space indicated additional
challenges for the spindles during nuclear division, explaining, at least in parts, the mitotic failure at 29°C.

Taken together, our quantifications and observations captured an embryo-wide increase in nuclear
displacement locally, along the AP- and AB-axis. In a fraction of embryos, the mitotic failures occurred in
great numbers and in close vicinity, leading to holes in the blastoderm epithelium at 29°C (Figure 4F, hollow
markers), and these embryos also exhibited greater displacement of nuclei during anaphase-telophase
(Figure 5H, hollow markers). Notably, however, our quantifications and observations of an increased nuclear
displacement could not reveal any correlation with nuclear loss at the level of individual nuclear groups
(Figure 5J). This result argues against the idea of increased dynamicity being the cause of nuclear division
defects at 29°C. Thus, although the formation of blastoderm holes (Figure S1) could be directly attributed to
clustered mitotic failures, it remained unclear why mitotic failures were stochastically distributed in most
embryos, yet locally clustered in the central region of a small fraction of embryos, producing blastoderm
holes.

Nuclei are crowded and divide with a time shift in embryos at elevated temperature.

To better understand the differences between stochastic mitotic failures and locally clustered failures, we
focused on cases with severe nuclear losses in embryos at 29°C. The most severe loss of nuclei in embryos at
29°C was often observed during cellularization (interphase 14) in the central domain of the embryo (Figure
2, 3 and S1). Thus, we wondered if the central regions of the embryo were somehow special, and if there

were characteristics in nuclear distribution or nuclear division, already during nuclear cycle 13, that rendered
the central regions to be distinct than the anterior/posterior poles.

The central region of the embryo had more tightly packed nuclei, indicating a defect in self-organized nuclear
reordering mechanisms[23]. To quantify this nuclear crowding, we focused on the end of interphase 13 (see
methods), as that's the time we observed maximum nuclear crowding, before the most intense nuclear loss.
To quantify the space available for cortical blastoderm nuclei, we segmented the nuclei and used Voronoi
tessellation to define a region around each nucleus, which we call pseudo-cells. The areas of pseudo-cells
were rather homogeneous in embryos developing at 22°C (Figure 6A, E, I). Already at 25°C (Figure 6B, F, J)
and 28°C (Figure 6C, G, K), the available space for cortical nuclei was no longer uniform across the embryo,
and it showed signs of local crowding in the central regions of the embryo. At 29°C, the space available for
nuclei along the anterior-posterior axis varied substantially and revealed strong nuclear crowding in the
central region of the embryo (Figure 6D, H, L). We quantified the extent of crowding (see methods), and
found a progressive increase in nuclear crowding (Figure 6M), with peak crowding at 29°C. To specifically test
whether the central region of the embryo was distinct from the poles in inter-nuclear spacing, we
quantitatively described elevated nuclear crowding at 29°C in the central region of the embryo (Figure 6N).

The central regions of the embryos at 29°C coincided with the "collision" of two waves of nuclear divisions
starting from the poles (see above, Figure S3). This observations was in stark contrast with embryos at 25°C,
where nuclear divisions are meta-synchronous[25], with anterior-posterior waves (Figure 6O). We could further
observe that in contrast to the meta-synchrony at 25°C, phases of nuclear division in the central region at
29°C were lagging behind those at the anterior/posterior pole of the embryos (Figure 6P). Accordingly, the
same phase of the nuclear division cycle could be observed at different time points in different regions of the
embryo (Figure S4A-B), indicating defects in the self-organised coordination of nuclear division cycles across
the syncytium[25]. Thus, we decided to quantify this asynchrony of nuclear division cycles at temperatures
22°C, 25°C, 28°C, and 29°C, as nuclei exited nuclear cycle 13 and entered cycle 14 (see methods). Our
quantifications show that the asynchrony significantly increases in embryos developing at 29°C as compared
to other temperatures (Figure 6Q). This indicated that the central region of the embryo is distinct from the
poles in nuclear division dynamics, with the phases of the nuclear cycle lagging (Figure 6R).

Taken together, in the central region of the *Drosophila* syncytial blastoderm embryo, nuclear division cycles
lag behind those at the anterior/posterior poles, and nuclear density is greater than that at the
anterior/posterior poles. Given that blastoderm holes also appear in the central regions of the embryo, we
tested whether the clustering of mitotic failures (leading to blastoderm holes) at 29°C correlated better with
the regions of highest nuclear crowding, or the most delayed nuclear cycles, or both.

Region of embryo with greater nuclear crowding and with delayed nuclear cycles are associated with
greater frequency of mitotic failures.

To test whether nuclear crowding and/or nuclear cycle asynchrony resulted in a predisposition towards a
local increase in mitotic failure, we first defined the regions of the embryo with high nuclear crowding (Figure
7A) at the end of interphase 13, or regions with substantially delayed nuclear division cycles (Figure 7B) at
the end of mitosis 13. In parallel, we marked the locations of mitotic failures occurring during cellularization
(interphase 14), which could be overlaid on any predefined region to calculate the fraction of mitotic failures
in that region. Then, to measure the fraction of mitotic failures associated with either nuclear crowding, or
nuclear cycle asynchrony, or both, we constructed the regions of the embryo with either high nuclear
crowding, i.e., crowded regions (Figure 7C); or substantial delay in nuclear division cycles, i.e., delayed regions
(Figure 7D); or a combination of crowded and delayed regions, i.e., combined region (Figure 7E) (see
methods).

In crowded regions, the fraction of mitotic failures increased progressively with higher temperatures, with
statistically significant differences between 22 and 29°C (Figure 7F). In delayed regions, the fraction of mitotic
failures showed an abrupt increase from 28 to 29°C, albeit with only marginal statistical significance (Figure
7G). In the combined regions, the fraction of mitotic failures was greater than those in the crowded or
delayed regions alone (Figure 7H). Additionally, the fraction of mitotic failures in all of the tested
temperatures could not be explained by the area fractions themselves (Figure 7I-K). This suggests that the
combination of nuclear crowding and nuclear cycle asynchrony contributes to a temperature-dependent
predisposition toward mitotic failure in the syncytial blastoderm, with nuclear crowding exerting the stronger
influence.

One plausible mechanism is that local nuclear crowding limits the space for proper mitotic spindle formation,
thereby increasing the likelihood of mitotic errors. In addition, crowding may locally elevate the Nuclear-to-
Cytoplasmic (N/C) ratio, a parameter that reflects the amount of cytoplasm available to each nucleus. The
N/C ratio is known to regulate the pace of nuclear division cycles, such that higher N/C ratios are associated
with slower nuclear division cycle[25–30], likely due to limited cytoplasmic resources required for the cell cycle.
Nuclear crowding would therefore slow nuclear cycle indirectly through its effect on N/C ratio. Consistent
with this interpretation, the reduced area of the 'combined regions' at 29°C (Figure 7K) indicates substantial
spatial overlap between crowded regions and delayed regions, particularly in embryos exhibiting blastoderm
holes. Thus, at elevated temperature, these combined regions likely reflect both the direct mechanical
consequences of crowding and its indirect effects mediated through altered nuclear division timing.

Taken together, the spatial overlap between nuclear crowding and nuclear cycle asynchrony helps explain
why clustered mitotic failures arise preferentially in the central region of the embryo, leading to blastoderm
holes. However, while this framework clarifies where and how mitotic failures are spatially organized, it does
not explain why elevated temperature enhances their frequency. This raises the question of how elevated
temperature perturbs mitosis at the molecular level.

Mitotic failures trigger DNA damage response

To distinguish whether the observed loss of nuclei were primarily caused by temperature-induced DNA
damage leading to mitotic failure, or instead by a temperature-induced failure of mitosis that subsequently
resulted in damaged DNA, we employed a previously described DNA damage reporter system[31]. Briefly, this
reporter relies on the nuclear localization of Stem-Loop Binding Protein (SLBP), which normally shuttles
between nucleus and cytoplasm during interphase, exporting the mRNAs for key early zygotic genes[31].
Following DNA damage, Chk2 phosphorylates SLPB at Serine 118, thus restricts nuclear localization of SLBP,
targeting the protein to degradation. Thus, exclusion of SLBP from the interphase nucleus can act as a live
marker for DNA damage in that nucleus (Figure 8A), provided that the nuclear envelope remains intact.

To determine the cause for the cortical loss of nuclei, we used the absence of SLBP nuclear accumulation as
an indicator for DNA damage and followed such nuclei over a nuclear division cycle. All nuclei displaying
evidence of DNA damage were mitotically arrested and were subsequently lost into the interior of the
embryo (Figure 8B). DNA damage therefore consistently coincided with mitotic failures and, just like the
mitotic failures, affected sister nuclei.

To quantify the kinetics of SLBP nuclear localization over the different phases of the nuclear division cycle,
we measured SLBP-GFP nuclear intensity from late interphase 13 through early interphase 14. SLBP showed
high nuclear localization at the end of interphase 13, was then excluded from the nuclei during mitosis, and
progressively re-accumulated in nuclei at the onset of interphase 14 (Figure 8C, F). This behavior was
comparable between 25 and 29°C during normal nuclear divisions. In contrast, following mitotic failure, we

observed a progressive reduction in SLBP nuclear localization within the first ~4 min of interphase 14,
followed by a more pronounced loss over the next ca. 10 minutes (Figure 8F), after which the nuclei were
typically expelled from the forming blastoderm. The absence of differences in SLBP localization during the
preceding interphase, together with indistinguishable behavior during mitosis, indicates that activation of
the DNA damage response occurs specifically during interphase 14 and only after the mitotic failure.

Mitotic failure can trigger a variety of signaling responses, raising the possibility that the observed loss of
SLBP nuclear localization reflects a response other than the DNA damage signaling. To address this, we
followed the nuclear localization of SLBP.S118A-GFP, a point mutant that cannot be phosphorylated at serine
118 and is therefore insensitive to Chk2-mediated DNA damage signaling[31]. Quantification of SLBP.S118A
nuclear localization revealed behavior comparable to wildtype SLBP in normally dividing nuclei (Figure 8D,
G). In nuclei with DNA damage, however, SLBP.S118A was retained in the nucleus during the early phase of
interphase 14, remaining detectable for approximately 4-5 minutes, before being progressively lost from the
nucleus at later time points (Figure 8G). These observations suggest that the loss of nuclear SLBP following
mitotic failure comprises at least two components; an early, DNA damage-dependent component and a later
component that is independent of SLBP phosphorylation at serine 118.

DNA damage can induce nuclear envelope destabilization, resulting in the release of nuclear proteins,
including SLBP. To contextualize the similarities and differences in the kinetics of SLBP and SLBP.S118A, we
therefore analyzed the behavior of a GFP reporter bearing a nuclear localization signal (NLS-GFP). In normally
dividing nuclei, NLS-GFP showed nuclear localization dynamics similar to those of both SLBP and SLBP.S118A
(Figure 8E, H). In contrast, following mitotic failure, the kinetics of NLS-GFP localization closely resembled
those of SLBP.S118A, but not wildtype SLBP (Figure 8F-H). This indicates that the altered nuclear localization
observed specifically during the early phase of interphase 14 reflects activation of the DNA damage response,
which is captured by SLBP-GFP but absent in both SLBP.S118A-GFP and NLS-GFP.

To quantify this DNA damage-dependent effect, we compared the rates of nuclear accumulation during the
initial phase of interphase 14. SLBP nuclear import was significantly slower in nuclei undergoing division
defects at 29°C compared to normally dividing nuclei at the same temperature (Figure 8I). In contrast, nuclear
import rates of both SLBP.S118A (Figure 8J) and NLS (Figure 8K) were unaffected by mitotic failure during this
early interphase period.

Taken together, these results indicate that DNA damage arises as a consequence of mitotic failure rather
than acting as its primary cause. Moreover, the DNA damage response is rapidly engaged, becoming
detectable within minutes after entry into interphase 14. This rapid onset highlights mitosis as a critical point
of vulnerability and underscores the importance of understanding the mechanisms underlying mitotic failure
in order to improve embryonic robustness at elevated temperatures.

Cytoskeletal dynamics during mitosis at elevated temperature

Our results so far link the vulnerability of mitosis at elevated temperature – and its central role in the
formation of blastoderm holes – with embryo lethality at elevated temperature. Given the essential roles of
the cytoskeletal elements, F-actin and microtubules, in orchestrating mitosis, we next asked whether
elevated temperature alters cytoskeletal dynamics during syncytial nuclear divisions. Specifically, we
quantified changes in cytoskeletal organization during mitosis at 29°C as compared to 25°C, and examined
how these dynamics differed between normally completing divisions and mitotic failures.

During syncytial divisions, cortical F-actin undergoes extensive remodeling across the nuclear division cycle,
most prominently supporting the formation of the transient pseudo-cleavage furrows during mitosis[15,24,32].
These furrows progressively deepen perpendicular to the embryo surface while elongating parallel to it,

encapsulating the microtubule spindle throughout division. The spindle axis aligns with the long axis of the
pseudo-cleavage furrow (Figure 9A), and proper spindle elongation is essential for faithful chromosome
segregation. Defects in pseudo-cleavage furrow formation and/or mitotic spindle organization are therefore
expected to compromise mitotic fidelity.

Based on these considerations, we visualized F-actin-based pseudo-cleavage furrows and microtubule-based
mitotic spindles simultaneously in the same embryos, and quantitatively compared their morphology
between 25°C and 29°C, as well as between normal divisions and mitotic failures.

During mitosis, F-actin caps are elongated and expand to fill the cortical space, thereby outlining the pseudo-
cleavage furrows. Using these outlines, we measured the area of the F-actin caps and estimated aspect ratio
by fitting an ellipse, which also allowed us to determine the orientation of the long axis (Figure 9A). In parallel,
to characterize microtubule spindle geometry, we measured spindle length by connecting the spindle poles
and extracted the orientation of the spindle axis (Figure 9A). Qualitatively, both F-actin caps and microtubule
spindles appeared intact and comparable in overall intensity and structure across conditions, with the
notable exception of their relative alignment (Figure 9B-E). This prompted a quantitative comparison of F-
actin cap and spindle geometry.

We found that F-actin caps were significantly larger at 29°C than at 25°C, both in normally completing
divisions and in divisions that subsequently failed (Figure 9F), while their aspect ratios remained comparable
across conditions (Figure 9G). In contrast, spindle length was selectively reduced in divisions that went on to
fail, irrespective of temperature (Figure 9H). Consistent with this observation, spindles associated with
mitotic failures at 29°C showed little to no net elongation during mitosis (Figure 9I).

Together with the increased F-actin cap area, the reduction in spindle length indicated that the distance
between spindle poles and the cortical F-actin network is increased at elevated temperature, especially in
case of mitotic failures. Under these conditions, spindles appeared less constrained by the cortex and were
frequently observed to rotate. To quantify this behavior, we measured changes in spindle orientation
between consecutive time points throughout metaphase and summed these angular displacements.
Cumulative spindle rotation was significantly higher at 29°C than at 25°C, both in control divisions and in
mitotic failures (Figure 9J). In addition, spindle axes were often misaligned with the long axis of the
corresponding F-actin caps. Quantification of the angular mismatch between spindle and cap orientations at
metaphase confirmed that this misalignment was increased at 29°C in both control divisions and mitotic
failures (Figure 9K).

Taken together, these analyses indicate that mitotic spindles are less securely anchored to cortical F-actin at
elevated temperature. We propose that this weakened coupling increases the susceptibility of spindles to
rotational instability, thereby elevating the probability of mitotic failure. In combination with the spatial bias
of mitotic failures toward central regions of the embryo and the tight coupling between mitotic failure and
subsequent DNA damage, these findings identify mitotic failure as the primary driver of developmental
vulnerability at elevated temperature.

Testing robustness of cell polarity establishment during syncytial divisions at elevated temperature

Our findings thus far identify mitosis in the early blastoderm as a temperature-labile process and link mitotic
failure to embryo lethality at elevated temperature. This raised the question of how a deeply conserved and
generally robust process such as mitosis could become destabilized over a narrow temperature range. We
therefore asked whether increasing the abundance of key components of the nuclear division machinery
could compensate for temperature-induced fragility and restore developmental robustness at 29°C.

A distinctive feature of syncytial divisions in the *Drosophila* blastoderm is their reliance on transient pseudo-
cleavage furrows, which are repeatedly assembled and disassembled at each nuclear cycle. These furrows
form through localized membrane insertion from late interphase to metaphase, support the alignment of
mitotic spindles parallel to the embryo surface during pro- and metaphase, and regress rapidly during
anaphase and telophase. As a consequence, each nuclear cycle requires the repeated reestablishment of
cortical polarity cues alongside extensive cytoskeletal reorganization[24]. This dependence on transient cortical
polarity represents a major deviation from canonical cell division in fully cellularized tissues and may
constitute an evolutionarily young and potentially labile feature of insect embryogenesis. Moreover, polarity
signaling is closely intertwined with orientation of mitotic spindles[33], and cytoskeletal organization in
epithelia[34]. We therefore tested whether temperature sensitivity during syncytial mitoses reflects insufficient
robustness of polarity establishment: we aimed to strengthen polarity signaling through overexpression of
two key polarity regulators active in the early blastoderm, Bazooka (baz; Par-3 homolog) and Discs-large
(dlg)[35]. Specifically, we asked whether either overexpressions could rescue embryo lethality at 29°C, and if
so, whether this rescue was accompanied by a reduction in mitotic failures in crowded and combined regions
of the blastoderm.

In embryos at 25°C, we did not observe a significant change in embryo lethality with either *baz* or *dlg*
overexpression as compared to control embryos with wildtype levels of baz and dlg (Figure S5A). At 29°C,
lethality increased in control embryos (Figure S5A), similar to results in previous experiments (Figure 1B). By
contrast, embryos with either *baz* or *dlg* overexpression performed notably better at 29°C, with lethality
reducing to about two-thirds (*baz* overexpression) and half (*dlg* overexpression) of the controls (Figure S5A).
We note however that the rescue of lethality at 29°C in *baz* overexpressing embryos was only marginally
significant, with a lot of variability between samples. Additionally, *dlg* overexpressing embryos at 25°C
showed higher variability in embryo lethality than their control counterparts. Consistent with these results,
baz overexpression did not rescue the increase in the fraction of mitotic failures, either in the crowded
regions or in the combined regions at 29°C (Figure S5B, C), and while the large blastoderm holes were absent
in these embryos, *baz* overexpression occasionally produced small gaps in blastoderm (Figure S5B, C). Then,
dlg overexpression rescued the increase in the fraction of mitotic failures only in the crowded region at 29°C
(Figure S5B, C), and while the large blastoderm holes were absent in these embryos, *dlg* overexpression
increased the fraction of mitotic failures at normal temperature in both, the crowded region and in the
combined region, occasionally producing small gaps in blastoderm (Figure S5B, C).

Taken together, these results indicated that polarity establishment during syncytial divisions is comparatively
robust to elevated temperature and that deviation from canonical polarity mechanisms are not, by
themselves, sufficient to explain temperature-induced mitotic failure. Instead, the partial rescue effects
suggest that the primary temperature-sensitive vulnerability lies downstream of polarity establishment, at
the level of cytoskeletal regulation required to anchor and stabilize the mitotic spindle. This led us to focus
on molecular interactions that directly link cortical F-actin to astral microtubules of the spindle during mitosis.

Identifying vulnerabilities in cytoskeletal interactions during mitosis at elevated temperature

Since strengthening polarity establishment was not sufficient to robustly suppress mitotic failures at elevated
temperature, we next turned to molecular interactions acting downstream of polarity cues that directly
execute spindle positioning and chromosome segregation. Proper assembly and anchoring of the mitotic
spindle at the cortex requires a dynamic interaction between microtubules and F-actin, which together
anchor chromatin at the cortex throughout nuclear division[24]. Faithful segregation of sister chromatids
therefore depends on tightly regulated interactions between the actin cortex and the astral microtubules of
the mitotic spindle[36,37].

In *Drosophila*, these critical actin-microtubule interactions have been associated with a tripartite protein complex consisting of α-Catenin (αCat), which associates with cortical F-actin; β-Catenin (βCat, or *armadillo* in *Drosophila*), which binds to α-Catenin; and Adenomatous polyposis coli 2 (Apc2), which links to βCat and binds to the astral microtubules of the mitotic spindle[38]. Within this complex, the interaction of αCat and βCat is direct and appears constitutive, whereas the interaction of βCat and Apc2 is mediated by a specific kinase, the *Drosophila* homolog of the Glycogen Synthase Kinase 3β, Shaggy (sgg)[39].

We reasoned that, if this tripartite protein complex was thermolabile, then our high temperature conditions should, at least in parts, phenocopy the loss of function of one of the components of the complex (αCat, βCat, or Apc2) or sgg. The loss of function of this protein complex has been associated with an "empty actin caps" phenotype, where cortical dome-like structures of F-actin are no longer associated with nuclei[38]. Thus, to test for the possibility that this tripartite protein complex is thermolabile during early embryonic development in *Drosophila*, we analyzed embryos that were fixed and stained to visualize nuclei and cortical F-actin (see methods), both at 25°C (Figure S6A) and 29°C (Figure S6B). At control temperature of 25°C, we could observe a tight association between nuclei and F-actin during interphases in the syncytial blastoderm, where cortical F-actin surrounds each nucleus, forming the actin cap. Notably, it was possible to identify gaps between neighboring actin-caps in cases of recently expelled nuclei (Figure S6C), or in cases where the spacing between nuclei is large enough (Figure S6D). Similar to embryos at 25°C, such gaps could also be observed between neighboring actin-caps in cases of recently expelled nuclei at 29°C (Figure S6E). In contrast to controls at 25°C, however, embryos developing at 29°C additionally exhibited the empty actin caps phenotype (Figure S6F), reminiscent of embryos mutant for *Apc2* or *βCat* (see Figure 3d-f in McCartney et al. 2001[38]). Thus, we hypothesized that stabilizing the activity of the complex, through elevated levels of αCat, βCat, Apc2, or sgg, could increase the fidelity of cortical mitoses at 29°C, possibly through a tighter control of actin and microtubule coupling. Such tighter coupling, in return, might decrease the fraction of mitotic failures in crowded and combined regions and ultimately attenuate embryonic lethality.

To test the effects of elevated levels of either αCat or sgg on embryo lethality at control and elevated temperatures, we overexpressed either gene and quantified embryo lethality. At 25°C, we did not observe a significant difference in embryo lethality between control embryos and those with *αCat* or *sgg* overexpression (Figure 10A). At 29°C, lethality was overall increased in controls, similar to results in similar experiments described above (Figure 1B, Figure S5A). By contrast, embryos with either *αCat* or *sgg* overexpression survived significantly better at 29°C, with lethality reducing to about half (*αCat* overexpression) or a third (*sgg* overexpression) relative to the controls (Figure 10A). Consistent with these results, large blastoderm holes were absent in these embryos, and the fraction of mitotic failures in crowded as well as in the combined regions was reduced compared to control embryos at 29°C (Figure 10B, C). These results suggest that overexpression either αCat or sgg strengthen the interaction between microtubule and F-actin, which in turn attenuates temperature-dependent destabilizations of the mitotic spindle. We therefore propose that the scaffolding activity of αCat, βCat, and Apc2, mediated by sgg, constitutes a thermosensitive controller of nuclear division fidelity in the syncytial *Drosophila* embryo.

To test whether thermosensitivity of nuclear division fidelity was dependent specifically on the scaffolding activity of αCat, βCat, and Apc2 rather than a common capacity of any regulator of mitotic spindle anchorage, we asked whether the elevated levels of other regulators could similarly increase robustness against elevated temperatures. To address this question, we focused on regulators known to dictate the orientation of the mitotic spindle during cell division[40]. Both, the activity of 'Mushroom body defect' (mud, NuMA homolog) and that of Inscuteable (insc) has been previously described to regulate the anchorage and orientation of the mitotic spindle during cell division. A similar activity might be present for nuclear divisions during the syncytial blastoderm stage, raising the possibility that an overexpression of *mud* or *insc* could rescue embryo

560 lethality, presumably by reducing mitotic failures. However, neither *mud* nor *insc* overexpression ameliorated
561 the high embryo lethality at 29°C (Figure S7), supporting the notion that elevated temperature specifically
562 challenges the core machinery of the mitotic spindle rather than being a non-specific failure that could be
563 rescued by any mitosis associated regulator.

Thermosensitive regulators of spindle dynamics exhibit natural variation along thermoclines.

While temperatures above 25°C are generally considered non-standard rearing conditions for *Drosophila* lab
strains, naturally occurring populations likely do experience temperatures above this presumed optimum
temperature. Since our analyses showed that elevated temperatures during the first hours of fly
development affect both, embryonic cell biology and organismal fitness, it may be tempting to predict that
natural populations could be adapting to temperature changes through standing genetic variation and
shifting allele frequencies in the same or similar genes that we identified as thermolabile regulators of nuclear
divisions. To address this possibility, we analyzed the distribution of standing genetic variations associated
with the regulators of spindle dynamics (Figure 10D) and asked whether genetic variation would be
distributed randomly with respect to the continent-scale gradients of annual mean temperature, i.e.
thermoclines. While most standing genetic variation associated with putative 'spindle dynamics' regulators
was indeed distributed without a clear thermoclinal pattern and did not allow for rejection of our null
hypothesis, (Figure 10E, Table S1, S2), *sgg* was a clear and notable outlier in this analysis and exhibited a
thermoclinal distribution of natural variations independently in populations in the North America, Europe,
and Australia. Additionally, the fraction of genes associated with 'spindle dynamics' that showed thermoclinal
pattern, was greater than the fraction of genes associated with the GO-term 'metabolic process' that showed
thermoclinal pattern (Figure 10E, Table S1, S2). These findings strongly suggest that the thermosensitivity
that *sgg* exhibits in lab conditions constitutes a biological property, which can be exposed and selected for
also in natural populations and during the course of evolution.

Discussion:

Early fly development is susceptible to elevated temperature

Our work demonstrates that processes of syncytial blastoderm formation and cellularization in *Drosophila*
embryonic development are vulnerable to elevated temperature. Embryos exposed to 29°C during these
stages exhibit defects in maintaining the bilateral symmetry of developmental processes, and in extreme
cases, enter a lethal trajectory during gastrulation (Figure 1). We argue that these extreme cases of
developmental failure likely coincide with structural defects in the blastoderm epithelium, which manifest as
regions devoid of nuclei/cells and discontinuities in this tissue (Figure 2, 3 and S1). The result is holes in the
blastoderm, which arise from a greater incidence of mitotic failures in a small region of the embryo during
syncytial nuclear divisions (Figure 4). Our analysis also establishes a link between mitotic failures and the
subsequent loss of nuclei, via DNA damage response (Figure 8). We show that mitotic failures trigger DNA
damage response in the defective daughter nuclei, which are then mitotically arrested, lose the integrity of
their nuclear envelop, and are "kicked-out" of the blastoderm. We note that the DNA damage repones is
active only in nuclei with DNA damage, and not in its normal neighbors. It remains to be determined whether
these lost nuclei deplete transcription factors required for local cell fate specification. This will, in turn, reveal
whether blastoderm holes cause gastrulation defects purely through mechanical discontinuity, or whether
loss of patterning information also contributes.

Our analysis shows that clustering of mitotic failures occurs in embryonic regions that lag behind in the
nuclear division cycle and exhibit nuclear crowding (Figure 6). While we treat these factors as independent
for most of their analysis, a combination of these factors – captured in 'combined regions' – predispose

particular regions of the embryo to increased mitotic failures at 29°C (Figure 7). Thus, our findings are
consistent with previous studies, which suggest that they are mechanistically coupled through the N/C
ratio[25–30]. These findings suggest a mechanism in which the dependence of nuclear cycle duration on nuclear
density could amplify the likelihood for mitotic failure at elevated temperature; whether a similar interplay
affects embryonic development under normal temperature conditions remains unclear.

Alternatively – or additionally – the source of vulnerability of early blastoderm formation could stem from
the non-deterministic distribution of nuclei, which relies on self-organization and is driven by the activity of
cytoskeletal elements such as F-actin and microtubules[23]. A related form of self-organization may be seen in
the progression of nuclear division cycles, which move across the surface of the embryo as non-deterministic
waves[25]. Both processes appear to be challenged at elevated temperatures, suggesting that developmental
mechanisms relying on biological self-organization may be particularly vulnerable at elevated temperature.
Given the widespread importance of self-organization in biology, it may be valuable to explore more
generally whether developmental processes that depend heavily on self-organizing components constitute
common points of vulnerability.

Potential impact of elevated temperature on essential protein-protein interactions

We show that mitosis itself is vulnerable at elevated temperatures due to weakened interactions between
microtubules and F-actin (Figure 9). The tripartite complex, involving αCat, βCat, and Apc2 (mediated by the
protein kinase sgg) has been implicated in mediating this interaction[38], at least partially, and in our
experiments overexpressing *αCat* or *sgg* rescue the developmental defects and embryo lethality at elevated
temperature (Figure 10A-C). These results support the idea that compromised coupling between
microtubules and F-actin represents a key molecular vulnerability at elevated temperature.

A mild temperature increase such as the one we studied here is unlikely to denature proteins, but may
instead weaken protein-protein interactions, such as the ones between components of the tripartite scaffold,
involving αCat, βCat, and Apc2 (mediated by the protein kinase sgg). Such weakening would effectively
deplete functional protein scaffolds, creating a loss-of-function-like scenario, as the equilibrium shifts
towards the dissociated state of the scaffold. Nuclear crowding can exacerbate this effect: in crowded
regions, individual nuclei experience a relative reduction in the local concentration of the scaffold owing to
an increase in the N/C ratio. Thus, the combination of elevated temperature and local crowding provides a
plausible mechanism for regional predisposition to mitotic failure and the formation of blastoderm holes.
This interpretation is consistent with the observation that overexpression of either *αCat* or *sgg* rescues these
defects (Figure 10A-C), as increasing even a single component should be sufficient to raise overall scaffold
abundance and counteract both temperature-dependent dissociation and crowding effects.

Independent support for this mechanistic framework comes from previous work in a different context. The
nuclear crowding defect we report here is reminiscent of a previous study, focusing on embryos with
increased maternal cyclin B, though at normal tempeatures[41]. While the impact of elevated temperature on
cyclin B activity remains to be characterized, this study found that genetic perturbations enhancing
microtubule-F-actin interactions suppress the phenotype. Even though the upstream perturbation differs
from heat stress, the convergence on the same interaction interface supports the view that microtubule-F-
actin coupling represents a common point of vulnerability.

Given that protein-protein interactions are omnipresent in biology, it is unlikely that microtubule-F-actin
coupling is the only process directly modulated by temperature, and other aspects of basic cell biology may
similarly be vulnerable. Because of their reliance on formation of protein scaffolds, it may be particularly
rewarding to assess potential temperature sensitivity in processes such as cellular signaling[42,43],

mitochondrial metabolism[44,45], cell polarity pathways[46,47], or the regulation of membrane trafficking[48], the
cell cycle[49], and cytoskeletal dynamics[50,51], all of which are conserved across animal kingdom.

Elevated temperature is distinct from heat shock effects

Previous studies in *Drosophila* have shown that temperatures of 33°C or higher trigger a classic heat-shock
response, characterized by *de novo* transcription heat-shock protein genes[52]. This response has been
observed across embryos[53], larvae, pupae[54,55], and adult flies[56], and overexpression of heat-shock proteins
such as hsp23 can increase embryo survival under acute heat stress[57]. Heat shock proteins, accordingly, are
often considered to act as capacitors, buffering cells against environmental and genetic variation[58,59].

However, the extreme temperatures that provoke a heat-shock response are only transiently encountered
in natural environments[55]. Consequently, these conditions may not impose strong selective pressure to
improve tolerance to sustained temperature increases associated with global warming. Supporting this, our
analyses of three continental datasets did not reveal consistent allelic variation in canonical heat-shock
genes. We can't exclude, however, that the lack of allelic variation stems from a limited standing genetic
variation, given their critical role in survival, precluding the evolution of thermoclinal distributions. In either
case, it is unlikely that heat-shock proteins would constitute a mechanism to improve developmental
vulnerability in face of global warming.

In contrast, we observed allelic variation in *sgg* along thermoclines, suggesting it has been shaped by natural
selection. Seasonal shifts in sgg allele frequencies provide additional evidence, whereas heat-shock genes
show no such pattern[60]. These observations suggest that while heat-shock proteins protect against acute
temperature fluctuations, they may not confer selective advantage under elevated average temperatures.
Instead, regulators of cytoskeletal interactions, such as sgg, could be targets of natural selection and serve
as mechanisms to mitigate the effects of gradual temperature increases.

Early embryonic vulnerability to elevated temperatures: implications for insect survival and evolution

Early developmental processes from fertilization to blastoderm formation are highly conserved across
insects[61,62], suggesting that the vulnerabilities we report in *Drosophila* embryos could apply to many other
species. Weakening of cytoskeletal interactions under elevated temperature leads to asymmetric nuclear
distributions, disrupting positional information for subsequent morphogenesis and increasing the risk of
embryo lethality. Such cellular vulnerabilities provide a mechanistic link between temperature stress and
developmental failure, one that may be broadly conserved across diverse insect taxa.

Flies and other insects are essential for ecosystem stability[63], and rising temperatures pose a tangible threat
to their survival. Reports of contemporary insect biodiversity loss[64,65] resonate with patterns of extinction
and diversification during past climatic fluctuations[66,67]. Although multiple abiotic factors contribute to these
historical trends[68–70], the current rapid rise in global temperature highlights heat stress as a unique relevant
driver. Limited data make it difficult to test whether past warming events directly drove innovation in
blastoderm architecture, but it is intriguing to speculate that early developmental processes may serve as
both vulnerabilities and substrates for evolutionary change.

By identifying the cellular and molecular "breaking points" of heat sensitivity during early development, we
can better predict which species and life stages are most at risk, guide conservation strategies, and even
explore genetic pathways that enhance developmental robustness. Understanding these mechanisms offers
a potential bridge from cell-biology to ecosystem-level predictions in a warming world.

Materials and methods:

<u>Fly lines and genetics</u>:

OregonR line was used as WT *Drosophila*, and *w1118* line was used for outcrosses.

Histone-RFP expresses RFP tagged version of His2Av. Insertion on 3rd chromosome. Bloomington Drosophila Stock Center (BDSC) #23650.

Jupiter-GFP expresses GFP tagged version of the microtubule-binding protein Jupiter. Insertion on 3rd chromosome. Always used in combination with *Histone-RFP*. Gift from Jan Felix Evers.

NLS-GFP expresses GFP tagged Nuclear Localization Signal peptide. Insertions of 2nd and 3rd chromosome. Always visualized alongside Histone-RFP. BDSC #1691.

UtrABD-GFP expresses GFP tagged version of Actin-Binding Domain of Utrophin. Insertion on 3rd chromosome. Always used in combination with *Histone-RFP* and *Jupiter-GFP*. Gift from Thomas Lecuit.

67-Gal4 (mat αTub-GAL4-VP16) is a maternal specific Gal4 driver. Insertion on 2nd chromosome. Always used in combination either with *Histone-RFP* or *Histone-RFP* and *Jupiter-GFP* in the overexpression experiments (see below). Gift from Thomas Lecuit.

SLBP- and *SLBP.S118A-GFP* expresses GFP tagged versions of SLBP and SLPB.S118A under UAS promoter. Insertions on 3rd chromosome. Gifts from Eric Lecuyer. For these overexpression experiments, we first crossed virgins from the Gal4 line '+; 67-Gal4; Histone-RFP' to the males of overexpression line '+; +; UAS-X' (X= *SLBP*, or *SLBP.S118A*). F1 virgin females from these crosses were backcrossed to males of the paternal genotype.

The following lines were used to drive the overexpression of various genes of interest in an attempt to rescue the effects of 29°C. *baz*, BDSC #17613; *dlg*, BDSC #15779; *αCat*, BDSC #65590; *sgg*, BDSC #15886; *mud*, BDSC #22371; *insc*, BDSC #39676. An outcross to *w1118* was used as control. For these overexpression experiments, we first crossed virgins from the Gal4 line '+; 67-Gal4; Jupiter-GFP, Histone-RFP' to the males of overexpression line 'UAS-Y; +; +' (Y= *baz*, *dlg*, *sgg*, or *mud*) or '+; +; UAS-Z' (Z= *αCat*, or *insc*), or to males of *w1118* as control. F1 virgin females from these crosses were backcrossed to males of the paternal genotype.

Genotypes associated with various figure panels are listed in Table S3.

<u>Embryo collection</u>: For any experiment, males and females of appropriate genotypes (Table S3) were placed into transparent cylindrical embryo collection cages (inner diameter ~5 cm). The cage top was closed with fine-wire mesh for air exchange, and bottom with apple-juice agar plates. We prepared the apple-juice agar plates from 1 l of solution: we boiled 24 gm of agar in 750 ml of water, adding 250 ml of fresh store-bought apple juice while stirring on a hot plate, adjusting Sucrose to final amount of 25 gm (accounting for the sugar content of the apple juice). This solution yielded ~80 petri dishes (6 cm diameter). After the agar solidified, the plates were stored at 4°C. Before the embryo collection, the plates were covered with a thin layer of yeast paste (to stimulate oviposition) and were pre-incubated to the embryo collection temperature for ≥30 min. Since *Drosophila* females can retain embryos long after fertilization, allowing early embryo development inside their abdomen, we performed a 1 h "sham" embryo collection at room temperature (24-25°C), and discarded these embryos. Following sham collections, we collected the embryos for experiments at specific experimental temperature.

<u>Embryo lethality experiment</u>: We chose embryo-to-larva transition as a conservative and technically simple read-out for embryo survival. This excludes any potential later-arising defects in viability, behavior, lifespan or fertility, meaning that our assay underestimates, rather than overestimates, the deleterious effects of

elevated temperature. For estimating the survival, embryos were exposed to three temperature regimes (Figure 1A). Depending on the regimen, we collected embryos for 1 hour either at 24-25°C (normal temperature) or 29-30°C (elevated temperature). The 24-25°C collections were done on the bench top. The 29-30°C collections were done in a water bath, with embryo collection plate bottom in thermal contact with the heated water. For NN or EN, we collected embryos for 1 hour at 24-25°C or 29-30°C, respectively, and then incubated for 2 hours at the same temperature. The embryos were then bleached for 10-15 seconds, rinsed thoroughly with tap water, aligned on a 24x60 mm coverslip, and covered with halocarbon oil. For NE, we collected embryos for 1 hour at 24-25°C, incubated for 1.5 hours at the same temperature, and within 30 min processed as above (bleached, rinsed, aligned, and covered), and subsequently incubated for 3 hours at 29-30°C. After these treatments, embryos from all regimes were incubated at 24-25°C for ~1 day, following which we counted the number of eclosed larvae and dead embryos. The embryonic lethality was calculated as,

$$embryo\ lethality\ = \frac{100 * number\ of\ dead\ embryos}{number\ of\ dead\ embryos\ +\ number\ of\ larvae}$$

excluding the unfertilized embryos from calculations.

<u>Imaging live embryos (after temperature treatment)</u>: Here we used 2 of the experimental temperature regimes used in the 'Embryo lethality experiment' above, namely NN and EN. In short, we collected embryos at 24-25°C (NN) or 29-30°C (EN) for 1 hour and then incubated the plate at the same temperature for another 2 hours. Then, the embryos were bleached for 35-40 sec, rinsed thoroughly with tap water, aligned with the dorsal side facing a 22x40 mm coverslip, covered with halocarbon oil, and immediately taken for microscopy. The embryos were recorded for at least 15 hours, at 25°C. Images were acquired on Leica SP8 DMi8 inverted confocal microscope using 20x glycerol objective (0.75NA), at 2 µm step-size in z, and a final x-y resolution of 0.568 µm/pixel. We recorded the time-lapse at 1 frame every 4 min and recorded up to 5 embryos in parallel.

<u>Imaging live embryos (at various temperatures)</u>: We used four temperature regimes. We collected embryos for 1 hour either at normal temperatures (22-23°C, 24-25°C, or 27-28°C) or elevated temperature (29-30°C). Collections at 22-23°C (22°C imaging) were done in the fly incubator; 24-25°C (25°C imaging) on the bench; 27-28°C (28°C imaging) or 29-30°C (29°C imaging) in a water bath, with embryo collection plate bottom in thermal contact with the heated water. Then, the embryos are bleached for 35-40 seconds, rinsed with tap water at the collection temperature, aligned on a 25 mm diameter coverslip, covered with halocarbon oil, and immediately transferred to the temperature controller setup from Warner Instruments (QR-40LP holder on QE-1HC heater/cooler, coupled to LCS-1 liquid cooling and CL-100 bipolar temperature controller, precision of ± 0.1°C), pre-equilibrated to the experimental temperatures (22, 25, 28 or 29°C). For the data in Figures 3-7, 10, S1-5, imaging was performed on Leica SP5 DMI6000CS inverted confocal microscope using 40x/1.1 NA water objective, 1x digital zoom, 2 µm z-steps (20 planes), and 0.379 µm/pixel x-y resolution; acquiring images at 1 frame every 30 sec, 1 min, or 2 min depending on whether 1, 2, or 3-4 embryos were recorded in parallel. For the data in Figures 8 and 9, the imaging was performed on Zeiss LSM 700 inverted confocal microscope using 63x/1.2 NA water objective, 0.5x digital zoom, 2 µm z-steps (12 planes), and 0.198 µm/pixel x-y resolution; acquiring images at 1 frame every 30 sec or 1 min depending on the signal strength for the genotype.

<u>Imaging fixed embryos</u>: We collected embryos for 3-4 hours either at 24-25°C (normal temperature) or 29-30°C (elevated temperature). Collections at 24-25°C were done on the bench top; 29-30°C done in a water bath, with embryo collection agar plate bottom in thermal contact with the heated water. Then, the embryos were bleached for 40-45 seconds, rinsed with tap water, and fixed immediately. We only image and analyze the embryos that are before the gastrulation stage. We performed two types of experiments; one, whole-

embryo nuclear imaging (Figure 2); and two, half-embryo imaging of nuclei and F-actin (Figure S6), using distinct fixation/staining protocols.

For **whole-embryo nuclear imaging**, we used heat fixation and stained the embryos with DRAQ5. Briefly, we transferred the processed embryos to 50 ml tubes, drained the excess water, added the boiled-to-turbidity heat-fix solution (28% 500 µl NaCl + 5% 200 µl TritonX100 + 19.3 ml water), incubated for ~20 sec with thorough mixing, and then immediately added water to cool down the mixture. Then, we transferred the fixed embryos to 2 ml microfuge tubes, sequentially added n-heptane and methanol in equal proportions, vigorously shook for ~30 sec to remove the vitelline membrane, and washed 3x with methanol. We started the staining immediately after (though long-term methanol storage is possible). The embryos were rehydrated with PBT (0.01% solution of Tween20 in PBS, 3 washes), stained with 1:1000 DRAQ5 for 1-hour, washed 6x with PBT, re-suspended in 3:1 glycerol:PBS solution, and mounted between 2 rectangular coverslips, with spacing optimized to immobilize the embryos w/o deforming them. Then we imaged the embryos through both coverslips and the two halves stitched in image analysis (see below). Images acquired on Leica SP8 DMi8 inverted confocal microscope using 20x/0.75 NA glycerol objective, 1.15x digital zoom, 1 µm z-step, and 0.247 µm/pixel x-y resolution.

For **half-embryo imaging of nuclei and F-actin**, we used formaldehyde fixation and stained the embryos with DAPI and Phallacidin. Briefly, we transferred the processed embryos to a 20 ml vial containing the fixation mix (3500 µl PBS + 37% 1000 µl formaldehyde + 4500 µl n-Heptane). The vial was shaken on a platform shaker for 25 minutes at 200 rpm. Then, we transferred the fixed embryos to 2 ml microfuge tubes, sequentially added n-heptane and 90% ethanol in equal proportions, vigorously shook for ~30 sec to remove the vitelline membrane, and washed 3x with 90% ethanol. We started the staining immediately after (though storage in 90% ethanol possible for ~1 week). The embryos were rehydrated with PBT (3 washes), stained with 1:1000 DAPI for 1 hour, 1:500 Phallacidin for 1-hour, washed 4x with PBT, re-suspended in 3:1 glycerol:PBS solution, and mounted between 2 rectangular coverslips for imaging. Images were acquired on Leica SP8 DMi8 inverted confocal microscope using 20x/0.75 NA glycerol objective, 1.2x digital zoom, 1 µm z-step, and 0.473 µm/pixel x-y resolution.

Image analysis and statistics: All image analysis was performed in FIJI. We conducted 11 types of analyses in live or fixed embryos: 1) Germband extension (live), 2) developmental landmarks analysis (live) 3) whole embryo reconstruction of nuclear distribution (fixed), 4) nuclear expulsions (live), 5) nuclear movements (live), 6) nuclear cycle asynchrony (live), 7) nuclear crowding (live), 8) fraction of mitotic failures (live), 9) SLBP nuclear localization (live), 10) F-actin and microtubule distribution (live), and 11) comparing the distribution of nuclei and F-actin (fixed). All the statistical analysis were performed in Prism (GraphPad), including curve fitting (see below, nuclear crowding, SLBP localization, and F-actin and microtubule distribution), non-parametric tests (non-parametric Kruskal-Wallis test with Dunn's uncorrected test for multiple comparisons), and estimating the 95% confidence intervals (CI) on the medians. The straight-line fits for analysis of nuclear movements were done using the curve fitting functions in ImageJ macro language.

For **Germband extension** (GBE), we measured the distance of the anterior end of the posterior mid-gut (PMG) from the embryo posterior end. The GBE was calculated as,

$$percent\ GBE = \frac{distance\ of\ PMG\ from\ posterior * 100}{length\ of\ the\ embryo}$$

Plotting percent GBE against time shows the fast and slow phases of GBE.

For **developmental landmarks analysis,** we noted down the time at which following developmental stages occurred. For *40% GBE*, we noted down the time when the anterior tip of the PMG was closest to 40% embryo

length from the posterior pole. For *Germband retraction begins* and *ends*, we noted down the time when the
anterior tip of the elongated ectoderm on the dorsal side first moved towards the posterior and then
stopped. For *head involution*, we noted down the time when the ectoderm meets in the anterior.

For **whole embryo reconstruction of nuclear distribution**, we used a custom-built image analysis pipeline to
fuse the two views of the embryo, and visualize subcortical nuclei; either isolated or in clusters. We refer to
each view as "half embryo", even though each spans about 70-75% of the embryo width, providing a
substantial overlap. The pipeline comprises, as outlined in the following, elements in FIJI for processing the
raw images, stitching the two halves, rendering a dorsal-ventral view, segmenting the embryo volume, and
then color-coding the nuclei by distance from the segmented embryo surface. We first processed the raw
images to improve visualization of the nuclei ('Minimum 3D' filter, radii x=y=1, z=3; then 'Median 3D' filter,
radii x=y=z=1). Then, we homogenized the brightness/contrast between the two half embryos. We then
manually aligned the halves along their AP axis and used the BigWarp[71] within BigDataViewer[72] to register
them using manually selected interest points (subcortical- or yolk-nuclei) visible in both halves. We tightly
cropped the output images to reduce the computational load in the steps to follow. Then, we used
BigStitcher[73] to further align the embryo, using the phase correlation method, to produce the "fused
embryo". In the case of many embryos, additional manual transforms were necessary to obtain an acceptable
fused embryo. The fused embryos were rotated and resliced to produce dorsal-ventral views. We then used
thresholding on nuclear signal to segment embryo volume, generating a binary embryo mask, and used '3D
Distance Map' (from '3DSuite' in ImageJ[74]) to compute the Euclidian distances from embryo surface for each
voxel inside the embryo. Voxels with distance <18 μm were classified as cortical, 18-45 μm as subcortical,
and >45μm as yolk region (Figure 2B); these distances were determined empirically, using measurements
from various fused embryos. The regions were then color-coded and presented as maximum projections and
cross-sections in Figure 2. For embryos with a blastoderm hole, we further localized the hole in the embryo
coordinates, and mapped these locations from multiple embryos onto a model embryo surface. The
maximum intensity projections of the dorsal and ventral halves of the model embryo show that the
blastoderm holes predominantly occur in the central region of the embryo.

For **nuclear expulsions**, we tracked expelled nuclei back in time to identify the mitosis when they were born
and to categorize the expulsion as isolated (Figure 4A, B) or pairwise (Figure 4E). A nucleus was defined as
expelled if its apical tip lay ≥12 μm away from the embryo surface. To simplify the identification of expelled
nuclei, we color-coded the Histone-RFP signal as a function of distance from embryo surface. This is similar
to above analyses, except we use the 'LocalZPojector' plugin[75] in FIJI, with embryo surface inferred from the
co-imaged Jupiter-GFP signal. After color-coding the nuclei, we manually identify and count the number of
pairwise or isolated nuclear expulsions.

For **nuclear movements**, we manually tracked the entire lineage for select few nuclei from the mother
nucleus in late interphase 10 to all 8 great-granddaughter nuclei in late interphase 13. Nuclei from a single
lineage were treated as a single "group", yielding a group center at each time point. We chose nuclei along
one continuous row at interphase 10 and only chose those groups whose entire progeny (all 8 great-
granddaughters) could be tracked, yielding ≥13 groups per embryo. For each group, we fitted a straight line
to the time series of the group-center coordinates (dashed lines in Figure 5B) to capture the global drift, and
then computed, at every time point, the distance of the group-center to this line. An average of these
distances during a specific period was used to estimate of group-center deviations over that period. Since
the groups move strongly from anaphase through the first few minutes of the subsequent interphase, we
analyzed "extensive-movement" and "low-movement" periods separately. Extensive movements
consistently began ≥1 timepoint (30 s) before the first telophase appeared in any tracked groups and largely
subsided within 3 minutes after all groups completed telophase. Thus, we defined the extensive-movement

window as 30 s before the first telophase nuclei to 3 min after the last telophase nuclei. With these
objectively defined windows, average group-center deviations were straightforward to compute. Then, at
29°C, we further followed the progeny of the 8 great-granddaughter nuclei to identify those nuclear groups
that have at least one mitotic failure during mitosis 13, leading to pair-wise nuclear expulsions in interphase
14.

For **nuclear cycle asynchrony**, we marked key time points in the timelapse recordings to follow nuclear cycle
progression in the anterior-, middle-, and posterior-5^{th} of the field of view. We focused on the transitions
from a telophase 13 to interphase 14. If these timepoints are 'T_a' (anterior), 'T_p' (posterior), and 'T_m' (middle),
the 'asynchrony' can be calculated as,

866
$$asynchrony = T_m - \left(\frac{T_a + T_p}{2}\right)$$

Thus, we assume that in a fully synchronous embryo T_m equals the average of T_a and T_p; deviation from this
expected value quantifies asynchrony.

For **nuclear crowding**, we analyzed nuclear distribution at the end of interphases, when the distribution is
stable and crowding is maximal, if any. We first segmented the nuclei using 'Find Maxima' to detect local
intensity peaks, generating an image of maxima. These maxima were then manually curated: we removed
the maxima not corresponding to nuclei, add missing ones, and corrected the maxima that were clearly off-
center. Then we used Voronoi tessellation to mark territories (pseudo-cells) around each nucleus. We
excluded territories that (1) touched the edge of the image, (2) had aberrant shapes (aspect ratio>2), or (3)
contained pixels with <5 units of intensity (typically poorly defined pseudo-cells at the edge of segmentation).
Each nucleus was thus associated with a pseudo-cell area ('Area', μm^2) and the x-coordinate of the pseudo-
cell center ('X', μm) with x-axis aligned the embryonic AP axis. We then fitted a parabola to the Area-X
relationship,

879
$$Area = aX^2 + bX + c$$

Greater nuclear crowding corresponds to stronger curvature (compare Figure 6I-L); therefore, the fit
parameter 'a', which dictates the curvature, was used a quantitative measure of crowding.

For **fraction of mitotic failures**, we tracked expelled nuclei back in time (see above nuclear expulsion), but
additionally recorded their positions, to map where mitotic failures occurred within the field-of-view. In
parallel, we generated regions of (1) maximum nuclear crowding, (2) maximum delay in nuclear divisions,
and (3) their combination. To construct underlined crowded regions, we drew on previous studies showing that syncytial
divisions arrest when a threshold nuclear content is reached at ~70% of the normal cellularization nuclear
density[26–29]. A previous study reported the nuclear domain (pseudo-cell) area to be 28.8 μm^2 at
cellularization[76], which we corroborated (data not shown). For simplicity, we assumed 6000 nuclei at
interphase 14, each with an area of 29 μm^2. At 70% density, this corresponds to 4200 nuclei with area 41.43
μm^2, approximated as 42 μm^2. Thus, we estimated that divisions should arrest when pseudo-cell area falls
below ~42 μm^2. At 29°C, many nuclei in the central regions had areas <42 μm^2 in interphase 13, so we used
this criterion to define regions likely to have reached the 70% density one division earlier, locally. At normal
temperatures (22, 25, and 28°C), nuclei rarely had areas <42 μm^2; therefore, we defined crowded regions as
nuclei in the lower quartile of the pseudo-cell areas. This construction is a practical rationalization and not a
mechanistic statement about how the effect of nuclear crowding might manifest. Then, to define delayed
region (maximally delayed nuclear divisions), we marked the areas containing the last ~20% of the nuclei to
complete nuclear cycle 13. This empirically chosen fraction yielded regions similar in size as the crowded
regions. The rationale, based on previous work[25], is that division control is coordinated at the embryo scale,

and shared cytoplasm implies that locally delayed cycles will conflict with the embryo-wide biochemical status. Finally, we obtained the combined region using the logical OR operator of crowded and delayed regions, to test the combinatorial effects of crowding and asynchrony. In the end, for each embryo we computed the fraction of mitotic failures in each region as,

$$fraction\ of\ mitotic\ failures = \frac{number\ of\ mitotic\ failures\ in\ a\ specific\ region}{total\ number\ of\ mitotic\ failures}$$

As we can imagine, the fraction of mitotic failures will also depend on the size of the specific region, as compared to the total field of view. As such, in case of both crowded and delayed regions, we empirically determined criteria for their construction, such that the size of these regions will be comparable: ~20% of the field of view. In case of combined regions, however, the size can vary from ~20% (crowded and delayed regions show complete overlap) to ~40% (crowded and delayed regions show no overlap).

There was a marginal increase in embryo lethality at 25°C in the controls for the overexpression experiments, which presumably happens due to the presence of additional transgenic insertions (UAS and Gal4). Due to this reason, to err on the side of caution, we chose to perform the imaging at 22°C, specifically for the Fraction of mitotic failures analysis. As such we don't expect any biologically significant difference between results at 22°C and 25°C terms of in embryo lethality, blastoderm holes, or other defects.

For **SLBP nuclear localization**, we tracked nuclei from late interphase 13 till early interphase 14 in embryos expressing GFP-tagged SLBP and Histone-RFP, developing at 25°C or 29°C. To measure the nuclear SLBP intensity, we drew ROIs (3 μm diameter) in the center of the nuclei in RFP channel, applying these ROIs to GFP channel. The same procedure was applied to embryos expressing GFP-tagged SLBP.S118A or NLS, along with Histone-RFP, to compare localization kinetics. After identifying the two components of SLBP loss during interphase 14 in nuclei with mitotic failure during mitosis 13, we fitted a straight line to nuclear intensities between 0 and 4 minutes of interphase 14.

For **F-actin and microtubule distribution**, we tracked F-actin caps and microtubule spindles around the same nuclei during mitosis 13 (late prophase to end of metaphase) in embryos developing at 25°C or 29°C, and followed the divisions to classify them as normal or mitotic failures. For F-actin caps, we manually outlined the cap and fitted an ellipse to this contour to obtain cross-sectional area, aspect ratio, and the orientation of the major axis. For microtubule spindles, we manually drew a line between spindle poles to measure the spindle length and the spindle-axis orientation. These measurements were used to plot F-actin cap area and aspect ratio at metaphase, spindle length at metaphase, and the relative angle between cap long axis and spindle axis. Cumulative spindle rotation was computed as the sum of absolute angular changes between consecutive time points. The rate of spindle-length change was estimated as a linear-fit to spindle length over time across mitosis.

For **comparing the distribution of nuclei and F-actin**, we used a pipeline similar to that for 'whole embryo reconstruction of nuclear distribution', but without fusing two half-embryos and starting directly from embryo-volume segmentation using the F-actin staining. A 3D distance map was then computed and nuclei classified into cortical, subcortical, or yolk based on their distance from the embryo surface. These nuclei were color-coded and presented as sum projections in Figure S6 (except for yolk region).

Analysis of thermoclinal genes: We sought genes showing variation across natural populations of *Drosophila melanogaster* along large-scale climatic gradients. At continental scale, we reasoned that climatic variation imposes selection gradient with distance from equator, generating geographic genomic variation, i.e., a thermocline, across a temperature range of at least 15°C. Previous studies reported such variation at different scales (whole genome, Quantitative Trait Loci, or individual genes/proteins, see Table S1 for the list

of studies). To be able to aggregate over these scales we chose "genes" as a level of abstraction. Consequently, we focus our analysis to genes that show consistent clinality across data sets, rather than to specific allelic change (coding vs. non-coding, position in the protein, etc.) that are involved. We thus extracted only genes reported to vary across geographically distant populations, and grouped studies by continent (North America, Europe, or Australasia), assuming these continent-scale gradients arose independently. The resulting "thermoclinal datasets" were then used to assign a score of 0 to 3 for a gene, counting the number of appearances in individual datasets.

To assemble genes relevant for spindle dynamics, we combined the genes associated with Gene Ontology (GO) terms cell polarity, spindle localization, and spindle organization; none of the individual GO terms alone capture all six candidate genes tested in this study. We queried the thermoclinal datasets with this combined list of genes related to spindle dynamics, assigning each gene a 0 to 3 score as above. For comparison, we applied the same procedure to genes associated with GO term 'metabolic process'. See also, Tables S1 and S2.

Figure and table legends:

Figure 1: Embryonic lethality and developmental defects in embryos exposed to elevated temperature.

(A, B) To assess developmental robustness, we exposed embryos to various temperature regimes and estimated the embryo lethality. (A) Schematic showing the temperature regimes in the embryo survival experiment. Here we have three conditions depending on exposure to normal or elevated temperature regimen in the 3-hour time windows as shown, normal-normal [NN], elevated-normal [EN], and normal-elevated [NE]. In all cases, the first time-window also includes 1 hour of embryo collection. Embryonic development beyond 6 hours continued at normal temperature. (B) The scatter plots show the distribution of the percentage of embryos that don't hatch into larvae. Each dot represents data from a sample with ≥26 embryos; n= 12 (NN), 10 (EN), 10 (NE) samples.

(C-E) Montages show embryos, expressing Histone-RFP, from early gastrulation till late gastrulation, in embryos at 25°C. Till late cellularization stage, these embryos were exposed to either 25°C, i.e., NN temperature regimen (C), or 29°C, i.e., EN temperature regimen, showing mild (D), or severe (E) defect. White dashed lines, dorsal midline drawn based on the egg shape; red dashed lines, the embryonic midline in the dorsal posterior ectoderm; orange dashed lines, anterior end of the posterior mid-gut (extent of GBE).

(F) Plot show extent of GBE: hollow markers, embryos with gastrulation defects, which are excluded from further analysis of developmental landmarks. (G) The analysis of Developmental landmarks shows the time at which various events can be seen in the embryos. Time points from individual embryos are plotted at the same y-value.

Scale bar, 50 μm. B show scatters with median, with whiskers showing 95% CI. Non-parametric Kruskal-Wallis test with Dunn's uncorrected test for multiple comparisons: *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$; ****, $p<0.0001$, non-significant comparisons are not shown.

Figure 2: Nuclear loss lacks left-right symmetry in embryos developing at elevated temperature.

(A) Pie charts show occurrence of blastoderm holes during cellularization in fixed embryos at normal temperature (n=68) or elevated temperature (n=77): blastoderm hole, a cluster of >10 subcortical nuclei with a corresponding discontinuity in the cortical layer.

(B) Schematic shows various layers of the embryo volume: green, cortical nuclei, <18 μm from embryo surface; magenta, sub-cortical nuclei, >18 μm and <45 μm from embryo surface; yellow, yolk nuclei (vitellophages), >45 μm from embryo surface.

(C-H) Distribution of nuclei in embryos during mid-cellularization (interphase 14), in fixed embryos that were
developing at normal temperature (C, F) or elevated temperature that shows either a mild defect (D, G) or a
severe defect (E, H). Top/bottom panels, maximum intensity projections in dorsal/ventral halves of the
embryos; white/grey dashed lines, midlines drawn based on the egg shape. The voxels are color-coded as in
B. Embryo orientation is consistently presented in all panels and is indicated in C. (C-E) Cortical and
subcortical nuclei are shown. Embryos at normal temperature (C) show rarely any gaps, while those at
elevated temperature occasionally show small gaps (D) or blastoderm holes (E), indicating discontinuities in
the distribution of cortical nuclei, corresponding also with the locations of subcortical nuclei. (F-H) The
corresponding subcortical nuclei are shown separately for improved visibility, along with yolk nuclei.

(I-N) Cross-sectional views long the orange and cyan dashed lines in C-H. Embryo orientation is consistently
presented in all panels and is indicated in I.

(O, P) Blastoderm holes from 11 embryos at elevated temperature were mapped onto a model embryo
surface (one blastoderm hole per embryo). Images show projections in dorsal (O) and ventral (P) halves of
the model embryo. *, blastoderm hole from embryo in E/H.

Scale bars: F, 50 μm; I, 25 μm.

Figure 3: Cortical nuclei are expelled in embryos at elevated temperature.

(A-E) Montages showing live embryos, expressing Histone-RFP, from early interphase 10 till late interphase
14, developing either at 22°C (A), 25°C (B), 28°C (C), or at 29°C showing mild (D), or severe (E) defect. The
nuclei were labelled as a function of their distance from the embryo surface. Green, cortical nuclei; magenta,
sub-cortical nuclei that are lost (see also schematic in Figure 4B). At elevated temperature, a lot more nuclei
are expelled, and as a result, we can see gaps between cortical nuclei, coinciding with subcortical nuclei:
arrowheads, small gaps; arrows, blastoderm hole.

Scale bar, 25 μm.

Figure 4: Nuclear loss as isolated or paired expulsions at elevated temperature.

Following isolated nuclear expulsions back in time, we found them to stem from both out-of-plane and in-
plane divisions. (A) Montages exemplify out-of-plane division during mitosis 10 at 25°C. Histone-RFP marks
nuclei (green), Jupiter-GFP marks microtubules (magenta). Arrowheads, in-plane divisions; arrows, out-of-
plane division; white lines, embryo surface. Each subsequent frame is 1 minute apart. (B) Montage
exemplifies in-plane divisions at 25°C, from telophase 12 to interphase 13. Histone-RFP marks nuclei: cortical
nuclei, green; subcortical nuclei, magenta; see schematic. The arrowheads/arrows track the daughter nucleus
that will be expelled/cortically retained. Inset, time since telophase 12. (C, D) Scatter plots show number of
out-of-plane divisions during mitosis 10 (C) and number of isolated nuclear expulsions during interphase 14.

To assess the pairwise loss of nuclei, we counted mitotic failures. (E) Montage exemplifies mitotic failure at
29°C, from late interphase 12 to late interphase 13. Histone-RFP marks nuclei: cortical nuclei, green;
subcortical (lost) nuclei, magenta; see the schematic in B. White arrows, normal nuclear division; white
arrowheads, mitotic failure; cyan arrowheads, lagging chromosomes. Inset, time since telophase 12. (F)
Scatter plot shows number of mitotic failures during interphase 14. (G) The schematic illustrates the
phenomenon of mitotic failures at elevated temperature.

Scale bars: A, 10 μm; B and E, 5 μm. C, D, and F show scatters with median, with whiskers showing 95% CI.
Large hollow markers, embryos with blastoderm holes; small hollow markers, embryos with small gaps in the
blastoderm. n= 10 (22°C), 7 (25°C), 9 (28°C), and 10 (29°C) embryos. Non-parametric Kruskal-Wallis test with

Dunn's uncorrected test for multiple comparisons: *, p<0.05; **, p<0.01; non-significant comparisons are not
shown.

Figure 5: Nuclear movement parallel to the embryo surface is amplified at elevated temperature.

(A-C) An example of nuclear movement tracking. Snapshots from embryo at 29°C, with Histone-RFP marking
nuclei, showing the mother nuclei at the end of interphase 10 (A) and the corresponding 8 great-
granddaughter nuclei, each, at the end of interphase 13 (C), connected via the group-center trajectories (B).
Cyan and yellow, neighbouring nuclear groups; thickening solid lines, trajectories of group-centers with
increasing number of nuclei due to divisions; dotted lines, linear fits to group-center trajectories; shaded
region, periods of greatest movement (see methods).

(D-I) To quantify the drift in the nuclear group centers, we compute the slope of the linear fits (dotted lines
in B). Plots show the distribution of embryo medians, for the drift along the AP axis (D) and DV axis (E). To
quantify the movements in nuclear group centers during interphase-metaphase vs. anaphase-telophase
(shaded vs. unshaded region in B), we compute the average of the deviations from the linear fits. Plots show
the distribution of embryo medians, for the fluctuations along the AP axis (F, H) and DV axis (G, I), during
interphase-metaphase (F, G) and anaphase-telophase (H, I). Large hollow markers, embryos with blastoderm
holes; small hollow markers, embryos with small gaps in the blastoderm. n= 10 (22°C), 7 (25°C), 9 (28°C), and
10 (29°C) embryos.

(J) To test whether nuclear groups with highest movement also correspond to mitotic failures, we plot the
deviations along AP-axis, now showing all of the tracked nuclear groups at 29°C. Hollow markers, nuclear
groups with at least one subsequent mitotic failure; underlined, embryo with blastoderm holes; insets,
number of tracked nuclear groups.

Scale bar, 25 μm. D to J show scatters with median, with whiskers showing 95% CI. Non-parametric Kruskal-
Wallis test with Dunn's uncorrected test for multiple comparisons: *, p<0.05; **, p<0.01; ***, p<0.001; non-
significant comparisons are not shown, statistical tests not performed on data in J.

Figure 6: Nuclear crowding and Nuclear cycle asynchrony increase at elevated temperature.

To assess the crowding of nuclei, we segmented the nuclei and generated pseudo-cells. (A-D) Snapshots from
embryos at the end of interphase 13 at indicated temperatures: Histone-RFP marks the nuclei; outline, region
of the image considered for quantifications. (E-H) The pseudo-cells for the corresponding embryos in A-D.
Color indicates pseudo-cell area in $μm^2$, LUT at the bottom. (I-L) Plots show distribution of pseudo-cell areas
as a function of AP position embryos in A-D. Dotted line, quadratic fit. (M) Scatter plots show distribution of
the coefficient of the quadratic term, with a larger quadratic coefficient indicating greater crowding. (N) The
schematic illustrates the crowding effect at elevated temperature: size of the black arrows, gradation in
nuclear density.

To assess the asynchrony in nuclear cycles, we identified the phases of the nuclear cycle along the AP axis.
(O, P) Image shows the entire anterior-posterior extent of the field of view, from embryos at 25°C (O) and
29°C (P). Histone-RFP marks the nuclei: green, cortical nuclei; magenta, subcortical (lost) nuclei (see also
schematic in Figure 4B). Arrows show the direction in which the wave of the nuclear division cycle will
progress. At 25°C, we see a single wave from the anterior, while at 29°C we see an anterior- and a posterior-
wave. (Q) Scatter plots show asynchrony in initiation of interphase 14. (R) The schematic illustrates the
progressive increase in asynchrony at elevated temperature.

Scale bars, 20 μm. M and Q shows scatters with median, with whiskers showing 95% CI. Large hollow markers,
embryos with blastoderm holes; small hollow markers, embryos with small gaps in the blastoderm. n= 10

(22°C), 7 (25°C), 9 (28°C), and 10 (29°C) embryos. Non-parametric Kruskal-Wallis test with Dunn's uncorrected
test for multiple comparisons: *, p<0.05; **, p<0.01; ***, p<0.001; non-significant comparisons are not
shown.

Figure 7: Nuclear crowding and nuclear cycle asynchrony combinatorially enhance mitotic failures.

(A, B) Snapshots from embryos at 29°C, at the end of interphase 13 (A) and 14 min afterwards (B). Histone-
RFP marks nuclei; green outline, region of the image with nuclear crowding; magenta outline, region
containing the most delayed nuclei, i.e., those that are most lagging in nuclear division cycles.

(C-E) The locations of lost pairs of nuclei during interphase 14 (mitotic failures during mitosis 13) are marked
as white dots. Then, various regions are defined in the embryo to illustrate the fraction of mitotic failures
occurring in that region. These regions are composed based on, the crowded region (C), the delayed region
(D), and the combined region (E).

(F-K) Scatter plots show distribution of the fraction of mitotic failures occurring in the crowded (F), delayed
(G), and combined (H) regions of the embryos, and the distribution of the area fractions in crowded (I),
delayed (J), and combined (K) regions of the embryos. In the case of the area fraction of the combined regions
(K), the increase in the overlap of crowded and delayed regions reduces the overall size of the combined
regions.

Scale bars, 20 μm. F-K show scatters with median, with whiskers showing 95% CI. Large hollow markers,
embryos with blastoderm holes; small hollow markers, embryos with small gaps in the blastoderm. n= 10
(22°C), 7 (25°C), 9 (28°C), and 10 (29°C) embryos. Non-parametric Kruskal-Wallis test with Dunn's uncorrected
test for multiple comparisons: *, p<0.05; **, p<0.01; ***, p<0.001; non-significant comparisons are not
shown, significance not tested for I and J (see methods).

Figure 8: Mitotic failures lead to DNA damage response.

(A) Schematic shows how the mislocalization of the Stem-Loop Binding Protein (SLBP) can be used as a
reporter for DNA damage. During interphase, SLBP normally localizes to the nuclei, while it is excluded in
nuclei with DNA damage.

(B) The nuclear localization of SLBP (green) allows us to distinguish nuclei with DNA damage (dashed circles)
vs. normal ones (circles). Following these nuclei over a mitosis shows that nuclei with DNA damage are
mitotically arrested, while the normal nuclei divide, giving rise to two daughter nuclei (smaller circles). One
of the nuclei with DNA damage is lost to embryo interior by the last timepoint. Inset: time since mitosis.

(C-E) Montages show representative embryos developing at 29°C from telophase 13 till early interphase 14,
in embryos expressing SLBP-GFP (C), SLBP.S118A-GFP (D), and NLS-GFP (E). Arrows, chromosome segregation
defect; asterisks, pair of defective daughter nuclei; dashed circles, reduction in GFP-signal in the nuclei with
mitotic failure, still localized cortically; inset, time since beginning of interphase 14.

(F-K) Quantifications showing the kinetics of nuclear localization of SLBP-GFP (F), SLBP.S118A-GFP (G), and
NLS-GFP (H) in control divisions in embryos at 25°C, control division in embryos at 29°C, and mitotic failures
in embryos at 29°C. Slope of a straight-line fit between timepoints 0 to 4 min in F-H is plotted respectively (I-
K). Insets, the number of embryos (N) used for pooling the number of nuclear divisions (n).

Scale bars; B, 10 μm; C, 5 μm. F-H plots median values over time, with shaded regions showing 95% CI. I-K
show scatters with median, with whiskers showing 95% CI. Non-parametric Kruskal-Wallis test with Dunn's
uncorrected test for multiple comparisons: ***, p<0.001; non-significant comparisons are not shown.

Figure 9: Weakened interaction between F-actin and microtubules precedes mitotic failures.

(A) Schematic depicts dramatic reorganisation of F-actin, microtubules, and chromatin as the nuclei transition from prophase to late metaphase (left). The images are then manually annotated (center) to facilitate automated quantitative analysis of F-actin caps and microtubule spindles (right).

(B-E) Montages from representative control nuclear divisions (B, C) and mitotic failures (D, E) from embryos developing at 25°C (B, D) and 29°C (C, E), during mitosis 13. Left panels show montage of UtrABD- and Jupiter-GFP signal: dashed contour, manually outlined F-actin cap; white dots, manually marked spindle poles. Right panels show montage of corresponding Histone-RFP signal: dashed green line, long axis of F-actin cap; solid cyan line, spindle position; insets, time till end of metaphase.

(F-K) Quantifications showing F-actin cap size (F) and aspect ratio (G) at metaphase, microtubule spindle length at metaphase (H), spindle extension rate (I) and cumulative rotation (J) during mitosis, and the angle mismatch between spindle axis and cap long-axis (K). n= 27 (25°C control division), 27 (29°C control division), 10 (25°C mitotic failure), 15 (29°C mitotic failure) nuclei, pooled from N= 9 (25°C) and 9 (29°C) embryos.

Scale bar, 5 μm. F-K show scatters with median, with whiskers showing 95% CI. Non-parametric Kruskal-Wallis test with Dunn's uncorrected test for multiple comparisons: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$; non-significant comparisons are not shown.

Figure 10: Rescue of embryonic lethality at elevated temperature through overexpressions, and clinal genes as "natural" candidates.

(A) Testing the rescue of embryonic lethality and the fraction of mitotic failures with overexpression of *αCat* or *sgg*. Scatter plot shows distribution of percent lethality in embryos from various genotypes, at normal (25°C) vs elevated temperature (29°C), calculated based on the fraction of embryos that don't hatch into larvae. Each dot represents data from a sample with ≥18 embryos: control, n= 10 (25°C), 11 (29°C) samples; αCat-OE, n= 10 (25°C), 12 (29°C) samples; sgg-OE, n= 8 (25°C), 7 (29°C) samples.

(B-C) Scatter plots show distribution of the fraction of mitotic failures in crowded regions (B), and in combined regions (C), in the embryos at normal (22°C) vs elevated temperature (29°C). Each dot represents data from one embryo: control, n= 8 (22°C), 9 (29°C) embryos; αCat-OE, n= 8 (22°C), 8 (29°C) embryos; sgg-OE, n= 7 (22°C), 7 (29°C) embryos.

(D, E) Analysis of "natural" candidate genes that vary along thermoclines. (D) Venn diagram illustrates how we constructed a non-redundant list of genes associated with 'spindle dynamics' through the union of genes associated with GO terms, cell polarity (n=203 genes), spindle localization (n=37 genes), and spindle organization (n=138 genes). (E) Pie charts show clinal distribution scores for various genes associated with 'spindle dynamics' as constructed in D (n=320 genes), or with GO term 'metabolic process' (n=6612 genes). ~1% of genes associated with 'spindle dynamics', while ~0.2% of the genes associated with 'metabolic process', are clinal in all three continental datasets. See also, Tables S1 and S2.

A-C shows scatters with median, with whiskers showing 95% CI. In B and C, large hollow markers, embryos with blastoderm holes; small hollow markers, embryos with small gaps in the blastoderm. Non-parametric Kruskal-Wallis test with Dunn's uncorrected test for multiple comparisons: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$; non-significant comparisons are not shown.

Figure S1: Blastoderm holes appear in a fraction of embryos at elevated temperature.

The figure shows the snapshots from live embryos, expressing Histone-RFP, from early interphase 14, in embryos at elevated temperature (29°C). The nuclei are labelled as a function of their distance from the embryo surface. Here, green indicates cortical, and magenta indicates sub-cortical nuclei that are expelled (see also schematic in Figure 4B), while arrows/arrowheads indicate large/small blastoderm holes. For

embryos with large blastoderm holes (embryos 05, and 08), we see many nuclei in magenta. In contrast, in the case of embryos with small blastoderm holes (embryos 04, and 09), we can see only a few nuclei in magenta per hole. We would argue that development in *Drosophila* is robust enough to buffer the defects due to the small gaps in the blastoderm, but fails at 29°C when large holes appear.

Scale bar, 25 μm.

Figure S2: Defective nuclei take different amounts of time to be expelled from the embryo cortex.

(A, B) Montages from live embryos, with a few representative nuclei tracked from late interphase 12 to interphase 13. Top panels, Histone-RFP signal marking nuclei; bottom panels, Jupiter-GFP signal marking microtubules. Nuclei tagged in green/magenta are the ones that arise from mitotic defects during mitosis 10/11. All of these are expelled during interphase 13. Notice the lack of spindle formation around the tagged nuclei. Also, note that the nuclei expelled by the 5th timepoint are not shown in the 6th timepoint. (A) An embryo developing at normal temperature (22°C), where 1 magenta and 1 green nuclei are followed. (B) An embryo developing at elevated temperature (29°C), where 6 magenta and 3 green nuclei are followed.

Scale bars, 20 μm. Insets, time in min:sec.

Figure S3: Nuclear movement along surface-yolk axis is amplified at elevated temperature.

Cross-sections from live embryos, expressing Histone-RFP, from late interphase 13 till early interphase 14, in embryos developing either at normal temperature (22°C, A) or at elevated temperature (29°C, B). Colored lines, basal tip of nuclei in the first frame; insets, time since initiation of anaphase 13 at the left/right extreme of the images (which is also the first time we see the shift from baseline in embryo at elevated temperature); arrowheads, local peak of the nuclear shift from the baseline; arrow, expelled nuclei.

Scale bars, 25 μm. Cross-sections are in the lateral region of the embryos, anterior to the left.

Figure S4: Nuclear cycle asynchrony at elevated temperature during embryo development.

To assess the asynchrony in nuclear cycles, we identified the phases of the nuclear cycle along the AP axis. (A, B) Montages show the transition from nuclear cycle 13 to early interphase 14, from live embryos at 25°C (A), and at 29°C (B). Histone-RFP marks the nuclei: green, cortical nuclei; magenta, subcortical (lost) nuclei (see also schematic in Figure 4B); inset, time in min:sec. The anterior, central, and posterior 1/5th of the field of view are as indicated. Time 00:00 is when the anterior 1/5th of the embryo enters interphase 14. Boxes highlight entry into interphase 14 in that region.

Scale bars, 20 μm.

Figure S5: Embryonic lethality at elevated temperature in *baz* or *dlg* overexpressions.

(A) Testing the rescue of embryonic lethality and the fraction of mitotic failures with overexpression of *baz* and *dlg*. Scatter plot shows distribution of percent lethality in embryos from various genotypes, at normal (25°C) vs elevated temperature (29°C), calculated based on the fraction of embryos that don't hatch into larvae. Each dot represents data from a sample with ≥12 embryos; control, n= 9 (25°C), 9 (29°C) samples; baz-OE, n= 11 (25°C), 10 (29°C) samples; dlg-OE, n= 12 (25°C), 10 (29°C) samples.

(B, C) Scatter plots show distribution of the fraction of mitotic failures in crowded regions (B), and in combined regions (C), in the embryos at normal (22°C) vs elevated temperature (29°C). Each dot represents data from one embryo: control, n= 7 (22°C), 9 (29°C) embryos; baz-OE, n= 7 (22°C), 7 (29°C) embryos; dlg-OE, n= 6 (22°C), 7 (29°C) embryos.

Plots show scatters with median, whiskers show 95% CI. In B and C, large hollow markers, embryos with blastoderm holes; small hollow markers, embryos with small gaps in the blastoderm. Non-parametric Kruskal-Wallis test with Dunn's uncorrected test for multiple comparisons: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; non-significant comparisons are not shown.

Figure S6: Elevated temperature disrupts the association between nuclei and F-actin caps.

Distribution of nuclei and F-actin in embryos during interphase 13, in fixed embryos that were developing either at normal temperatures (A, C, D), or elevated temperature (B, E, F). (A, B) The images show SUM projections of the DAPI-stained nuclei (top image) and Phallacidin-stained F-actin (bottom image) of representative embryos. In the top images, the nuclei are labelled as a function of their distance from the embryo surface, where green indicates cortical, and magenta indicates sub-cortical nuclei (see also schematic in Figure 2B). The yolk nuclei are not shown. Anterior is on the left, dorsal up. (C-F) The top and bottom images show a zoomed-in view of the indicated regions in panels A and B. Arrowheads, expelled nuclei (C, E top panels) and corresponding gaps between the actin-caps (C, E bottom panels); arrows, gaps between neighbouring nuclei (D, F top panels) and corresponding gaps between the actin-caps at normal temperature (D, bottom panel) and "empty actin caps" at elevated temperature (F, bottom panel).

Scale bars, 25μm. The boxed regions are 60x60 μm.

Figure S7: *mud* or *insc* overexpression does not ameliorate embryo survival at elevated temperature.

Scatter plot shows the distribution of percent lethality in embryos from various genotypes, at normal (25°C) vs elevated temperature (29°C), calculated based on the fraction of embryos that don't hatch into larvae. Each dot represents data from a sample with ≥21 embryos; control, n= 10 (22°C), 11 (29°C) samples; mud-OE, n= 11 (22°C), 12 (29°C) samples; insc-OE, n= 7 (22°C), 10 (29°C) samples. As we can see in the plot, neither *mud* nor *insc* overexpression rescues the embryo lethality at elevated temperature, with *mud* overexpression increasing embryo lethality even at normal temperature (25°C).

The plot shows scatters with a median, while whiskers show 95% CI. Non-parametric Kruskal-Wallis test with Dunn's uncorrected test for multiple comparisons: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; non-significant differences are omitted.

Table S1: This table contains genes that are identified as being clinal in different continents. Columns A to C contain lists of studies, and under the studies is a list of genes that are identified in these studies as being clinal. Columns A, B, and C respectively list studies that have data from the continents of Australia and/or Australasia (AUS) [77–86], North and/or Central America (USA) [77,84,94–96,86–93], and Europe and/or North Africa (EU) [77,83,97,98]. We collect all the genes associated with 'spindle dynamics' (Figure 9D), to compile a list of potential candidate genes in Column F. For each gene in column F, we count the number of times that gene appears in columns A to C, counting a maximum of once per column, to respectively populate columns G to I. For each gene in column F, column J contains the sum of values in G, H, and I, to indicate the number of continents in which a gene is clinal. Each gene gets a score from 0 to 3, and we consider the gene to be a relevant candidate if it is found in at least 2 columns, i.e., with a score of 2 or 3.

We collect all the genes associated with the GO term 'metabolic process', to compile a control list of candidate genes in Column L. For each gene in column L, we count the number of times that gene appears in columns A to C, counting a maximum of once per column, to respectively populate columns M to O. For each gene in column L, column P contains the sum of values in M, N, and O, to indicate the number of continents in which a gene is clinal. Thus, each gene gets a score from 0 to 3.

Table S2: Column A lists candidates from Table S1 column F that have a score of 2 or higher, i.e., those that show significant selectable variation in natural populations. For these genes, we query FlyBase (http://flybase.org/batchdownload) for additional details like gene name (column B), gene symbol (column C), and the level of expression during the first 2 hours of embryo development (column D). The level of expression is based on published transcriptomics data[99], and would correspond to a maternal pool of mRNAs, which is the most relevant mRNA source for this study. The candidate genes clinal in all 3 continents are highlighted in green. Columns F-J present a table with counts of genes that are associated with various categories of genes in Table S1 columns F and L, and a further breakdown of how many of these genes are clinal in how many continents. The breakdown is also plotted as pie charts in Figure 10E.

Table S3: This table lists various figure panels (columns A and B), and the related parental genotypes, i.e., the genotypes of females (column C) and males (column D) present in the embryo collection cages for the experiment related to a figure panel.

Additional information

Data and Code Availability: Data and codes developed for data analysis are available upon request from corresponding authors.

Author contributions: G.K. and S.L. planned the project. G.K. did the experiments with help from P.A. for baz and dlg overexpressions, JJ.D.-L. for co-staining of nuclei and F-actin. G.K. did the image analyses and statistical analyses. G.K. and S.L. wrote the manuscript, and all authors gave comments on the manuscript.

Acknowledgments: We would like to thank Lázaro Centanin, Nicolas Gompel, Marianthi Karageorgi, and Russ Hodge for insightful comments on various drafts of the manuscript, and Lemke lab members for discussions throughout the project. We are grateful to Jochen Wittbrodt for his enthusiasm and his continuous and generous support. We would also like to thank Ingrid Lohmann, Anja Nagel and Kristen Panfilio for *Drosophila* culture-related support, and Thomas Lecuit, Jan Felix Evers, and Eric Lecuyer for sharing fly lines. We sincerely appreciate the efforts taken by FlyBase to maintain a curated database and Bloomington Drosophila Stock Center for providing transgenic fly lines. This work was funded by the DFG (LE 2787/3-1) to S.L., while G.K. was supported through an HFSP Long-term postdoctoral fellowship (LT000597/2019-L).

Competing interests: The authors declare no competing financial interests.

References:

1.   Wiens, J. J. Climate-Related Local Extinctions Are Already Widespread among Plant and Animal Species. *PLOS Biol.* **14**, e2001104 (2016).

2.   Edwards, M. J. Review: Hyperthermia and fever during pregnancy. *Birth Defects Res. A. Clin. Mol. Teratol.* **76**, 507–16 (2006).

3.   Miller, M. W. & Church, C. C. Arrhenius thermodynamics and birth defects: chemical teratogen synergy. Untested, testable, and projected relevance. *Birth Defects Res. C. Embryo Today* **99**, 50–60 (2013).

4.   Graham, J. M. Update on the gestational effects of maternal hyperthermia. *Birth defects Res.* **112**, 943–952 (2020).

5.   Cockroft, D. L. & New, D. A. T. Abnormalities induced in cultured rat embryos by hyperthermia. *Teratology* **17**, 277–283 (1978).

6.   Germain, M. -A, Webster, W. S. & Edwards, M. J. Hyperthermia as a teratogen: parameters determining hyperthermia-induced head defects in the rat. *Teratology* **31**, 265–72 (1985).

7.   Schirone, R. C. & Gross, L. Effect of temperature on early embryological development of the zebra fish, Brachydanio rerio. *J. Exp. Zool.* **169**, 43–52 (1968).

8. Dorrity, M. W. *et al.* Proteostasis governs differential temperature sensitivity across embryonic cell types. *Cell* **186**, 5015-5027.e12 (2023).

9. LILLIE, F. R. & KNOWLTON, F. P. ON THE EFFECT OF TEMPERATURE ON THE DEVELOPMENT OF ANIMALS. *Zool. Bull.* **1**, 179–193 (1897).

10. Crapse, J. *et al.* Evaluating the Arrhenius equation for developmental processes. *Mol. Syst. Biol.* **17**, e9895 (2021).

11. Kuntz, S. G. & Eisen, M. B. Drosophila embryogenesis scales uniformly across temperature in developmentally diverse species. *PLoS Genet.* **10**, e1004293 (2014).

12. Chong, J., Amourda, C. & Saunders, T. E. Temporal development of Drosophila embryos is highly robust across a wide temperature range. *Interface* **15**, (2018).

13. Welte, M. A., Tetrault, J. M., Dellavalle, R. P. & Lindquist, S. L. A new method for manipulating transgenes: engineering heat tolerance in a complex, multicellular organism. *Curr. Biol.* **3**, 842–853 (1993).

14. Foe, V. E. & Alberts, B. M. Studies of nuclear and cytoplasmic behaviour during the five mitotic cycles that precede gastrulation in Drosophila embryogenesis. *J. Cell Sci.* **61**, 31–70 (1983).

15. Xie, Y. & Blankenship, J. T. Differentially-dimensioned furrow formation by zygotic gene expression and the MBT. *PLoS Genet.* **14**, e1007174 (2018).

16. Mazumdar, A. & Mazumdar, M. How one becomes many: Blastoderm cellularization in Drosophila melanogaster. *BioEssays* **24**, 1012–1022 (2002).

17. Martin, A. C. The Physical Mechanisms of Drosophila Gastrulation: Mesoderm and Endoderm Invagination. *Genetics* **214**, 543–560 (2020).

18. Lecuit, T. & Yap, A. S. E-cadherin junctions as active mechanical integrators in tissue dynamics. *Nat. Cell Biol.* **17**, 533–539 (2015).

19. Foe, V. E. Mitotic domains reveal early commitment of cells in Drosophila embryos. *Trends Genet.* **5**, 322 (1989).

20. Niemuth, J. & Wolf, R. Developmental asynchrony caused by steep temperature gradients does not impair pattern formation in the wasp, Pimpla turionellae L. *Roux's Arch. Dev. Biol.* **204**, 444–452 (1995).

21. Lucchetta, E. M., Lee, J. H., Fu, L. A., Patel, N. H. & Ismagilov, R. F. Dynamics of Drosophila embryonic patterning network perturbed in space and time using microfluidics. *Nature* **434**, 1134–1138 (2005).

22. Deneke, V. E. *et al.* Self-Organized Nuclear Positioning Synchronizes the Cell Cycle in Drosophila Embryos. *Cell* **177**, 925-941.e17 (2019).

23. Kanesaki, T., Edwards, C. M., Schwarz, U. S. & Grosshans, J. Dynamic ordering of nuclei in syncytial embryos: a quantitative analysis of the role of cytoskeletal networks. *Integr. Biol. (Camb).* **3**, 1112–9 (2011).

24. Schmidt, A. & Grosshans, J. Dynamics of cortical domains in early Drosophila development. *J. Cell Sci.* **131**, (2018).

25. Deneke, V. E., Melbinger, A., Vergassola, M. & Di Talia, S. Waves of Cdk1 Activity in S Phase Synchronize the Cell Cycle in Drosophila Embryos. *Dev. Cell* **38**, 399–412 (2016).

26. Edgar, B. A., Kiehle, C. P. & Schubiger, G. Cell cycle control by the nucleo-cytoplasmic ratio in early Drosophila development. *Cell* **44**, 365–72 (1986).

27. Blythe, S. A. & Wieschaus, E. F. Zygotic genome activation triggers the DNA replication checkpoint at the midblastula transition. *Cell* **160**, 1169–1181 (2015).

28. Lu, X., Li, J. M., Elemento, O., Tavazoie, S. & Wieschaus, E. F. Coupling of zygotic transcription to mitotic control at the Drosophila mid-blastula transition. *Development* **136**, 2101–10 (2009).

29. Hayden, L. *et al.* Cullin-5 mutants reveal collective sensing of the nucleocytoplasmic ratio in Drosophila embryogenesis. *Curr. Biol.* **32**, 2084-2092.e4 (2022).

30. Hayden, L., Hur, W., Vergassola, M. & Di Talia, S. Manipulating the nature of embryonic mitotic

waves. *Curr. Biol.* **32**, 4989-4996.e3 (2022).

31.   Iampietro, C. *et al.* Developmentally Regulated Elimination of Damaged Nuclei Involves a Chk2-
Dependent Mechanism of mRNA Nuclear Retention. *Dev. Cell* **29**, 468–481 (2014).

32.   Karr, T. L. & Alberts, B. M. Organization of the cytoskeleton in early Drosophila embryos. *J. Cell Biol.*
102, 1494–1509 (1986).

33.   Bergstralh, D. T., Haack, T. & St Johnston, D. Epithelial polarity and spindle orientation: intersecting
pathways. *Philos. Trans. R. Soc. B Biol. Sci.* **368**, 20130291 (2013).

34.   Buckley, C. E. & St Johnston, D. Apical–basal polarity and the control of epithelial form and function.
Nat. Rev. Mol. Cell Biol. **23**, 559–577 (2022).

35.   Harris, T. J. C. & Peifer, M. Adherens junction-dependent and -independent steps in the
establishment of epithelial cell polarity in Drosophila. *J. Cell Biol.* **167**, 135–47 (2004).

36.   Cao, J., Crest, J., Fasulo, B. & Sullivan, W. Cortical actin dynamics facilitate early-stage centrosome
separation. *Curr. Biol.* **20**, 770–6 (2010).

37.   Rizzelli, F., Malabarba, M. G., Sigismund, S. & Mapelli, M. The crosstalk between microtubules, actin
and membranes shapes cell division. *Open Biol.* **10**, 190314 (2020).

38.   McCartney, B. M. *et al.* Drosophila APC2 and Armadillo participate in tethering mitotic spindles to
cortical actin. *Nat. Cell Biol.* **3**, 933–8 (2001).

39.   Rim, E. Y., Clevers, H. & Nusse, R. The Wnt Pathway: From Signaling Mechanisms to Synthetic
Modulators. *Annu. Rev. Biochem.* **91**, 571–598 (2022).

40.   Mapelli, M. & Gonzalez, C. On the inscrutable role of Inscuteable: Structural basis and functional
implications for the competitive binding of NuMA and Inscuteable to LGN. *Open Biol.* **2**, (2012).

41.   Ji, J.-Y., Haghnia, M., Trusty, C., Goldstein, L. S. B. & Schubiger, G. A Genetic Screen for Suppressors
and Enhancers of the Drosophila Cdk1-Cyclin B Identifies Maternal Factors That Regulate
Microtubule and Microfilament Stability. *Genetics* **162**, 1179–1195 (2002).

42.   Cadigan, K. M. & Nusse, R. Wnt signaling: a common theme in animal development. *Genes Dev.* **11**,
3286–3305 (1997).

43.   Pires-daSilva, A. & Sommer, R. J. The evolution of signalling pathways in animal development. *Nat.*
Rev. Genet. **4**, 39–49 (2003).

44.   Long, Y. C., Chin Tan, T. M., Takao, I. & Tang, B. L. The biochemistry and cell biology of aging:
Metabolic regulation through mitochondrial signaling. *Am. J. Physiol. - Endocrinol. Metab.* **306**,
(2014).

45.   Bitto, A., Wang, A. M., Bennett, C. F. & Kaeberlein, M. Biochemical genetic pathways that modulate
aging in multiple species. *Cold Spring Harb. Perspect. Med.* **5**, 1–25 (2015).

46.   Hale, R. & Strutt, D. Conservation of Planar Polarity Pathway Function Across the Animal Kingdom.
Annu. Rev. Genet. **49**, 529–551 (2015).

47.   Wright, B. A., Kvansakul, M., Schierwater, B. & Humbert, P. O. Cell polarity signalling at the birth of
multicellularity: What can we learn from the first animals. *Front. Cell Dev. Biol.* **10**, 1–19 (2022).

48.   Schlacht, A., Herman, E. K., Klute, M. J., Field, M. C. & Dacks, J. B. Missing Pieces of an Ancient
Puzzle: Evolution of the Eukaryotic Membrane-Trafficking System. *Cold Spring Harb. Perspect. Biol.*
6, a016048–a016048 (2014).

49.   Harashima, H., Dissmeyer, N. & Schnittger, A. Cell cycle control across the eukaryotic kingdom.
Trends Cell Biol. **23**, 345–356 (2013).

50.   Erickson, H. P. Evolution of the cytoskeleton. *BioEssays* **29**, 668–677 (2007).

51.   Wickstead, B. & Gull, K. The evolution of the cytoskeleton. *J. Cell Biol.* **194**, 513–525 (2011).

52.   Vazquez, J., Pauli, D. & Tissières, A. Transcriptional regulation in Drosophila during heat shock: a
nuclear run-on analysis. *Chromosoma* **102**, 233–48 (1993).

53.   Leemans, R. *et al.* Quantitative transcript imaging in normal and heat-shocked Drosophila embryos

by using high-density oligonucleotide arrays. *Proc. Natl. Acad. Sci.* **97**, 12138–43 (2000).

54.   Feder, M. E., Cartaño, N. V., Milos, L., Krebs, R. A. & Lindquist, S. L. Effect of engineering Hsp70 copy number on Hsp70 expression and tolerance of ecologically relevant heat shock in larvae and pupae of Drosophila melanogaster. *J. Exp. Biol.* **199**, 1837–44 (1996).

55.   Feder, M. E., Blair, N. & Figueras, H. Natural thermal stress and heat-shock protein expression in Drosophila larvae and pupae. *Funct. Ecol.* **11**, 90–100 (1997).

56.   Sørensen, J. G., Nielsen, M. M., Kruhøffer, M., Justesen, J. & Loeschcke, V. Full genome gene expression analysis of the heat stress response in Drosophila melanogaster. *Cell Stress Chaperones* **10**, 312–28 (2005).

57.   Lockwood, B. L., Julick, C. R. & Montooth, K. L. Maternal loading of a small heat shock protein increases embryo thermal tolerance in Drosophila melanogaster. *J. Exp. Biol.* **220**, 4492–4501 (2017).

58.   Rutherford, S. L. & Lindquist, S. Hsp90 as a capacitor for morphological evolution. *Nature* **396**, 336–42 (1998).

59.   Sangster, T. A., Lindquist, S. & Queitsch, C. Under cover: causes, effects and implications of Hsp90-mediated genetic capacitance. *BioEssays* **26**, 348–62 (2004).

60.   Rudman, S. M. *et al.* Direct observation of adaptive tracking on ecological time scales in Drosophila. *Science (80-.).* **375**, (2022).

61.   Lynch, J. A., El-Sherif, E. & Brown, S. J. Comparisons of the embryonic development of Drosophila , Nasonia , and Tribolium. *Wiley Interdiscip. Rev. Dev. Biol.* **1**, 16–39 (2012).

62.   Lemke, S., Kale, G. & Urbansky, S. Comparing gastrulation in flies: Links between cell biology and the evolution of embryonic morphogenesis. *Mech. Dev.* **164**, 103648 (2020).

63.   Raguso, R. A. Don't forget the flies: dipteran diversity and its consequences for floral ecology and evolution. *Appl. Entomol. Zool.* **55**, 1–7 (2020).

64.   Harvey, J. A., Heinen, R., Gols, R. & Thakur, M. P. Climate change-mediated temperature extremes and insects: From outbreaks to breakdowns. *Glob. Chang. Biol.* **26**, 6685–6701 (2020).

65.   Wagner, D. L., Grames, E. M., Forister, M. L., Berenbaum, M. R. & Stopak, D. Insect decline in the Anthropocene: Death by a thousand cuts. *Proc. Natl. Acad. Sci.* **118**, 1–10 (2021).

66.   Condamine, F. L., Clapham, M. E. & Kergoat, G. J. Global patterns of insect diversification: towards a reconciliation of fossil and molecular evidence? *Sci. Rep.* **6**, 19208 (2016).

67.   Wiegmann, B. M. B. Episodic radiations in the fly tree of life. *Proc. …* **108**, 5690–5695 (2011).

68.   Clapham, M. E. & Karr, J. A. Environmental and biotic controls on the evolutionary history of insect body size. *Proc. Natl. Acad. Sci.* **109**, 10927–10930 (2012).

69.   Wang, B., Xu, C. & Jarzembowski, E. A. Ecological radiations of insects in the Mesozoic. *Trends Ecol. Evol.* **37**, 529–540 (2022).

70.   Chown, S. L. & Gaston, K. J. Body size variation in insects: a macroecological perspective. *Biol. Rev.* **85**, 139–169 (2010).

71.   Bogovic, J. A., Hanslovsky, P., Wong, A. & Saalfeld, S. Robust registration of calcium images by learned contrast synthesis. in *2016 IEEE 13th International Symposium on Biomedical Imaging (ISBI)* vols 2016-June 1123–1126 (IEEE, 2016).

72.   Pietzsch, T., Saalfeld, S., Preibisch, S. & Tomancak, P. BigDataViewer: visualization and processing for large image data sets. *Nat. Methods* **12**, 481–3 (2015).

73.   Preibisch, S., Saalfeld, S. & Tomancak, P. Globally optimal stitching of tiled 3D microscopic image acquisitions. *Bioinformatics* **25**, 1463–1465 (2009).

74.   Ollion, J., Cochennec, J., Loll, F., Escudé, C. & Boudier, T. TANGO: a generic tool for high-throughput 3D image analysis for studying nuclear organization. *Bioinformatics* **29**, 1840–1 (2013).

75.   Herbert, S. *et al.* LocalZProjector and DeProj: a toolbox for local 2D projection and accurate morphometrics of large 3D microscopy images. *BMC Biol.* **19**, 136 (2021).

76. Telley, I. A., Gáspár, I., Ephrussi, A. & Surrey, T. Aster migration determines the length scale of
nuclear separation in the Drosophila syncytial embryo. *J. Cell Biol.* **197**, 887–895 (2012).

77. Oakeshott, J. G., McKechnie, S. W. & Chambers, G. K. Population genetics of the metabolically
related Adh, Gpdh and Tpi polymorphisms in Drosophila melanogaster I. Geographic variation in
Gpdh and Tpi allele frequencies in different continents. *Genetica* **63**, 21–29 (1984).

78. McColl, G. & McKechnie, S. W. The Drosophila heat shock hsr-omega gene: An allele frequency cline
detected by quantitative PCR. *Mol. Biol. Evol.* **16**, 1568–1574 (1999).

79. Bettencourt, B. R., Kim, I., Hoffmann, A. A. & Feder, M. E. Response to natural and laboratory
selection at the Drosophila hsp70 genes. *Evolution (N. Y).* **56**, 1796–1801 (2002).

80. Frydenberg, J., Hoffmann, A. A. & Loeschcke, V. DNA sequence variation and latitudinal associations
in hsp23, hsp26 and hsp27 from natural populations of Drosophila melanogaster. *Mol. Ecol.* **12**,
2025–2032 (2003).

81. Umina, P. A., Weeks, A. R., Kearney, M. R., McKechnie, S. W. & Hoffmann, A. A. Evolution: A rapid
shift in a classic clinal pattern in Drosophila reflecting climate change. *Science (80-.).* **308**, 691–693
(2005).

82. Sawyer, L. A. *et al.* The period gene Thr-Gly polymorphism in Australian and African Drosophila
melanogaster populations: Implications for selection. *Genetics* **174**, 465–480 (2006).

83. Weeks, A. R., Mckechnie, S. W. & Hoffmann, A. A. In search of clinal variation in the period and clock
timing genes in Australian Drosophila melanogaster populations. *J. Evol. Biol.* **19**, 551–557 (2006).

84. Rand, D. M., Weinreich, D. M., Lerman, D., Folk, D. & Gilchrist, G. W. Three selections are better
than one: Clinal variation of thermal QTL from independent selection experiments in drosophila.
Evolution (N. Y). **64**, 2921–2934 (2010).

85. Kolaczkowski, B., Kern, A. D., Holloway, A. K. & Begun, D. J. Genomic differentiation between
temperate and tropical Australian populations of Drosophila melanogaster. *Genetics* **187**, 245–260
(2011).

86. Paaby, A. B., Bergland, A. O., Behrman, E. L. & Schmidt, P. S. A highly pleiotropic amino acid
polymorphism in the Drosophila insulin receptor contributes to life-history adaptation. *Evolution* **68**,
3395–409 (2014).

87. Vigue, C. L. & Johnson, F. M. Isozyme Variability in Species of the Genus Drosophila. VI. Frequency-
Property-Environment Relationships of Allelic Alcohol Dehydrogenases in D. melanogaster. *Biochem.*
Genet. **9**, 213–227 (1973).

88. Berry, A. & Kreitman, M. Molecular analysis of an allozyme cline: alcohol dehydrogenase in
Drosophila melanogaster on the east coast of North America. *Genetics* **134**, 869–93 (1993).

89. Duvernell, D. D. & Eanes, W. F. Contrasting molecular population genetics of four hexokinases in
Drosophila melanogaster, D. simulans and D. yakuba. *Genetics* **156**, 1191–1201 (2000).

90. Schmidt, P. S., Duvernell, D. D. & Eanes, W. F. Adaptive evolution of a candidate gene for aging in
Drosophila. *Proc. Natl. Acad. Sci.* **97**, 10861–10865 (2000).

91. Verrelli, B. C. & Eanes, W. F. Clinal variation for amino acid polymorphisms at the Pgm locus in
Drosophila melanogaster. *Genetics* **157**, 1649–1663 (2001).

92. Duvernell, D. D., Schmidt, P. S. & Eanes, W. F. Clines and adaptive evolution in the methuselah gene
region in Drosophila melanogaster. *Mol. Ecol.* **12**, 1277–1285 (2003).

93. Sezgin, E. *et al.* Single-locus latitudinal clines and their relationship to temperate adaptation in
metabolic genes and derived alleles in Drosophila melanogaster. *Genetics* **168**, 923–931 (2004).

94. Schmidt, P. S. & Paaby, A. B. Reproductive diapause and life-history clines in North American
populations of Drosophila melanogaster. *Evolution (N. Y).* **62**, 1204–1215 (2008).

95. Fabian, D. K. *et al.* Genome-wide patterns of latitudinal differentiation among populations of
Drosophila melanogaster from North America. *Mol. Ecol.* **21**, 4748–4769 (2012).

96. Machado, H. E. *et al.* Comparative population genomics of latitudinal variation in Drosophila

simulans and Drosophila melanogaster. *Mol. Ecol.* **25**, 723–40 (2016).

97.   Costa, R., Peixoto, A. A., Barbujani, G. & Kyriacou, C. P. A latitudinal cline in a Drosophila clock gene.
Proc. R. Soc. London 43–49 (1992).

98.   Kapun, M. *et al.* Genomic Analysis of European Drosophila melanogaster Populations Reveals
Longitudinal Structure, Continent-Wide Selection, and Previously Unknown DNA Viruses. *Mol. Biol.*
Evol. **37**, 2661–2678 (2020).

99.   Graveley, B. R. *et al.* The developmental transcriptome of Drosophila melanogaster. *Nature* **471**,
473–479 (2011).

1471

Figure 1

Increased embryonic lethality at elevated temperature

A   Temperature regimes during embryonic development

(N) Normal temperature = 25°   (E) Elevated temperature = 29°C

NN
EN
NE

0hr   3hr   6hr   30hr

Time since egg fertilization

B   Lethality at the end of embryo development

NN
EN
NE

5   10   15   20
% dead embryos

Gastrulation defects after exposure to elevated temperature

NN

EN (mild defect)

EN (severe defect)

20 min   40 min   60 min

time since onset of gastrulation

Quantification for developmental defects after exposure to elevated temperature

F   Germband extension (GBE)

GBE (% embryo length)

time since onset of gastrulation (min)

G   Developmental landmarks

EN

NN

time since onset of gastrulation (min)

• NN (12/12)

• EN (8/10) embryos continue post-gastrulation development (comparable to NN embryos in 'Developmental landmarks' analysis)

EN (2/10) embryos fail to complete gastrulation (excluded from 'Developmental landmarks' analysis)

■ Germband extends to 40% embryo length

◆ Germband retraction begins

✕ Germband retraction ends

+ head involution

Figure 2

The distribution of blastoderm holes is not bilaterally symmetrical

A Blastoderm holes during cellularization

Normal temperature

67 (C, F, I, J)

Elevated temperature

11 (E, H, M, N)

66 (D, G, K, L)

Blastoderm holes ● absent ● present

B

Schematic of embryo volume. Here colors represent the regions where cortical (green), subcortical (magenta), and yolk (yellow) nuclei are found.

18 μm
45 μm

Normal temperature (n=67/68 embryos)

C Anterior / Right / Posterior / Dorsal half / Left / Ventral half / Right

F

I Dorsal / Left / Right / Ventral

J

Elevated temperature (mild defect, n=66/77 embryos)

D

G

K

L

Elevated temperature (severe defect, n=11/77 embryos)

E

H

M

N

O Right / Dorsal half / Left

P Left / Ventral half / Right

Figure 3

Nuclear expulsion at elevated temperature

22°C (n=10/10)   25°C (n=7/7)   28°C (n=9/9)   29°C (mild defect, n=8/10)   29°C (severe defect, n=2/10)

Cortical and Subcortical (lost) nuclei

Figure 4

Isolated nuclear expulsion
out-of-plane divisions (Example from embryo at 25°C)

A

X-Y view

Y-Z view

Histone-RFP Jupiter-GFP

B

in-plane divisions (Example from embryo at 25°C)

Histone-RFP

0 min   4 min   7 min   12 min

Embryo cross-section

Subcortical (lost) nuclei

Cortical nuclei

Imaging depth

Cortical and Subcortical (lost) nuclei

C

out-of-plane divisions during mitosis-10

number of divisions

*

22°C   25°C   28°C   29°C

D

number of isolated nuclear expulsions during interphase 14

number of lost nuclei

22°C   25°C   28°C   29°C

Pairwise nuclear expulsion after in-plane divisions due to Mitotic failure
(Example from embryo at 29°C)

E   Histone-RFP

−05:00 min   00:00 min   00:30 min

05:00 min   10:00 min   15:00 min

Cortical and Subcortical (lost) nuclei

F

number of mitotic failures causing nuclear expulsions during interphase 14

number of lost pairs of nuclei

*
*
**

22°C   25°C   28°C   29°C

G   Mitotic failures increase at elevated temperature

Normal temperature   Elevated temperature

• • Normal mitosis

•-• Mitotic failure and pairwise loss of nuclei

Figure 5

Nuclear group-center drift and deviations in x-y plane

Figure 6

Increased nuclear crowding at elevated temperature

Histone-RFP

Pseudo-cells

Distribution of pseudo-cell areas

22°C

25°C

28°C

29°C

A B C D

E F G H

25 µm² ▬▬▬▬▬ 100 µm²

I J K L

Pseudo-cell areas (µm²)

Anterior-posterior position (µm)

M crowding at the end of interphase 13

Quadratic coefficient (AU×10⁻³)

22°C 25°C 28°C 29°C

N Nuclear crowding intensifies at elevated temperature

Normal temperature   Elevated temperature

• individual nuclei   ➜ Gradient of nuclear density

Asynchrony in the phases of nuclear division cycles at elevated temperature

Histone-RFP

O 25°C

Telophase → Anaphase → Metaphase

P 29°C

Telophase → Anaphase → Metaphase ← Anaphase

Cortical and Subcortical (lost) nuclei

Asynchrony in the initiation of interphase 14

Q

Asynchrony (min)

22°C 25°C 28°C 29°C

R Asynchrony in nuclear cycle increases at elevated temperature

Normal temperature   Elevated temperature

◯ Nuclear cycles **11**, **12**, and **13** in different regions of the embryo

⊢⋯⋯⊣ Asynchrony at the beginning of interphase 14

Figure 7
combinatorial effects of nuclear crowding and nuclear cycle asynchrony on distribution of mitotic failures at elevated temperature

Figure 8

DNA damage response after mitotic failures at elevated temperature

A — Stem-Loop Binding Protein (SLBP) mislocalization as a DNA damage reporter

B — Nuclei with DNA damage are mitotically arrested

Mitotic failure leads to DNA damage response

C — Wild-type SLBP

D — DNA damage insensitive SLBP

E — Nuclear Localization Signal (NLS)

(representative embryos at 29°C)

nuclear recruitment kinetics

F — SLBP-GFP

G — SLBP.S118A-GFP

H — NLS-GFP

I — SLBP-GFP

J — SLBP.S118A-GFP

K — NLS-GFP

— / · 25°C control division   — / · 29°C control division   --- / ○ 29°C mitotic failures

Figure 9

A analysis of cytoskeletal distribution during mitosis

prophase → metaphase

(manual annotation) → (automated analysis)

F-actin cap contour → cap area and aspect ratio → cap orientation

Spindle poles → Spindle length and orientation → Spindle vs. Cap angle mismatch

F-actin (cap), microtubules (spindle), and Chromatin

Interaction between F-actin cap and microtubule spindle at elevated temperature

B 25°C Control division
C 29°C Control division
D 25°C mitotic failure
E 29°C mitotic failure

-3 min, -2 min, -1 min, 0 min

UtrABD-GFP + Jupiter-GFP Histone-RFP

F F-actin cap size at metaphase

Cap Area (μm²)

25°C 29°C | 25°C 29°C
Control divisions | Mitotic failures

G F-actin cap aspect ratio at metaphase

Aspect ratio

25°C 29°C | 25°C 29°C
Control divisions | Mitotic failures

H Spindle length at metaphase

Spindle length (μm)

25°C 29°C | 25°C 29°C
Control divisions | Mitotic failures

I Spindle extension rate during mitosis

Extension rate (μm/min)

25°C 29°C | 25°C 29°C
Control divisions | Mitotic failures

J Spindle rotation during mitosis

Cumulative rotation (degrees)

25°C 29°C | 25°C 29°C
Control divisions | Mitotic failures

K Spindle-Cap angle mismatch at metaphase

Angle mismatch (degrees)

25°C 29°C | 25°C 29°C
Control divisions | Mitotic failures

Figure 10

Rescue of embryonic lethality at elevated temperature

A Embryo lethality

B fraction of mitotic failures in crowded region

C fraction of mitotic failures in combined region

"Natural" candidate genes varying along thermoclines

D

203 genes

37 genes

138 genes

{Spindle dynamics} = {Cell polarity} ∪
{Spindle localization} ∪ {Spindle organization}

E

Spindle
dynamics
(n=320)

1689

Metabolic
process
(n=6612)

4612

Genes clinal along   0   1   2   3 clines

Figure S1

Blastoderm holes at elevated temperature

Figure S2

Nuclear expulsion is a passive process

A Normal temperature (22°C)

Interphase-12 ⊢——————— mitosis-12 ———————⊣ Interphase-13 ——————→

B Elevated temperature (29°C)

Interphase-12 ⊢——————— mitosis-12 ———————⊣ Interphase-13 ——————→

● Cortical nuclei that failed to divide in mitosis 10   ● Cortical nuclei that failed to divide in mitosis 11

Figure S3

Nuclear movement along z axis at elevated temperature

A  Normal temperature (22°C)    B  Elevated temperature (29°C)

Figure S4

Asynchrony in the phases of nuclear division cycles at elevated temperature

Cortical and Subcortical (lost) nuclei

Figure S5

Rescue of embryonic lethality at elevated temperature

Figure S6

Empty F-actin caps at elevated temperature

Figure S7

Rescue of embryonic lethality at elevated temperature