1 # Mitochondrial genome analyses uncover intriguing population
2 # structure across the invasive trajectory of an iconic invader
3

Running title: Intriguing population structure of an invader

Cheung K[1,2*], Amos TG[2,3], Shine R[4], DeVore JL[5], Ducatez S[6], Edwards RJ[2,7 #] & Rollins LA[1 #]

[1]Evolution & Ecology Research Centre, School of Biological, Earth & Environmental Sciences,
University of New South Wales, Sydney, New South Wales, Australia

[2]School of Biotechnology & Biomolecular Sciences, University of New South Wales, Sydney, New
South Wales, Australia

[3]Garvan Institute of Medical Research, Sydney, New South Wales, Australia

[4]Department of Biological Sciences, Macquarie University, Sydney, New South Wales, Australia

[5]Univ. Polynésie Francaise, UMR 241 EIO (UPF, IRD, IFREMER, ILM) BP 6570 Faa'a, Tahiti, French
Polynesia

[6]Institut de Recherche pour le Développement (IRD), UMR 241 EIO (UPF, IRD, IFREMER, ILM) BP 6570
Faa'a, Tahiti, French Polynesia

[7]Ocean Genomes Laboratory, Minderoo OceanOmics Centre, The University of Western Australia,
Western Australia, Australia

[#]Joint last authors

*Corresponding author: kelton.cheung@unsw.edu.au

Abstract

Invasive species offer insights into rapid adaptations in novel environments. The iconic cane toad (*Rhinella marina*) is an excellent model for studying rapid adaptation during invasion. Previous research using the mitochondrial NADH dehydrogenase 3 (*ND3*) gene in the Hawai'ian and Australian invasive populations found a single haplotype, indicating an extreme genetic bottleneck following introduction. Nuclear genetic diversity also exhibited reductions across the genome in these two populations. Here, we investigated the mitochondrial genome diversity of cane toads across this invasion trajectory. We created the first reference mitochondrial genome of the cane toad with long-read sequencing and constructed a phylogeny of Anura full mitochondrial genomes. We used transcriptomic data of 125 individuals from the native (French Guiana) and introduced (Hawai'i and Australia) ranges to construct nearly-complete mitochondrial genomes for population genomics analyses. As expected, the cane toad belongs to family Bufonidae, distinct from genus *Bufo*. In agreement with previous investigations of these populations, we identified genetic bottlenecks in both Hawai'ian and Australian introduced populations, alongside evidence of population expansion in the invasive ranges. Although mitochondrial genetic diversity in introduced populations was reduced, our results revealed that it had been underestimated: we identified 45 mitochondrial haplotypes in Hawai'ian and Australian samples, none of which were found in the native range. Additionally, we identified two distinct groups of haplotypes from the native range, separated by a minimum of 110 base pairs (0.6%). These findings enhance our understanding of Anura phylogenetics and how invasion has shaped the genetic landscape of this species.

42 Keywords

43 *Rhinella marina*, population genomics, mitochondrial genome, haplotypes, invasion biology

Introduction

Invasive species are a substantial global concern, particularly due to likely range expansions under climate change (Bonebrake et al., 2018; Seebens et al., 2017), an increasing rate of anthropogenic introductions (Lowe, Browne, Boudjelas, & De Poorter, 2000), and their negative impacts on native biodiversity and the economy (Bradshaw et al., 2016; Mainka & Howard, 2010). Invasive species often rapidly adapt to their new environments (Rollins et al., 2013), creating an opportunity to study the process of rapid evolution. Biological invasions sometimes present a genetic paradox where, despite small founding population sizes that often reduce genetic diversity in the introduced range (i.e. genetic bottlenecks), they still maintain the ability to readily adapt to novel conditions (Allendorf & Lundquist, 2003; Estoup et al., 2016; Schrieber & Lachmuth, 2017).

Various genetic markers, including those from the nuclear and mitochondrial genomes, have been employed to investigate the introduction history and evolutionary processes of invasive species. Because of their distinct mode of inheritance and mutation rates, nuclear and mitochondrial markers may yield different insights into the evolutionary events underlying invasion (Toews & Brelsford, 2012). Genetic variation in invasive species is shaped by complex interplays among genetic drift, demography and selection. Small founding populations, common during an invasion, can promote genetic drift, leading to reduced genetic diversity. Furthermore, whilst genetic drift results in less efficient selection (Gravel, 2016), populations expanding their range into new environments may experience novel selection regimes (Sakai et al., 2001) and spatial sorting (Shine, Brown, & Phillips, 2011) that further reduce genetic diversity.

Mitochondrial DNA (mtDNA) is more susceptible to genetic drift than nuclear markers due

to its uniparental inheritance and small effective population size (Klucnika & Ma, 2019; Xiao,

Nguyen, Wu, & Hao, 2017). Nonetheless, mtDNA has been extensively used to study the

evolutionary history of different species or populations within the same species because of

its non-recombining nature and purported near-neutrality. However, the neutrality of

mtDNA has been challenged by evidence of selection, likely due to the important role of

mitochondria in ATP production and cellular activities (Meiklejohn, Montooth, & Rand,

2007). Deleterious mutations are rapidly removed by purifying selection to maintain the

function of the electron transport chain. Further, variation in mtDNA has been linked to

adaptation to high latitudes (Chen et al., 2020; Wang et al., 2021) and temperature (Baker

et al., 2019; Cheng, Liu, Yang, Sun, & Kong, 2013; Silva, Lima, Martel, & Castilho, 2014).

There exists a substantial repository of mtDNA data in public databases, enabling

comprehensive comparative genetic analyses, increasing its usefulness as a marker.

The cane toad (*Rhinella marina*) is one of the world's most infamous, and well-studied,

invasive species (Lowe et al., 2000). Cane toads are native to Central and South America

(Zug & Zug, 1979) and have been introduced to multiple Caribbean islands for biological

control of cane beetles, a parasite of sugarcane (Easteal, 1981). Cane toads were

transported from the Caribbean to Hawai'i in 1932 for cane beetle control and, in 1935, 101

Hawai'ian cane toads were translocated from Oahu to Queensland, Australia (Turvey, 2013).

However, they failed to control cane beetles in Australia and, facilitated by a high

reproductive rate, subsequently spread across the continent to arid Western Australia. This

species' inexorable spread throughout its introduced range in Australia has had catastrophic

effects on the naïve predators it has encountered (Chinchio et al., 2020). Freshwater

crocodiles, snakes, lizards, and quolls have been particularly badly affected by the cane toad

invasion in Australia (Shine, 2010). The colonization of the cane toad is still on-going; toads

occupy over 1.2 million km^2 of Australia (Urban, Phillips, Skelly, & Shine, 2008) including

tropical Queensland, the Northern Territory, and the Kimberley region of northern Western

Australia (Radford et al., 2019).

Like in many other invasions, the cane toads' translocation resulted in genetic bottlenecks.

Australian cane toads have very low MHC class I and class II (Lillie, Cui, Shine, & Belov, 2016;

Lillie, Dubey, Shine, & Belov, 2017; Lillie, Shine, & Belov, 2014) and microsatellite (Leblois,

Rousset, Tikel, Moritz, & Estoup, 2000) diversity. Previous analysis of the NADH

dehydrogenase 3 (*ND3*) mitochondrial gene from 31 individuals sampled from Hawai'i and

Australia identified a single haplotype, a substantial reduction of haplotype diversity

compared to the native range (Slade & Moritz, 1998), further supporting the presence of

genetic bottlenecks in the invasive ranges. Despite this low genetic variation, phenotypic

traits differ substantially across the cane toad's Australian range including immune function,

dispersal ability, and behaviour (Alford, Brown, Schwarzkopf, Phillips, & Shine, 2009; G. P.

Brown, Phillips, Dubey, & Shine, 2015; Gruber, Brown, Whiting, & Shine, 2018; Gruber,

Whiting, Brown, & Shine, 2017; Phillips, Brown, & Shine, 2010). Some of these traits have

been shown to be intergenerationally transmitted, consistent with genetic causes. For

example, toads from range-core and range-edge populations that were raised in a common-

garden experiment showed divergence of behavioural traits and morphology (Gruber,

Brown, Whiting, & Shine, 2017; Hudson, Brown, & Shine, 2016). The invasion success of this

110 species in Australia, coupled with evidence of evolutionary change since introduction make

111 it an ideal system for the study of rapid evolution during invasion.

112 Despite extensive research into toad evolution in the introduced range, we need improved

113 genetic resources to clarify the molecular mechanisms underlying these changes.

114 Transcriptome data from spleen, muscle and/or brain tissues of toads from French Guiana,

115 Hawai'i, and Australia, and liver tissue from Brazil exist (Richardson et al., 2018; Rollins,

116 Richardson, & Shine, 2015; Selechnik, Richardson, Shine, Brown, & Rollins, 2019; Yagound et

117 al., 2022) and a draft reference genome (Edwards et al., 2018) is available, but population

118 genomic analyses of cane toads across this invasion trajectory have thus far been restricted

119 to a single study of single nucleotide polymorphisms (SNPs) from the nuclear genome

120 (Selechnik, Richardson, Shine, DeVore, et al., 2019). Here, we present a complete cane toad

121 mitochondrial genome (mitogenome) and explore the species' phylogenetic relationships

122 with other Anurans. We then investigate evolution across the invasion trajectory using

123 mtDNA data derived from RNASeq and whole-genome resequencing. We compare our data

124 to those of a previously-published study of *ND3* on the same populations (Slade & Moritz,

125 1998), to determine whether our whole-mitochondrial genome analysis can extend

126 population-genetic inferences. We predicted that there would be decreasing genetic

127 diversity within the mitochondrial genome with increasing distance from the native range

128 due to genetic bottlenecks. We also predicted that we would identify more genetic diversity

129 within Australia than has been previously characterised, given the phenotypic evidence of

130 adaptation to this novel environment.

Materials and Methods

Reference mitochondrial genome

High-molecular-weight genomic DNA was extracted from the liver of a female cane toad

collected in the Kimberley region of Western Australia and sequenced on Illumina HiSeq X

Ten and PacBio RS II platforms as described previously (Edwards et al., 2018). The

mitogenome was assembled from 36 SMRT cells generated from the first two SMRTBell

libraries (ENA experiments ERX2389178 and ERX2389179).

The 17,757 basepair (bp) *Bufo japonicus* complete mitochondrial genome (gi:157939553)

was downloaded from NCBI on 29/3/2016 and used to identify mitochondrial subreads from

the raw sequencing data. GABLAM (Davey, Shields, & Edwards, 2006) was used to map the

B. japonicus mtDNA onto PacBio subreads using blastn (blast+ v2.2.31) and extract 143

subreads that had 75%+ bidirectional coverage, *e.g.* at least 75% of the subread mapped

onto the mtDNA and at least 75% of the mtDNA was covered by the subread. Any subreads

that mapped the negative strand were reverse complemented. The subread with the best

global hit to the *B. japonicus* mtDNA was selected as the core of the cane toad

mitochondrial genome. A second GABLAM search was performed of the 143 subreads

against this new query, retaining 29 sequences that had >80% global sequence identity

within local matches in the same orientation and order, excluding any "wrapping" due to

circularity. These 29 sequences were aligned using MAFFT v2.273 (Katoh & Standley, 2013)

with a very small gap penalty to account for the high indel rate (--localpair --maxiterate 1000

--op 0.1 --ep 0.1 --lop -0.1 --nuc). SeqSuite (Edwards, Paulsen, Aguilar Gomez, & Perez-

Bercoff, 2020) was used to generate a consensus sequence from the alignment using the

most frequent non-gap base call at each position. Any column with fewer than 5 non-gap

sequences was excluded from the consensus. All 143 raw subreads were mapped onto the

consensus using BLASR and the sequence was polished using Quiver (SMRT Analysis v2.3.0).

GABLAM was used to identify the overlapping ends of the sequence, which was then

circularised by trimming off the first 1957 bp for a final length of 18,152 bp. A final polishing

step was performed using the Illumina data (ENA experiment ERX2845325), mapped with

Bowtie v2.3.3 (Langmead & Salzberg, 2012) and error-corrected with Pilon v1.20 (Walker et

al., 2014).

The mitochondrial genome was annotated using MITOS online (Bernt et al., 2013) with the

vertebrate mtDNA genetic code (Genetic Code Table 02). Annotated protein-coding genes

were manually curated and extended to include stop codons where required. Where no

canonical in-frame stop codon was found, it was assumed that a partial (T or TA) stop codon

would be completed upon polyadenylation. Codon usage was calculated using

https://www.bioinformatics.org/sms2/codon_usage.html (Accessed: 2023-03-29). The final

annotated mitochondrial genome was uploaded to NCBI (Accession number: NC_066225.1).

The mitogenome map was drawn using the CGView server

(http://stothard.afns.ualberta.ca/cgview_server/).

Phylogenetic analysis

Complete mitochondrial genomes of 203 Anura, and 5 non-Anura outgroup species from

Caeciliidae and Caudata), were downloaded from NCBI BioProject database (Table S3,

Downloaded 2022-09-11). Each mitochondrial genome was re-circularised and edited to end

with the control region for subsequent analyses. The multiple sequence alignment was

conducted with the cane toad reference mitogenome using MAFFT with auto mode to find

the best alignment algorithm (Katoh & Standley, 2013). The resulting alignment was visually

inspected using AliView (Larsson, 2014) without further trimming. Maximum likelihood

phylogenetic analysis was conducted using IQ-Tree multicore version 2.2.2.7 (default

parameters, ultrafast bootstrap (UFBoot) and SH-aLRT branch test with 10000 replicates)

(Minh et al., 2020). Clades are considered reliable if SH-aLRT value >= 80% and UFboot value

>=95%. The best fit model was found by ModelFinder embedded in IQ-Tree

(Kalyaanamoorthy, Minh, Wong, von Haeseler, & Jermiin, 2017). The phylogenetic tree was

visualised and edited using R package *ggtree* (Yu, Smith, Zhu, Guan, & Lam, 2017).

Sequence data for population- and species-level analyses
Previously published cane toad RNASeq data from different tissues were retrieved and

downloaded from NCBI BioProject PRJNA510261 and PRJNA395127 (spleen), PRJNA479937

(brain) and PRJNA277985 (muscle) for analysis. A total of 125 individuals with 144 tissue

samples were included, sampled in French Guiana (n = 24), Hawai'i (n = 8) and Australia (n =

93) (Figure 1; Table S1). For some individuals, RNASeq data from both spleen and brain were

available; in such cases, we used data from brains. Raw reads were processed as described

previously with STAR v2.7.2b (Dobin et al., 2013) and Genome Analysis Toolkit (GATK)

v4.1.9.0 (McKenna et al., 2010), using the extracted mitogenome from the draft genome as

a reference (Selechnik, Richardson, Shine, DeVore, et al., 2019).

Genomic DNA was extracted from fifteen cane toad samples, including two from toe clips,

two from spleen, four from liver and seven from muscle tissues using a Gentra PureGene

DNA extraction kit (QIAGEN) (Table S1).

For two samples (RMH006 and RM265), paired-end sequencing libraries were prepared

using the TruSeq DNA PCR-free kit (Illumina). These libraries were sequenced on the

Illumina NovaSeq 6000 platform at the Ramaciotti Centre for Genomics, University of New

South Wales, Sydney, Australia (UNSW Sydney). The remaining thirteen samples had paired-

end sequencing libraries constructed according to the manufacturer's specifications for the

NEBNext Ultra DNA Library Prep Kit for Illumina (NEB). Three samples (RMF031, RMF042

and RMF044) were sequenced on the 10X Genomics Chromium platform at the Ramaciotti

Centre for Genomics, UNSW Sydney. The remaining libraries were sequenced on the

Illumina NovaSeq 6000 platform at Deakin Genomic Centre, Victoria, Australia with a target

depth of 20-fold coverage per genome. Raw sequencing reads from each sample underwent

initial processing to remove adapter sequences. Trimmomatic v0.38 (Bolger, Lohse, &

Usadel, 2014) was employed using the following parameters: ILLUMINACLIP:TruSeq3-

PE.fa:2:30:10:4 SLIDINGWINDOW:5:30 AVGQUAL:30 MINLEN:36.

Mitochondrial segments, *ND3* gene with tRNA-Gly and tRNA-Arg flanking sequence (*ND3*

dataset hereafter) were obtained from previous studies (Acevedo, Lampo, & Cipriani, 2016;

Slade & Moritz, 1998) (Table S1). We aligned our data with this *ND3* dataset using the R

package DECIPHER (Wright, 2016) and trimmed our data to match the existing sequences.

Construction of mtDNA genome and variant calling

GATK HaplotypeCaller (Poplin et al., 2018) was used to call SNPs and indels using the

following criteria: soft clipped bases were not included and variants with minimum Phred-

scaled confidence of 20 threshold were kept. To filter low quality variants, GATK

VariantFiltration (Van der Auwera, 2020) was used to remove variants in the resulting VCF

file: clusters of 3 SNPs within a window of 35 bases, variants with Fisher Strand (FS) greater

than 30.0, QualbyDepth (QD) less than 2.0, depth of coverage (DP) less than 20.0 and allelic

frequency (AF) less than 0.05. Due to the difficulty of mapping short-read RNASeq data to

sequences with multiple repeats, and the risk of introducing false-positive variant calls, we

excluded variants from the repetitive sequences in the hypervariable region 2 (HV2) of the

mitochondrial control region. We determined the consensus sequence of each sample using

FastaAlternateReferenceMaker (McKenna et al., 2010).

Consensus mitogenome sequences from individuals with both RNASeq and WGS data were

aligned against each other using DECIPHER R packages (Wright, 2016, 2020) using default

settings. The alignments were further visualised and analysed with AliView (Larsson, 2014).

NUMT identification
The presence of NUMTs in the cane toad's genome has not been documented previously.

The cane toad draft genome was retrieved and downloaded from NCBI BioProject

PRJEB24695. We used RepeatModeler (http://www.repeatmasker.org/RepeatModeler/) to

construct species-specific repeat library and masked the genome using RepeatMasker

(Tarailo-Graovac & Chen, 2009). NUMTFinder v0.5.3 (Edwards et al., 2021) was used to

identify potential NUMTs by searching the full and repeat-masked versions of the nuclear

genome with our new mtDNA assembly.

Population genetics analyses
The spatial distribution of haplotypes was examined by creating a median-joining haplotype

network using Network and postprocessed using the maximum parsimony calculation to

remove unnecessary median vectors and links (Polzin & Vahdati Daneshmand, 2003). The

final network was drawn using Network Publisher v2.1.2.5 (Fluxus Engineering, Clare, UK).

This same procedure also was used to analyse the *ND3* dataset.

We then investigated population genetic diversity and differentiation. The following

diversity indices were calculated using DnaSP v6 (Rozas et al., 2017): nucleotide diversity (π),

number of polymorphic sites, number of parsimony informative sites, number of haplotypes

(*H*) and haplotype diversity (*h*). Due to the uneven sample size across populations,

haplotype richness (*HR*) was calculated using FSTAT v2.9.4 (Goudet, 1995) to measure the

number of alleles independent of the sample size. DnaSP was also used to generate a

haplotype list for Arlequin software v3.5.2.2 (Excoffier & Lischer, 2010) and a Roehl data file

for use with Network software 10.2.0.0 (Bandelt, Forster, & Rohl, 1999; Bandelt, Forster,

Sykes, & Richards, 1995). To study the population structure, Arlequin was used to calculate

genetic differentiation among and between populations by pairwise fixation indices (*Fst*)

and Analysis of Molecular Variance (AMOVA) to estimate the genetic variation among and

within populations.

To study the demographic history of each population, we used both neutrality statistics and

mismatch distribution tests to infer population expansion. Fu's *Fs* statistics was produced

using Arlequin and Ramos-Onsins and Rozas R_2 statistics were produced using DnaSP. Fu's *Fs*

test calculates the probability of observing a number of haplotype similar or smaller than

the observed number of samples (Fu, 1997). Both tests are based on the infinite-site model

without recombination, which is suitable for analysing mitochondrial genomic data. A

significant negative value of *Fs* indicates an excess of rare alleles and haplotypes due to a

sudden population growth after a population bottleneck or genetic hitchhiking while

positive values indicate a lack of rare mutations, resulting from balancing selection or

population stability. Ramos-Onsins and Rozas R_2 statistic compares the difference between

the number of singleton mutations and the average number of segregating sites (Ramos-

Onsins & Rozas, 2002). The R_2 statistic is known to have better power than Fu's *Fs* when the

sample size is small (< 10) or the number of segregating sites is low (Ramirez-Soriano,

Ramos-Onsins, Rozas, Calafell, & Navarro, 2008). A small positive value of R_2 indicates

population expansion. The statistical power of these tests was validated using 10,000

simulations. Mismatch distributions of the frequency of pairwise differences between

haplotype pairs in each population were also examined using the sudden expansion and

spatial expansion models implemented in Arlequin (Excoffier, 2004; Schneider & Excoffier,

1999). The sum of square deviations (SSD) was used to test the differences between

observed and expected values while the raggedness index was used to explore the

smoothness of the observed mismatch distribution. The statistical significance of these tests

was validated with 500 replicates in Arlequin. A non-significant SSD value and a small

raggedness index value with a unimodal distribution indicates no deviation of the observed

data from the expectation under the expansion model.

Results
Summary of the cane toad reference mitochondrial genome

The mitogenome of *R. marina* (Figure 2) consists of 18,152 bp, including 22 tRNA genes, 2

rRNA genes and 13 protein-coding genes, with 28 genes located on the forward (+) strand

and 9 located on reverse (-) strand (Table S2). There is also a 2,753 bp GC-rich control region

containing the origin of replication. Two groups of tandem repeats were observed in the

latter part of the control region: 530 bps with 6-full repeats and 350 bps with 3-full repeats.

The gene order of the cane toad mitogenome was examined by aligning with the members

of its sister clades (*Bufo* and *Bufotes*) and we found no gene rearrangements. The base

composition of the mitogenome was A: 29%, T: 33%, G: 14% and C: 24%, with a bias with

62% of A + T. The codon usages of those with nucleotide G in the third codon position were

lower than those with A and T. GCG (Ala) was the least used codon and TGG was the only

codon used for encoding Tryptophan. All protein-coding genes used ATG as start codon.

Phylogenetics analysis
We produced a maximum likelihood phylogenetic tree using whole mitogenomes of Anura.

The best fit model was GTR+F+R9 according to lowest Bayesian information Criterion (BIC).

The location of *R. marina* in the phylogeny of Anura is shown in Figure 3. *Rhinella marina*

was grouped with *Anaxyrus americanus*, an American Bufonid and previously classified

under *Bufo* genus, with 100% statistical support in both UFBoot and SH-aLRT in branch tests.

Four species of *Bufotes* and four of *Bufo* were well resolved into two subclades. Even though

there was high branch support for differentiation between the genera *Rhinella* and

Anaxyrus, and high support for species within *Bufotes* and *Bufo*, the divergence among

genera was not significantly supported, which may indicate the signals in the whole

mitochondrial genomes were insufficient for resolving these clades or that wider taxonomic

coverage is required to resolve this phylogeny.

Verification of mtDNA genomes from RNASeq data
A near-complete mitochondrial genome with 17,261 to 17,270 bps (95.1% of reference

mitochondrial genome) for a total of 132 individuals was generated from the RNASeq and

WGS dataset. A sequence with length of 889 bases consisting of repeats in the HV2 section

of the control region was excluded due to difficulty in correctly mapping sequencing reads

on repeat regions. To determine the accuracy of our RNASeq-derived mtDNA sequences, we

then compared the sequences of the six individuals for which we had both RNASeq and

WGS data (Table S4). We identified four sites (site 2244, site, 5162, site 7162, site 7920) that

showed intra-individual variation across the two datasets in multiple individuals (i.e.

potential RNA editing). Two sites in coding regions (site 267, site 477) had intra-individual

differences in only one individual. Additionally, one individual had 12 sites in the control

region that differed across datasets (Table S4). We replaced the sites with intra-individual

variation in multiple individuals with Ns, minimizing variation for subsequent population

analyses that might be introduced by RNA editing.

NUMT analysis
We found 42 mtDNA fragments (38 – 140 bps in length) that mapped to 42 individual

contigs of the cane toad draft nuclear genome with over 80% identity (Table S5). All

fragments corresponded to the control region of the mitogenome. After the draft genome

was masked for repeat elements, no mtDNA fragments were found to map to the nuclear

genome.

Population Genetic Analyses
We identified 262 (1.30%) polymorphic sites in the RNASeq dataset, 198 of which were

parsimony-informative, including 25 from RNA genes, 145 from protein-coding genes, and

28 from non-coding regions and the control region (Table S2), which resulted in the

identification of 58 haplotypes (Table 1; Figure 4). The thirteen native samples contained

twelve unique haplotypes. Two haplotypes found in all introduced populations accounted

for nearly 50% of the sampled individuals. Within introduced populations, a "complex-star"

shape topology with mostly singleton mutants was observed and 41 of the 46 introduced

haplotypes (89%) were private to a single introduction (Hawai'i: 10; Australia: 31).

We also compared the nearly-complete mitochondrial genome dataset with the *ND3*

dataset to demonstrate the consistency of haplotype inference across studies, and to put

our data into the context of previous work. A total of 24 haplotypes were identified in this

analysis (Figure 5); six haplotypes were identified in French Guiana/Guyana, one haplotype

was identified in Indonesian samples, three in Hawai'ian and Australian samples, and the

remaining haplotypes were from other South America localities. Haplotypes from either side

of Andes clearly formed distinct lineages which is consistent to previous work (Acevedo et

al., 2016). The single Australian haplotype (Hap1) previously identified in (Slade & Moritz,

1998) was the most common haplotype in our Australian samples and the other two

haplotypes (Hap2 and Hap3) within Australia were a single nucleotide different to this

common haplotype.

Haplotype diversity was highest in the native range, with reduced diversity in both

Hawai'ian and Australian populations (*h*: 0.9782 vs 0.8800 and 0.8344, respectively) (Table

1, Figure 6A). Native toads had the highest nucleotide diversity (π: 0.00469) compared to

considerably lower diversity in Hawai'ian and Australian toads (π: 0.00010 and 0.00009,

respectively) (Table 1, Figure 6B). Queensland toads had similar haplotype diversity but

higher nucleotide diversity and haplotype richness to Hawai'ian toads. The population from

the Northern Territory had noticeably lower genetic diversity and haplotype richness than

other Australian populations (Table 1, Figure 6A, C). When the relationship between

haplotypes across the invasion trajectory is considered, the twelve haplotypes from French

Guiana were highly dissimilar to those from the Australian and Hawai'ian populations,

forming two distinct clades (Figure 4). These were separated from each other by a minimum

of 110 base pair changes and from the common invasive haplotype by a minimum of 38

base pair changes (Figure 4). Australian and Hawai'ian haplotypes were intermixed and most

of the haplotypes were one base pair different to other haplotypes, with invasive

haplotypes being separated from at least one other haplotype by no more than three base

pair changes.

AMOVA results showed that the overall genetic variation among populations (64.49%) was

larger than the variation within populations (36.33%) (Table 2). Pairwise Fst values were

significant between the native population and all introduced populations (Fst: 0.55 – 0.67;

Table 3). The Fst values between introduced populations were all less than 0.1, indicating

substantially less differentiation in the invasive range than in the native range. The

Hawai'ian population was significantly differentiated from Australian populations (Fst: 0.02,

p = 0.02), but this difference was driven by the Northern Territory (Fst: 0.05, p = 0.01 and

Western Australian populations (Fst: 0.03, p = 0.01); Queensland was not significantly

differentiated to the Hawai'i population (Fst: 0.01, p = 0.06). Within Australia, the

Queensland population showed significant differentiation to both the Northern Territory

and Western Australian populations (Fst: 0.04, p = 0.01 and 0.03, p = 0.003, respectively),

whereas the Northern Territory and Western Australian populations were not genetically

differentiated (Fst: -0.01, p = 0.72).

Two neutrality tests were implemented to study the demographic history of cane toad

populations: Fu's Fs and Ramos-Onsins and Rozas' R_2 statistic (Table 4). The native

population had non-significant positive Fu's Fs values and a large R_2 statistic, indicating the

population is not expanding or shrinking (i.e. is demographically stable). Fu's Fs values were

negative in all introduced populations, indicating an excess of rare haplotypes compared to

the expectation from neutrality and all these were statistically significant. R_2 statistics were

significant in Hawai'i, Queensland and Western Australia's populations.

Mismatch distribution graphs of all populations are shown in Figure S1. The native

population showed a multimodal distribution with non-significant SSD and raggedness

values, indicating support for spatial and demographic expansion (Table 4). Most introduced

populations showed a unimodal distribution for both expansion models with non-significant

SSD and raggedness indices (Table 4), supporting the presence of spatial and demographic

expansions. However, the Northern Territory population had significant or marginally

significant p-values for SSD and raggedness values for both expansion models indicating

poor support (spatial expansion: SSD p-value = 0.03, raggedness p-value = 0.05;

demographic expansion: SSD p-value = 0.07, raggedness p-value = 0.03).

Discussion

This study uses mitochondrial genome sequences to study the evolutionary and

phylogenetic relationships of the iconic invasive cane toad. Here, we constructed the first

reference mitochondrial genome of *R. marina* and placed this species within the order

Anura. We utilised RNASeq-derived mitogenomes to illustrate a greater number and

diversity of haplotypes in mtDNA present in Australia and Hawai'i, in contrast to the

surprisingly low genetic diversity previously described using the *ND3* gene alone. Our

conclusions are further supported by evidence of demographic and spatial expansion in

invasive cane toad populations. This study provides new insights into the diversity of native

cane toads and how they have evolved along their invasion trajectory.

Phylogenetic inference

The phylogenetic relationships of toads within Anura are understudied. More specifically,

the cryptic *Rhinella* species complex remains poorly understood (Pereyra et al., 2021). Zhang

et al. (2013) attempted to study the Anuran phylogeny using partial mitochondrial genomes.

However, no *Rhinella* species were included in their study. Currently, the evolutionary

relationship of *Rhinella* species in the order Anura has yet to be fully documented using

complete mitochondrial genome or even nuclear genes and/or mitochondrial genes. To

address this knowledge gap, we generated the first complete mitochondrial genome from

any *Rhinella* species and investigated the phylogenetic relationship of *Rhinella* to other

Anurans. Although the topology of our tree confirmed previous hypotheses (e.g., in line with

Feng et al. (2017), *Rhinella* formed a sister clade with *Anaxyrus*), it also yielded different

conclusions in some aspects. For example, Feng et al. (2017) built a tree for frog species

using 95 unlinked nuclear genes and suggested that the genus *Bufo* is monophyletic with

respect to *Duttaphrynus*. In contrast, our whole-mitochondrial genome phylogenetic

analysis found that the genus *Bufo* is a subclade of *Duttaphrynus*. Our study provides one of

the most comprehensive phylogenetic analyses of the relationships within the order Anura

to date and provides a starting point for future phylogenetic analyses within the genus

Rhinella.

Choosing to use a complete dataset of a few selected loci versus a large supermatrix with

missing data for phylogenetic inference remains a controversial issue (see for review

(Lemmon, Brown, Stanger-Hall, & Lemmon, 2009; Roure, Baurain, & Philippe, 2013; Wiens &

Morrill, 2011)). Typically, phylogenetic trees are generated using single or multiple nuclear

genes, single or multiple nuclear genes plus mitochondrial genes, or multiple nuclear genes

with morphological trait matrices, but the use of complete mitochondrial genomes for

phylogenetic inference is rare due to resource limitations. Missing data in phylogenetic

analyses can bias phylogenetic calculation and tree branch lengths (Lemmon et al., 2009;

Xia, 2014). Nonetheless, contrasting findings have also suggested that when a sufficient

number of loci and taxa are included, phylogenetic signals remain consistent regardless of

whether or not the dataset includes missing data (Molloy & Warnow, 2018; Shavit Grievink,

Penny, & Holland, 2013). Despite using complete mitochondrial genomes in this study, 11%

of branches remained statistically unsupported, and the genera *Bufo*, *Bufotes*, *Anaxyrus* and

Rhinella in the Bufonidae family are also not well-supported. This result suggests that the

resolution power of the complete mitochondrial genome may not be sufficient, and/or

more representative sampling of different anuran species may be needed to provide more

robust support for statistical calculations and improve our understanding of the phylogeny.

Comparison between RNASeq & WGS results
The use of transcriptomic data to build mitochondrial genomes has been suggested as a

cost-effective alternative to long PCRs or direct sequencing of total DNA (Forni et al., 2019),

but it is important to ensure that the inferred polymorphisms accurately reflect genetic

variation. In our study, comparing mtDNA data obtained from WGS and RNASeq, we found

discrepancies in multiple nucleotide positions, all of which were in rRNA, tRNA and the

control region, which could be the result of post-transcriptional modifications or RNA

editing which are well-recognised phenomena across multiple taxa (Levanon et al., 2004;

Porath, Knisbacher, Eisenberg, & Levanon, 2017; St Laurent et al., 2013). While common

substitutions such as adenosine to inosine (A-to-I) deamination and cytidine to uridine (C-to-

U) pyrimidine exchange were found, we also observed less common substitutions (e.g.

thymine to cytidine (T-to-C) that have been previously documented in another toad species,

Xenopus tropicalis (Zaranek, Levanon, Zecharia, Clegg, & Church, 2010). These processes are

important for creating multiple transcript isoforms and gene regulation (Tang, Fei, & Page,

2012) and are known to be associated with adaptation (Duan, Dou, Luo, Zhang, & Lu, 2017;

Duan et al., 2021). RNA editing of the tRNA anticodon recognition site would extend the

codon matching ability and further increase the variation of the resulting translated

proteins. Although we did not observe nucleotide differences in position 34 or position 37 of

tRNA, which are typically recognised for deamination of adenosine to inosine which can be

paired with A, C or U residues (Jackman & Alfonzo, 2013), we did find changes in the D arm

of tRNA (Aspartic acid and Lysine), the T arm of tRNA (Phenylalanine) and the acceptor stem

of tRNA (Tryptophan) (Figure S2), which are critical for the recognition of aminoacyl tRNA

synthetases to bind appropriate amino acids to its designated tRNA (Ganesh & Maerkl,

2022). Failure to form a tRNA-amino acid complex could inhibit the use of a specific amino

acid in subsequent protein translation. Further proteome analysis is required to determine if

these changes affect the protein translation machinery.

Post-transcriptional modifications were more diverse in the native populations than in the

invasive populations. In particular, one individual from French Guiana displayed

discrepancies in the control region between WGS and RNASeq data, all of which were

transitions. This observation might result from tissue-specific expression in spleen or RNA

editing. Studies of RNA editing in mtDNA have primarily focused on protein-coding genes

and tRNAs, with less research conducted on the control region of the mtDNA, which is

responsible for regulating RNA and DNA synthesis. While the implications of potential RNA

editing in cane toads remain unclear, these events may contribute to diversity at a

transcriptional and translational level. The low diversity of post-transcriptional modification

in the invasive populations might be the result of selective sweep for favourable haplotypes

with low activity of RNA-editing enzymes.

NUMTs, or nuclear insertions of mitochondrial DNA fragments (Lopez, Yuhki, Masuda, Modi,

& O'Brien, 1994), can potentially present another source of genetic variation and contribute

to false-positive signals of variation in nuclear data (Song, Buhay, Whiting, & Crandall, 2008).

NUMTs have been reported in many different taxa including vertebrates, plants and protists

(Hazkani-Covo, Zeller, & Martin, 2010). They vary in size, ranging from minimal insertions to

hundreds of kilobases and tend to be longer in species with larger nuclear genomes

(Hazkani-Covo et al., 2010; Ko & Kim, 2016). Despite NUMTs generally being perceived as

non-functional due to their ability to shift reading frames, the presence of in-frame stop

codons and the use of different genetic codes, the transcription of NUMTs into non-coding

RNA molecules can be captured by RNASeq and potentially falsely interpreted as

mitochondrial variation among or within individuals. NUMTs studies on amphibian species

are limited. In *Xenopus tropicalis*, over 10kb of NUMTs has been found in the nuclear

genome (Hazkani-Covo et al., 2010). Investigation of the presence/absence of NUMTs in

cane toad nuclear genome and the potential expression of NUMTs is required to distinguish

between genuine mtDNA variation within individuals and false-positive signals caused by

incorporating nuclear genetic data into mitochondrial genetic analyses.

A search for NUMTs in the published cane toad draft genome (Edwards et al., 2018) yielded

only matches with simple repeat regions within the mtDNA control region. In other species,

NUMTs originate across the entire mtDNA genome (Dayama, Zhou, Prado-Martinez,

Marques-Bonet, & Mills, 2020; Wei et al., 2022). Following the repeat-masking of the draft

nuclear genome, we found no evidence of NUMTs suggesting that the putative NUMTs were

just low-complexity regions in the nuclear genome. Similar to other gene-searching methods

such as BUSCO, the accuracy of the search algorithm heavily depends on the contiguity of

the genome and the divergence of the query from the database subjects. Given the high

number of contigs in the cane toad draft genome, potential NUMTs hits may have been

missed through breakpoints. Therefore, it is important to develop a cane toad genome

assembly with higher contiguity to improve our ability to investigate NUMTs. Additionally,

ancient NUMTs may be too divergent to be detected by the methods employed, but NUMTs

with such low identity would not interfere with mtDNA analysis.

It is worth noting that the consensus sequences generated from the bioinformatic pipelines

used in this study represented the most common nucleotide at positions where multiple

nucleotides were found. Although these discrepancies could result from sequencing errors,

our filtering is likely to have prevented this. As discussed above, RNA editing or NUMTs

could have resulted in putative polymorphisms within individuals in our RNASeq dataset. It

is also possible that in some cases, heteroplasmy could have caused these putative

polymorphisms. While determining the cause of this phenomenon is beyond the scope of

this study, it is unlikely that this has affected population-level statistics.

Population genetic analyses

Interestingly, both the haplotype network (Figure 4) and the multimodal mismatch

distribution (Figure S1) indicate the presence of multiple genetically divergent lineages

within native populations. Notably, individuals within each lineage are intermixed, rather

than being grouped by geographical locations. For instance, despite an 80 km geographical

separation, haplotypes H4 and H7 are found in the same lineage while haplotypes H12 and

H11 were collected only 9 m apart but belong to different lineages. Native-range cane toads

can travel 60-180 m per move for sheltering and foraging purposes (Shine et al., 2021).

However, the overall dispersal rates are quite low, as the toads ultimately tend to return to

the same shelters (DeVore, Shine, & Ducatez, 2021). Consequently, the possibility of

hybridisation between toads from these distinct lineages exists. However, a recent study in

South and Central America which utilised ddRAD sequencing has offered a different

perspective on the phylogenetic relationship of cane toads. This study revealed toads from

the coastal area formed two sister clades, distinct from the clade formed by toads from the

rainforest regions (Mittan-Moreau et al., 2022). It becomes crucial to ascertain the precise

populations of cane toads from the native range that were used as founding individuals,

especially given the limited comprehensive description in prior research (Easteal, 1981). In

Shine et al. (2021), cane toads from the range-edge regions (Northern Territory and

Western Australia) exhibited similar dispersal patterns to native toads in one of the coastal

sites. This suggests the adaptation of cane toads in hot and arid environments is likely

facilitated by the standing genetic variation(s) inherited from the progenitors utilising dry,

saline coastal beach environments (DeVore et al., 2021), rather than relying solely on the

genetic adaptation and selection of novel mutation(s). A more thorough investigation into

these distinct lineages becomes imperative for unravelling the mechanisms underlying the

adaptations of cane toads in Australia.

Our study challenges the assumption of low genetic diversity in Hawai'ian and Australian

populations. Slade and Moritz (1998) reported a unique haplotype at *ND3* within Australian

populations, in comparison with multiple haplotypes in native populations in Central and

South America. This result supported the presence of founder effects following the

introduction of cane toads into Hawai'i and Australia. Here, we revealed two additional,

previously undocumented, *ND3* haplotypes (Hap2 and Hap3) in Hawai'ian and Australian

populations, each of which differed by only one nucleotide from the previously identified

haplotype (Hap1) (Figure 5). In comparison to our analysis using nearly complete

mitogenomes discussed above, this analysis highlights that a single gene is not sufficient to

accurately infer genetic diversity.

The significant decrease in haplotype diversity and nucleotide diversity observed in both

Hawai'ian and Australian populations, compared to native populations, does support that

genetic bottlenecks and founder effects occurred post-introduction (Table 1, Figure 6). This

finding is consistent with previous studies on this species using nuclear genetic markers

including SNPs, microsatellite markers, and MHC class I and II loci (Lillie et al., 2016; Lillie et

al., 2014; Selechnik, Richardson, Shine, DeVore, et al., 2019). In this study, all three

neutrality indices showed strong signals of population expansion and the observed data

fitted to both demographic expansion and spatial expansion models in most invasive

sampling sites (Table 4). This conclusion is supported by the star-like shaped topology of the

haplotype network created using the nearly-complete mitochondrial genome dataset.

Despite reduced diversity in the introduced range, the degree of the decrease is less than

previously reported (Leblois et al., 2000). Additionally, the star-shaped topology of the

network containing the introduced haplotypes, the absence of native samples here, and the

significant negative Fu's *Fs* values indicate that many of the singleton haplotypes discovered

in Hawai'i and Australia may be the products of post-introduction mutations. Alternatively,

due to the limited sample sizes in the native range, it is possible that any introduced

haplotypes exist but were not included in our sample set for analysis.

We observed a curvilinear pattern of genetic diversity indices across Australian populations

(from Queensland to Western Australia), a pattern also found in spleen gene expression

(Selechnik, Richardson, Shine, Brown, et al., 2019), limb length (Hudson, McCurry, Lundgren,

McHenry, & Shine, 2016), spleen and body mass (Gregory P. Brown, Kelehear, Shilton,

Phillips, & Shine, 2015). The mechanism behind this reduction of genetic diversity in

Northern Territory toads remains unclear, but it may be due to demographic processes that

reduce the effective population size and increase genetic drift (Nei & Tajima, 1981), genetic

hitchhiking that removes neutral polymorphic sites under positive selection (Fay & Wu,

2000) or directional selection due to harsh climate (Selechnik, Richardson, Shine, DeVore, et

al., 2019), possibly in combination with dispersion-driven spatial sorting at the invasion front

(Clarke, Shine, & Phillips, 2019). Despite our finding that the Northern Territory and Western

Australia populations were not genetically differentiated, it is likely that they have

experienced different local evolutionary forces and demographic processes. Northern

Territory toads, unlike other populations, only exhibited demographic expansion and not

spatial expansion, as supported by lower levels in all statistical measures.

Our results demonstrate significant differentiation between native and introduced

populations (Table 3). We identified three genetic clusters: 1) native toads, 2) Hawai'i and

Queensland toads (*Fst*: 0.016, *p* = 0.056) and 3) Northern Territory and Western Australia

toads (*Fst*: -0.012, *p* = 0.72), consistent with previous results using SNPs from the nuclear

genome (Selechnik, Richardson, Shine, DeVore, et al., 2019). It is interesting that the

Queensland toads are not differentiated from those sampled in Hawai'i but differ from

toads sampled in the Northern Territory and Western Australia. Queensland toads have

been separated from their Hawai'ian conspecifics for almost 90 years but do share similar

environmental conditions (Selechnik, Richardson, Shine, DeVore, et al., 2019). Northern

Territory and Western Australian toads suffer from reduced genetic diversity compared to

Queensland toads, so they have likely experienced more genetic drift, which may explain

this difference. However, it is also possible that the striking reduction in precipitation and

increase in temperature experienced by Northern Territory and Western Australian toads

compared to those from Queensland has resulted in selective differences in the

mitogenome between these populations. In support of this possibility, significant negative

Fu's *Fs* can indicate the presence of selection.

Conclusion
591
Our study used the whole mtDNA genome as a genetic marker to investigate the evolution

of cane toads across their invasion trajectory from the native range to Australia, adding to

our knowledge of changes to genetic diversity as cane toads have been serially introduced

to multiple locations. Unsurprisingly, we found reduced genetic diversity in introduced

populations, suggesting that genetic bottlenecks have occurred in these populations.

Despite this reduction, our data indicate greater genetic diversity in introduced populations

than has been previously described using mtDNA. We identified signatures of demographic

and spatial expansion in most introduced localities we sampled, consistent with ongoing

invasion. We described 46 haplotypes in introduced populations, mostly consisting of single

base-pair changes, suggesting evolution following introduction. By the comparison of WGS-

and RNASeq-derived mitogenomes, we have identified the presence of post-transcriptional

modification or RNA editing, which may supply additional genetic diversity upon which

selection can act.

Acknowledgement

We thank Mark Richardson and Yini Peng Lee for assistance with genomic sequencing and

processing of genomic data and the Ramaciotti Centre for Genomics, University of New

South Wales (UNSW) and Deakin Genomic Centre, Deakin University who performed the

sequencing in this study. This research includes computations using the computational

cluster Katana supported by Research Technology Services at UNSW Sydney. This study was

supported by the Australian Research Council (grants DP160102991, DP190100507, to RS

and LAR), the UNSW Scientia Program (to LAR), and the UNSW Scientia PhD Scholarship (to

KC).

Author Contributions

KC, RS, RJE, and LAR developed this project. JLD and SD conducted field work. LAR

conducted DNA extractions. KC, TGA, RJE, and LAR analysed the data. All authors

contributed to data interpretation and writing.

Data availability

The haplotype sequences will be available on the NCBI public database once the manuscript is
accepted.

References

Acevedo, A. A., Lampo, M., & Cipriani, R. (2016). The cane or marine toad, Rhinella marina (Anura, Bufonidae): two genetically and morphologically distinct species. *Zootaxa, 4103*(6), 574-586. doi:10.11646/zootaxa.4103.6.7

Alford, R., Brown, G., Schwarzkopf, L., Phillips, B., & Shine, R. (2009). Comparisons through time and space suggest rapid evolution of dispersal behavior in an invasive species. *Wildlife Research, 36*, 23-28. doi:10.1071/wr08021

Allendorf, F. W., & Lundquist, L. L. (2003). Introduction: Population Biology, Evolution, and Control of Invasive Species. *Conservation Biology, 17*(1), 24-30. doi:https://doi.org/10.1046/j.1523-1739.2003.02365.x

Baker, E. P., Peris, D., Moriarty, R. V., Li, X. C., Fay, J. C., & Hittinger, C. T. (2019). Mitochondrial DNA and temperature tolerance in lager yeasts. *Sci Adv, 5*(1), eaav1869. doi:10.1126/sciadv.aav1869

Bandelt, H. J., Forster, P., & Rohl, A. (1999). Median-joining networks for inferring intraspecific phylogenies. *Mol Biol Evol, 16*(1), 37-48. doi:10.1093/oxfordjournals.molbev.a026036

Bandelt, H. J., Forster, P., Sykes, B. C., & Richards, M. B. (1995). Mitochondrial portraits of human populations using median networks. *Genetics, 141*(2), 743-753. doi:10.1093/genetics/141.2.743

Bernt, M., Donath, A., Juhling, F., Externbrink, F., Florentz, C., Fritzsch, G., . . . Stadler, P. F. (2013). MITOS: improved de novo metazoan mitochondrial genome annotation. *Mol Phylogenet Evol, 69*(2), 313-319. doi:10.1016/j.ympev.2012.08.023

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics, 30*(15), 2114-2120. doi:10.1093/bioinformatics/btu170

Bonebrake, T. C., Brown, C. J., Bell, J. D., Blanchard, J. L., Chauvenet, A., Champion, C., . . . Pecl, G. T. (2018). Managing consequences of climate-driven species redistribution requires integration of ecology, conservation and social science. *Biological Reviews, 93*(1), 284-305. doi:10.1111/brv.12344

Bradshaw, C. J., Leroy, B., Bellard, C., Roiz, D., Albert, C., Fournier, A., . . . Courchamp, F. (2016). Massive yet grossly underestimated global costs of invasive insects. *Nat Commun, 7*, 12986. doi:10.1038/ncomms12986

Brown, G. P., Kelehear, C., Shilton, C. M., Phillips, B. L., & Shine, R. (2015). Stress and immunity at the invasion front: a comparison across cane toad (Rhinella marina) populations. *Biological Journal of the Linnean Society, 116*(4), 748-760. doi:10.1111/bij.12623

654 Brown, G. P., Phillips, B. L., Dubey, S., & Shine, R. (2015). Invader immunology: invasion history alters
655   immune system function in cane toads (Rhinella marina) in tropical Australia. *Ecol Lett,*
656   *18*(1), 57-65. doi:10.1111/ele.12390

657 Chen, Y., Gong, L., Liu, X., Chen, X., Yang, S., & Luo, Y. (2020). Mitochondrial DNA genomes revealed
658   different patterns of high-altitude adaptation in high-altitude Tajiks compared with Tibetans
659   and Sherpas. *Sci Rep, 10*(1), 10592. doi:10.1038/s41598-020-67519-z

660 Cheng, Y. T., Liu, J., Yang, L. Q., Sun, C., & Kong, Q. P. (2013). Mitochondrial DNA content contributes
661   to climate adaptation using Chinese populations as a model. *PLoS One, 8*(11), e79536.
662   doi:10.1371/journal.pone.0079536

663 Chinchio, E., Crotta, M., Romeo, C., Drewe, J. A., Guitian, J., & Ferrari, N. (2020). Invasive alien
664   species and disease risk: An open challenge in public and animal health. *PLoS Pathog, 16*(10),
665   e1008922. doi:10.1371/journal.ppat.1008922

666 Clarke, G. S., Shine, R., & Phillips, B. L. (2019). May the (selective) force be with you: Spatial sorting
667   and natural selection exert opposing forces on limb length in an invasive amphibian. *J Evol*
668   *Biol, 32*(9), 994-1001. doi:10.1111/jeb.13504

669 Davey, N. E., Shields, D. C., & Edwards, R. J. (2006). SLiMDisc: short, linear motif discovery, correcting
670   for common evolutionary descent. *Nucleic Acids Res, 34*(12), 3546-3554.
671   doi:10.1093/nar/gkl486

672 Dayama, G., Zhou, W., Prado-Martinez, J., Marques-Bonet, T., & Mills, R. E. (2020). Characterization
673   of nuclear mitochondrial insertions in the whole genomes of primates. *NAR Genom*
674   *Bioinform, 2*(4), lqaa089. doi:10.1093/nargab/lqaa089

675 DeVore, J. L., Shine, R., & Ducatez, S. (2021). Spatial ecology of cane toads (Rhinella marina) in their
676   native range: a radiotelemetric study from French Guiana. *Sci Rep, 11*(1), 11817.
677   doi:10.1038/s41598-021-91262-8

678 Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., . . . Gingeras, T. R. (2013). STAR:
679   ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21.
680   doi:10.1093/bioinformatics/bts635

681 Duan, Y., Dou, S., Luo, S., Zhang, H., & Lu, J. (2017). Adaptation of A-to-I RNA editing in Drosophila.
682   *PLoS Genet, 13*(3), e1006648. doi:10.1371/journal.pgen.1006648

683 Duan, Y., Dou, S., Porath, H. T., Huang, J., Eisenberg, E., & Lu, J. (2021). A-to-I RNA editing in
684   honeybees shows signals of adaptation and convergent evolution. *iScience, 24*(1), 101983.
685   doi:10.1016/j.isci.2020.101983

686 Easteal, S. (1981). The history of introductions of Bufo marinus (Amphibia: Anura); a natural
687   experiment in evolution. *Biological Journal of the Linnean Society, 16*(2), 93-113.
688   doi:10.1111/j.1095-8312.1981.tb01645.x

689 Edwards, R. J., Field, M. A., Ferguson, J. M., Dudchenko, O., Keilwagen, J., Rosen, B. D., . . . Ballard, J.
690   W. O. (2021). Chromosome-length genome assembly and structural variations of the primal
691   Basenji dog (Canis lupus familiaris) genome. *BMC Genomics, 22*(1), 188. doi:10.1186/s12864-
692   021-07493-6

693 Edwards, R. J., Paulsen, K., Aguilar Gomez, C. M., & Perez-Bercoff, A. (2020). Computational
694   Prediction of Disordered Protein Motifs Using SLiMSuite. *Methods Mol Biol, 2141*, 37-72.
695   doi:10.1007/978-1-0716-0524-0_3

696 Edwards, R. J., Tuipulotu, D. E., Amos, T. G., O'Meally, D., Richardson, M. F., Russell, T. L., . . . White,
697   P. A. (2018). Draft genome assembly of the invasive cane toad, Rhinella marina. *Gigascience,*
698   *7*(9). doi:10.1093/gigascience/giy095

699 Estoup, A., Ravigné, V., Hufbauer, R., Vitalis, R., Gautier, M., & Facon, B. (2016). Is There a Genetic
700   Paradox of Biological Invasion? *Annual Review of Ecology, Evolution, and Systematics, 47*(1),
701   51-72. doi:10.1146/annurev-ecolsys-121415-032116

702   Excoffier, L. (2004). Patterns of DNA sequence diversity and genetic structure after a range
703   expansion: lessons from the infinite-island model. *Mol Ecol, 13*(4), 853-864.
704   doi:10.1046/j.1365-294x.2003.02004.x

705   Excoffier, L., & Lischer, H. E. (2010). Arlequin suite ver 3.5: a new series of programs to perform
706   population genetics analyses under Linux and Windows. *Mol Ecol Resour, 10*(3), 564-567.
707   doi:10.1111/j.1755-0998.2010.02847.x

708   Fay, J. C., & Wu, C. I. (2000). Hitchhiking under positive Darwinian selection. *Genetics, 155*(3), 1405-
709   1413. doi:10.1093/genetics/155.3.1405

710   Feng, Y. J., Blackburn, D. C., Liang, D., Hillis, D. M., Wake, D. B., Cannatella, D. C., & Zhang, P. (2017).
711   Phylogenomics reveals rapid, simultaneous diversification of three major clades of
712   Gondwanan frogs at the Cretaceous-Paleogene boundary. *Proc Natl Acad Sci U S A, 114*(29),
713   E5864-E5870. doi:10.1073/pnas.1704632114

714   Forni, G., Puccio, G., Bourguignon, T., Evans, T., Mantovani, B., Rota-Stabelli, O., & Luchetti, A.
715   (2019). Complete mitochondrial genomes from transcriptomes: assessing pros and cons of
716   data mining for assembling new mitogenomes. *Sci Rep, 9*(1), 14806. doi:10.1038/s41598-
717   019-51313-7

718   Fu, Y. X. (1997). Statistical tests of neutrality of mutations against population growth, hitchhiking and
719   background selection. *Genetics, 147*(2), 915-925. doi:10.1093/genetics/147.2.915

720   Ganesh, R. B., & Maerkl, S. J. (2022). Biochemistry of Aminoacyl tRNA Synthetase and tRNAs and
721   Their Engineering for Cell-Free and Synthetic Cell Applications. *Front Bioeng Biotechnol, 10*,
722   918659. doi:10.3389/fbioe.2022.918659

723   Goudet, J. (1995). FSTAT (Version 1.2): A Computer Program to Calculate F-Statistics. *Journal of
724   Heredity, 86*(6), 485-486. doi:10.1093/oxfordjournals.jhered.a111627

725   Gravel, S. (2016). When Is Selection Effective? *Genetics, 203*(1), 451-462.
726   doi:10.1534/genetics.115.184630

727   Gruber, J., Brown, G., Whiting, M. J., & Shine, R. (2017). Is the behavioural divergence between
728   range-core and range-edge populations of cane toads (Rhinella marina) due to evolutionary
729   change or developmental plasticity? *R Soc Open Sci, 4*(10), 170789. doi:10.1098/rsos.170789

730   Gruber, J., Brown, G., Whiting, M. J., & Shine, R. (2018). Behavioural divergence during biological
731   invasions: a study of cane toads (Rhinella marina) from contrasting environments in Hawai'i.
732   *R Soc Open Sci, 5*(4), 180197. doi:10.1098/rsos.180197

733   Gruber, J., Whiting, M. J., Brown, G., & Shine, R. (2017). The loneliness of the long-distance toad:
734   invasion history and social attraction in cane toads (Rhinella marina). *Biol Lett, 13*(11).
735   doi:10.1098/rsbl.2017.0445

736   Hazkani-Covo, E., Zeller, R. M., & Martin, W. (2010). Molecular poltergeists: mitochondrial DNA
737   copies (numts) in sequenced nuclear genomes. *PLoS Genet, 6*(2), e1000834.
738   doi:10.1371/journal.pgen.1000834

739   Hudson, C. M., Brown, G. P., & Shine, R. (2016). It is lonely at the front: contrasting evolutionary
740   trajectories in male and female invaders. *R Soc Open Sci, 3*(12), 160687.
741   doi:10.1098/rsos.160687

742   Hudson, C. M., McCurry, M. R., Lundgren, P., McHenry, C. R., & Shine, R. (2016). Constructing an
743   Invasion Machine: The Rapid Evolution of a Dispersal-Enhancing Phenotype During the Cane
744   Toad Invasion of Australia. *PLoS One, 11*(9), e0156950. doi:10.1371/journal.pone.0156950

745   Jackman, J. E., & Alfonzo, J. D. (2013). Transfer RNA modifications: nature's combinatorial chemistry
746   playground. *Wiley Interdiscip Rev RNA, 4*(1), 35-48. doi:10.1002/wrna.1144

747   Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., & Jermiin, L. S. (2017).
748   ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods, 14*(6),
749   587-589. doi:10.1038/nmeth.4285

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol, 30*(4), 772-780.
doi:10.1093/molbev/mst010
Klucnika, A., & Ma, H. (2019). A battle for transmission: the cooperative and selfish animal
mitochondrial genomes. *Open Biol, 9*(3), 180267. doi:10.1098/rsob.180267
Ko, Y. J., & Kim, S. (2016). Analysis of Nuclear Mitochondrial DNA Segments of Nine Plant Species:
Size, Distribution, and Insertion Loci. *Genomics Inform, 14*(3), 90-95.
doi:10.5808/GI.2016.14.3.90
Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods,*
9(4), 357-359. doi:10.1038/nmeth.1923
Larsson, A. (2014). AliView: a fast and lightweight alignment viewer and editor for large datasets.
Bioinformatics, 30(22), 3276-3278. doi:10.1093/bioinformatics/btu531
Leblois, R., Rousset, F., Tikel, D., Moritz, C., & Estoup, A. (2000). Absence of evidence for isolation by
distance in an expanding cane toad (Bufo marinus) population: an individual-based analysis
of microsatellite genotypes. *Mol Ecol, 9*(11), 1905-1909. doi:10.1046/j.1365-
294x.2000.01091.x
Lemmon, A. R., Brown, J. M., Stanger-Hall, K., & Lemmon, E. M. (2009). The effect of ambiguous data
on phylogenetic estimates obtained by maximum likelihood and Bayesian inference. *Syst*
Biol, 58(1), 130-145. doi:10.1093/sysbio/syp017
Levanon, E. Y., Eisenberg, E., Yelin, R., Nemzer, S., Hallegger, M., Shemesh, R., . . . Jantsch, M. F.
(2004). Systematic identification of abundant A-to-I editing sites in the human
transcriptome. *Nat Biotechnol, 22*(8), 1001-1005. doi:10.1038/nbt996
Lillie, M., Cui, J., Shine, R., & Belov, K. (2016). Molecular characterization of MHC class II in the
Australian invasive cane toad reveals multiple splice variants. *Immunogenetics, 68*(6-7), 449-
460. doi:10.1007/s00251-016-0919-9
Lillie, M., Dubey, S., Shine, R., & Belov, K. (2017). Variation in Major Histocompatibility Complex
diversity in invasive cane toad populations. *Wildlife Research, 44*(7), 565.
doi:10.1071/wr17055
Lillie, M., Shine, R., & Belov, K. (2014). Characterisation of major histocompatibility complex class I in
the Australian cane toad, Rhinella marina. *PLoS One, 9*(8), e102824.
doi:10.1371/journal.pone.0102824
Lopez, J. V., Yuhki, N., Masuda, R., Modi, W., & O'Brien, S. J. (1994). Numt, a recent transfer and
tandem amplification of mitochondrial DNA to the nuclear genome of the domestic cat. *J*
Mol Evol, 39(2), 174-190. doi:10.1007/BF00163806
Lowe, S., Browne, M., Boudjelas, S., & De Poorter, M. (2000). 100 of the World's Worst Invasive Alien
Species: A Selection From the Global Invasive Species Database. *Published by The Invasive*
Species Specialist Group (ISSG) a specialist group of the Species Survival Commission (SSC) of
the World Conservation Union (IUCN), 12pp. First published as special lift-out in Aliens, 12.
Mainka, S. A., & Howard, G. W. (2010). Climate change and invasive species: double jeopardy. *Integr*
Zool, 5(2), 102-111. doi:10.1111/j.1749-4877.2010.00193.x
McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., . . . DePristo, M. A.
(2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Res, 20*(9), 1297-1303.
doi:10.1101/gr.107524.110
Meiklejohn, C. D., Montooth, K. L., & Rand, D. M. (2007). Positive and negative selection on the
mitochondrial genome. *Trends Genet, 23*(6), 259-263. doi:10.1016/j.tig.2007.03.008
Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler, A., &
Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference
in the Genomic Era. *Mol Biol Evol, 37*(5), 1530-1534. doi:10.1093/molbev/msaa015

Mittan-Moreau, C. S., Kelehear, C., Toledo, L. F., Bacon, J., Guayasamin, J. M., Snyder, A., & Zamudio,
K. R. (2022). Cryptic lineages and standing genetic variation across independent cane toad
introductions. *Molecular Ecology, 31*(24), 6440-6456. doi:10.1111/mec.16713
Molloy, E. K., & Warnow, T. (2018). To Include or Not to Include: The Impact of Gene Filtering on
Species Tree Estimation Methods. *Syst Biol, 67*(2), 285-303. doi:10.1093/sysbio/syx077
Nei, M., & Tajima, F. (1981). Genetic drift and estimation of effective population size. *Genetics,*
98(3), 625-640. doi:10.1093/genetics/98.3.625
Pereyra, M., Blotto, B. L., Baldo, D., Chaparro, J. C., Ron, S., Elias-Costa, A., . . . Faivovich, J. (2021).
Evolution in the Genus Rhinella: A Total Evidence Phylogenetic Analysis of Neotropical True
Toads (Anura: Bufonidae). *Bulletin of the American Museum of Natural History, 447*, 1–156.
doi:10.1206/0003-0090.447.1.1
Phillips, B. L., Brown, G. P., & Shine, R. (2010). Evolutionarily accelerated invasions: the rate of
dispersal evolves upwards during the range advance of cane toads. *J Evol Biol, 23*(12), 2595-
2601. doi:10.1111/j.1420-9101.2010.02118.x
Polzin, T., & Vahdati Daneshmand, S. (2003). On Steiner Trees and Minimum Spanning Trees in
Hypergraphs. *Operations Research Letters, 31*, 12-20. doi:10.1016/S0167-6377(02)00185-2
Poplin, R., Ruano-Rubio, V., DePristo, M. A., Fennell, T. J., Carneiro, M. O., Auwera, G. A. V. d., . . .
Banks, E. (2018). Scaling accurate genetic variant discovery to tens of thousands of samples.
bioRxiv, 201178. doi:10.1101/201178
Porath, H. T., Knisbacher, B. A., Eisenberg, E., & Levanon, E. Y. (2017). Massive A-to-I RNA editing is
common across the Metazoa and correlates with dsRNA abundance. *Genome Biol, 18*(1),
185. doi:10.1186/s13059-017-1315-y
Radford, I. J., Woolley, L.-A., Dickman, C. R., Corey, B., Trembath, D., & Fairman, R. (2019). Invasive
species-driven trophic cascades: Are cane toads indirectly contributing to small mammal
collapses across tropical Australia? *bioRxiv*, 616771. doi:10.1101/616771
Ramirez-Soriano, A., Ramos-Onsins, S. E., Rozas, J., Calafell, F., & Navarro, A. (2008). Statistical power
analysis of neutrality tests under demographic expansions, contractions and bottlenecks
with recombination. *Genetics, 179*(1), 555-567. doi:10.1534/genetics.107.083006
Ramos-Onsins, S. E., & Rozas, J. (2002). Statistical properties of new neutrality tests against
population growth. *Mol Biol Evol, 19*(12), 2092-2100.
doi:10.1093/oxfordjournals.molbev.a004034
Richardson, M. F., Sequeira, F., Selechnik, D., Carneiro, M., Vallinoto, M., Reid, J. G., . . . Rollins, L. A.
(2018). Improving amphibian genomic resources: a multitissue reference transcriptome of
an iconic invader. *Gigascience, 7*(1), 1-7. doi:10.1093/gigascience/gix114
Rollins, L. A., Moles, A. T., Lam, S., Buitenwerf, R., Buswell, J. M., Brandenburger, C. R., . . . Sherwin,
W. B. (2013). High genetic diversity is not essential for successful introduction. *Ecology and*
Evolution, 3(13), 4501-4517. doi:https://doi.org/10.1002/ece3.824
Rollins, L. A., Richardson, M. F., & Shine, R. (2015). A genetic perspective on rapid evolution in cane
toads (Rhinella marina). *Mol Ecol, 24*(9), 2264-2276. doi:10.1111/mec.13184
Roure, B., Baurain, D., & Philippe, H. (2013). Impact of missing data on phylogenies inferred from
empirical phylogenomic data sets. *Mol Biol Evol, 30*(1), 197-214.
doi:10.1093/molbev/mss208
Rozas, J., Ferrer-Mata, A., Sanchez-DelBarrio, J. C., Guirao-Rico, S., Librado, P., Ramos-Onsins, S. E., &
Sanchez-Gracia, A. (2017). DnaSP 6: DNA Sequence Polymorphism Analysis of Large Data
Sets. *Mol Biol Evol, 34*(12), 3299-3302. doi:10.1093/molbev/msx248
Sakai, A. K., Allendorf, F. W., Holt, J. S., Lodge, D. M., Molofsky, J., With, K. A., . . . Weller, S. G. (2001).
The Population Biology of Invasive Species. *Annual Review of Ecology and Systematics, 32*(1),
305-332. doi:10.1146/annurev.ecolsys.32.081501.114037

Schneider, S., & Excoffier, L. (1999). Estimation of past demographic parameters from the
distribution of pairwise differences when the mutation rates vary among sites: application to
human mitochondrial DNA. *Genetics, 152*(3), 1079-1089. doi:10.1093/genetics/152.3.1079
Schrieber, K., & Lachmuth, S. (2017). The Genetic Paradox of Invasions revisited: the potential role of
inbreeding x environment interactions in invasion success. *Biol Rev Camb Philos Soc, 92*(2),
939-952. doi:10.1111/brv.12263
Seebens, H., Blackburn, T. M., Dyer, E. E., Genovesi, P., Hulme, P. E., Jeschke, J. M., . . . Essl, F. (2017).
No saturation in the accumulation of alien species worldwide. *Nat Commun, 8*, 14435.
doi:10.1038/ncomms14435
Selechnik, D., Richardson, M. F., Shine, R., Brown, G. P., & Rollins, L. A. (2019). Immune and
environment-driven gene expression during invasion: An eco-immunological application of
RNA-Seq. *Ecol Evol, 9*(11), 6708-6721. doi:10.1002/ece3.5249
Selechnik, D., Richardson, M. F., Shine, R., DeVore, J. L., Ducatez, S., & Rollins, L. A. (2019). Increased
Adaptive Variation Despite Reduced Overall Genetic Diversity in a Rapidly Adapting Invader.
Front Genet, 10, 1221. doi:10.3389/fgene.2019.01221
Shavit Grievink, L., Penny, D., & Holland, B. R. (2013). Missing data and influential sites: choice of
sites for phylogenetic analysis can be as important as taxon sampling and model choice.
Genome Biol Evol, 5(4), 681-687. doi:10.1093/gbe/evt032
Shine, R. (2010). The ecological impact of invasive cane toads (Bufo marinus) in Australia. *Q Rev Biol,*
85(3), 253-291. doi:10.1086/655116
Shine, R., Alford, R. A., Blennerhasset, R., Brown, G. P., DeVore, J. L., Ducatez, S., . . . Phillips, B. L.
(2021). Increased rates of dispersal of free-ranging cane toads (Rhinella marina) during their
global invasion. *Sci Rep, 11*(1), 23574. doi:10.1038/s41598-021-02828-5
Shine, R., Brown, G. P., & Phillips, B. L. (2011). An evolutionary process that assembles phenotypes
through space rather than through time. *Proc Natl Acad Sci U S A, 108*(14), 5708-5711.
doi:10.1073/pnas.1018989108
Silva, G., Lima, F. P., Martel, P., & Castilho, R. (2014). Thermal adaptation and clinal mitochondrial
DNA variation of European anchovy. *Proc Biol Sci, 281*(1792). doi:10.1098/rspb.2014.1093
Slade, R. W., & Moritz, C. (1998). Phylogeography of Bufo marinus from its natural and introduced
ranges. *Proc Biol Sci, 265*(1398), 769-777. doi:10.1098/rspb.1998.0359
Song, H., Buhay, J. E., Whiting, M. F., & Crandall, K. A. (2008). Many species in one: DNA barcoding
overestimates the number of species when nuclear mitochondrial pseudogenes are
coamplified. *Proc Natl Acad Sci U S A, 105*(36), 13486-13491. doi:10.1073/pnas.0803076105
St Laurent, G., Tackett, M. R., Nechkin, S., Shtokalo, D., Antonets, D., Savva, Y. A., . . . Reenan, R. A.
(2013). Genome-wide analysis of A-to-I RNA editing by single-molecule sequencing in
Drosophila. *Nat Struct Mol Biol, 20*(11), 1333-1339. doi:10.1038/nsmb.2675
Tang, W., Fei, Y., & Page, M. (2012). Biological Significance of RNA Editing in Cells. *Molecular*
Biotechnology, 52(1), 91-100. doi:10.1007/s12033-012-9498-7
Tarailo-Graovac, M., & Chen, N. (2009). Using RepeatMasker to identify repetitive elements in
genomic sequences. *Curr Protoc Bioinformatics, Chapter 4*, 4 10 11-14 10 14.
doi:10.1002/0471250953.bi0410s25
Toews, D. P., & Brelsford, A. (2012). The biogeography of mitochondrial and nuclear discordance in
animals. *Mol Ecol, 21*(16), 3907-3930. doi:10.1111/j.1365-294X.2012.05664.x
Turvey, N. (2013). *Cane toads: a tale of sugar, politics and flawed science*: Sydney University Press.
Urban, M. C., Phillips, B. L., Skelly, D. K., & Shine, R. (2008). A toad more traveled: the heterogeneous
invasion dynamics of cane toads in Australia. *Am Nat, 171*(3), E134-148. doi:10.1086/527494
Van der Auwera, G. A. (2020). *Genomics in the cloud : using Docker, GATK, and WDL in Terra* (First
edition. ed.). Sebastopol, CA: Sebastopol, CA : O'Reilly Media.

Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., . . . Earl, A. M. (2014).
Pilon: an integrated tool for comprehensive microbial variant detection and genome
assembly improvement. *PLoS One, 9*(11), e112963. doi:10.1371/journal.pone.0112963
Wang, X., Zhou, S., Wu, X., Wei, Q., Shang, Y., Sun, G., . . . Zhang, H. (2021). High-altitude adaptation
in vertebrates as revealed by mitochondrial genome analyses. *Ecol Evol, 11*(21), 15077-
15084. doi:10.1002/ece3.8189
Wei, W., Schon, K. R., Elgar, G., Orioli, A., Tanguy, M., Giess, A., . . . Chinnery, P. F. (2022). Nuclear-
embedded mitochondrial DNA sequences in 66,083 human genomes. *Nature, 611*(7934),
105-114. doi:10.1038/s41586-022-05288-7
Wiens, J. J., & Morrill, M. C. (2011). Missing data in phylogenetic analysis: reconciling results from
simulations and empirical data. *Syst Biol, 60*(5), 719-731. doi:10.1093/sysbio/syr025
Wright, E. S. (2016). Using DECIPHER v2.0 to analyze big biological sequence data in R. *R Journal, 8*,
352-359.
Wright, E. S. (2020). RNAconTest: comparing tools for noncoding RNA multiple sequence alignment
based on structural consistency. *RNA, 26*(5), 531-540. doi:10.1261/rna.073015.119
Xia, X. (2014). *Phylogenetic Bias in the Likelihood Method Caused by Missing Data Coupled with*
Among-Site Rate Variation: An Analytical Approach (Vol. 8492).
Xiao, S., Nguyen, D. T., Wu, B., & Hao, W. (2017). Genetic Drift and Indel Mutation in the Evolution of
Yeast Mitochondrial Genome Size. *Genome Biol Evol, 9*(11), 3088-3099.
doi:10.1093/gbe/evx232
Yagound, B., West, A. J., Richardson, M. F., Selechnik, D., Shine, R., & Rollins, L. A. (2022). Brain
transcriptome analysis reveals gene expression differences associated with dispersal
behaviour between range-front and range-core populations of invasive cane toads in
Australia. *Mol Ecol, 31*(6), 1700-1715. doi:10.1111/mec.16347
Yu, G., Smith, D. K., Zhu, H., Guan, Y., & Lam, T. T. Y. (2017). ggtree: an R package for visualization
and annotation of phylogenetic trees with their covariates and other associated data.
Methods in Ecology and Evolution, 8(1), 28-36. doi:10.1111/2041-210x.12628
Zaranek, A. W., Levanon, E. Y., Zecharia, T., Clegg, T., & Church, G. M. (2010). A survey of genomic
traces reveals a common sequencing error, RNA editing, and DNA editing. *PLoS Genet, 6*(5),
e1000954. doi:10.1371/journal.pgen.1000954
Zhang, P., Liang, D., Mao, R. L., Hillis, D. M., Wake, D. B., & Cannatella, D. C. (2013). Efficient
sequencing of Anuran mtDNAs and a mitogenomic exploration of the phylogeny and
evolution of frogs. *Mol Biol Evol, 30*(8), 1899-1915. doi:10.1093/molbev/mst091
Zug, G. R., & Zug, P. B. (1979). The marine toad, Bufo marinus : a natural history resumé of native
populations. doi:https://doi.org/10.5479/si.00810282.284

Figure 1: Sampling locations of the cane toad (*Rhinella marina*) in (A) Australia and (B) French Guiana (the native range) and (C) Hawai'i. Red
dots in French Guiana, Hawai'i and Australia indicate sample collection sites.

Figure 2: Mitochondrial genome structure of *R. marina*. Genes, GC skew and GC content
were shown in the inner circle, middle circle and outer circle respectively. The innermost
circle indicates the negative strand and the outer circle indicates the positive strand. Genes
are coloured by type: red, tRNA; pink, rRNA; blue, coding sequence. Starting and stop
positions of each gene are described in Table S3.

Figure 3: Maximum-likelihood phylogenetic tree of the complete mitochondrial genomes of 203 Anuran species downloaded from the NCBI

BioProject public database. Five non-Anuran species from Caudata and Caeciliidae were used as outgroup (Table S3). The dendrogram was

constructed using IQ-Tree multicore version 2.2.2.6. Branch lengths represent genetic distances. Ultrafast bootstrap and SH-aLRT branch tests

were performed with 10,000 replications. Branches with SH-aLRT values ≤ 80% or UFboot values ≤ 95% were labelled with red dots. The

leftmost and rightmost panels display zoomed upper and lower parts of the dendrogram, respectively. The cane toad (*Rhinella marina*)

labelled with a red arrow.

Figure 4: The haplotype network illustrating the relationships among 132 individuals from French Guinea, Hawai'i, and Australia based on near-
complete mitochondrial genomes. Each sample location is represented by a different colour. A black solid circle represents a putative
haplotype that was not directly sampled. The red number adjacent to a line indicates the number of nucleotide substitutions between
haplotypes, while the absence of a number indicates a single nucleotide substitution.

Figure 5: Haplotype network of a 510 bp segment of ND3 with flanking tRNAs, a total of 223 samples, including samples from this study, Slade

and Moritz (1998) and Acevedo et al. (2016). Sample locations are represented by different colours. A black solid circle represents a putative

haplotype that was not directly sampled. The red number adjacent to a line indicates the number of nucleotide substitutions between

haplotypes, while the absence of a number indicates a single nucleotide substitution. Samples are separated by geographical locations on each

side of the Andes Mountains in South America (Acevedo et al., 2016). Toads on the west side of the Andes are now known as *Rhinella*

horribilis.

Figure 6: Haplotype diversity (A) and nucleotide diversities (B and C) of native population and introduced populations.

Table 1: Population genetics statistics of native range population (French Guiana) and introduced populations (Hawai'i and Australia).

	N	Polymorphic sites	*H*	π ± SD	*h* ± SD	HR
French Guiana	13	205	12	0.00469 ± 0.00037	0.987 ± 0.035	12
Hawai'i	25	11	13	0.00010 ± 0.00002	0.880 ± 0.051	10.7
Australia	94	40	36	0.00009 ± 0.00001	0.834 ± 0.035	12.5
Queensland	42	27	23	0.00012 ± 0.00001	0.890 ± 0.040	13.5
Northern Territory	20	5	7	0.00005 ± 0.00001	0.711 ± 0.089	6.4
Western Australia	32	16	13	0.00008 ± 0.00001	0.809 ± 0.06	9.6

N: Number of samples; *H*: Number of haplotypes; π: nucleotide diversity; *h*: haplotype diversity; HR; haplotype richness.

964   Table 2: Analysis of molecular variance (AMOVA) of all populations.
965

Source of variation	*df*	Sum of squares	% of variation	Fixation indices
All populations				
Among groups	2	482.96	64.49	
Among populations within groups	2	2.91	-0.82	
Within populations	127	579.35	36.33	$F_{ST} = 0.64$***

***: $p < 0.001$

967 Table 3: Population differentiation (pairwise *Fst*) between native range and introduced ranges (Hawai'i and Australia) (below the diagonal line)
968 The associated *p*-values were shown above the diagonal line.

	French Guiana	Hawai'i	Queensland	Northern Territory	Western Australia	Australia
French Guiana		< 0.001	< 0.001	< 0.001	< 0.001	< 0.001
Hawai'i	0.58		0.06	0.01	0.01	0.02
Queensland	0.67	0.02		0.009	0.003	0.34
Northern Territory	0.55	0.05	0.04		0.72	0.54
Western Australia	0.63	0.03	0.03	-0.01		0.8
Australia	0.79	0.02	0.002	-0.004	-0.006	

970 Table 4: Demographic estimators of native range population (French Guiana) and introduced populations (Hawai'i and Australia).

	Fu's *Fs*	R_2	Demographic expansion		Spatial expansion	
			SSD	Raggedness	SSD	Raggedness
French Guiana	1.29 (p = 0.66)	0.18 (p = 0.891)	0.03 (p = 0.19)	0.03 (p = 0.39)	0.03 (p = 0.08)	0.03 (p = 0.55)
Hawai'i	-8.78 (p < 0.001)	0.08 (p = 0.02)	0.01 (p = 0.44)	0.080 (p = 0.24)	0.01 (p = 0.31)	0.08 (p = 0.21)
Australia	-27.92 (p < 0.001)	0.02 (p < 0.001)	0.002 (p = 0.30)	0.067 (p = 0.16)	0.002 (p = 0.20)	0.07 (p = 0.14)
Queensland	-23.04 (p < 0.001)	0.03 (p < 0.001)	0.001 (p = 0.70)	0.05 (p = 0.38)	0.002 (p = 0.66)	0.05 (p = 0.42)
Northern Territory	-3.91 (p < 0.001)	0.10 (p = 0.072)	0.03 (p = 0.07)	0.21 (p = 0.03)	0.03 (p = 0.03)	0.21 (p = 0.05)
Western Australia	-8.950 (p < 0.001)	0.05 (p = 0.001)	0.003 (p = 0.58)	0.08 (p = 0.34)	0.003 (p= 0.35)	0.08 (p = 0.26)