Massive-scale single-cell chromatin accessibility sequencing using combinatorial fluidic indexing

Xuan Zhang[1], Alexandre P Marand[1,2], Haidong Yan[1], Robert J Schmitz[1*]

[1]Department of Genetics, University of Georgia, Athens, GA, USA.

[2]Current address: Department of Molecular, Cellular, and Development Biology, University of Michigan, Ann Arbor, MI, USA

* Correspondence

Contact:
Xuan Zhang: xuanzhang@uga.edu
Alexandre P Marand: amarand@umich.edu
Haidong Yan: hy17471@uga.edu
Robert J Schmitz: schmitz@uga.edu

Abstract

Single-cell ATAC-seq has emerged as a powerful approach for revealing candidate *cis*-regulatory elements genome-wide at cell-type resolution. However, current single-cell methods suffer from limited throughput and high costs. Here, we present a novel technique called single-cell combinatorial fluidic indexing ATAC-sequencing ("scifi-ATAC-seq"), which combines a barcoded Tn5 pre-indexing step with droplet-based single-cell ATAC-seq using a widely commercialized microfluidics platform (10X Genomics). With scifi-ATAC-seq, up to 200,000 nuclei across multiple samples in a single emulsion reaction can be indexed, representing a ~20-fold increase in throughput compared to the standard 10X Genomics workflow.

Main

ATAC-seq (Assay for Transposase-Accessible Chromatin using sequencing) [1] has emerged as a popular method for chromatin accessibility profiling to unveil genome-wide candidate *cis*-regulatory elements, which govern precise gene expression patterns for specifying distinct cell types or cell states. Recently, ATAC-seq was further modified to profile chromatin accessibility at single-cell resolution (scATAC-seq), which was used to generate several *cis*-regulatory atlases for animals and plants [2-7]. Most single-cell methods have the capability to generate data for hundreds to thousands of cells/nuclei simultaneously and are categorized as either droplet-based or combinatorial indexed techniques [8-11]. Droplet-based approaches are commonly implemented using microfluidics platforms, such as the commercialized products from companies like 10X Genomics, which offer straightforward handling and consistent data quality compared with combinatorial indexing methods [12].

In droplet-based assays the single-cell suspension is loaded into the microfluidics device at a significantly lower concentration compared to the barcoded beads to minimize the occurrence of cell/nuclei doublets that lead to false cell states. For instance, typically the loading cell/nuclei number is only about 1-10% of the total number of Gel Bead-in-Emulsions (GEMs) from the 10X Genomics Chromium Controller (Fig. 1a), which uses reagents inefficiently, leading to

limited throughput and higher costs. To address the inefficiency of droplet-based approaches,
a chimeric single-cell strategy has been developed, in which nuclei were pre-indexed prior to
overloading of the microfluidics device. With this strategy, the technique known as
"dsciATAC-seq" was developed, which combined pre-indexing with a microfluidics system
from Bio-Rad, and recovered about 25k nuclei with 100k nuclei loaded in one experiment [13].
More recently, similar approaches have been used for scRNA-seq with the 10X Genomics
platform, referred to as scifi-RNA-seq, which recovered around 150k cells upon loading 380k
cells, and further significantly increased the throughput [14]. However, there is currently no
massive-scale scATAC-seq protocol for use with the 10X Genomics platform, which is most
commonly used for scATAC-seq data generation.

In this study, we present a method for profiling massive-scale single-cell chromatin
accessibility sequencing using the 10X Genomics microfluidics system. We refer to this
method as single-cell combinatorial fluidic indexing ATAC-sequencing (scifi-ATAC-seq), as
it was initially inspired by the scifi-RNA-seq design [14]. In scifi-ATAC-seq, the nuclei are
pre-indexed in a 96-well plate with a two-sided barcoded Tn5 (96 distinct barcode
combinations), which is based on our previous sci-ATAC-seq design [15]. Then, a standard
scATAC-seq library preparation is performed, except that an overloaded number of nuclei are
used in the microfluidics system. Approximately 100-200k nuclei per channel instead of the
recommended maximum of 15.3k are loaded. Compared to the one-sided barcoded Tn5 [13],
the two-sided barcode design offers several advantages: (i) It requires fewer Tn5 adapter oligos
for Tn5 barcoding, and readily accommodates scaling up the index complexity when necessary.
Only 20 oligos (8 rows x 12 columns) are needed to create 96 unique barcode combinations.
(ii) It requires less Tn5 for Tn5 assembly. Only 280 uL of Tn5 (15 uL in 8 rows and 10 uL in
12 columns) is necessary, whereas the one-sided barcode would require over 1,000 uL of Tn5,
assuming a minimal assembly volume of 10 uL per well. (iii) The barcode in the s5 end helps
to distinguish index hopping reads and reduces index hopping contamination (Extended Data
Fig. 1 a,b).

To evaluate the performance of scifi-ATAC-seq, we generated two scifi-ATAC-seq libraries
using 100k and 200k overloaded nuclei from frozen tissue with mixed genotypes (B73 and
Mo17) of *Zea mays* (maize) seedling tissue. We compared the scifi-ATAC-seq data with two
scATAC-seq libraries generated from the same tissue type using the standard 10X Genomics
workflow (referred to as scATAC-seq from this point onward) [4]. All datasets were processed
with the same parameters (see Methods). We assessed various quality control metrics,
including the proportion of reads around the transcription start site (TSS), fraction of reads in
peaks (FRiP), unique Tn5 insertions per cell, fragment size distribution, and organelle DNA
contamination (see Fig. 1c-e and Extended Data Fig. 1c). The scifi-ATAC-seq data showed
similar or superior data quality compared to the other datasets. Particularly, scifi-ATAC-seq
successfully recovered approximately 35k and 70k clean cells for the 100k and 200k input
nuclei, respectively, which represents a 9-fold to 18-fold increase in throughput compared to
the scATAC-seq (Fig. 1f). As expected, we observed a higher number of nuclei per droplet
when overloading the nuclei, resulting in an average of 1.57 nuclei per droplet for the 100k
library and 2.02 nuclei per droplet for the 200k library (Extended Data Fig. 1 d,f). These data
show that scifi-ATAC-seq produces high-quality libraries while increasing the number of
nuclei profiled.

Collecting multiple nuclei within a single droplet increases the risk of cell barcode collisions
and potential index hopping contamination (Extended Data Fig 1a). Next, we compared the
barcode collision rate for scATAC-seq and scifi-ATAC-seq with the same approach [4]. The

identified barcode collision rate is 4.87% and 5.68% for the 16k scATAC-seq and 100k scifi-
ATAC-seq, respectively (Fig. 1g,h). The total barcode collision should be around 10% for both
datasets considering the collisions of the same genotype, which is similar with the collision
rate reported in other studies [4]. The throughput increased about 9-fold with the 100k scifi-
ATAC-seq experiment with a similar barcode collision rate. The identified collision rate was
9.52% for the 200k scifi-ATAC-seq, which is higher than the 16k scATAC-seq dataset.
Although, the total recovered nuclei number is further increased to 69,302, which reflects an
18-fold increase in throughput (Fig. 1i). As expected, the total number of Tn5 insertions in
nuclei with barcode collisions is significantly higher than non-collision nuclei (see Extended
Data Fig 1e, g, q-value < 10e-16). Barcode collisions are typically removed using an array of
doublet detection tools [16-19], minimizing the effect of clustering and cell-type identification.
Lastly, the cross cell contamination was 1.47% and 1.69% for both scifi-ATAC-seq datasets,
which is much lower than the 5.63% observed in the 16k scATAC-seq dataset indicating this
could benefit from the double-side barcode design implemented in our method (Extended Data
Fig. 1 a,b). Collectively, these data show that scifi-ATAC-seq has similar barcode collision
rates as the standard scATAC-seq workflow, while producing data for significantly greater
numbers of nuclei.

To evaluate whether the differences in data quality and cell number could affect clustering, we
clustered all the nuclei from the scATAC-seq and scifi-ATAC-seq datasets together with
identical parameters using *Socrates* [4] and annotated the clusters using previously reported
methods. In total, we identified 14 clusters representing the major cell types in the maize
seedling (Fig. 1j, Extended Data Fig. 2a). All the resulting cell types identified were consistent
and found across multiple techniques (Extended Data Fig. 2b-d). The ability to identify and
characterize rare cell populations in detail is a significant advantage of single-cell technologies,
which requires profiling a substantial number of cells within each tissue. Here, we profiled
more than 100k nuclei in the seedling tissue with scifi-ATAC-seq, which provides an
opportunity to study rarer cell types. In maize seedlings, bundle sheath and mesophyll represent
major cell types for photosynthesis and typically represent ~20% of cell types in seedling tissue,
whereas vascular cells, such as phloem, procambium, and xylem, responsible for nutrient
transport, are present in much lower proportions (Extended Data Fig. 3a-d). We observed a
similar profile for bundle sheath and mesophyll cells in both scifi-ATAC-seq and the scATAC-
seq data (Extended Data Fig. 3e, f). However, for vascular cells, the resolution of accessible
chromatin was difficult to discern in the scATAC-seq data, but were clearly visible in both
scifi-ATAC-seq datasets (Fig. 1k, l, Extended Data Fig. 3g-i). Scifi-ATAC-seq data leads to
greater number of profiled nuclei per sample increasing the chances of studying rarer cell
populations.

To assess the robustness of scifi-ATAC-seq in profiling single-cell chromatin accessibility
across multiple samples within a single reaction, we implemented a multiplexing strategy using
maize seedlings from eight samples with seven different genotypes in a 96-well plate (Fig. 2a).
With seven different genotypes in this assay, the likelihood of barcode collisions within the
same genotype is expected to be low, as most barcode collisions occurred between distinct
genotypes and can be identified through computational methods. Thus, increasing the number
of loaded nuclei, while maintaining a low barcode collision rate within the same genotype, can
further enhance throughput. To test this, a total of 300k nuclei were used to prepare the scifi-
ATAC-seq library. We applied the same nuclei quality control criteria as before and
successfully recovered 199,212 high-quality nuclei. We identified 133,524 singlet nuclei with
a low index hopping contamination rate of 1.93% (Fig. 2b, c, Extended Data Fig. 4). The
average identified barcode collision rate was 31.55%, indicating that the final barcode collision

rate with the same genotype would be less than 5.26% (Fig. 2c, Extended Data Fig. 4d-k,
Methods). These results demonstrate that the throughput was further enhanced with scifi-
ATAC-seq by approximately 27-fold compared to the scATAC-seq method, while maintaining
a similar barcode collision rate and data quality.

Subsequently, we clustered all the singlet nuclei from this assay and annotated the clusters
using the same method as before. We identified all the cell types observed in the previous
dataset and observed that there was no significant bias in cell-type proportions among the
genotypes (Fig. 2d, e, Extended Data Fig. 5d). With this dataset, we identified 79,092
accessible chromatin regions, with approximately 30.35% showing cell-type-specific patterns
and 12.67% showing genotype-specific patterns (Fig. 2f, i). More specifically, vascular cells
(procambium, phloem, and xylem) were consistently identified in all samples (Fig. 2g,
Extended Data Fig. 6a-c), and genotype-specific accessible chromatin regions at cell-type
resolution were discernible (Fig. 2j, Extended Data Fig. d).

In summary, these results collectively show that scifi-ATAC-seq provides a robust, efficient,
and flexible approach for massive-scale single-cell chromatin accessibility profiling using the
widely available 10X Genomics Chromium systems. We anticipate that this method will
facilitate the utilization of ATAC-seq to identify candidate *cis*-regulatory elements at cell-type
resolution in greater numbers of tissues, time point, genotypes and facilitate the study of rarer
cell types.

Material and Methods.

Plant material and growth conditions
The maize kernels used in this study were obtained from the USDA National Plant Germplasm System
(https://npgsweb.ars-grin.gov). The seedlings were grown in Sungro Horticulture professional growing
mix (Sungro Horticulture Canada) under controlled conditions. The soil was saturated with tap water
and the seedlings were exposed to a mixture of 4,100 K (Sylvania Supersaver Cool White Delux
F34CWX/SS, 34 W) and 3,000 K (GE Ecolux with starcoat, F40CX30ECO, 40 W) light, with a
photoperiod of 16 hours of light and 8 hours of darkness. The temperature was maintained at
approximately 25°C during the light hours, and the relative humidity was approximately 54%. The
above-ground seedling tissues were harvested between 8 and 9 AM, six days after sowing. Flash-frozen
seedling tissue was used to generate scifi-ATAC-seq libraries for B73 and Mo17 mixed genotypes. And
fresh seedling tissue were used for scifi-ATAC-seq library with seven different maize genotypes.
170
171 **scifi-ATAC-seq protocol**
172 A detailed step-by-step sci-ATAC-seq protocol with lists of necessary reagents and equipment are
173 included in the Supplementary Methods. The Tn5 expression and purification steps were carried out
174 according to the method described by Tu et al. (2020) [20], and the plasmids were obtained from
175 Addgene (accession number 127916).
176
Assembly of indexed Tn5 transposase complexes
To generate indexed Tn5 transposase complexes, we modified the Tn5-ME-A and Tn5-ME-B by adding
a 5-nucleotide barcode (with 12 distinct barcodes for Tn5-ME-A and 8 distinct barcodes for Tn5-ME-
B; see Supplementary Methods Supplementary Table 1 for barcode sequences). Each indexed
oligonucleotide was annealed to a 19-bp complementary mosaic-end oligonucleotide (Tn5-ME-rev, 5′
phosphorylated, Supplementary Methods). The annealing reaction was carried out by mixing the
oligonucleotides at a 1:1 molar ratio (Tn5-ME-A or Tn5-ME-B: Tn5-ME-rev) at a final concentration
of 100 μM. The mixture was heated to 95 °C, cooled gradually to 20 °C at a rate of -1 °C per minute,

and then held at 20 °C. The annealed oligonucleotides were then mixed at a 1:1 molar ratio (Tn5-ME-
A: Tn5-ME-B). To assemble the transposase, 10 μL of Tn5 transposase was added to 0.143 μL of the
annealed adapter mixture, mixed well by pipetting slowly, incubated at room temperature for 60 minutes
and then storing it at -20°C until the tagmentation reactions were performed [21].
189
Nuclei isolation with quick purification
Approximately 3-4 maize seedlings were chopped on ice for about 2 minutes in 600 μL of pre-chilled
Nuclei Isolation Buffer (NIB cutting, 10 mM MES-KOH pH 5.4, 10 mM NaCl, 250 mM sucrose, 0.1
mM spermine, 0.5 mM spermidine, 1 mM DTT, 1% BSA, 0.5% TritonX-100), which was modified
from the original buffer composition [22]. After chopping, the total mixture was filtered with a 40-μm
cell strainer and then centrifuged at 500 rcf for 5 minutes at 4 ˚C. The supernatant was carefully removed,
and the pellet was resuspended in 500 μL of NIB wash buffer (10 mM MES-KOH, pH 5.4, 10 mM
NaCl, 250 mM sucrose, 0.1 mM spermine, 0.5 mM spermidine, 1 mM DTT, and 1% BSA). The sample
was filtered with a 20-μm cell strainer and then carefully loaded onto the surface of 1 mL 35% Percoll
buffer (made by mixing 35% Percoll and 65% NIB wash buffer) in a 1.5-mL centrifuge tube. The nuclei
were centrifuged at 500 rcf for 10 minutes at 4 ˚C. After centrifugation, the supernatant was carefully
removed, and the pellets were washed once in 100 μL TAPS buffer (25 mM TAPS-NaOH, pH 8.0, and
12.5 mM MgCl$_2$) and then resuspended in 30 μL of 2.5x TAPS buffer. About 5 μL of nuclei were
diluted 10 times and stained with DAPI (Sigma Cat. D9542). The nuclei quality and density were
evaluated with a hemocytometer under a microscope. Finally, equal nuclei number of B73 and Mo17
were mixed together and the nuclei density was adjusted to 0.5k~1k/μL with TAPS buffer.
206
Indexed Tn5 tagmentation and pooling
To generate a combination of 96 indexed transposases, 1.5 μL of Tn5-ME-A with 12 distinct barcodes
were dispensed by rows, and 1.5 μL of Tn5-ME-B with 8 distinct barcodes were dispensed by columns
in a 96-well plate. Each well had a unique combination of A and B indexed Tn5. To each well, 10 μL
of nuclei in TAPS buffer with 0.1% Tween 20 and 0.01% digitonin was added, and the plate was sealed.
The tagmentation reaction was carried out for 60 minutes at 37 ˚C. The reaction was stopped by adding
12 μL of stop buffer (10 mM Tris-HCl pH 7.8, 20 mM EDTA, pH 8.0, 2% BSA) supplemented with
ethylenediaminetetraacetic acid (EDTA) to quench the Mg2+. All nuclei were transferred to a reservoir
and then divided into two 1.5 mL centrifuge tubes. The nuclei were pelleted, resuspended in 200 μL
diluted nuclei buffer (DNB, 10x Genomics Cat#2000207), filtered with a 40 um strainer, pooled into
one PCR tube, and centrifuged at 500 rcf for 2 min at 4˚C. After centrifugation, the supernatant was
carefully removed, and the nuclei (approximately 3 μL) were resuspended in 5 μL of DNB and 7 μL
ATAC buffer B (10x Genomics Cat#2000193).
220
Library preparation and sequencing
scATAC-seq libraries were prepared using the Chromium scATAC v1.1 (Next GEM) kit from
10xGenomics,   following   the   manufacturer's   instructions.   (10xGenomics,
CG000209_Chromium_NextGEM_SingleCell_ATAC_ReagentKits_v1.1_UserGuide_RevE).   The
leftover nuclei after loading to ChIP-H were diluted, stained with DAPI, and nuclei quality and density
was evaluated with a hemocytometer under a microscope. The final libraries were sequenced using an
Illumina NovaSeq 6000 S4 in the dual-index mode using custom sequencing primer sets
(Supplementary Methods). To balance the nucleotide distribution at the beginning of the forward and
reverse reads, the proportion of the scifi-ATAC-seq library in a lane should be less than 50% or extra
spike-in library (e.g., PhiX control from Illumina) should be added to the lane.
231
232
Raw reads processing and alignment
During the preprocessing of all single-cell ATAC-seq data, the 16-bp i5 beads barcode was added to
the read names of the paired-end reads using the extract function from UMItools v.1.01 [23]. The
customization parameter '--bc-pattern=NNNNNNNNNNNNNNNN' was used for this process.
Moreover, for scifi-ATAC-seq data, the inline Tn5 barcode was demultiplexed and subsequently
appended to the read names using cutadapt v3.4 [24]. Next, the processed reads were aligned to the *Zea*
mays reference genome v5 [25] using BWA-MEM v0.7.17 [26]. To obtain high-quality, properly paired,

and unique alignments, the view function from samtools v1.9 [27] was applied with the parameters '-q
10 -f 3.' Additionally, reads with XA tags were filtered out. Subsequently, the cell barcodes were
included in the alignments using the CB tag and BC tag for the 10X Genomics scATAC-seq and
scifiATAC-seq datasets, respectively. To eliminate duplicate reads, Picard Tools v.2.21.6
(http://broadinstitute.github.io/picard/) was employed while considering the cell barcode. Finally, the
alignments were converted to single base-pair Tn5 integration sites in a BED format by adjusting the
start coordinates of the forward and reverse strands by +4 and -5, respectively. Only unique Tn5
insertion sites within a cell were retained for downstream analysis.
248
Nuclei calling and quality control
The R package *Socrates* [4] was utilized for nuclei identification and quality control. In summary, the
BED file containing single base-pair Tn5 integration sites was imported into *Socrates* along with the
Zea mays v5 GFF gene annotation and the genome index file. The scaffolds ('scaf_23', 'scaf_34', and
'scaf_36') were considered as organelle genomes. To identify bulk-scale ACRs (Accessible Chromatin
Regions) in *Socrates*, the callACRs function was employed with the following parameters: genome size
= 8.5e8, shift = -75, extsize = 150, and FDR = 0.1. This step allowed us to estimate the fraction of Tn5
integration sites located within ACRs for each nucleus. Metadata for each nucleus were collected using
the buildMetaData function, using a TSS (Transcription Start Site) window size of 2 kb (tss.window =
2000). Subsequently, sparse matrices were generated with the generateMatrix function, using a window
size of 500. High-quality nuclei were identified based on the following criteria: a minimum of 1,000
Tn5 insertion sites per nucleus, at least 20% of Tn5 insertions within 2 kb of TSSs, and at least 20% of
Tn5 insertions within ACRs across all datasets. Additionally, a maximum of 30% of Tn5 insertions in
organelle genomes was allowed.
263
Cell barcode collision detection
Cell barcode collisions were identified using a previously described approach [4]. Specifically, the
known genotype data were obtained from Panzea[28] and lifted over to v5 genome [25] coordinates
using CrossMap (v0.5.1) [29]. Only homozygous biallelic SNPs were retained for further analysis. The
Souporcell (git version 6872d88) pipeline [30] was used to count the number of reads for each genotype,
using known common variants and specific non-default parameters (--min_alt 50, --min_ref 50, --
max_loci 25000000, --skip_remap TRUE). In the dataset comprising a mixture of two genotypes (B73
and Mo17), genotype identification was performed by modeling allele counts as a binomial distribution,
accounting for a conservative sequencing error rate of 0.05. Posterior probabilities were estimated via
Bayes theorem to assign the genotype or identify cell barcode collisions (i.e., mixtures of genotypes)
with the highest probability. The index hopping contamination was calculated as the proportion of reads
that did not match the assigned genotype among all reads covering the biallelic SNPs. In the dataset
featuring a mixture of two genotypes, cell doublets can originate from either two cells of the same
genotype (A x A or B x B) or cells from different genotypes (A x B or B x A). However, the observed
collisions consist entirely of cell doublets from different genotypes, representing only approximately
half of the collision events that actually take place in the experiment. Therefore, there should be an
equal proportion of doublets mixed with nuclei of the same genotype.
For the 7-genotype-mixed scifi-ATAC-seq data, expected genotypes were assigned by matching known
Tn5 barcodes. Any reads that did not match the target genotype were considered as belonging to another
genotype. The same genotype calling approach described above was then used to assign nuclei to their
target genotype, identify mixtures of genotypes resulting from cell barcode collisions, and estimate the
index hopping contamination. Similarly, In the 7-genotype-mixed experiment, cell doublets can arise
from either two cells of the same genotype (A x A) or cells from different genotypes (A x 6 non-As).
The observed collisions involve different genotypes, which should account for approximately 6/7
(85.71%) of the collision events occurring in the experiment. Consequently, there should be a 1/7
(14.29%) proportion of doublets mixed with nuclei of the same genotype.
290
Integrated clustering analysis
The integrated clustering analysis of the four datasets, combining scifi-ATAC-seq and 10X Genomics
scATAC-seq data, was performed using the R package *Socrates* [4]. In brief, firstly, the ACRs were
identified by treating each library as a traditional bulk ATAC-seq library with function callACRs

(genomesize=8.5e8, shift= -75, extsize=150, fdr=0.1). Then a binary nucleus x ACR matrix was
generated with the function generateMatrix (peaks=T). The ACRs that were accessible in less than 0.15%
of all nuclei, and nuclei with less than 100 accessible ACRs were removed. Then the filtered nucleus x
ACR matrix were normalized with the term-frequency inverse-document-frequency (TF-IDF)
algorithm(doL2=T). The dimensionality of the normalized accessibility scores was reduced using the
function reduceDims (method="SVD", n.pcs = 25, cor.max =0.5). The reduced embedding was
visualized as a UMAP embedding using projectUMAP (k.near = 15). Approximately 5% of potential
cell doublets were identified and filtered by performing a modified version of the *Socrates* workflow
on each library separately with the function detectDoublets and filterDoublets (removeDoublets = T).
To address genotype and batch effects, we used the R package Harmony with non-default parameters
(do_pca=F, vars_use=c("library", "genotype"), tau=c(5), lambda=c(0.1, 0.1), nclust=50,
max.iter.cluster=100, max.iter.harmony=30). The dimensionality of the nuclei embedding was further
reduced with Uniform Manifold Approximation Projection (UMAP) via the R implementation of umap
(n_neighbors=30, metric = "cosine", a = 1.95, b = 0.75, ret_model=T). Finally, the nuclei were clustered
with function callClusters (res=0.4, k.near = 30, cl.method = 4, m.clust = 100).
A similar clustering process was applied to the 7-genotype-mixed scifi-ATAC-seq dataset with minor
modifications. Specifically, we removed only the genotype effect using Harmony, and the final clusters
were identified at a resolution of 0.5.
313
Cell-type annotation
To assign cell types for each cluster, we used a combination of marker gene-based annotation and gene
set enrichment analysis. Initially, we compiled a list of known cell-type-specific marker genes for maize
seedlings through an extensive literature review, primarily referring to Marrand et al., 2021
(Supplementary Table 5) [4]. Firstly, the gene chromatin accessibility score was calculated using the
Tn5 integration number in the gene body, 500-bp upstream and 100-bp downstream region, then the
raw counts were normalized with cpm function in edgeR. The Z-score was calculated for each marker
gene across all cell types with scale function in R, and key cell types were assigned based on the most
enriched marker genes with highest Z-score. Ambiguous clusters displaying similar patterns to key cell
types were assigned to the same cell type as the key cell types, reflecting potential variations in cell
states within a cell type. For gene set enrichment analysis, we used the R package fgsea [31], following
a methodology described previously[4]. Firstly, we constructed a reference panel by uniformly
sampling nuclei from each cluster, with the total number of reference nuclei set to the average number
of nuclei per cluster. Subsequently, we aggregated the read counts across nuclei in each cluster for each
gene and identified the differential accessibility profiles for all genes between each cluster and the
reference panel using the R package edgeR. For each cluster, we generated a gene list sorted in
decreasing order of the log2 fold-change value compared to the reference panel and utilized this list for
gene set enrichment analysis. We excluded GO terms with gene sets comprising less than 10 or greater
than 600 genes from the analysis and GO terms were considered significantly enriched at an FDR <
0.05 with 10,000 permutations. The cell type annotation was additionally validated by identifying the
top enriched GO terms that align with the expected cell type functions (Supplementary Table 6,7).
335
ACR identification
Following cell clustering and annotation, ACRs were further identified using all Tn5 integration sites
for each cell type and genotype with running MACS2 [32] with non-default parameters: --extsize 150
--shift -75 --nomodel --keep-dup all --qvalue 0.05. Then the cell type based ACRs for each genotype
were further redefined as 500-bp windows centered on the ACR coverage summit. To consolidate
information across all clusters and genotypes, we concatenated all ACRs into a unified master list using
a custom script, as previously described by Marrand et al., 2021 [4], calculated the ACR chromatin
accessibility score based on the Tn5 integration count within the ACR region and then normalized it
using the 'cpm' function in edgeR [33].
345
346 **Data availability**
347 All data supporting the results of this study are available within the article and supplementary
348 information files. The scifi-ATAC-seq data generated in this study have been deposited in the National

Center for Biotechnology Information Short Reads Archive (PRJNA996051).

Code availability

Acknowledgments

We would like to acknowledge Dr. Pedro Augusto Braga Dos Reis, Dr. Yinxin Dong, Yangyang Xu, for HS-Tn5 transposase purification; the GACRC for providing valuable assistance; and the Duke University School of Medicine for the use of the Sequencing and Genomic Technologies Shared Resource, which provided illumina sequencing service. This study was funded with support from the National Science Foundation (IOS-1856627 and IOS-2026554). This material is based upon work supported by the U.S. Department of Energy, Office of Science, Biological and Environmental Research Program under Award Number DE-SC0023338.

Author contributions

R.J.S. and X.Z. designed the research. X.Z. performed the experiments. A.P.M., H.Y., X.Z., and R.J.S. analyzed the data. R.J.S. and X.Z. wrote the manuscript. The authors read and approved the final manuscript.

Competing interests

R.J.S. is a co-founder of REquest Genomics, LLC, a company that provides epigenomic services. The remaining authors declare no competing interests.

1. Buenrostro, J.D., et al., *Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position.* Nat Methods, 2013. **10**(12): p. 1213-8.
2. Cusanovich, D.A., et al., *A Single-Cell Atlas of In Vivo Mammalian Chromatin Accessibility.* Cell, 2018. **174**(5): p. 1309-1324 e18.
3. Domcke, S., et al., *A human cell atlas of fetal chromatin accessibility.* Science, 2020. **370**(6518).
4. Marand, A.P., et al., *A cis-regulatory atlas in maize at single-cell resolution.* Cell, 2021. **184**(11): p. 3041-3055 e21.
5. Zhang, K., et al., *A single-cell atlas of chromatin accessibility in the human genome.* Cell, 2021. **184**(24): p. 5985-6001 e19.
6. Farmer, A., et al., *Single-nucleus RNA and ATAC sequencing reveals the impact of chromatin accessibility on gene expression in Arabidopsis roots at the single-cell level.* Mol Plant, 2021. **14**(3): p. 372-383.
7. Dorrity, M.W., et al., *The regulatory landscape of Arabidopsis thaliana roots at single-cell resolution.* Nat Commun, 2021. **12**(1): p. 3334.
8. Marand, A.P. and R.J. Schmitz, *Single-cell analysis of cis-regulatory elements.* Curr Opin Plant Biol, 2022. **65**: p. 102094.
9. Satpathy, A.T., et al., *Massively parallel single-cell chromatin landscapes of human immune cell development and intratumoral T cell exhaustion.* Nat Biotechnol, 2019. **37**(8): p. 925-936.
10. Cusanovich, D.A., et al., *Multiplex single cell profiling of chromatin accessibility by combinatorial cellular indexing.* Science, 2015. **348**(6237): p. 910-4.
11. Vitak, S.A., et al., *Sequencing thousands of single-cell genomes with combinatorial indexing.* Nat Methods, 2017. **14**(3): p. 302-308.
12. Ding, J., et al., *Systematic comparison of single-cell and single-nucleus RNA-sequencing methods.* Nat Biotechnol, 2020. **38**(6): p. 737-746.
13. Lareau, C.A., et al., *Droplet-based combinatorial indexing for massive-scale single-cell chromatin accessibility.* Nat Biotechnol, 2019. **37**(8): p. 916-924.
14. Datlinger, P., et al., *Ultra-high-throughput single-cell RNA sequencing and perturbation screening*

401   *with combinatorial fluidic indexing.* Nat Methods, 2021. **18**(6): p. 635-642.

402   15.   Tu, X., et al., *A combinatorial indexing strategy for low-cost epigenomic profiling of plant single*
403   *cells.* Plant Commun, 2022. **3**(4): p. 100308.

404   16.   Granja, J.M., et al., *ArchR is a scalable software package for integrative single-cell chromatin*
405   *accessibility analysis.* Nat Genet, 2021. **53**(3): p. 403-411.

406   17.   Germain, P.L., et al., *Doublet identification in single-cell sequencing data using scDblFinder.*
407   F1000Res, 2021. **10**: p. 979.

408   18.   Thibodeau, A., et al., *AMULET: a novel read count-based method for effective multiplet detection*
409   *from single nucleus ATAC-seq data.* Genome Biol, 2021. **22**(1): p. 252.

410   19.   Xi, N.M. and J.J. Li, *Benchmarking Computational Doublet-Detection Methods for Single-Cell RNA*
411   *Sequencing Data.* Cell Syst, 2021. **12**(2): p. 176-194 e6.

412   20.   Tu, X., et al., *Reconstructing the maize leaf regulatory network using ChIP-seq data of 104*
413   *transcription factors.* Nat Commun, 2020. **11**(1): p. 5089.

414   21.   Lu, Z., et al., *Combining ATAC-seq with nuclei sorting for discovery of cis-regulatory regions in*
415   *plant genomes.* Nucleic Acids Res, 2017. **45**(6): p. e41.

416   22.   Sikorskaite, S., et al., *Protocol: Optimised methodology for isolation of nuclei from leaves of species*
417   *in the Solanaceae and Rosaceae families.* Plant Methods, 2013. **9**: p. 31.

418   23.   Smith, T., A. Heger, and I. Sudbery, *UMI-tools: modeling sequencing errors in Unique Molecular*
419   *Identifiers to improve quantification accuracy.* Genome Res, 2017. **27**(3): p. 491-499.

420   24.   Martin, M., *Cutadapt removes adapter sequences from high-throughput sequencing reads.*
421   EMBnet. journal, 2011. **17**(1): p. 10-12.

422   25.   Hufford, M.B., et al., *De novo assembly, annotation, and comparative analysis of 26 diverse maize*
423   *genomes.* Science, 2021. **373**(6555): p. 655-662.

424   26.   Li, H. and R. Durbin, *Fast and accurate short read alignment with Burrows-Wheeler transform.*
425   Bioinformatics, 2009. **25**(14): p. 1754-60.

426   27.   Danecek, P., et al., *Twelve years of SAMtools and BCFtools.* Gigascience, 2021. **10**(2).

427   28.   Canaran, P., et al., *Panzea: an update on new content and features.* Nucleic Acids Res, 2008.
428   **36**(Database issue): p. D1041-3.

429   29.   Zhao, H., et al., *CrossMap: a versatile tool for coordinate conversion between genome assemblies.*
430   Bioinformatics, 2014. **30**(7): p. 1006-7.

431   30.   Heaton, H., et al., *Souporcell: robust clustering of single-cell RNA-seq data by genotype without*
432   *reference genotypes.* Nat Methods, 2020. **17**(6): p. 615-620.

433   31.   Korotkevich, G., et al., *Fast gene set enrichment analysis.* BioRxiv, 2016: p. 060012.

434   32.   Zhang, Y., et al., *Model-based analysis of ChIP-Seq (MACS).* Genome Biol, 2008. **9**(9): p. R137.

435   33.   Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor package for differential*
436   *expression analysis of digital gene expression data.* Bioinformatics, 2010. **26**(1): p. 139-40.

Figure 1. scifi-ATAC-seq combines pre-indexing with droplet-based scATAC-seq

(a) Schematic of regular droplet-based 10X Genomics scATAC-seq experimental workflow.

(b) Schematic of scifi-ATAC-seq experimental workflow.

(c) Distributions of the proportion of Tn5 integration sites within the promoter regions, encompassing the 2-kb flanking regions around gene transcription start sites (TSSs).

(d) Distributions of the proportion of Tn5 integration sites within peaks per nucleus.

(e) Distribution of unique Tn5 integration sites per nucleus.

(f) Number of nuclei that passed quality control thresholds.

(g-i) Scatterplot displaying SNP counts per cell for B73 and Mo17, color-coded by genotype classification. g, 16k input B73/Mo17 scATAC-seq; h, 100k scifi-ATAC-seq;
  i, 200k scifi-ATAC-seq. Median Contamination Rate: The median cross-contamination rate, attributed to index hopping, among all predicted singlets.

(j) UMAP of all nuclei (n = 98,424). Nuclei are colored by their predicted cell type.

(k) Pseudobulk cell type Tn5 integration site coverage around the phloem precursor marker ZmSMXL3. Vas. par. precursor: Vascular parenchyma precursor.

(l) Pseudobulk cell type Tn5 integration site coverage for and UMAP embeddings overlaid with gene chromatin accessibility around the *ZmSMXL3* gene across all datasets.
  N: Number of phloem nuclei.

Figure 2. Multiplexing eight samples with scifi-ATAC-seq

(a) Well assignment showing the multiplexing of primary samples and genotypes.

(b) Distribution of proportion of SNPs in target genotypes. Colored dots: singlet. Grey dots: identified cell barcode collision.
Median Contamination Rate: The median cross-contamination rate, attributed to index hopping, among all predicted singlets.

(c) Number of profiled nuclei for each genotype. Est. Collision: Estimated cell barcode collisions within the same genotype.

(d) UMAP of all nuclei (n = 124,656). Nuclei are colored by their predicted cell type. Vas. par. precursor: Vascular parenchyma precursor,
Procambium phloem pre.: Procambium phloem precursor.

(e) UMAP of all nuclei across the eight samples.

(f) Chromatin accessibility of cell-type-specific accessible chromatin regions (ctACRs, Z-score > 2) among all cell types.

(g) Pseudobulk cell type Tn5 integration site coverage around the phloem precursor marker *ZmSMXL3*.

(h) Pseudobulk cell type Tn5 integration site coverage and UMAP embeddings overlaid with gene chromatin accessibility around the *ZmSMXL3*.
N: Nuclei number of procambium phloem precursor.

(i) Chromatin accessibility of genotype specific accessible chromatin regions (gtACRs) among seven genotypes.

(j) Pseudobulk cell type Tn5 integration site coverage with gene chromatin accessibility around the *Zm00001eb280310* across all samples
in procambium phloem precursor cells.

Extended Data Figure 1. Evaluation and quality control of scifi-ATAC-seq, related to Figure1.

(a-b) Illustration of index hopping occurring during linear amplification in 10X Genomics Gel Beads-in-emulsion (GEMs) for scifi-ATAC-seq with

    (a) distinct Tn5-A barcoded nuclei, ensuring no contamination between nuclei within the droplet;

    or (b) same Tn5 barcoded nuclei, resulting in contamination between nuclei within the droplet.

(c) Distributions of the proportion of Tn5 integration sites within organelle genomes.

(d) Number of nuclei per droplet in 100k scifi-ATAC-seq.

(e) Distributions of unique Tn5 integration sites across predicted droplet type in 100k scifi-ATAC-seq.

(f) Number of nuclei per droplet in 200k scifi-ATAC-seq.

(g) Distributions of unique Tn5 integration sites across predicted droplet type in 200k scifi-ATAC-seq.

(h) Distributions of nuclei number per droplet for non-collision droplet and collision droplet in 100k and 200k scifi-ATAC-seq.

Extended Data Figure 2. Clustering and cell-type annotation of scifi-ATAC-seq and scATAC-seq, related to Figure 1.

(a) UMAP of all nuclei (n = 98,424). Nuclei are colored by clusters and labeled with predicted cell types.

   Vas. par. precursor: Vascular parenchyma precursor.

(b) UMAP of nuclei from the two scATAC-seq dataset. Nuclei are colored by the library.

(c) UMAP of nuclei from the two scifi-ATAC-seq datasets. Nuclei are colored by the library.

(d) Proportions of nuclei derived from the four datasets for each cell type. Color scheme is the same with b and c.

   There was no significant bias in cell-type proportions between the two technologies (Spearman correlation: 0.62, p-value < 0.05).

(e) UMAP embeddings overlaid with gene chromatin accessibility for representative marker genes.

(f) Relative chromatin accessibility for the marker genes across all cell types.

Extended Data Figure 3. Characterization of vascular cells with scifi-ATAC-seq, related to Figure 1.

(a) Overall distribution of nuclei number derived from each dataset across all cell types.

(b) Distributions of nuclei number derived from each dataset across vascular cells, e.g. Phloem, Procambium and Xylem.
The total nuclei number was labeled on the top of each bar.

(c) Overall distribution of aggregated Tn5 integration numbers derived from each dataset across all cell types.

(d) Distribution of aggregated Tn5 integration number derived from each dataset across vascular cells, e.g. Phloem, Procambium and Xylem.
The total Tn5 integration numbers were labeled on the top of each bar.

(e-i) Pseudobulk cell-type Tn5 integration site coverage at representative marker genes for vascular cells across four datasets.
e, Bundle sheath; f, Mesophyll, g, Phloem; h, Procambium, i. Xylem. N: The total nuclei number according to cell type.

Extended Data Figure 4. Evaluation and quality control of multiplexed scifi-ATAC-seq, related to Figure 2.

(a) Identification of high-quality barcodes using a knee-plot.

(b) Density scatterplots of log10 transformed barcode read depths (x axis) by the fraction of Tn5 integration sites mapping to within 2-kb of transcription start sites (TSSs). Dashed red lines indicate the threshold of two standard deviations from the mean used to filter lower quality barcodes.

(c) Distributions of nuclei number per droplet in 300K input multiplexed scifi-ATAC-seq.

(d-k) Distributions of proportion of SNPs in target genotypes for eight samples. Colored dots: singlet. Grey dots: identified cell barcode collision.
Median Contamination Rate: The median cross-contamination rate, attributed to index hopping, among all predicted singlets.

(l) Distributions of the proportion of Tn5 integration sites within the promoter regions, encompassing the 2-kb flanking regions around gene transcription start sites (TSSs).

(m) Distributions of the proportion of Tn5 integration sites within peaks per nucleus.

(n) Distributions of the proportion of Tn5 integration sites within organelle genomes.

(o) Distributions of unique Tn5 integration sites per nucleus.

Extended Data Figure 5. Clustering and cell-type annotation of multiplexed scifi-ATAC-seq, related to Figure 2.
(a) UMAP of all nuclei (n = 124,656). Nuclei are colored by clusters and labeled with predicted cell types.
    Vas. par. precursor: Vascular parenchyma precursor. Procabium phloem pre.: procambium phloem precursor.
(b) Proportion of nuclei derived from the eight samples for each cell type.
(c) UMAP embeddings overlaid with gene chromatin accessibility for representative marker genes.
(d) Relative chromatin accessibility for the marker genes across all cell types.

Extended Data Figure 6. Characterization of vascular cells and genotype-specific accessible chromatin regions with demultiplexed scifi-ATAC-seq, related to Figure 2.

(a-c) Pseudobulk cell-type Tn5 integration site coverage at representative marker genes for vascular cells across all samples.
  a, Phloem; b, Procambium, c. Xylem. N: The total nuclei number according to cell type.

(e) Pseudobulk cell-type Tn5 integration site coverage at representative genotype-specific accessible chromatin regions in procambium phloem precursor cells.