Life-stage specific effects of ocean temperatures on the hatchery chum salmon

Shuichi Kitada*, and Hirohisa Kishino[†,‡]

Short running title: Causal effects for declining chum salmon

* Tokyo University of Marine Science and Technology, Tokyo, Japan

[†] Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, Japan

[‡] Research and Development Initiative, Chuo University, Tokyo, Japan

Correspondence: Shuichi Kitada

Email: kitada@kaiyodai.ac.jp

ORCID

Shuichi Kitada https://orcid.org/0000-0001-5838-0374

Hirohisa Kishino https://orcid.org/0000-0002-3244-359X

CONFLICT OF INTEREST

The authors have declared that no conflict of interests exist.

Abstract

Pacific salmon (*Oncorhynchus* spp., Salmonidae) are exposed to environmental and anthropogenic stresses due to their wide geographic distribution and complex life history. We investigated the causal effects on return rates of Japanese chum salmon (*O. keta*, Salmonidae), focusing on the period of sharp decline since the early 2000s, using more than 40 years of fishery and hatchery release data and high-resolution sea surface temperature (SST) datasets along the Japanese coasts and SST along the Japanese chum salmon migration route. We examined the geographic distribution of chum salmon allele frequencies using published isoenzyme datasets and a single nucleotide polymorphism (SNP) dataset collected from the range, focusing on lactate dehydrogenase (LDH), growth hormone, gonadotropin-releasing hormone and mitochondrial DNA loci. The effect of SST increases or decreases varied depending on the life history stage of the chum salmon. Winter SST in overwintering areas had a positive effect on return rates, whereas summer SST in coastal areas had a negative effect. The spike in 2022 was driven by an increase in age 4 fish, and the shift in age structure towards younger age at maturity occurred in all areas, being consistent with recent studies throughout Alaska. Allele frequencies varied with geographical patterns, with those in Japan significantly different from the overall trend, and those involved in metabolism and growth were close to Alaskan populations. We hypothesise that relaxed selection in hatcheries has altered allele frequencies, resulting in better growth in the first and second winters, thereby promoting a younger age at maturity.

KEYWORDS

Anthropogenic selection, climate change, countergradient variation, growth and maturation, metabolism, younger age at maturity

1 | INTRODUCTION

Pacific salmon are widely distributed in the North Pacific and adjacent waters, and their life history is complex, with large variations in time spent in freshwater, coastal marine habitats, and the international waters (high seas) (Groot & Margolis, 1991; Quinn, 2018). Juvenile chum and pink salmon (*O. gorbuscha*, Salmonidae) spend a short time in freshwater before migrating to the ocean, while other Pacific salmon spend months to several years in freshwater, depending on their specific life history (Beamish, 2022). Salmon populations often consist of hundreds of subpopulations with diverse life history traits and local adaptations to environmental variation in spawning and rearing habitats (Hilborn et al., 2003). Their wide geographic range, long migrations for growth and reproduction, and complex life histories expose salmon populations to environmental and anthropogenic stressors at both local and North Pacific-wide scales (Cline et al., 2019).

The effects of climate change, ocean acidification, and large-scale hatchery releases on the productivity of Pacific salmon populations and body size decline have received increasing attention in recent years (Jeffrey et al., 2017; Cunningham et al., 2018; Cline et al., 2019; Losee et al., 2019; Connors et al., 2020; Ohlberger et al., 2022). Declines in body size, primarily due to a shift in age structure towards younger ages at maturity, have been found in chum, Chinook (*O. tshawytscha*, Salmonidae), coho (*O. kisutch*, Salmonidae), and sockeye (*O. nerka*, Salmonidae) salmon from large-scale analyses across Alaska and have been linked to climate change, increased inter- and intra-specific competition due to an overabundance of salmon in the high seas (Lewis et al., 2015; Oke et al., 2020; Ohlberger et al., 2023). This represents a trade-off between fish abundance and body size reduction, and the consequences for fisheries and ecosystems could be substantial (Oke et al., 2020; Ohlberger et al., 2023).

Pacific salmon abundance in the ocean is currently at an all-time high, and the most abundant species in the commercial catch is pink salmon, followed by the larger-bodied chum salmon (Myers et al., 2016; Beamish, 2022). In contrast, abundance in Japan and British Columbia, Canada, is at its lowest level on record, and the commercial chum salmon fishery was closed off the central coast of British Columbia in 2021 (Atlas et al., 2022). The abundance of salmon in the ocean is thought to be initially regulated by predation soon after the fish enter the ocean and is related to the size of the fish, the abundance and type of predators, and the density of juveniles; the second phase of mortality is a consequence of growth rate in the early ocean period and the ability of a fish to survive the first ocean winter (Beamish & Mahnken, 2001; Beamish, 2022). The effects of climate change on the winter ecology of salmon are importance for predicting coastal salmon fisheries and spawning biomass (Myers et al., 2016). Warm winter SST has been associated with improved growth, while summer SST has been associated with reduced growth of sockeye salmon in Bristol Bay, Alaska (Ohlberger et al., 2023).

Pacific salmon have the longest history of artificial propagation, beginning in the mid-1800s, and management of these species comprises the largest set of hatchery release programmes (Naish et al., 2007).

The hatchery release of chum salmon constitutes the largest marine stock enhancement and sea-ranching
programme in the world (Amoroso et al., 2017; Kitada, 2018). Japan is located at the southern limit of the
geographic distribution of chum salmon (Salo, 1991) and runs the world's largest hatchery release programme
of chum salmon. The Japanese chum salmon hatchery release began in 1888 (Kobayashi, 1980) and has
continued for 135 years (=27 generations, assuming a generation time of 5 years). Juvenile chum salmon
releases from Japan have increased remarkably since the 1970s to ~2 billion per year. The number of returning
chum salmon to Japan has increased dramatically since the 1970s. However, decreases in mean body weight
and 'older' age at maturity occurred in 1970s-1980s, possibly due to the density-dependent effect in the high
seas (Kaeriyama, 1989; Ishida et al., 1991; Hayashizaki & Ida, 1993; Morita & Fukuwaka, 2007). Despite
continued hatchery releases, the number of returning chum salmon declined sharply after the early 2000s.
Previous studies have reported that higher spring SSTs in the coastal zone contributed to better survival of
juvenile chum salmon through improved feeding, growth and swimming ability (Saito & Nagasawa, 2009;
Kuroda et al., 2020; Honda et al., 2021). Conversely, shorter inshore stays associated with warming SSTs
could lead to lower juvenile survival along the Honshu Pacific coast (Kaeriyama, 2022). It is generally
accepted that a warming climate may cause a decline in return rates of chum salmon. However, the underlying
mechanisms leading to the decline of Japanese chum salmon remain largely unknown because few studies
have used data at both the coastal and North Pacific-wide scales, from ocean entry to migration route, to infer
causal effects on survival or return rates.
99
The number of chum salmon returning to Japan (Table S1) reached an all-time low of 19 million fish in 2021,
but unexpectedly increased in 2022 (169% of the previous year in all of Japan), especially in the Hokkaido
Sea of Japan (182%) and the Sea of Okhotsk (179%), while remaining very low in the Honshu Pacific (112%),
the former main production area. In the Chitose River in the Sea of Japan, where Japan's oldest hatchery
operates, 587,000 fish were returned in 2022, surpassing the historical record of 551,000 in 1995 (Chitose
Aquarium, 2023). The sudden increase in 2022 came as a big surprise to fishers, hatchery managers, local
people and biologists. This unusual phenomenon may provide us with an opportunity to understand the
underlying mechanisms of chum salmon population dynamics. Several regions in the USA and Russia also
showed rebounds in chum salmon catches in 2022 (Figures S1, S2). The mean body weight (BW) of Japanese
chum salmon in Hokkaido and Honshu has decreased since 2000 (Yamaguchi et al., 2021; Watanabe et al.,
2022). The mean BW of chum salmon returning to Japan in 2022 was 2.83 kg in Hokkaido and 2.76 kg in
Honshu, which was the historical low after 1989 and 1994, respectively (Fishery Research Agency, 2023).
Selection in hatcheries may cause a reduction in body size, as allele frequencies of several SNPs were
significantly differentiated beyond the neutral population structure in Japanese chum salmon populations
(Kitada & Kishino, 2021).
115
116 In this study, we investigate the effects of changing climate, stocking practices, predators, salmon abundance
117 (chum and pink), and selection in hatcheries on Japanese chum salmon population dynamics, focusing on the

period of sharp decline since the early 2000s. We estimated size at release and time of release for 31 million juveniles based on 58 thousand release events in 355 rivers in Hokkaido and Honshu (1998-2020). We calculated mean SSTs for the time of release based on high-resolution monthly mean SSTs in 30 coastal areas of Japan (1982-2020), and age composition based on 1,853 surveys of 1.2 million adult fish returned to Japan (1999-2019). We identified potential predators of juvenile chum salmon using catch data from Japanese fisheries covering 70 species (1980-2020). For the high seas environment, we defined the seasonal migration routes of Japanese chum salmon based on the results of the previous high seas studies (Myers et al., 2007; Urawa et al., 2018) and calculated the monthly mean SST at summer feeding and wintering grounds in the North Pacific (1982-2022). We estimated return rates for releases from 2003 to 2019 based on age composition and the number of juveniles released and adults returning. Using these data, we inferred causal factors for the decline of Japanese chum salmon through multiple regression analysis of return rates on climate, stocking practices, predation and competition at both local and North Pacific scales. Finally, we inferred the geographic distribution of chum salmon gene allele frequencies focusing on metabolism, growth, maturation, and oxygen consumption, using the published LDH allozyme data set (18,892 individuals from 147 populations), with re-examination of SNP data set (53 loci from 10,458 individuals from 114 populations) of adult chum salmon collected from the Pacific Rim.

2 | METHODS

2.1 | Data sets

We compiled data on the release and catch of Japanese chum salmon. The catch (coastal catch plus hatchery broodstock catch) and release numbers of chum salmon from 1970 to 2022 in Hokkaido were obtained from the Hokkaido Salmon Enhancement Association (HSEA, temporal translation). Those in the prefectures in Honshu were obtained from FRA. For Aomori Prefecture (in the northernmost part of Honshu), which has both Pacific and Sea of Japan coastlines, catch and release data by reporting region were obtained from the Aomori Prefectural Government (APG). They were divided into the Pacific (Pacific Coast, Tsugaru Strait and Mutsu Bay) and Sea of Japan coasts because the Tushima Current in Tsugaru Strait flows from the Sea of Japan to the Pacific Ocean (Kuroda et al., 2020), through which released juveniles recruit in the Pacific Ocean. We summarised the catch and release data of chum salmon for the seven management areas (Table S1, Figure S3): Sea of Okhotsk, Nemuro Strait, Hokkaido Eastern Pacific, Hokkaido Western Pacific, Hokkaido Sea of Japan, Honshu Pacific, and Honshu Sea of Japan (Saito & Nagasawa, 2009; Miyakoshi et al., 2013). The management areas coincided with the genetic structure of the population (Beacham et al., 2008).

Datasets on size at release and month of release from 1998 to 2020 and age composition from 1997 to 2019 in rivers in Hokkaido and Honshu were downloaded from the FRA Salmon Database (http://salmon.fra.affrc.go.jp/zousyoku/fri_salmon_dept/sdb.html) (Data S1, S2). Annual commercial catch data for all marine species in Japan from 1980 to 2020 were also downloaded from the Ministry of Agriculture,

Forestry and Fisheries (MAFF, https://www.jfa.maff.go.jp/j/kikaku/toukei/). Catch (1925-2022) and release (1952-2022) data for chum and pink salmon in the North Pacific were obtained from North Pacific Anadromous Fish Commission statistics (NPAFC, https://npafc.org/statistics/).

We downloaded the global 0.25°×0.25° monthly mean SST data file from 1982 to 2022 ("sst.mnmean.nc") from the NOAA OISST high-resolution dataset (https://downloads.psl.noaa.gov/Datasets/noaa.oisst.v2/). For the Japanese coasts, we downloaded the 0.25°×0.25° daily mean SST datasets from 1982 to 2021 for 18 areas in Hokkaido (Data S3) and 12 areas in Honshu (Data S4) from the Japan Meteorological Agency (JMA). The names of the SST observation areas and the map with the JMA sources are shown in Table S2.

2.2 | Size and time at release, and age composition

The Japanese chum salmon hatchery programme recommended a release size greater than 1 g (BW) and a release time when the coastal SST ranged from 5 to 13 °C (Honda et al., 2020). Increasing body weight (BW) at release has been associated with increased return rates of chum salmon (Saito & Nagasawa, 2009), and survival of Chinook and coho salmon (James et al., 2023). We selected release events with no missing measurements of BW and fork length (FL) for Hokkaido and Honshu from the FRA salmon database (Data S1). The number of release events was 37,886 in 184 rivers in Hokkaido and it was 19,605 in 171 rivers in Honshu between 1998 and 2020. Since the number of individuals released (n_l) varied among release events ($l = 1, ..., L$), we calculated the weighted means size at release for BW (1998-2020) as $\overline{BW} = (\sum_{l=1}^{L} BW_l\, n_l)/\sum_{l=1}^{L} n_l$, where $L = 37,886$, $\sum_{l=1}^{L} n_l = 21,332,093$ for Hokkaido and $L = 19,605$, $\sum_{l=1}^{L} n_l = 9,588,413$ for Honshu. Weighted means were also calculated for FL and Fulton's condition factor ($K = BW(g)/FL(cm)^3 \times 100$) in a similar manner.

Juvenile chum salmon spend only a short time in freshwater before migrating to the ocean (Groot & Margolis, 1991; Quinn, 2018; Beamish, 2020). In Japan, most rivers where chum salmon are released are short (less than 50 km in Hokkaido) and released fish spend only ~10 days or less before entering the ocean (Kobayashi, 1980). Therefore, the time of release can be related to the critical period in the coastal environment (Beamish & Mahnken, 2001; Beamish, 2022). To estimate the distribution of release times in Japan, we counted the number of release events by month.

Age composition is necessary to estimate the number of fish returning from each stocking. We compiled age composition data of chum salmon returns (1997-2019) with sample size in each survey (Data S2), from which we calculated the weighted mean age composition as $\hat{p}_a = (\sum_{h=1}^{H} p_{la}\, n_h)/\sum_{h=1}^{H} n_h$, where h refers to each survey ($H = 1,853$), n_h is the sample size (individuals) ($\sum_{h=1}^{H} n_h = 1,170,817$) and a refers to age at return (a=2,…,8<). These calculations were made for each year by management area. We summarised the compiled data for size at release (Table S3, Figures S4-S6), month of release (Figure S7) and age composition (Table S4, Figure S8).

2.3 | Long-term SST changes across the range of Pacific salmon

Pacific salmon are distributed throughout the high seas of the North Pacific, Bering Sea, Gulf of Alaska and Sea of Okhotsk (Walker et al. 2000; Myers et al. 2007; Urawa et al., 2018). Early juvenile chum salmon have recently been found in the North American Arctic (Dunmall et al., 2022). Environmental conditions such as SST at regional spatial scales largely determine salmon survival and are better predictors of survival than large-scale climate indices such as PDO (Mueter et al., 2002). To understand the long-term changes in SST in the salmon distribution area, we created the monthly mean 1°×1° SST isoclines in the North Pacific Ocean for 1900, 2000, 2010 and 2022 using the Plot COBE Sea Surface Temperature system at the NOAA Physical Sciences Laboratory (https://psl.noaa.gov/data/gridded/data.cobe.html), based on the COBE SST dataset from the Japan Meteorological Agency (JMA). The 1°×1° monthly mean SST isoclines around Japan for 1980, 2000 and 2022 are also generated in a similar way.

2.4 | SST at release

To approximate the SST values near the estuaries that juvenile salmonids are most likely to experience after entering the ocean (Oke et al., 2020), we calculated the monthly mean SSTs from 1982 to 2020 for the 18 areas in the Hokkaido coast and the 12 areas in the Honshu coast based on the 0.25°×0.25° daily SST data from the JMA (Table S2). Based on the monthly mean SSTs of the areas, we calculated the monthly mean SSTs from 1982 to 2020 in the seven chum salmon management areas (Figure S3). The monthly mean SSTs for the Sea of Okhotsk were obtained by averaging the monthly mean SSTs in the Abashiri, Monbetsu and Soya regions. For the Nemuro region, the monthly mean SSTs of the Nemuro Strait were used. Monthly mean SSTs for the Hokkaido Eastern Pacific were calculated by averaging those over Nemuro Pacific, Kushiro, and Tokachi, and for the Hokkaido Western Pacific over Hidaka, Iiburi, Uchiura Bay, western Tsugaru Strait, and Tsugaru Strait. For the Hokkaido Sea of Japan, we averaged the monthly mean SSTs of northern and southern Rumoi, Ishikari, western Shiribeshi and Hiyama. For the Honshu Sea of Japan, monthly mean SSTs were averaged from Aomori, northern and southern Iwate, Miyagi, Fukushima and northern Ibaraki. For the Honshu Sea of Japan, monthly mean SSTs were averaged from Aomori, Akita, Yamagata, northern and southern Niigata, and Toyama Bay. Finally, the mean SSTs for the months of release (Fig. S7) were calculated by management area: Sea of Okhotsk (Apr-Jun), Nemuro (Mar-Jun), Hokkaido Eastern Pacific (Mar-Jun), Hokkaido Western Pacific (Mar-May), Hokkaido Sea of Japan (Mar-May), Honshu Pacific (Feb-May), and Honshu Sea of Japan (Feb-Apr) (Table S5).

2.5 | Japanese chum salmon migration route and SST

The seasonal migration of Japanese chum salmon was modelled based on a series of high seas studies, including genetic stock identification (Myers et al., 2007; Urawa et al., 2018). According to these studies, juvenile chum salmon (age 0) are distributed in the Sea of Okhotsk until autumn (Aug-Nov) and migrate to a narrow area in the western North Pacific (hereafter North Pacific) to overwinter. After the first

overwintering period, age 1 chum salmon migrated to the Bering Sea to feed until the following summer.

Immature salmon began migrating to the Gulf of Alaska in late autumn for the second overwintering period

(age 2). The following year, they return to the Bering Sea and again migrate to the Gulf of Alaska to

overwinter (age 3). The migration between the Bering Sea and the Gulf of Alaska is repeated until they return

to Japan through the Bering Sea.

Juvenile chum salmon born in rivers along the Pacific coast of Japan migrate to the Sea of Okhotsk (Honda

et al., 2017). Juvenile chum salmon from all regions of Japan are found in the Sea of Okhotsk, and there is a

significant positive correlation between the abundance of juvenile Japanese chum salmon and adult returns

to Japan (Urawa & Bugaev, 2021). Juvenile chum salmon were abundant in coastal areas of the Sea of

Okhotsk from May to June when SST were between 8 and 13°C, while they disappeared from coastal waters

after late June when SST exceeded 13°C (Nagata et al., 2007). Based on these results, we assumed that

juvenile chum salmon released in all areas would migrate to the Sea of Okhotsk and disperse offshore in July.

Based on information obtained from field surveys (Urawa et al., 2018; Urawa et al., 2022), we defined the

locations of four main areas of Japanese chum salmon migration: 1. Sea of Okhotsk Japan Coast (45.2°N-

46.5°N, 142.5°E-148.5°E), 2. North Pacific Ocean (NPO, 42°N-45°N, 158°E-165°E), 3. Bering Sea (53°N-

58°N, 175°E-170°W), 4. Gulf of Alaska (48°N-56°N, 140°W-150°W). Using the NOAA OISST 0.25°×0.25°

data "sst.mnmean.nc", we calculated the monthly mean SSTs from 1982 to 2022 for the migration areas using

the   `fldmean`   function   in   the   `cdo-2.1.1`   package   (Climate   Data   Operators,

https://code.mpimet.mpg.de/projects/cdo) (Data S5). We described the areas and migration routes on the map

using the `sf` package in R. Trends in the monthly mean summer and winter SSTs (Table S6) were tested

using the `MannKendall` function in R.

2.6 | Catch of marine species in Japan

Predation immediately after fish enter the ocean is considered a major cause of salmon mortality, depending

on fish size, juvenile density, and predator abundance and species (Beamish, 2022). To identify potential

predators of juvenile chum salmon, we summarised commercial catches of all marine species landed in Japan

with no missing data from 1980 to 2020. Marine mammals were excluded because they are subject to strict

catch limits. This resulted in catch data for 56 out of 70 species, including 39 fish, six crustaceans, five

shellfish, four animals and two algae (Table S7). To characterise temporal changes in catch, we performed

principal components analysis (PCA) on the catch data for the 56 marine species using the `prcomp` function

in R. Changes in chum salmon catch and species with increasing catch in the chum salmon range were plotted

on the map.

2.7 | Return rates

Return rate has been widely used as a general measure of the effectiveness of enhancement of marine stocks,

including salmonids (Kitada, 2018, 2020). It has been used as a measure of relative survival for chum salmon
(Kaeriyama, 1999; Saito & Nagasawa, 2009), Chinook and coho salmon (James et al., 2023). It is generally
accepted that most chum salmon in Japan are hatchery-reared fish (Hiroi, 1998; Kaeriyama, 1999; Kaeriyama
& Edpalina, 2004; Kobayashi, 1980; reviewed by Iida et al., 2018). We assumed that all returning chum
salmon were hatchery-released. Return rates were estimated for each management area. The total number of
fish returning from releases in year t (Y_t, coastal commercial catch plus river catch for hatchery broodstock)
from releases in year t is estimated as

273
$$\hat{Y}_t = \sum_{a=A_{min}}^{A_{max}} y_{t+a-1}\, \hat{p}_{a,t+a-1} = \sum_{a=A_{min}}^{A_{max}} \hat{Y}_{a,t}$$

where y_{t+a-1} is the observed number of fish returned in year $t + a - 1$ from juveniles released in year t,
and $\hat{p}_a$ is the estimated proportion of age a fish among the fish returned in year $t + a - 1$, The return rate of
juveniles released in year t was estimated by $\hat{R}_t = \hat{Y}_t/X_t$, where X_t is the number of hatchery-reared
juveniles released in year t (known). The age-specific return rate was estimated as $\hat{R}_{a,t} = \hat{Y}_{a,t}/X_t$. We
summed the number of fish returned from ages 2 to 5 ($A_{min} = 2$, $A_{max} = 5$) because the main ages of return
were ages 4 and 5, and there were very few returns older than Age 6 (Table S4).

We estimated the age composition from Data S2 (1997-2019), but it was not available for 2020-2023, which
was estimated as a moving average of the previous three years. The number of fish returning in 2023 was
currently unknown (September 2023) and we estimated it as an average of those in 2021 and 2022 to predict
only the number of age 5 fish returning in 2023 from the 2019 release. Based on the age composition, we
calculated the number of returning fish (ages 2-5) for each release year, from which we estimated return rates
and age-specific return rates from 2003 to 2019. We summarised the return rates by management area used
for the regression analysis (Table S8).

The distances between return rates in the management areas were measured by $1 - r_{jk}$ and visualised by a
neighbour-joining (NJ) tree in the R package `ape`, where r_{jk} is the correlation coefficient between pairs of
return rates. We visualised the relationship between geographical distance and return rates by management
area. Great circle geographical distances between hatcheries and the central coast of the Sea of Okhotsk (off
Monbetsu, 44.5°N, 143.5°E) were calculated using the distm function in the R package `geosphere`,
assuming 6 waypoints representing reasonable migration paths from the hatcheries to off Monbetsu: Fukaura
(40.6°N, 139.9°E) and Wakkanai (45.5°N, 141.6°E) on the Sea of Japan coast, and Shimokita (41.4°N,
141.5°E), Erimo (41.9°N, 143.2°E), Nemuro (43.4°N, 145.8°E), and Shiretoko (44.3°N, 145.3°E) on the
Pacific coast. We used 254 of the 262 hatchery locations identified in our previous study (Kitada, 2020),
excluding 8 hatcheries whose locations could not be identified.

2.8 | Regression analysis of return rates

Our model describes the relationship between return rates across all ages ($\hat{R}_t$) and explanatory variables such

as SSTs, size at release and catch of potential predators and competitors (See next section for details). We

transformed the return rate to the log scale for model fitting in a multiple linear regression:

$$\log(\hat{R}_t) = \log \beta_0 + \sum_{i=1}^{s} \beta_i \, z_i + \varepsilon_t$$

where β_0 is the intercept, the explanatory variables are denoted by $\boldsymbol{z}$. The egression coefficients were

denoted by $\boldsymbol{\beta}$ and $\varepsilon_t \sim N(0, \sigma^2)$ is the error which was assumed to be normally distributed.

We performed a regression analysis by areas that had a similar trend in return rates based on the NJ tree,

using a model selection using AICc (Sugiura, 1978), which is a small sample correction of the Akaike

Information Criterion (AIC, Akaike, 1973). We used the results of the best model rather than model averaging,

which may not be reliable in the presence of multicollinearity between explanatory variables (Cade, 2015).

The use of an unstandardised measure is usually preferred to a standardised measure if the units of

measurement are useful as effect sizes at a practical level (Wilkinson, 1999). We therefore used the regression

coefficient for unstandardised explanatory variables as the effect size: $\exp(\hat{\beta}_i)$ for coefficient i, visualised

using the `forestplot` function in R.

We used 14 explanatory variables (s=14), and the hypothesised mechanisms were summarised in Table 1. We

selected the most parsimonious model (hereafter, best model) from 2^{14} (=16,384) models with the minimum

AICc value using the `dredge` function in the `MuMIn` package in R. In our model, size at release and SST at

release were the area-specific, while July SST in the Sea of Okhotsk was common to all management areas

because juvenile chum salmon from all regional populations in Japan are distributed in the Sea of Okhotsk

in summer (Urawa et al., 2018). Yellowtail (*Seriola quinqueradiata*, Carangidae) has been identified as a

potential predator of juvenile chum salmon (see Section 3.5 for details). Winter SSTs in the North Pacific

and Gulf of Alaska were used to infer effects during the first and second overwintering at ages 1 and 2, while

summer SSTs in the Bering Sea were used to infer effects during feeding of age 1 fish (Myers et al., 2007;

Urawa et al., 2018). The stomach contents of chum and pink salmon were similar in the Bering Sea, Gulf of

Alaska and Western Subarctic Gyre (Davis et al., 1998; Kaeriyama et al., 2004; Qin & Kaeriyama, 2016;

Daly et al., 2019), and the catches of Russian, US and Canadian chum and pink salmon (Figures S9, S10)

were used as explanatory variables to infer intra and interspecific competition for food. The age composition

of Japanese chum salmon showed that Age 4 fish were dominant (Table S4, Figure S8), so the catches of

Russian, US and Canadian chum salmon in year t+3 were used as explanatory variables to measure

intraspecific competition with Japanese chum salmon released in year t, as they return in year t+3 as Age 4

fish. They were assumed to spend 3 years in the shared environment of the North Pacific and adjacent waters.

Pink salmon have a fixed life span of 2 years (Groot & Margolis, 1991; Quinn, 2018), and Russian, US and

Canadian catches in year t+1 were used as explanatory variables to measure interspecific competition with

Japanese chum salmon released in year t. They were assumed to spend 1 year in the shared environment. We

have summarised the explanatory variables specific to each management area (Table S9) and common to all management areas (Table S10).

2.9 | Gene frequencies for metabolism, growth, and maturation

First, we focused on LDH-A_4 and LDH-B_4 as isoenzyme markers (Markert & Møller, 1959) that are potentially useful for understanding physiological thermal adaptation and evolution in fish (Merritt, 1972; Place & Powers, 1979; Powers et al., 1991; Fields & Somero, 1998; Somero, 2004; Somero, 2010; Somero, 2022). The LDH-A_4 isozyme is commonly found in vertebrate skeletal muscle and the LDH-B_4 isozyme is predominantly found in the heart (Powers et al., 1991; Somero, 2004). The metabolism of white muscle (skeletal muscle) is predominantly anaerobic and expresses the LDH-A_4 locus, whereas red muscle (heart) has significant aerobic metabolism and expresses almost exclusively the LDH-B_4 locus (Powers et al., 1991). The LDH-A_4 locus mainly involved in the conversion of pyruvate to lactate (i.e., anaerobic glycolysis), whereas the LDH-B_4 locus mainly converts lactate to pyruvate (i.e., gluconeogenesis and aerobic metabolism) (Somero, 2004, 2010; Powers et al., 1991). The lactate is resynthesized to glycogen in the liver and used as an energy source, while pyruvate produce adenosine triphosphate (ATP) for new energy, which are critical fuel metabolites during exercise (Liang et al., 2016).

Second, we re-examined the geographic distribution of chum salmon SNP allele frequencies (Seeb et al., 2011), focusing on growth hormone (GHII3129), gonadotropin-releasing hormone (GnRH373), and the combined mitochondrial DNA3 (mtDNA3) locus of chum salmon SNPs. Growth hormone (GH) is an important regulator of somatic growth in salmonids (Björnsson, 1997). Gonadotropin-releasing hormone (GnRH) regulates reproduction in vertebrates including salmonids (Khakoo et al., 1994). mtDNA plays a key role in aerobic ATP production and contributes to the ability of endurance exercise (reviewed by Stefàno et al., 2019). The mtDNA control region has functions in Dntp metabolism (Nicholls & Minczuk, 2014) and oxygen consumption (Kong et al., 2020).

We organised the allele frequencies of the LDH-A1 (LDH-A_4) and LDH-B2 (LDH-B_4) locus in chum salmon in the Pacific Rim (Supplementary Methods). To avoid the problems associated with standardization of electrophoretic bands obtained in different laboratories, we only used data generated using the same protocol (Aebersold et al., 1987) and followed the American Fisheries Society genetic nomenclature (Shaklee et al., 1990). These were collected in Japan (Sato & Urawa, 2015), Japan and the northern Russia (Winans et al., 1994), Sakhalin (Salmenkova et al., 2008) and the Russian Far East (Ivankova & Efremov, 2009). For populations in Alaska, the Alaska Peninsula, Southeast Alaska, British Columbia and Washington, allele frequencies were summarised from Seeb et al. (1995). As done in Seeb et al. (1999), we confirmed that the equal allele frequencies of LDH-A1 *100 (LDH-A_4^a) were obtained in the same rivers in Japan and Russia. All Japanese samples were caught at weirs in hatchery-reared rivers and hatcheries and were therefore hatchery-reared fish and/or wild-born hatchery progeny. In contrast, the northern Russian samples were

collected from natural streams; these fish were assumed to be wild, as most Russian chum salmon are
naturally maintained (Winans et al., 1994). The North American samples included 3 hatchery samples out of
42 (Seeb et al., 1995). The Russian samples from Sakhalin and the Russian Far East included 12 and 6
samples of hatchery fish out of 19 and 14 samples, respectively. We used the most common alleles, LDH-
A1*100, and LDH-B2*100 in our analysis of allele frequencies. We obtained allele frequencies in 147 chum
salmon populations collected throughout the range in 1986-2005 (n = 18,892) (Data S6).
380
For the nuclear SNPs (Seeb et al., 2011), we converted the original genotype data to Genepop format (Rousset,
2008) and loaded the data using the read.GENEPOP function in the R package `FinePop2`. We used minor
alleles that had a mean allele frequency across populations of less than 0.5 (Kitada & Kishino, 2021). We
summarised the SNP allele frequencies of GHII3129, GnRH373, and mtDNA3 in 114 chum salmon
populations collected throughout the range in 1989-2006 (n = 10,458) (Data S7). The mtDNA3 locus had
five alleles, and the major allele (second one) was used for the analysis. The sample from South Korea (n =
96) was excluded from the analysis, due to our focus on Japanese chum.
388
389 We recorded the approximate longitude and latitude of the sampling locations based on the names of the
390 rivers and/or areas and the maps of the original studies using Google Map (Data S6 and S7). We plotted the
391 locations on the map using the `sf` package in R. Sampling points were visualized by a color gradient of allele
392 frequencies.
393

3 | RESULTS

3.1 | Long-term SST changes across the range of Pacific salmon

The mean SST isoclines show that SSTs are generally lower on the Asian side than on the American side
(Figure 1, Figures S11-S13). Until 2010 (during 110 years) the increase in SST was relatively slow in all
seasons, while after 2010 a rapid increase in summer was observed. Summer SST (June-August) increased
in the Bering Sea, the Sea of Okhotsk, the western, central and eastern North Pacific, and the Gulf of Alaska,
while spring (April-May), autumn (October-November) and winter (January) SST were stable throughout the
Pacific salmon range until 2010, with relatively slow increases over the past decade. Around Japan, SST
increased gradually but continuously in May, June and July (Figure S14). The Pacific side of Japan has denser
SST contours than the Sea of Japan side. Higher SSTs were observed along the Sea of Japan coast than in the
Honshu Pacific for the same latitude.
405

3.2 | Japanese chum salmon migration route and SST

The map shows the migration route of Japanese chum salmon until their return, and the box plot shows the
changes in SST along the seasonal migration of Japanese chum salmon (Figure 2). In the Sea of Okhotsk, the
SST (mean ± standard deviation) in July was very high (12.3 ± 1.2°C). In the North Pacific, SST remained
relatively high even in December (8.0 ± 0.9°C), but decreased significantly by 2.0°C in January (6.0 ± 0.7°C,

$t = 10.9, p = 0.0000$, Welch t-test). SSTs were lower but stable in February (5.0 ± 0.6℃), March (4.7 ±

0.5℃), and April (5.0 ± 0.5℃), and increased in May (6.2 ± 0.6℃) to the same level as in January. In the

Bering Sea, the SST remained low in June (5.7 ± 0.6℃). It increased in July (8.0 ± 0.7℃) and was the highest

in August (9.7 ± 0.8℃) and remained high in September (9.3 ± 0.8℃). It decreased significantly in October

(7.4 ± 0.6℃) and November (5.4 ± 0.5℃). In the Gulf of Alaska, SSTs were 1.5℃ higher in December (6.9

± 0.8℃) than in November in the Bering Sea, and then decreased significantly in January (6.0 ± 0.7℃, $t =$

5.2, $p = 0.0000$). SSTs were lower but stable in February (5.6 ± 0.7℃), March (5.4 ± 0.7℃), and April (5.7

± 0.6℃), and increased in May (6.9 ± 0.8℃) to a similar level in December.

Interestingly, the mean January SST was just the same in both wintering areas. However, the February-April

mean SST was significantly lower than the January SST in the North Pacific ($t = 8.3$, $p = 0.0000$) and in

the Gulf of Alaska ($t = 3.3$, $p = 0.0012$), and was significantly lower in the North Pacific (4.9 ± 0.5℃,

$t = -5.3$, $p = 0.0000$) than in the Gulf of Alaska (5.5 ± 0.6℃). January SST and February-April mean SST

were positively correlated in the North Pacific ($r = 0.78, t = 7.9, p = 0.0000$) and Gulf of Alaska ($r =$

$0.91, t = 13.6, p = 0.0000$), but the linear relationship was weaker in the North Pacific (Figure S15). From

the characteristics of winter SSTs between and within wintering areas, we chose both the January SST and

the February-April mean SST as explanatory variables, leaving the choice of which to use to model selection.

For the summer SST, the Sea of Okhotsk July SST and the Bering Sea August-September mean SST were

used as explanatory variables (Table 1).

July SST in the Sea of Okhotsk showed an increasing trend (Mann-Kendall trend test, $\tau = 0.38, p =$

0.0005), and the recent increase was remarkable (Figure 3). August-September SST in the Bering Sea

increased $\tau = 0.43, p = 0.0001$), but decreased with variability after a strong increase in 2013. Winter SST

trends in the North Pacific were insignificant for both January ($\tau = 0.21, p = 0.0506$) and February-April

($\tau = 0.21, p = 0.0562$). In the Gulf of Alaska, winter SST trends were also insignificant both for January

($\tau = -0.06, p = 0.5821$) and February-April ($\tau = -0.03, p = 0.7875$).

3.3 | Chum salmon enhancement and SST at release

From 1970 to 2021, a total of 86,357 million juvenile fish were released and 2,266 million fish (2.6%) were

returned (Figure 4). As the number of fish released increased, the total number of fish returned increased

significantly until 1996, when the historical maximum of 88 million fish was reached. Despite the relatively

stable trend in the number of fish released, the number of fish returned has continued to decline since 2004

and will reach the four-decade minimum of 19 million in 2021. In particular, the number of fish returned

increased in 2022, especially in the Sea of Okhotsk and the Hokkaido Sea of Japan. The trend across Japan

was synchronised in the two main production areas, Nemuro Strait and Honshu Pacific, as shown by the

dashed lines. Hokkaido Eastern Pacific and Hokkaido Western Pacific showed a similar trend after the mid-

2000s. In contrast, the other major production area, the Sea of Okhotsk, showed a unique trend where the

number of fish returned increased with large fluctuations, reaching the historical maximum in 2013, with a
large decrease until 2019, but recovering significantly in 2022. The Hokkaido Sea of Japan also had a unique
trend, but showed a similar significant increase in 2022. It is worth noting that the recovery in 2022 was
observed in all areas, although the recovery in 2022 was highly variable.

SST at release (Figure 4) increased in the Sea of Okhotsk ($5.5 \pm 0.4°C$, $\tau = 0.37, p = 0.001$), Nemuro ($3.9 \pm$
$0.6°C$, $\tau = 0.36, p = 0.001$), the Hokkaido Sea of Japan ($7.5 \pm 0.4°C$, $\tau = 0.35, p = 0.002$), and the
Honshu Sea of Japan ($9.9 \pm 0.5°C$, $\tau = 0.28, p = 0.016$), while the trends were insignificant in the Hokkaido
Eastern Pacific ($4.4 \pm 0.4°C$, $\tau = 0.06, p = 0.611$) and the Hokkaido Western Pacific ($6.2 \pm 0.6°C$, $\tau =$
$0.06, p = 0.578$), and the Honshu Pacific ($10.1 \pm 0.8°C$, $\tau = -0.09, p = 0.421$).

3.4 | Return rate and weight at release
Return rates had common local peaks for the 2004, 2006, 2012 and 2015 releases (Figure 5). The return rate
for ages 2-5 was highest in the Sea of Okhotsk ($6.5 \pm 2.2\%$). The recent increase was synchronised in the
Hokkaido Sea of Japan ($1.2 \pm 0.6\%$). Age-specific return rates revealed that the increase in return rates for
the 2019 release in the Sea of Okhotsk and the Hokkaido Sea of Japan was driven by an increase in those at
age 4, while the return rate at age 5 decreased slightly. Conversely, return rates remained low in other areas,
with a slight increase in the 2019 release: Nemuro ($3.3 \pm 2.0\%$), Hokkaido Eastern ($1.7 \pm 1.4\%$) and Hokkaido
Western Pacific ($2.0 \pm 1.0\%$), Honshu Pacific ($1.1 \pm 0.7\%$) and Hoshu Sea of Japan ($0.6 \pm 0.2\%$). The younger
ages at maturity was also observed in these areas, although the increases were relatively small.

Higher return rates were obtained in the management areas that were geographically closer to the central Sea
of Okhotsk (Figure 6a,b). The weighted mean size at release (Figure 6c) was above the recommended
threshold (BW > 1g) in all management areas. The mean weight at release was maintained over the two
decades in the Hokkaido Western Pacific ($1.3 \pm 0.03g$, $\tau = 0.01, p = 0.967$), while decreasing trends were
found in other areas: the Sea of Okhotsk ($1.2 \pm 0.06g$, $\tau = -0.43, p = 0.019$), Nemuro Strait ($1.2 \pm 0.09g$,
$\tau = -0.50, p = 0.006$), Hokkaido Sea of Japan ($1.1 \pm 0.05g$, $\tau = -0.43, p = 0.019$), Honshu Pacific (1.3
$\pm 0.16g$, $\tau = -0.65, p = 0.000$) and Honshu Sea of Japan ($1.0 \pm 0.20g$, $\tau = -0.50, p = 0.006$). In the
Hokkaido Eastern Pacific, it was relatively high, but the trend was insignificant with large variations ($1.5 \pm$
$0.13g$, $\tau = -0.15, p = 0.434$).

The NJ tree (Figure 6d), which measures pairwise differences in return rates between management areas,
showed that the Hokkaido Sea of Japan, Sea of Okhotsk and Honshu Sea of Japan were distinct, while the
Nemuro Strait, Hokkaido Eastern Pacific and Hokkaido Western Pacific and Honshu Pacific had a similar
trend in return rates and were very close to each other. In subsequent analyses, we used the 4 geographical
areas: Sea of Okhotsk, Hokkaido Sea of Japan, Honshu Sea of Japan, and Hokkaido and Honshu Pacific
(combining the Nemuro Strait, Hokkaido Eastern Pacific, and Hokkaido Western Pacific and Honshu Pacific).

For the Hokkaido and Honshu Pacific area, simple averages were calculated for SST and weighted size at

release over the four areas. The number of returned and released chum salmon was also summed over the

areas and the return rate was calculated.

3.5 | A possible marine predator of juvenile chum salmon

The biplot of the PCA (Figure 7a) identified differences in the magnitude of catches as a primary component

(PC1, 51% variance), while the second component (PC2, 17% variance) corresponded to the trends in catches

in 1980-1998 and 1999-2020. The PCA confirmed that the changes in commercial catches of 56 marine

species in Japan after 2000 were mainly 15ertilized15ed by the 4 species that increased in recent decades:

yellowtail, Japanese Spanish mackerel (*Scomberomorus niphonius*, Scombridae), Japanese scallop

(*Mizuhopecten yessoensis*, Pectinidae) and round herring (*Etrumeus micropus*, Dussumieriidae) (Figure S16).

Among these, yellowtail and Japanese Spanish mackerel are strong piscivores. The map of changes in catches

of chum salmon and the two piscivores by prefecture (Figure 7b) shows the northward shift of yellowtail

landings into the chum salmon distribution areas, Honshu North Pacific, Hokkaido Pacific, Hokkaido Sea of

Japan and the Sea of Okhotsk. In fact, there were significant increases in these areas (Figure S17). On the

other hand, the increase in catches of Japanese Spanish mackerel remained in the Eastern Japan area, with

slight increases in the Honshu North Pacific. Based on these results, we defined yellowtail as a possible

predator of juvenile chum salmon, and catches of yellowtail were also summed over the areas.

3.6 | Causal effects on chum salmon return rates

Table 2 summarises the best models from the regression analysis. 75% of the variation in return rates from

the Sea of Okhotsk was explained by the positive effects of winter SST (Jan) in the Gulf of Alaska and

Russian chum salmon, and the negative effect of yellowtail. In the Hokkaido and Honshu Pacific, the positive

effect of weight at release explained 43% of the variation in return rates. In the Hokkaido Sea of Japan, 49%

of the variation in return rates was explained by the largest positive effect of winter SST (Feb-Apr) in the

North Pacific and the negative effect of summer SST (July) in the Sea of Okhotsk. Weight at release also had

a negative effect on the return rate. The 69% of the variation in return rates from the Honshu Sea of Japan

was explained by the positive effects of weight at release and winter SST (Jan) in the Gulf of Alaska, and the

negative effect of SST at release.

The correlation between predicted and observed return rates were high in the Sea of Okhotsk ($r = 0.89, t =$

$7.73, p = 0.0000$) and Honshu Sea of Japan ($r = 0.86, t = 6.63, p = 0.0000$), while moderate in Hokkaido

and Honshu Pacific with a wider 95% prediction interval ($r = 0.68, t = 3.61, p = 0.0026$) and Hokkaido

Sea of Japan ($r = 0.77, t = 4.6, p = 0.0003$) (Figure S18). Our regression models predicted the trends in

the observed return rates in the 4 areas well (Figure 8), and all observed values were within the 95%

prediction intervals.

The effect size for each regression coefficient (Figure 9), measured by $\exp(\hat{\beta}_i)$, represents the rate of change

per unit. Winter SST in the North Pacific and the Gulf of Alaska had a positive effect in 3 areas. A 1℃

increase in winter SST in the Gulf of Alaska increased the return rate by 36% in the Sea of Okhotsk, and 25%

in the Honshu Sea of Japan. In particular, the positive effect of winter SST was greatest in the Hokkaido Sea

of Japan, where a 1℃ increase in the North Pacific increased the return rate by 98%. In contrast, summer

SST in the Sea of Okhotsk reduced the return rate for the Hokkaido Sea of Japan by 30%. In the Honshu Sea

of Japan, SST at release had a significant negative effect, reducing the return rate by 41%. Yellowtail also

had a negative effect in the Sea of Okhotsk, reducing the return rate by 18%. On the other hand, weight at

release had a positive effect in 2 areas, and the effect was much greater in the Hokkaido and Honshu Pacific,

where a 0.1g increase in size at release increased the return rate by 82%, than in the Honshu Sea of Japan,

where the increase was 10%. In the Hokkaido Sea of Japan, weight at release had a negative effect, reducing

the return rate by 34%. The abundance of chum salmon in Russia had a positive effect on chum salmon in

the Sea of Okhotsk, but it was small, increasing the return rate by 3% per million chum salmon caught.

3.7| Geographical patterns of allele frequencies

LDH-A1*100 allele frequencies (Figure 10) were distributed along a north-to-south gradient. A significant

latitudinal gradient was observed in the Russian and American samples, while the allele frequencies in

Etorofu Island (Figure S3) and Japan were distributed outside the geographic gradient and were similar to

those in the northernmost western Alaska and Yukon River populations. LDH-A1*100 allele frequencies

were higher in warmer areas and lower in colder environments. On the other hand, the frequency of LDH-

B2*100 allele was 1.0 or close to 1.0 for all samples (Figure S19).

Except for Japan, GHII3129 SNP allele frequencies (Figure 10) also showed a latitudinal gradient from north

to south. In contrast, GnRH373 allele frequencies (Figure S20) were at relatively similar levels across the

Pacific, but those of Japan were the highest among the populations. The allele of mtDNA3 (Figure S20) was

fixed at close to 1.0 in American and Russian populations, whereas the allele frequencies were significantly

lower in Japanese populations.

4 | DISCUSSION

We provide initial evidence that the sudden increase in the number of returning chum salmon in 2022 is due

to an increase in age 4 fish, leading to higher return rates for the 2019 release. This shift in age composition

towards younger age at maturity is observed in all areas, particularly pronounced in the Hokkaido Sea of

Japan and the Sea of Okhotsk. Our results are consistent with evidence of body size decline primarily

resulting from younger age at maturity from large-scale analyses throughout Alaska (Lewis et al., 2015; Oke

et al., 2020; Ohlberger et al., 2023). North Pacific February-April SST during the first overwintering period

had the largest positive effect on return rates, and Gulf of Alaska January SST during the second

overwintering period also had a positive effect on return rates. In coastal areas, summer SST, SST at release

and a northward expansion of piscivorous fish had a negative effect. Our regression analysis did not infer
causal effects of younger age at maturity (age-specific return rates) on the environmental variables but did
infer causal effects of total return rates. LDH-A1*100 (LDH-A$_4^a$) and GHII3129 allele frequencies in Japan
were outside the latitudinal gradient and close to those in western Alaska and the Yukon, while the allele
frequencies of the GnRH373 and mtDNA3 SNPs were significantly higher and lower than other populations.
In contrast, LDH-B2*100 allele frequencies are fixed at or near 1.0 in all populations.

On Etorofu Island, near Shiretoko Peninsula, Hokkaido, LDH-A1*100 allele frequencies (Figure 10) were at
the similar level to those in Japan. Japan started salmon hatcheries on Etorofu Island in the 1890s and operated
10 hatcheries before World War II (Yagisawa, 1970). Between 1920 and 1944, the Etorofu hatchery, the most
representative, produced 654 million fertilised eggs from 495 thousand parent chum salmon and released 602
million fries. Currently, Russia operates the largest chum salmon hatchery programme on this island, with 14
salmon hatcheries producing 240-250 million juveniles per year, and almost all chum salmon returning to the
island are hatchery fish (Klovach et al., 2021). Japanese salmon seed production methods and facilities have
been inherited in the island. The LDH-A1*100 allele frequencies obtained from hatchery and wild chum
salmon may be due to intensive hatchery releases in the smaller population.

In Japanese hatcheries, fertilised eggs are generally kept in incubators for about 2 months until hatching, and
hatched larvae are reared in indoor rearing ponds for about 2 months until emergence, both in a very dark
environment with calm water flow (Salmon and Freshwater Research Institute, 2023). Metabolic activity in
Atlantic salmon (*Salmo salar*, Salmonidae) was significantly higher in daylight than in darkness, and the
daily oxygen consumption was correlated with sunrise and day length (Maxime et al., 1989). The blind, cave-
dwelling form of the Mexican tetra (*Astyanax mexicanus*, Characidae), which lives in complete darkness,
consumes far less oxygen than the surface morph, and uses ~27% less energy per day (reviewed by Boggs &
Gross, 2021). The evidence suggests that the rearing environment from eggs to emergence (calm water flow
in the dark in indoor ponds for about 4 months) may reduce metabolic activity and oxygen consumption in
hatchery fish, leading to selection at the LDH-A1*100 and mtDNA3 loci. Hatchery steelhead (*O. mykiss*,
Salmonidae) grow significantly faster in hatcheries (Blouin et al., 2021). Hatchery reared chum salmon are
fed daily formulated diets and must grow with reduced metabolic activity where growth hormone may have
a significant effect, potentially altering GHII3129 allele frequencies. Efforts to increase the early-run
population since the early 1980s have induced changes in run timing distribution and the late-run populations
have almost disappeared in Hokkaido, the main production area (Miyakoshi et al., 2013). The strong selection
of early-run salmon could result in GnRH373 allele frequencies above Pacific Rim levels. On the contrary,
LDH-B2*100, which is associated with cardiac activity (Powers et al., 1991; Somero, 2004), may be
conservative even under anthropogenic selection (Figure S19). The deviation of Japanese chum salmon allele
frequencies in LDH-A1*100, GHII3129, GnRH373 and mtDNA3 (Figures 10, S20) from the geographic
pattern could occur during hatchery rearing by relaxing natural selection (Heath et al., 2003; Araki et al.,

2008). Although LDH and SNP allele frequencies were estimated from old samples collected in 1986-2006,

allele frequencies are expected to be similar or more influenced as hatchery releases continue at similar levels

(Figure 4).

The LDH-A1*100 and GHII3129 allele frequencies show countergradient variation (CnGV) (Conover, 1995).

CnGV is a negative covariance between genotypic and environmental effects on phenotype (Cov (G,E) < 0)

and occurs when the genotypic influence on the phenotype counteracts the environmental influence, so that

phenotypic variation between environments is minimised (Conover, 1995; Albecker et al., 2022). CnGV has

been reported for several traits in several species (Conover, 1995), including Atlantic salmon for growth and

digestion (Nicieza et al., 1994) and maturation (Åsheim et al., 2023). CnGV of growth rate is one of the most

common examples (Wikipedia, 2023). Adaptation to cold environments often requires an increase in growth

rate to compensate for the slowdown in growth and metabolism that occurs at lower temperatures (Åsheim

et al., 2023). In other words, during the short growing season, northern fish must grow much faster than

southern fish (Conover, 1990). CnGV adjusts metabolism and growth in the latitudinal cline, minimising

variation in salmon body size. Salmon are highly migratory and populations from different countries spend

several years in the high seas with similar water temperatures. CnGV in metabolism (LDH-A1*100) and

growth (GHII3129) could therefore be generated during their early life stages in freshwater and coastal waters.

LDH-A1*100 (LDH-A_4^a) and GHII3129 allele frequencies outside the latitudinal gradient and close to those

in western Alaska and the Yukon (Figure 10) suggest that Japanese chum salmon may have similar metabolic

and growth characteristics to Alaskan chum salmon, where younger age at maturity has been found in large-

scale analyses (Lewis et al., 2015; Oke et al., 2020; Ohlberger et al., 2023). LDH-A_4 is cold-adapted (Fields

& Somero, 1998). In a common environment experiment in the temperate freshwater fish, the fathead

minnow (*Pimephales promelas*, Cyprinidae) (Supplementary Methods), Merritt (1972) was found that the

homozygote of the northern allele LDH-A_4^{bb} had significantly higher Michaelis constant of pyruvate, K_m^{PYR}

(lower substrate affinities=lower catalytic efficiency) than the homozygote of the southern allele LDH-A_4^{aa}

and the heterozygote LDH-A_4^{ab} at 25°C and above. The result shows that individuals homozygous for the

northern allele LDH-A_4^{bb} require more energy (food) to grow at warmer temperatures. Although we could

not find an experimental study in salmonids to assess genotype effects on catalytic efficiency, Merritt's result

suggests that contemporary Japanese chum salmon, which may have similar metabolic efficiency to Alaskan

chum salmon, should have lower catalytic efficiency and consume more energy at warmer temperatures along

the Japanese coast. The release of Japanese chum salmon is equivalent to the release of Alaskan chum salmon

in a southern environment. This results in a common environmental experiment for growth and metabolism

with higher water temperatures in terms of LDH-A1 and GHII3129 loci. As a result, contemporary Japanese

chum salmon grow faster but are less metabolically efficient than native Japanese populations with original

allele frequencies adapted to the Japanese environment (now unobserved). In the Sanriku Coast, Honshu

Pacific, Iino et al. (2022) found that juvenile chum salmon reared under low food availability (1% body

weight) allocated less than half of their energy to growth at high temperatures (14°C) compared to juveniles reared under high food availability (4% body weight). The similar LDH-A1*100 and GHII3129 allele frequencies to Alaskan chum salmon may require much food to grow at high temperatures. The long distance migration from the Pacific coast of Honshu to the Sea of Okhotsk (Figure 6b) may also have a negative effect on juvenile growth and survival. This may explain the negative effect of SST at release and summer SST along the coast of the Sea of Okhotsk.

In a warmer climate, the body size of adult salmon will decrease, mainly due to a younger age at maturity (Jonsson, 2023). Younger maturing salmon grow much faster than older maturing salmon (Ricker, 1981). Growth history of salmon was more strongly associated with the probability of maturation than age or age-specific body size, and the most recent growth condition was the most important factor in determining whether a fish matured in the following breeding season (Morita & Fukuwaka, 2006). The growth history of Japanese chum salmon at early life stages with increased growth rate may result in better growth and younger age at maturity during warm winter SST in the wintering grounds, as seen in Alaskan sockeye salmon (Ohlberger et al., 2023). In the North Pacific and Gulf of Alaska, mean January SST was the same at $6.0 \pm 0.7°C$, and mean February SST was slightly lower at $5.6 \pm 0.7°C$ in the Gulf of Alaska (Figures 2,3). The optimum temperature for chum salmon growth has been estimated to be 8-12°C (Kaeriyama, 1986; Urawa et al., 2019; Kaeriyama, 2022). Warm winter SST in the wintering grounds (Figure 3) may result in better growth of Japanese chum salmon, which have similar metabolic and growth characteristics to Alaska salmon, when food is abundant. The effect of food abundance remains unknown and is left for future study.

In the Sea of Okhotsk, yellowtail reduced the return rate by 18% (Figure 9). Young yellowtail and mahi-mahi (*Coryphaena hippurus*) were found in chum salmon landings in the Sea of Okhotsk in September 2012 (Figure S21). Since 2014, yellowtail have been caught along the coast of the Sea of Okhotsk in May-June (Hoshino & Fujioka, 2021), coinciding with the release period. Another candidate predator is a seabird, the rhinoceros auklet (*Cerorhinca monocerata*), which preys on juvenile chum salmon off the Pacific coast of Hokkaido from late June to mid-July, with an estimated consumption of 317 tonnes (31.7 million juveniles, assuming 10 g BW per fish) in 2014 (Okado et al., 2020). In freshwater, brown trout (*Salmo trutta*) collected near the hatchery and ~10 km downstream predated 49-481 chum salmon juveniles in the Chitose River, Hokkaido Sea, Japan (Honda et al., 2023). Piscivorous salmonids chase schools of juvenile fish, and small juvenile chum salmon are easily predated due to their slow swimming speed to escape predators (Hasegawa et al., 2021). Before 1945, wild-born chum salmon fry were too agile to be netted in the rivers of Etrofu Island, where Japan operated salmon hatcheries (Yagisawa, 1970). A significant reduction in mtDNA3 SNP allele frequency in Japanese chum salmon (Figure S20) could reduce the efficiency of aerobic fitness and swimming endurance, energy metabolism and oxygen consumption (Kitada & Kishino, 2021). The similar LDH-A1*100 allele frequencies to Alaskan chum salmon may reduce burst swimming speed in warming SST along Japanese coasts, as LDH-A is predominantly found in skeletal (white) muscle (Powers et al., 1991;

Somero, 2004). The effect of predation by piscivorous fish can be significant when many slow swimming

juveniles are released at the same time. Predation by large marine animals, such as killer whales (Ohlberger

et al., 2019), may also occur in the high seas.

The positive effect of weight at release was greatest in the Hokkaido and Honshu Pacific, where a 0.1 g

increase in weight at release increased the return rate by 82% (Figure 9). In the Honshu Sea of Japan, the

effect of weight at release was smaller; a 0.1 g increase in weight at release increased the return rate by 10%,

while a 1°C increase in SST at release reduced the return rate by 41%. The positive effect of weight at release

could be reduced by the effect of increasing SST. Unexpectedly, the effect of weight at release was negative

in the Hokkaido Sea of Japan, where a 0.1 g increase in weight at release reduced the return rate by 34%,

suggesting that the current weight at release (1.04 ± 0.03 g, average 2016-2020) is sufficient in this region.

Our regression analysis assumed that all chum salmon returning to Japan are hatchery-reared. On the other

hand, field studies provide evidence that natural reproduction of chum salmon occurs in 31%-38% of 238

non-enhanced rivers in Hokkaido (Miyakoshi et al., 2012), and in 94% of 47 hatchery-enhanced rivers and

75% of 47 non-enhanced rivers along the northern Honshu Sea coast of Japan (Iida et al., 2018; Iida et al.,

2021). Escapement of returned fish is not a common practice in Hokkaido, although all broodstock collection

weirs have been redesigned to allow passage of fish for natural spawning in the Kitami region, Sea of Okhotsk

since 2005 (Kitada, 2014). In the Honshu Sea of Japan, about half of the streams from which hatchery

broodstock are collected use small nets or hooks that do not interfere with the upstream migration of chum

salmon (Iida et al., 2018). Hatchery fish are stocked in most large rivers (Miyakoshi et al., 2013), while

naturally spawning chum salmon populations persist in many small (short) rivers in Hokkaido, the main

production area (Kitada, 2014). The very high gene flow between populations allows genetic admixture of

hatchery-born fish and wild-born hatchery progeny throughout the Japanese populations (Kitada & Kishino,

2021). The relative reproductive success (RRS) of hatchery-reared salmon is approximately 50% of that of

wild-origin fish spawning in the wild (Araki et al., 2007; Christie et al., 2014; Shedd et al., 2022), with

considerable variation (Kitada et al., 2011). Moreover, reduced RRS can be carried over to the next generation

(Araki et al., 2009). Egg-to-size-at-release survival of wild-born fish in natural streams should be very low.

Although the magnitude of recruitment from natural reproduction was unknown, it may not be substantial

compared to the number of juveniles released (1-1.5 billion juveniles throughout Japan).

Using the world's largest hatchery release programme, we show how salmon populations are exposed to

environmental and anthropogenic stressors at both local and North Pacific-wide scales. The geographic

pattern of gene allele frequencies suggests that Japanese chum salmon have similar metabolic and growth

characteristics to Alaskan chum salmon, with unique features of maturation, aerobic endurance, energy

metabolism and oxygen consumption. We hypothesise that intentional and unintentional hatchery selection

on Japanese chum salmon have relaxed natural selection and altered allele frequencies, reducing survival in

coastal Japan, while improving growth in the first and second winters in warming climate, thereby promoting
a younger age at maturity. The apparent increase in return rates of Japanese chum salmon is due to their
returning one year earlier. This is a trade-off between fish abundance and body size decline. However, the
underlying mechanisms for the decline in body size and younger age at maturity are largely unknown and
remain the subject of future research. Although more research is clearly needed to fully understand the
underlying mechanisms of Japanese chum salmon population dynamics, our findings provide a new insight
into salmon conservation and hatchery release.
714

ACKNOWLEDGMENTS
We sincerely acknowledge the efforts of thousands of biologists, fishery and hatchery managers, and fishers
who contributed to the collection of the data we analyse in this manuscript. This study would not have been
possible without the efforts of APG, FRA, HSEA, JMA, MAFF, NOAA, and NPAFC in creating the databases
for open use of the data. This study was supported by Japan Society for the Promotion of Science Grants-in-
Aid for Scientific Research KAKENHI No. 18K0578116 for SK and 22K11950 for HK.
721

722 **DATA AVAILABILITY STATEMENT**
723 The data generated or analysed in this study are fully described in the published article and its Supplementary
724 Information including Supplementary Figures S1-S21. Supplementary Tables S1-S10, Supplementary Data
725 S1-S7 and the analysis scripts and data are available at https://doi.org/10.5281/zenodo.8394732.
726

REFERENCES
Aebersold, P. B., Winans, G. A., Teel, D. J., Milner, G. B., & Utter, F. M. (1987). Manual for starch gel
electrophoresis: a method procedures for detection of genetic variation. *NOAA Technical Reports*, 61,
19 p.
Akaike, H. (1973). Information theory as an extension of the maximum likelihood principle. In B. N. Petrov
& F. Csaki (Eds.), *Second international symposium on information theory* (pp. 267–281). Budapest:
Akademiai Kiado.
Albecker, M. A., Trussell, G. C., & Lotterhos, K. E. (2022). A novel analytical framework to quantify co-
gradient and countergradient variation. *Ecology Letters*, 25, 1521–1533.
https://doi.org/10.1111/ele.14020
Amoroso, R. O., Tillotson, M. D., Hilborn, R. (2017). Measuring the net biological impact of fisheries
enhancement: Pink salmon hatcheries can increase yield, but with apparent costs to wild populations.
Canadian Journal of Fisheries and Aquatic Sciences, 74, 1233–1242. https://doi.org/10.1139/cjfas-
2016-0334
Araki, H., Cooper, B., & Blouin, M. S. (2007). Genetic effects of captive breeding cause a rapid, cumulative
fitness decline in the wild. *Science*, 318, 100–103. https://doi.org/10.1126/science.1145621
Araki, H., Berejikian, B. A., Ford, M. J., & Blouin, M. S. (2008). Fitness of hatchery-reared salmonids in the

wild. *Evolutionary Applications*, 1, 342–355. https://doi.org/10.1111/j.1752-4571.2008.00026.x

Araki, H., Cooper, B., & Blouin, M. S. (2009). Carry-over effect of captive breeding reduces reproductive

fitness of wild-born descendants in the wild. *Biology Letters*, 5, 621–624.

https://doi.org/10.1098/rsbl.2009.0315

Åsheim, E. R., Debes, P. V., House, A., Liljeström, P., Niemelä, P. T., Siren, J. P., … & Primmer, C. R. (2023).

Atlantic salmon (*Salmo salar*) age at maturity is strongly affected by temperature, population and age-

at-maturity genotype. *Conservation Physiology*, 11, coac086. https://doi.org

/10.1093/conphys/coac086

Atlas, W. I., Wilson, K. L., Whitney, C. K., Moody, J. E., Service, C. N., Reid, M., & Sloat, M. R. (2022).

Quantifying regional patterns of collapse in British Columbia Central Coast chum salmon

(*Oncorhynchus keta*) populations since 1960. *Canadian Journal of Fisheries and Aquatic Sciences*, 79,

2072–2086. https://doi.org /10.1139/cjfas-2022-0013

Beacham, T. D., Sato, S., Urawa, S., Le, K. D., & Wetklo, M. (2008). Population structure and stock

identification of chum salmon Oncorhynchus keta from Japan determined by microsatellite DNA

variation. *Fisheries Science*, 74, 983–994. https://doi.org/10.1111/j.1444-2906.2008.01616.x

Beamish, R. (2022). The need to see a bigger picture to understand the ups and downs of Pacific salmon

abundances. *ICES Journal of Marine Science*, 79, 1005–1014. https://doi.org/10.1093/icesjms/fsac036

Beamish, R. J., & Mahnken, C. (2001). A critical size and period hypothesis to explain natural regulation of

salmon abundance and the linkage to climate and climate change. *Progress in Oceanography*, 49,

423–437. https://doi.org/10.1016/S0079-6611(01)00034-9

Björnsson, B. T. (1997). The biology of salmon growth hormone: from daylight to dominance. Fish

Physiology and Biochemistry, 17, 9–24. https://doi.org /10.1023/A:1007712413908

Blouin, M. S., Wrey, M. C., Bollmann, S. R., Skaar, J. C., Twibell, R. G., & Fuentes, C. (2021). Offspring of

first-generation hatchery steelhead trout (*Oncorhynchus mykiss*) grow faster in the hatchery than

offspring of wild fish, but survive worse in the wild: Possible mechanisms for inadvertent

domestication and fitness loss in hatchery salmon. *Plos One*, 16, e0257407.

https://doi.org/10.1371/journal.pone.0257407

Boggs, T., & Gross, J. (2021). Reduced oxygen as an environmental pressure in the evolution of the blind

Mexican cavefish. *Diversity*, 13, 26. https://doi.org /10.3390/d13010026

Cade, B. S. (2015). Model averaging and muddled multimodel inferences. *Ecology*, 96, 2370–2382.

https://doi.org/10.1890/14-1639.1

Chitose Aquarium (2023, August, 8) Capture Information of chum salmon. Retrieved from https://chitose-

aq.jp/

Christie, M. R., Ford, M. J., & Blouin, M. S. (2014). On the reproductive success of early-generation hatchery

fish in the wild. *Evolutionary Applications*, 7, 883–896. https://doi.org/10.1111/eva.12183

Cline, T. J., Ohlberger, J., & Schindler, D. E. (2019). Effects of warming climate and competition in the ocean

for life-histories of Pacific salmon. *Nature Ecology & Evolution*, 3, 935–942.

781   https://doi.org/10.1038/s41559-019-0901-7

782 Connors, B., Malick, M. J., Ruggerone, G. T., Rand, P., Adkison, M., Irvine, J. R. et al. (2020). Climate and
783   competition influence sockeye salmon population dynamics across the Northeast Pacific Ocean.
784   *Canadian Journal of Fisheries and Aquatic Sciences*, 77, 943–949. https://doi.org/10.1139/cjfas-2019-
785   0422

786 Conover, D. O. (1990). The relation between capacity for growth and length of growing season: evidence for
787   and implications of countergradient variation. *Transactions of the American Fisheries Society*, 119,
788   416–430.https://doi.org/10.1577/1548-8659(1990)119%3C0416:TRBCFG%3E2.3.CO;2

789 Conover, D. O., & Schultz, E. T. (1995). Phenotypic similarity and the evolutionary significance of
790   countergradient variation. *Trends in Ecology & Evolution*, 10, 248–252. https://doi.org
791   /10.1016/S0169-5347(00)89081-3

792 Cunningham, C. J., Westley, P. A., & Adkison, M. D. (2018). Signals of large scale climate drivers, hatchery
793   enhancement, and marine factors in Yukon River Chinook salmon survival revealed with a Bayesian
794   life history model. *Global Change Biology*, 24, 4399–4416. https://doi.org/10.1111/gcb.14315

795 Daly, E. A., Moss, J. H., Fergusson, E., & Debenham, C. (2019). Feeding ecology of salmon in eastern and
796   central Gulf of Alaska. Deep Sea Research Part II: *Topical Studies in Oceanography*, 165, 329–339.

797 Davis, N. D., Myers, K. W., & Ishida, Y. (1998). Caloric value of high-seas salmon prey organisms and
798   simulated salmon ocean growth and prey consumption. *North Pacific Anadromous Fish Commission*
799   *Bulletin*, 1, 146–162.

800 Dunmall, K. M., McNicholl, D. G., Zimmerman, C. E., Gilk-Baumer, S. E., Burril, S., & von Biela, V. R.
801   (2022). First juvenile chum salmon confirms successful reproduction for Pacific salmon in the North
802   American Arctic. *Canadian Journal of Fisheries and Aquatic Sciences*, 79, 703–707.
803   https://doi.org/10.1139/cjfas-2022-0006

804 Fields, P. A., & Somero, G. N. (1998). Hot spots in cold adaptation: localized increases in conformational
805   flexibility in lactate dehydrogenase A4 orthologs of Antarctic notothenioid fishes. *Proceedings of the*
806   *National Academy of Sciences*, 95, 11476–11481. https://doi.org /10.1073/pnas.95.19.11476

807 Fisheries Research Agency (FRA). (2023, July, 7). Salmon return in 2022 (7[th] report: as of 28 Feb) Salmon
808   age composition and body size (Tentative translation). (in Japanese) Retrieved from
809   http://salmon.fra.affrc.go.jp/zousyoku/salmon/

810 Groot, C., & Margolis, L. (1991). *Pacific Salmon Life Histories*. Vancouver: University of British Columbia
811   Press,.

812 Hasegawa, K., Honda, K., Yoshiyama, T., Suzuki, K., & Fukui, S. (2021). Small biased body size of salmon
813   fry preyed upon by piscivorous fish in riverine and marine habitats. *Canadian Journal of Fisheries and*
814   *Aquatic Sciences*, 78, 631-638. https://doi.org/10.1139/cjfas-2020-0339

815 Hayashizaki, K. I., & Ida, H. (1993). Size decrease of chum salmon, Oncorhynchus keta. Tohoku districts,
816   Japan. *International Symposium on Biological Interactions of Enhanced and Wild Salmonids*. In B.
817   Riddell (Ed.), *Biological Interactions of Enhanced and Wild Salmonids*. Canadian Special Publication

of Fisheries and Aquatic Sciences.

Heath, D. D., Heath, J. W., Bryden, C. A., Johnson, R. M., & Fox, C. W. (2003). Rapid evolution of egg size

in captive salmon. *Science*, 299, 1738–1740. https://doi.org/10.1126/science.1079707

Hilborn, R., Quinn, T. P., Schindler, D. E., & Rogers, D. E. (2003). Biocomplexity and fisheries sustainability.

Proceedings of the National Academy of Sciences, 100, 6564–6568.

https://doi.org/10.1073/pnas.1037274100

Hiroi, O. (1998). Historical trends of salmon fisheries and stock conditions in Japan. *North Pacific*

Anadromous Fish Commission Bulletin, 1, 23–27

Honda, K., Kawakami, T., Suzuki, K., Watanabe, K., & Saito, T. (2017). Growth rate characteristics of

juvenile chum salmon *Oncorhynchus keta* originating from the Pacific coast of Japan and reaching

Konbumori, eastern Hokkaido. *Fisheries Science*, 83, 987–996. https://doi.org/10.1007/s12562-017-

1137-6

Honda, K., Shirai, K., Komatsu, S., & Saito, T. (2020). Sea-entry conditions of juvenile chum salmon

Oncorhynchus keta that improve post-sea-entry survival: a case study of the 2012 brood-year stock

released from the Kushiro River, eastern Hokkaido, Japan. *Fisheries Science*, 86, 783–792.

https://doi.org/10.1007/s12562-020-01442-0

Honda, K., Sato, T., Kuroda, H., & Saito, T. (2021). Initial growth characteristics of poor-return stocks of

chum salmon *Oncorhynchus keta* originating from the Okhotsk and Nemuro regions in Hokkaido on

the basis of scale analysis. *Fisheries Science*, 87, 653–663. https://doi.org/10.1007/s12562-021-01538-

1

Honda, K., Hasegawa, K., Ono, I., & Miyashita, K. (2023). Piscivorous brown trout Salmo trutta does not

migrate from distant downstream habitats to a massive release site for chum salmon *Oncorhynchus*

keta fry in the Chitose River, northern Japan. *Environmental Biology of Fishes*, 106, 707–715.

https://doi.org/10.1007/s10641-023-01402-4

Hoshino, N., & Fujioka, T. (2021). Characteristics of yellowtail catch in Hokkaido, Japan in the 2010s.

Scientific Report of Hokkaido Fisheries Research Institute, 100, 71–82. (in Japanese)

Iida, M., Yoshino, K., & Katayama, S. (2018). Current status of natural spawning of chum salmon

Oncorhynchus keta in rivers with or without hatchery stocking on the Japan Sea side of northern

Honshu, Japan. *Fisheries science*, 84, 453–459. https://doi.org/10.1007/s12562-018-1192-7

Iida, M., Yagi, Y., & Iseki, T. (2021). Occurrence of wild chum salmon fry in the surf zone, and spawning

and emergence timing in the adjacent nonstocked river in Niigata Prefecture, Japan. *Fisheries Science*,

87, 549–557. https://doi.org/10.1007/s12562-021-01535-4

Iino, Y., Kitagawa, T., Abe, T. K., Nagasaka, T., Shimizu, Y., Ota, K., ... & Kawamura, T. (2022). Effect of

food amount and temperature on growth rate and aerobic scope of juvenile chum salmon. *Fisheries*

Science, 8, 397–409. https://doi.org/10.1007/s12562-022-01599-w

Ishida, Y., Ito, S. O., Kaeriyama, M., McKinnell, S., & Nagasawa, K. (1993). Recent changes in age and size

of chum salmon (Oncorhynchus keta) in the North Pacific Ocean and possible causes. *Canadian*

Journal of Fisheries and Aquatic Sciences, 50, 290–295. https://doi.org/10.1139/f93-033

Ivankova, E. V., & Efremov, V. V. (2009). Geographic variation and temporal population differentiation of

chum salmon Oncorhynchus keta from some regions of the Russian Far East. *Russian Journal of*

Genetics, 45, 715–725. https://doi.org/10.1134/S1022795409060118

James, S. E., Doherty, B., Cox, S. P., Pearsall, I. A., & Riddell, B. (2023). Size and timing of hatchery releases

influence juvenile-to-adult survival rates of British Columbia Chinook (*Oncorhynchus tshawytscha*)

and coho (*Oncorhynchus kisutch*) salmon. *Canadian Journal of Fisheries and Aquatic Sciences*, 80,

700–718. https://doi.org/10.1139/cjfas-2022-0121

Jeffrey, K. M., Côté, I. M., Irvine, J. R., & Reynolds, J. D. (2017). Changes in body size of Canadian Pacific

salmon over six decades. *Canadian Journal of Fisheries and Aquatic Sciences*, 74, 191–201.

https://doi.org /10.1139/cjfas-2015-0600

Jonsson, B. (2023). Thermal effects on eEcological traits of salmonids. *Fishes*, 8, 337. https://doi.org

/10.3390/fishes8070337

Kaeriyama, M. (1989). Aspects of salmon ranching in Japan. *Physiology & Ecology, Japan*, Special Volume

1, 625–638.

Kaeriyama, M. (1999). Hatchery programmes and stock management of salmonid populations in Japan. In

B. R. Howell, E. Moksness, & T. Svåsand (Eds.), *Stock enhancement and sea ranching* (pp. 153–167).

Oxford, UK: Blackwell.

Kaeriyama, M. (2022). Warming climate impacts on production dynamics of southern populations of Pacific

salmon in the North Pacific Ocean. *Fisheries Oceanography*, 32, 121–132.

https://doi.org/10.1111/fog.12598

Kaeriyama, M., & Edpalina, R. R. (2004). Evaluation of the biological interaction between wild and hatchery

populations for sustainable fisheries and management of Pacific salmon. In K. Leber, S. Kitada, H. L.

Blankenship, & T. Svåsand (Eds.), *Stock enhancement and sea ranching, developments pitfalls and*

opportunities (pp. 247–259). Oxford, UK: Blackwell.

Kaeriyama, M., Nakamura, M., Edpalina, R., Bower, J. R., Yamaguchi, H., Walker, R. V., & Myers, K. W.

(2004). Change in feeding ecology and trophic dynamics of Pacific salmon (*Oncorhynchus* spp.) in the

central Gulf of Alaska in relation to climate events. *Fisheries Oceanography*, 13, 197–207.

https://doi.org/10.1111/j.1365-2419.2004.00286.x

Karabanov, D. P. & Kodukhova, Y. V. (2018). Biochemical polymorphism and intraspecific structure in

populations of Kilka *Clupeonella cultriventris* (Nordmann, 1840) from natural and invasive parts of

its range. *Inland Water Biology*, 11, 496–500. https://doi.org /10.1134/S1995082918040107

Khakoo, Z., Bhatia, A., Gedamu, L., & Habibi, H. R. (1994). Functional specificity for salmon gonadotropin-

releasing hormone (GnRH) and chicken GnRH-II coupled to the gonadotropin release and subunit

messenger ribonucleic acid level in the goldfish pituitary. *Endocrinology*, 134, 838–847.

https://doi.org/10.1210/en.134.2.838

Kitada, S. (2014). Japanese chum salmon stock enhancement: current perspective and future challenges.

Fisheries Science, 80, 237–249. https://doi.org/10.1007/s12562-013-0692-8

Kitada, S. (2018). Economic, ecological and genetic impacts of marine stock enhancement and sea ranching:

A systematic review. *Fish and Fisheries*, 19, 511–532. https://doi.org/10.1111/faf.12271

Kitada, S. (2020). Lessons from Japan marine stock enhancement and sea ranching programmes over 100

years. *Reviews in Aquaculture*, 12, 1944–1969. https://doi.org/10.1111/raq.12418

Kitada, S., & Kishino, H. (2021). Population structure of chum salmon and selection on the markers collected

for stock identification. *Ecology and Evolution*, 11, 13972–13985. https://doi.org/10.1002/ece3.8102

Kitada, S., Kishino, H., & Hamasaki, K. (2011). Bias and significance of relative reproductive success

estimates based on steelhead trout (*Oncorhynchus mykiss*) data: a Bayesian meta-analysis. *Canadian*

Journal of Fisheries and Aquatic Sciences, 68, 1827–1835. https://doi.org/10.1139/f2011-087

Klovach, N., Leman, V., & Gordeev, I. (2021). The relative importance of enhancement to the production of

Salmon on Iturup Island (Kuril Islands, Russia). *Reviews in Aquaculture*, 13, 664–675. https://doi.org

/10.1111/raq.12493

Kobayashi, T. (1980). Salmon propagation in Japan. In J. E. Thorpe (Ed.), *Salmon ranching* (pp. 91–107).

London, UK: Academic Press.

Kong, M., Xiang, H., Wang, J., Liu, J., Zhang, X., & Zhao, X. (2020). Mitochondrial DNA Haplotypes

Influence Energy Metabolism across Chicken Transmitochondrial Cybrids. *Genes*, 11, 100.

https://doi.org /10.3390/genes 11010100

Kuroda, H., Saito, T., Kaga, T., Takasuka, A., Kamimura, Y., Furuichi, S., & Nakanowatari, T. (2020).

Unconventional sea surface temperature regime around Japan in the 2000s-2010s: Potential influences

on major fisheries resources, *Frontiers in Marine Science*, 7, 574904.

https://doi.org/10.3389/fmars.2020.574904

Lewis, B., Grant, W. S., Brenner, R. E., & Hamazaki, T. (2015). Changes in size and age of Chinook salmon

Oncorhynchus tshawytscha returning to Alaska. *Plos One*, 10, e0130184.

https://doi.org/10.1371/journal.pone.0130184

Liang, X., Liu, L., Fu, T., Zhou, Q., Zhou, D., Xiao, L., … & Gan, Z. (2016). Exercise inducible lactate

dehydrogenase B regulates mitochondrial function in skeletal muscle. *Journal of Biological Chemistry*,

291, 25306–25318. https://doi.org /10.1074/jbc.M116.749424

Losee, J. P., Kendall, N. W., & Dufault, A. (2019). Changing salmon: An analysis of body mass, abundance,

survival, and productivity trends across 45 years in Puget Sound. *Fish and Fisheries*, 20, 934–951.

https://doi.org/10.1111/faf.12385

Markert, C. L., & Møller, F. (1959). Multiple forms of enzymes: tissue, ontogenetic, and species specific

patterns. *Proceedings of the National Academy of Sciences of the United States of America*, 45, 753–

763. https://doi.org /10.1073/pnas.45.5.753

Maxime, V., Boeuf, G., Pennec, J. P., & Peyraud, C. (1989). Comparative study of the energetic metabolism

of Atlantic salmon (Salmo salar) parr and smolts. *Aquaculture*, 82(1–4), 163–171. https://doi.org

/10.1016/0044-8486(89)90405-5

Merritt, R. B. (1972). Geographic distribution and enzymatic properties of lactate dehydrogenase allozymes in the fathead minnow, *Pimephales promelas*. *American Naturalists*, 106, 173–184. https://doi.org /10.1086/282760

Miyakoshi, Y., Urabe, H., Saneyoshi, H., Aoyama, T., Sakamoto, H., Ando et al. (2012). The occurrence and run timing of naturally spawning chum salmon in northern Japan. *Environmental Biology of Fishes*, 94, 197–206. https://doi.org/10.1007/s10641-011-9872-5

Miyakoshi, Y., Nagata, M., Kitada, S., & Kaeriyama, M. (2013). Historical and current hatchery programs and management of chum salmon in Hokkaido, northern Japan. *Reviews in Fisheries Science*, 21, 469–479. https://doi.org/10.1080/10641262.2013.836446

Morita, K., & Fukuwaka, M. A. (2006). Does size matter most? The effect of growth history on probabilistic reaction norm for salmon maturation. *Evolution*, 60, 1516–1521. https://doi.org //10.1111/j.0014-3820.2006.tb01230.x

Morita, K., & Fukuwaka, M. A. (2007). Why age and size at maturity have changed in Pacific salmon. Marine Ecology Progress Series, 335, 289-294. https://doi.org /10.3354/meps335289

Mueter, F. J., Peterman, R. M., & Pyper, B. J. (2002). Opposite effects of ocean temperature on survival rates of 120 stocks of Pacific salmon (*Oncorhynchus* spp.) in northern and southern areas. *Canadian Journal of Fisheries and Aquatic Sciences*, 59, 456–463. https://doi.org/10.1139/f02-020

Myers, K. W., Klovach, N. V., Gritsenko, O. F., Urawa, S., & Royer, T. C. (2007). Stock-specific distributions of Asian and North American salmon in the open ocean, interannual changes, and oceanographic conditions. *North Pacific Anadromous Fish Commission Bulletin*, 4, 159–177.

Myers, K. W., Irvine, J. R., Logerwell, E. A., Urawa, S., Naydenko, S. V., Zavolokin, A. V., & Davis, N. D. (2016). Pacific salmon and steelhead: life in a changing winter ocean. *North Pacific Anadromous Fish Commission Bulletin*, 6, 113–138.

Nagata, M., Miyakoshi, Y., Ando, D., Fujiwara, M., Sawada, M., Shimada, H., & Asami, H. (2007). Influence of coastal seawater temperature on the distribution and growth of juvenile chum salmon, with recommendations for altered release strategies. *North Pacific Anadromous Fish Commission Bulletin*, 4, 223–235.

Naish, K. A., Taylor III, J. E., Levin, P. S., Quinn, T. P., Winton, J. R., Huppert, D., & Hilborn, R. (2007). An evaluation of the effects of conservation and fishery enhancement hatcheries on wild populations of salmon. *Advances in Marine Biology*, 53, 61–194. https://doi.org/10.1016/S0065-2881(07)53002-6

Nicieza, A. G., Reiriz, L., & Braña, F. (1994). Variation in digestive performance between geographically disjunct populations of Atlantic salmon: countergradient in passage time and digestion rate. *Oecologia*, 99, 243–251. https://doi.org /10.1007/BF00627736

Nicholls, T. J., & Minczuk, M. (2014). In D-loop: 40 years of mitochondrial 7S DNA. *Experimental Gerontology*, 56, 175–181. https://doi.org /10.1016/j.exger.2014.03.027

Ohlberger, J., Schindler, D. E., Ward, E. J., Walsworth, T. E., & Essington, T. E. (2019). Resurgence of an apex marine predator and the decline in prey body size. *Proceedings of the National Academy of*

Sciences, 116, 26682–26689. https://doi.org/10.1073/pnas.1910930116

Ohlberger, J., Ward, E. J., Brenner, R. E., Hunsicker, M. E., Haught, S. B., Finnoff, D. et al. (2022). Non-stationary and interactive effects of climate and competition on pink salmon productivity. *Global Change Biology*, 28, 2026–2040. https://doi.org/10.1111/gcb.16049

Ohlberger, J., Cline, T. J., Schindler, D. E., & Lewis, B. (2023). Declines in body size of sockeye salmon associated with increased competition in the ocean. *Proceedings of the Royal Society B*, 290, 20222248. https://doi.org/10.1098/rspb.2022.2248

Okado, J., Koshino, Y., Kudo, H., & Watanuki, Y. (2020). Consumption of juvenile chum salmon by a seabird species during early sea life. *Fisheries Research*, 222, 105415. https://doi.org/10.1016/j.fishres.2019.105415

Oke, K. B., Cunningham, C. J., Westley, P. A. H., Baskett, M. L., Carlson, S. M., Clark et al. (2020). Recent declines in salmon body size impact ecosystems and fisheries. *Nature Communications*, 11, 4155. https://doi.org/10.1038/s41467-020-17726-z

Place, A. R., and Powers, D. A. (1979). Genetic variation and relative catalytic efficiencies: lactate dehydrogenase B allozymes of *Fundulus heteroclitus*. *Proceedings of the National Academy of Sciences*, 76: 2354–2358. https://doi.org /10.1073/pnas.76.5.2354

Powers, D. A., Lauerman, T., Crawford, D., & DiMichele, L. (1991). Genetic mechanisms for adapting to a changing environment. *Annual Review of Genetics*, 25: 629–660. https://doi.org /10.1146/annurev.ge.25.120191.003213

Qin, Y., & Kaeriyama, M. (2016). Feeding habits and trophic levels of Pacific salmon (*Oncorhynchus* spp.) in the North Pacific Ocean. *North Pacific Anadromous Fish Commission Bulletin*, 6, 469–481.

Quinn, T. P. (2018). *The behavior and ecology of Pacific salmon and trout*. Second edition, Seattle: University of Washington Press, In association with American Fisheries Society, Bethesda, Maryland.

Ricker, W. E. (1981). Changes in the average size and average age of Pacific salmon. *Canadian Journal of Fisheries and Aquatic Sciences*, 38, 1636–1656. https://doi.org/10.1139/f81-213

Rousset, F. (2008). Genepop'007: A complete re-implementation of the genepop software for Windows and Linux. Molecular Ecology Resources, 8, 103–106. https://doi.org /10.1111/j.1471-8286.2007.01931x.

Saito, T., & Nagasawa, K. (2009). Regional synchrony in return rates of chum salmon (*Oncorhynchus keta*) in Japan in relation to coastal temperature and size at release. *Fisheries Research*, 95, 14–27. https://doi.org/10.1016/j.fishres.2008.07.004

Salmenkova, E. A., Omel'chenko, V. T., Afanas' ev, K. I., & Rubtsova, G. A. (2008). Genetic diversity of Asian chum salmon *Oncorhynchus keta* (Salmonidae, Salmoniformes) and the dynamics of gene pools in Sakhalin populations under artificial reproduction. *Journal of Ichthyology*, 48, 313–325. https://doi.org/10.1134/S0032945208040048

Salmon and Freshwater Research Institute. (SFRI) (2023, October, 9). Salmon hatchery release. (Tentative translation). (in Japanese) Retrieved from https://www.hro.or.jp/list/fisheries/research/hatch/section/shigen/td6oqn000000087r.html

Salo, E. O. (1991). Life history of chum salmon (*Oncorhynchus keta*). In: C. Groot & L. Margolis (Eds.), *Pacific salmon life histories* (pp. 231–309). Vancouver, Canada: UBC Press.

Sato, S., & Urawa, S. (2015). Genetic structure of chum salmon populations in Japan. *Bulletin of Fishery Research Agency*, 39, 21–47. (in Japanese)

Seeb, L. W., Crane, P. A. & Gates, R. B. (1995). Progress Report of Genetic Studies of Pacific Rim Chum Salmon, and Preliminary Analysis of the 1993 and 1994 South Unimak June Fisheries. Anchorage, USA: Alaska Department of Fish and Game.

Seeb, L. W., & Crane, P. A. (1999). High genetic heterogeneity in chum salmon in western Alaska, the contact zone between northern and southern lineages. Transactions of the American Fisheries Society, 128, 58–87. https://doi.org/10.1577/1548-8659(1999)128<0058:HGHICS>2.0.CO;2

Seeb, L. W., Templin, W. D., Sato, S., Abe, S., Warheit, K., Park, J. Y., & Seeb, J. E. (2011). Single nucleotide polymorphisms across a species'range: Implications for conservation studies of Pacific salmon. *Molecular Ecology Resources*, 11(Suppl. 1), 195–217. https://doi.org/10.1111/j.1755-0998.2010.02966.x

Shaklee, J. B., Allendorf, F. W., Morizot, D. C., & Whitt, G. S. (1990). Gene nomenclature for protein-coding loci in fish. *Transactions of the American Fisheries Society*, 119, 2–15. https://doi.org/10.1577/1548-8659(1990)119<0002:GNFPLI>2.3.CO;2

Shedd, K. R., Lescak, E. A., Habicht, C., Knudsen, E. E., Dann, T. H., Hoyt, H. A., … & Templin, W. D. (2022). Reduced relative fitness in hatchery-origin Pink Salmon in two streams in Prince William Sound, Alaska. *Evolutionary Applications*, 15, 429–446. https://doi.org/10.1111/eva.13356

Somero, G. N. (2004). Adaptation of enzymes to temperature: searching for basic "strategies". *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 139, 321–333. https://doi.org/10.1016/j.cbpc.2004.05.003

Somero, G. N. (2010). The physiology of climate change: how potentials for acclimatization and genetic adaptation will determine 'winners' and 'losers'. *Journal of Experimental Biology*, 213, 912–920. https://doi.org/10.1242/jeb.037473

Somero, G. N. (2022). The Goldilocks Principle: A unifying perspective on biochemical adaptation to abiotic stressors in the sea. *Annual Review of Marine Science*, 14, 1-23. https://doi.org/10.1146/annurev-marine-022521-102228

Stefàno, E., Marsigliante, S., Vetrugno, C., & Muscella, A. (2019). Is mitochondrial DNA profiling predictive for athletic performance? *Mitochondrion*, 47, 125–138. https://doi.org/10.1016/j. mito.2019.06.004

Sugiura, N. (1978). Further analysis of the data by Akaike's information criterion and the finite corrections. *Communications in Statistics, Theory and Methods*, A7, 13–26. https://doi.org/10.1080/03610927808827599

Urawa, S., & Bugaev, A. V. (2021). Survival of Japanese chum salmon during early ocean life in 2011–2017. *North Pacific Anadromous Fish Commission Technical Report*, 17, 60–62.

Urawa, S., Beacham, T. D., Fukuwaka, M., & Kaeriyama, M. (2018). Ocean ecology of chum salmon. In R.

J. Beamish (Ed.), *The Ocean Ecology of Pacific Salmon and Trout* (pp. 161–317). Bethesda, Maryland:

American Fisheries Society. https://doi.org/10.47886/9781934874455.ch3

Urawa, S., Beacham, T., Sutherland, B., & Sato, S. (2022). Winter distribution of chum salmon in the Gulf

of Alaska: a review. *North Pacific Anadromous Fish Commission Technical Report*, 18, 83–87.

Walker, R. V., Myers, K. W., Davis, N. D., Aydin, K. Y., Friedland, K. D., Carlson, H. R., ... & Anma, G.

(2000). Diurnal variation in thermal environment experienced by salmonids in the North Pacific as

indicated by data storage tags. *Fisheries Oceanography*, 9, 171–186. https://doi.org/10.1046/j.1365-

2419.2000.00131.x

Watanabe, K., Mizumoto, H., Honda, K., & Sato, S. (2022). *2022 status of Japanese chum salmon stocks*

(Tentative translation). Fisheries Research and Education Agency. (in Japanese) Retrieved from

https://kokushi.fra.go.jp/R04/R04_61_CHU.pdf

Wikipedia (2023, October, 10). Countergradient variation. Retrieved from

https://en.wikipedia.org/wiki/Countergradient_variation

Wilkinson, L. (1999). Statistical methods in psychology journals: Guidelines and explanations. *American*

Psychologist, 54, 594. https://psycnet.apa.org/doi/10.1037/0003-066X.54.8.594

Winans, G. A., Aebersold, P. B., Urawa, S., & Varnavskaya, N. V. (1994). Determining continent of origin

of chum salmon (*Oncorhynchus keta*) using genetic stock identification techniques: status of

allozyme baseline in Asia. *Canadian Journal of Fisheries and Aquatic Sciences*, 51(S1), 95–113.

https://doi.org/10.1139/f94-298

Yagisawa, Y. (1970). Salmon hatchery enhancement program on Etorofu Island (Tentative translation), *Fish*

and eggs, Salmon Management Center Technical Report, 133, 35–79. (in Japanese)

http://salmon.fra.affrc.go.jp/kankobutu/tech_repo/fe04.htm

Yamaguchi, F., Nakamura, T., & Urabe, H. (2021). Temporal and spatial variations in body size of chum

salmon in Hokkaido. *North Pacific Anadromous Fish Commission. Technical Report*, 17, 23–24.

1064

TABLE 1 Explanatory variables included in the regression model and hypothesised mechanism.

Explanatory variables	Spatial scale	Age	Time scale	Unit	Mechanism
Size at release in year t (BW)	Japan coasts	Age 0	After release	0.1g	Larger juveniles survive better than smaller ones.
Size at release in year t (Condition factor)	Japan coasts	Age 0	After release		Juveniles with higher condition factors survive better than those with lower factors.
Catch of yellowtail in year t	Japan coasts	Age 0	After release	100 tons	Yellowtail feed on juvenile chum salmon during their extended northward migration.
Mean SST at release in year t	Japan coast	Age 0	After release	°C	SSTs at and after release affect survival of juvenile chum salmon.
Summer SST in the Sea of Okhotsk in year t	Off the Sea of Okhotsk	Age 0	Jul.	°C	SSTs affect survival of juvenile chum salmon after leaving the Sea of Okhotsk coast.
Winter SST in the North Pacific in year t+1 when Age 1 fish over winter	Western north Pacific	Age 1	Jan. and/or Feb.-Apr.	°C	During the first overwintering, SSTs affect the survival of young chum salmon.
Summer SST in the Bering Sea in year t+1 when Age 1 fish feed	Bering Sea	Age 1	Aug.	°C	During the feeding migration, SSTs affect the survival of young chum salmon.
Winter SST in the Gulf of Alaska in year t+2 when Age 2 fish over winter	Gulf of Alaska	Age 2	Jan. and/or Feb.-Apr.	°C	During the second overwintering, SSTs affect the survival of chum salmon.
Russian chum salmon catch in year t+3 (relative recruitment in year t)	Sea of Okhotsk and North Pacific	Age 0 - Age 4	3 years and months	million fish	Intraspecific competition in the Sea of Okhotsk and the North Pacific until return.
USA and Canadian chum salmon catch in year t+3	North Pacific	Age 1 - Age 4	2 years and months	million fish	Intraspecific competition in the North Pacific until return.
Russian pink salmon catch in year t+1	Sea of Okhotsk and North Pacific	Age 0 - Age 2	1 year and months	million fish	Interspecific competition in the Sea of Okhotsk and the North Pacific.
USA and Canadian pink salmon catch in year t+1	North Pacific	Age 1 - Age 2	1 year	million fish	Interspecific competition in the North Pacific.

1065 All data including the response variable (return rate from release in year t), are available in the Supplementary Tables and Supplementary Data.

TABLE 2 Summary of the best fit models with minimum AICc values out of 2^{14} (=16,384) models for each geographical area.

Area	Explanatory variable	Estimate	SE	t-value	Pr (>$|t|$)	Adjusted R^2	p-value
Sea of Okhotsk	(Intercept)	-4.937	0.672	-7.34	**0.000**	0.75	**0.000**
	Gulf of Alaska winter SST [†]	0.304	0.095	3.21	**0.007**		
	Russian chum salmon	0.030	0.008	3.57	**0.003**		
	Yellowtail	-0.202	0.033	-6.14	**0.000**		
Hokkaido and Honshu Pacific	(Intercept)	-12.267	2.220	-5.53	**0.000**	0.43	**0.003**
	Size at release	0.601	0.166	3.61	**0.003**		
Hokkaido Sea of Japan	(Intercept)	1.182	2.499	0.47	0.644	0.49	**0.008**
	Size at release	-0.419	0.186	-2.25	**0.043**		
	North Pacific winter SST [‡]	0.682	0.213	3.20	**0.007**		
	Sea of Okhotsk summer SST [§]	-0.360	0.110	-3.28	**0.006**		
Honshu Sea of Japan	(Intercept)	-2.257	1.375	-1.64	0.125	0.69	**0.000**
	Size at release	0.096	0.037	2.59	**0.023**		
	Gulf of Alaska winter SST [†]	0.223	0.081	2.76	**0.016**		
	SST at release	-0.529	0.105	-5.05	**0.000**		

1066   Bold indicates significance.   [†] Jan.,   [‡] Feb.-Apr. (mean),   [§] Jul.

JUL   **JAN**

1067

FIGURE 1 Long-term changes in the summer and winter SST isocline in the North Pacific. Plotted from
Plot: COBE Sea Surface Temperature, NOAA Physical Sciences Laboratory. Changes in April, May, June,
August, October and November SSTs are shown in Figures S11-13.

1071

1072 **FIGURE 2** Migration routes of Japanese chum salmon and critical SSTs. Map shows migration routes
1073 estimated from the long-term field surveys, redrawn from Urawa et al. (2018). Box plots show the change in
1074 mean summer and winter SSTs of feeding and wintering areas, calculated from the NOAA 1/4° daily OISST
1075 climate record. The gray boxes show the summer and winter SSTs included in the explanatory variables.

1076

1077 **FIGURE 3** Summer and winter mean SSTs in the Japanese chum salmon migration route in Figure 2. SSTs
1078 are calculated from the NOAA 1/4° daily OISST climate record.

1079

1080 **FIGURE 4** Number of chum salmon released and returned in Japan and SST at release. The dashed lines

1081 show the historical maximum and the minimum after the maximum. Mean SSTs for the release months are

1082 calculated from the 0.25°×0.25° daily mean SST datasets from the JMA (see Methods).

1083

1084   **FIGURE 5** Age-specific return rate of chum salmon by management area.

1085

1086 **FIGURE 6** Chum salmon return rate and size at release by management area. (a) Map shows salmon

1087 hatcheries in the seven management areas in different colors and waypoints (×) to off Monbetsu, Sea of

1088 Okhotsk (+). (b) Geographic distance from hatcheries to off Monbetsu vs. return rate (Ages 2-5). (c) Mean

1089 size at release weighted by number of juveniles released. (d) Neighbor-joining tree showing pairwise

1090 differences of return rates between management areas.

1091

1092   **FIGURE 7** Changes in commercial catches of 56 marine species in Japan over four decades (1980-2020).

1093   (a) A biplot of the principal component analysis (PCA) based on the catch statistics. (b) Two decadal shifts

1094   in the commercial catch of chum salmon and two piscivores; yellowtail and Japanese Spanish mackerel.

1095

FIGURE 8 Predicted (gray line) and observed return rates (black dots) from Age 2 to Age 5 for the four
geographic areas. The 95% prediction intervals are shown as black lines.

Explanatory variable	Effect size (95%CI)	Unit
Sea of Okhotsk		
a1. Gulf of Alaska Jan. SST	1.36 (1.13−1.63)	1°C
a2. Russian chum salmon	1.03 (1.01−1.05)	million fish
a3. Yellowtail	0.82 (0.77−0.87)	100 tons
Hokkaido and Honshu Pacific		
b1. Size at release	1.82 (1.32−2.53)	0.1 g
Hokkaido Sea of Japan		
c1. Size at release	0.66 (0.46−0.95)	0.1 g
c2. North Pacific Feb.−Apr. SST	1.98 (1.30−3.00)	1°C
c3. Sea of Okhotsk Jul. SST	0.70 (0.56−0.86)	1°C
Honshu Sea of Japan		
d1. Size at release	1.10 (1.02−1.18)	0.1 g
d2. Gulf of Alaska Jan. SST	1.25 (1.07−1.47)	1°C
d3. SST at release	0.59 (0.48−0.72)	1°C

Effect size (Rate of change per unit)

1098

FIGURE 9 Effect size of the significant explanatory variables of the best models. The effect size indicates the rate of change per unit, calculated as the exponential of the regression coefficient.

FIGURE 10 Geographic distributions of LDH-A1 and growth hormone 2 SNP (GHII3129) allele frequencies. LDH-A1 from 147 populations ($n = 18,892$) and SNPs from 114 populations ($n = 10,458$). The dots on the maps show sampling locations and allele frequencies with a continuous color gradient from blue to red for the lowest and highest allele frequencies.