Hatcheries to high seas: climate change connections to salmon marine survival

Shuichi Kitada[1*], Katherine W. Myers[2], and Hirohisa Kishino[3,4]

Short running title: Climate connections to salmon survival

[1] Tokyo University of Marine Science and Technology, Tokyo, Japan

[2] School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98195, USA.

[3] Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, Japan

[4] Research and Development Initiative, Chuo University, Tokyo, Japan

*correspondence: email: kitada@kaiyodai.ac.jp

Abstract

We investigated variations in the marine survival of Japanese hatchery chum salmon (*Oncorhynchus keta*) during 25 years of climate change (1998-2023). Japan is the world's largest producer of hatchery salmon, and is located near the global southern distribution limit of chum salmon. Our goal was to identify local and context-specific metrics related to salmon marine survival for use in the adaptive management of hatchery salmon resources. We hypothesized multiple metrics in three categories of stressors: hatchery carryovers, ocean conditions, and predators and competitors. The hatchery carryovers are stressors related to hatchery rearing that affect survival at a different life stage. We collected, processed, and collated large publicly available datasets of metrics into a comprehensive open-access database encompassing the life cycle of Japanese chum salmon, from eggs to adult spawners. Multivariate regression models showed associations between stressor metrics and adult salmon return rate (marine survival) varied by coastal management region, salmon life stage, and seasonal high seas distribution area. Generalized additive models identified a shift to younger age at maturity and egg size decline in all management regions. Our analysis suggested that maternal traits (egg size, fecundity) and the quality of fry hatchery salmon (body size at release) are particularly important for survival at later ocean stages. Our results support the hypothesis that whole life-cycle processes from anthropogenic selection in hatcheries to climate-driven changes in coastal predation, inter- and intra-specific high seas competition, and maternal effects during egg maturation at sea are responsible for the observed coastwide decline in survival of Japanese hatchery chum salmon over the past two decades. Our study highlights the need for an experimental approach to hatchery practices, including monitoring and analyses with updated information, leading to effective management decisions and policies for future sustainability and conservation of salmon resources.

KEYWORDS age at maturity, competition, egg size decline, fry size at release, Japanese hatchery chum salmon, predation

1 | INTRODUCTION

Climate change influences the survival and abundance of many economically important marine and anadromous fish species, including Pacific salmon (*Oncorhynchus* spp.) (Springer & van Vliet, 2014; Cheung & Frölicher, 2020; Cheung et al., 2021; Crozier et al., 2021; Crozier & Siegel, 2023). Pacific salmon populations often consist of hundreds of subpopulations with diverse life history traits and local adaptations to environmental variation (Groot & Margolis, 1991; Hilborn et al., 2003; Larson et al., 2014; Quinn, 2018). Pacific salmon species are distributed in coastal marine waters and throughout the high seas (international waters) of the North Pacific Ocean and Bering Sea, where Asian and North American populations intermingle during seasonal feeding migrations (Myers et al., 2007; Beamish, 2018; Quinn, 2018). Their wide geographic range, long ocean migrations, and complex life histories expose salmon populations to multiple climate-driven stressors at both local and North Pacific-wide scales (Cline et al., 2019; Crozier et al., 2021). Global warming of 1.5°C above pre-industrial levels is projected to shift the ranges of many marine species to higher latitudes and increase damage to marine ecosystems, such as rising temperatures, ocean acidification, heatwaves, decreases in sea ice, habitat loss, disruptions in food webs, and declining fish populations (Dahlke et al., 2018; Frölicher et al., 2018; Masson-Delmotte, 2022).

The effects of climate change, ocean acidification, and large-scale hatchery releases on the abundance and productivity of Pacific salmon populations have received increasing attention in recent years (Lewis et al., 2015; Jeffrey et al., 2017; Cunningham et al., 2018; Cline et al., 2019; Losee et al., 2019; Connors et al., 2020, 2024; Oke et al., 2020; Ohlberger et al., 2022, 2023; Malick et al., 2023). Declines in body size, primarily due to a shift in age structure towards younger ages at maturity, were found in chum, Chinook (*O. tshawytscha*), coho (*O. kisutch*), and sockeye (*O. nerka*) salmon from large-scale analyses across Alaska and were linked to climate change and increased inter- and intra-specific competition due to an overabundance of salmon in the high seas (Lewis et al., 2015; Oke et al., 2020; Ohlberger et al., 2023). Pink salmon (*O. gorbuscha*) abundance increased since the mid-1970s to record highs in the 2000s, supported by climate change effects on the ocean environment and production processes, and may have affected the abundance of salmon, other species, and their ecosystems (Springer & van Vliet, 2014; Connors et al., 2020; Ruggerone et al., 2023).

Ocean mortality of salmon is thought to be regulated initially by predation on fry soon after ocean entry, and is related to body size and density of salmon and abundance and type of predators. A later mortality phase is a consequence of growth rate in the early-ocean period and the ability of juvenile fish to survive the first ocean winter (Ricker, 1976; Beamish & Mahnken, 2001). Fisheries-oceanographic surveys in the western North Pacific, however, indicated that

survival of salmon is related to ocean conditions (not growth rate) during the first fall and winter in the open ocean (Naydenko & Temnykh, 2016; Naydenko et al., 2016; Bugaev & Gerlits, 2023). Natural mortality of immature and maturing high seas salmon is generally thought to be low (Ricker, 1976). Hypotheses about the mechanisms underlying high-seas salmon distribution and abundance in winter are varied (reviewed by Myers et al., 2016). Historical survey data indicated seasonal high-seas salmon distribution was related to sea surface temperature (SST) and salinity (Manzer et al., 1965; Welch et al., 1998; Azumaya et al., 2007; Abdul-Aziz et al., 2011; Langan et al., 2024). Salmon survey data in the Sea of Okhotsk and western North Pacific indicated the seasonal distribution of high-seas salmon reflects the climate-driven landscape (abiotic and biotic) of pelagic ecosystems, and interannual variation in high-seas salmon abundance is determined at earlier life stages (Naydenko & Temnykh, 2016; Naydenko et al., 2016; Bugaev & Gerlits, 2023). Rapid seasonal climate-change events such as marine heatwaves are driving the natural selection and adaptation of organisms in altered ecosystems (Bradshaw & Holzapfel, 2006).

Pacific salmon have a long history of artificial propagation beginning in the mid-1800s, and their management comprises the largest set of hatchery release programs (Naish et al., 2007), including chum salmon hatcheries in Japan that constitute the world's largest salmon hatchery release program (Amoroso et al., 2017; Kitada, 2018). Hatchery releases of fry chum in Japan increased rapidly in the 1970s, but gradually declined from a peak of 2 billion per year in 1983 to 1.3 billion per year in 2021 because of poor returns of adult spawners. Adult returns of Japanese hatchery chum increased rapidly since the 1970s, but declined sharply after the late 1990s from the historical maximum of 88 million in 1996 to the minimum of 19 million in 2021. Previous studies have reported higher spring SSTs in the coastal zone contributed to better survival of juvenile chum through improved feeding, growth, and swimming ability (Saito & Nagasawa, 2009; Kuroda et al., 2020; Honda et al., 2021). Conversely, shorter inshore stays associated with warming SSTs could lead to lower juvenile survival and insufficient growth (Kaeriyama, 2022; Honda et al., 2025). Although it is generally accepted that a warming climate is affecting marine survival of salmon, the mechanisms underlying recent declines in Japanese hatchery chum salmon are largely unknown. The need for adaptive management is advocated for the long-term sustainability of chum salmon (Kaeriyama, 2008, 2022; Kaeriyama et al., 2012).

The development of adaptive management plans to address climate-change impacts on fish, fisheries, aquaculture, and associated habitats and ecosystems requires knowledge of "local and context-specific indicators associated with climate stressors" (FAO, 2022). Adaptive management is the design of management actions as experiments with feedback through

monitoring, enabling an iterative process of learning and adapting to uncertainties and risks (Walters & Hilborn, 1978; Hilborn & Sibert, 1988; Bahri et al., 2021; Bryndum-Buchholz et al., 2021). A comprehensive review of literature addressed strengths and weaknesses of climate-change research by life stage and showed the complexity of whole life cycle approaches to adaptive management strategies for anadromous salmonids (Crozier & Siegel, 2023). Here, we investigated associations between adult return rates (an index of marine survival) of Japanese hatchery chum salmon by life stage and a suite of hypothesized climate-change stressors. In our study, we expected survival of all life stages of chum salmon to be associated with climate change impacts on coastal and open ocean ecosystems. Our primary objective was to identify potential stressors along known coastal and high seas migration routes of Japanese chum salmon for use in regional adaptive management of salmon resources.

2 | MATERIALS AND METHODS

2.1 Study area

The ecological setting of our study encompasses the known geographic distribution and seasonal migration patterns of Japanese chum salmon in the North Pacific Ocean and adjacent seas (Figure 1; reviewed by Fisheries Agency of Japan, 1955; Sano, 1966, 1967; Shepard et al., 1968; Neave et al., 1976; Salo, 1991; Urawa et al., 2018). Chum salmon fry are released from more than 260 coastal hatcheries (Kitada, 2020) in northern Honshu and Hokkaido in winter and spring and migrate rapidly through river estuaries (Figure 1a; Kobayashi, 1980; Kaeriyama, 1989; Saito & Nagasawa, 2009). After ocean entry, fry migrate northward in shallow coastal waters before moving offshore to summer feeding grounds in the southern Sea of Okhotsk (Figure 1b; Fukuwaka & Suzuki, 1998; Urawa et al., 2006; Kasugai et al., 2016; Shubin & Akinicheva, 2016; Honda et al., 2025). Japanese chum salmon spend their first ocean winter (age 1) in the western North Pacific Ocean and in spring move northeastward to summer feeding grounds in the Bering Sea (Figure 1b; Mayama & Ishida, 2003; Urawa et al., 2018). At subsequent life stages (ages 2-4; immature and maturing), they make seasonal migrations between the Gulf of Alaska in winter and the Bering Sea in summer (Figure 1b; Beacham et al., 2009; Urawa et al., 2009, 2016). In spring, the northward migration of maturing chum salmon precedes that of immature fish (Neave et al., 1976; Nagasawa & Azumaya, 2009). In late summer and fall, maturing fish return to Japan from the Bering Sea through the western North Pacific (Figure 1b; Yonemori, 1975; Urawa et al., 2018).

2.2 Data sources, sample periods, sample sizes

We assembled available time-series datasets during 25 years of climate change (1998-2023) from public open-data websites. We assembled longer data time series to aid in interpreting results.

We obtained annual (1970-2023) numbers of adult salmon catches (coastal fishery and hatchery

broodstock) and hatchery chum salmon fry releases from the Hokkaido Salmon Enhancement

Association (HSEA, 2024), Fisheries Research and Education Agency (FRA, 2024), and the

Aomori Prefectural Government (APG, 2024). FRA (2023) biological and reproductive trait

data included time series of fry size at release and release month. We selected hatchery release

events with fry body weight (BW) and fork length (FL) data (1998-2020, Hokkaido: 184 rivers,

37,886 release events, n = 21,332,093 fish; Honshu: 141 rivers, 19,605 release events, n =

9,588,413 fish). The data also included adult age composition at return (1997-2020, Hokkaido:

52 rivers; Honshu: 102 rivers; 1,902 surveys, n = 1,210,717 fish). Adult salmon trait data

included egg weight, fecundity, FL, age at return, and gonad weight (1997-2020, Hokkaido: 10

rivers; Honshu: 14 rivers; n = 37,533 fish), and adult age at return (counts of annular marks on

fish scales), including six age groups: age 2 (n = 25), age 3 (n = 2,892), age 4 (n = 20,643), age

5 (n = 13,118), age 6 (n = 846), and age 7 (n = 9). We used trait data such as FL, egg weight,

and fecundity of age–4 and age–5 fish for our analysis.

We obtained annual commercial catch data for all marine species in Japan (1980-2020) from

the Japan Ministry of Agriculture, Forestry and Fisheries (MAFF, 2023). Catch statistics (1998-

2022) for Canada, Russia, and USA pink and chum salmon were available from the North

Pacific Anadromous Fish Commission (NPAFC, 2023). For SST analyses, we downloaded

Japan coastal 0.25°×0.25° daily mean SST data (1982 to 2020) from Hokkaido (18 areas) and

Honshu (12 areas) from the Japan Meteorological Agency (JMA, 2022) and global 0.25°×0.25°

monthly mean SST data (1982 to 2022) from the NOAA Optimum Interpolation (OI) SST V2

high-resolution dataset (NOAA, 2023a).

2.3 Catch, release, and reproductive traits

We summarized Japanese chum salmon catch, release, and reproductive traits data by seven

management regions in northern Honshu and Hokkaido, Japan (Figure 1a). To estimate the

distribution of release times in each region, we counted the number of release events by month.

We also calculated the mean values of size at release, biological traits, and age composition by

management region. Since the number of individuals released (n_k) varied among release events

(k=1,…,K), we calculated the weighted mean size at release for BW (g) (1998-2020) as $\overline{BW} =$

$(\sum_{k=1}^{K} BW_k n_k)/\sum_{k=1}^{K} n_k$. Here, the suffix for a region was omitted for simplicity. We

calculated the mean values of the traits and the relative fecundity (eggs/$FL(\text{cm})^3 \times 10^5$) of the

age 4 and age 5 fish from 1997 to 2020. The weighted mean proportion of age a fish was

calculated as $\hat{p}_a = (\sum_{h=1}^{H} p_{ah} n_h)/\sum_{h=1}^{H} n_h$, where h refers to each survey, p_{ah} is the

proportion of age a fish in survey h, n_h is the sample size (individuals), and a refers to age at

return ($a = 2,\ldots,8$).

2.4 Adult return rates (marine survival index)

We assumed all adult chum salmon returning to Japan were hatchery reared, and adult return

rates are survival rates from release to return. In this study, we define the adult return rate as an

index of marine survival of chum salmon (James et al. 2023, see in discussion section 4.1).

Hatchery programs have been managed in five management regions in Hokkaido and by

prefecture in Honshu, and previous studies have evaluated return rates in the management

regions (Saito & Nagasawa, 2009; Miyakoshi et al., 2013; Kitada, 2014). The management

regions coincide with the genetic structure of the population (Beacham et al., 2008). We

estimated return rates for the current hatchery management regions (Figure 1a). We estimated

the total number of fish returning from releases in year t (Y_t, coastal commercial catch plus river

catch for hatchery broodstock) by

203
$$\hat{Y}_t = \sum_{a=A_{min}}^{A_{max}} y_{t+a-1}\, \hat{p}_{a,t+a-1} = \sum_{a=A_{min}}^{A_{max}} \hat{Y}_{a,t}$$

where y_{t+a-1} is the observed number of fish returned in year $t + a - 1$ from fry released in

year t, and $\hat{p}_a$ is the estimated proportion of age-a fish among the fish returned in year $t + a -$

1. We estimated the return rate of fry released in year t by $\hat{R}_t = \hat{Y}_t/X_t$, where X_t is the number

of hatchery-reared fry released in year t. We estimated the age-specific return rate as

$\hat{R}_{a,t} = \hat{Y}_{a,t}/X_t$. We summed the number of fish returned from ages 2–5 ($A_{min} = 2$, $A_{max} = 5$).

Most returns were ages 2–5 fish, and few returns were age 6. There were no age–7 and age–8

fish in returns to any region except Hokkaido Okhotsk, where returns of these age groups were

very low. We estimated the age composition of returning adult chum salmon from available

(1997-2020) data (Figure S1). We estimated the age composition in 2021-2023 as a moving

average of the previous three years. Based on age composition, we calculated the number of

returning fish (ages 2–5) for each release year. Then, we estimated return rates and age-specific

return rates from releases in 1998-2019. We measured distances between return rates in the

management regions by $1 - r_{uv}$ and illustrated distances by a neighbor-joining (NJ) tree

(Saitou & Nei, 1987) using the "MEGA11" package (Tamura et al., 2021), where r_{uv} is the

correlation coefficient between pairs of return rates in regions u and v.

2.5 Age at maturity and body size

We calculated the percentage of ages 2-4 chum salmon in the total number of adult returns (ages

2–5), used as an index of younger age at maturity, because the preliminary analysis showed that

ages 2-4 chum salmon increased after around 2010 in all management regions (Figure S1).

Excluding older fish (age 6 or older) should have a negligible effect on estimates of mean ages

at maturity (Figure S1). To assess and visualize continuous changes in the percentage of ages 2–4 chum salmon, return rates, body sizes, and other biological traits over the year, we fit general additive models (GAMs) to the data using the 'gam' function in the "mgcv" package (Wood, 2017) in R (R Core Team, 2024). Year was included as a nonlinear smoothed term (explanatory variable) because return rates and other biological traits varied across years. The trend of the traits over the years is captured by the non-linear and linear year effects (Wood, 2004, 2011).

2.6 Long-term changes in chum salmon thermal habitat

To understand long-term changes in SST in the chum salmon habitats covered by our regression model (see section 2.9), we mapped monthly mean 1°×1° SST isoclines in the North Pacific Ocean for 1990, 2000, 2010, and 2022, using the Plot COBE Sea Surface Temperature system (NOAA, 2023b), based on the COBE2 SST dataset from JMA. The 1°×1° monthly mean SST isoclines around Japan for 2000, 2010, and 2022 were generated similarly. We mapped the available temperature range of chum salmon (4-13℃), based on previous predictions: optimal growth and feeding (8–12℃), available habitat for feeding (5–13℃), and overwintering temperature range (4–6℃) (Kaeriyama et al., 2012; Urawa et al., 2018).

2.7 Japanese chum salmon migration route and SST

We calculated monthly mean SSTs for 30 coastal areas in Japan (JMA, 2022). From these, we calculated monthly mean SSTs for the seven management regions (Figure 1a), and for release periods of chum salmon fry (Figure S2): Hokkaido Okhotsk (Apr-Jun), Hokkaido Nemuro (Mar-Jun), Hokkaido Pacific East (Mar-Jun), Hokkaido Pacific West (Mar-May), Hokkaido Sea of Japan (Mar-May), Honshu Pacific (Feb-May), and Honshu Sea of Japan (Feb-Apr).

Chum salmon fry born in rivers along the Pacific coast of Japan migrate to the Sea of Okhotsk (Honda et al., 2017, 2025). Juvenile chum salmon from all regions of Japan are found in the Sea of Okhotsk, and abundance of juveniles is significantly correlated (positive) to adult returns to Japan (Urawa & Bugaev, 2021). Juvenile chum salmon were abundant in coastal areas of the Sea of Okhotsk from May to June when SST was between 8 and 13°C, and disappeared from coastal waters after late June when SST exceeded 13°C (Nagata et al., 2007). Based on these results, we assumed that chum fry released in all areas would migrate to the Sea of Okhotsk and disperse offshore in July (Figure 1a).

We identified the locations of four main offshore and high seas areas of Japanese chum salmon distribution (Figure 1b) based on information from previous studies (Myers et al., 2007; 2016; Nagata et al., 2007; Urawa et al., 2018, 2022). Using "sst.mnmean.nc" data (NOAA, 2023a),

we calculated the monthly mean SSTs (1982-2022) for salmon distribution areas using the 'fldmean' function in the "cdo-2.1.1" package (Max Planck Institute, 2023). We mapped areas and migration routes using the "sf" package in R. We used monthly mean SSTs (1998-2022) in our regression model. To assess and visualize continuous changes in SSTs over the year, we fit GAMs to these variables by migration area using the 'gam' function in the "mgcv" package in R.

2.8 Potential predators of chum salmon fry

To characterize temporal changes in catches of marine fish predators, we performed principal components analysis (PCA) on the annual commercial catch data for the marine species in 1980-2020 (MAFF, 2023) using the 'prcomp' function in R. Because catches vary by geographic area and season, we obtained catch data of potential marine fish predator species by reporting area in Hokkaido and by prefecture in Honshu (MAFF, 2023), which should reflect their fishing seasons and migration patterns. Changes in chum salmon catch and catches of potential marine fish predators, identified by increasing catch in the chum salmon coastal distribution range, were mapped.

2.9 Regression analysis of adult return rates

Our regression model describes the relationship between return rates of chum salmon fry released in year t across all ages ($\hat{R}_t$) and explanatory variables such as SSTs, size at release, and catch of potential predators and competitors. We transformed the return rate to the log scale for model fitting in a multiple linear regression:

$$\log(\hat{R}_t) = \log\beta_0 + \sum_{i=1}^{s} \beta_i z_i + \varepsilon_t$$

where β_0 is the intercept, and the explanatory variables are denoted by $\boldsymbol{z}$. The regression coefficients were denoted by $\boldsymbol{\beta}$ and $\varepsilon_t \sim N(0, \sigma^2)$ is the error that we assumed to be independently and normally distributed. We did not include interactions between explanatory variables. We selected the explanatory variables by AICc (Sugiura, 1978), which is the small sample correction of the Akaike Information Criterion (AIC, Akaike, 1973).

We estimated regression coefficients based on standardized explanatory variables, which allow for consistent comparison of different explanatory variables across management regions as done in salmon studies (e.g., Oke et al., 2020; Ohlberger et al., 2022). The use of an unstandardized measure (regression coefficient) is preferred to a standardized measure if the units of measurement (e.g., fry size at release, SST, and abundance of competitors) are used to describe effect sizes at a "practical" or "real world" level (Wilkinson et al., 1999). Therefore,

we also estimated the regression coefficients using unstandardized explanatory variables, as done in Malick et al. (2023). In the latter, we define effect size as the effect of the unit change of an explanatory variable on the change in return rate. The effect size and 95% confidence intervals were calculated by $\exp\left(\hat{\beta}_i \pm 1.96 \times \text{SE}(\hat{\beta}_i)\right)$ for coefficient i. We predicted return rates by region using the 'predict' function in R based on regression coefficients using the unstandardized measure and compared them to the observed return rates.

When many explanatory variables are included, the model selection procedure may select an inappropriate model as the best model due to complex multicollinearity. To avoid this problem, we created three subset models of stressors (see below), performed variable selection within each subset model, and compared the best models for combinations of the subset models and the full model, which included all subset models. In each model, we used the variance inflation factor (VIF) to diagnose multicollinearity among explanatory variables. We determined variables that could be included in a model by sequentially removing each explanatory variable with the highest VIF value until all remaining VIFs were less than the threshold of 3.0 (Zuur et al., 2010), using the 'VIF' function in the "car" package in R.

The dependent variable is adult return rates of fry released in 1998-2019. In the terminology of adaptive management, "a performance indicator is a variable that quantifies how well a fishery is managed, in relation to specified objectives, and typically has an audit function" (Rochet & Trenkel, 2009). Our dependent variable (an index of marine survival) is a "performance indicator" of marine survival in the context of adaptive management. We created three subset models for our regression analysis of adult return rates and explanatory variables. We used 17 hypothesized metrics (covariates) for explanatory variables ($s = 17$) and: (1) hatchery carryover (three metrics), (2) ocean condition (five metrics), and (3) predators and competitors (nine metrics) (Table 1). The hypothesized hatchery carryover metrics included relative fecundity and egg size of females in year t-1, where t is the year of fry release, and fry size at release. Ocean-condition metrics included SST at fry release and in Japanese coastal management regions; summer SST in the southern Sea of Okhotsk and central Bering Sea, and winter SST in the western North Pacific and central Gulf of Alaska (see Figure 1). Predator and competitor metrics included predation in coastal Japan regions and competition in high seas areas (Table 1). We identified yellowtail (*Seriola quinqueradiata*), also known as Japanese amberjack, as a potential predator of chum salmon fry (see Section 3.5). Age–4 chum salmon were the dominant age group in adult returns to Japan (Figure S1). We used the annual commercial salmon catch (total number of fish) of chum salmon in Russia, the United States, and Canada in year t+3 as indices of intraspecific competition with Japanese chum salmon released in year t, as they

return in year $t+3$ as age–4 fish. They were assumed to spend three years in the high seas and to compete when sharing the same environment. Pink salmon have a fixed life span of two years (Quinn, 2018). We used annual commercial catches in Russia and North America (US and Canada, combined totals) in years $t+1$, $t+2$, and $t+3$ as indices of interspecific competition with Japanese chum salmon released in year t. The ocean distribution of chum and pink salmon overlaps throughout the North Pacific (Langan et al., 2024). We assumed cohorts of pink salmon compete with chum during the first ($t+1$), second ($t+2$), and third ($t+3$) ocean years.

We performed regression analysis for each subset model and combinations of subset models with explanatory variables less than 3.0 VIF. This procedure reduces the number of parameters to be estimated and increases the information available for inference, resulting in robust inference. In each analysis, we selected the most parsimonious model (hereafter, the best model) with the minimum AICc using the "MuMIn" package in R. We compared all models by their minimum AICc values. To summarize the result, we presented the regression coefficients of the best model for the selected subset models. In three of the four regions, the minimum of the minimum AICc values was achieved by multiple hierarchical subset models. In such cases, we chose the most parsimonious subset models. The estimated regression coefficients were compared with the results obtained by model averaging over all models with ΔAICc values less than 2 ΔAIC < 2, which suggests substantial support for the model (Burnham & Anderson, 2002). A practical model averaging provides the weighted averages of the regression coefficients over the models. The weights represent the contributions of the models and are proportional to $\exp(-1/2 \times \text{AICc})$. All data and the R script used for the regression analysis are available (see data availability statement).

3 | RESULTS

3.1 Adult return rate (marine survival index)

For Japan as a whole, the number of hatchery-released fry increased until the early 1980s, reaching 2 billion, and then decreased to about 1.3 billion at present (Figure 2). In the Honshu Pacific, there was a steady decline in releases since the mid-1990s and a rapid decline since 2021. Decreasing numbers of released fry were also observed in Hokkaido Pacific East and Honshu Sea of Japan, while stable numbers were observed in Hokkaido Nemuro, Hokkaido Pacific West, and Hokkaido Sea of Japan. Only in Hokkaido Okhotsk did the number of fry released increase slightly but continuously. Releases were largest in Honshu Pacific, while other regions had similar release levels.

The total number of adult chum salmon returns throughout Japan increased rapidly until 1996 (88 million fish), and then declined to the current level of 20 million fish (Figure 2). The highest

regional returns were in the Honshu Pacific in 1996 (31 million), and then declined continuously to the historical regional minimum (~10 thousand). Among other regions, adult returns were highest in northern Hokkaido regions (Okhotsk and Nemuro) and lowest in Sea of Japan regions (Hokkaido and Honshu). Since approximately 2008, adult returns decreased in all management regions except Hokkaido Okhotsk and Hokkaido Sea of Japan, and the decrease was most severe in Hokkaido and Honshu Pacific regions.

The mean adult return rate ± standard deviation (including ages 2–5) for 1998-2019 releases was highest in Hokkaido Okhotsk (6.9 ± 2.4%), followed by Hokkaido Nemuro (4.5 ± 3.0%), Hokkaido Pacific East (2.3 ± 1.7%) and Hokkaido Pacific West (2.3 ± 1.2%), Hokkaido Sea of Japan (1.4 ± 0.7%), Honshu Pacific (1.2 ± 0.7%), and Honshu Sea of Japan (0.6 ± 0.2%) (Figure S3a). The GAMs showed that return rates decreased over time in all management regions, except in Hokkaido Okhotsk and Hokkaido Sea of Japan, where return rates increased recently (Figure S4). The nonlinear and/or linear year effect was significant in all management regions except in Honshu Sea of Japan (Table S1).

The NJ tree analysis showed that Hokkaido Okhotsk, Honshu Sea of Japan, and Hokkaido Sea of Japan adult return rates were distinct, and the Hokkaido Nemuro, Hokkaido Pacific East and West, and Honshu Pacific return rates were similar (Figure 4). As a result, we used four coastal regions in Japan in our regression analysis: (1) Hokkaido Okhotsk, (2) Hokkaido and Honshu Pacific (combined returns to Hokkaido Nemuro, Hokkaido Pacific East and West, and Honshu Pacific), (3) Hokkaido Sea of Japan, and (4) Honshu Sea of Japan.

3.2 Age at maturity and body size

The percentage of age 2–4 adults in the total number of adult returns from 1998-2019 fry releases was very high in all management regions: Honshu Sea of Japan (81.8 ± 7.1%), Hokkaido Sea of Japan (73.4 ± 7.6%), Honshu Pacific (71. 3 ± 9.7%), Hokkaido Pacific West (70.7 ± 10.9%), Hokkaido Nemuro (66.7 ± 7.1%), Hokkaido Okhotsk (64.4 ± 9.4%), and Hokkaido Pacific East (59.7 ± 8.0%) (Figure S3b). The GAMs confirmed that after approximately 2008, the percentage of returning age 2-4 adults increased over time in all management regions (Figure 3). This trend was strongest after around 2008, and the rates of change differed among regions. The trend was significant in all regions, except Hokkaido Nemuro and Honshu Sea of Japan (Table S1).

The GAMs confirmed that body size (FL) of age–4 and age–5 hatchery chum salmon generally remained at similar levels after 2008 in all regions, though the variation was particularly large in age–5 fish in the Hokkaido Sea of Japan region (Figures S5, S6).

3.3 Long-term changes in chum salmon thermal habitat

Decadal-scale maps of SST isoclines showed that until 2000, the summer and winter thermal habitat of chum salmon was stable (Figure 5). After 2010, optimal summer habitat for chum salmon growth and feeding (especially, 8–11°C isotherms) shifted north and contracted longitudinally, and winter habitat remained relatively stable. Evaluation of intra-seasonal changes in spring and summer thermal habitat revealed a progressive northward shift in optimal habitat, from a slight shift in early spring (March and April) to a larger shift in early summer (May and June), northward longitudinal contraction in late summer (August and September), and a slight northward shift in fall (October and November) (Figures S7–S10). In coastal Japan, optimal thermal habitat for chum salmon fry migrating to the Sea of Okhotsk shifts northward in spring and early summer (April-June), especially in the Sea of Japan (Figure S11).

3.4 Japanese chum salmon migration route and SST

In Hokkaido Okhotsk and Hokkaido Sea of Japan, the mean SST at hatchery fry release was relatively stable from 1998 to 2019 (Figure S12, Table S1). In Hokkaido Nemuro, the mean SST increased. Hokkaido and Honshu Pacific showed a similar pattern; the mean SST decreased until around 2010 and then slightly increased. In the Honshu Sea of Japan, the mean SST remained stable until around 2010 and then increased.

The summer and winter SSTs used in our regression analysis (1998-2022) as common explanatory variables for all offshore (southern Sea of Okhotsk) and high seas areas (Figure 1b) were southern Sea of Okhotsk July SST, western North Pacific January-April SST, central Bering Sea August-September SST, and central Gulf of Alaska January-April SST (Figure S13a). A GAM (Figure S13b) showed that July SST in the southern Sea of Okhotsk increased after the early 2000s, and a strong increase was observed after around 2015. In the western North Pacific, winter SST in January-April increased after around 2015. In the central Bering Sea, August-September SST continued to increase since the late 1990s but decreased after around 2015. In the central Gulf of Alaska, a strong increasing trend in January-April SST was observed after around 2010, but SST decreased after around 2015. Similar trends were observed in the western and eastern North Pacific.

Summer SST (August–September) in the Bering Sea (range, 8.3–11.4°C) was within the optimal growth and feeding range, while winter SST (January–April) increased after 2010 in the western North Pacific (range, 4.6–6.1°C) and the Gulf of Alaska (range, 4.6–7.2°C) (Figure S13b). These results are consistent with the increasing trend in younger age at maturity of Japanese chum salmon after 2008 (Figure 3).

3.5 Potential predators of chum salmon fry

We summarized commercial catches of all marine species landed in Japan with no missing data from 1980 to 2020. We excluded marine mammals because catches are subject to strict limits, and catches were not recorded by species but by total marine mammals. This resulted in catch data for 56 of 70 species, including 39 fish, six crustaceans, five shellfish, two squids, an octopus, a sea urchin, and two algae. The PCA biplot identified differences in commercial fishery catch levels as the primary component (PC1, 51% variance), and the second component (PC2, 17% variance) corresponded to catch trends before and after 1998 (Figure 6a). The PCA confirmed that changes in commercial catches of 56 marine species in Japan after 1998 were largely characterized by three species that had positive PC1 values and increased in recent decades: yellowtail and Japanese Spanish mackerel (*Scomberomorus niphonius*) and Japanese scallop (*Mizuhopecten yessoensis*). We visualized trends in catches of 56 species (1980-2020) and confirmed that catches increased only in the three species identified by the PCA analysis (unpublished figure, S. Kitada). Among these, yellowtail and Japanese Spanish mackerel are strong piscivores. There was a northward shift of yellowtail landings into the range of Japanese chum salmon (Figure 6b). In contrast, increased landings of Japanese Spanish mackerel occurred only in the eastern area of Japan, with slight increases in the Honshu North Pacific area. Based on these results, we identified yellowtail as a possible new predator of chum salmon fry.

3.6 Regression analysis of adult return rates

Readers are cautioned that our statistical modeling results are correlational, not causal. All models used explanatory variables with a VIF less than the threshold of 3.0. Model selection results for all subset models are shown in Table S2. The full model results and the best subset model results were the same for all management regions except for the Hokkaido Sea of Japan (Table 2; Tables S2 and S3). The results of the model averaging (Table S4) were qualitatively the same as the best models (Table 2) for all regions. The explanatory variables selected in the best models were region-specific or common to some areas, or both. For ocean conditions, SST at release (year t) and summer SST (July) in the Sea of Okhotsk (year t) and the Bering Sea (August-September) (year $t+1$) were negatively associated with return rates. In contrast, winter SST (January-April) in the western North Pacific (year $t+1$) and the Gulf of Alaska (year $t+2$) were positively associated with return rates. For predators and competitors, we found negative associations between Japanese chum salmon return rates and yellowtail catch in the year of fry release. Russian chum salmon catch in year $t+3$, and Russian pink salmon (adult, age 1) catch in years $t+2$ and $t+3$ were negative. No associations between Japanese chum salmon return rates and US and Canadian chum and pink salmon catches were found. For hatchery carryovers, fry

size at release was positively associated with salmon return rates, while relative fecundity was negatively associated with return rate.

The GAMs indicated size at release has continued to decrease in all regions after around 2010, except Hokkaido Pacific West, and the decline in size at release was substantial in the Honshu Pacific region (Figure S14, Table S1). The relative fecundity of age–4 salmon had a smaller AICc (30.0) than age–5 salmon (38.2). We used the relative fecundity of age–4 salmon to identify the best model. Relative fecundity of age–4 salmon was negatively correlated with egg size in all management regions except in the Hokkaido Sea of Japan region (unpublished figure, S. Kitada). We found that relative fecundity of parent females was negatively associated with return rates in the Honshu Pacific region. GAMs confirmed that egg size for 4-year-old fish has decreased over the last two decades in all regions, except Hokkaido Okhotsk (Figure 7, Table S1). The trend was significant in all regions, except Honshu Sea of Japan. Similar decreasing trends in egg size for 5-year-old fish were also observed (Figure S15, Table S1).

The standardized effects of explanatory variables on adult returns allow comparison of the magnitude of the effects across the four regions (Figure 8). The largest positive effect was winter SST (January-April) in the Gulf of Alaska (year $t+2$), followed by size at release and winter SST (January-April) in the western North Pacific (year $t+1$). On the other hand, yellowtail catch had the largest negative effect, followed by SST at release, summer SST in the Bering Sea (August-September) (year $t+1$) and in the Sea of Okhotsk (July) (year t), and parental relative fecundity (year $t-1$). The negative effect of intraspecific and interspecific competition was relatively small and similar in all areas.

The practical effect size in a regression model describes how much of a change in the dependent variable is expected for a given change in the independent variable. The effect size for each regression coefficient, measured by $\exp(\hat{\beta}_i)$, represents the rate of change per practical unit in each coastal area (Figure S16), which showed a similar pattern to the standardized effects of the explanatory variables (Figure 8). The area-specific practical effect sizes showed that a 1℃ increase in winter SST in the Gulf of Alaska increased the return rate by 50% ($\exp(0.405) = 1.5$) in Hokkaido Sea of Japan, and a 1℃ increase in winter SST in the western North Pacific increased the return rate by 30% in Honshu Sea of Japan. In contrast, summer SST in the southern Sea of Okhotsk decreased return rate by 27% in Hokkaido Sea of Japan, and by 13% in Honshu Sea of Japan. SST at release reduced the return rate by 38% in Honshu Sea of Japan. An increase of 100 eggs (relative fecundity) of parent females (age 4) reduced the return rate by 61%, while fry size at release had a positive effect; a 0.1 g increase in hatchery fry BW increased the return rate by 63% in Hokkaido and Honshu Pacific. The average return rate over

the last five years was 0.74% in this region, and a 0.1g increase in size at release would be

expected to increase the return rate to 1.2% (= 0.74 × (1+0.63)) from current levels. An increase

in biomass of yellowtail landings of 100 tons predicted a 19% reduction in the adult return rate

of chum salmon in Hokkaido Okhotsk. An increase in Russian pink salmon abundance (100

million fish) in year t+2 reduced return rates by 12% in Hokkaido Okhotsk, 27% in Hokkaido

and Honshu Pacific, and 13% in Honshu Sea of Japan areas. An increase in abundance of

Russian pink salmon (100 million fish) in year t+3 reduced return rates by 20% in Hokkaido

and Honshu Pacific. An increase in abundance of Russian chum salmon (10 million fish)

reduced return rates by 14% in Hokkaido Sea of Japan.

The correlation between predicted and observed adult return rates was high in all four coastal

regions: Hokkaido Okhotsk ($r = 0.79, t = 5.75, p = 0.0000$), Hokkaido and Honshu Pacific

($r = 0.91, t = 9.98, p = 0.0000$), Hokkaido Sea of Japan ($r = 0.79, t = 5.78, p = 0.0003$),

and Honshu Sea of Japan ($r = 0.81, t = 6.17, p = 0.0000$). Our best models with few (2–4)

explanatory variables provided good fits to the four coastal regions and the annual variations in

chum return rates, although the 95% prediction intervals are large (Figure 9).

4 | DISCUSSION

We investigated relationships between marine survival of Japanese hatchery chum salmon and

multiple hypothesized climate-change stressors. While previous studies have focused largely

on climate change impacts on survival during the early marine (coastal, juvenile) life history

stage, we hypothesized that marine survival is related to climate change effects during all life

stages (from eggs to adults). We identified potential climate-driven stressors along known

coastal and high seas migration routes of Japanese chum in the North Pacific Ocean and

adjacent seas. Associations between stressors and marine survival varied by coastal region,

salmon life stage, and seasonal high seas distribution area. Fry size at release was positively

associated with marine survival of Japanese chum salmon, while relative fecundity of parent

fish was negatively associated with marine survival. We identified yellowtail as a new potential

marine fish predator of chum salmon fry, whose distribution was shifting northward. The SST

at release and summer SST in the Sea of Okhotsk and the Bering Sea were negatively associated

with marine survival, while winter SST in the western North Pacific and the Gulf of Alaska was

positively associated with marine survival. Negative associations between Russian chum and

Russian pink salmon catches and Japanese chum salmon marine survival were found, while no

associations were identified between US and Canadian chum and pink salmon catches. We

found a trend toward younger age at maturity in all hatchery management regions of the country

after around 2008, while adult body size generally remained at similar levels. Moreover, egg

size decreased over the past two decades in all regions.

4.1 Adult return rates (marine survival index)

Adult return rates (marine survival index) of Japanese hatchery chum salmon were correlated with multiple climate-driven stressors that varied by coastal management region, salmon life stage, and seasonal high seas distribution area. Whether regional variation in our results is related to genetics or environment, or both, is unknown. The history of transplantation of Japanese chum salmon (Beacham et al., 2008; Kaeriyama & Qin, 2014) and very high gene flow (Kitada, 2020) may facilitate genetic admixture of hatchery and naturally produced (hereafter, wild) fish throughout the Japanese population. However, the estimated population structure (Nei & Chesser, 1983; Kitada et al., 2021) based on microsatellite markers (Beacham et al., 2008) showed weak but persistent genetic structure and some nested populations (Kitada et al., 2021; Figure S17). The signatures of adaptation to local environmental selection in a population's history can only be captured by integrating all sources of resilience, such as environmental factors, genome-wide single nucleotide polymorphisms (SNPs), and gene expressions (Kishino et al., 2023).

Our adult return rate estimates should be regarded as indices of marine survival because little is known about freshwater mortality of hatchery chum salmon fry after release or mixtures of regional populations in landed commercial catches in Japan (Nagata et al., 2016). Genetic stock identification showed that 80-98% in the mixed samples of chum salmon caught in September-early October along the western coast of the Sea of Okhotsk originated from Hokkaido rivers, while 53-71% originated from Honshu rivers in late October (Saito et al., 2020). The mixed-stock fishery could bias the return rates in each management region in Hokkaido and Honshu. It is generally accepted that most chum salmon in Japan are hatchery-reared fish (Hiroi, 1998; Kaeriyama, 1999; Kobayashi, 1980; Iida et al., 2018; Kaeriyama & Sakaguchi, 2023). We assumed all chum salmon returning to Japan were hatchery-reared. The total proportion of wild chum salmon in Japan is largely unknown. Field studies provide evidence that natural reproduction of chum salmon occurs in Hokkaido (Miyakoshi et al., 2012) and in the Honshu Sea coast of Japan (Iida et al., 2018; Iida et al., 2021). Hatchery salmon have a relative reproductive success (RRS, number of offspring a salmon produces compared to other salmon in the population) that is approximately 50% of the RRS of wild fish (Araki et al., 2007; Christie et al., 2014; Shedd et al., 2022), with considerable variation (Kitada et al., 2011). Reduced RRS can be carried over to the next generation (Araki et al., 2009). We expected low freshwater survival from egg to ocean entry of wild chum salmon in natural streams in Japan. Our marine survival indices may be overestimated depending on the magnitude of recruitment of wild fish, which we expected to be very low compared to the number of hatchery-released fish (100-200 million fry per year in each management region, Figure 2). Our regression analysis

overestimated the intercept proportional to the magnitude of natural recruitment, but regression coefficients were expected to be robust if natural recruitment and genetic mixture were constant over the period of our analysis.

4.2 Egg size and body size of fry at release

Our finding of a nationwide decline in egg size of age-4 and age-5 Japanese hatchery chum salmon over the past two decades is striking. Egg size is an important fitness trait, and larger eggs have a positive effect on juvenile survival and growth in salmonids (Hutchings, 1991; Einum & Fleming, 1999, 2000). Negative relationships between egg size and fecundity have been reported for Pacific salmon (Beacham & Murray, 1993; Fleming & Gross, 1990). Our results suggested that higher relative fecundity (smaller eggs) could contribute to the reduced survival rate of chum salmon in the Hokkaido and Honshu Pacific region. Hasegawa et al. (2021a) showed that the egg size of 5-year-old Japanese chum salmon in 2010 was smaller in more northeasterly locations than in 1994 and 2001, and hypothesized that domestication selection in hatcheries affected by global warming may have caused the temporal shifts in egg size. Selection in hatcheries could impact fecundity and egg size (Heath et al., 2003), although egg size or fecundity was not found to differ between hatchery and wild fish in other populations (Beacham & Murray, 1993; Quinn et al., 2004; Beacham, 2010). An alternative process might be maternal effects, whereby environmental variation (e.g., ocean temperature during egg development; maternal diet) experienced by the mother is translated into phenotypic variation in reproductive traits of offspring (e.g., egg size, age at maturation) (reviewed by Mousseau & Fox, 1998; Thorn et al., 2019).

Our regression model coefficients (practical effect size) suggested that size at release and quality of hatchery fish are particularly important for survival at later ocean stages for Honshu Pacific region chum salmon. The reasons for the continuous decline in fry size at the time of release are unknown. Due to slower swimming speed, small chum salmon fry are more susceptible to nearshore predation than large fry (Hasegawa et al., 2021b). After release, Honshu Pacific region chum fry migrate long distances northward against the Oyashio Current (Figure 1a) to their summer feeding area in the Sea of Okhotsk. On the Sanriku Coast, Honshu Pacific region, where the decline in abundance was most severe (Figure 2), chum salmon fry reared under low food availability (1% body weight) allocated less than half of their energy to growth at high temperatures (14℃) compared to fry reared under high food availability (4% body weight; Iino et al., 2022). In the high food availability group, 32–47% of the energy consumed by the fish should be used for daily growth (Iino et al., 2022). Japanese hatchery chum salmon larger than 2 g (BW) significantly increased long-distance swimming performance and reduced the metabolic cost of swimming against the southward current at 8-

12°C SST (Iino et al., 2024).

4.3 Predators of fry during coastal migrations

Predation is well-accepted as the proximate cause of salmon fry mortality during their coastal

migrations. There are many known fish, bird, and marine mammal predators of Japanese chum

salmon (Okado et al., 2020; Hasegawa et al., 2021b; Okado & Hasegawa, 2024). Long-term

climate change and marine heatwaves are projected to result in poleward shifts in the

distribution of many North Pacific marine species by 2050 (Cheung et al. 2015, Cheung &

Frölicher, 2020). Commercial catch data for coastal marine species in Japan was a useful metric

for identifying poleward shifts in potential predators of chum salmon fry.

We identified yellowtail as a potential marine fish predator of chum salmon fry. Yellowtail are

a subtropical species distributed from Japan and the eastern Korean Peninsula to the Hawaiian

Islands (Robins et al., 1991), and are one of the most commercially important marine fish

species in Japan. A climate regime shift (warming) during the late 1980s was associated with

the northward extension of yellowtail distribution in the Tsushima Warm Current off

northwestern Honshu and Hokkaido (Figure 1a). Global warming was projected to further

expand the suitable thermal habitat of yellowtail northward (Tian et al., 2012). Since 2014,

yellowtail have been caught along the coast of the Sea of Okhotsk in May-December by a setnet

fishery targeting yellowtail, adult salmon (September-December), and other marine fish species

(Hoshino & Fujioka, 2021). The rapid increase in salmon setnet catches of yellowtail led to the

identification of this species as a potential indicator of human (fishing) pressure in Japan

(Hunsicker et al., 2022). In May 2023, the peak month of hatchery releases of chum salmon fry

in the Hokkaido Okhotsk region (Figure S2), schools of yellowtail were often observed on the

sea surface near the central Sea of Okhotsk coast (pers. comm., Takashi Ichikawa). Adult

yellowtail are well-known to feed on small pelagic fish and invertebrates (Robins et al., 1991),

but stomach contents surveys are necessary to confirm yellowtail predation on chum salmon

fry.

Our regression model results indicated yellowtail catches are an important metric of fry chum

survival in the year of release, particularly in the Hokkaido Okhotsk region. Annual estimates

of stock abundance (biomass) or relative abundance (catch per unit effort) would be more

appropriate metrics when considering many potential predator species (e.g., Pauly et al., 2013;

Thorson et al., 2020; Punt, 2023), but were not available for the period of our study. The use of

stock assessment data is a promising approach for developing new indices of climate-driven

predator exposure, especially if predator diet information is available. A similar approach for

high seas migration areas may be possible through future scientific collaborations among

international treaty organizations (PICES-NPFC, 2019).

4.4 Inter- and intra-specific competition during high seas migrations

669 Indirect evidence of competition between Russian pink salmon and many other marine species, including chum salmon, is substantial (see review by Ruggerone et al., 2023). Our results suggest that Russian pink and chum salmon compete for food, space, or other resources with Japanese chum salmon (ages 2-3) in the central Bering Sea in summer and the Gulf of Alaska in winter, while competition with immature (age 1) chum salmon could be weak or negligible in the Sea of Okhotsk and western North Pacific (Naydenko & Temnykh, 2016; Naydenko et al., 2016; Bugaev & Gerlits, 2023). A similar regression analysis supported our finding of competition (negative) with pink salmon in the Bering Sea (Kaeriyama et al., 2025). Inter- and intra-specific competition for prey resources may not be a limiting factor for salmon in the western subarctic North Pacific because the climatic connections to the ecosystem differ from those in the Gulf of Alaska (reviewed by Radchenko et al., 2018; Somov et al., 2024). Annual Russian surveys of juvenile pink and chum salmon biomass and field data on ocean conditions in the Sea of Okhotsk (e.g., Radchenko et al., 2013; Shuntov et al., 2019) could provide information on changes in survival of Japanese chum during their first summer at sea, but we did not have access to these data.

685 Field data on interannual variation in the biomass, relative abundance, or overlap in high seas distribution and diets of Japanese, Russian, and North American chum and pink salmon corresponding to the period of our study are limited due to infrequent and costly high seas salmon research. Analysis of Japanese salmon research vessel data (1972-1998, April-September) indicated density-dependent interactions between pink and chum salmon, including shifts in chum salmon distribution, westward in years of low pink salmon abundance and southeastward in years of high pink salmon abundance (Azumaya & Ishida, 2000). A reanalysis of coastal and high seas salmon research survey data (1953-2022) confirmed the results of earlier studies showing overlap in distributions of chum, pink, and sockeye salmon throughout the North Pacific (Langan et al., 2024). A meta-analysis of salmon trophic ecology data showed similar diet composition of chum, pink, and sockeye salmon throughout the North Pacific, with a diet overlap of 31.8% between chum and pink and 30.9% between chum and sockeye, and diet overlap was highest between pink and sockeye salmon at 46.6% (Graham et al., 2021). The substantial decrease in abundance of Japanese chum salmon after the mid-2000s may have reduced inter- and intra-specific competition of age-1 chum salmon for prey resources. The lack of an association between Russian pink salmon catches and survival of age-1 Japanese chum salmon might also indicate improved growth in the Sea of Okhotsk.

Competition with other salmon is considered an important driver of declines in body size (Oke et al., 2020; Ohlberger et al., 2023). When wild and hatchery salmon populations are abundant, increased competition at sea could lead to reductions in body size by reducing the availability or quality of food resources (Bigler et al., 1996; Jeffrey et al., 2017). The estimated proportion of Asian and North American hatchery chum salmon was 60% of the total abundance of maturing chum salmon during 1990-2015 (Ruggerone & Irvine, 2018). The substantial decline in abundance of Japanese chum salmon after the mid-2000s may have reduced inter- and intra-specific competition for prey resources with Russian chum and pink salmon, resulting in relatively stable body sizes in all regions after about 2008 (Figures S5, S6). However, even when catches of Japanese chum salmon were low, their survival (age–2 and age–3 fish) was negatively correlated with Russian chum and pink salmon catches, suggesting that Russian chum and pink salmon have a competitive advantage over Japanese chum salmon.

Directional swimming to preferred thermal habitat seems to play an important role in the seasonal northward migrations of salmon (Azumaya & Ishida, 2004). If immature Japanese salmon are growing faster with warming ocean temperatures, we would expect their larger body size to increase swimming performance (migration speed and endurance) and to improve their ability to compete with Russian salmon. Alternatively, Russian pink and chum salmon may have a competitive advantage simply because they do not have as far to swim from their overwintering habitat (central North Pacific and Aleutian Islands) as Japanese chum salmon (Gulf of Alaska) to their shared optimal summer feeding habitat in the central Bering Sea. If Russian salmon are first to arrive in the central Bering Sea in summer, they might occupy most of the optimal thermal habitat and reduce the amount of available preferred prey, causing a trophic cascade (Batten et al., 2018). Models project that the optimal thermal habitat of chum and pink salmon will continue to contract and shift northwestward under future climate-change scenarios (e.g., Abdul-Aziz et al., 2011), and we expect Russian salmon to continue to have a competitive advantage over Japanese chum salmon.

We hypothesize that inferior athletic performance in a warming climate may be related to the long-term decline in marine survival of Japanese hatchery chum salmon. Wild-born chum salmon fry were too agile to be netted prior to 1945 in a river on Etrofu Island where Japan operated salmon hatcheries (Yagisawa, 1970). Currently, piscivorous salmonids chase schools of hatchery chum salmon fry, and small fry are easily predated due to their slow swimming speed to escape predators (Hasegawa et al., 2021b). Water temperature and oxygen are important regulators of physiological processes in fish (Powers et al., 1991; Somero, 2022). Swim tunnel experiments show that temperature and body size influence swimming performance in Japanese hatchery chum fry (Iino et al., 2024). Swimming performance has

been least studied in adult chum salmon and Chinook salmon (reviewed by Kraskura et al.,
2024).

Our analysis indicated marine survival of Japanese chum salmon was not related to abundance
of North American pink and chum salmon. This may reflect inter- and intra-specific differences
in ocean distribution of Japanese chum salmon and North American pink and chum salmon. For
example, limited data from salmon research vessel surveys in winter indicate Japanese chum
salmon are distributed farther to the south in the Gulf of Alaska and northeastern North Pacific
than North American chum salmon (reviewed by Myers et al., 2016; Urawa et al., 2016). Past
research indicated negative intraspecific relationships between the abundance of Japanese
hatchery chum salmon and the abundance and annular scale growth of local and regional stock
groups of North American chum salmon (Ruggerone et al., 2011; Agler et al., 2013; Frost et al.,
2021). All our best regional models included North American chum salmon as an explanatory
variable with VIF < 3 (Table S2), but were excluded by our model selection procedure. Our
regression analysis detected a negative association between return rates of Hokkaido Sea of
Japan chum and Russia chum catch (r=-0.44, t=-2.22, p=0.038), while correlations between
regional return rates and North American chum catch were too weak to detect North American
chum salmon as a competitor.

4.5 Ocean conditions in coastal and high seas habitats
The Sea of Japan is a global hotspot for long-term declines in marine fisheries productivity
(Free et al., 2019). After a brief (1998-1999) natural cooling event, anthropogenic climate
change became the principal driver of extreme warming events in the Sea of Japan, particularly
in southern regions (Hayashi et al., 2022). Hatchery chum salmon fry were released mainly in
February-March along the Honshu coast and March-April along the Hokkaido Sea of Japan
(Mar-Apr) (Figure S2). The peak release month in the Hokkaido Okhotsk region was May,
when SSTs were below ~6°C (Figures S11, S12). After release, chum salmon fry migrate in the
direction of the northbound Tsushima Warm Current to the Sea of Okhotsk (Figure 1a). The
negative association of survival with July SST may be related to lengthy exposure to warm
coastal SSTs in the Sea of Japan and delayed arrival at their nursery area in the Sea of Okhotsk.
Juvenile chum salmon were abundant in the coastal Okhotsk Sea from May to June when SST
was between 8 and 13°C, and disappeared from coastal waters after late June when SST
exceeded 13°C (Nagata et al., 2007). Late arrival of juveniles released from the Sea of Japan to
the Sea of Okhotsk may reduce survival during the first summer. Our finding of a negative
effect of summer SST in the Sea of Okhotsk was consistent with that of Kaeriyama et al. (2025).
In contrast, their estimate of the regression coefficient of winter SST in the Gulf of Alaska
(negative) was opposite to ours (positive). The apparent inconsistency seemed to be explained

by the effect of the fry quality and the number of fry released (data not shown). Our regression
model effects of SST (positive or negative) were inferred only for fry released in the Hokkaido
and Honshu Sea of Japan management regions. The causes were unknown.
780
Marine heatwaves and long-term climate change are projected to result in poleward shifts in
the distribution of many North Pacific pelagic fish species by 2050 (Cheung et al. 2015,
Cheung & Frölicher, 2020). Our maps of Japanese chum salmon thermal habitat indicated a
summer northward shift and contraction since 2010; winter habitat remained relatively stable
with a slight northeastward shift in 2017 and 2018 (unpublished figure, S. Kitada). The
frequency of annual marine heatwaves increased in the western and central Bering Sea, and
heatwaves in the eastern Bering were longer and more intense in the past decade (2010–2019)
(Carvalho et al., 2021). Marine heatwaves in the Gulf of Alaska (2013-2016, 2018-2020)
were accompanied by increased acidity and low oxygen, resulting in cascading ecosystem
effects (Hauri et al., 2024). Our analysis showed that SST in the Gulf of Alaska increased in
2014-2015 and 2019-2020 (Figure S13b), which is consistent with the occurrence of marine
heat waves.
793
Ranges of chum salmon thermal habitat used in our analysis were narrower (thermal limits,
4–13°C) than previously published estimates, based largely on historical high seas salmon
survey data collected before the winter 1976-1977 climate regime shift (preferred spring-
summer SST range, 1.3–12.6°C; Langan et al., 2024). The pre-regime shift period (1948-
1975) was characterized by low ocean productivity, cold winter SSTs in the Gulf of Alaska,
and a low abundance of salmon populations that overwinter in the Gulf of Alaska, including
Japanese chum salmon. During periods of low ocean productivity, optimal temperatures for
salmon growth decline (Beauchamp, 2009). Prey quality and energy density of western
Alaska juvenile chum salmon were negatively associated with summer SST in the eastern
Bering Sea during a warm period (marine heat wave) in 2014-2019, potentially decreasing
survival during the critical first winter (Farley et al., 2024). We found that survival of Honshu
Sea of Japan chum salmon was negatively related to SST during the first winter in the western
North Pacific, supporting the hypothesis that juvenile salmon survival during the first fall and
winter in the open ocean is related to ocean conditions (Naydenko & Temnykh, 2016;
Naydenko et al., 2016; Bugaev & Gerlits, 2023).
809
810   **4.6 Adult body size and age at maturity**
811   Adult body size of Japanese hatchery chum salmon declined from the 1970s to the 1980s, likely
812   due to density-dependent effects related to the carrying capacity of Japanese chum salmon in
813   the open ocean (Ishida et al., 1993; Hayashizaki & Ida, 1993; Kaeriyama, 1998; Morita &

Fukuwaka, 2007). Analysis of high seas salmon density and mean FLs indicated declining growth of age 2-4 chum salmon from 1972 to 1998 was directly related to intra-specific competition in the Bering Sea, and was influenced indirectly by abundance of pink salmon (Azumaya & Ishida, 2000). After 1992, the body size of adult Japanese chum salmon increased, possibly the result of natural selection and phenotypic plasticity induced by environmental changes after the termination of Asian high-seas driftnet fisheries (Fukuwaka & Morita, 2008). Our analysis showed that adult body size (age-specific FL) of Japanese chum salmon increased from 1997 to around 2003, and then decreased until the mid-2000s (Figures S5, S6). Until the mid-2000s, the decreasing trends in adult body size (age–4 and age–5) of Japanese hatchery chum salmon agreed with those of Alaskan chum, coho, and sockeye salmon (Oke et al., 2020; Ohlberger et al., 2023). The body size of Japanese hatchery chum salmon was relatively stable in all regions after about 2008, although there were fluctuations (Figures S5, S6), while the body size of Alaskan salmon continued to decline (Figure 2 in Oke et al., 2020). After 2003, the positive effect of the relaxation of high-seas fishing pressure on adult body size might have been masked or reduced by density-dependent effects related to climate change effects on oceanic thermal habitats of salmon.

We observed a trend toward younger age at maturity (increase in age-4 fish) and relatively constant trends in adult body size after 2008, suggesting an increase in growth rates of Japanese chum salmon. A study of annular growth history on the scales of adult (age-4) Japanese hatchery chum salmon showed that growth during the 2^{nd} and 3^{rd} years is more strongly associated with the probability of maturation than age or age-specific body size, which is affected by environmental conditions (Morita & Fukuwaka, 2006). Summer SSTs in the Bering Sea and winter SSTs in the Gulf of Alaska, where immature age-2 and age-3 Japanese chum salmon are distributed, were warmer after 2010 within the optimal growth and feeding range (Figure S13b), possibly resulting in increased growth rates of immature age-2 and age-3 fish. Our results are consistent with improved growth of adult Alaskan sockeye salmon that was associated with warm winter SST (Ohlberger et al., 2023). However, warm summer SST was associated with reduced growth of Alaskan Chinook and sockeye salmon (Yasumiishi et al., 2020; Ohlberger et al., 2023).

In 2022, the number of chum salmon returning to Japan unexpectedly increased in the Hokkaido Sea of Japan and the Sea of Okhotsk regions, while remaining very low in the Honshu Pacific, the former main production area. In the Chitose River in the Hokkaido Sea of Japan, where Japan's oldest hatchery operates, 587,000 fish were returned, surpassing the historical record of 551,000 in 1995 (Chitose Aquarium, 2023). The sudden increase in 2022 surprised fishermen, hatchery managers, local people, and biologists. However, returning chum numbers declined

again in all geographic regions in 2023, reaching historic lows (a total of about 18 million in Japan) in 2024. In the Hokkaido Sea of Japan and Okhotsk regions, the peak in adult returns in 2022 (Figure 2) was related to an increase in age–4 fish released as fry in 2019 and increased total return rates of all age groups (Figure S18), indicating better growth and survival of the 2019 year class. In contrast, total return rates decreased in other regions (Figure S18), despite the generally increasing proportion of age–2, age–3, and age–4 fish in all regions after approximately 2008 (Figure 3). Thermal habitat (8–10°C) within the optimal thermal habitat range for growth and feeding (8–12°C) of chum salmon was widely distributed in the Sea of Okhotsk in July 2019 and in the central Bering Sea in September 2020 (Figure S19), where age-1 Japanese chum salmon are distributed during their second summer growing season. Expansion of optimal thermal habitat during the first and second summers likely improved the growth and survival of immature chum salmon in 2020 and 2021, and may have resulted in the anomalous peak returns of age-4 fish in 2022.

4.7 Management implications

Hatchery experiments to test appropriate fry size and time of release and nearshore monitoring of SSTs might provide an adaptive approach for improving adult returns of Japanese chum salmon (Saito & Nagasawa, 2009; Nagata et al., 2016; Iida et al., 2021; Saito, 2022; Iida, 2024; Honda et al., 2025). For example, in the most successful management region (Hokkaido Okhotsk), the peak time of release is in May (SST, 4.7–6.2°C). In the Sea of Japan region, SST at release and summer SST (July) in the Sea of Okhotsk were negatively associated with return rate. Adjustment of release timing to allow fry to arrive in the Sea of Okhotsk in May or early June or both might improve nearshore survival. In the Pacific region of Japan, parental fecundity (negative) and fry size at release (positive) were associated with salmon survival. Size at release declined in most regions after around 2010. Our results suggest that hatchery fry quality (including size at release) is particularly important for survival of later ocean stages. Experiments to assess fry quality, metabolism, and swimming performance of Japanese chum salmon (e.g., Iino et al. 2022, 2024; Abe et al., 2024) should provide information useful to management.

Investigations of many other potentially important stressors beyond the scope of our study remain to be addressed. For example, relationships between the marine survival of salmon and climate-driven changes in marine fisheries operating along high seas salmon migration routes and within the 200-mile EEZs of other nations. In Alaska, genetic analyses show that Asian (Russia and Japan) populations dominate early (June) catches by salmon fisheries targeting sockeye salmon along the south side of the Alaska Peninsula in early (June), when chum salmon are migrating from winter (North Pacific) to summer (Bering Sea) feeding grounds (Dann et al.,

2023). Asian stocks of chum salmon also dominate prohibited species bycatch of salmon in the US walleye pollock (*Gadus chalcogrammus*) fishery in the eastern Bering Sea in early summer (Barry et al., 2022). Bycatches of Asian chum salmon (ocean-age-3 and age-4; unknown maturity stage) by both fisheries in 2021 were unexpectedly high (Barry et al., 2022; Dann et al., 2023). The Alaska bycatch of Asian chum salmon preceded the spike in adult returns to Japan in 2022. Seeking predictive (vs. retrospective) indices would benefit preseason forecasting of adult salmon returns. In the absence of quantitative marine stock assessment surveys of salmon, potentially predictive indices might be used in a qualitative "stoplight" approach to pre-season forecasting (Caddy, 2002; NOAA, 2024). This method requires an adaptive approach because relationships between salmon survival and index variables change over time (non-stationarity; Litzow et al., 2018; Ohlberger et al., 2022).

5 | SUMMARY AND CONCLUSIONS

We used long-term data series collected from the world's largest hatchery salmon program to investigate potential stressors of hatchery practices, ocean conditions (SST), and predators and competitors related to the 25-year population dynamics of Japanese chum salmon (1998-2023). We assessed stressors on survival along known coastal and open ocean migration routes in the context of climate change. Our results highlighted the regional differences in stressors associated with the decline in marine survival of Japanese chum salmon. Parental fecundity and fry body size at release were associated with survival in the Pacific region, where the decline in abundance was most severe. The northward migration shift of a piscivorous marine fish indicated a possible new predator in Hokkaido Okhotsk, the most successful management region. Effects of sea temperature were inferred for fry released in the Sea of Japan, while no effects of sea temperature were found at any life stage for hatchery releases in other regions. Russian chum and pink salmon abundance was negatively associated with survival of Japanese chum salmon in all regions. Importantly, we observed a trend toward younger age at maturity in all hatchery management regions of the country after around 2008, while adult body size generally remained at similar levels, indicating that growth rates are increasing. Moreover, egg size decreased over the past two decades in all regions. Age at maturity, fecundity, and egg size are key demographic parameters that contribute to the productivity of salmon populations, and the decadal-scale changes observed herein could result in significant ecosystem, fishery, and societal losses. The results of our study highlight the need for monitoring and analysis with updated information to rethink the role of hatcheries and develop adaptive management plans for future sustainability and conservation of salmon in a changing climate.

AUTHOR CONTRIBUTIONS

Shuichi Kitada: Conceptualization (lead); data curation; formal analysis (lead); funding

acquisition; investigation; methodology; software; validation; visualization (lead); writing – original draft; writing – review and editing. Katherine W. Myers: Conceptualization; writing – review and editing (lead). Hirohisa Kishino: Conceptualization; formal analysis; funding acquisition; methodology (lead); validation (lead); visualization; writing – review and editing. All authors approved the final version of the manuscript.

ACKNOWLEDGMENTS
We sincerely acknowledge the thousands of biologists, fishery and hatchery managers, and fishers who contributed to the collection of data analyzed in our manuscript. The efforts of APG, FRA, HSEA, JMA, MAFF, NOAA, and NPAFC to create public databases made our study possible. We thank Brian Riddell and the reviewers for their constructive comments, which significantly improved earlier versions of the manuscript. Takashi Ichikawa provided information on chum salmon and yellowtail. This study was supported by Japan Society for the Promotion of Science Grants-in-Aid for Scientific Research KAKENHI No. 22K11950 for HK and 18K0578116 for SK.

CONFLICT OF INTEREST STATEMENT
The authors declare that no conflict of interest exists.

DATA AVAILABILITY STATEMENT
The data generated or analyzed in this study are fully described in the published article and its Supplementary Information. Analysis scripts and data will be available at https://doi.org/10.5281/zenodo.14868156 on acceptance.

ORCID
Shuichi Kitada https://orcid.org/0000-0001-5838-0374
Hirohisa Kishino https://orcid.org/0000-0002-3244-359X

REFERENCES
Abdul-Aziz, O. I., Mantua, N. J., & Myers, K. W. (2011). Potential climate change impacts on thermal habitats of Pacific salmon (*Oncorhynchus* spp.) in the North Pacific Ocean and adjacent seas. *Canadian Journal of Fisheries and Aquatic Sciences*, 68, 1660–1680. doi.org/10.1139/f2011-079

Abe, T. K., Kitagawa, T., Iino, Y., Ito, M., & Sato, K. (2024). Ecological features of upriver migration in Kitakami River chum salmon and their connection to aerobic thermal performance. *Conservation Physiology*, 12, coae087. https://doi.org/10.1093/conphys/coae087

Agler, B. A., Ruggerone, G. T., Wilson, L. I., & Mueter, F. J. (2013). Historical growth of Bristol Bay and Yukon River, Alaska chum salmon (Oncorhynchus keta) in relation to climate and inter-and intraspecific competition. *Deep Sea Research Part II: Topical Studies in Oceanography*, 94, 165–177. doi.org/10.1016/j.dsr2.2013.03.028

Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. In B. N. Petrov & F. Csaki (Eds.), *Second International Symposium on Information Theory* (pp. 267–281). Budapest: Akademiai Kiado. Reprinted in S. Kotz & N. L. Johnson, Eds. (1992) *Breakthroughs in Statistics Volume I, Springer Series in Statistics*, 610-624. New York: Springer-Verlag. doi.org/10.1007/978-1-4612-0919-5_38

Amoroso, R. O., Tillotson, M. D., & Hilborn, R. (2017). Measuring the net biological impact of fisheries enhancement: Pink salmon hatcheries can increase yield, but with apparent costs to wild populations. *Canadian Journal of Fisheries and Aquatic Sciences*, 74, 1233–1242. doi.org/10.1139/cjfas-2016-0334

APG (Aomori Prefectural Government) (2024). *Chum salmon catch statistics*, Aomori: APG. (in Japanese) www.pref.aomori.lg.jp/soshiki/nourin/sshinko/sake.html (accessed 20 November 2024).

Araki, H., Cooper, B., & Blouin, M. S. (2007). Genetic effects of captive breeding cause a rapid, cumulative fitness decline in the wild. *Science*, 318, 100–103. doi.org/10.1126/science.1145621

Araki, H., Cooper, B., & Blouin, M. S. (2009). Carry-over effect of captive breeding reduces reproductive fitness of wild-born descendants in the wild. *Biology Letters*, 5, 621–624. doi.org/10.1098/rsbl.2009.0315

Azumaya, T., & Ishida, Y. (2004). An evaluation of the potential influence of SST and currents on the oceanic migration of juvenile and immature chum salmon (*Oncorhynchus keta*) by a simulation model. *Fisheries Oceanography*, 13, 10–23. https://doi.org/10.1046/j.1365-2419.2003.00271.x

Azumaya, T., Nagasawa, T., Temnykh, O. S., & Khen, G. V. (2007). Regional and seasonal differences in temperature and salinity limitations of Pacific Salmon (*Oncorhynchus* spp.). *North Pacific Anadromous Fish Commission Bulletin*, 4, 179–187. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-4/179-187Azumaya.pdf

Bahri, T., Vasconcellos, M., Welch, D.J., Johnson, J., Perry, R.I., Ma, X. & Sharma, R., Eds. (2021). Adaptive management of fisheries in response to climate change. *FAO Fisheries and Aquaculture Technical Paper* 667, Rome: FAO. doi.org/10.4060/cb3095en

Barry, P. D., Kondzela, C. M., Whittle, J. A., Watson, J. T., Karpan, K., D'Amelio, K., & Larson, W. A. (2022). Genetic stock composition analysis of chum salmon from the prohibited species catch of the 2021 Bering Sea walleye pollock trawl fishery. U.S.

999   Department of Commerce, *NOAA Technical Memorandum NMFS-AFSC-*
1000   *454*.repository.library.noaa.gov/view/noaa/47923/noaa_47923_DS1.pdf

1001 Batten, S. D., Ruggerone, G. T., & Ortiz, I. (2018). Pink salmon induce a trophic cascade in
1002   plankton populations in the southern Bering Sea and around the Aleutian Islands.
1003   *Fisheries Oceanography*, 27, 548-559. https://doi.org/10.1111/fog.12276

1004 Beacham, T. D. (2010). Revisiting trends in the evolution of egg size in hatchery-enhanced
1005   populations of Chinook salmon from British Columbia. *Transactions of the American*
1006   *Fisheries Society*, 139, 579–585. doi.org/10.1577/T09-093.1

1007 Beacham, T. D., & Murray, C. B. (1993). Fecundity and egg size variation in North American
1008   Pacific salmon (*Oncorhynchus*). *Journal of Fish Biology,* 42, 485–508.
1009   doi.org/10.1111/j.1095-8649.1993.tb00354.x

1010 Beacham, T. D., Sato, S., Urawa, S., Le, K. D., & Wetklo, M. (2008). Population structure and
1011   stock identification of chum salmon *Oncorhynchus keta* from Japan determined by
1012   microsatellite DNA variation. *Fisheries Science*, 74, 983–994. doi.org/10.1111/j.1444-
1013   2906.2008.01616.x

1014 Beacham, T. D., Candy, J. R., Sato, J.S., Urawa, S., Le, K. D., & Wetklo, M. (2009). Stock
1015   origin of chum salmon (*Oncorhynchus keta*) in the Gulf of Alaska during winter as
1016   estimated with microsatellites. *North Pacific Anadromous Fish Commission Bulletin*, 5,
1017   15–23. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-5/NPAFC_Bull_5_015-
1018   023Beacham.pdf

1019 Beamish, R. J., Ed. (2018). *The Ocean Ecology of Pacific Salmon and Trout*. Bethesda:
1020   American Fisheries Society. doi.org/10.47886/9781934874455

1021 Beamish, R. J., & Mahnken, C. (2001). A critical size and period hypothesis to explain natural
1022   regulation of salmon abundance and the linkage to climate and climate change.
1023   *Progress in Oceanography*, 49, 423–437. doi.org/10.1016/S0079-6611(01)00034-9

1024 Beauchamp, D. (2009). Bioenergetic ontogeny: Linking climate and mass specific feeding to
1025   life-cycle growth and survival of salmon. In C. Krueger & C. E. Zimmerman (Eds.),
1026   *Pacific salmon: Ecology and Management of Western Alaska's Populations*
1027   (Symposium 70, pp. 53–72). Bethesda: American Fisheries Society.
1028   doi.org/10.47886/9781934874110

1029 Bigler, B. S., Welch, D. W., & Helle, J. H. (1996). A review of size trends among North
1030   Pacific salmon (*Oncorhynchus* spp.). *Canadian Journal of Fisheries and Aquatic*
1031   *Sciences*, 53, 455–465. doi.org/10.1139/f95-181

1032 Bradshaw, W. E., & Holzapfel, C. M. (2006). Evolutionary response to rapid climate change.
1033   *Science*, 312, 1477-1478. doi.org/10.1126/science.1127000

1034 Bryndum-Buchholz, A., Tittensor, D. P., & Lotze, H. K. (2021). The status of climate change
1035   adaptation in fisheries management: Policy, legislation and implementation. *Fish and*

Fisheries, 22, 1248–1273. doi.org/10.1111/faf.12586

Bugaev, A. V., & Gerlits, A. I. (2023). The characteristics of feeding migrations of hatchery-reared Pacific salmon juveniles in the Sea of Okhotsk Basin and adjacent Pacific waters in the autumn–winter period (regional identification, abundance, distribution of catches, biological characteristics, and mortality rate estimates). *Russian Journal of Marine Biology*, 49, 546–568. doi.org/10.1134/S1063074023070039

Burnham, K.P., & Anderson, D.R. (2002). *Model Selection and Multimodel Inference: A Practical Information- Theoretic Approach*, *2nd edition*. New York: Springer-Verlag. link.springer.com/book/10.1007/b97636

Caddy, J. F. (2002). Limit reference points, traffic lights, and holistic approaches to fisheries management with minimal stock assessment input. *Fisheries Research*, 5, 133–137. doi.org/10.1016/S0165-7836(01)00343-5

Carvalho, K. S., Smith, T. E., & Wang, S. E. (2021). Bering Sea marine heatwaves: Patterns, trends and connections with the Arctic. *Journal of Hydrology*, 600, 126462. doi.org/10.1016/j.jhydrol.2021.126462

Cheung, W. W., & Frölicher, T. L. (2020). Marine heatwaves exacerbate climate change impacts for fisheries in the northeast Pacific. *Scientific Reports*, 10, 6678. doi.org/10.1038/s41598-020-63650-z

Cheung, W. W., Brodeur, R. D., Okey, T. A., & Pauly, D. (2015). Projecting future changes in distributions of pelagic fish species of Northeast Pacific shelf seas. *Progress in Oceanography*, 130, 19-31. doi.org/10.1016/j.pocean.2014.09.003

Cheung, W. W., Frölicher, T. L., Lam, V. W., Oyinlola, M. A., Reygondeau, G., Sumaila, U. R., ... & Wabnitz, C. C. (2021). Marine high temperature extremes amplify the impacts of climate change on fish and fisheries. *Science Advances*, 7, eabh0895. doi.org/10.1126/sciadv.abh0895

Chitose Aquarium (2023). Capture Information of chum salmon. Chitose Aquarium, Chitose. https://chitose-aq.jp/ (accessed, 8 August 2023).

Christie, M. R., Ford, M. J., & Blouin, M. S. (2014). On the reproductive success of early-generation hatchery fish in the wild. *Evolutionary Applications*, 7, 883–896. doi.org/10.1111/eva.12183

Cline, T. J., Ohlberger, J., & Schindler, D. E. (2019). Effects of warming climate and competition in the ocean for life-histories of Pacific salmon. *Nature Ecology & Evolution*, 3, 935–942. doi.org/10.1038/s41559-019-0901-7

Connors, B., Malick, M. J., Ruggerone, G. T., Rand, P., Adkison, M., Irvine, J. R., ... & Gorman, K. (2020). Climate and competition influence sockeye salmon population dynamics across the Northeast Pacific Ocean. *Canadian Journal of Fisheries and Aquatic Sciences*, 77, 943–949. doi.org/10.1139/cjfas-2019-0422

Connors, B., Ruggerone, G. T., & Irvine, J. R. (2024). Adapting management of Pacific salmon to a warming and more crowded ocean. *ICES Journal of Marine Science*, fsae135. doi.org/10.1093/icesjms/fsae135

Crozier, L. G., & Siegel, J. E. A. (2023). Comprehensive review of the impacts of climate change on salmon: strengths and weaknesses of the literature by life stage. *Fishes*, 8, 1–50. doi.org/10.3390/fishes8060319

Crozier, L. G., Burke, B. J., Chasco, B. E., Widener, D. L., & Zabel, R. W. (2021). Climate change threatens Chinook salmon throughout their life cycle. *Communications Biology*, 4, 222. doi.org/10.1038/s42003-021-01734-w

Cunningham, C. J., Westley, P. A., & Adkison, M. D. (2018). Signals of large scale climate drivers, hatchery enhancement, and marine factors in Yukon River Chinook salmon survival revealed with a Bayesian life history model. *Global Change Biology*, 24, 4399–4416. doi.org/10.1111/gcb.14315

Dahlke, F. T., Butzin, M., Nahrgang, J., Puvanendran, V., Mortensen, A., Pörtner, H. O., & Storch, D. (2018). Northern cod species face spawning habitat losses if global warming exceeds 1.5 C. *Science Advances*, 4, eaas8821. doi.org/10.1126/sciadv.aas8821

Dann, T. H., Hoyt, H. A., Lee, E. M., Fox, E. K. C., & Foster, M. B. (2023). Genetic stock composition of chum salmon harvested in commercial salmon fisheries of the South Alaska Peninsula, 2022. *Alaska Department of Fish and Game, Special Publication*, 23-07. www.adfg.alaska.gov/FedAidPDFs/SP23-07.pdf

Einum, S., & Fleming, I. A. (1999). Maternal effects of egg size in brown trout (*Salmo trutta*): Norms of reaction to environmental quality. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 266, 2095–2100. doi.org/10.1098/rspb.1999.0893

Einum, S., & Fleming, I. A. (2000). Highly fecund mothers sacrifice offspring survival to maximize fitness. *Nature*, 405, 565–567. doi.org/10.1038/35014600

FAO (United Nations Food and Agriculture Organization) (2022). *The State of World Fisheries and Aquaculture: 2022: Towards Blue Transformation*. Rome: FAO. digitallibrary.un.org/record/3978694?v=pdf

Farley Jr, E. V., Yasumiishi, E. M., Murphy, J. M., Strasburger, W., Sewall, F., Howard, K., ... & Moss, J. H. (2024). Critical periods in the marine life history of juvenile western Alaska chum salmon in a changing climate. *Marine Ecology Progress Series*, 726, 149–160. https://doi.org/10.3354/meps14491

Fisheries Agency of Japan (1955). On the salmon in waters adjacent to Japan. *International North Pacific Fisheries Commission Bulletin*, 1–2. www.npafc.org/wp-content/uploads/Bulletin1FRA.pdf

Fleming, I. A., & Gross, M. R. (1990). Latitudinal clines: A trade-off between egg number and size in Pacific salmon. *Ecology*, 71, 1–11. doi.org/10.2307/1940241

FRA (Japan Fisheries Research and Education Agency) (2023). *Salmon Data Base.* Yokohama: FRA. (in Japanese) www.fra.go.jp/shigen/salmon/sdb.html (accessed 20 November 2024).

FRA (Japan Fisheries Research and Education Agency) (2024). *Chum salmon catch and release statistics.* Yokohama: FRA. (in Japanese) www.fra.go.jp/shigen/salmon/index.html (accessed 5 March 2024).

Free, C. M., Thorson, J. T., Pinsky, M. L., Oken, K. L., Wiedenmann, J., & Jensen, O. P. (2019). Impacts of historical warming on marine fisheries production. *Science*, 363, 979–983. doi.org/10.1126/science.aau1758

Frölicher, T. L., Fischer, E. M., & Gruber, N. (2018). Marine heatwaves under global warming. *Nature*, 560, 360–364. doi.org/10.1038/s41586-018-0383-9

Frost, T. J., Yasumiishi, E. M., Agler, B. A., Adkison, M. D., & McPhee, M. V. (2021). Density-dependent effects of eastern Kamchatka pink salmon (*Oncorhynchus gorbuscha*) and Japanese chum salmon (*O. keta*) on age-specific growth of western Alaska chum salmon. *Fisheries Oceanography*, 30, 99-109. doi.org/10.1111/fog.12505

Fukuwaka, M., & Suzuki, T. (1998). Role of a riverine plume as a nursery area for chum salmon *Oncorhynchus keta*. *Marine Ecology Progress Series*,173, 289–297. doi.org/10.3354/meps173289

Fukuwaka, M., & Morita, K. (2008). Increase in maturation size after the closure of a high seas gillnet fishery on hatchery-reared chum salmon *Oncorhynchus keta*. *Evolutionary Applications*,1, 376–387. doi.org/10.1111/j.1752-4571.2008.00029.x

Graham, C., Pakhomov, E. A., & Hunt, B. P. (2021). Meta-analysis of salmon trophic ecology reveals spatial and interspecies dynamics across the North Pacific Ocean. *Frontiers in Marine Science*, 8, 618884. doi.org/10.3389/fmars.2021.618884

Groot, C., & Margolis, L. (1991). *Pacific Salmon Life Histories*. Vancouver: University of British Columbia Press. www.ubcpress.ca/pacific-salmon-life-histories

Hasegawa, K., Okamoto, Y., & Morita, K. (2021a). Temporal trends in geographic clines of chum salmon reproductive traits associated with global warming and hatchery programmes. *Ecological Solutions and Evidence*, 2, e12107. doi.org/10.1002/2688-8319.12107

Hasegawa, K., Honda, K., Yoshiyama, T., Suzuki, K., & Fukui, S. (2021b). Small biased body size of salmon fry preyed upon by piscivorous fish in riverine and marine habitats. *Canadian Journal of Fisheries and Aquatic Sciences*, 78, 631–638. doi.org/10.1139/cjfas-2020-0339

Hauri, C., Pages, R., Hedstrom, K., Doney, S. C., Dupont, S., Ferriss, B., & Stuecker, M. F. (2024). More than marine heatwaves: A new regime of heat, acidity, and low oxygen compound extreme events in the Gulf of Alaska. *AGU Advances*, 5, e2023AV001039.

Hayashi, M., Shiogama, H., & Ogura, T. (2022). The contribution of climate change to increasing extreme ocean warming around Japan. *Geophysical Research Letters*, 49, e2022GL100785. doi.org/10.1029/2022GL100785

Hayashizaki, K. I., & Ida, H. (1993). Size decrease of chum salmon, *Oncorhynchus keta* in Tohoku districts, Japan. In A. Thompson (Ed.), *International Symposium on Biological Interactions of Enhanced and Wild Salmonids* (p. 45*)*. Canadian Special Publication of Fisheries and Aquatic Sciences. Ottawa: Department of Fisheries and Oceans. waves-vagues.dfo-mpo.gc.ca/library-bibliotheque/217398.pdf

Hilborn, R., & Sibert, J. (1988). Adaptive management of developing fisheries. *Marine Policy*, 12, 112–121. doi.org/10.1016/0308-597X(88)90039-5

Hilborn, R., Quinn, T. P., Schindler, D. E., & Rogers, D. E. (2003). Biocomplexity and fisheries sustainability. *Proceedings of the National Academy of Sciences*, 100, 6564–6568. doi.org/10.1073/pnas.1037274100

Hiroi, O. (1998). Historical trends of salmon fisheries and stock conditions in Japan. *North Pacific Anadromous Fish Commission Bulletin*, 1, 23–27. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-1/page-23-27Hiroi.pdf

Honda, K., Kawakami, T., Suzuki, K., Watanabe, K., & Saito, T. (2017). Growth rate characteristics of juvenile chum salmon *Oncorhynchus keta* originating from the Pacific coast of Japan and reaching Konbumori, eastern Hokkaido. *Fisheries Science*, 83, 987–996. doi.org/10.1007/s12562-017-1137-6

Honda, K., Sato, T., Kuroda, H., & Saito, T. (2021). Initial growth characteristics of poor-return stocks of chum salmon *Oncorhynchus keta* originating from the Okhotsk and Nemuro regions in Hokkaido on the basis of scale analysis. *Fisheries Science*, 87, 653–663. doi.org/10.1007/s12562-021-01538-1

Honda, K., Shirai, K., Morishita, T., & Saito, T. (2025). Characteristics of juvenile chum salmon carrying out inverse migrations after ocean entry from rivers along the Pacific coast of Hokkaido, Japan. *Fisheries Oceanography*, 34, e12698. doi.org/10.1111/fog.12698

Hoshino, N., & Fujioka, T. (2021). Characteristics of yellowtail catch in Hokkaido, Japan in the 2010s. *Scientific Reports of Hokkaido Fisheries Research Institutes*, 100, 71–82. (in Japanese)

HSEA (Hokkaido Salmon Enhancement Association) (2024). Chum salmon catch and release statistics, Sapporo: HSEA. (in Japanese) www.sake-masu.or.jp (accessed 31 January 2024).

Hunsicker, M. E., Boldt, J. L., Blasiak, R., Hazen, E. L., Kulik, V. V., Ryu, J., ... & Wakita, K., Eds. (2022). Report of Working Group 36 on Common Ecosystem Reference Points across PICES Member Countries. *PICES Scientific Report*, (64), 155 pp. meetings.pices.int/publications/scientific-reports/Report64/Rpt64.pdf

Hutchings, J. A. (1991). Fitness consequences of variation in egg size and food abundance in brook trout *Salvelinus fontinalis*. *Evolution*, 45, 1162–1168. doi.org/10.1111/j.1558-5646.1991.tb04382.x

Iida, M. (2024). Relationship between in-river return rate of hatchery-origin chum salmon Oncorhynchus keta and coastal water temperature and body weight at release on the Japan Sea side of Honshu, Japan. *Environmental Biology of Fishes*, 107, 1247–1261. https://doi.org/10.1007/s10641-024-01633-z

Iida, M., Yoshino, K., & Katayama, S. (2018). Current status of natural spawning of chum salmon *Oncorhynchus keta* in rivers with or without hatchery stocking on the Japan Sea side of northern Honshu, Japan. *Fisheries science*, 84, 453–459. doi.org/10.1007/s12562-018-1192-7

Iida, M., Yagi, Y., & Iseki, T. (2021). Occurrence of wild chum salmon fry in the surf zone, and spawning and emergence timing in the adjacent nonstocked river in Niigata Prefecture, Japan. *Fisheries Science*, 87, 549–557. doi.org/10.1007/s12562-021-01535-4

Iino, Y., Kitagawa, T., Abe, T. K., Nagasaka, T., Shimizu, Y., Ota, K., ... & Kawamura, T. (2022). Effect of food amount and temperature on growth rate and aerobic scope of juvenile chum salmon. *Fisheries Science*, 88, 397–409. doi.org/10.1007/s12562-022-01599-w

Iino, Y., Abe, T. K., Shimizu, Y., Nagasaka, T., & Kitagawa, T. (2024). Body size- and temperature-related metabolic traits of juvenile chum salmon during northward migration. *Canadian Journal of Fisheries and Aquatic Sciences*, 81, 1514–1527. doi.org/10.1139/cjfas-2023-0334

Ishida, Y., Ito, S. O., Kaeriyama, M., McKinnell, S., & Nagasawa, K. (1993). Recent changes in age and size of chum salmon (*Oncorhynchus keta*) in the North Pacific Ocean and possible causes. *Canadian Journal of Fisheries and Aquatic Sciences*, 50, 290–295. doi.org/10.1139/f93-033

James, S. E., Doherty, B., Cox, S. P., Pearsall, I. A., & Riddell, B. (2023). Size and timing of hatchery releases influence juvenile-to-adult survival rates of British Columbia Chinook and coho salmon. *Canadian Journal of Fisheries and Aquatic Sciences*, 80, 700–718. doi.org/10.1139/cjfas-2022-0121

Jeffrey, K. M., Côté, I. M., Irvine, J. R., & Reynolds, J. D. (2017). Changes in body size of Canadian Pacific salmon over six decades. *Canadian Journal of Fisheries and Aquatic*

Sciences, 74, 191–201. doi.org/10.1139/cjfas-2015-0600

JMA (Japan Meteorological Agency) (2022). *Coastal sea surface temperature information.*

Tokyo: JMA. (in Japanese) www.jma-net.go.jp (accessed 23 August 2022).

Kaeriyama, M. (1989). Aspects of salmon ranching in Japan. *Physiology & Ecology, Japan*,

Special Volume 1, 625–638. www.researchgate.net/publication/303362061

Kaeriyama, M. (1998). Dynamics of chum salmon, *Oncorhynchus keta*, populations released

from Hokkaido, Japan. *North Pacific Anadromous Fish Commission Bulletin*, 1, 90–

102. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-1/page-90-

102Kaeriyama.pdf

Kaeriyama, M. (1999). Hatchery programmes and stock management of salmonid populations

in Japan. In B. R. Howell, E. Moksness, & T. Svåsand (Eds.), *Stock Enhancement and*

Sea Ranching (pp. 153–167). Oxford: Blackwell.

www.researchgate.net/publication/246360103

Kaeriyama, M. (2008). Ecosystem-based sustainable conservation and management of Pacific

salmon. In K. Tsukamoto, T. Kawamura, T. Takeuchi, T. D. Bard, Jr., & M.J. Kaiser

(Eds.), *Fisheries for Global Welfare and Environment* (pp. 371–380). Tokyo: Terrapub.

wcfs.fisheries.org/proceedings-of-the-fifth-world-fisheries-congress/

Kaeriyama, M. (2022). Warming climate impacts on production dynamics of southern

populations of Pacific salmon in the North Pacific Ocean. *Fisheries Oceanography*, 32,

121–132. doi.org/10.1111/fog.12598

Kaeriyama, M., & Qin, Y. X. (2014). Biological interactions between wild and hatchery-

produced Pacific salmon. In P. T. K. Woo, D. J. Noakes (Eds.), *Salmon: Biology,*

Ecological Impacts and Economic Importance (pp. 223–238). New York: Nova Science

Publishers.

Kaeriyama, M., & Sakaguchi, I. (2023). Ecosystem-based sustainable management of chum

salmon in Japan's warming climate. *Marine Policy*, 157, 105842.

doi.org/10.1016/j.marpol.2023.105842

Kaeriyama, M., Alabia, I. D., & Urawa, S. (2025). Production trend of Hokkaido chum

salmon estimated by multivariable models incorporating environmental factors and

biological interactions in the North Pacific Ocean. *North Pacific Anadromous Fish*

Commission Technical Report, 23, 34–39. doi.org/10.23849/npafctr23/4bb8ty

Kaeriyama, M., Seo, H., Kudo, H., & Nagata, M. (2012). Perspectives on wild and hatchery

salmon interactions at sea, potential climate effects on Japanese chum salmon, and the

need for sustainable salmon fishery management reform in Japan. *Environmental*

Biology of Fishes, 94, 165–177. doi.org/10.1007/s10641-011-9930-z

Kaeriyama, M., Seo, H., Kudo, H., & Nagata, M. (2012). Perspectives on wild and hatchery

salmon interactions at sea, potential climate effects on Japanese chum salmon, and the

need for sustainable salmon fishery management reform in Japan. *Environmental Biology of Fishes*, 94, 165–177. doi.org/10.1007/s10641-011-9930-z

Kasugai, K., Saneyoshi, H., Aoyama, T., Shinriki, Y., Iijima, A., & Miyakoshi, Y. (2016). Early marine migration of juvenile chum salmon along the Pacific coast of eastern Hokkaido. *North Pacific Anadromous Fish Commission Bulletin*, 6, 61–72. doi.org/10.23849/npafcb6/61.72

Kishino, H., Nakamichi, R., & Kitada, S. (2023). Genetic adaptations in the population history of *Arabidopsis thaliana*. *G3: Genes, Genomes, Genetics*, 13, jkad218. doi.org/10.1093/g3journal/jkad218

Kitada, S. (2014). Japanese chum salmon stock enhancement: Current perspective and future challenges. *Fisheries Science*, 80, 237–249. doi.org/10.1007/s12562-013-0692-8

Kitada, S. (2018). Economic, ecological and genetic impacts of marine stock enhancement and sea ranching: A systematic review. *Fish and Fisheries*, 19, 511–532. doi.org/10.1111/faf.12271

Kitada, S. (2020). Lessons from Japan marine stock enhancement and sea ranching programmes over 100 years. *Reviews in Aquaculture*, 12, 1944–1969. doi.org/10.1111/raq.12418

Kitada, S., Kishino, H., & Hamasaki, K. (2011). Bias and significance of relative reproductive success estimates based on steelhead trout (*Oncorhynchus mykiss*) data: A Bayesian meta-analysis. *Canadian Journal of Fisheries and Aquatic Sciences*, 68, 1827–1835. doi.org/10.1139/f2011-087

Kitada, S., Nakamichi, R., & Kishino, H. (2021). Understanding population structure in an evolutionary context: population-specific F_{ST} and pairwise F_{ST}. G3, 11(11), jkab316. doi.org/10.1093/g3journal/jkab316

Kobayashi, T. (1980). Salmon propagation in Japan. In J.E. Thorpe (Ed.), *Salmon Ranching* (pp. 91–107). London: Academic Press. archive.org/details/salmonranching0000unse/page/446/mode/2up

Kraskura, K., Patterson, D. A., & Eliason, E. J. (2024). A review of adult salmon maximum swim performance. *Canadian Journal of Fisheries and Aquatic Sciences*, 81, 1174–1216. doi.org/10.1139/cjfas-2023-0246

Kuroda, H., Saito, T., Kaga, T., Takasuka, A., Kamimura, Y., Furuichi, S., & Nakanowatari, T. (2020). Unconventional sea surface temperature regime around Japan in the 2000s-2010s: Potential influences on major fisheries resources, *Frontiers in Marine Science*, 7, 574904. doi.org/10.3389/fmars.2020.574904

Langan, J. A., Cunningham, C. J., Watson, J. T., & McKinnell, S. (2024). Opening the black box: New insights into the role of temperature in the marine distributions of Pacific salmon. *Fish and Fisheries*. doi.org/10.1111/faf.12825

Larson, W. A., Seeb, J. E., Dann, T. H., Schindler, D. E., & Seeb, L. W. (2014). Signals of
heterogeneous selection at an MHC locus in geographically proximate ecotypes of
sockeye salmon. *Molecular Ecology*, 23, 5448–5461. doi.org/10.1111/mec.12949

Lewis, B., Grant, W. S., Brenner, R. E., & Hamazaki, T. (2015). Changes in size and age of
Chinook salmon *Oncorhynchus tshawytscha* returning to Alaska. *Plos One*, 10,
e0130184. doi.org/10.1371/journal.pone.0130184

Litzow, M. A., Ciannelli, L., Puerta, P., Wettstein, J. J., Rykaczewski, R. R., & Opiekun, M.
(2018). Non-stationary climate–salmon relationships in the Gulf of Alaska.
Proceedings of the Royal Society B, 285, 20181855. doi.org/10.1098/rspb.2018.1855

Losee, J. P., Kendall, N. W., & Dufault, A. (2019). Changing salmon: An analysis of body
mass, abundance, survival, and productivity trends across 45 years in Puget Sound.
Fish and Fisheries, 20, 934–951. doi.org/10.1111/faf.12385

MAFF (Ministry of Agriculture, Forestry and Fisheries) (2023). *Annual statistics of fisheries*
and aquaculture production. Tokyo: MAFF. (in Japanese)
www.maff.go.jp/j/tokei/kouhyou/kaimen_gyosei/index.html (accessed 11 January 2023)

Malick, M. J., Losee, J. P., Marston, G., Agha, M., Berejikian, B. A., Beckman, B. R., &
Cooper, M. (2023). Fecundity trends of Chinook salmon in the Pacific Northwest. *Fish*
and Fisheries, 24, 454–465. doi.org/10.1111/faf.12738

Manzer, J. I., Ishida, T., Peterson, A. E., & Hanavan, M. G. (1965). Salmon of the North
Pacific Ocean: Part V-offshore distribution of salmon. *International North Pacific*
Fisheries Commission Bulletin, 15, 452 p. www.npafc.org/wp-
content/uploads/Bulletin15.pdf

Masson-Delmotte, V., Zhai, P., Pörtner, H. O., Roberts, D., Skea, J., & Shukla, P. R. (2022).
Global Warming of 1.5 C: IPCC special report on impacts of global warming of 1.5 C
above pre-industrial levels in context of strengthening response to climate change,
sustainable development, and efforts to eradicate poverty. Cornwall: TJ Books
Limited. www.ipcc.ch/sr15/

Max Planck Institute (2023). *Climate Data Operators*. Hamburg: Max Planck Institute für
Meteorologie. code.mpimet.mpg.de/projects/cdo (accessed 14 June 2023).

Mayama, H., & Ishida, Y. (2003). Japanese Studies on the early ocean life of juvenile salmon.
North Pacific Anadromous Fish Commission Bulletin, 3, 41–67. www.npafc.org/wp-
content/uploads/Bulletins/Bulletin-3/Pages-41-68-JapanReview.pdf

Miyakoshi, Y., Urabe, H., Saneyoshi, H., Aoyama, T., Sakamoto, H., Ando, D., ... & Nagata,
M. (2012). The occurrence and run timing of naturally spawning chum salmon in
northern Japan. *Environmental Biology of Fishes*, 94, 197–206.
doi.org/10.1007/s10641-011-9872-5

Miyakoshi, Y., Nagata, M., Kitada, S., & Kaeriyama, M. (2013). Historical and current

hatchery programs and management of chum salmon in Hokkaido, northern Japan. *Reviews in Fisheries Science*, 21, 469–479. doi.org/10.1080/10641262.2013.836446

Mousseau, T. A., & Fox, C. W. (1998) The adaptive significance of maternal effects. *Trends in Ecology & Evolution*, 13, 403–407. doi.org/10.1016/S0169-5347(98)01472-4

Morita, K., & Fukuwaka, M. A. (2006). Does size matter most? The effect of growth history on probabilistic reaction norm for salmon maturation. *Evolution*, 60, 1516–1521. doi.org/10.1111/j.0014-3820.2006.tb01230.x

Morita, K., & Fukuwaka, M. A. (2007). Why age and size at maturity have changed in Pacific salmon. *Marine Ecology Progress Series*, 335, 289-294. doi.org/10.3354/meps335289

Myers, K. W., Klovach, N. V., Gritsenko, O. F., Urawa, S., & Royer, T. C. (2007). Stock-specific distributions of Asian and North American salmon in the open ocean, interannual changes, and oceanographic conditions. *North Pacific Anadromous Fish Commission Bulletin*, 4, 159–177. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-4/159-177Myers.pdf

Myers, K. W., Irvine, J., Logerwell, E., Urawa, S., Naydenko, S., Zavolokin, A., & Davis, N. (2016). Pacific salmon and steelhead: Life in a changing winter ocean. *North Pacific Anadromous Fish Commission Bulletin*, 6, 113–138. doi.org/10.23849/npafcb6/113.138

Nagasawa, T., & Azumaya, T. (2009). Distribution and CPUE trends of Pacific salmon, especially in sockeye salmon in the Bering Sea and adjacent waters from 1972 to mid 2000's. *North Pacific Anadromous Fish Commission Bulletin*, 5, 1–13. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-5/NPAFC_Bull_5_001-013Nagasawa.pdf

Nagata, M., Miyakoshi, Y., Ando, D., Fujiwara, M., Sawada, M., Shimada, H., & Asami, H. (2007). Influence of coastal seawater temperature on the distribution and growth of juvenile chum salmon, with recommendations for altered release strategies. *North Pacific Anadromous Fish Commission Bulletin*, 4, 223–235. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-4/223-235Nagatachum.pdf

Nagata, M., Miyakoshi, Y., Fujiwara, M., Kasugai, K., Ando, D., Torao, M., ... & Irvine, J. R. (2016). Adapting Hokkaido hatchery strategies to regional ocean conditions can improve chum salmon survival and reduce variability. *North Pacific Anadromous Fish Commission Bulletin*, 6, 73-85. doi.org/10.23849/npafcb6/73.85

Naish, K. A., Taylor III, J. E., Levin, P. S., Quinn, T. P., Winton, J. R., Huppert, D., & Hilborn, R. (2007). An evaluation of the effects of conservation and fishery enhancement hatcheries on wild populations of salmon. *Advances in Marine Biology*, 53, 61–194. doi.org/10.1016/S0065-2881(07)53002-6

Naydenko, S. V., & Temnykh, O. S. (2016). The survival of Pacific salmon in the North Pacific in the winter and spring. *Russian Journal of Marine Biology*, 42, 527–550.

doi.org/10.1134/S1063074016070087

Naydenko, S. V., Temnykh, O. S., & Figurkin, A. L. (2016). Is winter the critical period in the marine life history of Pacific salmon? *North Pacific Anadromous Fish Commission Bulletin*, 6, 139-152. doi.org/10.23849/npafcb6/139.152

Neave, F., Yonemori, T., & Bakkala, R. G. (1976). Distribution and origin of chum salmon in offshore waters of the North Pacific Ocean. *International North Pacific Fisheries Commission Bulletin*, 35. www.npafc.org/wp-content/uploads/Bulletin35.pdf

Nei, M., & Chesser, R. K. (1983). Estimation of fixation indices and gene diversities. Annals of Human Genetics, 47, 253–259. doi.org/10.1111/j.1469-1809.1983.tb00993.x

NOAA (National Oceanic and Atmospheric Administration, Physical Sciences Laboratory) (2023a). *NOAA Optimum Interpolation SST V2 high-resolution dataset (OISST V2)*. Boulder: NOAA. psl.noaa.gov/data/gridded/data.noaa.oisst.v2.highres.html (accessed 14 June 2023).

NOAA (National Oceanic and Atmospheric Administration, Physical Sciences Laboratory) (2023b). *Plot: COBE sea surface temperature*. Boulder: NOAA. psl.noaa.gov/mddb2/makePlot.html?variableID=1570 (accessed 18 March 2024).

NOAA (National Oceanic and Atmospheric Administration, Northwest Fisheries Science Center) (2024). *Ocean ecosystem indicators of Pacific salmon marine survival in the Northern California Current*. Seattle: NOAA. www.fisheries.noaa.gov/west-coast/science-data/ocean-ecosystem-indicators-pacific-salmon-marine-survival-northern (accessed 9 November 2024).

NPAFC (North Pacific Anadromous Fish Commission) (2023). *NPAFC statistics: Pacific salmonid catch and hatchery release data*. Vancouver: North Pacific Anadromous Fish Commission. www.npafc.org/statistics/ (accessed 7 August 2023).

Ohlberger, J., Ward, E. J., Brenner, R. E., Hunsicker, M. E., Haught, S. B., Finnoff, D., ... & Hauri, C. (2022). Non-stationary and interactive effects of climate and competition on pink salmon productivity. *Global Change Biology*, 28, 2026–2040. doi.org/10.1111/gcb.16049

Ohlberger, J., Cline, T. J., Schindler, D. E., & Lewis, B. (2023). Declines in body size of sockeye salmon associated with increased competition in the ocean. *Proceedings of the Royal Society B*, 290, 20222248. doi.org/10.1098/rspb.2022.2248

Okado, J., Koshino, Y., Kudo, H., & Watanuki, Y. (2020). Consumption of juvenile chum salmon by a seabird species during early sea life. *Fisheries Research*, 222, 105415. doi.org/10.1016/j.fishres.2019.105415

Okado, J., & Hasegawa, K. (2024). Exploring predators of Pacific salmon throughout their life history: The case of Japanese chum, pink, and masu salmon. *Reviews in Fish Biology and Fisheries*, 34, 895–917. doi.org/10.1007/s11160-024-09858-y

Oke, K. B., Cunningham, C. J., Westley, P. A. H., Baskett, M. L., Carlson, S. M., Clark, J., ... & Palkovacs, E. P. (2020). Recent declines in salmon body size impact ecosystems and fisheries. *Nature Communications*, 11, 4155. doi.org/10.1038/s41467-020-17726-z

PICES–NPFC (North Pacific Marine Science Organization, PICES,–North Pacific Fisheries Commission, NPFC) (2019). NPFC–PICES Framework for Enhanced Scientific Collaboration in the North Pacific. *North Pacific Fisheries Commission*, Document number NPFC-2019-SC04-WP02. www.npfc.int/npfc-pices-framework-enhanced-scientific-collaboration-north-pacific

Pauly, D., Hilborn, R., & Branch, T. A. (2013). Fisheries: Does catch reflect abundance?. *Nature*, 494(7437), 303-306. doi.org/10.1038/494303a

Powers, D. A., Lauerman, T., Crawford, D., & DiMichele, L. (1991). Genetic mechanisms for adapting to a changing environment. *Annual Review of Genetics*, 25, 629–660. doi.org/10.1146/annurev.ge.25.120191.003213

Punt, A. E. (2023). Those who fail to learn from history are condemned to repeat it: A perspective on current stock assessment good practices and the consequences of not following them. *Fisheries Research*, 261, 106642. doi.org/10.1016/j.fishres.2023.106642

Quinn, T. P. (2018). *The behavior and ecology of Pacific salmon and trout, 2nd edition.* Seattle: University of Washington Press. uwapress.uw.edu/book/9780295743332/the-behavior-and-ecology-of-pacific-salmon-and-trout/

Quinn, T. P., Vøllestad, L. A., Peterson, J., & Gallucci, V. (2004). Influences of freshwater and marine growth on the egg size-egg number tradeoff in coho and Chinook salmon. *Transactions of the American Fisheries Society*, 133, 55–65. doi.org/10.1577/T03-032

R Core Team (2024). R: A language and environment for statistical computing. *R Foundation for Statistical Computing.* https://www.R-project.org/

Radchenko, V. I., Temnykh, O. S., & Zavolokin, A. V. (2013). Review of studies on Asian juvenile Pacific salmon stocks, 2006–2012. *North Pacific Anadromous Fish Commission Technical Report*, 9, 1–10. https://www.npafc.org/wp-content/uploads/TechReport9.pdf

Radchenko, V. I., Beamish, R. J., Heard, W. R., and Temnykh, O.S. (2018) Ocean ecology of pink salmon. In R. J. Beamish (Ed.), *The Ocean Ecology of Pacific Salmon and Trout* (pp. 15–160). Bethesda: American Fisheries Society. https://doi.org/10.47886/9781934874455.ch1

Ricker, W. E. (1976) Review of the rate of growth and mortality of Pacific Salmon in salt water, and noncatch mortality caused by fishing. *Journal of the Fisheries Board of Canada*, 33. doi.org/10.1139/f76-191

Robins, C. R., Bailey, R.M., Bond, C. E., Brooker, J. R., Lacher, E. A., Lea, R. N., & Scott,

W. B. (1991). *World fishes important to North Americans, exclusive of species from the continental waters of the United States and Canada*. American Fisheries Society Special Publication, 21. Bethesda: American Fisheries Society. fisheries.org/bookstore/all-titles/special-publications/x51017pxm/

Rochet, M. J., & Trenkel, V. M. (2009). Why and how could indicators be used in an ecosystem approach to fisheries management?. In Beamish, R.J., & Rothschild, B.J. (Eds.), *The Future of Fisheries Science in North America* (pp. 153–167). Dordrecht: Springer. doi.org/10.1007/978-1-4020-9210-7_12

Ruggerone, G. T., Agler, B. A., & Nielsen, J. L. (2011). Evidence for competition at sea between Norton Sound chum salmon and Asian hatchery chum salmon. *Environmental Biology of Fishes*, 94, 149–163. doi.org/10.1007/s10641-011-9856-5

Ruggerone, G. T., & Irvine, J. R. (2018). Numbers and biomass of natural-and hatchery-origin pink salmon, chum salmon, and sockeye salmon in the North Pacific Ocean, 1925–2015. *Marine and Coastal Fisheries*, 10, 152–168. doi.org/10.1002/mcf2.10023

Ruggerone, G. T., Springer, A. M., van Vliet, G. B., Connors, B., Irvine, J. R., Shaul, L. D., ... & Atlas, W. I. (2023). From diatoms to killer whales: Impacts of pink salmon on North Pacific ecosystems. *Marine Ecology Progress Series*, 719, 1–40. doi.org/10.3354/meps14402

Saito, T. (2022). Optimum timing and size at release of chum salmon: Improving survival by modifying hatchery practices. *Fisheries Oceanography*, 31, 416–428. https://doi.org/10.1111/fog.12585

Saito, T., & Nagasawa, K. (2009). Regional synchrony in return rates of chum salmon (*Oncorhynchus keta*) in Japan in relation to coastal temperature and size at release. *Fisheries Research*, 95, 14–27. doi.org/10.1016/j.fishres.2008.07.004

Saito, T., Honda, K., Sasaki, K., Watanabe, K., Suzuki, K., Hirabayashi, Y., ... & Sato, S. (2020). Stock composition of adult chum salmon *Oncorhynchus keta* caught in a setnet fishery estimated using genetic identification, scale patterns, and otolith thermal marking. Fisheries Science, 86, 271–286. doi.org/10.1007/s12562-020-01401-9

Saitou, N., & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4, 406–425. doi.org/10.1093/oxfordjournals.molbev.a040454

Salo, E.O. (1991). Life history of chum salmon (*Oncorhynchus keta*). In C. Groot, & L. Margolis (Eds.) *Pacific Salmon Life Histories* (pp. 231–310). Vancouver: University of British Columbia Press. www.ubcpress.ca/pacific-salmon-life-histories

Sano, S. (1966). Chum Salmon in the Far East. *International North Pacific Fisheries Commission Bulletin*, 18-3, 41–57. www.npafc.org/wp-content/uploads/Bulletin18-3.pdf

Sano, S. (1967). Chum salmon in the Far East. *International North Pacific Fisheries Commission Bulletin*, 23–3: 23-41. www.npafc.org/wp-content/uploads/Bulletin23-3.pdf

Shedd, K. R., Lescak, E. A., Habicht, C., Knudsen, E. E., Dann, T. H., Hoyt, H. A., … & Templin, W. D. (2022). Reduced relative fitness in hatchery-origin pink salmon in two streams in Prince William Sound, Alaska. *Evolutionary Applications*, 15, 429–446. doi.org/10.1111/eva.13356

Shepard, M. P., Hartt, A. C., & Yonemori, T. (1968). Chum salmon in offshore waters. *International North Pacific Fisheries Commission Bulletin*, 25. www.npafc.org/wp-content/uploads/Bulletin-25.pdf

Shubin, A. O., & Akinicheva, E. G. (2016). Origin of juvenile chum salmon *Oncorhynchus keta* (Salmonidae) in the Sea of Okhotsk coastal waters off south Sakhalin. *Journal of Ichthyology*, 56, 728–737. doi.org/10.1134/S0032945216050131

Shuntov, V. P., Temnykh, O. S., & Naydenko, S. V. (2019). More on the factors that limit the abundance of Pacific salmon (*Oncorhynchus* spp., family Salmonidae) during the ocean phase of their life history. *Russian Journal of Marine Biology*, 45, 511–524. doi.org/10.1134/S106307401907006X

Somero, G. N. (2022). The Goldilocks principle: A unifying perspective on biochemical adaptation to abiotic stressors in the sea. *Annual Review of Marine Science*, 14, 1–23. doi.org/10.1146/annurev-marine-022521-102228

Somov, A., Farley, E. V., Yasumiishi, E. M., & McPhee, V. (2024) Comparison of juvenile Pacific salmon abundance, distribution, and body condition between western and eastern Bering Seaming spatiotemporal models. *Fisheries Research*, 278:107086. doi.org/10.1016/j.fishres.2024.107086

Springer, A. M., & van Vliet, G. B. (2014). Climate change, pink salmon, and the nexus between bottom-up and top-down forcing in the subarctic Pacific Ocean and Bering Sea. *Proceedings of the National Academy of Sciences*, 111, E1880–E1888. doi.org/10.1073/pnas.1319089111

Sugiura, N. (1978). Further analysis of the data by Akaike's information criterion and the finite corrections. *Communications in Statistics - Theory and Methods*, 7, 13–26. doi.org/10.1080/03610927808827599

Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: Molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution*, 38, 3022–3027. doi.org/10.1093/molbev/msab120

Thorn, M. W., Dick, M. F., Oviedo, L., Guglielmo, C. G., & Morbey, Y. E. (2019). Transgenerational effects of egg nutrients on the early development of Chinook salmon (*Oncorhynchus tshawytscha*) across a thermal gradient. *Canadian Journal of Fisheries*

and Aquatic Sciences, 76, 1253-1262. doi.org/10.1139/cjfas-2018-0013

Thorson, J. T., Maunder, M. N., & Punt, E. (2020). The development of spatio-temporal

models of fishery catch-per-unit-effort data to derive indices of relative abundance.

Fisheries Research, 230, 105611. doi.org/10.1016/j.fishres.2020.105611

Tian, Y., Kidokoro, H., Watanabe, T., Igeta, Y., Sakaji, H., & Ino, S. (2012). Response of

yellowtail, *Seriola quinqueradiata*, a key large predatory fish in the Japan Sea, to sea

water temperature over the last century and potential effects of global warming. *Journal*

of Marine Systems, 91, 1-10. doi.org/10.1016/j.jmarsys.2011.09.002

Urawa, S., & Bugaev, A. V. (2021). Survival of Japanese chum salmon during early ocean life

in 2011–2017. *North Pacific Anadromous Fish Commission Technical Report*, 17, 60–

62. doi.org/10.23849/npafctr17

Urawa, S., Seki, J., Kawana, M., Saito, T., Crane, P. A., Seeb, L. W., Fukuwaka, M., &

Akinicheva, E. (2006). Origins of juvenile chum salmon caught in the southwestern

Okhotsk Sea during the fall of 2000. *Bulletin of National Salmon Resources Center*, 8,

9–16. salmon.fra.affrc.go.jp/kankobutu/bulletin/BNSRC08-2.pdf

Urawa, S., Sato, S., Crane, P. A., Agler, B., Josephson, R., & Azumaya, T. (2009). Stock-

specific ocean distribution and migration of chum salmon in the Bering Sea and North

Pacific Ocean. *North Pacific Anadromous Fish Commission Bulletin*, 5:131–146.

www.npafc.org/wp-content/uploads/Bulletins/Bulletin-5/NPAFC_Bull_5_131-

146Urawa.pdf

Urawa, S., Beacham, T., Sato, S., Kaga, T., Agler, B. A., Josephson, R., & Fukuwaka, M.

(2016). Stock-specific abundance of chum salmon in the central Gulf of Alaska

during winter. *North Pacific Anadromous Fish Commission Bulletin*, 6, 153–160.

doi.org/10.23849/npafcb6/153.160

Urawa, S., Beacham, T. D., Fukuwaka, M., & Kaeriyama, M. (2018). Ocean ecology of chum

salmon. In R. J. Beamish (Ed.), *The Ocean Ecology of Pacific Salmon and Trout* (pp.

161–317). Bethesda: American Fisheries Society.

doi.org/10.47886/9781934874455.ch3

Urawa, S., Beacham, T., Sutherland, B., & Sato, S. (2022). Winter distribution of chum

salmon in the Gulf of Alaska: A review. *North Pacific Anadromous Fish Commission*

Technical Report, 18, 83–87. doi.org/10.23849/npafctr18

Walters, C. J., & Hilborn, R. (1978). Ecological optimization and adaptive management.

Annual review of Ecology and Systematics, 9, 157–188. www.jstor.org/stable/2096747

Welch, D. W., Ishida, Y., & Nagasawa, K. (1998). Thermal limits and ocean migrations of

sockeye salmon (*Oncorhynchus nerka*): Long-term consequences of global warming.

Canadian Journal of Fisheries and Aquatic Sciences, 55, 937–948.

doi.org/10.1139/f98-023

Wilkinson, L., & Task Force on Statistical Inference, American Psychological Association, Science Directorate (1999). Statistical methods in psychology journals: Guidelines and explanations. *American Psychologist*, 54, 594–604. doi.org/10.1037/0003-066X.54.8.594

Wood, S. N. (2004). Stable and efficient multiple smoothing parameter estimation for generalized additive models. *Journal of the American Statistical Association*, 99, 673–686. doi.org/10.1198/016214504000000980

Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society B*, 73, 3–36. doi.org/10.1111/j.1467-9868.2010.00749.x

Wood, S. N. (2017). *Generalized additive models*. New York: Chapman and Hall/CRC. doi.org/10.1201/9781315370279

Yagisawa, Y. (1970). Salmon hatchery enhancement program on Etorofu Island, Fish and eggs, *Salmon Management Center Technical Report*, 133, 35–79. (in Japanese) http://salmon.fra.affrc.go.jp/kankobutu/tech_repo/fe04.htm

Yasumiishi, E. M., Farley Jr, E. V., Maselko, J., Aydin, K. Y., Kearney, K. A., Hermann, A. J., ... & Strasburger, W. W. (2020). Differential north–south response of juvenile Chinook salmon (*Oncorhynchus tshawytscha*) marine growth to ecosystem change in the eastern Bering Sea, 1974-2010. *ICES Journal of Marine Science*, 77, 216–229. doi.org/10.1093/icesjms/fsz166

Yonemori, T. (1975). A trial analysis of the results obtained from tagging on chum salmon origination from Hokkaido. *International North Pacific Fisheries Commission Bulletin*, 32, 130–160. www.npafc.org/wp-content/uploads/Bulletin-32.pdf

Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid common statistical problems. *Methods in Ecology and Evolution*, 1, 3–14. doi.org/10.1111/j.2041-210X.2009.00001.x

1580

1581

FIGURE 1 Japanese chum salmon management regions, coastal migration routes, and high

seas distribution areas. (a) Seven management regions (delineated by black bars) and

hatcheries (black dots) in Japan. Migration paths of hatchery-released fry were modified from

Saito and Nagasawa (2009, Figure 1). Schematic sea surface currents were redrawn from

Kuroda et al. (2020, Figure 1). (b) principal areas of high seas distribution (boxes), month(s),

age group(s), and seasonal direction of wintering, feeding and homing migrations (arrows),

numbered areas are: 1.Southern Sea of Okhotsk (45.2°N-46.5°N, 142.5°E-148.5°E), 2.

Western North Pacific (42°N-45°N, 158°E-165°E), 3. Central Bering Sea (53°N-58°N,

175°E-170°W), 4. Central Gulf of Alaska (48°N-56°N, 140°W-150°W). Migration routes of

Japanese chum salmon estimated from long-term field surveys, modified from Myers et al.

(2007, Figure 1) and Urawa et al. (2018, Figure 37).

1593

1594 **FIGURE 2** Number of hatchery chum salmon fry released (red line) and adult returns (coastal
1595 catch plus hatchery broodstock catch; black line) in seven management regions in Japan (1970-
1596 2023; Figure 1a).

1597

FIGURE 3 Percentage of age 2 - 4 Japanese chum salmon in the total number of adult returns for 1998-2019 hatchery releases generally increased in all regions after around 2008, as shown by the nonlinear and linear year effect from GAMs. The trend was significant in all regions, except Hokkaido Nemuro and Honshu Sea of Japan (Table S1).

1602

1603 **FIGURE 4** Neighbor-joining tree based on correlation coefficients of chum salmon adult return

1604 rates between pairs of seven management regions in Japan, 1998-2019.

1605

FIGURE 5 Sea surface temperature (SST) isotherms in recent decades (1990-2022) showed a
northward shift and longitudinal contraction of chum salmon summer (July) thermal habitat in
the North Pacific Ocean, while winter (January) habitat was relatively stable. Blue, green and
orange show the range of Japanese chum salmon thermal limits (4-13℃): optimal growth and
feeding habitat (8-12℃), swimming and feeding habitat (5-13℃), and overwintering habitat
(4-6℃) (Kaeriyama et al., 2012; Urawa et al., 2018). Red (>13℃) and purple (<4℃) indicate
critical habitat, lost to ocean production of Japanese chum salmon.

1613

1614 **FIGURE 6** Northward shifts of two marine piscivorous fishes in Japan. (a) Principal
1615 component analysis biplot based on catch statistics of 56 marine species (1980 - 2020) showed
1616 increasing catches of yellowtail and Japanese Spanish mackerel. (b) Two-decadal shifts in
1617 commercial catches of chum salmon and two piscivores.

1618

FIGURE 7 Egg size of age–4 chum salmon has generally decreased in all regions, as shown
by the nonlinear and linear year effect from GAMs (1998-2019). The trend was significant in
all regions, except Honshu Sea of Japan. Open circles show observed values. The black lines
show predicted values with shaded areas of 2×standard errors.

1623

FIGURE 8 Standardized effects of hypothesized explanatory variables on Japanese chum salmon marine survival (adult return rates across regions). Regression coefficients of best regression models with 95% confidence intervals in four coastal regions in Japan, estimated using standardized explanatory variables.

FIGURE 9 Predicted and observed chum salmon marine survival (adult return rates) by management region in Japan. Predicted return rates (grey line), observed return rates (black dots), and 95% prediction intervals (black lines). Note that scales on return rate axis differ by region.

TABLE 1 Three subset models of stressors related to chum salmon marine survival (adult return rate) of fry released in year t (1998-2019) and explanatory variables considered in a regression model and hypothesized directional relationship. Sea surface temperature (SST); Western North Pacific (WN Pacific); Gulf of Alaska (GOA).

Subset model	Explanatory variable (year)	Age[†]	Spatial scale[‡]	Hypothesized relationship to chum salmon survival (source)
Hatchery carryovers	Parental relative fecundity (t -1)	0	Japan coast	Survival related (negative) to higher fecundity (Hasegawa et al., 2021a)
	Egg size (t)	0	Japan coast	Survival related (positive) to larger eggs (Hasegawa et al., 2021a)
	Fry size at release (t)	0	Japan coast	Survival related (positive) to size at release (Hasegawa et al., 2021b)
Ocean conditions	SST at release (t)	0	Japan coast	Survival related (negative) to SST (Nagata et al., 2007)
	Okhotsk coast Jul SST (t)	0	Okhotsk	1st summer survival related (negative) to SST (Nagata et al., 2007)
	WN Pacific Jan-Apr SST (t+1)	1	WN Pacific	1st winter survival related (positive) to SST (this study)
	Bering Sea Aug-Sep SST (t+1)	1	Bering Sea	2nd summer survival related (negative) to SST (this study)
	GOA Jan-Apr SST (t+2)	2	Gulf of Alaska	2nd winter survival related (positive) to SST (this study).
Predators and competitors	Yellowtail catch (t)	0	Japan coast	1st year survival related (negative) to predation (Beamish, 2018)
	Russia chum catch (t+3)	0-4	Okhotsk, North Pacific	1st - 3rd year survival related (negative) to Russian chum (this study).
	Russia pink catch (t+1)	0-1	Okhotsk, WN Pacific	1st year survival related (negative) to Russian pink (this study)
	Russia pink catch (t+2)	1-2	Bering Sea	2nd year survival related (negative) to Russian pink (Ruggerone et al., 2023
	Russia pink catch (t+3)	2-3	Gulf of Alaska, Bering Sea	3rd year survival related (negative) to Russian pink (Ruggerone et al., 2023)
	US & Canada chum catch (t+3)	1-4	North Pacific	2nd - 3rd year survival related (negative) to US & Canadia chum (this study)
	US &Canada pink catch (t+1)	0-1	Okhotsk, WN Pacific	1st year survival related (negative) to US & Canada pink (this study)
	US & Canada pink catch (t+2)	1-2	Bering Sea	2nd year survival related (negative) to US & Canada pink (this study)
	US and Canada pink catch (t+3)	2-3	Gulf of Alaska, Bering Sea	3rd year survival related (negative) to US & Canada pink (this study)

[†] Age of Japanese chum salmon released in year t (=1998,…,2019). [‡] See Japanese salmon migration route (Fig. 1).

TABLE 2 Best regression models of Japanese chum salmon survival in four coastal regions in Japan and regression coefficients with their p-values in parentheses. Bold indicates significant relationship ($\alpha = 0.05$). t = ocean entry year of chum salmon fry. SST = sea surface temperature. See Figure 1a for management region locations.

Subset model	Explanatory variable (year)	Management region			
		Hokkaido Okhotsk	Hokkaido & Honshu Pacific	Hokkaido Sea of Japan	Honshu Sea of Japan
Hatchery carryovers	Relative fecundity[†][‡] (t-1)	–	**−0.931 (0.0009)**	–	–
	Egg size[†] (t-1)	–	–	–	–
	Fry size at release (t)	–	**0.486 (0.0009)**	–	–
Ocean conditions	SST at release (t)	–	–		**-0.483 (0.0033)**
	Okhotsk coast Jul SST (t)	–	–	**-0.315 (0.0042)**	**-0.138 (0.0452)**
	WN Pacific Jan-Apr SST (t+1)	–	–	–	0.259 (0.0647)
	Bering Sea Aug-Sep SST (t+1)	–	–	**-0.383 (0.0153)**	–
	GOA Jan-Apr SST (t+2)	–	–	**0.405 (0.0051)**	–
Predators and competitors	Yellowtail catch in year (t)	**-0.093 (0.0001)**	–	–	–
	Russia chum catch (t+3)	–	–	-0.151 (0.0504)	–
	Russia pink catch (t+1)	–	–	–	–
	Russia pink catch (t+2)	-0.127 (0.0559)	**-0.311 (0.0025)**	–	-0.139 (0.0605)
	Russia pink catch (t+3)	–	**-0.225 (0.0138)**	–	–
	USA & Canada chum catch (t+3)				
	USA & Canada pink catch (t+1)	–	–	–	–
	USA & Canada pink catch (t+2)	–	–	–	–
	USA & Canada pink catch (t+3)	–	–	–	–
	Adjusted R^2 (p-value)	**0.58**	**0.79**	**0.54**	**0.57**

The three subset models were fitted to log return rates for 1998-2019 releases and variable selection was performed using the Akaike Information Criterion corrected for small sample sizes (AICc). Multicollinearity of explanatory variables was checked using the variance inflation factor (VIF), and those with VIF < 3.0 were used for variable selection.

[†] Egg sizes and relative fecundity of age 4 and age 5 fish were examined. [‡] Age 4 had a smaller AICc (30.0) than that of age 5 (38.2).