Hatcheries to high seas: climate change connections to salmon marine survival

Shuichi Kitada[1*], Katherine W. Myers[2], and Hirohisa Kishino[3,4]

Short running title: Climate connections to salmon survival

[1] Tokyo University of Marine Science and Technology, Tokyo, Japan
[2] School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98195, USA.
[3] Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, Japan
[4] Research and Development Initiative, Chuo University, Tokyo, Japan

*correspondence

Abstract

We investigated variations in the marine survival of Japanese hatchery chum salmon (*Oncorhynchus keta*) during 25 years of climate change (1998–2023). Japan is the world's largest producer of hatchery salmon, and is located near the global southern distribution limit of chum salmon. Our goal was to identify local and context-specific metrics related to the observed coastwide decline in salmon marine survival over the past two decades. We hypothesized multiple metrics in three categories of stressors: hatchery carryovers, ocean conditions, and predators and competitors. The hatchery carryovers are stressors related to hatchery rearing that affect survival at a different life stage. We collected, processed, and collated large publicly available datasets into a comprehensive open-access database encompassing the life cycle of Japanese chum salmon, from eggs to adult spawners. Multivariate regression models showed associations between stressors and adult salmon return rate (marine survival) varied by coastal management region, salmon life stage, and seasonal high-seas distribution area. In the early marine life history stage, parental egg size and fry size-at-release had the largest positive model effects on marine survival. The sea surface temperature (SST) at time of fry release and a predator of fry had significant negative effects. In the offshore and high-seas life stages, summer SST had negative effects, while winter SST had positive effects. Russian chum and/or pink salmon abundance had negative effects, while no effect was found for North American pink and chum salmon abundance. Generalized additive models (GAMs) identified a nationwide decline in egg size and fry size-at-release. Our study highlights the need for an experimental approach to hatchery practices, including monitoring and analyses with updated information, leading to effective management decisions and policies for future sustainability and conservation of salmon resources.

KEYWORDS competition, egg size, fry size-at-release, Japanese hatchery chum salmon, predation

1 | INTRODUCTION

Climate change influences the survival and abundance of many economically important marine and anadromous fish species, including Pacific salmon (*Oncorhynchus* spp.) (Springer & van Vliet, 2014; Cheung & Frölicher, 2020; Cheung et al., 2021; Crozier et al., 2021; Crozier & Siegel, 2023). Pacific salmon populations often consist of hundreds of subpopulations with diverse life history traits and local adaptations to environmental variation (Groot & Margolis, 1991; Hilborn et al., 2003; Larson et al., 2014; Quinn, 2018). Pacific salmon species are distributed in coastal marine waters and throughout the high seas (international waters) of the North Pacific Ocean and Bering Sea, where Asian and North American populations intermingle during seasonal feeding migrations (Myers et al., 2007; Beamish, 2018; Quinn, 2018). Their wide geographic range, long ocean migrations, and complex life histories expose salmon populations to multiple climate-driven stressors at both local and North Pacific-wide scales (Cline et al., 2019; Crozier et al., 2021). Global warming of 1.5°C above pre-industrial levels is projected to shift the ranges of many marine species to higher latitudes and increase damage to marine ecosystems, such as rising temperatures, ocean acidification, heatwaves, decreases in sea ice, habitat loss, disruptions in food webs, and declining fish populations (Dahlke et al., 2018; Frölicher et al., 2018; Masson-Delmotte, 2022).

The effects of climate change, ocean acidification, and large-scale hatchery releases on the abundance and productivity of Pacific salmon populations have received increasing attention in recent years (Lewis et al., 2015; Jeffrey et al., 2017; Cunningham et al., 2018; Cline et al., 2019; Losee et al., 2019; Connors et al., 2020, 2024; Oke et al., 2020; Ohlberger et al., 2022, 2023; Malick et al., 2023). Declines in body size, primarily due to a shift in age structure towards younger ages at maturity, were found in chum, Chinook (*O. tshawytscha*), coho (*O. kisutch*), and sockeye (*O. nerka*) salmon from large-scale analyses across Alaska and were linked to climate change and increased inter- and intra-specific competition due to an overabundance of salmon in the high seas (Lewis et al., 2015; Oke et al., 2020; Ohlberger et al., 2023). Pink salmon (*O. gorbuscha*) abundance increased since the mid-1970s to record highs in the 2000s, supported by climate change effects on the ocean environment and production processes, and may have affected the abundance of salmon, other species, and their ecosystems (Springer & van Vliet, 2014; Connors et al., 2020; Ruggerone et al., 2023).

Pacific salmon have a long history of artificial propagation beginning in the mid-1800s, and their management comprises the largest set of hatchery release programs (Naish et al., 2007). Chum salmon hatcheries in Japan constitute the world's largest salmon hatchery release program (Amoroso et al., 2017; Kitada, 2018), continuing for over 130 years (Miyakoshi et al., 2013). Hatchery releases of fry chum in Japan increased rapidly in the 1970s, but gradually declined from a peak of 2 billion per year in 1983 to 1.3 billion per year in 2021 because of poor returns of adult spawners. Adult returns of Japanese hatchery chum increased rapidly since the 1970s, but declined sharply after the late 1990s from the historical maximum of 88 million in 1996 to the minimum of 19 million in 2021. Previous studies have reported higher spring SSTs in the coastal zone contributed to better survival of juvenile chum through improved feeding, growth, and swimming ability (Saito & Nagasawa, 2009; Kuroda et al., 2020; Honda et al., 2021). Conversely, shorter inshore stays associated with warming SSTs could lead to lower juvenile survival and insufficient growth (Kaeriyama, 2022; Honda et al., 2025). Although it is generally accepted that a warming climate is affecting marine survival of salmon, the mechanisms underlying recent declines in Japanese hatchery chum salmon are largely unknown. The need for adaptive management is advocated for the long-term sustainability of

90   chum salmon (Kaeriyama, 2008, 2022; Kaeriyama et al., 2012).
91
The development of adaptive management plans to address climate-change impacts on fish,
fisheries, aquaculture, and associated habitats and ecosystems requires knowledge of "local and
context-specific indicators associated with climate stressors" (FAO, 2022). Adaptive
management is the design of management actions as experiments with feedback through
monitoring, enabling an iterative process of learning and adapting to uncertainties and risks
(Walters & Hilborn, 1978; Hilborn & Sibert, 1988; Bahri et al., 2021; Bryndum-Buchholz et
al., 2021). A comprehensive review of literature addressed strengths and weaknesses of climate-
change research by life stage and showed the complexity of whole life cycle approaches to
adaptive management strategies for anadromous salmonids (Crozier & Siegel, 2023). Here, we
investigated associations between adult return rates (an index of marine survival) of Japanese
hatchery chum salmon by life stage and a suite of hypothesized climate-change stressors. In our
study, we expected survival of all life stages of chum salmon to be associated with climate
change impacts on coastal and open ocean ecosystems. Our primary objective was to identify
potential stressors along known coastal and high-seas migration routes of Japanese chum
salmon for use in regional adaptive management of salmon resources.
107
2 | MATERIALS AND METHODS
2.1 Ecological setting, study design, and methods overview
The ecological setting of our study encompasses the known geographic distribution and
seasonal migration patterns of Japanese chum salmon in the North Pacific Ocean and adjacent
seas (Figure 1; reviewed by Fisheries Agency of Japan, 1955; Sano, 1966, 1967; Shepard et al.,
1968; Neave et al., 1976; Salo, 1991; Urawa et al., 2018). Chum salmon fry are released from
more than 260 coastal hatcheries in northern Honshu and Hokkaido in winter and spring (Kitada,
2020, Figure 1a) and migrate rapidly through river estuaries (Kobayashi, 1980; Kaeriyama,
1989; Saito & Nagasawa, 2009). After ocean entry, fry migrate northward in shallow coastal
waters before moving offshore to summer feeding grounds in the southern Sea of Okhotsk
(Figure 1b; Fukuwaka & Suzuki, 1998; Urawa et al., 2006; Kasugai et al., 2016; Shubin &
Akinicheva, 2016; Honda et al., 2025). Japanese chum salmon spend their first ocean winter
(age-1) in the western North Pacific Ocean and in spring move northeastward to summer
feeding grounds in the Bering Sea (Figure 1b; Mayama & Ishida, 2003; Urawa et al., 2018). At
subsequent life stages (ages 2–4; immature and maturing), they make seasonal migrations
between the Gulf of Alaska in winter and the Bering Sea in summer (Figure 1b; Beacham et al.,
2009; Urawa et al., 2009, 2016). In spring, the northward migration of maturing chum salmon
precedes that of immature fish (Neave et al., 1976; Nagasawa & Azumaya, 2009). In late
summer and fall, maturing fish return to Japan from the Bering Sea through the western North
Pacific (Figure 1b; Yonemori, 1975; Urawa et al., 2018).
128
A brief overview of our study design and methodology for evaluating relationships between
salmon marine survival and hypothesized stressors is provided below (see details in sections
2.2–2.8). Our study design is observational (a retrospective case study). We identified multiple
a priori hypotheses about potential stressors driving marine survival of Japanese hatchery chum
salmon at specific life stages and locations along their coastal and high-seas migration routes
(Table 1). We grouped hypothesized stressors into three broad categories: hatchery carryovers,
ocean conditions (SST), and predators and competitors. Hatchery carryovers are stressors
related to hatchery rearing that affect survival at a different life stage. We assembled available
data sets for use as measures of hypothesized stressors in each category. We summarized time

series data on reproductive and biological traits of hatchery salmon and adult return rates by coastal management region (Figure 1a). We created isotherm maps to illustrate decadal-scale trends in SST throughout the known ocean range of Japanese chum salmon. We estimated SSTs at ocean entry and along coastal and high-seas migratory routes (Figure 1b). We developed measures of potential coastal marine fish predators and high-seas salmon competitors of Japanese chum salmon. Investigation of coastal mammal and bird predators and high-seas predators was beyond the scope of our analysis due to inadequate time-series data covering the period of our study. We estimated annual adult return rates (marine survival), stratified by salmon age group and year. We used multivariate regression models to evaluate relationships between the marine survival of salmon and hypothesized categories of stressors by coastal management region in Japan and seasonal high-seas habitats in the Sea of Okhotsk, western North Pacific, central Bering Sea, and Gulf of Alaska. Readers are cautioned that our statistical modeling results are correlational, not causal.

2.2 Data sources, sample periods, sample sizes
We assembled available time-series datasets during 25 years of climate change (1998–2023) from public open-data websites. We assembled longer data time series to aid in interpreting results.

We obtained annual (1970–2023) numbers of adult salmon catches (coastal fishery and hatchery broodstock) and hatchery chum salmon fry releases from the Hokkaido Salmon Enhancement Association (HSEA, 2024), Fisheries Research and Education Agency (FRA, 2024), and the Aomori Prefectural Government (APG, 2024). FRA (2023) biological and reproductive trait data included time series of fry size-at-release and release month. We selected hatchery release events with fry body weight (BW) and fork length (FL) data (1998–2020, Hokkaido: 184 rivers, 37,886 release events, $n = 21,332,093$ fish; Honshu: 141 rivers, 19,605 release events, $n = 9,588,413$ fish). The data also included adult age composition at return (1997–2020, Hokkaido: 52 rivers; Honshu: 102 rivers; 1,902 surveys, $n = 1,210,717$ fish). Adult salmon trait data included egg weight, fecundity, FL, age-at-return, and gonad weight (1997–2020, Hokkaido: 10 rivers; Honshu: 14 rivers; $n = 37,533$ fish), and adult age-at-return (counts of annular marks on fish scales), including six age groups: age-2 ($n = 25$), age-3 ($n = 2,892$), age-4 ($n = 20,643$), age-5 ($n = 13,118$), age-6 ($n = 846$), and age-7 ($n = 9$). We used trait data such as FL, egg weight, and fecundity of age-4 and age-5 fish for our analysis.

We obtained annual commercial catch data for all marine species in Japan (1980–2020) from the Japan Ministry of Agriculture, Forestry and Fisheries (MAFF, 2023). Catch statistics (1998–2022) for Canada, Russia, and USA pink and chum salmon were available from the North Pacific Anadromous Fish Commission (NPAFC, 2023). For SST analyses, we downloaded Japan coastal 0.25°×0.25° daily mean SST data (1982–2020) from Hokkaido (18 areas) and Honshu (12 areas) from the Japan Meteorological Agency (JMA, 2022) and global 0.25°×0.25° monthly mean SST data (1982 to 2022) from the NOAA Optimum Interpolation (OI) SST V2 high-resolution dataset (NOAA, 2023a).

2.3 Catch, release, and reproductive traits
We summarized Japanese chum salmon catch, release, and reproductive traits data by seven management regions in northern Honshu and Hokkaido, Japan (Figure 1a). To estimate the distribution of release times in each region, we counted the number of release events by month. We also calculated the mean values of size-at-release, biological traits, and age composition by

management region. Since the number of individuals released (n_k) varied among release events
($k=1,\ldots,K$), we calculated the weighted mean size-at-release for BW (g) (1998–2020) as $\overline{BW} = $
$(\sum_{k=1}^{K} BW_k \, n_k)/\sum_{k=1}^{K} n_k$. Here, the suffix for a region was omitted for simplicity. We
calculated the mean values of the traits and the relative fecundity (eggs/$FL(\text{cm})^3 \times 10^3$) of the
age-4 and age-5 fish from 1997 to 2020. The weighted mean proportion of age-a fish was
calculated as $\hat{p}_a = (\sum_{h=1}^{H} p_{ah} \, n_h)/\sum_{h=1}^{H} n_h$, where h refers to each survey, p_{ah} is the
proportion of age-a fish in survey h, n_h is the sample size (individuals), and a refers to age at
return ($a = 2,\ldots,8$).
194

2.4 Adult return rates (marine survival index)
We assumed all adult chum salmon returning to Japan were hatchery reared (see Discussion,
section 4.1), and adult return rates are survival rates from release to return. In this study, we
defined the adult return rate as an index of the marine survival of chum salmon (James et al.
2023). Hatchery programs have been managed in five management regions in Hokkaido and by
prefecture in Honshu, and previous studies evaluated return rates by management region (Saito
& Nagasawa, 2009; Miyakoshi et al., 2013; Kitada, 2014). The management regions coincide
with the genetic structure of the population (Beacham et al., 2008). We estimated return rates
for the current hatchery management regions (Figure 1a). We estimated the total number of fish
returning from releases in year t (1998–2019) (Y_t, coastal commercial catch plus river catch for
hatchery broodstock) by

206
$$\hat{Y}_t = \sum_{a=A_{min}}^{A_{max}} y_{t+a-1} \, \hat{p}_{a,t+a-1} = \sum_{a=A_{min}}^{A_{max}} \hat{Y}_{a,t}$$

where y_{t+a-1} is the observed number of fish returned in year $t + a - 1$ from fry released in
year t, and $\hat{p}_a$ is the estimated proportion of age-a fish among the fish returned in year $t +$
$a - 1$. We estimated the return rate of fry released in year t by $\hat{R}_t = \hat{Y}_t/X_t$, where X_t is the
number of hatchery-reared fry released in year t. We estimated the age-specific return rate as
$\hat{R}_{a,t} = \hat{Y}_{a,t}/X_t$. We summed the number of fish returned from ages 2–5 ($A_{min} = 2$, $A_{max} = 5$).
Most returns were ages 2–5 fish, and few returns were age-6. There were no age-7 and age-8
fish in returns to any region except Hokkaido Okhotsk, where returns of these age groups were
very low. We estimated the age composition of returning adult chum salmon from available
(1997–2020) data (Figure S1). We estimated the age composition in 2021–2023, where no data
were available, as a moving average of the previous three years to account for the recent trend
of increasing numbers of young fish. Based on age composition, we calculated the number of
returning fish (ages 2–5) for each release year. Then, we estimated return rates and age-specific
return rates from releases in 1998–2019. We measured distances between return rates in the
management regions by $1 - r_{uv}$ and illustrated distances by a neighbor-joining (NJ) tree
(Saitou & Nei, 1987) using the "MEGA11" package (Tamura et al., 2021), where r_{uv} is the
correlation coefficient between pairs of return rate time series in regions u and v.
223

To assess and visualize continuous changes in log-transformed return rates and covariates over
the years, we fit GAMs to the data using the 'gam' function in the "mgcv" package (Wood,
2017) in R (R Core Team, 2024), assuming normal errors. Year was included as a nonlinear
smoothed term (explanatory variable) because these covariates varied across years. The trend
of the traits over the years is captured by the non-linear and/or linear year effects (Wood, 2004,
2011).
230

2.5 Long-term changes in chum salmon thermal habitat

To understand long-term changes in SST in the chum salmon habitats covered by our regression model (see section 2.8), we mapped monthly mean 1°×1° SST isoclines in the North Pacific Ocean for 1990, 2000, 2010, and 2022, using the Plot COBE Sea Surface Temperature system (NOAA, 2023b), based on the COBE2 SST dataset from JMA. The 1°×1° monthly mean SST isoclines around Japan for 2000, 2010, and 2022 were generated similarly. We mapped the available temperature range of chum salmon (4–13°C), based on previous predictions: optimal growth and feeding (8–12°C), available habitat for feeding (5–13°C), and overwintering temperature range (4–6°C) (Kaeriyama et al., 2012; Urawa et al., 2018). Langan et al. (2024) found that chum salmon in the North Pacific Ocean were frequently caught at SSTs below 4°C in spring and summer. Their analysis was based on unknown population mixtures of chum salmon in historical high-seas research vessel catches. It is quite possible that southern populations such as Japanese chum salmon are less cold tolerant than northern populations.

2.6 Japanese chum salmon migration route and SST

We calculated monthly mean SSTs for 30 coastal areas in Japan (JMA, 2022). From these, we calculated monthly mean SSTs for the seven management regions (Figure 1a), and for release periods of chum salmon fry (Figure S2): Hokkaido Okhotsk (Apr–Jun), Hokkaido Nemuro (Mar–Jun), Hokkaido Pacific East (Mar–Jun), Hokkaido Pacific West (Mar–May), Hokkaido Sea of Japan (Mar–May), Honshu Pacific (Feb–May), and Honshu Sea of Japan (Feb–Apr).

Chum salmon fry born in rivers along the Pacific coast of Japan migrate to the Sea of Okhotsk (Honda et al., 2017, 2025). Juvenile chum salmon from all regions of Japan are found in the Sea of Okhotsk, and the abundance of juveniles is significantly correlated (positive) to adult returns to Japan (Urawa & Bugaev, 2021). Juvenile chum salmon were abundant in coastal areas of the Sea of Okhotsk from May to June when SST was between 8 and 13°C, and disappeared from coastal waters after late June when SST exceeded 13°C (Nagata et al., 2007). Based on these results, we assumed that chum fry released in all areas would migrate to the Sea of Okhotsk and disperse offshore in July (Figure 1a).

We identified the locations of four main offshore and high-seas areas of Japanese chum salmon distribution (Figure 1b) based on information from previous studies (Myers et al., 2007; 2016; Nagata et al., 2007; Urawa et al., 2018, 2022): 1. Southern Sea of Okhotsk (45.2°N–46.5°N, 142.5°E–148.5°E), 2. Western North Pacific (42°N–45°N, 158°E–165°E), 3. Central Bering Sea (53°N–58°N, 175°E–170°W), 4. Central Gulf of Alaska (48°N–56°N, 140°W–150°W). Using "sst.mnmean.nc" data (NOAA, 2023a), we calculated the monthly mean SSTs (1982–2022) for salmon distribution areas (Figure 1b) using the 'fldmean' function in the "cdo-2.1.1" package (Max Planck Institute, 2023). We mapped areas and migration routes using the "sf" package in R. We used monthly mean SSTs (1998–2021) in our regression model.

2.7 Potential predators of chum salmon fry

We expected that climate-driven changes in abundance and distribution of piscivorous marine fish predators are related to decreases in the marine survival of Japanese chum salmon fry in coastal waters. We used principal components analysis (PCA) to reduce a large number of potential predator variables to a smaller number that explained most of the variance in the predator data. To characterize temporal changes in abundance of marine fish for the first stage of screening for potential predators, we performed PCA on the annual commercial catch data for the marine species in 1980-2020 (MAFF, 2023) using the 'prcomp' function in R. For the

second stage of screening for potential predators, we visualized trends in catches of the marine
species analyzed in PCA. Because catches vary by geographic area and season, we obtained
catch data of potential marine fish predator species identified by PCA by reporting area in
Hokkaido and by prefecture in Honshu (MAFF, 2023), which should reflect their fishing
seasons and migration patterns. Commercial catches of potential marine predators identified by
PCA and chum salmon were mapped in the coastal range of Japanese chum salmon to illustrate
decadal-scale shifts in predator-prey abundance and distribution.
286
287 **2.8 Regression analysis of adult return rates**
288 Our regression model describes the relationship between return rates of chum salmon fry
289 released in year t across all ages ($\hat{R}_t$) and explanatory variables (i=1,…,s) such as SSTs, size-
290 at-release, and catch of potential predators and competitors. We transformed the return rate to
291 the log scale for model fitting in a multiple linear regression:

292
$$\log(\hat{R}_t) = \log\beta_0 + \sum_{i=1}^{s} \beta_i\, z_i + \varepsilon_t$$

293 where β_0 is the intercept, and the explanatory variables are denoted by $\boldsymbol{z}$. The regression
294 coefficients were denoted by $\boldsymbol{\beta}$ and $\varepsilon_t \sim N(0, \sigma^2)$ is the error that we assumed to be
295 independently and normally distributed. We did not include interactions between explanatory
296 variables. We selected the explanatory variables by AICc (Sugiura, 1978), which is the small
297 sample correction of the Akaike Information Criterion (AIC, Akaike, 1973).
298
We estimated regression coefficients based on standardized explanatory variables, which allow
for consistent comparison of different explanatory variables across management regions as
done in salmon studies (e.g., Oke et al., 2020; Ohlberger et al., 2022). The use of an
unstandardized measure (regression coefficient) is preferred to a standardized measure if the
units of measurement (e.g., fry size-at-release, SST, and abundance of competitors) are used to
describe effect sizes at a "practical" or "real world" level (Wilkinson et al., 1999). Therefore,
we also estimated the regression coefficients using unstandardized explanatory variables, as
done in Malick et al. (2023). In the latter, we define effect size as the effect of the unit change
of an explanatory variable on the change in return rate. The effect size and 95% confidence
intervals were calculated by $\exp\left(\hat{\beta}_i \pm 1.96 \times \mathrm{SE}(\hat{\beta}_i)\right)$ for coefficient i. We predicted return
rates by region using the 'predict' function in R based on regression coefficients using the
unstandardized measure and compared them to the observed return rates.
311
When many explanatory variables are included, the model selection procedure may select an
inappropriate model as the best model due to complex multicollinearity. To avoid this problem,
we created three subset models of stressors (see below), performed variable selection within
each subset model, and compared the best models for combinations of the subset models and
the full model, which included all subset models. In each model, we used the variance inflation
factor (VIF) to diagnose multicollinearity among explanatory variables. We determined
variables that could be included in a model by sequentially removing each explanatory variable
with the highest VIF value until all remaining VIFs were less than the threshold of 3.0 (Zuur et
al., 2010), using the 'VIF' function in the "car" package in R. Our model selection procedure
examined all combinations of explanatory variables (2^s, in our case s = 17 without subset
models). Analysis using subset models and a strict VIF threshold should help reduce the risk of
selecting incorrect models due to multicollinearity.
324

The dependent variable is the adult return rates of fry released in 1998–2019. We used 17 hypothesized stressors (covariates) for explanatory variables ($s = 17$) and grouped them into three broad categories of subset models: (1) hatchery carryover (three metrics), (2) ocean condition (five metrics), and (3) predators and competitors (nine metrics) (Table 1). The hypothesized hatchery carryover covariates included relative fecundity and egg size of females in year t-1, where t is the year of fry release, and fry size-at-release. Ocean-condition covariates included SSTs at fry release and in Japanese coastal management regions, summer SSTs in the southern Sea of Okhotsk and central Bering Sea, and winter SSTs in the western North Pacific and central Gulf of Alaska (see Figure 1). Predator and competitor covariates included predation in coastal Japan regions and competition in high-seas areas (Table 1). We identified yellowtail (*Seriola quinqueradiata*), also known as Japanese amberjack, as a potential predator of chum salmon fry (see section 3.4). Age-4 chum salmon were the dominant age group in adult returns to Japan (Figure S1). We used the annual commercial salmon catch (total number of fish) of chum salmon in Russia, the United States, and Canada in year t+3 as indices of intraspecific competition with Japanese chum salmon released in year t, as they return in year t+3 as age-4 fish. The correlation between chum and pink salmon catches in Russia and US/Canada was very weak ($r = 0.0005$ for chum and $r = -0.18$ for pink), and we used Russia and US/Canada catches as covariates. Chum salmon were assumed to spend three years in the high seas and to compete when sharing the same environment. Pink salmon have a fixed life span of two years (Quinn, 2018). We used annual commercial catches in Russia and North America (US and Canada, combined totals) in years t+1, t+2, and t+3 as indices of interspecific competition with Japanese chum salmon released in year t. The ocean distribution of chum and pink salmon overlaps throughout the North Pacific (Langan et al., 2024). We assumed cohorts of pink salmon compete with chum during the first (t+1), second (t+2), and third (t+3) ocean years.

We performed regression analysis independently for each subset model and combinations of subset models (including the full model) with explanatory variables less than 3.0 VIF (see, Table S2). This procedure reduces the number of parameters to be estimated and increases the information available for inference, resulting in robust inference. In each analysis, we selected the most parsimonious model (hereafter, the best model) with the minimum AICc using the "MuMIn" package in R. We compared all models by their minimum AICc values. To summarize the results, we presented the regression coefficients of the best model for the selected subset models. We expected there may be differences in covariate effects among management regions, probably due to differences in the quality and size of fry at release and the coastal environments. Therefore, we modeled the management regions separately (see section 3.1). In three of the four regions, the minimum of the minimum AICc values was achieved by multiple hierarchical subset models. In such cases, we chose the most parsimonious subset models. The estimated regression coefficients were compared with the results obtained by model averaging over all models with ΔAICc values less than 2 (ΔAIC < 2), which suggests substantial support for the model (Burnham & Anderson, 2002). Practical model averaging provides the weighted averages of the regression coefficients over the models. The weights represent the contributions of the models and are proportional to $\exp(-1/2 \times \text{AICc})$. All data and the R script used for the regression analysis are available (see data availability statement).

3 | RESULTS
3.1 Adult return rate (marine survival index)
For Japan as a whole, the number of hatchery-released fry increased until the early 1980s, reaching 2 billion, and then decreased to about 1.3 billion at present (Figure 2). In the Honshu

Pacific, there has been a steady decline in releases since the mid-1990s and a rapid decline since 2021. Decreasing numbers of released fry were also observed in Hokkaido Pacific East and Honshu Sea of Japan, while stable numbers were observed in Hokkaido Nemuro, Hokkaido Pacific West, and Hokkaido Sea of Japan. Only in Hokkaido Okhotsk did the number of fry released increase slightly but continuously. Releases were largest in Honshu Pacific, while other regions had similar release levels.

The total number of adult chum salmon returns throughout Japan increased rapidly until 1996 (88 million fish), and then declined to the current level of 20 million fish (Figure 2). The highest regional returns were in the Honshu Pacific in 1996 (31 million), and then declined continuously to the historical regional minimum (~10 thousand). Among other regions, adult returns were highest in northern Hokkaido regions (Okhotsk and Nemuro) and lowest in Sea of Japan regions (Hokkaido and Honshu). Since approximately 2008, adult returns decreased in all management regions except the Hokkaido Okhotsk and Hokkaido Sea of Japan, and the decrease was most severe in the Hokkaido and Honshu Pacific regions.

The mean adult return rate with the interquartile range of the 5th and 95th percentiles (including ages 2–5) for 1998–2019 releases was highest in Hokkaido Okhotsk (6.9 [3.0, 10.4]%), followed by Hokkaido Nemuro (4.5 [1.1, 9.5]%), Hokkaido Pacific East (2.3 [0.6, 5.5]%) and Hokkaido Pacific West (2.3 [0.6, 4.5]%), Hokkaido Sea of Japan (1.4 [0.5, 2.9]%), Honshu Pacific (1.2 [0.1, 2.3]%), and Honshu Sea of Japan (0.6 [0.4, 1.4]%) (Figure S3). The GAMs showed that log-transformed return rates decreased over time in all management regions, except in the Hokkaido Okhotsk and the Hokkaido Sea of Japan, where return rates increased recently (Figure S4). The nonlinear and/or linear year effect was significant in all management regions except in the Hokkaido Sea of Japan and the Honshu Sea of Japan (Table S1).

The NJ tree analysis showed that the Hokkaido Okhotsk, Honshu Sea of Japan, and Hokkaido Sea of Japan adult return rates were distinct, and the Hokkaido Nemuro, Hokkaido Pacific East and West, and Honshu Pacific return rates were similar (Figure 3). As a result, we used four coastal regions in Japan in our regression analysis: (1) Hokkaido Okhotsk, (2) Hokkaido and Honshu Pacific (combined returns to Hokkaido Nemuro, Hokkaido Pacific East and West, and Honshu Pacific), (3) Hokkaido Sea of Japan, and (4) Honshu Sea of Japan.

3.2 Long-term changes in chum salmon thermal habitat
Decadal-scale maps of SST isoclines showed that until 2000, the summer and winter thermal habitat of chum salmon was stable (Figure 4). After 2010, the optimal summer habitat for chum salmon growth and feeding (especially, 8–11℃ isotherms) shifted north and contracted longitudinally, and the winter habitat remained relatively stable. Evaluation of intra-seasonal changes in spring and summer thermal habitat revealed a progressive northward shift in optimal habitat, from a slight shift in early spring (March and April) to a larger shift in early summer (May and June), northward longitudinal contraction in late summer (August and September), and a slight northward shift in fall (October and November) (Figures S5–S8). In coastal Japan, the optimal thermal habitat for chum salmon fry migrating to the Sea of Okhotsk shifts northward in spring and early summer (April–June), especially in the Sea of Japan (Figure S9).

3.3 Japanese chum salmon migration route and SST
In the Hokkaido Nemuro region, the mean SST increased significantly, while the Hokkaido Pacific West and Honshu Pacific regions showed a similar pattern; the mean SST decreased

421 until around 2010 and then slightly increased (Figure S10, Table S1). In the other regions, the
422 year effects were insignificant.
423
The summer and winter SSTs used in our regression analysis (1998–2022) as common
explanatory variables for all offshore (southern Sea of Okhotsk) and high-seas areas (Figure
1b) were southern Sea of Okhotsk July SST, western North Pacific January–April SST, central
Bering Sea August–September SST, and central Gulf of Alaska January–April SST (Figure
S11a). A GAM (Figure S11b) showed that July SST in the southern Sea of Okhotsk (range,
10.8–14.2°C) increased after the early 2000s, and a strong increase was observed after around
2015. In the western North Pacific, winter SST in January–April (range, 4.6–6.1°C) increased
after around 2015. In the central Bering Sea, August–September SST (range, 8.3–11.4°C)
continued to increase since the late 1990s but decreased after around 2015. In the central Gulf
of Alaska, a strong increasing trend in January–April SST (range, 4.6–7.2°C) was observed after
around 2010, but SST decreased after around 2015. Similar trends were observed in the western
and eastern North Pacific.
436
3.4 Potential predators of chum salmon fry
We summarized the landed commercial catches of all marine species in Japan with no missing
data from 1980 to 2020 (56 species). We excluded marine mammals because catches are subject
to strict limits, and catches were not recorded by species but by total marine mammals. We also
excluded species known not to be predators of salmon (e.g., sardine, smelt, algae, shellfish, and
crustaceans). This resulted in catch data for 30 fish species. The PCA biplot identified
differences in commercial fishery catch levels as the primary component (PC1, 46% variance),
and the second component (PC2, 18% variance) corresponded to catch trends before and after
1998 (Figure 5a). The PCA confirmed that changes in commercial catches of 30 marine fishes
in Japan after 1998 were largely characterized by two species that had positive PC1 values and
increased in recent decades: yellowtail and Japanese Spanish mackerel (*Scomberomorus*
niphonius). We visualized trends in catches of the 30 species (1980–2020) and confirmed that
catches increased after 1998 only in the two species identified by the PCA analysis (unpublished
figure, S. Kitada). Yellowtail and Japanese Spanish mackerel are strong piscivores. There was
a northward shift of yellowtail landings into the range of Japanese chum salmon (Figure 5b). In
contrast, increased landings of Japanese Spanish mackerel occurred only in the eastern area of
Japan, with slight increases in the Honshu North Pacific area. Based on these results, we
identified yellowtail as a possible new predator of chum salmon fry.
455
3.5 Regression analysis of adult return rates
All models used explanatory variables with a VIF less than the threshold of 3.0. Model selection
results for all subset models are shown in Table S2. Egg size and relative fecundity of age-4
and age-5 fish were examined, but these explanatory variables were selected only for Hokkaido
and Honshu Pacific, where age-4 fish had a smaller AICc (30.0) and larger R^2 (0.79) than age-
5 fish (AICc = 38.2 and R^2 = 0.61). Therefore, we used egg size and relative fecundity of age-
4 fish in our regression analysis. The full model results and the best subset model results were
the same for all management regions except for the Hokkaido Sea of Japan (Table 2; Tables S2
and S3). The results of the model averaging (Table S4) were qualitatively the same as the best
models (Table 2) for all regions. The explanatory variables selected in the best models were
region-specific or common to some areas, or both.
467
For ocean conditions, the effects of SST were found only for fry released in the Hokkaido Sea

of Japan and Honshu Sea of Japan regions (Table 2). SST at release (year t) and summer SST
(July) in the Sea of Okhotsk (year t) and the Bering Sea (August-September) (year t+1) were
negatively associated with Japanese chum salmon return rates. In contrast, winter SST (January-
April) in the western North Pacific (year t+1) and the Gulf of Alaska (year t+2) were positively
associated with return rates. For predators and competitors, we found negative associations
between return rates and yellowtail catch in the year of fry release only in the Hokkaido Okhotsk
region, while negative effects of Russian pink and/or chum salmon catches were found in all
regions. Russian chum salmon catch in year t+3, and Russian pink salmon (adult, age-1) catch
in years t+2 and t+3 were negative. No associations between Japanese chum salmon return rates
and US and Canadian chum and pink salmon catches were found. For hatchery carryovers, fry
size-at-release was positively associated with salmon return rates and relative fecundity was
negatively associated with return rate only in the Hokkaido and Honshu Pacific region.

The GAMs indicated a similar trend in fry size-at-release in Hokkaido Okhotsk, Hokkaido Sea
of Japan, and Honshu Pacific, where fry size-at-release increased or remained at similar levels
until around 2010 and then declined (Figure S12, Table S1). The decline in fry size-at-release
was remarkable in the Honshu Pacific and Honshu Sea of Japan regions. In contrast, the fry
size-at-release increased after about 2015 in Hokkaido Nemuro. Fry size-at-release varied
among regions (Figure S13) and the mean was smallest in the Hokkaido Sea of Japan, while
the Honshu Sea of Japan had the greatest variation with a recent minimum.

We found that the relative fecundity of parent females was negatively associated with return
rates in the Honshu Pacific region. Relative fecundity was negatively correlated with egg size
in all management regions except in the Hokkaido Sea of Japan region (unpublished figure, S.
Kitada). GAMs confirmed that egg size for age-4 fish has decreased over the last two decades
in all regions, except Hokkaido Okhotsk (Figure 6, Table S1). The trend was significant in all
regions, except Honshu Sea of Japan. Similar decreasing trends in egg size for age-5 fish were
also observed (Figure S14, Table S1). Egg size varied among the management regions, and was
smallest in the Hokkaido Sea of Japan region (Figure S15).

The standardized model effect size of explanatory variables on adult returns allowed
comparison of the magnitude of the effects across the four regions (Figure 7). The largest
positive effect was fry size-at-release (year t), which was equal to winter SST (January–April)
in the Gulf of Alaska (year t+2), followed by winter SST (January–April) in the western North
Pacific (year t+1). On the other hand, parental relative fecundity (year t-1) had the largest
negative effect, followed by Russia pink and/or chum salmon catches which varied among
regions. The negative effect of summer SST in the Bering Sea (August–September) (year t+1)
was the second largest, followed by summer SST in the Sea of Okhotsk (July) (year t) and SST
at the time of release (year t). The negative effect of yellowtail (year t) was similar to summer
SST in the Bering Sea.

The practical effect size in a regression model describes how much of a change in the dependent
variable is expected for a given change in the independent variable in each region. The effect
size for each regression coefficient, measured by $\exp(\hat{\beta}_i)$, represents the rate of change per
practical unit in each management region (Figure S16), which showed a similar pattern to the
standardized effects of the explanatory variables (Figure 7). The area-specific practical effect
sizes ($\exp(0.486) = 1.625$) showed that a 0.1 g increase in hatchery fry size increased the
return rate by 62.5% (($1.625-1) \times 100$) in the Hokkaido and Honshu Pacific region (Table 3).

The average return rate over the last five years (2015–2019 releases) was 0.74% in this region, and a 0.1g increase in size-at-release would be expected to increase the return rate to 1.2% (= $0.74 \times (1+0.625)$) from current levels. A 1°C increase in winter SST in the Gulf of Alaska increased the return rate by 49.9% in the Hokkaido Sea of Japan region. On the contrary, an increase of 100 eggs (relative fecundity) of parent females (age-4) (= egg size reduction) reduced the return rate by 60.6% in the Hokkaido and Honshu Pacific regions, and a 1°C increase in SST at release reduced the return rate by 38.3% in the Honshu Sea of Japan region. An increase in biomass of yellowtail landings of 100 tons predicted an 8.9% reduction in the adult return rate of chum salmon in the Hokkaido Okhotsk region.

The correlation between predicted and observed adult return rates was high in all four coastal regions: Hokkaido Okhotsk ($r = 0.79, t = 5.75, p = 0.0000$), Hokkaido and Honshu Pacific ($r = 0.91, t = 9.98, p = 0.0000$), Hokkaido Sea of Japan ($r = 0.79, t = 5.78, p = 0.0003$), and Honshu Sea of Japan ($r = 0.81, t = 6.17, p = 0.0000$). Our best models with few (2–4) explanatory variables provided good fits to the four coastal regions and the annual variations in chum return rates, although the 95% prediction intervals are large (Figure 8).

4 | DISCUSSION
We investigated relationships between the marine survival of Japanese hatchery chum salmon and multiple hypothesized stressors during 25 years of climate change. While previous studies of Japanese chum salmon have focused largely on climate change impacts on survival during the early marine (coastal, fry) life history stage, we hypothesized that marine survival is related to climate change effects during all life stages (from eggs to adults). We identified potential climate-driven stressors along known coastal and high-seas migration routes of Japanese chum in the North Pacific Ocean and adjacent seas. Associations between stressors and marine survival varied by coastal region, salmon life stage, and seasonal high-seas distribution area. The stressors exhibiting the largest model effects were fry size-at-release (positive) and parental relative fecundity (negative), and were found only in the Hokkaido and Honshu Pacific where the decline in chum salmon catches was most severe. The negative correlation between relative fecundity and egg size of age-4 fish was significant in this region (*r* = -0.60, *t* = -3.34, *p* = 0.003), indicating that egg size had the largest positive effect. We identified yellowtail, whose distribution was shifting northward, as a new potential marine fish predator of chum salmon fry in the Hokkaido Okhotsk. The model effects of SST were inferred only for fry released in the Hokkaido Sea of Japan and Honshu Sea of Japan regions, and the directional effects of SST were consistent for summer (negative) and winter (positive) in both regions. The SST at release and summer SST in the Sea of Okhotsk and the Bering Sea were negatively associated with marine survival, while winter SST in the western North Pacific and the Gulf of Alaska was positively associated with marine survival. Negative associations between Russian chum and/or Russian pink salmon catches and Japanese chum salmon marine survival were found in all regions, while no associations were identified between US and Canadian chum and pink salmon catches. Moreover, we found a nationwide decline in egg size in Japanese chum salmon over the past two decades.

4.1 Adult return rates (marine survival index)
Our adult return rate estimates should be regarded as indices of marine survival because little is known about freshwater mortality of hatchery chum salmon fry after release (e.g., Hasegawa et al., 2021b) or mixtures of regional populations in landed commercial catches in Japan (Nagata et al., 2016). Genetic stock identification showed that 80–98% of the mixed samples

565 of chum salmon caught in September– early October along the western coast of the Sea of
566 Okhotsk originated from Hokkaido rivers, while 53−71% originated from Honshu rivers in late
567 October (Saito et al., 2020). The mixed-stock fishery could bias the return rates in each
568 management region in Hokkaido and Honshu.
569
It is generally accepted that most chum salmon in Japan are hatchery-reared fish (Hiroi, 1998;
Kaeriyama, 1999; Kobayashi, 1980; Iida et al., 2018; Kaeriyama & Sakaguchi, 2023). We
assumed all chum salmon returning to Japan were hatchery reared. The total proportion of wild
chum salmon in Japan is largely unknown. Field studies provide evidence that natural
reproduction of chum salmon occurs in Hokkaido (Miyakoshi et al., 2012) and in the Honshu
Sea of Japan (Iida et al., 2018; Iida et al., 2021). Hatchery salmon have a relative reproductive
success (RRS, number of offspring a salmon produces compared to other salmon in the
population) that is approximately 50% of the RRS of wild fish (Araki et al., 2007; Christie et
al., 2014; Shedd et al., 2022), with considerable variation (Kitada et al., 2011). Reduced RRS
can be carried over to the next generation (Araki et al., 2009). We expected low freshwater
survival from egg to ocean entry of wild chum salmon in natural streams in Japan. Our marine
survival indices may be overestimated depending on the magnitude of recruitment of wild fish,
which we expected to be very low compared to the number of hatchery-released fish (100–200
million fry per year in each management region, Figure 2). Our regression analysis
overestimated the intercept proportional to the magnitude of natural recruitment, but regression
coefficients were expected to be robust if natural recruitment and genetic mixture were constant
over the period of our analysis.
587
4.2 Regression model assumption
Our linear regression model of the log-transformed return rate assumes that the error is
independent and normally distributed. We checked the model residuals to see if the normality
assumption of our linear regression model held for our data. The qq plots of the residuals from
the best models (Table 2) based on the real scale data showed that most of the studentized
residuals from our linear model were within the 95% confidence intervals against the theoretical
quantiles of a normal distribution (Figure S17). The result indicates that there are no serious
problems with our regression model. This is probably because the catches were not small in
number, although the return rates were very small. Return rates are proportions that can also be
modeled with a beta distribution with a logit link function. Beta linear regression is a topic for
future research. We inferred year effects for logit-transformed return rates with a beta error
(Figure S18), which were very similar to those for log-transformed return rates with a normal
error (Figure S4), showing that the log-normal model visualizes the year effects of return rates.
601
4.3 Egg size and body size of fry at release
Our finding of a nationwide decline in egg size of age-4 and age-5 Japanese hatchery chum
salmon over the past two decades is striking. Egg size is an important fitness trait, and larger
eggs have a positive effect on juvenile survival and growth in salmonids (Hutchings, 1991;
Einum & Fleming, 1999, 2000). Negative relationships between egg size and fecundity have
been reported for Pacific salmon (Beacham & Murray, 1993; Fleming & Gross, 1990). Our
regression model coefficients (practical effect size, Table 2) suggested that higher relative
fecundity (smaller eggs) could contribute to the reduced survival rate of chum salmon in the
Hokkaido and Honshu Pacific region. In fact, the correlation between egg sizes and return rates
in this region was significant for age-4 ($r = 0.76$, $t = 5.17$, $p = 0.0000$) and for age-5 ($r = 0.81$,
$t = 6.13$, $p = 0.0000$) chum salmon, even though egg size was excluded from the explanatory

variables due to multicollinearity (Table S2). Selection in hatcheries could impact fecundity
and egg size (Heath et al., 2003), although egg size or fecundity were not found to differ
between hatchery and wild fish in other populations (Beacham & Murray, 1993; Quinn et al.,
2004; Beacham, 2010). Hasegawa et al. (2021a) showed that the egg size of 5-year-old Japanese
chum salmon in 2010 was smaller in more northeasterly locations than in 1994 and 2001, and
hypothesized that domestication selection in hatcheries affected by global warming may have
caused the temporal shifts in egg size. An alternative process might be maternal effects,
whereby environmental variation (e.g., ocean temperature during egg development; maternal
diet) experienced by the mother is translated into phenotypic variation in reproductive traits of
offspring (e.g., egg size, age at maturation) (reviewed by Mousseau & Fox, 1998; Thorn et al.,
2019).

Our regression model coefficients (Table 2) suggested that fry size-at-release and quality of
hatchery fish are particularly important for survival at later ocean stages for the Hokkaido and
Honshu Pacific region chum salmon. After release, Honshu Pacific region chum fry migrate
long distances northward against the Oyashio Current (Figure 1a) to their summer feeding area
in the Sea of Okhotsk. On the Sanriku Coast, Honshu Pacific region, where the decline in
abundance was most severe (Figure 2), chum salmon fry reared under low food availability (1%
body weight) allocated less than half of their energy to growth at high temperatures (14℃)
compared to fry reared under high food availability (4% body weight; Iino et al., 2022). In the
high food availability group, 32–47% of the energy consumed by the fish should be used for
daily growth (Iino et al., 2022). Japanese hatchery chum salmon larger than 2 g (BW)
significantly increased long-distance swimming performance and reduced the metabolic cost of
swimming against the southward current at 8–12°C SST (Iino et al., 2024). Juvenile chum
salmon in western Alaska consumed lower quality prey during warm periods than during cold
periods, with the proportion of lower quality prey increasing during warm periods, and the
energy density of juvenile chum salmon was lower at warmer SST (Farley et al., 2024).

4.4 Predators of fry during coastal migrations
641
Predation is well-accepted as the proximate cause of salmon fry mortality during their coastal
migrations. There are many known fish, bird, and marine mammal predators of Japanese chum
salmon (Okado et al., 2020; Hasegawa et al., 2021b; Okado & Hasegawa, 2024). Long-term
climate change and marine heatwaves are projected to result in poleward shifts in the
distribution of many North Pacific marine species by 2050 (Cheung et al. 2015, Cheung &
Frölicher, 2020). Commercial catch data for coastal marine species in Japan was a useful metric
for identifying poleward shifts in the distribution of potential piscivorous fish predators of chum
salmon fry.

We identified yellowtail as a potential marine fish predator of chum salmon fry. Yellowtail are
a subtropical species distributed from Japan and the eastern Korean Peninsula to the Hawaiian
Islands (Robins et al., 1991), and are one of the most commercially important marine fish
species in Japan. A climate regime shift (warming) during the late 1980s was associated with
the northward extension of yellowtail distribution in the Tsushima Warm Current off
northwestern Honshu and Hokkaido (Figure 1a), and global warming was projected to further
expand the suitable thermal habitat of yellowtail northward (Tian et al., 2012). Since 2014,
yellowtail have been caught along the coast of the Sea of Okhotsk in May–December by a setnet
fishery targeting adult salmon (September–December), and other marine fish species (May–
August) (Hoshino & Fujioka, 2021). The rapid increase in salmon setnet catches of yellowtail

led to the identification of this species as a potential indicator of human (fishing) pressure in Japan (Hunsicker et al., 2022). In May 2023, the peak month of hatchery releases of chum salmon fry in the Hokkaido Okhotsk region (Figure S2), schools of yellowtail were often observed on the sea surface near the central Sea of Okhotsk coast (pers. comm., Takashi Ichikawa). Adult yellowtail are well-known to feed on small pelagic fish and invertebrates (Robins et al., 1991), but stomach contents surveys are necessary to confirm yellowtail predation on chum salmon fry.

Our regression model results indicated yellowtail catches are an important covariate of fry chum survival in the year of release only in the Hokkaido Okhotsk region, although yellowtail catches were higher in the more southerly management regions such as Hokkaido Sea of Japan and Honshu Pacific (Figure 5b). Yellowtail spawning grounds cover a wide area from the East China Sea to western Kyushu Island, and the main spawning season is from February to June (Tian et al., 2012). After spawning, adult yellowtail migrate north for feeding. Yellowtail fry also migrate north on drifting seaweed using the Tsushima Warm Current in the Sea of Japan and the Kuroshio Current off the Pacific coast of Honshu (Figure 1a). Yellowtail (age 0−3 years and older) are caught in Hokkaido from May to August (Hoshino & Fujioka, 2021), while hatchery chum fry are released mainly in March-April in the Hokkaido Sea of Japan region (Figure S2). On the Pacific coast of Honshu, yellowtail (age unknown) are caught in commercial fisheries from late July to mid-September (Iwate Prefecture, unpublished figure) and from June to December (Miyagi Prefectural Government, 2022), while hatchery chum fry were released mainly in February–April (Figure S2). The earlier timing of chum fry release compared to the timing of the yellowtail migration may be a reason for the exclusion of yellowtail catches from the explanatory variables in these regions.

Annual estimates of stock abundance (biomass) or relative abundance (catch per unit effort) would be more appropriate covariates than annual catches when considering many potential predator species (e.g., Pauly et al., 2013; Thorson et al., 2020; Punt, 2023), but were not available for the period of our study. The use of stock assessment data is a promising approach for developing new indices of climate-driven predator exposure, especially if predator diet information is available. A similar approach for high-seas migration areas may be possible through future scientific collaborations among international treaty organizations (PICES-NPFC, 2019).

4.5 Inter- and intra-specific competition during high-seas migrations
Indirect evidence of competition between Russian pink salmon and many other marine species, including chum salmon, is substantial (see review by Ruggerone et al., 2023). Our results suggest that Russian pink and chum salmon compete for food, space, or other resources with Japanese chum salmon (ages 2−3) in the central Bering Sea in summer and the Gulf of Alaska in winter, while competition with immature (age-1) chum salmon could be weak or negligible in the Sea of Okhotsk and western North Pacific (Naydenko & Temnykh, 2016; Naydenko et al., 2016; Bugaev & Gerlits, 2023). Our analysis found no relationship between Japanese chum salmon return rates and North American pink and chum salmon catches. This may reflect inter- and intra-specific differences in the ocean distribution of Japanese chum salmon and North American pink and chum salmon. For example, limited data from salmon research vessel surveys in winter indicate Japanese chum salmon are distributed farther to the south in the Gulf of Alaska and northeastern North Pacific than North American chum salmon (reviewed by

Myers et al., 2016; Urawa et al., 2016). Past research indicated negative intraspecific
relationships between the abundance of Japanese hatchery chum salmon and the abundance and
annular scale growth of local and regional stock groups of North American chum salmon
(Ruggerone et al., 2011; Agler et al., 2013; Frost et al., 2021). All our best regional models
included North American chum salmon as an explanatory variable with VIF < 3 (Table S2), but
were excluded by our model selection procedure. Our regression analysis (Table 2) detected a
negative association between return rates of Hokkaido Sea of Japan chum and Russia chum
catch (r = -0.44, t = -2.22, p = 0.038), while correlations between regional return rates and North
American chum catch were too weak to detect North American chum salmon as a competitor.
717
A reanalysis of coastal and high-seas salmon research survey data (1953–2022) confirmed the
results of earlier studies showing the overlap in distributions of chum, pink, and sockeye salmon
throughout the North Pacific (Langan et al., 2024). A meta-analysis of salmon trophic ecology
data showed similar diet composition of chum, pink, and sockeye salmon throughout the North
Pacific, with a diet overlap of 31.8% between chum and pink and 30.9% between chum and
sockeye, and diet overlap was highest between pink and sockeye salmon at 46.6% (Graham et
al., 2021). The estimated proportion of Asian and North American hatchery chum salmon was
60% of the total abundance of maturing chum salmon during 1990–2015 (Ruggerone & Irvine,
2018). The substantial decline in abundance of Japanese chum salmon after the mid-2000s may
have reduced inter- and intra-specific competition for prey resources with Russian chum and
pink salmon. The lack of an association between Russian pink salmon catches (t+1) and survival
of age-1 Japanese chum salmon might also indicate improved growth and survival in the Sea
of Okhotsk. However, even when catches of Japanese chum salmon were low, their survival
(age-2 and age-3 fish) was negatively correlated with Russian chum and pink salmon catches,
suggesting that Russian chum and pink salmon have a competitive advantage over Japanese
chum salmon.
734
Directional swimming to preferred thermal habitat seems to play an important role in the
seasonal northward migrations of salmon (Azumaya & Ishida, 2004). Russian pink and chum
salmon may have a competitive advantage simply because they do not have as far to swim from
their overwintering habitat (central North Pacific and Aleutian Islands) as Japanese chum
salmon (Gulf of Alaska) to their shared optimal summer feeding habitat in the central Bering
Sea. If Russian salmon are first to arrive in the central Bering Sea in summer, they might occupy
most of the optimal thermal habitat and reduce the amount of available preferred prey, causing
a trophic cascade (Batten et al., 2018). Models project that the optimal thermal habitat of chum
and pink salmon will continue to contract and shift northwestward under future climate-change
scenarios (e.g., Abdul-Aziz et al., 2011, also see Figures 4, S7), and we expect Russian salmon
to continue to have a competitive advantage over Japanese chum salmon.
746
We hypothesize that inferior athletic performance in a warming climate may be related to the
long-term decline in the marine survival of Japanese hatchery chum salmon. Wild-born chum
salmon fry were too agile to be netted before 1945 in a river on Etrofu Island where Japan
operated salmon hatcheries (Yagisawa, 1970). Currently, piscivorous salmonids chase schools
of hatchery chum salmon fry, and small fry are easily predated due to their slow swimming
speed to escape predators (Hasegawa et al., 2021b). Water temperature and oxygen are
important regulators of physiological processes in fish (Powers et al., 1991; Somero, 2022).
Swim tunnel experiments show that temperature and body size influence swimming
performance in Japanese hatchery chum fry (Iino et al., 2024). Swimming performance has

been least studied in adult chum salmon and Chinook salmon (reviewed by Kraskura et al.,
2024).
758
4.6 Ocean conditions in coastal and high-seas habitats
Ranges of chum salmon thermal habitat used in our analysis were narrower (thermal limits, 4–
13°C) than previously published estimates, based largely on historical high-seas salmon survey
data collected before the winter 1976–1977 climate regime shift (preferred spring-summer SST
range, 1.3–12.6°C; Langan et al., 2024). The pre-regime shift period (1948–1975) was
characterized by low ocean productivity, cold winter SSTs in the Gulf of Alaska, and a low
abundance of salmon populations that overwinter in the Gulf of Alaska, including Japanese
chum salmon. During periods of low ocean productivity, optimal temperatures for salmon
growth decline (Beauchamp, 2009).
768
Marine heatwaves and long-term climate change are projected to result in poleward shifts in
the distribution of many North Pacific pelagic fish species by 2050 (Cheung et al. 2015, Cheung
& Frölicher, 2020). Our maps of Japanese chum salmon thermal habitat indicated a summer
northward shift and contraction since 2010; winter habitat remained relatively stable with a
slight northeastward shift in 2017 and 2018 (unpublished figure, S. Kitada). The frequency of
annual marine heatwaves increased in the western and central Bering Sea, and heatwaves in the
eastern Bering were longer and more intense in the past decade (2010–2019) (Carvalho et al.,
2021). Marine heatwaves in the Gulf of Alaska (2013–2016, 2018–2020) were accompanied by
increased acidity and low oxygen, resulting in cascading ecosystem effects (Hauri et al., 2024).
Our analysis showed increases in SST in the central Bering Sea increased after 2010 and SST
in the Gulf of Alaska in 2014–2015 and 2019–2020 (Figure S11b) that are consistent with the
occurrence of marine heat waves.
781
Japanese chum salmon from all regions would experience similar environmental effects from
common coastal and high-seas habitat conditions, but the effects of SST were inferred only for
fry released in the Hokkaido Sea of Japan and Honshu Sea of Japan regions. The Sea of Japan
is a global hotspot for long-term declines in marine fisheries productivity (Free et al., 2019).
After a brief (1998–1999) natural cooling event, anthropogenic climate change became the
principal driver of extreme warming events in the Sea of Japan, particularly in southern regions
(Hayashi et al., 2022). Our regression analysis found that SST at release was negatively
associated with return rates in the Honshu Sea of Japan. Summer SST (July) on the Hokkaido
Okhotsk coast was also negatively associated with return rates for fry released in the Sea of
Japan regions. Summer SST (July) on the Hokkaido Okhotsk coast (range, 10.8–14.2°C) was
higher than published temperatures for optimal growth and feeding (8–12°C) and available
habitat for feeding (5–13°C) of Japanese chum salmon (Kaeriyama et al., 2012; Urawa et al.,
2018). Hatchery chum salmon fry were released mainly in February–March along the Honshu
Sea of Japan coast and March–April along the Hokkaido Sea of Japan coast (Figure S2). After
release, chum salmon juveniles migrate in the direction of the northbound Tsushima Warm
Current to the Sea of Okhotsk (Figure 1a). The negative association of survival with July SST
may be related to lengthy exposure to warm coastal SSTs in the Sea of Japan and delayed arrival
at their nursery area in the Sea of Okhotsk. Juvenile chum salmon were abundant in the coastal
Okhotsk Sea from May to June when SST was between 8 and 13°C, and disappeared from
coastal waters after late June when SST exceeded 13°C (Nagata et al., 2007). Hokkaido
Okhotsk is the most successful management region (Figure 2), and the peak release time is in
May (SST, 4.7–6.2°C). Late arrival of juveniles released from the Honshu and Hokkaido Sea

of Japan regions to the Sea of Okhotsk may reduce survival during the first summer. The results
suggest that release timing is important to avoid mortality at release and summer mortality on
the Hokkaido Okhotsk coast.
807
In high-seas habitats, summer SST (August–September) in the Bering Sea (range, 8.3–11.4°C)
was within the published optimal growth and feeding range of Japanese chum salmon (8–12°C;
Kaeriyama et al., 2012; Urawa et al., 2018). The negative effect of summer SST in the Bering
Sea was found only for fry released in the Hokkaido Sea of Japan. In this region, egg size was
the smallest in Japan (Figure S15). These results suggest a hypothesis that the upper thermal
tolerance is lower for hatchery salmon produced in this region. On the contrary, the model
effects of winter SST in the western North Pacific and in the Gulf of Alaska were consistently
positive for fry released in the Sea of Japan regions. Winter SST (January–April) increased after
2010 in the western North Pacific (range, 4.6–6.1°C) and the Gulf of Alaska (range, 4.6–7.2°C)
(Figure S11b); these were similar or slightly higher than the published overwintering
temperature range (4–6°C; Kaeriyama et al., 2012; Urawa et al., 2018). This result supports the
hypothesis that Japanese chum salmon in the Sea of Japan regions affected by the Tsushima
Warm Current are better able to survive increasing winter SST (Figure S11b).
821
In 2022, the number of chum salmon returning to Japan unexpectedly increased in the Hokkaido
Sea of Japan and the Hokkaido Okhotsk management regions (Figure 2). In the Chitose River
in the Hokkaido Sea of Japan, where Japan's oldest hatchery operates, 587,000 fish returned,
surpassing the historical record of 551,000 in 1995 (Chitose Aquarium, 2023). The sudden
increase in 2022 surprised fishermen, hatchery managers, local people, and biologists. However,
returning chum numbers declined again in all geographic regions in 2023, reaching historic
lows (a total of about 18 million in Japan) in 2024. In the Hokkaido Sea of Japan and Okhotsk
regions, the peak in adult returns in 2022 (Figure 2) was mainly caused by an increase in age-4
fish released as fry in 2019 (Figure S19), indicating better survival of the 2019 year class.
Thermal habitat (8–10°C) within the published optimal thermal habitat range for growth and
feeding (8–12°C) of Japanese chum salmon was widely distributed in the Sea of Okhotsk in
July 2019 and in the central Bering Sea in September 2020 (Figure S20), where age-1 Japanese
chum salmon are distributed during their second summer growing season. Expansion of optimal
thermal habitat during the first and second summers likely improved the survival of immature
chum salmon in 2020 and 2021, and may have resulted in the anomalous peak returns of age-4
fish in 2022. In contrast, total return rates decreased in other regions (Honshu Sea of Japan and
the Pacific regions; Figure S19). The results suggest that coastal mortality of fry released from
these regions was higher and substantial.
840
4.7 Management implications
The largest positive model effect identified by our regression analysis was fry size-at-release
and egg size (relative fecundity, negative), suggesting that hatchery fry quality (size-at-release)
and release timing are particularly important for survival in later ocean stages. Hatchery
experiments to test appropriate fry size and time of release and nearshore monitoring of SSTs
might provide an adaptive approach for improving adult returns of Japanese chum salmon
(Saito & Nagasawa, 2009; Nagata et al., 2016; Iida et al., 2021; Saito, 2022; Iida, 2024; Honda
et al., 2025). Adjustment of release timing to allow fry to arrive in the Sea of Okhotsk in May
or early June or both might improve nearshore survival. Experiments to assess fry quality,
metabolism, and swimming performance of Japanese chum salmon (e.g., Iino et al. 2022, 2024;
Abe et al., 2024) should provide information useful for improving hatchery fry quality.

International comparisons of the swimming performance of hatchery and wild chum salmon may provide crucial information for Japanese chum salmon fitness. Moreover, our analysis found a nationwide decline in egg size over the past two decades. The mechanisms of egg size reduction in Japanese chum salmon are unknown and an important topic for further study. A nationwide evaluation of current hatchery techniques including parent selection, egg incubation, and fry rearing and release strategies would be useful for understanding the underlying mechanisms of the decline of Japanese chum salmon abundance over the past two decades.

Investigations of many other potentially important stressors beyond the scope of our study remain to be addressed. For example, relationships between the marine survival of salmon and climate-driven changes in marine fisheries operating along high-seas salmon migration routes and within the 200-mile EEZs of other nations. In Alaska, genetic analyses show that Asian (Russia and Japan) chum salmon populations dominate early (June) catches by salmon fisheries targeting sockeye salmon along the south side of the Alaska Peninsula in early (June), when chum salmon are migrating from winter (North Pacific) to summer (Bering Sea) feeding grounds (Dann et al., 2023). Asian stocks of chum salmon also dominate prohibited species bycatch of salmon in the US walleye pollock (*Gadus chalcogrammus*) fishery in the eastern Bering Sea in early summer (Barry et al., 2022). Bycatches of Asian chum salmon (ocean-age-3 and age-4; unknown maturity stage) by both fisheries in 2021 were unexpectedly high (Barry et al., 2022; Dann et al., 2023). The Alaska bycatch of Asian chum salmon preceded the spike in adult returns to Japan in 2022. The chum salmon bycatch in the Bering Sea decreased substantially in 2022 and 2023 (NPFMC, 2025), coinciding with the decrease in adult returns to Japan in 2023 and 2024. Seeking predictive (vs. retrospective) indices would benefit preseason forecasting of adult salmon returns. In the absence of quantitative marine stock assessment surveys of salmon, potentially predictive indices might be used in a qualitative "stoplight" approach to pre-season forecasting (Caddy, 2002; NOAA, 2024). This method requires an adaptive approach because relationships between salmon survival and index variables change over time (non-stationarity; Litzow et al., 2018; Ohlberger et al., 2022).

5 | CONCLUSIONS

Our results from the world's largest long-term salmon hatchery program indicate that maternal traits (egg size and fecundity), hatchery fry quality, fry body size-at-release, and release timing are particularly important for survival at later ocean stages. Egg size and parental fecundity are key demographic parameters that contribute to the fitness and productivity of salmon populations, and the decadal-scale changes observed herein could result in significant ecosystem, fishery, and societal losses. Investigation of the mechanisms of egg size reduction is one of the most important topics for further study in Japanese chum salmon. Our study highlights the need for an experimental approach to hatchery practices, including monitoring and analyses with updated information, leading to effective management decisions and policies for future sustainability and conservation of salmon resources.

AUTHOR CONTRIBUTIONS

Shuichi Kitada: Conceptualization (lead); data curation (lead); formal analysis (lead); funding acquisition (lead); investigation (lead); methodology (equal); software (lead); validation (equal); visualization (lead); writing–original draft (lead); writing–review and editing (equal). Katherine W. Myers: Conceptualization (equal); writing–review and editing (lead). Hirohisa Kishino: Conceptualization (equal); formal analysis (support); funding acquisition (equal); methodology (lead); validation (lead); visualization (support); writing–review and editing

(equal). All authors approved the final version of the manuscript.

ACKNOWLEDGMENTS
We sincerely acknowledge the thousands of biologists, fishery and hatchery managers, and fishers who contributed to the collection of data analyzed in our manuscript. The efforts of APG, FRA, HSEA, JMA, MAFF, NOAA, and NPAFC to create public databases made our study possible. We thank Brian Riddell and the reviewers for their insightful and constructive comments, which significantly improved the earlier version of the manuscript. Takashi Ichikawa provided information on chum salmon and yellowtail. This study was supported by Japan Society for the Promotion of Science Grants-in-Aid for Scientific Research KAKENHI No. 18K0578116 for SK and No. 22K11950 for HK.

CONFLICT OF INTEREST STATEMENT
The authors declare that no conflict of interest exists.

DATA AVAILABILITY STATEMENT
The data generated or analyzed in this study are fully described in the published article and its Supplementary Information. Analysis scripts and data will be available at https://doi.org/10.5281/zenodo.14868156 on acceptance.

ORCID
Shuichi Kitada https://orcid.org/0000-0001-5838-0374
Hirohisa Kishino https://orcid.org/0000-0002-3244-359X

REFERENCES
Abdul-Aziz, O. I., Mantua, N. J., & Myers, K. W. (2011). Potential climate change impacts on thermal habitats of Pacific salmon (*Oncorhynchus* spp.) in the North Pacific Ocean and adjacent seas. *Canadian Journal of Fisheries and Aquatic Sciences*, 68, 1660–1680. doi.org/10.1139/f2011-079

Abe, T. K., Kitagawa, T., Iino, Y., Ito, M., & Sato, K. (2024). Ecological features of upriver migration in Kitakami River chum salmon and their connection to aerobic thermal performance. *Conservation Physiology*, 12, coae087. https://doi.org/10.1093/conphys/coae087

Agler, B. A., Ruggerone, G. T., Wilson, L. I., & Mueter, F. J. (2013). Historical growth of Bristol Bay and Yukon River, Alaska chum salmon (Oncorhynchus keta) in relation to climate and inter-and intraspecific competition. *Deep Sea Research Part II: Topical Studies in Oceanography*, 94, 165–177. doi.org/10.1016/j.dsr2.2013.03.028

Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. In B. N. Petrov & F. Csaki (Eds.), *Second International Symposium on Information Theory* (pp. 267–281). Budapest: Akademiai Kiado. Reprinted in S. Kotz & N. L. Johnson, Eds. (1992) *Breakthroughs in Statistics Volume I, Springer Series in Statistics*, 610-624. New York: Springer-Verlag. doi.org/10.1007/978-1-4612-0919-5_38

Amoroso, R. O., Tillotson, M. D., & Hilborn, R. (2017). Measuring the net biological impact of fisheries enhancement: Pink salmon hatcheries can increase yield, but with apparent costs to wild populations. *Canadian Journal of Fisheries and Aquatic Sciences*, 74, 1233–1242. doi.org/10.1139/cjfas-2016-0334

APG (Aomori Prefectural Government) (2024). *Chum salmon catch statistics*, Aomori: APG. (in Japanese) www.pref.aomori.lg.jp/soshiki/nourin/sshinko/sake.html (accessed 20

November 2024).

Araki, H., Cooper, B., & Blouin, M. S. (2007). Genetic effects of captive breeding cause a
rapid, cumulative fitness decline in the wild. *Science*, 318, 100–103.
doi.org/10.1126/science.1145621

Araki, H., Cooper, B., & Blouin, M. S. (2009). Carry-over effect of captive breeding reduces
reproductive fitness of wild-born descendants in the wild. *Biology Letters*, 5, 621–624.
doi.org/10.1098/rsbl.2009.0315

Azumaya, T., & Ishida, Y. (2004). An evaluation of the potential influence of SST and
currents on the oceanic migration of juvenile and immature chum salmon
(*Oncorhynchus keta*) by a simulation model. *Fisheries Oceanography*, 13, 10–23.
https://doi.org/10.1046/j.1365-2419.2003.00271.x

Bahri, T., Vasconcellos, M., Welch, D.J., Johnson, J., Perry, R.I., Ma, X. & Sharma, R., Eds.
(2021). Adaptive management of fisheries in response to climate change. *FAO*
Fisheries and Aquaculture Technical Paper 667, Rome: FAO.
doi.org/10.4060/cb3095en

Barry, P. D., Kondzela, C. M., Whittle, J. A., Watson, J. T., Karpan, K., D'Amelio, K., &
Larson, W. A. (2022). Genetic stock composition analysis of chum salmon from the
prohibited species catch of the 2021 Bering Sea walleye pollock trawl fishery. U.S.
Department of Commerce, *NOAA Technical Memorandum NMFS-AFSC-*
454.repository.library.noaa.gov/view/noaa/47923/noaa_47923_DS1.pdf

Batten, S. D., Ruggerone, G. T., & Ortiz, I. (2018). Pink salmon induce a trophic cascade in
plankton populations in the southern Bering Sea and around the Aleutian Islands.
Fisheries Oceanography, 27, 548-559. https://doi.org/10.1111/fog.12276

Beacham, T. D. (2010). Revisiting trends in the evolution of egg size in hatchery-enhanced
populations of Chinook salmon from British Columbia. *Transactions of the American*
Fisheries Society, 139, 579–585. doi.org/10.1577/T09-093.1

Beacham, T. D., & Murray, C. B. (1993). Fecundity and egg size variation in North American
Pacific salmon (*Oncorhynchus*). *Journal of Fish Biology,* 42, 485–508.
doi.org/10.1111/j.1095-8649.1993.tb00354.x

Beacham, T. D., Sato, S., Urawa, S., Le, K. D., & Wetklo, M. (2008). Population structure and
stock identification of chum salmon *Oncorhynchus keta* from Japan determined by
microsatellite DNA variation. *Fisheries Science*, 74, 983–994. doi.org/10.1111/j.1444-
2906.2008.01616.x

Beacham, T. D., Candy, J. R., Sato, J.S., Urawa, S., Le, K. D., & Wetklo, M. (2009). Stock
origin of chum salmon (*Oncorhynchus keta*) in the Gulf of Alaska during winter as
estimated with microsatellites. *North Pacific Anadromous Fish Commission Bulletin*, 5,
15–23. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-5/NPAFC_Bull_5_015-
023Beacham.pdf

Beamish, R. J., Ed. (2018). *The Ocean Ecology of Pacific Salmon and Trout*. Bethesda:
American Fisheries Society. doi.org/10.47886/9781934874455

Beauchamp, D. (2009). Bioenergetic ontogeny: Linking climate and mass specific feeding to
life-cycle growth and survival of salmon. In C. Krueger & C. E. Zimmerman (Eds.),
Pacific salmon: Ecology and Management of Western Alaska's Populations
(Symposium 70, pp. 53–72). Bethesda: American Fisheries Society.
doi.org/10.47886/9781934874110

Bryndum-Buchholz, A., Tittensor, D. P., & Lotze, H. K. (2021). The status of climate change
adaptation in fisheries management: Policy, legislation and implementation. *Fish and*
Fisheries, 22, 1248–1273. doi.org/10.1111/faf.12586

Bugaev, A. V., & Gerlits, A. I. (2023). The characteristics of feeding migrations of hatchery-reared Pacific salmon juveniles in the Sea of Okhotsk Basin and adjacent Pacific waters in the autumn–winter period (regional identification, abundance, distribution of catches, biological characteristics, and mortality rate estimates). *Russian Journal of Marine Biology*, 49, 546–568. doi.org/10.1134/S1063074023070039

Burnham, K.P., & Anderson, D.R. (2002). *Model Selection and Multimodel Inference: A Practical Information- Theoretic Approach, 2nd edition*. New York: Springer-Verlag. link.springer.com/book/10.1007/b97636

Caddy, J. F. (2002). Limit reference points, traffic lights, and holistic approaches to fisheries management with minimal stock assessment input. *Fisheries Research*, 5, 133–137. doi.org/10.1016/S0165-7836(01)00343-5

Carvalho, K. S., Smith, T. E., & Wang, S. E. (2021). Bering Sea marine heatwaves: Patterns, trends and connections with the Arctic. *Journal of Hydrology*, 600, 126462. doi.org/10.1016/j.jhydrol.2021.126462

Cheung, W. W., & Frölicher, T. L. (2020). Marine heatwaves exacerbate climate change impacts for fisheries in the northeast Pacific. *Scientific Reports*, 10, 6678. doi.org/10.1038/s41598-020-63650-z

Cheung, W. W., Brodeur, R. D., Okey, T. A., & Pauly, D. (2015). Projecting future changes in distributions of pelagic fish species of Northeast Pacific shelf seas. *Progress in Oceanography*, 130, 19-31. doi.org/10.1016/j.pocean.2014.09.003

Cheung, W. W., Frölicher, T. L., Lam, V. W., Oyinlola, M. A., Reygondeau, G., Sumaila, U. R., ... & Wabnitz, C. C. (2021). Marine high temperature extremes amplify the impacts of climate change on fish and fisheries. *Science Advances*, 7, eabh0895. doi.org/10.1126/sciadv.abh0895

Chitose Aquarium (2023). Capture Information of chum salmon. Chitose Aquarium, Chitose. https://chitose-aq.jp/ (accessed, 8 August 2023).

Christie, M. R., Ford, M. J., & Blouin, M. S. (2014). On the reproductive success of early-generation hatchery fish in the wild. *Evolutionary Applications*, 7, 883–896. doi.org/10.1111/eva.12183

Cline, T. J., Ohlberger, J., & Schindler, D. E. (2019). Effects of warming climate and competition in the ocean for life-histories of Pacific salmon. *Nature Ecology & Evolution*, 3, 935–942. doi.org/10.1038/s41559-019-0901-7

Connors, B., Malick, M. J., Ruggerone, G. T., Rand, P., Adkison, M., Irvine, J. R., ... & Gorman, K. (2020). Climate and competition influence sockeye salmon population dynamics across the Northeast Pacific Ocean. *Canadian Journal of Fisheries and Aquatic Sciences*, 77, 943–949. doi.org/10.1139/cjfas-2019-0422

Connors, B., Ruggerone, G. T., & Irvine, J. R. (2024). Adapting management of Pacific salmon to a warming and more crowded ocean. *ICES Journal of Marine Science*, fsae135. doi.org/10.1093/icesjms/fsae135

Crozier, L. G., & Siegel, J. E. A. (2023). Comprehensive review of the impacts of climate change on salmon: strengths and weaknesses of the literature by life stage. *Fishes*, 8, 1–50. doi.org/10.3390/fishes8060319

Crozier, L. G., Burke, B. J., Chasco, B. E., Widener, D. L., & Zabel, R. W. (2021). Climate change threatens Chinook salmon throughout their life cycle. *Communications Biology*, 4, 222. doi.org/10.1038/s42003-021-01734-w

Cunningham, C. J., Westley, P. A., & Adkison, M. D. (2018). Signals of large scale climate drivers, hatchery enhancement, and marine factors in Yukon River Chinook salmon survival revealed with a Bayesian life history model. *Global Change Biology*, 24,

1044   4399–4416. doi.org/10.1111/gcb.14315

Dahlke, F. T., Butzin, M., Nahrgang, J., Puvanendran, V., Mortensen, A., Pörtner, H. O., &
            Storch, D. (2018). Northern cod species face spawning habitat losses if global
            warming exceeds 1.5 C. *Science Advances*, 4, eaas8821.
            doi.org/10.1126/sciadv.aas8821

Dann, T. H., Hoyt, H. A., Lee, E. M., Fox, E. K. C., & Foster, M. B. (2023). Genetic stock
            composition of chum salmon harvested in commercial salmon fisheries of the South
            Alaska Peninsula, 2022. *Alaska Department of Fish and Game, Special Publication*,
            23-07. www.adfg.alaska.gov/FedAidPDFs/SP23-07.pdf

Einum, S., & Fleming, I. A. (1999). Maternal effects of egg size in brown trout (*Salmo trutta*):
            Norms of reaction to environmental quality. *Proceedings of the Royal Society of
            London. Series B: Biological Sciences*, 266, 2095–2100.
            doi.org/10.1098/rspb.1999.0893

Einum, S., & Fleming, I. A. (2000). Highly fecund mothers sacrifice offspring survival to
            maximize fitness. *Nature*, 405, 565–567. doi.org/10.1038/35014600FAO (United
            Nations Food and Agriculture Organization) (2022). *The State of World Fisheries and
            Aquaculture: 2022: Towards Blue Transformation*. Rome: FAO.
            digitallibrary.un.org/record/3978694?v=pdf

FAO (United Nations Food and Agriculture Organization) (2022). *The State of World
            Fisheries and Aquaculture: 2022: Towards Blue Transformation*. Rome: FAO.
            digitallibrary.un.org/record/3978694?v=pdf

Farley Jr, E. V., Yasumiishi, E. M., Murphy, J. M., Strasburger, W., Sewall, F., Howard, K., ...
            & Moss, J. H. (2024). Critical periods in the marine life history of juvenile western
            Alaska chum salmon in a changing climate. *Marine Ecology Progress Series*, 726, 149–
            160. https://doi.org/10.3354/meps14491

Fisheries Agency of Japan (1955). On the salmon in waters adjacent to Japan. *International
            North Pacific Fisheries Commission Bulletin*, 1–2. www.npafc.org/wp-
            content/uploads/Bulletin1FRA.pdf

Fleming, I. A., & Gross, M. R. (1990). Latitudinal clines: A trade-off between egg number and
            size in Pacific salmon. *Ecology*, 71, 1–11. doi.org/10.2307/1940241

FRA (Japan Fisheries Research and Education Agency) (2023). *Salmon Data Base.*
            Yokohama: FRA. (in Japanese) www.fra.go.jp/shigen/salmon/sdb.html (accessed 20
            November 2024).

FRA (Japan Fisheries Research and Education Agency) (2024). *Chum salmon catch and
            release statistics.* Yokohama: FRA. (in Japanese)
            www.fra.go.jp/shigen/salmon/index.html (accessed 5 March 2024).

Free, C. M., Thorson, J. T., Pinsky, M. L., Oken, K. L., Wiedenmann, J., & Jensen, O. P.
            (2019). Impacts of historical warming on marine fisheries production. *Science*, 363,
            979–983. doi.org/10.1126/science.aau1758

Frölicher, T. L., Fischer, E. M., & Gruber, N. (2018). Marine heatwaves under global
            warming. *Nature*, 560, 360–364. doi.org/10.1038/s41586-018-0383-9

Frost, T. J., Yasumiishi, E. M., Agler, B. A., Adkison, M. D., & McPhee, M. V. (2021).
            Density-dependent effects of eastern Kamchatka pink salmon (*Oncorhynchus
            gorbuscha*) and Japanese chum salmon (*O. keta*) on age-specific growth of western
            Alaska chum salmon. *Fisheries Oceanography*, 30, 99-109.
            doi.org/10.1111/fog.12505

Fukuwaka, M., & Suzuki, T. (1998). Role of a riverine plume as a nursery area for chum
            salmon *Oncorhynchus keta*. *Marine Ecology Progress Series*,173, 289–297.

doi.org/10.3354/meps173289

Graham, C., Pakhomov, E. A., & Hunt, B. P. (2021). Meta-analysis of salmon trophic ecology
reveals spatial and interspecies dynamics across the North Pacific Ocean. *Frontiers in*
Marine Science, 8, 618884. doi.org/10.3389/fmars.2021.618884

Groot, C., & Margolis, L. (1991). *Pacific Salmon Life Histories*. Vancouver: University of
British Columbia Press. www.ubcpress.ca/pacific-salmon-life-histories

Hasegawa, K., Okamoto, Y., & Morita, K. (2021a). Temporal trends in geographic clines of
chum salmon reproductive traits associated with global warming and hatchery
programmes. *Ecological Solutions and Evidence*, 2, e12107. doi.org/10.1002/2688-
8319.12107

Hasegawa, K., Honda, K., Yoshiyama, T., Suzuki, K., & Fukui, S. (2021b). Small biased body
size of salmon fry preyed upon by piscivorous fish in riverine and marine habitats.
Canadian Journal of Fisheries and Aquatic Sciences, 78, 631–638.
doi.org/10.1139/cjfas-2020-0339

Hauri, C., Pages, R., Hedstrom, K., Doney, S. C., Dupont, S., Ferriss, B., & Stuecker, M. F.
(2024). More than marine heatwaves: A new regime of heat, acidity, and low oxygen
compound extreme events in the Gulf of Alaska. *AGU Advances*, 5, e2023AV001039.

Hayashi, M., Shiogama, H., & Ogura, T. (2022). The contribution of climate change to
increasing extreme ocean warming around Japan. *Geophysical Research Letters*, 49,
e2022GL100785. doi.org/10.1029/2022GL100785

Hilborn, R., & Sibert, J. (1988). Adaptive management of developing fisheries. *Marine*
Policy, 12, 112–121. doi.org/10.1016/0308-597X(88)90039-5

Hilborn, R., Quinn, T. P., Schindler, D. E., & Rogers, D. E. (2003). Biocomplexity and
fisheries sustainability. *Proceedings of the National Academy of Sciences*, 100, 6564–
6568. doi.org/10.1073/pnas.1037274100

Hiroi, O. (1998). Historical trends of salmon fisheries and stock conditions in Japan. *North*
Pacific Anadromous Fish Commission Bulletin, 1, 23–27. www.npafc.org/wp-
content/uploads/Bulletins/Bulletin-1/page-23-27Hiroi.pdf

Honda, K., Kawakami, T., Suzuki, K., Watanabe, K., & Saito, T. (2017). Growth rate
characteristics of juvenile chum salmon *Oncorhynchus keta* originating from the Pacific
coast of Japan and reaching Konbumori, eastern Hokkaido. *Fisheries Science*, 83, 987–
996. doi.org/10.1007/s12562-017-1137-6

Honda, K., Sato, T., Kuroda, H., & Saito, T. (2021). Initial growth characteristics of poor-
return stocks of chum salmon *Oncorhynchus keta* originating from the Okhotsk and
Nemuro regions in Hokkaido on the basis of scale analysis. *Fisheries Science*, 87, 653–
663. doi.org/10.1007/s12562-021-01538-1

Honda, K., Shirai, K., Morishita, T., & Saito, T. (2025). Characteristics of juvenile chum
salmon carrying out inverse migrations after ocean entry from rivers along the Pacific
coast of Hokkaido, Japan. *Fisheries Oceanography*, 34, e12698.
doi.org/10.1111/fog.12698

Hoshino, N., & Fujioka, T. (2021). Characteristics of yellowtail catch in Hokkaido, Japan in
the 2010s. *Scientific Reports of Hokkaido Fisheries Research Institutes*, 100, 71–82. (in
Japanese)

HSEA (Hokkaido Salmon Enhancement Association) (2024). Chum salmon catch and release
statistics, Sapporo: HSEA. (in Japanese) www.sake-masu.or.jp (accessed 31 January
2024).

Hunsicker, M. E., Boldt, J. L., Blasiak, R., Hazen, E. L., Kulik, V. V., Ryu, J., ... & Wakita, K.,
Eds. (2022). Report of Working Group 36 on Common Ecosystem Reference Points

across PICES Member Countries. *PICES Scientific Report*, (64), 155 pp. meetings.pices.int/publications/scientific-reports/Report64/Rpt64.pdf

Hutchings, J. A. (1991). Fitness consequences of variation in egg size and food abundance in brook trout *Salvelinus fontinalis*. *Evolution*, 45, 1162–1168. doi.org/10.1111/j.1558-5646.1991.tb04382.x

Iida, M. (2024). Relationship between in-river return rate of hatchery-origin chum salmon Oncorhynchus keta and coastal water temperature and body weight at release on the Japan Sea side of Honshu, Japan. *Environmental Biology of Fishes*, 107, 1247–1261. doi.org/10.1007/s10641-024-01633-z

Iida, M., Yoshino, K., & Katayama, S. (2018). Current status of natural spawning of chum salmon *Oncorhynchus keta* in rivers with or without hatchery stocking on the Japan Sea side of northern Honshu, Japan. *Fisheries science*, 84, 453–459. doi.org/10.1007/s12562-018-1192-7

Iida, M., Yagi, Y., & Iseki, T. (2021). Occurrence of wild chum salmon fry in the surf zone, and spawning and emergence timing in the adjacent nonstocked river in Niigata Prefecture, Japan. *Fisheries Science*, 87, 549–557. doi.org/10.1007/s12562-021-01535-4

Iino, Y., Kitagawa, T., Abe, T. K., Nagasaka, T., Shimizu, Y., Ota, K., ... & Kawamura, T. (2022). Effect of food amount and temperature on growth rate and aerobic scope of juvenile chum salmon. *Fisheries Science*, 88, 397–409. doi.org/10.1007/s12562-022-01599-w

Iino, Y., Abe, T. K., Shimizu, Y., Nagasaka, T., & Kitagawa, T. (2024). Body size- and temperature-related metabolic traits of juvenile chum salmon during northward migration. *Canadian Journal of Fisheries and Aquatic Sciences*, 81, 1514–1527. doi.org/10.1139/cjfas-2023-0334

James, S. E., Doherty, B., Cox, S. P., Pearsall, I. A., & Riddell, B. (2023). Size and timing of hatchery releases influence juvenile-to-adult survival rates of British Columbia Chinook and coho salmon. *Canadian Journal of Fisheries and Aquatic Sciences*, 80, 700–718. doi.org/10.1139/cjfas-2022-0121

Jeffrey, K. M., Côté, I. M., Irvine, J. R., & Reynolds, J. D. (2017). Changes in body size of Canadian Pacific salmon over six decades. *Canadian Journal of Fisheries and Aquatic Sciences*, 74, 191–201. doi.org/10.1139/cjfas-2015-0600

JMA (Japan Meteorological Agency) (2022). *Coastal sea surface temperature information*. Tokyo: JMA. (in Japanese) www.jma-net.go.jp (accessed 23 August 2022).

Kaeriyama, M. (1989). Aspects of salmon ranching in Japan. *Physiology & Ecology, Japan*, Special Volume 1, 625–638. www.researchgate.net/publication/303362061

Kaeriyama, M. (1999). Hatchery programmes and stock management of salmonid populations in Japan. In B. R. Howell, E. Moksness, & T. Svåsand (Eds.), *Stock Enhancement and Sea Ranching* (pp. 153–167). Oxford: Blackwell. www.researchgate.net/publication/246360103

Kaeriyama, M. (2008). Ecosystem-based sustainable conservation and management of Pacific salmon. In K. Tsukamoto, T. Kawamura, T. Takeuchi, T. D. Bard, Jr., & M.J. Kaiser (Eds.), *Fisheries for Global Welfare and Environment* (pp. 371–380). Tokyo: Terrapub. wcfs.fisheries.org/proceedings-of-the-fifth-world-fisheries-congress/

Kaeriyama, M. (2022). Warming climate impacts on production dynamics of southern populations of Pacific salmon in the North Pacific Ocean. *Fisheries Oceanography*, 32, 121–132. doi.org/10.1111/fog.12598

Kaeriyama, M., & Sakaguchi, I. (2023). Ecosystem-based sustainable management of chum

salmon in Japan's warming climate. *Marine Policy*, 157, 105842. doi.org/10.1016/j.marpol.2023.105842

Kaeriyama, M., Seo, H., Kudo, H., & Nagata, M. (2012). Perspectives on wild and hatchery salmon interactions at sea, potential climate effects on Japanese chum salmon, and the need for sustainable salmon fishery management reform in Japan. *Environmental Biology of Fishes*, 94, 165–177. doi.org/10.1007/s10641-011-9930-z

Kasugai, K., Saneyoshi, H., Aoyama, T., Shinriki, Y., Iijima, A., & Miyakoshi, Y. (2016). Early marine migration of juvenile chum salmon along the Pacific coast of eastern Hokkaido. *North Pacific Anadromous Fish Commission Bulletin*, 6, 61–72. doi.org/10.23849/npafcb6/61.72

Kitada, S. (2014). Japanese chum salmon stock enhancement: Current perspective and future challenges. *Fisheries Science*, 80, 237–249. doi.org/10.1007/s12562-013-0692-8

Kitada, S. (2018). Economic, ecological and genetic impacts of marine stock enhancement and sea ranching: A systematic review. *Fish and Fisheries*, 19, 511–532. doi.org/10.1111/faf.12271

Kitada, S. (2020). Lessons from Japan marine stock enhancement and sea ranching programmes over 100 years. *Reviews in Aquaculture*, 12, 1944–1969. doi.org/10.1111/raq.12418

Kitada, S., Kishino, H., & Hamasaki, K. (2011). Bias and significance of relative reproductive success estimates based on steelhead trout (*Oncorhynchus mykiss*) data: A Bayesian meta-analysis. *Canadian Journal of Fisheries and Aquatic Sciences*, 68, 1827–1835. doi.org/10.1139/f2011-087

Kobayashi, T. (1980). Salmon propagation in Japan. In J.E. Thorpe (Ed.), *Salmon Ranching* (pp. 91–107). London: Academic Press. archive.org/details/salmonranching0000unse/page/446/mode/2up

Kraskura, K., Patterson, D. A., & Eliason, E. J. (2024). A review of adult salmon maximum swim performance. *Canadian Journal of Fisheries and Aquatic Sciences*, 81, 1174–1216. doi.org/10.1139/cjfas-2023-0246

Kuroda, H., Saito, T., Kaga, T., Takasuka, A., Kamimura, Y., Furuichi, S., & Nakanowatari, T. (2020). Unconventional sea surface temperature regime around Japan in the 2000s-2010s: Potential influences on major fisheries resources, *Frontiers in Marine Science*, 7, 574904. doi.org/10.3389/fmars.2020.574904

Langan, J. A., Cunningham, C. J., Watson, J. T., & McKinnell, S. (2024). Opening the black box: New insights into the role of temperature in the marine distributions of Pacific salmon. *Fish and Fisheries*. doi.org/10.1111/faf.12825

Larson, W. A., Seeb, J. E., Dann, T. H., Schindler, D. E., & Seeb, L. W. (2014). Signals of heterogeneous selection at an MHC locus in geographically proximate ecotypes of sockeye salmon. *Molecular Ecology*, 23, 5448–5461. doi.org/10.1111/mec.12949

Lewis, B., Grant, W. S., Brenner, R. E., & Hamazaki, T. (2015). Changes in size and age of Chinook salmon *Oncorhynchus tshawytscha* returning to Alaska. *Plos One*, 10, e0130184. doi.org/10.1371/journal.pone.0130184

Litzow, M. A., Ciannelli, L., Puerta, P., Wettstein, J. J., Rykaczewski, R. R., & Opiekun, M. (2018). Non-stationary climate–salmon relationships in the Gulf of Alaska. *Proceedings of the Royal Society B*, 285, 20181855. doi.org/10.1098/rspb.2018.1855

Losee, J. P., Kendall, N. W., & Dufault, A. (2019). Changing salmon: An analysis of body mass, abundance, survival, and productivity trends across 45 years in Puget Sound. *Fish and Fisheries*, 20, 934–951. doi.org/10.1111/faf.12385

MAFF (Ministry of Agriculture, Forestry and Fisheries) (2023). *Annual statistics of fisheries*

and aquaculture production. Tokyo: MAFF. (in Japanese)
www.maff.go.jp/j/tokei/kouhyou/kaimen_gyosei/index.html (accessed 11 January 2023)

Malick, M. J., Losee, J. P., Marston, G., Agha, M., Berejikian, B. A., Beckman, B. R., & Cooper, M. (2023). Fecundity trends of Chinook salmon in the Pacific Northwest. *Fish and Fisheries*, 24, 454–465. doi.org/10.1111/faf.12738

Masson-Delmotte, V., Zhai, P., Pörtner, H. O., Roberts, D., Skea, J., & Shukla, P. R. (2022). *Global Warming of 1.5 C*: IPCC special report on impacts of global warming of 1.5 C above pre-industrial levels in context of strengthening response to climate change, sustainable development, and efforts to eradicate poverty. Cornwall: TJ Books Limited. www.ipcc.ch/sr15/

Max Planck Institute (2023). *Climate Data Operators*. Hamburg: Max Planck Institute für Meteorologie. code.mpimet.mpg.de/projects/cdo (accessed 14 June 2023).

Mayama, H., & Ishida, Y. (2003). Japanese Studies on the early ocean life of juvenile salmon. *North Pacific Anadromous Fish Commission Bulletin*, 3, 41–67. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-3/Pages-41-68-JapanReview.pdf

Miyagi Prefectural Government. (2022) Commercial fishing landings in four major ports. (in Japanese) www.pref.miyagi.jp/soshiki/suishin/mizuage.html (accessed 16 October 2022)

Miyakoshi, Y., Urabe, H., Saneyoshi, H., Aoyama, T., Sakamoto, H., Ando, D., ... & Nagata, M. (2012). The occurrence and run timing of naturally spawning chum salmon in northern Japan. *Environmental Biology of Fishes*, 94, 197–206. doi.org/10.1007/s10641-011-9872-5

Miyakoshi, Y., Nagata, M., Kitada, S., & Kaeriyama, M. (2013). Historical and current hatchery programs and management of chum salmon in Hokkaido, northern Japan. *Reviews in Fisheries Science*, 21, 469–479. doi.org/10.1080/10641262.2013.836446

Mousseau, T. A., & Fox, C. W. (1998) The adaptive significance of maternal effects. *Trends in Ecology & Evolution*, 13, 403–407. doi.org/10.1016/S0169-5347(98)01472-4

Myers, K. W., Klovach, N. V., Gritsenko, O. F., Urawa, S., & Royer, T. C. (2007). Stock-specific distributions of Asian and North American salmon in the open ocean, interannual changes, and oceanographic conditions. *North Pacific Anadromous Fish Commission Bulletin*, 4, 159–177. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-4/159-177Myers.pdf

Myers, K. W., Irvine, J., Logerwell, E., Urawa, S., Naydenko, S., Zavolokin, A., & Davis, N. (2016). Pacific salmon and steelhead: Life in a changing winter ocean. *North Pacific Anadromous Fish Commission Bulletin*, 6, 113–138. doi.org/10.23849/npafcb6/113.138

Nagasawa, T., & Azumaya, T. (2009). Distribution and CPUE trends of Pacific salmon, especially in sockeye salmon in the Bering Sea and adjacent waters from 1972 to mid 2000's. *North Pacific Anadromous Fish Commission Bulletin*, 5, 1–13. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-5/NPAFC_Bull_5_001-013Nagasawa.pdf

Nagata, M., Miyakoshi, Y., Ando, D., Fujiwara, M., Sawada, M., Shimada, H., & Asami, H. (2007). Influence of coastal seawater temperature on the distribution and growth of juvenile chum salmon, with recommendations for altered release strategies. *North Pacific Anadromous Fish Commission Bulletin*, 4, 223–235. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-4/223-235Nagatachum.pdf

Nagata, M., Miyakoshi, Y., Fujiwara, M., Kasugai, K., Ando, D., Torao, M., ... & Irvine, J. R. (2016). Adapting Hokkaido hatchery strategies to regional ocean conditions can improve chum salmon survival and reduce variability. *North Pacific Anadromous Fish*

Commission Bulletin, 6, 73-85. doi.org/10.23849/npafcb6/73.85

Naish, K. A., Taylor III, J. E., Levin, P. S., Quinn, T. P., Winton, J. R., Huppert, D., & Hilborn, R. (2007). An evaluation of the effects of conservation and fishery enhancement hatcheries on wild populations of salmon. *Advances in Marine Biology*, 53, 61–194. doi.org/10.1016/S0065-2881(07)53002-6

Naydenko, S. V., & Temnykh, O. S. (2016). The survival of Pacific salmon in the North Pacific in the winter and spring. *Russian Journal of Marine Biology*, 42, 527–550. doi.org/10.1134/S1063074016070087

Naydenko, S. V., Temnykh, O. S., & Figurkin, A. L. (2016). Is winter the critical period in the marine life history of Pacific salmon? *North Pacific Anadromous Fish Commission Bulletin*, 6, 139-152. doi.org/10.23849/npafcb6/139.152

Neave, F., Yonemori, T., & Bakkala, R. G. (1976). Distribution and origin of chum salmon in offshore waters of the North Pacific Ocean. *International North Pacific Fisheries Commission Bulletin*, 35. www.npafc.org/wp-content/uploads/Bulletin35.pdf

NOAA (National Oceanic and Atmospheric Administration, Physical Sciences Laboratory) (2023a). *NOAA Optimum Interpolation SST V2 high-resolution dataset (OISST V2)*. Boulder: NOAA. psl.noaa.gov/data/gridded/data.noaa.oisst.v2.highres.html (accessed 14 June 2023).

NOAA (National Oceanic and Atmospheric Administration, Physical Sciences Laboratory) (2023b). *Plot: COBE sea surface temperature*. Boulder: NOAA. psl.noaa.gov/mddb2/makePlot.html?variableID=1570 (accessed 18 March 2024).

NOAA (National Oceanic and Atmospheric Administration, Northwest Fisheries Science Center) (2024). *Ocean ecosystem indicators of Pacific salmon marine survival in the Northern California Current*. Seattle: NOAA. www.fisheries.noaa.gov/west-coast/science-data/ocean-ecosystem-indicators-pacific-salmon-marine-survival-northern (accessed 9 November 2024).

NPAFC (North Pacific Anadromous Fish Commission) (2023). *NPAFC statistics: Pacific salmonid catch and hatchery release data*. Vancouver: North Pacific Anadromous Fish Commission. www.npafc.org/statistics/ (accessed 7 August 2023).

NPFMC (North Pacific Fishery Management Council) (2025). *Salmon bycatch*. North Pacific Fishery Management Council. www.npfmc.org/fisheries-issues/bycatch/salmon-bycatch/ (accessed 7 April 2025).

Ohlberger, J., Ward, E. J., Brenner, R. E., Hunsicker, M. E., Haught, S. B., Finnoff, D., ... & Hauri, C. (2022). Non-stationary and interactive effects of climate and competition on pink salmon productivity. *Global Change Biology*, 28, 2026–2040. doi.org/10.1111/gcb.16049

Ohlberger, J., Cline, T. J., Schindler, D. E., & Lewis, B. (2023). Declines in body size of sockeye salmon associated with increased competition in the ocean. *Proceedings of the Royal Society B*, 290, 20222248. doi.org/10.1098/rspb.2022.2248

Okado, J., Koshino, Y., Kudo, H., & Watanuki, Y. (2020). Consumption of juvenile chum salmon by a seabird species during early sea life. *Fisheries Research*, 222, 105415. doi.org/10.1016/j.fishres.2019.105415

Okado, J., & Hasegawa, K. (2024). Exploring predators of Pacific salmon throughout their life history: The case of Japanese chum, pink, and masu salmon. *Reviews in Fish Biology and Fisheries*, 34, 895–917. doi.org/10.1007/s11160-024-09858-y

Oke, K. B., Cunningham, C. J., Westley, P. A. H., Baskett, M. L., Carlson, S. M., Clark, J., ... & Palkovacs, E. P. (2020). Recent declines in salmon body size impact ecosystems and fisheries. *Nature Communications*, 11, 4155. doi.org/10.1038/s41467-020-17726-z

PICES–NPFC (North Pacific Marine Science Organization, PICES, – North Pacific Fisheries Commission, NPFC) (2019). NPFC–PICES Framework for Enhanced Scientific Collaboration in the North Pacific. *North Pacific Fisheries Commission*, Document number NPFC-2019-SC04-WP02. www.npfc.int/npfc-pices-framework-enhanced-scientific-collaboration-north-pacific

Pauly, D., Hilborn, R., & Branch, T. A. (2013). Fisheries: Does catch reflect abundance?. *Nature*, 494(7437), 303-306. doi.org/10.1038/494303a

Powers, D. A., Lauerman, T., Crawford, D., & DiMichele, L. (1991). Genetic mechanisms for adapting to a changing environment. *Annual Review of Genetics*, 25, 629–660. doi.org/10.1146/annurev.ge.25.120191.003213

Punt, A. E. (2023). Those who fail to learn from history are condemned to repeat it: A perspective on current stock assessment good practices and the consequences of not following them. *Fisheries Research*, 261, 106642. doi.org/10.1016/j.fishres.2023.106642

Quinn, T. P. (2018). *The behavior and ecology of Pacific salmon and trout, 2nd edition*. Seattle: University of Washington Press. uwapress.uw.edu/book/9780295743332/the-behavior-and-ecology-of-pacific-salmon-and-trout/

Quinn, T. P., Vøllestad, L. A., Peterson, J., & Gallucci, V. (2004). Influences of freshwater and marine growth on the egg size-egg number tradeoff in coho and Chinook salmon. *Transactions of the American Fisheries Society*, 133, 55–65. doi.org/10.1577/T03-032

R Core Team (2024). R: A language and environment for statistical computing. *R Foundation for Statistical Computing*. www.R-project.org/

Robins, C. R., Bailey, R.M., Bond, C. E., Brooker, J. R., Lacher, E. A., Lea, R. N., & Scott, W. B. (1991). *World fishes important to North Americans, exclusive of species from the continental waters of the United States and Canada*. American Fisheries Society Special Publication, 21. Bethesda: American Fisheries Society. fisheries.org/bookstore/all-titles/special-publications/x51017pxm/

Ruggerone, G. T., Agler, B. A., & Nielsen, J. L. (2011). Evidence for competition at sea between Norton Sound chum salmon and Asian hatchery chum salmon. *Environmental Biology of Fishes*, 94, 149–163. doi.org/10.1007/s10641-011-9856-5

Ruggerone, G. T., & Irvine, J. R. (2018). Numbers and biomass of natural-and hatchery-origin pink salmon, chum salmon, and sockeye salmon in the North Pacific Ocean, 1925–2015. *Marine and Coastal Fisheries*, 10, 152–168. doi.org/10.1002/mcf2.10023

Ruggerone, G. T., Springer, A. M., van Vliet, G. B., Connors, B., Irvine, J. R., Shaul, L. D., ... & Atlas, W. I. (2023). From diatoms to killer whales: Impacts of pink salmon on North Pacific ecosystems. *Marine Ecology Progress Series*, 719, 1–40. doi.org/10.3354/meps14402

Saito, T. (2022). Optimum timing and size at release of chum salmon: Improving survival by modifying hatchery practices. *Fisheries Oceanography*, 31, 416–428. doi.org/10.1111/fog.12585

Saito, T., & Nagasawa, K. (2009). Regional synchrony in return rates of chum salmon (*Oncorhynchus keta*) in Japan in relation to coastal temperature and size at release. *Fisheries Research*, 95, 14–27. doi.org/10.1016/j.fishres.2008.07.004

Saito, T., Honda, K., Sasaki, K., Watanabe, K., Suzuki, K., Hirabayashi, Y., ... & Sato, S. (2020). Stock composition of adult chum salmon *Oncorhynchus keta* caught in a setnet fishery estimated using genetic identification, scale patterns, and otolith thermal marking. Fisheries Science, 86, 271–286. doi.org/10.1007/s12562-020-01401-9

Saitou, N., & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing

phylogenetic trees. *Molecular Biology and Evolution*, 4, 406–425. doi.org/10.1093/oxfordjournals.molbev.a040454

Salo, E.O. (1991). Life history of chum salmon (*Oncorhynchus keta*). In C. Groot, & L. Margolis (Eds.) *Pacific Salmon Life Histories* (pp. 231–310). Vancouver: University of British Columbia Press. www.ubcpress.ca/pacific-salmon-life-histories

Sano, S. (1966). Chum Salmon in the Far East. *International North Pacific Fisheries Commission Bulletin*, 18-3, 41–57. www.npafc.org/wp-content/uploads/Bulletin18-3.pdf

Sano, S. (1967). Chum salmon in the Far East. *International North Pacific Fisheries Commission Bulletin*, 23–3: 23-41. www.npafc.org/wp-content/uploads/Bulletin23-3.pdf

Shedd, K. R., Lescak, E. A., Habicht, C., Knudsen, E. E., Dann, T. H., Hoyt, H. A., … & Templin, W. D. (2022). Reduced relative fitness in hatchery-origin pink salmon in two streams in Prince William Sound, Alaska. *Evolutionary Applications*, 15, 429–446. doi.org/10.1111/eva.13356

Shepard, M. P., Hartt, A. C., & Yonemori, T. (1968). Chum salmon in offshore waters. *International North Pacific Fisheries Commission Bulletin*, 25. www.npafc.org/wp-content/uploads/Bulletin-25.pdf

Shubin, A. O., & Akinicheva, E. G. (2016). Origin of juvenile chum salmon *Oncorhynchus keta* (Salmonidae) in the Sea of Okhotsk coastal waters off south Sakhalin. *Journal of Ichthyology*, 56, 728–737. doi.org/10.1134/S0032945216050131

Somero, G. N. (2022). The Goldilocks principle: A unifying perspective on biochemical adaptation to abiotic stressors in the sea. *Annual Review of Marine Science*, 14, 1–23. doi.org/10.1146/annurev-marine-022521-102228

Springer, A. M., & van Vliet, G. B. (2014). Climate change, pink salmon, and the nexus between bottom-up and top-down forcing in the subarctic Pacific Ocean and Bering Sea. *Proceedings of the National Academy of Sciences*, 111, E1880–E1888. doi.org/10.1073/pnas.1319089111

Sugiura, N. (1978). Further analysis of the data by Akaike's information criterion and the finite corrections. *Communications in Statistics - Theory and Methods*, 7, 13–26. doi.org/10.1080/03610927808827599

Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: Molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution*, 38, 3022–3027. doi.org/10.1093/molbev/msab120

Thorn, M. W., Dick, M. F., Oviedo, L., Guglielmo, C. G., & Morbey, Y. E. (2019). Transgenerational effects of egg nutrients on the early development of Chinook salmon (*Oncorhynchus tshawytscha*) across a thermal gradient. *Canadian Journal of Fisheries and Aquatic Sciences*, 76, 1253–1262. doi.org/10.1139/cjfas-2018-0013

Thorson, J. T., Maunder, M. N., & Punt, E. (2020). The development of spatio-temporal models of fishery catch-per-unit-effort data to derive indices of relative abundance. *Fisheries Research*, 230, 105611. doi.org/10.1016/j.fishres.2020.105611

Tian, Y., Kidokoro, H., Watanabe, T., Igeta, Y., Sakaji, H., & Ino, S. (2012). Response of yellowtail, *Seriola quinqueradiata*, a key large predatory fish in the Japan Sea, to sea water temperature over the last century and potential effects of global warming. *Journal of Marine Systems*, 91, 1–10. doi.org/10.1016/j.jmarsys.2011.09.002

Urawa, S., & Bugaev, A. V. (2021). Survival of Japanese chum salmon during early ocean life in 2011–2017. *North Pacific Anadromous Fish Commission Technical Report*, 17, 60–62. doi.org/10.23849/npafctr17

Urawa, S., Seki, J., Kawana, M., Saito, T., Crane, P. A., Seeb, L. W., Fukuwaka, M., & Akinicheva, E. (2006). Origins of juvenile chum salmon caught in the southwestern Okhotsk Sea during the fall of 2000. *Bulletin of National Salmon Resources Center*, 8, 9–16. salmon.fra.affrc.go.jp/kankobutu/bulletin/BNSRC08-2.pdf

Urawa, S., Sato, S., Crane, P. A., Agler, B., Josephson, R., & Azumaya, T. (2009). Stock-specific ocean distribution and migration of chum salmon in the Bering Sea and North Pacific Ocean. *North Pacific Anadromous Fish Commission Bulletin*, 5:131–146. www.npafc.org/wp-content/uploads/Bulletins/Bulletin-5/NPAFC_Bull_5_131-146Urawa.pdf

Urawa, S., Beacham, T., Sato, S., Kaga, T., Agler, B. A., Josephson, R., & Fukuwaka, M. (2016). Stock-specific abundance of chum salmon in the central Gulf of Alaska during winter. *North Pacific Anadromous Fish Commission Bulletin*, 6, 153–160. doi.org/10.23849/npafcb6/153.160

Urawa, S., Beacham, T. D., Fukuwaka, M., & Kaeriyama, M. (2018). Ocean ecology of chum salmon. In R. J. Beamish (Ed.), *The Ocean Ecology of Pacific Salmon and Trout* (pp. 161–317). Bethesda: American Fisheries Society. doi.org/10.47886/9781934874455.ch3

Urawa, S., Beacham, T., Sutherland, B., & Sato, S. (2022). Winter distribution of chum salmon in the Gulf of Alaska: A review. *North Pacific Anadromous Fish Commission Technical Report*, 18, 83–87. doi.org/10.23849/npafctr18

Walters, C. J., & Hilborn, R. (1978). Ecological optimization and adaptive management. *Annual review of Ecology and Systematics*, 9, 157–188. www.jstor.org/stable/2096747

Wilkinson, L., & Task Force on Statistical Inference, American Psychological Association, Science Directorate (1999). Statistical methods in psychology journals: Guidelines and explanations. *American Psychologist*, 54, 594–604. doi.org/10.1037/0003-066X.54.8.594

Wood, S. N. (2004). Stable and efficient multiple smoothing parameter estimation for generalized additive models. *Journal of the American Statistical Association*, 99, 673–686. doi.org/10.1198/016214504000000980

Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society B*, 73, 3–36. doi.org/10.1111/j.1467-9868.2010.00749.x

Wood, S. N. (2017). *Generalized additive models*. New York: Chapman and Hall/CRC. doi.org/10.1201/9781315370279

Yagisawa, Y. (1970). Salmon hatchery enhancement program on Etorofu Island, Fish and eggs, *Salmon Management Center Technical Report*, 133, 35–79. (in Japanese) salmon.fra.affrc.go.jp/kankobutu/tech_repo/fe04.htm

Yonemori, T. (1975). A trial analysis of the results obtained from tagging on chum salmon origination from Hokkaido. *International North Pacific Fisheries Commission Bulletin*, 32, 130–160. www.npafc.org/wp-content/uploads/Bulletin-32.pdf

Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid common statistical problems. *Methods in Ecology and Evolution*, 1, 3–14. doi.org/10.1111/j.2041-210X.2009.00001.x

1471
1472
FIGURE 1 Japanese chum salmon management regions, coastal migration routes, and high-
seas distribution areas. (a) Seven management regions (delineated by black bars) and
hatcheries (black dots) in Japan. Migration paths of hatchery-released fry were modified from
Saito and Nagasawa (2009, Figure 1). Schematic sea surface currents were redrawn from
Kuroda et al. (2020, Figure 1). (b) principal areas of high-seas distribution (boxes), month(s),
age group(s), and seasonal direction of wintering, feeding and homing migrations (arrows),
numbered areas are: 1.Southern Sea of Okhotsk (45.2°N–46.5°N, 142.5°E–148.5°E), 2.
Western North Pacific (42°N–45°N, 158°E–165°E), 3. Central Bering Sea (53°N–58°N,
175°E–170°W), 4. Central Gulf of Alaska (48°N–56°N, 140°W–150°W). Migration routes of
Japanese chum salmon estimated from long-term field surveys, modified from Myers et al.
(2007, Figure 1) and Urawa et al. (2018, Figure 37).

1484

1485 **FIGURE 2** Number of hatchery chum salmon fry released (red line) and adult returns (coastal
1486 catch plus hatchery broodstock catch; black line) in seven management regions in Japan (1970–
1487 2023; Figure 1a).

1488
1489 **FIGURE 3** Neighbor-joining tree based on correlation coefficients of chum salmon adult return
1490 rates between pairs of seven management regions in Japan, 1998–2019 (Figure 1a).

FIGURE 4 Sea surface temperature (SST) isotherms in recent decades (1990-2022) showed a northward shift and longitudinal contraction of chum salmon summer (July) thermal habitat in the North Pacific Ocean, while winter (January) habitat was relatively stable. Blue, green and orange show the range of Japanese chum salmon thermal limits (4–13℃): optimal growth and feeding habitat (8–12℃), swimming and feeding habitat (5–13℃), and overwintering habitat (4–6℃) (Kaeriyama et al., 2012; Urawa et al., 2018). Red (>13℃) and purple (<4℃) indicate critical habitat, lost to ocean production of Japanese chum salmon.

1499

1500 **FIGURE 5** Northward shifts of two marine piscivorous fishes in Japan. (a) Principal
1501 component analysis biplot based on catch statistics of 30 marine fish species (1980 - 2020)
1502 showed increasing catches of yellowtail and Japanese Spanish mackerel. (b) Two-decadal shifts
1503 in commercial catches of chum salmon and two piscivores.

1504
1505 **FIGURE 6** Egg size of age–4 chum salmon has generally decreased in all regions, except
1506 Hokkaido Okhotsk as shown by the nonlinear and/or linear year effect from GAMs (1998-
1507 2019). The trend was significant in all regions, except Honshu Sea of Japan (Table S1). Open
1508 circles show observed values. The black lines show predicted values with shaded areas of
1509 2×standard errors.

FIGURE 7 Standardized effects of hypothesized explanatory variables on Japanese chum salmon marine survival (adult return rates across regions). Regression coefficients of best regression models with 95% confidence intervals in four coastal regions in Japan, estimated using standardized explanatory variables. t = hatchery release year of chum salmon fry. Sea surface temperature (SST). See Figure 1a for management region locations.

FIGURE 8 Predicted and observed chum salmon marine survival (adult return rates) by management region in Japan. Predicted return rates (grey line), observed return rates (black dots), and 95% prediction intervals (black lines). Note that scales on return rate axis differ by region. See Figure 1a for management region locations.

TABLE 1 Three subset models of stressors related to chum salmon marine survival (adult return rate) of fry released in year t (1998–2019) and explanatory variables considered in a regression model and hypothesized directional relationship. Sea surface temperature (SST); Western North Pacific (WN Pacific); Gulf of Alaska (GOA).

Subset model	Explanatory variable (year)	Age[†]	Spatial scale[‡]	Hypothesized relationship to chum salmon survival (source)
Hatchery carryovers	Parental relative fecundity (t -1)	0	Japan coast	Survival related (negative) to higher fecundity (Hasegawa et al., 2021a)
	Egg size (t)	0	Japan coast	Survival related (positive) to larger eggs (Hasegawa et al., 2021a)
	Fry size at release (t)	0	Japan coast	Survival related (positive) to size at release (Hasegawa et al., 2021b)
Ocean conditions	SST at release (t)	0	Japan coast	Survival related (negative) to SST (Nagata et al., 2007)
	Okhotsk coast Jul SST (t)	0	Okhotsk	1st summer survival related (negative) to SST (Nagata et al., 2007)
	WN Pacific Jan-Apr SST (t+1)	1	WN Pacific	1st winter survival related (positive) to SST (this study)
	Bering Sea Aug-Sep SST (t+1)	1	Bering Sea	2nd summer survival related (negative) to SST (this study)
	GOA Jan-Apr SST (t+2)	2	Gulf of Alaska	2nd winter survival related (positive) to SST (this study).
Predators and competitors	Yellowtail catch (t)	0	Japan coast	1st year survival related (negative) to predation (Beamish, 2018)
	Russia chum catch (t+3)	0–4	Okhotsk, North Pacific	1st - 3rd year survival related (negative) to Russian chum (this study).
	Russia pink catch (t+1)	0–1	Okhotsk, WN Pacific	1st year survival related (negative) to Russian pink (this study)
	Russia pink catch (t+2)	1–2	Bering Sea	2nd year survival related (negative) to Russian pink (Ruggerone et al., 2023
	Russia pink catch (t+3)	2–3	Gulf of Alaska, Bering Sea	3rd year survival related (negative) to Russian pink (Ruggerone et al., 2023)
	US & Canada chum catch (t+3)	1–4	North Pacific	2nd - 3rd year survival related (negative) to US & Canadia chum (this study)
	US &Canada pink catch (t+1)	0–1	Okhotsk, WN Pacific	1st year survival related (negative) to US & Canada pink (this study)
	US & Canada pink catch (t+2)	1–2	Bering Sea	2nd year survival related (negative) to US & Canada pink (this study)
	US and Canada pink catch (t+3)	2–3	Gulf of Alaska, Bering Sea	3rd year survival related (negative) to US & Canada pink (this study)

[†] Age of Japanese chum salmon released in year t (=1998,…,2019). [‡] See Japanese salmon migration route (Figure 1).

TABLE 2 Best regression models of Japanese chum salmon survival in four coastal regions in Japan and regression coefficients with their *p*-values in parentheses. Bold indicates significant relationship ($\alpha = 0.05$). *t* = hatchery release year of chum salmon fry. Sea surface temperature (SST); Western North Pacific (WN Pacific); Gulf of Alaska (GOA). See Figure 1a for management region locations.

Subset model	Explanatory variable (year)	Management region			
		Hokkaido Okhotsk	Hokkaido & Honshu Pacific	Hokkaido Sea of Japan	Honshu Sea of Japan
Hatchery carryovers	Relative fecundity[†][‡] (*t*-1)	–	**−0.931 (0.0009)**	–	–
	Egg size[†] (*t*-1)	–	–	–	–
	Fry size at release (*t*)	–	**0.486 (0.0009)**	–	–
Ocean conditions	SST at release (*t*)	–	–		**-0.483 (0.0033)**
	Okhotsk coast Jul SST (*t*)	–	–	**-0.315 (0.0042)**	**-0.138 (0.0452)**
	WN Pacific Jan-Apr SST (*t*+1)	–	–	–	0.259 (0.0647)
	Bering Sea Aug-Sep SST (*t*+1)	–	–	**-0.383 (0.0153)**	–
	GOA Jan-Apr SST (*t*+2)	–	–	**0.405 (0.0051)**	–
Predators and competitors	Yellowtail catch in year (*t*)	**-0.093 (0.0001)**	–	–	–
	Russia chum catch (*t*+3)	–	–	-0.151 (0.0504)	–
	Russia pink catch (*t*+1)	–	–	–	–
	Russia pink catch (*t*+2)	-0.127 (0.0559)	**-0.311 (0.0025)**	–	-0.139 (0.0605)
	Russia pink catch (*t*+3)	–	**-0.225 (0.0138)**	–	–
	USA & Canada chum catch (*t*+3)				
	USA & Canada pink catch (*t*+1)	–	–	–	–
	USA & Canada pink catch (*t*+2)	–	–	–	–
	USA & Canada pink catch (*t*+3)	–	–	–	–
	Adjusted R^2 (*p*-value)	**0.58**	**0.79**	**0.54**	**0.57**

The three subset models were fitted to log return rates for 1998-2019 releases and variable selection was performed using the Akaike Information Criterion corrected for small sample sizes (AICc). Multicollinearity of explanatory variables was checked using the variance inflation factor (VIF), and those with VIF < 3.0 were used for variable selection.

[†] Egg size and relative fecundity of age 4 and age 5 fish were examined. [‡] Age 4 had a larger R^2 value (0.79) than that of age 5 (0.61), so the result for age 4 fish is presented.

TABLE 3 Regression coefficients and practical effect sizes on Japanese chum salmon return rates in four coastal regions in Japan. t = hatchery release year of chum salmon fry. Sea surface temperature (SST); Western North Pacific (WN Pacific); Gulf of Alaska (GOA).

Region	Explanatory variable (year)	Estimate $\hat{\beta}_i$	Standard error	Effect size [†]	95% lower	95% upper	Unit	Practical effect (%) [‡]
Sea of Okhotsk	Russia pink catch (t+2)	-0.127	0.062	0.881	0.779	0.995	100 million fish	-11.9
	Yellowtail catch (t)	-0.093	0.020	0.911	0.877	0.947	100 tons	-8.9
Hokkaido and Honshu Pacific	Relative fecundity (t-1)	-0.931	0.232	0.394	0.250	0.620	100 eggs [§]	-60.6
	Russia pink catch (t+2)	-0.311	0.088	0.732	0.617	0.870	100 million fish	-26.8
	Russia pink catch (t+3)	-0.225	0.082	0.799	0.680	0.938	100 million fish	-20.1
	Fry size at release (t)	0.486	0.121	1.625	1.283	2.059	0.1g	62.5
Hokkaido Sea of Japan	Bering Sea Aug-Sep SST (t+1)	-0.383	0.142	0.682	0.516	0.901	1°C	-31.8
	GOA Jan-Apr SST (t+2)	0.405	0.126	1.499	1.171	1.918	1°C	49.9
	Okhotsk coast Jul SST (t)	-0.315	0.096	0.730	0.605	0.880	1°C	-27.0
	Russia chum catch (t+3)	-0.151	0.072	0.859	0.746	0.990	10 million fish	-14.1
Honshu Sea of Japan	WN Pacific Jan-Apr SST (t+1)	0.259	0.131	1.295	1.002	1.675	1°C	29.5
	Okhotsk coast Jul SST (t)	-0.138	0.064	0.871	0.769	0.987	1°C	-12.9
	Russia pink catch (t+2)	-0.139	0.069	0.870	0.760	0.997	100 million fish	-13.0
	SST at release (t)	-0.483	0.142	0.617	0.467	0.814	1°C	-38.3

[†] $\exp(\hat{\beta}_i)$, [‡] $(\exp(\hat{\beta}_i) - 1) \times 100$, [§] Number of eggs /FL $(cm)^3 \times 10^3$