Subcortical volumes, frontal cortical thickness, and pro-inflammatory cytokines in schizophrenia versus methamphetamine-induced psychosis

Blake, Lauren.[*a,c], Williams, Kimberley Clare.[a], Uhlmann, Anne Alex.[b], Temmingh, Henk.[a], Burger, Antoinette.[a,e], Stein, Dan Joseph.[a,c,d], Naudé, Petrus J. W.[a,c]

Running title: MRI and inflammation in psychotic disorders

[a] Department of Psychiatry and Mental Health, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa
[b] Department of Child and Adolescent Psychiatry and Psychotherapy, Faculty of Medicine, Technische Universität Dresden, Dresden, Germany
[c] Neuroscience Institute, University of Cape Town, South Africa
[d] South African Medical Research Council Unit on Risk and Resilience in Mental Disorders, University of Cape Town, South Africa.
[e] Brown University, Providence, Rhode Island, United States of America

* LBlake4759@hotmail.com, +27735344164
Department of Psychiatry and Mental Health
Groote Schuur Hospital, Neuroscience Institute
University of Cape Town
Main Road, Observatory
Cape Town, South Africa
7935

Word count:

Abstract: 248

Text body: 2940

Abstract

Background and Hypothesis: Clinical features of schizophrenia and methamphetamine-induced psychosis (MAP) have significant overlap. Schizophrenia is associated with alterations in subcortical volumes and cortical thickness and pro-inflammatory cytokines, that may correlate with clinical features. However, analogous work on MAP is lacking. We hypothesise decreased frontal cortical thickness, amygdala and hippocampal volumes in schizophrenia and MAP compared to controls, with increased basal ganglia in schizophrenia compared to MAP.

Study Design: Participants included n=36 schizophrenia, n=27 MAP and 32 age- and sex-matched healthy controls. Diagnosis and symptom severity were determined with the Structured Clinical Interview for Axis I Disorders, and the Positive and Negative Syndrome Scale, respectively. Structural T1-weighted images were acquired using a 3-Tesla magnetic resonance imaging scanner. Serum peripheral cytokine concentrations were measured using a multiplex bead array.

Study Results: Schizophrenia and MAP participants showed decreased left amygdala volumes and frontal cortical thickness compared to controls. Schizophrenia participants had increased bilateral caudate, putamen, and nucleus accumbens (NAcc) volumes compared to controls, and greater right globus pallidus and NAcc volumes compared to MAP. Decreased left amygdala volumes and frontal cortical thickness were significantly associated with longer illness duration in schizophrenia and methamphetamine use in MAP. No significant differences were found in cytokine levels between groups or associations with neuroimaging measures.

Conclusions: This study highlights overlaps and differences in the psychobiology of schizophrenia and MAP.The novel discovery of increased globus pallidus and NAcc volumes in schizophrenia compared with MAP may show important distinctions in the neurobiology between these two conditions.

Keywords: Basal ganglia, amygdala, hippocampus, cortical thickness, psychotic disorders, neuroinflammation

1. Introduction

There is considerable overlap in the clinical features of schizophrenia and methamphetamine-induced psychosis (MAP). Both schizophrenia and MAP are characterized by positive symptoms of psychosis. Thus, it has been suggested that MAP may serve as an important model for understanding the neurobiology of schizophrenia. Nevertheless, few studies have directly compared the biosignatures of schizophrenia and MAP.

Structural magnetic resonance imaging (MRI) studies have shown differences in subcortical gray matter volumes, in schizophrenia, which have been associated with clinical symptom profiles [1-5]. In schizophrenia, decreased hippocampal volume has been associated with negative symptoms and hallucinations [4,5], with decreased amygdala-hippocampal complex associated with negative symptoms of psychosis [5]. Furthermore, increased basal ganglia volumes have been consistently observed in schizophrenia compared to healthy controls [6-8]. Alterations in basal ganglia volumes have also been associated with clinical symptom profiles [1-5] including positive and negative symptoms [9]. Differences in cortical thickness of the frontal and temporal lobes have also been observed in individuals with schizophrenia [10-15], with the frontal cortex being most affected [15]. While structural neuroimaging studies of MAP are limited, there is some evidence of increased cortical thickness in MAP compared to individuals with methamphetamine dependence [16] and decreased hippocampal and amygdala volumes in MAP compared to healthy controls [17]. However, no studies have directly compared subcortical volumes and cortical thickness in schizophrenia to that in MAP.

Schizophrenia is further characterized by a number of alterations in immune biomarkers (Prestwood et al., 2021). Alterations of peripheral blood inflammatory cytokines (e.g. TNF)-α, IL-1β, IL-6, IL-8, IL-10, IL-12, and IFN-γ) have been associated with clinical outcomes in schizophrenia [18,19]. However, few studies have investigated immune markers in MAP, and none have directly compared immune markers in schizophrenia to those in MAP. Given findings of associations between neuroimaging measures and cytokine levels in schizophrenia [20], and evidence of neuro-inflammation in laboratory models of MA use [21,22], investigation on associations between neuroimaging measures and cytokines in MAP can advance our understanding on the potential functions of neuro-inflammation in schizophrenia and MAP.

This study aims to compare subcortical volumes, cortical thickness, and cytokine levels in individuals with schizophrenia, MAP, and healthy controls. We also assessed associations of neuroimaging

measures with illness duration, symptom severity, and pro-inflammatory cytokines in schizophrenia and MAP, and with duration of methamphetamine use in MAP.

2. Materials and methods

2.1. Participants

In this exploratory study, participants between the ages of 20-40 years, were recruited from Valkenberg Hospital and Groote Schuur Hospital in Cape Town, and included 36 (8 females: 28 males) participants with schizophrenia, 27 (6 females: 21 males) participants with MAP and 32 (13 females: 19 males) healthy controls, matched for age and sex, were recruited from the general population in Cape Town.

Individuals with schizophrenia and MAP had clinical diagnosis confirmed with the Structured Clinical Interview for DSM-IV (SCID-I), and symptom severity measured with the Positive and Negative Syndrome Scale (PANSS) [23]. The SCID-I was also given to healthy controls to include only those without Axis-I disorders. Participants were excluded if they had a history of 1) chronic medical illness that required medical care or prescription medication, including severe brain trauma and intellectual disability; 2) were recently or currently pregnant or lactating; 3) were unable to undergo MRI because of metal implants, claustrophobia, or other reasons; 4) substance dependence other than nicotine dependence.

Written consent was obtained from all participants included in the study. The work adhered to human research guidelines as stipulated in the Declaration of Helsinki [24]. The protocol underwent ethical review and approval with the University of Cape Town's Human Research Ethics Committee (UCT HSF HREC: 413/2016).

2.2. Imaging procedures

Structural magnetic resonance imaging (sMRI) was conducted in the morning on the second day of the study at the Cape Universities Body Imaging Centre (CUBIC), using a 3T Siemens Skyra MRI scanner (Siemens Medical Systems, Erlangen, Germany). A 32-channel head coil was used. A high resolution, T1-weighted scans were acquired with a multi-echo MPRAGE structural scan with the following scan

parameters: TR= 2530ms; graded TE= 1.69, 3.55, 5.41 and 7.27ms; flip angle= 7°; field of view (FOV)= 256mm; voxel size = 1x1x1mm; 6:39min.

2.3. Image processing

Processing of all MRIs was performed using FreeSurfer v6.0 [25,26] (http://surfer.nmr.mgh.harvard.edu/) using the Desikan-Killiany atlas, for cortical reconstruction and volume segmentation [27]. The processing pipeline included motion correction, skull-stripping, talairach transformation, segmentation of subcortical white matter and deep grey matter volumetric structures, intensity normalization, tessellation of the grey matter/white matter boundary, automated topology correction, and surface deformation and registration to the Desikan-Killiany atlas [27]. Reconstruction of the grey and white matter boundary and pial surface was performed to extract cortical thickness measurements [28-30]. Data and statistical summaries were generated using ENIGMA protocols (http://enigma.ini.usc.edu/protocols/imaging-protocols) and RStudio v 1.2.5054 [31].

2.4. Inflammatory markers

Blood samples were collected via venepuncture into serum tubes in the morning of the first day of the study. The tubes were kept at room temperature at for approximately 30 minutes to allow clotting and were subsequently centrifuged at 1,800 × g for 10 min. The separated serum was then aliquoted into cryovials and stored at -80 °C until analyses. The Luminex Milliplex® MAP human cytokine / chemokine magnetic bead panel (HSTCMAG28SK07; Merck) was used to detect cytokines; TNF-α, IL-1β, IL-6, IL-8, IL-10, IL-12p, and IFN-γ. The intra-assay coefficients of variation for each cytokine were all below 14%. All plates were analysed on the same day. Serum measures for IL-6 were below the detectable limit and were consequently excluded from statistical analysis.

2.5. Statistical analyses

A priori power analysis, to determine the minimum required sample size (n) required, was conducted with G*Power 3 [32] as a function of the required power level (1 – β error probability = 0.8), the pre-specified significance level (α = 0.05), and the population effect size (d = 0.5) with an allocation ratio of N2/N1 = 1. The *a priori* power analysis yielded a required total sample size of n = 102, with statistical power of 0.81.

To compare brain measures across schizophrenia (n = 36), MAP (n = 27), and healthy controls (n = 32), we employed a multiple analysis of covariance (MANCOVA) for all brain regions, with intracranial volume (ICV) [33-35], sex, and age [36-38] as covariates for subcortical volume measurements and, sex and age as covariates for frontal cortical thickness measurements. Thereafter, a multiple analysis of variance (MANOVA) for parametric data and Kruskal Wallis analysis of variance (ANOVA) for non-parametric data to determine whether there were group differences. MANOVA was followed by Bonferroni post-hoc tests to determine which groups differed from each other. Effect sizes for group comparisons were calculated using partial eta squared for all parametric ANOVAs and eta squared for Kruskal Wallis ANOVAs at a confidence interval of 90% [39]. Spearman's Rank correlations (for non-normally distributed data) and Pearson's correlations (normally distributed data) were performed within each group to determine associations of neuroimaging measures with symptom severity, illness duration, duration of methamphetamine use and cytokine concentrations. Significance levels were set at Rho of ± 0.5 and $p < 0.01$ (where the critical value for r = 0.449 when $p = 0.01$) to reduce Type I Error and increase stringency and strength of the associations between variables.

In a sensitivity analysis to determine whether there were differences between the schizophrenia participants who reported methamphetamine (n = 14) use to those that did not report methamphetamine use (n = 22), we employed t-tests for parametric brain regions and Mann Whitney U tests for non-parametric brain regions (Supplementary Tables 1, 2 and 3). Effect sizes were calculated using Cohen's D for parametric data at a 95% confidence interval and biserial correlations r at $p < 0.05$.

3. Results

3.1. Demographic and clinical variables

Table 1 shows that there were no significant group differences for age and sex, lower levels of school education were found in schizophrenia ($H_{(2,94)}$= 19.827, $p = 0.002$) and in MAP ($H_{(2,94)}$= 19.827, $p = 0.00046$), compared to healthy controls. Duration of illness was significantly longer in schizophrenia compared to MAP ($z = 3.271$, $p = 0.0008$) and the number of psychotic episodes were greater in schizophrenia than MAP ($p = 0.038$). Significant differences were found for the PANSS positive ($H_{(2,92)}$= 40.21, $p < 0.0001$), negative ($H_{(2,92)}$= 43.86, $p < 0.0001$), general ($H_{(2,92)}$= 32.99, $p < 0.0001$) and total

scores ($H_{(2,92)}$= 46.64, p < 0.0001), with significant differences found between controls and schizophrenia (p < 0.0001) and, controls and MAP ($p \leq 0.00012$).

3.2. Subcortical volumes

There were decreased left amygdala volumes in schizophrenia ($F_{(2,91)}$ = 6.37, p = 0.012) and in MAP ($F_{(2,91)}$ = 6.37, p = 0.0054) compared to healthy controls (Table 2). There were increased bilateral caudate ($F_{(2,91)}$ = 4.11, p = 0.03; $F_{(2,91)}$ = 3.95, p = 0.05), putamen ($F_{(2,91)}$ = 6.85, p = 0.016; $F_{(2,91)}$ = 5.27, p = 0.0072) and NAcc ($F_{(2,91)}$ = 18.69, p < 0.0001; $F_{(2,91)}$ = 7.83, p = 0.014) volumes in schizophrenia compared to healthy controls. Decreased bilateral NAcc ($F_{(2,91)}$ = 18.69; p < 0.0001; $F_{(2,91)}$ = 7.83, p = 0.0017) and right globus pallidus ($F_{(2,91)}$ = 3.81, p = 0.021) volumes were observed in MAP compared to schizophrenia (Table 2).

No significant differences in subcortical volumes were observed between the schizophrenia group which reported methamphetamine use and the schizophrenia group who did not report methamphetamine use (Supplementary Table 1).

3.3 Frontal cortical thickness

There was decreased frontal cortical thickness in schizophrenia compared to healthy controls for all frontal regions ($F_{(2,91)} > 10.5$, $p < 0.0004$) except for the right lateral orbitofrontal, left and right medial orbitofrontal and right rostral middle frontal regions (Table 3).

Decreased frontal cortical thickness in MAP compared to healthy controls for all frontal regions (All, $F_{(2,91)} > 10.5$, $p < 0.013$) was found except for the left lateral orbitofrontal, medial orbitofrontal, right rostral middle frontal and superior frontal and the left and right pars triangularis, and increased cortical thickness in MAP compared to healthy controls in the right medial orbitofrontal ($H_{(2,94)} = 7.21$, $p = 0.042$) and right lateral orbitofrontal ($F_{(2,91)} = 10.7$, $p = 0.028$).

Increased frontal cortical thickness was observed in MAP compared to schizophrenia for the right lateral orbitofrontal ($F_{(2,91)} = 10.7$, $p < 0.0001$), left pars opercularis ($F_{(2,91)}) = 82.5$, $p < 0.0001$), left pars triangularis ($F_{(2,91)} = 29.2$, $p < 0.0001$), left lateral orbitofrontal ($H_{(2,94)} = 60.76$, $p < 0.0001$), left medial orbitofrontal ($H_{(2,94)} = 12.22$, $p = 0.0014$), left frontal pole ($H_{(2,94)} = 82.30$, $p < 0.0001$) and left and right pars triangularis ($F_{(2,91)} = 29.2$, $p < 0.0001$). There was increased frontal cortical thickness was observed in schizophrenia compared to MAP for the right rostral middle frontal ($H_{(2,94)} = 10.78$, $p = 0.0032$) and the right pars opercularis ($F_{(2,91)} = 31.6$, $p = 0.0001$).

Increased frontal cortical thickness was observed in schizophrenia group which used methamphetamine for the left lateral orbitofrontal ($U = 64.5$, $p = 0.039$) and right medial orbitofrontal ($U = 67.5$, $p = 0.05$) compared to the schizophrenia group who reported no methamphetamine use. Decreased frontal cortical thickness was observed in the schizophrenia group who reported methamphetamine use for the left frontal pole ($U = 178.5$, $p = 0.039$) compared to the schizophrenia group who reported no methamphetamine use (Supplementary Tables 2 and 3).

3.4. Cytokines

There were no significant group differences in cytokine concentrations for IFN-γ ($H_{2,94} = 2.81$; $p = 0.245$), IL-10 ($H_{2,94} = 0.93$, $p = 0.627$), IL-12 ($H_{2,94} = 4.90$, $p = 0.0862$), IL-1β ($H_{2,94} = 1.27$, $p = 0.531$), IL-8 ($H_{2,94} = 0.23$; $p = 0.892$) and TNF-α ($H_{2,94} = 1.01$, $p = 0.605$).

3.5. Associations of neuroimaging measures with pro-inflammatory cytokines, illness duration and symptom severity

There were no significant associations between neuroimaging measures and either symptom severity with cytokine concentrations.

In schizophrenia, longer illness duration was associated with decreased left amygdala volumes ($R_{(n=36)}$ = -0.43, p = 0.0088), and decreased frontal cortical thickness (All, $R_{(n=36)} < -0.51$, $p < 0.0013$), except for the left frontal pole ($R_{(n=36)}$ = 0.48, p = 0.0029) and right frontal pole ($R_{(n=36)}$= 0.37, p = 0.025).

In MAP, increased duration of methamphetamine use was associated with decreased left amygdala ($r_{(n=27)}$ = -0.51, p = 0.0069) volumes, left lateral orbitofrontal ($R_{(n=27)}$ = -0.54, p = 0.0030), left medial orbitofrontal ($R_{(n=27)}$ = -0.55; p = 0.0030), right medial orbitofrontal ($R_{(n=27)}$ = -0.55, p = 0.0030), right rostral middle frontal ($R_{(n=27)}$ = -0.55, p = 0.0030) and left frontal pole ($R_{(n=27)}$ = -0.55, p = 0.0030).

4. Discussion

The findings of this study include decreased left amygdala volumes and frontal cortical thickness in schizophrenia and MAP compared to controls; increased volumes in bilateral caudate, putamen, and nucleus accumbens (NAcc) volumes in schizophrenia compared to controls; and increased right globus pallidus and bilateral NAcc volumes in schizophrenia compared to MAP. There were no significant differences across groups for cytokine levels and their associations with neuroimaging measures. Lastly, decreased left amygdala volumes and frontal cortical thickness were significantly associated with longer illness duration in schizophrenia and with longer duration of methamphetamine use in MAP.

The present data are consistent with previous findings that decreased left amygdala volumes and frontal cortical thinning are present in both schizophrenia [40,41] and MAP [16,17] compared to healthy controls, and are associated with duration of illness in schizophrenia [42,43]. This study is the first to our knowledge showing decreased amygdala volume and cortical thickness in MAP. Taken together, these findings suggest important overlaps in the neurobiology of schizophrenia and MAP. Decreased cortical thickness and thus pyramidal neuron layers may reflect decreased neural density and functional dysconnectivity to subcortical and other cortical brain regions, which has been reported to underlie the symptoms of schizophrenia [44-46] and therefore may occur in MAP. However, further studies are needed to elucidate whether psychotic symptoms in MAP are indeed due to loss of neural connectivity from the frontal cortical neurons to subcortical and cortical structures.

Increased caudate volume was observed in schizophrenia but not in MAP, which is a consistent finding in schizophrenia and has been attributed to typical antipsychotic use [6,47-50]. The finding of increased basal ganglia volumes has been replicated in several studies, including putamen volumes in chronic schizophrenia [41,49,51-53]. The finding that globus pallidus and NAcc volume were increased in schizophrenia compared with MAP is a novel one, and may also highlight important differences in the neurobiology between schizophrenia and MAP. However, it is notable that increased globus pallidus volume was associated with increased duration of illness in schizophrenia, and that increased volume of these structures has been attributed to antipsychotic use. Given that use of antipsychotic agents is typically of lower duration in MAP, we cannot exclude the possibility that this finding points to differences in treatment between schizophrenia and MAP.

Decreased frontal cortical thickness was observed in the superior, middle and inferior frontal brain regions in schizophrenia compared to healthy controls, in addition to that of the left lateral OFC, widely reported finding in schizophrenia [10,11,15,16,54-60], this was also observed for the schizophrenia group who did not report methamphetamine use. Cortical thickness has been reported to decrease more rapidly throughout the course of the illness [57,61] compared to healthy controls and has also been observed in chronic schizophrenia [14,62-64]. In MAP we observed significant decreases for the right medial OFC and lateral OFC cortical thickness compared to healthy controls, whilst the schizophrenia group showed significant decreases in cortical thickness of the lateral OFC and left medial OFC compared to MAP, suggesting a deleterious effect of schizophrenia on these regions in the left hemisphere. However, sensitivity analysis suggests that these findings were not observed in the schizophrenia group who reported no methamphetamine use, therefore this finding may be attributed to methamphetamine use instead of the structural differences we observed between the two psychotic disorders. This finding suggests decreased frontal cortical thickness in the right hemisphere in MAP compared to schizophrenia, occurring in the left hemisphere, which may aid in distinguishing MAP from schizophrenia. This is the first study to report decreased cortical thickness of the OFC regions in MAP, which has been reported in antipsychotic naïve schizophrenia individuals [65]. We observed increased left frontal pole in the schizophrenia group compared to MAP as well as in the schizophrenia group who did not report methamphetamine use compared to those who reported methamphetamine use.

In MAP we observed significant decreases for the right lateral and medial OFC cortical thickness compared to healthy controls, whilst the schizophrenia group showed significant decreases in cortical thickness compared to that of MAP, particularly the lateral and left medial OFC. We speculate that the considerable duration of illness in the schizophrenia participants of this study may contribute to the differences between schizophrenia to MAP. In this respect, longer illness duration has been attributed to cortical thinning in patients with schizophrenia, including the PFC [57,61]. Furthermore, increased duration of methamphetamine use was significantly associated with decreased frontal cortical thickness in MAP participants. Therefore, our results suggest a distinct deleterious effect of schizophrenia on the pathophysiology of reduced cortical thickness in the lateral and medial OFC regions in the left hemisphere.

We found no significant differences in cytokine concentrations across groups. Furthermore, no significant associations were observed between neuroimaging measures and cytokines levels. There

is some inconsistency in the literature on immune markers on schizophrenia; our findings are consistent with some previous work [67-71]. However, we cannot rule out the possibility that our negative findings are due to insufficient power. Moreover, the anti-inflammatory effects of antipsychotic medication in both schizophrenia and MAP groups may have ameliorated the inflammatory status in these study groups [70,72,73].

A number of limitations deserve emphasis. First, the use of antipsychotics in both schizophrenia and MAP patients may have confounded the results reported in this study. Second, although our sample size is consistent with previous work on MAP, the small sample size warrants replication in a larger study. Third, self-reported methamphetamine abstinence was not supported by an objective confirmation. Fourth, sex differences were not examined due to the small sample of female participants. Fifth, methamphetamine use in some participants in the schizophrenia group may have confounded the comparisons with the MAP group, however our sensitivity analysis provides some reassurance that this is not the case.

4.1 Conclusions

This study shows distinctive structural variations in basal ganglia and amygdala volumes, in addition to frontal cortical thickness, in individuals with schizophrenia and MAP, suggesting that these structures play a pivotal role in the pathophysiology of these disorders. Differences in basal ganglia volumes between schizophrenia and MAP may be attributed by longer illness duration in schizophrenia. The novel discovery of increased globus pallidus and NAcc volumes in schizophrenia compared with MAP may show crucial distinctions in the neurobiology between these two conditions. To deepen our understanding of the neurobiological underpinnings of MAP, future investigations should employ larger sample sizes, incorporate longitudinal study designs, and integrate magnetic resonance spectroscopy which may show important neurometabolic signatures in these brain regions in MAP.

Acknowledgements We thank all study staff and staff at the Cape Universities Body Imaging Centre (CUBIC) of the University of Cape Town. We also wish to thank all who participated in this study.

Funding: The study was supported by the Medical Research Council and National Research Foundation. L.B. also acknowledges her funding sources, administered by the University of Cape Town, National Research Foundation and Deutscher Akademischer Austauschdienst.

Conflict of interest: The authors have nothing to disclose.

5. References

1.	Fan F, Xiang H, Tan S, et al. Subcortical structures and cognitive dysfunction in first episode schizophrenia. *Psychiatry Res Neuroimaging*. Apr 30 2019;286:69-75. doi:10.1016/j.pscychresns.2019.01.003

2.	Koshiyama D, Fukunaga M, Yamashita F, et al. *Role of subcortical structures on cognitive and social function in schizophrenia*. Vol. 8. 2018:1183. *Scientific Reports*. 2045-2322. 2018/01/19. https://doi.org/10.1038/s41598-017-18950-2

3.	Makowski C, Bodnar M, Shenker JJ, et al. Linking persistent negative symptoms to amygdala-hippocampus structure in first-episode psychosis. *Transl Psychiatry*. Aug 8 2017;7(8):e1195. doi:10.1038/tp.2017.168

4.	Duan X, He C, Ou J, et al. Reduced Hippocampal Volume and Its Relationship With Verbal Memory and Negative Symptoms in Treatment-Naive First-Episode Adolescent-Onset Schizophrenia. *Schizophr Bull*. Jan 23 2021;47(1):64-74. doi:10.1093/schbul/sbaa092

5.	Rajarethinam R, DeQuardo JR, Miedler J, et al. Hippocampus and amygdala in schizophrenia: assessment of the relationship of neuroanatomy to psychopathology. *Psychiatry Res*. Nov 30 2001;108(2):79-87. doi:10.1016/s0925-4927(01)00120-2

6.	Chakos MH, Lieberman JA, Bilder RM, et al. Increase in caudate nuclei volumes of first-episode schizophrenic patients taking antipsychotic drugs. *The American journal of psychiatry*. Oct 1994;151(10):1430-6. doi:10.1176/ajp.151.10.1430

7.	Lauer M, Senitz D, Beckmann H. Increased volume of the nucleus accumbens in schizophrenia. *J Neural Transm (Vienna)*. 2001;108(6):645-60. doi:10.1007/s007020170042

8.	Corson PW, Nopoulos P, Miller DD, Arndt S, Andreasen NC. Change in basal ganglia volume over 2 years in patients with schizophrenia: typical versus atypical neuroleptics. *The American journal of psychiatry*. Aug 1999;156(8):1200-4. doi:10.1176/ajp.156.8.1200

9.	Li Y, Li WX, Xie DJ, Wang Y, Cheung EFC, Chan RCK. Grey matter reduction in the caudate nucleus in patients with persistent negative symptoms: An ALE meta-analysis. *Schizophr Res*. Feb 2018;192:9-15. doi:10.1016/j.schres.2017.04.005

10.	Rimol LM, Hartberg CB, Nesvag R, et al. Cortical thickness and subcortical volumes in schizophrenia and bipolar disorder. *Biological psychiatry*. Jul 1 2010;68(1):41-50. doi:10.1016/j.biopsych.2010.03.036

11.	Rimol LM, Nesvag R, Hagler DJ, Jr., et al. Cortical volume, surface area, and thickness in schizophrenia and bipolar disorder. *Biological psychiatry*. Mar 15 2012;71(6):552-60. doi:10.1016/j.biopsych.2011.11.026

12.	Sugihara G, Oishi N, Son S, Kubota M, Takahashi H, Murai T. Distinct Patterns of Cerebral Cortical Thinning in Schizophrenia: A Neuroimaging Data-Driven Approach. *Schizophr Bull*. Jul 1 2017;43(4):900-906. doi:10.1093/schbul/sbw176

13.	Nenadic I, Yotter RA, Sauer H, Gaser C. Patterns of cortical thinning in different subgroups of schizophrenia. *Br J Psychiatry*. Jun 2015;206(6):479-83. doi:10.1192/bjp.bp.114.148510

14.	Kuperberg GR, Broome MR, McGuire PK, et al. Regionally localized thinning of the cerebral cortex in schizophrenia. *Arch Gen Psychiatry*. Sep 2003;60(9):878-88. doi:10.1001/archpsyc.60.9.878

15.	van Erp TGM, Walton E, Hibar DP, et al. Cortical Brain Abnormalities in 4474 Individuals With Schizophrenia and 5098 Control Subjects via the Enhancing Neuro Imaging Genetics Through Meta Analysis (ENIGMA) Consortium. *Biological psychiatry*. Nov 1 2018;84(9):644-654. doi:10.1016/j.biopsych.2018.04.023

16.   Uhlmann A, Fouche JP, Koen N, Meintjes EM, Wilson D, Stein DJ. Fronto-temporal alterations and affect regulation in methamphetamine dependence with and without a history of psychosis. *Psychiatry Res Neuroimaging*. Feb 28 2016;248:30-8. doi:10.1016/j.pscychresns.2016.01.010

17.   Orikabe L, Yamasue H, Inoue H, et al. Reduced amygdala and hippocampal volumes in patients with methamphetamine psychosis. *Schizophr Res*. Nov 2011;132(2-3):183-9. doi:10.1016/j.schres.2011.07.006

18.   Kose M, Pariante CM, Dazzan P, Mondelli V. The Role of Peripheral Inflammation in Clinical Outcome and Brain Imaging Abnormalities in Psychosis: A Systematic Review. Systematic Review. *Frontiers in Psychiatry*. 2021-February-19 2021;12doi:10.3389/fpsyt.2021.612471

19.   Pillinger T, Osimo EF, Brugger S, Mondelli V, McCutcheon RA, Howes OD. A Meta-analysis of Immune Parameters, Variability, and Assessment of Modal Distribution in Psychosis and Test of the Immune Subgroup Hypothesis. *Schizophr Bull*. Sep 11 2019;45(5):1120-1133. doi:10.1093/schbul/sby160

20.   Wu D, Lv P, Li F, et al. Association of peripheral cytokine levels with cerebral structural abnormalities in schizophrenia. *Brain Research*. 2019/12/01/ 2019;1724:146463. doi:https://doi.org/10.1016/j.brainres.2019.146463

21.   Davidson M, Mayer M, Habib A, et al. Methamphetamine Induces Systemic Inflammation and Anxiety: The Role of the Gut-Immune-Brain Axis. *Int J Mol Sci*. Sep 23 2022;23(19)doi:10.3390/ijms231911224

22.   Shi S, Chen T, Zhao M. The Crosstalk Between Neurons and Glia in Methamphetamine-Induced Neuroinflammation. *Neurochem Res*. Apr 2022;47(4):872-884. doi:10.1007/s11064-021-03513-9

23.   Thompson PM, Hayashi KM, Simon SL, et al. Structural Abnormalities in the Brains of Human Subjects Who Use Methamphetamine. *The Journal of Neuroscience*. 2004;24(26):6028.

24.   World Medical A. World Medical Association Declaration of Helsinki: Ethical Principles for Medical Research Involving Human Subjects. *JAMA*. 2013;310(20):2191-2194. doi:10.1001/jama.2013.281053

25.   Fischl B. FreeSurfer. *NeuroImage*. Aug 15 2012;62(2):774-81. doi:10.1016/j.neuroimage.2012.01.021

26.   *FreeSurfer*. Version 2.1. Harvard University; 1999.

27.   Desikan RS, Segonne F, Fischl B, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*. Jul 1 2006;31(3):968-80. doi:10.1016/j.neuroimage.2006.01.021

28.   Fischl B, Dale AM. Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc Natl Acad Sci U S A*. Sep 26 2000;97(20):11050-5. doi:10.1073/pnas.200033797

29.   Fischl B, Liu A, Dale AM. Automated manifold surgery: constructing geometrically accurate and topologically correct models of the human cerebral cortex. *IEEE Trans Med Imaging*. Jan 2001;20(1):70-80. doi:10.1109/42.906426

30.   Segonne F, Pacheco J, Fischl B. Geometrically accurate topology-correction of cortical surfaces using nonseparating loops. *IEEE Trans Med Imaging*. Apr 2007;26(4):518-29. doi:10.1109/TMI.2006.887364

31.   *RStudio*. Version 1.2.5054. Free Software Foundation, Incorporated; 2009-2020.

32.   Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. May 2007;39(2):175-91. doi:10.3758/bf03193146

33.   Mathalon DH, Sullivan EV, Rawles JM, Pfefferbaum A. Correction for head size in brain-imaging measurements. *Psychiatry Res*. Jun 1993;50(2):121-39. doi:10.1016/0925-4927(93)90016-b

34.   O'Brien LM, Ziegler DA, Deutsch CK, Frazier JA, Herbert MR, Locascio JJ. Statistical adjustments for brain size in volumetric neuroimaging studies: some practical implications in methods. *Psychiatry Res*. Aug 30 2011;193(2):113-22. doi:10.1016/j.pscychresns.2011.01.007

35.   Sanfilipo MP, Benedict RH, Zivadinov R, Bakshi R. Correction for intracranial volume in analysis of whole brain atrophy in multiple sclerosis: the proportion vs. residual method. *NeuroImage*. Aug 2004;22(4):1732-43. doi:10.1016/j.neuroimage.2004.03.037

36.   Barnes J, Ridgway GR, Bartlett J, et al. Head size, age and gender adjustment in MRI studies: a necessary nuisance? *NeuroImage*. 2010;53(4):1244-1255.

37.   Crowley S, Huang H, Tanner J, et al. Considering total intracranial volume and other nuisance variables in brain voxel based morphometry in idiopathic PD. *Brain Imaging and Behavior*. 2018;12:1-12.

38.   Smith CD, Chebrolu H, Wekstein DR, Schmitt FA, Markesbery WR. Age and gender effects on human brain anatomy: a voxel-based morphometric study in healthy elderly. *Neurobiol Aging*. Jul 2007;28(7):1075-87. doi:10.1016/j.neurobiolaging.2006.05.018

39.   Jané M, Xiao Q, Yeung SK, et al. *Guide to Effect Sizes and Confidence Intervals*. 2024.

40.   Niu L, Matsui M, Zhou SY, et al. Volume reduction of the amygdala in patients with schizophrenia: a magnetic resonance imaging study. *Psychiatry Res*. Nov 15 2004;132(1):41-51. doi:10.1016/j.pscychresns.2004.06.002

41.   van Erp TG, Hibar DP, Rasmussen JM, et al. Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Molecular psychiatry*. Apr 2016;21(4):547-53. doi:10.1038/mp.2015.63

42.   Bustillo JR, Lauriello J, Rowland LM, et al. Effects of chronic haloperidol and clozapine treatments on frontal and caudate neurochemistry in schizophrenia. *Psychiatry Res*. Oct 1 2001;107(3):135-49. doi:10.1016/s0925-4927(01)00102-0

43.   Molina V, Martin C, Ballesteros A, de Herrera AG, Hernandez-Tamames JA. Optimized voxel brain morphometry: association between brain volumes and the response to atypical antipsychotics. *European archives of psychiatry and clinical neuroscience*. Sep 2011;261(6):407-16. doi:10.1007/s00406-010-0182-2

44.   Li S, Hu N, Zhang W, et al. Dysconnectivity of Multiple Brain Networks in Schizophrenia: A Meta-Analysis of Resting-State Functional Connectivity. Systematic Review. *Frontiers in Psychiatry*. 2019-July-12 2019;10doi:10.3389/fpsyt.2019.00482

45.   Li T, Wang Q, Zhang J, et al. Brain-Wide Analysis of Functional Connectivity in First-Episode and Chronic Stages of Schizophrenia. *Schizophrenia Bulletin*. 2016;43(2):436-448. doi:10.1093/schbul/sbw099

46.   Lewis DA, Curley AA, Glausier JR, Volk DW. Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends Neurosci*. Jan 2012;35(1):57-67. doi:10.1016/j.tins.2011.10.004

47.   Lang DJ, Kopala LC, Vandorpe RA, et al. Reduced basal ganglia volumes after switching to olanzapine in chronically treated patients with schizophrenia. *The American journal of psychiatry*. Oct 2004;161(10):1829-36. doi:10.1176/ajp.161.10.1829

48. Breier A, Buchanan RW, Elkashef A, Munson RC, Kirkpatrick B, Gellad F. Brain morphology and schizophrenia. A magnetic resonance imaging study of limbic, prefrontal cortex, and caudate structures. *Arch Gen Psychiatry*. Dec 1992;49(12):921-6. doi:10.1001/archpsyc.1992.01820120009003

49. Emsley R, du Plessis S, Phahladira L, et al. Antipsychotic treatment effects and structural MRI brain changes in schizophrenia. *Psychological Medicine*. 2023;53(5):2050-2059. doi:10.1017/S0033291721003809

50. Jørgensen KN, Nesvåg R, Gunleiksrud S, Raballo A, Jönsson EG, Agartz I. First- and second-generation antipsychotic drug treatment and subcortical brain morphology in schizophrenia. *European archives of psychiatry and clinical neuroscience*. 2016/08/01 2016;266(5):451-460. doi:10.1007/s00406-015-0650-9

51. Hokama H, Shenton ME, Nestor PG, et al. Caudate, putamen, and globus pallidus volume in schizophrenia: a quantitative MRI study. Article. *Psychiatry Res*. Nov 10 1995;61(4):209-29. doi:10.1016/0925-4927(95)02729-h

52. Mamah D, Wang L, Barch D, de Erausquin GA, Gado M, Csernansky JG. Structural analysis of the basal ganglia in schizophrenia. *Schizophr Res*. Jan 2007;89(1-3):59-71. doi:10.1016/j.schres.2006.08.031

53. Kitabayashi Y, Narumoto J, Otakara C, Hyungin C, Fukui K, Yamada K. Schizophrenia-like psychosis following right putaminal infarction. *The Journal of neuropsychiatry and clinical neurosciences*. Fall 2006;18(4):561-2. doi:10.1176/jnp.2006.18.4.561

54. Ahmed M, Cannon DM, Scanlon C, et al. Progressive Brain Atrophy and Cortical Thinning in Schizophrenia after Commencing Clozapine Treatment. *Neuropsychopharmacology*. Sep 2015;40(10):2409-17. doi:10.1038/npp.2015.90

55. Narr KL, Bilder RM, Toga AW, et al. Mapping cortical thickness and gray matter concentration in first episode schizophrenia. *Cereb Cortex*. Jun 2005;15(6):708-19. doi:10.1093/cercor/bhh172

56. Sprooten E, Papmeyer M, Smyth AM, et al. Cortical thickness in first-episode schizophrenia patients and individuals at high familial risk: a cross-sectional comparison. *Schizophr Res*. Dec 2013;151(1-3):259-64. doi:10.1016/j.schres.2013.09.024

57. van Haren NE, Schnack HG, Cahn W, et al. Changes in cortical thickness during the course of illness in schizophrenia. *Arch Gen Psychiatry*. Sep 2011;68(9):871-80. doi:10.1001/archgenpsychiatry.2011.88

58. Walton E, Hibar DP, van Erp TGM, et al. Prefrontal cortical thinning links to negative symptoms in schizophrenia via the ENIGMA consortium. *Psychol Med*. Jan 2018;48(1):82-94. doi:10.1017/S0033291717001283

59. Xiao Y, Lui S, Deng W, et al. Altered cortical thickness related to clinical severity but not the untreated disease duration in schizophrenia. *Schizophr Bull*. Jan 2015;41(1):201-10. doi:10.1093/schbul/sbt177

60. Zugman A, Gadelha A, Assunção I, et al. Reduced dorso-lateral prefrontal cortex in treatment resistant schizophrenia. *Schizophr Res*. Aug 2013;148(1-3):81-6. doi:10.1016/j.schres.2013.05.002

61. DeLisi LE, Sakuma M, Tew W, Kushner M, Hoff AL, Grimson R. Schizophrenia as a chronic active brain process: a study of progressive brain structural change subsequent to the onset of schizophrenia. Article. *Psychiatry Research: Neuroimaging*. 1997;74(3):129-140. doi:10.1016/s0925-4927(97)00012-7

62. Nesvag R, Lawyer G, Varnas K, et al. Regional thinning of the cerebral cortex in schizophrenia: effects of diagnosis, age and antipsychotic medication. *Schizophr Res*. Jan 2008;98(1-3):16-28. doi:10.1016/j.schres.2007.09.015

63.	Goldman AL, Pezawas L, Mattay VS, et al. Widespread reductions of cortical thickness in schizophrenia and spectrum disorders and evidence of heritability. *Arch Gen Psychiatry*. May 2009;66(5):467-77. doi:10.1001/archgenpsychiatry.2009.24

64.	Madre M, Canales-Rodriguez EJ, Fuentes-Claramonte P, et al. Structural abnormality in schizophrenia versus bipolar disorder: A whole brain cortical thickness, surface area, volume and gyrification analyses. *Neuroimage Clin*. 2020/01/01/ 2020;25:102131. doi:10.1016/j.nicl.2019.102131

65.	Venkatasubramanian G, Jayakumar PN, Gangadhar BN, Keshavan MS. Automated MRI parcellation study of regional volume and thickness of prefrontal cortex (PFC) in antipsychotic-naive schizophrenia. *Acta Psychiatr Scand*. Jun 2008;117(6):420-31. doi:10.1111/j.1600-0447.2008.01198.x

66.	Jung WH, Kim JS, Jang JH, et al. Cortical thickness reduction in individuals at ultra-high-risk for psychosis. *Schizophr Bull*. Jul 2011;37(4):839-49. doi:10.1093/schbul/sbp151

67.	Wilke I, Arolt V, Rothermundt M, Weitzsch C, Hornberg M, Kirchner H. Investigations of cytokine production in whole blood cultures of paranoid and residual schizophrenic patients. *European archives of psychiatry and clinical neuroscience*. 1996;246(5):279-84. doi:10.1007/BF02190280

68.	Potvin S, Stip E, Sepehry AA, Gendron A, Bah R, Kouassi E. Inflammatory cytokine alterations in schizophrenia: a systematic quantitative review. *Biological psychiatry*. Apr 15 2008;63(8):801-8. doi:10.1016/j.biopsych.2007.09.024

69.	Kaminska T, Wysocka A, Marmurowska-Michalowska H, Dubas-Slemp H, Kandefer-Szerszen M. Investigation of serum cytokine levels and cytokine production in whole blood cultures of paranoid schizophrenic patients. *Arch Immunol Ther Exp (Warsz)*. 2001;49(6):439-45.

70.	Miller BJ, Buckley P, Seabolt W, Mellor A, Kirkpatrick B. Meta-analysis of cytokine alterations in schizophrenia: clinical status and antipsychotic effects. *Biological psychiatry*. Oct 1 2011;70(7):663-71. doi:10.1016/j.biopsych.2011.04.013

71.	Lv MH, Tan YL, Yan SX, et al. Decreased serum TNF-alpha levels in chronic schizophrenia patients on long-term antipsychotics: correlation with psychopathology and cognition. *Psychopharmacology*. Jan 2015;232(1):165-72. doi:10.1007/s00213-014-3650-y

72.	Fraguas D, Diaz-Caneja CM, Ayora M, et al. Oxidative Stress and Inflammation in First-Episode Psychosis: A Systematic Review and Meta-analysis. *Schizophr Bull*. Jun 18 2019;45(4):742-751. doi:10.1093/schbul/sby125

73.	Flatow J, Buckley P, Miller BJ. Meta-analysis of oxidative stress in schizophrenia. *Biological psychiatry*. Sep 15 2013;74(6):400-9. doi:10.1016/j.biopsych.2013.03.018

Tables

Table 1: Demographics, duration of illness, clinical scales and methamphetamine use variables for control, schizophrenia and methamphetamine-induced psychotic groups

	CON (n = 32) 13 females: 19 males			SCHIZOPHRENIA (n = 36) 8 females: 28 males			MAP (n = 27) 6 females: 21 males				
	Mean	SD	Range	Mean	SD	Range	Mean	SD	Range	Test statistics	Group differences
Age on day of study	30.12	4.8	23 - 40	29.53	5.36	22 - 39	27.89	4.74	20 – 37	$H_{(2,94)}= 3.09, p = 0.21$	
School education (years)	11.65	0.91	9 – 13[a]	10.11	1.94	4 - 12	9.56	2.55	1 - 12	$H_{(2,94)}= 19.83, p < 0.01$	CON vs. schizophrenia, $p < 0.01$; CON vs. MAP, $p < 0.01$
Post-school education (years)	0.94	1.35	0 – 4	0.25	0.88	0 – 5	0.61	1.14	0 - 4	$H_{(2,94)}= 8.44, p = 0.02$	n.s.
Total education duration (years)	12.58	1.85	9 – 16[b]	10.37	2.28	4 - 17	10.17	3.16	1 - 16	$H_{(2,94)}= 18.829, p < 0.01$	CON vs. schizophrenia, $p < 0.01$; CON vs. MAP, $p < 0.01$
Duration of illness (months)	-	-	-	102.31	52.61	14 – 228[c]	60	48.34	2 – 180	$U =250, z = 3.271, p < 0.01$	
No. of psychotic episodes	-	-	-	3.65	1.62	1 - 9	2.32	1.08	1 - 5	$U = 336.5, z =-2.07, p = 0.038$	
Antipsychotic use variables											
Chlorpromazine equivalents	-	-	-	337.72	289.02	0-1100	210.74	198.76	0 -750		
Methamphetamine use measures											
Duration of use (months)	-	-	-	64.80	46.16	24-144	104.56	51.64	24 – 228		
	Mean	SD	Range	Mean	SD	Range	Mean	SD	Range		
Abstinence (months)	-	-	-	40.41	46.793	0.03 – 132	8.01	13.47	0.03 – 48		
Psychotic symptomatology PANSS											
Positive scale	7.10	0.41	7 – 9	12.16	4.88	7 – 25[d]	10.59	5.23	1 – 23	$H_{(2,92)}= 40.21, p < 0.01$	CON vs. schizophrenia, $p < 0.01$; CON vs. MAP, $p < 0.01$
Negative scale	7.07	0.26	7 – 8	14.89	6.41	7– 28[d]	14.30	10.02	7 – 45	$H_{(2,92)}= 43.86, p < 0.01$	CON vs. schizophrenia, $p < 0.01$; CON vs. MAP, $p < 0.01$
General psychopathology	16.31	0.71	16 – 19	21.81	7.08	16 – 45[d]	20.74	6.21	16 – 37	$H_{(2,92)}= 32.99, p < 0.01$	CON vs. schizophrenia, $p < 0.01$; CON vs. MAP, $p < 0.01$
Total score	30.48	1.12	30 -35	48.86	15.56	30 – 91[e]	45.63	18.68	30 – 97	$H_{(2,92)}= 46.64, p < 0.01$	CON vs. schizophrenia, $p < 0.01$; CON vs. MAP, $p < 0.01$

a CON longer duration of school education compared to schizophrenia and MAP

b CON longer duration total education than schizophrenia and MAP

c schizophrenia greater duration of illness compared to MAP

d schizophrenia higher scores than CON and MAP

e schizophrenia lower scores than MAP

Abbreviations: CON – healthy controls; MAP – methamphetamine-induced psychosis. PANSS – Positive and Negative Syndrome Scale; SD – standard deviation.

Table 2: Subcortical volume group differences across schizophrenia, methamphetamine-induced psychosis and health controls

Regions	All groups			SCZ-CON	MAP-CON	MAP-SCZ
	Effect size $\eta\rho^2$	CI: 90%	p	p	p	p
Left hippocampus	0.012	[0, 0.056]	0.57	n.s.	n.s.	n.s.
Right hippocampus	0.051	[0, 0.13]	0.090	n.s.	n.s.	n.s.
Left amygdala	0.012	[0.028, 0.22]	0.0026	0.012	0.054	n.s.
Right amygdala	0.029	[0, 0.091]	0.26	n.s.	n.s.	n.s.
Left caudate	0.082	[0.0075, 0.17]	0.020	0.03	n.s.	n.s.
Right caudate	0.08	[0.0062, 0.17]	0.023	0.05	n.s.	n.s.
Left putamen	0.13	[0.033, 0.23]	0.0017	0.016	n.s.	n.s.
Right putamen	0.10	[0.017, 0.2]	0.0068	0.0072	n.s.	n.s.
Left globus pallidus	0.035	[0, 0.1]	0.19	n.s.	n.s.	n.s.
Right globus pallidus	0.076	[0.052, 0.16]	0.026	n.s.	n.s.	0.021
Left nucleus accumbens	0.29	[0.16, 0.39]	< 0.0001	< 0.0001	n.s.	< 0.0001
Right nucleus accumbens	0.14	[0.043, 0.25]	0.0007	0.014	n.s.	0.0017

Effect size: for all group differences. *MANOVA (F(2, 91)) was performed for subcortical volumes in methamphetamine-induced psychosis (MAP), schizophrenia (SCZ) and healthy controls (CON) groups, with p-values of <0.05 considered statistically significant.

n.s : non-significant. Effect sizes: partial eta squared (Confidence Interval: 90%)

Table 3: Frontal cortical thickness group differences

Regions	All groups			SCZ-CON	MAP-CON	MAP-SCZ
	Effect size[*#]	*CI: 90%*	*p*	*p*	*p*	*p*
Left caudal middle frontal*	0.44	[0.31, 0.53]	< 0.0001	< 0.0001	< 0.0001	n.s.
Right caudal middle frontal*	0.28	[0.15, 0.39]	< 0.0001	< 0.0001	< 0.0001	n.s.
Left lateral orbitofrontal#	0.65	[0.64, 065]	< 0.0001	< 0.0001	n.s.	< 0.0001
Right lateral orbitofrontal*	0.19	[0.074, 0.29	< 0.0001	n.s.	0.028	< 0.0001
Left medial orbitofrontal#	0.11	[0.11, 0.12]	0.022	n.s.	n.s.	0.0014
Right medial orbitofrontal#	0.057	[0.05, 0.064]	0.027	n.s.	0.042	n.s.
Left rostral middle frontal*	0.47	[0.34, 0.56]	< 0.0001	<0.0001	< 0.0001	n.s.
Right rostral middle frontal#	0.096	[0.091, 0.10]	0.0046	n.s.	n.s.	0.0032
Left superior frontal*	0.19	[0.071, 0.29]	< 0.0001	< 0.0001	0.013	n.s.
Right superior frontal*	0.085	[0.0091, 0.18]	0.016	0.015	n.s.	n.s.
Left frontal pole#	0.88	[0.87, 0.89]	< 0.0001	< 0.0001	0.0002	< 0.0001
Right frontal pole#	0.69	[0.67, 0.70]	< 0.0001	< 0.0001	< 0.0001	n.s.
Left pars opercularis*	0.64	[0.54, 0.71]	< 0.0001	< 0.0001	< 0.0001	< 0.0001
Right pars opercularis*	0.41	[0.27, 0.5]	< 0.0001	< 0.0001	< 0.0001	0.0001
Left pars orbitalis*	0.19	[0.071, 0.29]	< 0.0001	< 0.0001	0.0068	n.s.
Right pars orbitalis*	0.23	[0.11, 0.34]	< 0.0001	< 0.0001	0.0002	n.s.
Left pars triangularis*	0.39	[0.25, 0.49]	< 0.0001	< 0.0001	n.s.	< 0.0001
Right pars triangularis#	0.31	[0.30, 0.32]	< 0.0001	0.0004	n.s.	< 0.0001

Effect size for all group differences. *MANOVA (F(2,91)) and #Kruskal Wallis ANOVA (H(2,94)) performed for frontal cortical thickness in methamphetamine-induced psychosis (MAP), schizophrenia (SCZ) and healthy controls (CON) groups, with *p*-values of <0.05 considered statistically significant. n.s : results which did not survive the MANOVA and Kruskal Wallis ANOVA. Effect sizes, *partial eta squared (Confidence Interval: 90%) and #epsilon squared (Confidence Interval: 90%)