1   **Coral species-specific loss and physiological legacy effects are elicited by**

2   **extended marine heatwave**

4   Strand, E.L.[1*], Wong, K.H.[1,2], Farraj, A.[1], Gray, S.[1], McMenamin, A.[1], Putnam, H.M.[1]
5
[1] University of Rhode Island
[2] Department of Marine Biology and Ecology, University of Miami Rosenstiel School of Marine
and Atmospheric Science, 4600 Rickenbacker Causeway, Miami, FL, 33149, USA.
[*]corresponding author

Key Words

• Physiological legacy
• Double-stressor experiment
• Thermal and ocean acidification tolerance
• Coral-algal symbiosis
• Multivariate statistics
• Symbiodiniaceae communities
• Stress and recovery periods
19

Abstract

Marine heatwaves are increasing in frequency and intensity, with potentially catastrophic

consequences for marine ecosystems such as coral reefs. An extended heatwave and recovery

time-series that incorporates multiple stressors and is environmentally realistic can provide

enhanced predictive capacity for performance under climate change conditions. We exposed

common reef-building corals in Hawaiʻi, *Montipora capitata* and *Pocillopora acuta*, to a 2-month

period of high temperature and high pCO_2 conditions or ambient conditions in a factorial design,

followed by 2 months of ambient conditions. High temperature, rather than high pCO_2, drove

multivariate physiology shifts through time in both species, including decreases in respiration

rates and endosymbiont densities. *Pocillopora acuta* exhibited more significantly negatively

altered physiology, and substantially higher bleaching and mortality than *M. capitata*. The

sensitivity of *P. acuta* appears to be driven by higher baseline rates of photosynthesis paired with

lower host antioxidant capacity, creating an increased sensitivity to oxidative stress. Thermal

tolerance of *M. capitata* may be partly due to harboring a mixture of *Cladocopium* and

Durusdinium spp., whereas *P. acuta* was dominated by other distinct *Cladocopium* spp. Only *M.*

capitata survived the experiment, but physiological state in heatwave exposed *M. capitata*

remained significantly diverged at the end of recovery relative to individuals that experienced

ambient conditions. In future climate scenarios, particularly marine heatwaves, our results indicate

a species-specific loss of corals that is driven by baseline host and symbiont physiological

differences as well as Symbiodiniaceae community compositions, with the surviving species

experiencing physiological legacies that are likely to influence future stress responses.

Introduction

Coral holobionts, a meta-organism composed of an animal coral host, endosymbiont

dinoflagellates and associated microbial community (Rohwer et al., 2002), make up the structure

of one of the most ecologically diverse and economically valuable ecosystems in the world: coral

reefs. The function and health of the coral holobiont, and therefore coral reefs, is dependent on

this multipartner symbiotic relationship that cycles nutrients and provides each partner with

otherwise limited or unavailable nutritional resources (Bourne et al., 2016). However, climate

change is impacting the marine environment through increasing ocean temperatures (IPCC,

2018), specifically marine heatwaves (Oliver et al., 2021) and ocean acidification (Gattuso et al.,

2015), among other local stressors, challenging the symbiotic function, physiological processes

and, ultimately, the survival of reef-building corals. Although ocean acidification will place a

chronic energetic strain on coral calcification (Mollica et al., 2018) and cellular acid–base balance

(Tresguerres et al., 2017; Gibbin et al., 2014), the dominant physiological stress is currently in the

form of increasingly frequent, intense marine heatwaves (Oliver et al., 2018; Sweet and Brown,

2016). This thermal stress triggers dysbiosis known as 'coral bleaching', or the disruption of the

symbiotic relationship between a coral host and their photosynthetic endosymbionts,

Symbodinicaeae. This results in the coral host expelling the endosymbionts, leaving a stark white

('bleached') appearance and the coral host depleted of their primary carbon source (Gates et al.,

1992; Boilard et al., 2020). Mass coral bleaching events resulting from marine heatwaves have

increased in frequency and intensity in recent years (Oliver et al., 2021), and this trend is on track

to continue, with catastrophic consequences projected for coral reef ecosystems (Heron et al.,

2016; Le Nohaïc et al., 2017; van Woesik and Kratochwill, 2022).

The current understanding of the mechanistic cascade of bleaching is linked to the

response of the host and/or endosymbionts to increased reactive oxygen species (ROS) levels

produced under higher temperature that consequently increase oxidative stress and cause a

breakdown of photosynthetic processes within the symbiont cells (Weis, 2008; Gates et al., 1992).

This breakdown causes the symbiont cells to be either digested or expelled via several

mechanisms including exocytosis, apoptosis, necrosis and host cell detachment (Weis, 2008;

Lesser, 2004; Gates et al., 1992). A review of recent studies has shown that oxidative stress can

occur in the host first (Oakley and Davy, 2018), suggesting that host antioxidant capacity may be

critical in tolerating higher levels of temperature stress. However, host response to changing

environment conditions has been shown to be highly species-specific (Bahr et al., 2016). In

Kāneʻohe Bay, Oʻahu, Hawaiʻi, two common reef-building coral species display differential thermal

tolerance: *Montipora capitata* is more resistant and *Pocillopora acuta* is more sensitive to

environmental stress (Bahr et al., 2015; Gibbin et al., 2015). The underlying mechanisms to

differential tolerance are known to be complex, ranging from genetic influences, epigenetic

patterns, and associated endosymbiont and bacterial communities (Putnam, 2021), to host

morphological structure and tissue thickness differences (Loya et al., 2001), but are not fully

understood. Thus, multivariate analyses of key processes and traits can provide the capacity to

quantify temporal variation in host, symbiont and holobiont responses, and are necessary to more

fully elucidate the bleaching cascade (Gardner et al., 2017; Wall et al., 2021; McLachlan et al.,

2021).

Although extensive research has been conducted on the effects of increased temperature

on coral (Mydlarz et al., 2010; Sweet and Brown, 2016; McLachlan et al., 2020; Cziesielski et al.,

2019; Lesser, 2011; van Oppen and Lough, 2018), there are fewer experimental studies that

simultaneously address critical environmentally relevant aspects that reflect a coral's natural

conditions. Specifically, there are few studies that concurrently: (1) mimic daily and seasonal

environmental fluctuations (van Woesik et al., 2022; Ziegler et al., 2021; Putnam and Edmunds,

2011), while (2) accounting for simultaneous, multivariate stressors (Pendleton et al., 2016; van

Woesik et al., 2022), (3) sampling with high frequency to capture short and long-term temporal

stress and recovery dynamics (Claar et al., 2020; Gardner et al., 2017), (4) during environmentally

realistic times of the year (vanWoesik et al., 2022; Ziegler et al., 2021) and (5) tracking the

survivors following the stress exposure (Claar et al., 2020; Gardner et al., 2017). These factors

are particularly important, as exposure to diurnal temperature and pCO_2 fluctuations elicit different

responses than stable, or less variable conditions (Putnam and Edmunds, 2011; Dufault et al.,

2012; Schoepf et al., 2022; Barshis et al., 2013; Wall et al., 2021). Further, high-frequency

sampling of a variety of variables was able to elucidate that symbiont expulsion and bleaching

precedes severe holobiont physiological responses and health decline in later stress time points

(Gardner et al., 2017). Additionally, seasonal studies of influences on coral physiology (Jurriaans

and Hoogenboom, 2020; Scheufen et al., 2017; Thornhill et al., 2011; Edmunds and Putnam,

2020) highlight the variation in traits based on recent environmental conditions, supporting the

need to study bleaching under seasonally realistic conditions to more accurately predict an

organism's stress response (e.g. conducting bleaching studies at the timing of the peak of

temperature for that region). Thus, high-frequency sampling of coral multivariate phenotypes

through time and following bleaching stress is necessary to understand the mechanisms and

sequential timing of the symbiotic state during stress and subsequent recovery.

Here, we tested the effect of an environmentally realistic extended heatwave in the context

of diurnally fluctuating temperature and ocean acidification on coral physiology to improve our

understanding of coral stress biology and forecasting reef futures. Specifically,we conducted a 4-

month multi-stressor time-series experiment during the fall warming season in Kāneʻohe Bay,

Oʻahu, Hawaiʻi, to assess the temporal dynamics during exposure to increased temperature and

partial pressure of CO_2 (pCO_2) in addition to seasonal change, to compare differential stress

response in a resilient and a sensitive coral species, and to investigate temporal dynamics during

recovery and the potential for physiological legacies after an environmental perturbation.

Methods

Coral Collections

Coral fragments (one ~3×3 cm fragment per colony from 75 colonies of *Montipora*

capitata Dana 1846 and 75 colonies of Pocillopora acuta Lamarck 1816) were collected via

clippers at ~3–4.5 m depth from each of six reef areas ranging across the north to south span of

Kāneʻohe Bay including both fringing and patch reefs [Lilipuna Fringe: 21°25'45.9"N

157°47'28.0"W; Hawaiʻi Institute of Marine Biology (HIMB): 21°26′ 09.8″ N 157°47′ 12.7″ W;

Reef.11.3: 21°27′ 02.9″ N 157°47′ 41.8″ W; Reef.18: 21°27′ 02.9″ N 157°48′ 40.1″ W;

Reef.35.36: 21°28′ 26.0″ N 157°50′ 01.2″ W; Reef.42.43: 21°28′ 37.9″ N 157°49′ 36.8″

W] under Hawaiʻi Department of Aquatic Resources Special Activity Permit 2019-60, between 4

and 10 September 2018 (Fig. 1A,B). This collection approach, ensuring each fragment was >10

m apart from each other, was designed to target a diverse collection of genotypes that represent

the coral population for *M. capitata* and *P. acuta* in Kāneʻohe Bay. Coral fragments were

brought back to HIMB and affixed in an upright position to individually numbered plastic coral

fragment mounting plugs by hot gluing the base of the skeleton to the plug (McMaster-Carr, Hot-

Melt Glue #7518A54). Coral fragments were allowed to acclimate for ~14 days in outdoor

mesocosm tanks and then were randomly allocated in an even distribution (n=405 *P. acuta* and

n=440 *M. capitata*) to 12 replicate tanks for 2 months (weeks 1–8) of temperature and pCO_2

stress exposure and 2 months of ambient condition recovery (weeks 9–16) as described below.

The goal of our coral collection design was to examine the population of corals in Kāneʻohe Bay.

Therefore, the coral fragments were randomly allocated to the tanks, so the site of collection

was not included statistically.

Experimental Timeline

Coral fragments were exposed to one of four experimental conditions – (1) ambient

temperature+ambient pCO_2 (ATAC), (2) ambient temperature+high pCO_2 (ATHC), (3) high

temperature+ambient pCO_2 (HTAC) and (4) high temperature+high pCO_2 (HTHC) – for 2

months (22 September–17 November 2018; Fig. 1C). Treatment conditions (n=3 replicate tanks

per treatment) were randomly assigned to 12 outdoor flow-through mesocosm tanks

(122×122×30 cm; 510 l) supplied with sand-filtered seawater from Kāneʻohe Bay. Flow rates,

measured daily with a graduated cylinder and timer, averaged 84.36±1.20 ml s^{-1} (n=826),

providing full mesocosm tank turnover every ~2 h. Mesocosm tanks were 60% shaded from full

irradiance, and photosynthetically active radiation (PAR) was measured continuously with an

Apex cosine corrected PAR Sensor (PMK, Neptune Systems, accuracy =±5%) that was cross-

calibrated to the Li-Cor cosine corrected PAR sensor (LI-193 spherical underwater quantum

sensor, LI-COR Biosciences). Additionally, light values (roughly 100–700 PAR) from six different

positions in each tank were compared to determine spatial differences within each tank. There

was no significant difference in light between positions in each tank (n=4 per position per tank;

position P=0.978; tank P=0.143). Based on these results, light was measured in the center of

the tank approximately daily for the duration of the experiment using the LI-193 spherical

underwater quantum sensor. To further reduce any potential position effects with respect to

incoming water, heater position or bubble stream, the positions of the coral fragments in the

tank were changed weekly. During this process, coral fragments were gently cleaned with

toothbrushes and tanks were scrubbed to remove excess algal growth.

Experimental temperature conditions

Temperature treatment conditions were programmed to mimic the natural daily

fluctuations of the environment at the collection sites in Kāneʻohe Bay, Hawaiʻi (NOAA Moku o

Loʻe Buoy data from September 2018, https://www.ndbc.noaa.gov/station_page.php?

station=mokh1; Fig. 1G; Fig. S1). Based on these data, the high temperature treatment

fluctuated between ~29 and 30°C to reflect previous marine heatwaves in Kāneʻohe Bay,

Hawaiʻi (+2°C above ambient temperature). Temperature was monitored with Apex Extended

Life Temperature Probes (accuracy =±0.05°C, Neptune Systems) and temperature loggers

(HOBO Water Temp Pro v2, accuracy=±0.21°C, resolution=0.02°C, Onset Computer Corp.) that

were placed in each tank at the same height as the coral fragments for the duration of the

experiment and logged temperature at 10 min intervals. Temperature was separately controlled

by submersible heaters (ProHeat D-1500 Heater Controllers, precision ±1°C). Ambient

temperature treatments reflected the natural conditions of Kāneʻohe Bay owing to the flow-

through nature of the tanks.

Experimental pCO_2 conditions

Experimental pCO_2 treatment conditions were based on measurements of ambient

conditions in Kāneʻohe Bay, Hawaiʻi (Drupp et al., 2011), and projected future lower pH

conditions in embayments, which can have lower mean pH and higher diel fluctuations owing to

calcification, photosynthesis and respiration of the benthic community and increased residence

time in the embayment (Jury et al., 2013; Shaw et al., 2016). The target pH range was based

on values reported from low pH measured on the HIMB patch reef in Kāneʻohe Bay (Guadayol

et al., 2014). The daily fluctuating pH levels in the high pCO_2 treatment tanks were maintained

between 7.6 and 7.7 (pH NBS scale) with an independent pH-stat feedback system in each

tank, and ambient conditions fluctuated between 7.9 and 8.0 (pH NBS scale; Fig. 1E,F). To

generate the high pCO_2 treatment, two 99.99% food-grade CO_2 cylinders were connected to an

automatic gas cylinder changeover system (Assurance Valve Systems, Automatic Gas

Changeover Eliminator Valves #6091) to prevent an abrupt shortage of CO_2 supply. CO_2 was

added into the seawater on-demand through gas flow solenoids (Milwaukee MA955), based on

the pH reading (NBS) of a lab-grade pH probe (Apex pH, Neptune Systems) in each tank. CO_2

was delivered via airlines plumbed into a Venturi injector (Forfuture-go G1/2 Garden Irrigation

Device Venturi Fertilizer Injector), which was connected to a water circulating pump

(Pondmaster Pond-mag Magnetic Drive Water Pump Model 5). Gas injected into the system

was either CO_2 or ambient air depending on treatment, and was injected through a Venturi

injector connected to a pressure-driven pump, resulting in constant bubbling and moving

water. An Apex AquaController (Neptune Systems) environmental control system with a wifi

base unit (Apex Controller Base Unit, Neptune Systems) was linked to 12 individual monitoring

units (Apex PM1 pH/ORP Probe Module, Neptune Systems) with Apex lab-grade pH Probes

(accuracy=±0.01 pH NBS scale, Neptune Systems), which were used for microprocessor-

control of a power strip (Apex Energy Bar 832, Neptune Systems) containing 12 individual

solenoids. This pH-stat feedback system constantly monitored seawater temperature and pH

conditions. Once sampling and treatments were completed, the pCO_2 injections were shut off

immediately in all tanks and the temperature gradually decreased over the course of a week

(Fig. 1G), and fragments were exposed to ambient temperature and ambient pCO_2 conditions

for~2months (18November 2018–12 January 2019), identified here as the recovery period.

Total Alkalinity and Carbonate Chemistry

Tank parameters [temperature (°C), pH (total scale) and salinity (psu)] were measured

2–3 times daily using a handheld digital thermometer (Fisherbrand, Traceable Platinum Ultra-

Accurate Digital Thermometer, accuracy=±0.05°C, resolution=0.001°C) and a portable

multiparameter meter (Thermo Fisher Scientific, Orion Star A series A325). A pH probe (Mettler

Toledo, InLab Expert Pro pH probe #51343101; accuracy=±0.2 mV, resolution=0.1 mV) and

conductivity probe (Orion, DuraProbe 4-Electrode Conductivity Cell Model 013010MD;

accuracy=0.5% of psu reading, resolution=0.01 psu) were used with an Orion A Star meter to

measure pH (mV) and salinity (psu), respectively. pH (total scale) was calculated from standard

curves of pH (mV) across a range of temperature (°C) in a tris standard (Dickson Laboratory,

Tris batch T27 bottles 269 and 236, and batch T26 bottle 198). Water samples (125 ml) were

taken from each tank twice a week to measure total alkalinity and, along with the pH on the total

scale, as described above, to calculate carbonate chemistry. An automated titrator (Mettler

Toledo T50) was used to titrate water samples with salinity-adjusted 0.1 mol l−1 hydrochloric

acid (Dickson Laboratory Titrant A3, A14). A non-linear, least-squares procedure of the Gran

approach (SOP 3b; Dickson et al., 2007) was used to calculate total alkalinity (TA; $\mu mol\ kg^{-1}$

seawater). Accuracy of the titrations was determined using certified reference material [Dickson

Laboratory CO_2 Certified Reference Material (CRM) batch 132 (salinity=33.441,

TA=2229.24±1.03 $\mu mol\ kg^{-1}$), batch 137 (salinity=33.607, TA=2231.59±0.62 $\mu mol\ kg^{-1}$), batch

176 (salinity=33.532, TA=2226.38±0.53 $\mu mol\ kg^{-1}$); NOAA Ocean Carbon and Acidification

Data System]. Tank total alkalinity values were only accepted if the total alkalinity value of the

CRM from that day was within 1%of the reported CRM. Carbonate values, including aragonite

saturation (Ω), carbon dioxide (CO_2), carbonate (CO_2^{-3}), dissolved inorganic carbon (DIC),

bicarbonate (HCO^{3-}), partial pressure of carbon dioxide (pCO_2) and pH, were calculated using

SEACARB (v3.2.16 in R Studio; https://CRAN.R-project.org/package=seacarb) with the

following parameters: calculated total pH, calculated TA, flag=8 [pH and ALK given], P=0, Pt=0,

Sit=0, pHscale='T', kf='pf', k1k2='l' and ks='d'. The equilibrium constant of hydrogen fluoride

(mol kg−1) (kf='pf') was set to 'pf' for salinities ranging from 10 to 40 psu and temperature

ranging from 9 to 33°C (Perez and Fraga, 1987) and recommended by Dickson et al. (2007).

The first (K1) and second (K2) dissociation constants of carbonic acid (mol kg−1) were set to 'l'

(k1k2='l') for salinities ranging from 19 to 43 psu and temperature ranging from 2 to 35°C

(k1k2='l'; Lueker et al., 2000). The stability constant of hydrogen sulfate (mol kg−1) (Ks) was set

to 'd' (Dickson, 1990) for salinities ranging from 5 to 45 psu and temperature ranging from 0 to

45°C. Salinity and temperature inputs were the measurements taken at the time of the water

sample collection.

Physiological Response Variables

Physiological sampling occurred at the following time points: 1, 2, 4, 6, 8, 12 and 16 weeks

(Fig. 1D). Physiological samples were immediately snap-frozen in liquid nitrogen after

photosynthetic and respiration rates were measured (described below; n=2 per tank×3 tanks per

treatment=6 fragments per treatment per species per time point). All samples were stored at

−80°C until subsequent processing.

Survivorship, growth rates, and color score

Survivorship was tracked daily throughout the 2-month exposure and 2-month recovery

periods using a binary score for dead versus alive. Alive was defined as any fragment with live

tissue. Growth over time was determined by the buoyant weight technique (Davies, 1989; Jokiel

et al., 1978) using air, freshwater and saltwater standard curves as a function of temperature.

Each coral fragment was buoyant weighed in its respective treatment temperature and pCO_2

conditions. Wet buoyant mass was converted to dry mass using the aragonite density of 2.03 g

cm^{-3} for Montipora spp. (Anthony and Hoegh-Guldberg, 2003) and 2.93 g cm^{-3} for *P. acuta*

(Jokiel et al., 1978). Rates were normalized to surface area, determined by a single-dip wax

dipping technique (Veal et al., 2010), and reported in units of mg $CaCO_3$ cm^{-2} day^{-1}. To assess

tissue color change over time as a proxy for cell density and pigmentation sensu Edmunds et al.

(2003), each coral fragment was photographed once a week (16 time points across the 4-month

experiment) with a red, green, blue color standard. ImageJ was used to extract the mean red,

green and blue color score for each coral fragment and these values were normalized to the

mean of the red, green and blue color standards, respectively. The color scorewas quantified as

principal component 1 (PC1; using the princomp function in R) from the principal component

analysis (PCA) (sensu Edmunds et al., 2003).

Photosynthesis and Respiration Rates

Prior to experimental exposures, 10 coral fragments (5 per species) were used to generate

a photosynthetic irradiance (PI) curve to determine saturating irradiance for assessing rates of

photosynthesis. Fragments were exposed to 10 light levels: 0, 15, 30, 60, 91, 136, 227, 416, 529

and 756 μmol photons m^{-2} s^{-1} (as measured by the Apex PAR sensor described above) generated

by two LEDs (Aqua Illumination Hydra FiftyTwo) hung above the incubation chambers

(described below). Rates of oxygen consumption or evolution were extracted using local linear

regressions in the LoLinR package (Olito et al., 2017) in μmol O_2 l^{-1} s^{-1}. Rates were corrected to

chamber volume including coral displacement. Rates of the blank chambers were subtracted from

the rates of the coral fragments within each run to account for background photosynthesis or

respiration owing to phytoplankton, zooplankton and bacteria. Rates of oxygen flux were

normalized to surface area for final units of μmolO_2 cm^{-2} h^{-1}. Curve fitting of a non-linear least

squares fit for a non-rectangular hyperbola (Marshall and Biscoe, 1980; Heberling and Fridley,

2013) was used to identify PI curve characteristics of each species. To calculate net

photosynthesis (Pnet; μmol O_2 h^{-1}), the following PI curve fit equation was used:

$$Pnet = \frac{\phi PPFD + \sqrt{(\phi PPFD + P_{max})2 - 4\Theta \phi PPFD + P_{max}}}{2\Theta} - Rd$$ [1]

where PPFD (photosynthetic photon flux density) is the irradiance (μmol photons m^{-2} s^{-1}), areal

Pnet is net photosynthesis normalized to tissue area covered (cm^2) and cellular Pnet is net

photosynthesis normalized to symbiont cell density (number of cells), Pmax is the maximum

gross photosynthetic rate (μmol O2 cm^{-2} h^{-1}), φ is the apparent quantum yield (mol O_2 mol

photons^{-1}) and Rd is the light-enhanced dark respiration rate (Edmunds and Davies, 1988)

(μmol O2 cm^{-2} h^{-1}).

Photoinhibition, as represented by Ik,was calculated as Pmax divided by the apparent

quantum yield. PI curves were performed four times throughout the experiment to ensure the

coral fragments were not reaching photoinhibition (Fig. S4). Fragments were placed in individual

respiration chambers (~610 ml), with individual temperature (Pt1000 temperature sensor,

PreSens Precision Sensing GmbH) and fiber-optic oxygen probes (Oxygen Dipping Probes

DPPSt7, accuracy=±0.05% O_2, PreSens Precision Sensing GmbH) connected to a 10-channel

oxygen meter (OXY-10 ST, accuracy= ±1.0°C, resolution=0.1°C, PreSens Precision Sensing

GmbH) to evaluate photosynthesis (under saturating light conditions, as determined by PI

curves described above) and light-enhanced dark respiration (Edmunds and Davies, 1988). The

respirometry setup consisted of 10 chambers with stir bars and respective treatment water.

Samples were measured in a series of runs that consisted of eight fragments (n=4 per species)

and two blank chambers per run exposed to PAR irradiance of ~500 μmol photons m^{-2} s^{-1} for

15 min to assess photosynthetic rates. Immediately afterwards, these fragments were exposed

to dark conditions (0 μmol photons m^{-2} s^{-1}) for 20min to assess light-enhanced dark respiration.

Following respirometry, fragments were immediately snap-frozen in liquid nitrogen and stored at

−80°C. Raw oxygen flux data (temperature corrected) were thinned every 40 s and rates of

oxygen flux were calculated with local linear regression as described above. Net photosynthesis

was calculated by subtracting the oxygen change of the dark runs (a

negative value) from the oxygen change of the light runs (a positive value), to isolate

photosynthetic rates of the symbionts from the light enhanced dark respiration of the coral

holobiont.

Tissue removal, surface area, and endosymbiont density

Coral tissue was removed from the coral skeletons by airbrushing (Iwata Eclipse HP-

BCS) with ice-cold 1× phosphate buffered saline (PBS) solution, and the total volumewas

recorded. The tissue slurry was then homogenized for 15 s at high speed (PRO Scientific Bio-

Gen PRO200 Homogenizer, 10 mm Homogenizer Probe). Of the total tissue slurry, aliquots of

500 μl were taken for endosymbiont density, 500 μl for host soluble protein and host total

antioxidant capacity (TAC), 1 ml for chlorophyll a (*chl a*) concentration assays, and 5 ml for ash

free dry weight (AFDW). The skeleton was rinsed in freshwater and soaked in 20% bleach for

24 h and dried at room temperature for 24 h. Symbiodiniaceae cell concentration was quantified

with six technical separately loaded replicate counts on a hemocytometer (Hausser Scientific,

Bright-Line Counting Chamber) after homogenizing the tissue slurry for 1 min. The host

tissue was homogenized so that the symbiont cells were free of host tissue and clearly visible

on the counting slide. The average count (cells ml^{-1}) was standardized to the airbrushed

homogenate volume and surface area to produce cell density values in cells cm^{-2}.

Host soluble protein and total antioxidant capacity

To quantify host soluble protein and host TAC, the host tissue was isolated from

symbionts by centrifugation at 10,000 g for 10 min at 4°C. Soluble host protein of the

supernatant was measured using the Pierce BCA Protein Assay Kit (Pierce Biotechnology,

Waltham, MA, USA, catalog no. 23225) with two duplicates per sample measured by a 96-well

microplate reader (BioTek Synergy HTX Multi-Mode Reader Model S1LFA) at a wavelength of

562 nm. Resulting values were calculated from a bovine serum albumin standard curve and

standardized to homogenate volume and surface area to obtain host soluble protein values (mg

cm^{-2}). Host TAC was quantified in duplicate per sample with the Cell BioLabs OxiSelect Total

Antioxidant Capacity (TAC) Assay Kit (catalog no. STA-360) according to the manufacturer's

instructions and compared with a uric acid standard curve. Antioxidants in the sample reduce

Copper II to Copper I, resulting in positive correlation between copper reducing equivalents

(CRE, mmol l^{-1}) and antioxidant levels. CRE values were standardized to total soluble protein

for final values in units of CRE µmol mg^{-1}.

Chlorophyll-a concentration

To isolate symbiont cells for *chl a* concentration measurements, a 1 ml aliquot of tissue

homogenate was centrifuged at 13,000 g for 3 min and the supernatant was discarded. Then, 1

ml of 100% acetone was added to the symbiont pellet, mixed for 15 s on a lab vortex mixer, and

incubated in the dark at 4°C for 24 h. Samples and acetone blanks were read in duplicate (200

µl each) on a 96-well microplate reader at λ=630 and 663 nm. *Chl a* concentrations were

calculated with the following dinoflagellate equation for 100% acetone: *chl a*=11.43E663–

0.64E630 (Jeffrey and Humphrey, 1975). Concentration values were multiplied by 0.584 to

correct for the path length of the plate for units of µg ml^{-1} and standardized to the airbrushed

homogenate volume, normalized by surface area (µg cm^{-2}) and per symbiont cell (µg $cell^{-1}$).

Tissue Biomass

Tissue biomass was quantified using the AFDW method. Specifically, 5 ml of tissue

slurry was centrifuged at 13,000 g for 3 min. A volume of 4 ml of the supernatant host tissue

was removed and placed in preburned aluminum pan weighed on an analytical balance (Mettler

Toledo, model ML203T/00) and the symbiont pellet was resuspended in 1 ml PBS solution and

added to separate pre-weighed burned aluminum pans. The samples were placed in a drying

oven (Thermo Fisher Scientific, Heratherm General Protocol Oven, catalog no. 51028112) for

24 h at 60°C, weighed, and then placed in a muffle furnace for 4 h at 500°C (Thermo Fisher

Scientific, Lindberg Blue M Muffle Furnace, catalog no. BF51728C-1). AFDW (mg cm^{-2}) of the

host and symbiont fractions was calculated as the post-drying oven mass (dry mass)–post-

muffle furnace mass, and final values were standardized to the airbrushed homogenate volume

and normalized to surface area. Symbiont to host (S:H) biomass ratios were calculated by

dividing the symbiont AFDW by the host AFDW.

Univariate statistical analysis

370   Owing to the near-complete mortality during the recovery period in *P. acuta*, only the

exposure period was analyzed for the following physiological variables: tissue biomass, *chl a*,

endosymbiont density, respiration rates, photosynthetic rates, host TAC and host soluble

protein. *Montipora capitata* responses were analyzed in the exposure and recovery period

separately to assess the effect of treatment on physiological state and the effect of treatment

history on physiological state during a recovery period. Net photosynthetic rates per cm^2 from

the exposure period in *M. capitata* and color score for both species were square root

transformed to meet assumptions of analysis. Gross photosynthetic rates per cm^2 and growth

rates in *M. capitata*, and growth rates and *chl a* per cell in *P. acuta* were log10 transformed for

the exposure periods. Endosymbiont density, symbiont AFDW per cell, host soluble protein and

host TAC were log transformed in both *M. capitata* and *P. acuta*. All physiological responses

(tissue biomass, *chl a* concentration, endosymbiont density, respiration rates, photosynthetic

rates, host TAC and host soluble protein) were analyzed with linear mixed-effects models (lme4

package version 1.1-27.1; Bates et al., 2015) with time, temperature and pCO$_2$ as fixed factors,

tank number as a random factor, and sum effect coding (Kugler et al., 2018) to test for

interaction effects. The goal of our coral collection design was to examine the population of

corals in Kāneʻohe Bay. Therefore, the coral fragments were randomly allocated to the tanks, so

site of collection was not included statistically, as there was not a balanced design across the

tanks. Color score and growth rates were analyzed using linear mixed-effects models with time,

temperature and pCO_2 as fixed factors and plug ID nested within tank number as a random

repeated measures factor. Survivorship probability was calculated with a Cox proportional

hazards mixed effects model fit by maximum likelihood using the coxpH function in the coxme

package (version 2.2-16; Therneau and Grambsch, 2013) with temperature and pCO_2

treatments as fixed effects and plug ID as a random factor. Type III Wald χ2 ANOVA tests were

performed on all models to assess significance of fixed effects. The normality and variance

assumptions of the ANOVA test were assessed with quantile–quantile plots and Levene's test.

To determine baseline differences at week 1 in physiology at between the two species,

treatment replicates were grouped for each response variable (n=6 per treatment×4

treatments=24 per species), prior to significant physiological change, and a Welch two-sample t-

test was used for each univariate measurement to assess significance.

Multivariate statistical analysis

A correlation matrix was calculated using the 'corrplot' function from the corrplot R package

(https://CRAN.R-project.org/package=corrplot) to determine which univariate physiology

variables were correlated and thus what to include in multivariate analyses. Gross and net

photosynthetic rates were highly correlated and as a result, only net photosynthetic rates were

used in downstream analyses as measure of symbiont photophysiology without the influence of

cellular respiration. Multivariate physiological matrices were centered and scaled prior to

principal components analyses that were performed using the 'prcomp' function from the stats

package in R (https://www.r-project.org/) to examine multivariate physiological states for each

species in the exposure period, as well as the recovery period for *M. capitata*. Permutational

multivariate analysis of variance (PERMANOVA) tests were used to assess significant

differences in multivariate physiology with time, temperature and pCO_2 as fixed effects ('adonis'

function in the R package vegan; https://CRAN.Rproject.org/package=vegan) separately for

exposure and recovery periods for each species.

Molecular response variables

Molecular sampling occurred at the following time points: 0, 6, 12 and 30 h, and 1, 2, 4,

6, 8, 12 and 16weeks (Fig. 1D). At each time point, one fragment was selected at random from

each tank of each species (n=1 per tank×3 tanks per treatment=3 fragments per treatment per

species) and placed in sterile whirlpak bags and snap-frozen in liquid nitrogen at 13:00 h each

day (with the exception of 6, 12 and 30 h time points).

DNA and RNA Extractions

DNA and RNA were extracted simultaneously from the coral fragments that had been

snap-frozen and stored at −80°C. A small piece (∼0.5–1 cm) was clipped off using clippers

sterilized in 10% bleach, deionized water, isopropanol and RNAse free water, and then placed in

2 ml microcentrifuge tube containing 0.5 mm glass beads (Fisher Scientific, catalog no. 15-340-

152) with 1000 µl of DNA/RNA shield. A two-step extraction protocol was used to extract RNA

and DNA for RNASeq and DNA methylation, respectively, with the first step as a 'soft'

homogenization to reduce shearing of RNA or DNA. Tubes were vortexed at high speed for 1 and

2 min for *P. acuta* and *M. capitata* fragments, respectively. The supernatant was removed and

designated as the 'soft extraction'. In the second step, 500 µl of DNA/RNA shield was added to

the bead tubes and placed in a Qiagen TissueLyser for 1 min at 20 Hz. The supernatant was

removed and designated as the 'hard extraction'. Subsequently, 300 µl of samples from both soft

and hard homogenate were extracted individually with the Zymo Quick-DNA/RNA Miniprep Plus

Kit (Zymo Research, catalog no. D7003T) following the protocol instructions, with the following

modifications. For the DNA elution step, the first elution of 10 µl of warmed 10 mmol l^{-1} Tris HCl

was added to DNA columns for a 15 min room temperature incubation and centrifuged at 16,000

g for 30 s. A second elution of 100 µl of warmed 10 mmol l^{-1} Tris HCl was added for a 5 min

incubation and centrifuged at 16,000 g for 30 s. Broad-range DNA quantity (ng µl^{-1}) was measured

with a Thermo Fisher Qubit Fluorometer, and DNA quality was assessed using gel

electrophoresis. Only the second elution of the 'hard extraction' DNA elution was used for

downstream internal transcribed spacer 2 (ITS2) sequencing protocols and analysis.

Internal Transcribed Spacer 2 (ITS2) Sequencing

Custom primers were ordered from Integrated DNA Technologies with ITS2 region

locus-specific sequences based on Lajeunesse and Trench (2000) and Illumina adapters (in

italics) to create our ITS2 forward primer (5′ *GAATTG CAG AAC TCC GTG TCG TCG GCA*

GCGTCA GATGTG TATAAG AGACAG 3′) and ITS2 reverse primer (5′ *GGGATC*

CATATGCTTAAG TTCAGCGGGTGTCTCGTG GGC TCG GAG ATG TGT ATA AGA GAC AG

3′). PCR master mix solution included (per sample values): 16.6 µl of 2× Phusion HiFi

Mastermix (Thermo Scientific, catalog no. F531s), 14 µl of DNase/RNase freewater, 0.6 µl of 10

µmol l^{-1} ITS2 forward primer, 0.6 µl of 10 µmol l^{-1} ITS2 reverse primer and 1 µl of 10 ng µl^{-1}

DNA sample or 1 µl ofDNase/RNase freewater as a negative control. To avoid PCR

amplification bias, each sample was run in triplicate on 96-well plates with 32 µl of master mix

and 1 µl of 10 ng µl^{-1} DNA sample in each well. Samples were run on the following PCR

settings: 3 min at 95°C, then 35 cycles of 30 s at 95°C, 30 s at 52°C and 30 s at 72°C, followed

by 2 min at 72°C and held indefinitely at 4°C. The triplicate PCR products for each sample were

pooled and stored at −20°C. The PCR products were run on a 2% agarose gel to confirm 360–

390 bp pair fragment length (330–360 bp ITS2 region and locus-specific primers+33 bp adapter

length). Then, 45 µl of PCR product was sent to the RI-INBRE Molecular Informatics Core at the

University of Rhode Island (RI, USA) for the index PCR and sequencing. There, PCR products

were cleaned with NucleoMag beads (Takara Bio, catalog no. 744503) and then visualized

using gel electrophoresis. A second round of PCR was performed with 50 ng DNA input for 8

cycles to attach Nextera XT indices and adapters using the Illumina Nextera XT Index Kit

(Illumina, catalog no. FC-131-200×) and Phusion HF PCR master mix (Thermo Fisher Scientific,

catalog no. F531L). Products were then cleaned with NucleoMag beads and visualized using

gel electrophoresis (1.5% agarose gel, 110 V, 35 min). The quality of samples was assessed

using an Agilent BioAnalyzer DNA1000 chip (Agilent, catalog no. 5067-1505) and quantification

was performed using a Qubit fluorometer with a High Sensitivity Kit (Invitrogen, catalog no.

Q32584) prior to pooling all samples. The final pooled library was quantified using qPCR (Roche

LightCycler480) with the KAPA Biosystems Illumina Kit (KAPA Biosystems). Samples were

analyzed using 2×250 bp paired-end sequencing on an Illumina MiSeq (Illumina).

Bioinformatic analysis

ITS2 sequences fromrawFASTQ fileswere analyzed using SymPortal (symportal.org;

Hume et al., 2019) on theUniversity of Rhode Island's High Performance Cluster. This workflow

used Miniconda3 v4.9.2 and SymPortal v0.3.21-foss-2020b to assign sequences as defined

intragenomic variants (DIV) and 'ITS2 Type Profiles'. A raw counts matrix of the ITS2 Type Profile

output from SymPortal was exported and analyzed locally using R Software Version 4.0.3 in

RStudio Version 1.4.1103. Relative abundance and a Bray–Curtis distance matrix were calculated

using the 'vegan' package (Dixon, 2003). A PERMANOVA was performed for each species to

assess significant effects of time point, treatment and their interaction.

Results

At the end of the ~14-day acclimation period, 89.8% of *P. acuta* (n=405) and 97.8% of

M. capitata (n=440) fragments were alive. During the exposures, treatments showed no

significant variation in TA across the 16-week experiment (treatment P=0.776; ATAC=2159.06

±5.88, ATHC=2162.46±0.12, HTAC=2155.58±5.25, HTHC= 2156.79±0.058; Fig. S2).

Additionally, light conditions were not significantly different between treatments (treatment

P=0.558; ATAC=179.70±9.97, ATHC=186.54±10.20, HTAC=176.64±10.84,

HTHC=165.12±10.41; Fig. S3). High temperature treatments during the 8-week stress period

were maintained above 29°C (HTAC=29.47 ±0.004, HTHC=29.59±0.004; n=7691), whereas

ambient conditions, dictated by flow-through water, steadily declined from 28°C to 26°C

(Fig. 1G). Recovery temperature conditions reflected natural conditions of Kāneʻohe Bay with

declines from 26°C to 23°C by week 16 in mid-January (Fig. 1G,H).

High pCO_2 treatments (ATHC, HTHC) were maintained between 7.6 and 7.7 pH total

scale (ATAC=8.03±0.011, ATHC=7.67±0.014, HTAC=8.01±0.011, HTHC=7.64±0.019). During

the stress period, high pCO_2 treatments were characterized by significant decreases

in aragonite saturation (Ω; treatment×time P<0.0001), carbonate (CO_3^{2-}; treatment×time

P<0.0001) and pH (treatment×time P<0.0001), as well as increases in carbon dioxide (CO_2;

treatment P<0.0001), dissolved inorganic carbon (DIC; treatment×time P<0.0001), bicarbonate

(HCO^{3-}; treatment×time P<0.0001) and pCO_2 (treatment P<0.0001; Fig. 1E,F, Fig. S2). During

recovery, all treatments reflected the natural conditions of Kāneʻohe Bay, ~8 pH total scale (Fig.

1E). Additional univariate figures and summary statistical tables can be found at Open Science

Framework (OSF) doi:10.17605/OSF.IO/Q9HMC.

High temperature drives bleaching responses and low P. acuta survivorship

In high temperature treatments (HTAC and HTHC), both species displayed significant

color change relative to ambient, starting at 4 weeks in *P. acuta* and 6 weeks in *M. capitata*.

High temperature and ambient pCO_2 (HTAC) and high temperature high pCO_2 (HTHC)

treatments drove this decrease in color score in the exposure period in *M. capitata*

(time×temperature×pCO$_2$ P=0.0004) and *P. acuta* (time×temperature P<0.0001;

time×temperature×pCO$_2$ P<0.0001). Additionally, *P. acuta* color score decreased significantly

through the autumn season (time P<0.0001) and in high pCO$_2$ treatments over time

(time×pCO$_2$ P<0.0001; Fig. S6). Following this visual bleaching response in the exposure

period, both high temperature (HTHC, HTAC) and pCO$_2$ treatments (ATHC) elicited decreased

survivorship probability in both *M. capitata* (temperature P <0.0001; pCO$_2$ P=0.0007) and *P.*

acuta (temperature×pCO$_2$ P<0.0001; Fig. S5). In the recovery period, *M. capitata* fragments that

experienced high temperature treatments (HTHC and HTAC) increased in color score to

ambient values by week 16 (time×temperature×pCO$_2$ P=0.0124; Fig. S6). The decreases in

survivorship probability for *M. capitata* stopped once treatment conditions were removed.

However, *P. acuta* fragments did not recover color, and survivorship probability continued

to sharply decrease (Fig. S5).

High temperature decreased metabolic rates in both species, but P. acuta displays higher

baseline photosynthetic rates

There were significant decreases during the stress period (September to November)

regardless of treatment in both species for respiration (*M. capitata* time P=0.0005; *P. acuta* time

P<0.0001) and photosynthetic rates (*M. capitata* Pnet cm^{-2} time P=0.0060; *P. acuta* Pnet cm^{-2}

time P<0.0001; *P. acuta* Pnet cell−1 time P <0.0001). However, *P. acuta* displayed significantly

higher baseline (as defined by week 1) areal (Pnet cm^{-2} P=0.0002; Pgross cm^{-2} P=0.0013) and

per cell photosynthetic rates (Pnet cell^{-1} P=0.0045), as well as higher photosynthesis:respiration

ratios (P:R P=0.0023) than *M. capitata* (Fig. 2). During the exposure period, high temperature,

rather than high pCO$_2$, caused a decrease in respiration rates in *M. capitata* (temperature

P=0.0203) and *P. acuta* (time×temperature P=0.0412). Similarly, high temperature drove

significant decreases in net photosynthetic rates in *M. capitata* (time×temperature P=0.0079)

and *P. acuta* (time×temperature P<0.0001).When net photosynthetic rates were normalized per

symbiont cell, treatments did not affect *M. capitata* fragments (time×temperature×pCO$_2$

P=0.1833), but *P. acuta* fragments in high temperature treatments displayed a significant

decrease by week 6 (time×temperature P=0.0011). Concomitantly, by week 6 (out of 8

exposure weeks), *P. acuta* fragments also displayed a significant decrease in P:R in high

temperature treatments (time×temperature P=0.0139) and overall changes in metabolic rates

were more drastic in *P. acuta* fragments.

High pCO$_2$ stress depresses seasonal increase in growth rates in M. capitata and high

temperature decreases growth rates in both species

547   In both species, growth rates (mg CaCO$_3$ cm^{-2} day^{-1}) peaked in at week 4 of exposure

(October) and gradually decreased by week 8 (November) of the exposure period (*M. capitata*

time P<0.0001; *P. acuta* time P<0.0001) regardless of treatment. High temperature treatment

decreased growth rates in later exposure time points (time×temperature×pCO$_2$ P<0.0001) in

both species. Similarly, both pCO$_2$ and temperature treatments exhibited lower growth rates

through the entire exposure period (time×temperature×pCO$_2$ P<0.0001) in *P. acuta* fragments.

In *M. capitata*, corals in high pCO$_2$ treatments displayed significantly lower growth rates over

time (Time× pCO$_2$ P=0.0003). However, during the recovery period, *M. capitata* fragments that

experienced only high pCO$_2$ (ATHC) displayed increasing growth rates through week 16 (time×

pCO$_2$ P=0.0017), whereas fragments that experienced the high temperature treatments

(HTAC and HTHC) gradually increased to ambient levels by week 16 (time×temperature

P<0.0001).

Host physiology: biomass remained stable while temperature stress increased TAC and

decreased protein levels

On average, *M. capitata* exhibited higher baseline TAC than *P. acuta* (P=0.0008; Fig. 2). Host

TAC (CRE µmol mg^{-1}) remained constant in *M. capitata* through the exposure period regardless

of treatment (time×temperature×pCO$_2$ P=0.7028) but increased by week 8 in *P. acuta* under

high temperature in HTAC and HTHC (time×temperature P=0.0277). During recovery, *M.*

capitata fragments previously exposed to high temperature treatments showed reduced TAC at

week 12, which returned to ambient treatment values by week 16 (time×temperature P=0.0237).

Host soluble protein (mg cm^{-2}) was not significantly different between species at baseline, week

1 (P=0.29), but significantly decreased through time from September to January in both *M.*

capitata (time P<0.0001) and *P. acuta* (time P<0.0001). At week 6, protein concentration in *M.*

capitata was significantly greater in both high pCO$_2$ treatments (pCO$_2$ P=0.0002) in comparison

to HTAC and ATAC. Additionally, corals in high temperature treatments (HTHC and HTAC in

comparison with ATAC and ATHC) displayed a decrease in protein over time (temperature×time

P=0.0463). In *P. acuta* fragments, high temperature treatments decreased in protein at week 8

(time×temperature P=0.0490). Host AFDW (mg cm^{-2}) was not significantly different at baseline,

week 1 (P=0.22), or affected by treatment, but gradually and significantly decreased throughout

the stress and exposure period from September to January in *M. capitata* (time P=0.0050) and

P. acuta (time P=0.0188).

Symbiont and holobiont physiology: stress treatments caused an increase in symbiont

biomass but decreased chl a content and endosymbiont density

Symbiont biomass (mg AFDW cm^{-2}) was unaffected by either temperature or pCO$_2$

treatment (time×temperature×pCO$_2$; *M. capitata* P=0.3839, *P. acuta* P=0.3790), but significantly

decreased through the autumn season in both *M. capitata* (P=0.0013) and *P. acuta* (P=0.0028).

Despite no treatment effect on symbiont and host biomass (mg AFDW cm^{-2}), for *P. acuta* the

symbiont:host biomass ratio significantly decreased in high temperature treatments

(temperature P=0.0180) and decreased also over time (Time P=0.0090). *Montipora capitata*

fragments exhibited significantly higher baseline week 1 biomass (mg cm^{-2} P=0.0145; mg cell^{-1}

P=0.0079; Fig. 2) than *P. acuta*. In *M. capitata*, symbiont cellular biomass (mg AFDW cell^{-1})

significantly increased in corals in the high temperature treatments during exposure

(temperature×pCO$_2$ P=0.0133) and then returned to ambient levels by week 16

(time×temperature P=0.0041). In *P. acuta*, symbiont per cell biomass decreased from

September to January (P=0.0141), but significantly increased under both temperature and pCO$_2$

exposure (time×temperature P<0.0001; time×pCO$_2$ P=0.0146). Holobiont *chl a* (cm^{-2})

increased over the autumn season in *M. capitata* (time P=0.0003; Fig. 3G), but high

temperature exposure resulted in decreased *chl a* over time (time×temperature P=0.0089) that

remained low in the recovery period (temperature P<0.0001). In *P. acuta*, *chl a* decreased over

time (time P<0.0001) and in high temperature treatments over time (time×temperature

P<0.0001; Fig. 3H). In *M. capitata*, only time significantly affected cellular *chl a* cell^{-1} (time

P=0.0347), and in all fragments, regardless of treatment, *chl a* cell^{-1} increased from September

to January. However, in *P. acuta*, *chl a* per cell was significantly affected by temperature and

time (time×temperature P=0.0764). Endosymbiont density (cells cm^{-2}) decreased in high

temperature and pCO$_2$ treatments over time in *M. capitata* (time×temperature P=0.0100;

time×pCO$_2$ P=0.0249) and the magnitude of decline was greater in high temperature in *P. acuta*

(time×temperature P<0.0001). *Pocillopora acuta* endosymbiont density decreased by week 4 in

all treatments (time P<0.0001; Fig. 3F), but all *M. capitata* fragments displayed an increase in

densities in the same time period, followed decreased values by week 8 (time P<0.0001; Fig.

3E). By the end of the recovery period, *M. capitata* fragments that had experienced high

temperature treatments did not fully recover in symbiont cell densities (temperature P=0.0070),

despite being statistically indistinguishable in color score.

Multivariate physiology: high temperature exacerbates seasonal changes during stress

and drives physiological states during recovery

In both species, treatments exacerbated existing seasonal shifts in multivariate

physiological states (Fig. 4). In *P. acuta*, physiological state shifted over time (P<0.001) and in

high temperature treatments (time×temperature P<0.001). Similarly, *M. capitata* fragments

experienced a shift in physiological state over time (time P<0.001) owing to temperature

(temperature P=0.005), and this was different between pCO_2 treatments over time (time×pCO_2

P=0.038). PC1 explained 39.83% of the variance and separated the samples between ambient

temperature and high temperature for *P. acuta*, with changes in respiration rates and host TAC

levels contributing most strongly to the separation (Fig. 4A,B). The PC2 axis explained 18.40%

of the variation and the separation between treatment groups was characterized by changes in

endosymbiont density and cellular symbiont biomass. In *M. capitata*, the PC1 axis explained

30.51% of the variation and was characterized by changes in the host (protein, biomass,

respiration) and holobiont physiology (areal net photosynthetic rates, areal symbiont AFDW and

holobiont *chl a* content) that contribute to separation primarily between time points. The PC2

axis explained 25.81% of the variation and was characterized by symbiont functions (symbiont

AFDW mg cell^{-1}, net photosynthetic rates and *chl a* content) that separate primarily treatment

groups (Fig. 4C,D). During the recovery period, both time (P<0.001) and high temperature

treatment (P<0.001) significantly affected physiological shifts in *M. capitata* (Fig. 5C,D). PC1

explained 37.61% of the variation and was characterized by host (biomass, protein) and

holobiont traits (mg AFDW cm^{-2}, *chl a* cm^{-2}) that separate between the history of high

temperature treatment groups in *M. capitata*. PC2 explained 30.11% of the variation and

changes in endosymbiont density and symbiont traits (*chl a* cell^{-1}, S:H biomass ratio, biomass)

contributed strongly to the separation of treatment and time groups.

Montipora capitata and P. acuta harbor distinctly different Symbiodiniaceae community

compositions that do not change significantly during stress or recovery

The ITS2 sequencing generated 5,807,744 total raw reads from 251 samples, with an

average of 22,775 reads per sample. The SymPortal analysis resulted in 4,902,971 total

cleaned sequences, which excludes week 16 in *P. acuta* because of low sample sizes.

Montipora capitata fragments harbored mixed Symbiodiniaceae communities composed of ITS2

symbiont type profiles with majority sequences 'D1/D4' and variations of 'C17d/No-Name'

(Durusdinium and Cladocopium spp.; LaJeunesse et al., 2018; Fig. 6). *Pocillopora acuta*

communities were dominated by type profiles in clade C with majority groups 'C1d/C42.2' and

'C1d'. Endosymbiont communities were significantly different between species (PERMANOVA

P<0.001) and differed only between two time points in *M. capitata* (12 h versus 24 h P=0.045)

but did not significantly change in exposure treatments (P=1.000).

Discussion

Extended heatwave scenario induces species-specific mortality of corals and creates

altered physiological legacies for the survivors

Scaling up from controlled single variable lab experiments to environmentally relevant

multi-stressor experiments is essential to capture the effects of stressors in the context of

seasonal physiological and thus the time-dependent and environmentally relevant sensitivity

of corals to global change stressors. Our examination of the effects of high pCO_2 and an

extended heatwave at the peak of seasonal temperatures on *M. capitata* and *P. acuta* reinforce

that increased temperatures are the more substantial threat to coral physiological

performance (Putnam et al., 2013), survivorship (Klein et al., 2022) and physiological legacy

effects (Wall et al., 2021) than high pCO_2 conditions. We found that high pCO_2 concentrations

alone can elicit potentially beneficial effects in the more resilient *M. capitata* such as higher

endosymbiont density and thus higher photosynthetic rates (cm^{-2}) during exposure, and higher

growth rates during recovery periods. However, these effects are not seen when high pCO$_2$

conditions occur simultaneously with higher temperatures. The extent of physiological shifts,

decreased survival probability and increased bleaching were all greater in *P. acuta* than in *M.*

capitata, reinforcing the outcome of species-specific susceptibility and loss of corals on future

reefs (Loya et al., 2001; van Woesik et al., 2011). Our data indicate that a combination of

Symbiodiniaceae community and baseline levels of host antioxidant capacity and cellular

photosynthetic rates are contributing to species-specific stress responses. Further, our findings

of different physiology trajectories of *M. capitata* during the recovery period, specifically in those

exposed to high temperature, support prior work that high temperature conditions leave lasting

physiological legacies several months into recovery (Rodríguez-Troncoso et al., 2016; Schoepf

et al., 2015; Rodrigues and Grottoli, 2007), and those legacies can have cumulative effects that

shift physiological baselines leading into a subsequent stress event (Wall et al., 2021; Safaie et

al., 2018; Ainsworth et al., 2016).

Symbiodiniaeceae communities and high levels of antioxidants relative to cellular

photosynthetic rates contribute to M. capitata thermal tolerance

Symbiodiniaceae communities aid in a coral holobiont's tolerance and survival (Baskett,

et al., 2009; Oliver and Palumbi, 2011); specifically, studies have documented that the presence

of Durusdinium spp. (Silverstein et al., 2015; Guest et al., 2016) and/or mixed endosymbiont

communities (Abbott et al., 2021) can be beneficial for coral holobiont survival. Our results

support these hypotheses, as the mixture of Durusdinimum spp. (D1/D4; ~60–80%) and

Cladocopium spp. (C17/C31; ~20–40%) in *M. capitata* was associated with higher thermal

tolerance, compared with *P. acuta*'s Cladocopium spp. (C1; 100%) dominated communities.

Although there was no evidence of Symbiodiniaceae community shuffling by treatment or

through time, a holobiont's community prior to an exposure event likely plays a crucial role in

their ability to handle changing environmental conditions. Through these community

composition differences, symbiont physiology influences holobiont tolerance (Hoadley et al.,

2019; Ziegler et al., 2015), thus indicating that both the host and symbiont physiological states

are critical to understanding the stress response of the holobiont (Bellantuono et al., 2012).

Differential photophysiology, carbon assimilation and translocation to the host (Stat et al., 2008)

and metabolite profiles (Matthews et al., 2017, 2018) have been seen between

Symbiodiniaceae genera and thus influence holobiont tolerance (Wall et al., 2020; Matthews et

al., 2018; Roach et al., 2021).

For both host and symbiont partners, a core physiological response to stress is the

activity of antioxidant systems to neutralize ROS molecules that build up during daily

metabolism and in response to stressful conditions such as increased temperature (Baird et al.,

2009; Downs et al., 2000). These excess ROS molecules can leak into the host to cause a

downstream cascade of cellular damage and eventually coral bleaching and apoptosis (Weis,

2008). A mismatch between the amount of symbiont-derived ROS molecules and the host's

antioxidant capacity may lead to higher susceptibility for coral bleaching (e.g. Cunning and

Baker, 2013). We hypothesize that *P. acuta*'s naturally higher cellular photosynthetic rates, but

lower host antioxidant capacity per unit protein, may lead to a faster build-up of ROS damage

and consequently higher mortality rates. Thus, *P. acuta* are challenged under thermal stress

because the symbiotic community is operating closer to the threshold between productivity and

damage. In comparison, *M. capitata*'s naturally lower photosynthetic rates and higher host

antioxidant capacity levels may be providing a greater buffer, allowing them capacity to support

and balance their higher endosymbiont densities during November (Fig. 3), resulting in a more

muted bleaching response when faced with stressful conditions. Further, *M. capitata* have the

benefit of thicker tissues (Loya et al., 2001; Magnusson et al., 2007) with greater energy

reserves (Rodrigues and Grottoli, 2007) in addition to their symbiont community composition. By

the end of the exposure period, *P. acuta* host antioxidant capacity levels increased, and cellular

photosynthetic rates decreased, but visual bleaching responses preceded this cellular

physiological change, indicating that the response was insufficient. *Montipora capitata* may be

well equipped to acclimatize to changing environmental conditions because of baseline

holobiont physiological states and existing Symbiodiniaceae communities.

Thermal exposure influences bleaching recovery trajectories in M. capitata

721   Recent studies have emphasized the importance of recovery time in between marine

heatwaves (Osborne et al., 2017; Neal et al., 2017; Grottoli et al., 2014). Recovery time can be

highly variable, with species taking anywhere from 1.5 months (Rodrigues and Grottoli, 2007), 2

months (Cunning et al., 2016), 3 months (Wall et al., 2019) up to 2 years (Matsuda et al., 2020).

We found that in our experimental recovery period of 2 months following exposure, physiological

metrics that decreased in chronic high temperature conditions stabilized as *M. capitata* fragments

began to recover pigmentation (color score, symbiont density and *chl a*). Despite full recovery in

color score and growth rates by week 16, *chl a* and endosymbiont densities did not fully recover

to the same extent as the ambient condition corals, with differential physiological states and

trajectories throughout the recovery period (Fig. 5). Our results indicate that extended high

temperature exposure at the peak of seasonal temperature leaves a physiological legacy that

lasts beyond 2 months. Even in this resistant and resilient species, legacy effects of bleaching on

organismal state can be as long as 1–2 years (Wall et al., 2019, Roach et al., 2021; Thomas and

Palumbi, 2017).

735   These legacy effect results are consistent with those of other studies that suggest that

environmental conditions shift coral phenotypes, ultimately eliciting a lasting legacy that could

have either a beneficial or detrimental priming effect for response in future stress events (Wall et

al., 2021; Hackerott et al., 2021; Putnam et al., 2017; Brown et al., 2000). However, recovery time

is imperative to understand whether the lasting effects of prior thermal events can be beneficial.

Sublethal temperatures that induce a bleaching response 10 days prior to a larger intensity

bleaching event have been shown to be beneficial (termed a 'protective trajectory'; Ainsworth et

al., 2016). Although beneficial effects were documented in long-term recovery periods of 11

months (Schoepf et al., 2015) following thermal perturbation, not all species recovery is on the

same timeline, with Porites divaricata and Orbicella faveolata recovering faster than Porites

astreoides (Schoepf et al., 2015). This potential for 'environmental memory' based on previous

environmental conditions is clear in corals (Brown et al., 2000, 2022). Utilizing this to elicit

purposeful stress hardening or environmental priming (Hackerott et al., 2021; Putnam et al., 2017;

Brown et al., 2000; DeMerlis et al., 2022) is a proposed restoration technique (van Oppen et al.,

2015), but it is clear there are limits of recovery time. *Montipora capitata* is likely to survive future

thermal stress events, but back-to-back bleaching events and decreased recovery time may

reduce their buffer and push their physiology closer to a threshold. More multi-year studies to

track differential bleaching patterns over time are needed to elucidate the threshold between

eliciting beneficial 'environmental memory' versus accumulated physiological damage and

energetic deficits (Schoepf et al., 2015).

Conclusion

This study highlighted the divergent thermal tolerance and recovery potential of *M.*

capitata and *P. acuta*, indicating a potential for species-specific susceptibility to stress and loss

of corals in future, more frequent and higher intensity bleaching events. *Montipora capitata*

elevated stress tolerance may be due in part to mixed Cladocopium and Durusdinium

Symbiodiniaceae communities, higher antioxidant capacity and elevated photosynthetic rates at

baseline, and stability of metabolic and symbiont physiology during chronic high temperature

conditions. In the 2 months of ambient recovery conditions that followed 2 months of elevated

temperature and pCO_2 conditions, *M. capitata* multivariate physiology did not fully recover. High

temperature treatment history elicited a lasting multivariate physiological legacy that set

recovering *M. capitata* on a different trajectory than those that experienced ambient conditions.

This study illuminates and emphasizes the importance of environmentally and seasonally

realistic time-series experiments to further understand the impacts of pre-existing seasonal

dynamics on thermal stress response. Multi-year repeated bleaching event studies are needed

to disentangle howa physiological legacy may be adaptive or maladaptive under predicted

climate change scenarios.

Contributions

Conceptualization: E.L.S., H.M.P.; Methodology: E.L.S., K.H.W., A.F., S.G., A.M., H.M.P.;

Software: E.L.S., H.M.P.; Validation: E.L.S., H.M.P.; Formal analysis: E.L.S., H.M.P.;

Investigation: E.L.S., H.M.P.; Resources: E.L.S., H.M.P.; Data curation: E.L.S., K.H.W., A.F.,

S.G., A.M., H.M.P.; Writing - original draft: E.L.S., H.M.P.; Writing - review & editing: E.L.S.,

K.H.W., A.F., S.G., A.M., H.M.P.; Visualization: E.L.S., H.M.P.; Supervision: E.L.S., H.M.P.;

Project administration: E.L.S., H.M.P.; Funding acquisition: H.M.P.

Data Availability

All data and code used to produce results and complete analyses are available at

https://github.com/hputnam/Acclim_Dynamics. Laboratory protocols and processing can be

found on Emma Strand's Open Laboratory Notebook at

https://emmastrand.github.io/EmmaStrand_Notebook/. ITS2 Sequencing FASTQ files are

uploaded to NCBI under Project #PRJNA761780.

Acknowledgements

We thank the Hawaiʻi Institute of Marine Biology and the Coral Resilience Laboratory for

hosting us to conduct the experiment on Coconut Island, and Chris Suchoki and Adam Helbig

for assistance with field work. Ariana Huffmyer provided valuable feedback on an earlier version

of the manuscript. As guests, we recognize and give thanks for the land and water resources of

the ʻā ina and the traditional owners of the land, kānaka ʻō iwi, on which this experimental work

was conducted in the Kāneʻohe Ahupuaʻa and the islands of Hawaiʻi. The authors acknowledge

use of the computational resources of the URI Center for Computational Research for this work.

References

Abbott, E., Dixon, G. and Matz, M. (2021). Shuffling between Cladocopium and Durusdinium
extensively modifies the physiology of each symbiont without stressing the coral host. Mol. Ecol.
30, 6585-6595. doi:10.1111/mec.16190
797
798   Ainsworth, T. D., Heron, S. F., Ortiz, J. C., Mumby, P. J., Grech, A., Ogawa, D., Mark Eakin, C.
799   and Leggat, W. (2016). Climate change disables coral bleaching protection on the Great Barrier
800   Reef. Science 352, 338-342. doi:10.1126/science.aac7125
801
802   Anthony, K. R. N. and Hoegh-Guldberg, O. (2003). Variation in coral photosynthesis, respiration
803   and growth characteristics in contrasting light microhabitats: an analogue to plants in forest
804   gaps and understoreys? Funct. Ecol. 17, 246-259. doi:10.1046/j.1365-2435.2003.00731.x
805
806   Bahr, K. D., Jokiel, P. L. and Rodgers, K. S. (2015). The 2014 coral bleaching and freshwater
807   flood events in Kāneʻohe Bay, Hawaiʻi. PeerJ 3, e1136. doi:10.7717/peerj.1136
808
809   Bahr, K. D., Jokiel, P. L. and Rodgers, K. S. (2016). Relative sensitivity of five Hawaiian coral
810   species to high temperature under high-pCO_2 conditions. Coral Reefs 35, 729-738.
811   doi:10.1007/s00338-016-1405-4
812
813   Baird, A. H., Bhagooli, R., Ralph, P. J. and Takahashi, S. (2009). Coral bleaching: the role of the
814   host. Trends Ecol. Evol. 24, 16-20. doi:10.1016/j.tree.2008.09.005
815
816   Barshis, D. J., Ladner, J. T., Oliver, T. A., Seneca, F. O., Traylor-Knowles, N. and Palumbi, S.
817   R. (2013). Genomic basis for coral resilience to climate change. Proc. Natl. Acad. Sci. U.S.A.
818   110, 1387-1392. doi:10.1073/pnas.1210224110
819

820 Baskett, M. L., Gaines, S. D. and Nisbet, R. M. (2009). Symbiont diversity may help coral reefs
821 survive moderate climate change. Ecol. Appl. 19, 3-17. doi:10.1890/08-0139.1
822
823 Bates, D., Ma¨chler, M., Bolker, B. and Walker, S. (2015). Fitting linear mixed effects models
824 using lme4. J. Stat. Softw. 67, 1-48. doi:10.18637/jss.v067.i01
825
826 Bellantuono, A. J., Hoegh-Guldberg, O. and Rodriguez-Lanetty, M. (2012). Resistance to
827 thermal stress in corals without changes in symbiont composition. Proc. Biol. Sci. R. Soc. 279,
828 1100-1107. doi:10.1098/rspb.2011.1780
829
830 Boilard, A., Dube´, C. E., Gruet, C., Mercie`re, A., Hernandez-Agreda, A. and Derome, N.
831 (2020). Defining coral bleaching as a microbial dysbiosis within the coral holobiont.
832 Microorganisms 8, 1682. doi:10.3390/microorganisms8111682
833
834 Bourne, D. G., Morrow, K. M. and Webster, N. S. (2016). Insights into the coral microbiome:
835 underpinning the health and resilience of reef ecosystems. Annu. Rev. Microbiol. 70, 317-340.
836 doi:10.1146/annurev-micro-102215-095440
837
838 Brown, B. E., Dunne, R. P., Goodson, M. S. and Douglas, A. E. (2000). Bleaching patterns in
839 reef corals. Nature 404, 142-143. doi:10.1038/35004657
840
841 Brown, K. T., Mello-Athayde, M. A., Sampayo, E. M., Chai, A., Dove, S. and Barott, K. L. (2022).
842 Environmental memory gained from exposure to extreme pCO variability promotes coral cellular
843 acid-base homeostasis. Proc. Biol. Sci. R. Soc. 289, 20220941. doi:10.1098/rspb.2022.0941
844
845 Claar, D. C., Starko, S., Tietjen, K. L., Epstein, H. E., Cunning, R., Cobb, K. M., Baker, A. C.,
846 Gates, R. D. and Baum, J. K. (2020). Dynamic symbioses reveal pathways to coral survival
847 through prolonged heatwaves. Nat. Commun. 11, 6097. doi:10.1038/s41467-020-19169-y
848
849 Cunning, R. and Baker, A. C. (2013). Excess algal symbionts increase the susceptibility of reef
850 corals to bleaching. Nat. Clim. Change. 3, 259-262. doi:10.1038/nclimate1711
851
852 Cunning, R., Ritson-Williams, R. and Gates, R. D. (2016). Patterns of bleaching and recovery of
853 *Montipora capitata* in Kāne'ohe Bay, Hawai'i, USA. Mar. Ecol. Prog. Ser. 551, 131-139.
854 doi:10.3354/meps11733
855
856 Cziesielski, M. J., Schmidt-Roach, S. and Aranda, M. (2019). The past, present, and future of
857 coral heat stress studies. Ecol. Evol. 9, 10055-10066. doi:10.1002/ece3.5576
858
859 Davies, S. P. (1989). Short-term growth measurements of corals using an accurate buoyant
860 weighing technique. Mar. Biol. 101, 389-395.
861
862 DeMerlis, A., Kirkland, A., Kaufman, M. L., Mayfield, A. B., Formel, N., Kolodziej, G., Manzello,
863 D. P., Lirman, D., Traylor-Knowles, N. and Enochs, I. C. (2022). Pre-exposure to a variable
864 temperature treatment improves the response of Acropora cervicornis to acute thermal stress.
865 Coral Reefs. 41, 435-445. doi:10.1007/s00338-022-02232-z
866
Dickson, A. G. (1990). Standard potential of the reaction: AgCl(s)+12H2(g)=Ag(s)+HCl(aq), and
the standard acidity constant of the ion HSO4 − in synthetic sea water from 273.15 to 318.15 K.
J. Chem. Thermodyn. 22, 113-127. doi:10.1016/0021-9614(90)90074-Z

Dickson, A. G., Sabine, C. L. and Christian, J. R. (ed.) (2007). Guide to Best Practices for
Ocean CO2 Measurements. PICES Special Publication 3. Sidney, BC: North Pacific Marine
Science Organization.

Dixon, P. (2003). VEGAN, a package of R functions for community ecology. J. Veg. Sci. 14,
927-930. doi:10.1111/j.1654-1103.2003.tb02228.x

Downs, C. A., Mueller, E., Phillips, S., Fauth, J. E. and Woodley, C. M. (2000). A molecular
biomarker system for assessing the health of coral (Montastraea faveolate) during heat stress.
Mar. Biotechnol. 2, 533-544. doi:10.1007/s101260000038

Drupp, P., De Carlo, E. H., Mackenzie, F. T., Bienfang, P. and Sabine, C. L. (2011). Nutrient
inputs, phytoplankton response, and CO2 variations in a semienclosed subtropical embayment,
Kaneohe Bay, Hawaii. Aquat. Geochem. 17, 473-498. doi:10.1007/s10498-010-9115-y

Dufault, A. M., Cumbo, V. R., Fan, T. Y. and Edmunds, P. J. (2012). Effects of diurnally
oscillating pCO_2 on the calcification and survival of coral recruits. Proc. R. Soc. B 279, 2951-
2958. doi:10.1098/rspb.2011.2545

Edmunds, P. J. and Davies, P. S. (1988). Post-illumination stimulation of respiration rate in the
coral Porites porites. Coral Reefs 7, 7-9. doi:10.1007/BF00301975

Edmunds, P. J. and Putnam, H. M. (2020). Science-based approach to using growth rate to
assess coral performance and restoration outcomes. Biol. Lett. 16, 20200227.
doi:10.1098/rsbl.2020.0227

Edmunds, P. J., Gates, R. D. and Gleason, D. F. (2003). The tissue composition of Montastraea
franksi during a natural bleaching event in the Florida Keys. Coral Reefs 22, 54-62.
doi:10.1007/s00338-003-0278-5

Gardner, S. G., Raina, J.-B., Nitschke, M. R., Nielsen, D. A., Stat, M., Motti, C. A., Ralph, P. J.
and Petrou, K. (2017). A multi-trait systems approach reveals a response cascade to bleaching
in corals. BMC Biol. 15, 117. doi:10.1186/s12915-017-0459-2

Gates, R. D., Baghdasarian, G. and Muscatine, L. (1992). Temperature stress causes host cell
detachment in symbiotic cnidarians: implications for coral bleaching. Biol. Bull. 182, 324-332.
doi:10.2307/1542252

Gattuso, J.-P., Magnan, A., Billeˊ, R., Cheung, W. W. L., Howes, E. L., Joos, F., Allemand, D.,
Bopp, L., Cooley, S. R., Eakin, C. M. et al. (2015). Contrasting futures for ocean and society
from different anthropogenic CO2 emissions scenarios. Science 349, 6243.
doi:10.1126/science.aac4722

Gibbin, E. M., Putnam, H. M., Davy, S. K. and Gates, R. D. (2014). Intracellular pH and its
response to CO2-driven seawater acidification in symbiotic versus nonsymbiotic coral cells. J.
Exp. Biol. 217, 1963-1969. doi:10.1242/jeb.099549

Gibbin, E. M., Putnam, H. M., Gates, R. D., Nitschke, M. R. and Davy, S. K. (2015). Species-
specific differences in thermal tolerance may define susceptibility to intracellular acidosis in reef
corals. Mar. Biol. 162, 717-723. doi:10.1007/s00227-015-2617-9

Grottoli, A. G., Warner, M. E., Levas, S. J., Aschaffenburg, M. D., Schoepf, V., McGinley, M.,
Baumann, J. and Matsui, Y. (2014). The cumulative impact of annual coral bleaching can turn
some coral species winners into losers. Glob. Change Biol. 20, 3823-3833.
doi:10.1111/gcb.12658

Guadayol, O., Silbiger, N. J., Donahue, M. J. and Thomas, F. I. (2014). Patterns in temporal
variability of temperature, oxygen and pH along an environmental gradient in a coral reef. PLoS
ONE 9, e85213. doi:10.1371/journal.pone.0085213

Guest, J. R., Low, J., Tun, K., Wilson, B., Ng, C., Raingeard, D., Ulstrup, K. E., Tanzil, J. T. I.,
Todd, P. A., Toh, T. C. et al. (2016). Coral community response to bleaching on a highly
disturbed reef. Sci. Rep. 6, 20717. doi:10.1038/srep20717

Hackerott, S., Martell, H. A. and Eirin-Lopez, J. M. (2021). Coral environmental memory:
causes, mechanisms, and consequences for future reefs. Trends Ecol. Evol. 36, 1011-1023.
doi:10.1016/j.tree.2021.06.014

Heberling, J. M. and Fridley, J. D. (2013). Resource-use strategies of native and invasive plants
in eastern North American forests. New Phytol. 200, 523-533. doi:10.1111/nph.12388

Heron, S. F., Maynard, J. A., van Hooidonk, R. and Mark Eakin, C. (2016). Warming trends and
bleaching stress of the World's coral reefs 1985–2012. Sci. Rep. 6, 1-14. doi:10.1038/s41598-
016-0001-8

Hoadley, K. D., Lewis, A. M., Wham, D. C., Tye Pettay, D., Grasso, C., Smith, R., Kemp, D. W.,
LaJeunesse, T. C. and Warner, M. E. (2019). Host–symbiont combinations dictate the photo-
physiological response of reef-building corals to thermal stress. Sci. Rep. 9, 9985.
doi:10.1038/s41598-019-46412-4

Hume, B. C. C., Smith, E. G., Ziegler, M., Warrington, H. J. M., Burt, J. A., LaJeunesse, T. C.,
Wiedenmann, J. and Voolstra, C. R. (2019). SymPortal: A novel analytical framework and
platform for coral algal symbiont next-generation sequencing ITS2 profiling. Mol. Ecol. Res. 19,
1063-1080. doi:10.1111/1755-0998.13004

IPCC (2018). Summary for policymakers. In: Global Warming of 1.5°C. An IPCC Special Report
on the Impacts of Global Warming of 1.5°C above Pre-Industrial Levels and Related Global
Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to
the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty (ed.
Masson-Delmotte, T. W. V., Zhai, P., Pö rtner, H. O., Roberts, D., Skea, J., Shukla, P. R.,
Pirani, A. et al.), pp. 3-24. Cambridge University Press. doi:10.1017/9781009157940.001

Jeffrey, S. W. and Humphrey, G. F. (1975). New spectrophotometric equations for determining
chlorophylls a, b, c1 and c2 in higher plants, algae and natural phytoplankton. Biochem. Physiol.
Pflanzen 167, 191-194. doi:10.1016/s0015-3796(17)30778-3

Jokiel, P. L., Maragos, J. E. and Franzisket, L. (1978). Coral growth: buoyant weight technique.
In Coral Reefs: Research Methods (ed. D. R. Stoddart and R. E. Johannes), pp. 529-541. Paris:
UNESCO.

Jurriaans, S. and Hoogenboom, M. O. (2020). Seasonal acclimation of thermal performance in
two species of reef-building corals. Mar. Ecol. Prog. Ser. 635, 55-70. doi:10.3354/meps13203
973
974   Jury, C. P., Thomas, F. I. M., Atkinson, M. J. and Toonen, R. J. (2013). Buffer capacity,
975   ecosystem feedbacks, and seawater chemistry under global change. WATER 5, 1303-1325.
976   doi:10.3390/w5031303
977
978   Klein, S. G., Geraldi, N. R., Anton, A., Schmidt-Roach, S., Ziegler, M., Cziesielski, M. J., Martin,
979   C., Ra¨decker, N., Fro¨ licher, T. L., Mumby, P. J. et al. (2022). Projecting coral responses to
980   intensifying marine heatwaves under ocean acidification. Glob. Change Biol. 28, 1753-1765.
981   doi:10.1111/gcb.15818
982
983   Kugler, K. C., Dziak, J. J. and Trail, J. (2018). Coding and interpretation of effects in analysis of
984   data from a factorial experiment. In Optimization of Behavioral, Biobehavioral, and Biomedical
985   Interventions. Statistics for Social and Behavioral Sciences (ed. L. Collins and K. Kugler), pp.
986   175-205. Cham: Springer. doi:10.1007/978-3-319-91776-4_6
987
988   Lajeunesse, T. C. and Trench, R. K. (2000). Biogeography of two species of Symbiodinium
989   (Freudenthal) Inhabiting the intertidal sea anemone Anthopleura elegantissima (Brandt). Biol.
990   Bull. 199, 126-134. doi:10.2307/1542872
991
992   LaJeunesse, T. C., Parkinson, J. E., Gabrielson, P. W., Jeong, H. J., Reimer, J. D., Voolstra, C.
993   R. and Santos, S. R. (2018). Systematic revision of Symbiodiniaceae highlights the antiquity
994   and diversity of coral endosymbionts. Curr. Biol. 28, 2570-2580. doi:10.1016/j.cub.2018.07.008
995
996   Le Nohaïc, M., Ross, C. L., Cornwall, C. E., Comeau, S., Lowe, R., McCulloch, M. T. and
997   Schoepf, V. (2017). Marine heatwave causes unprecedented regional mass bleaching of
998   thermally resistant corals in Northwestern Australia. Sci. Rep. 7, 14999. doi:10.1038/s41598-
999   017-14794-y
1000
1001   Lesser, M. P. (2004). Experimental biology of coral reef ecosystems. J. Exp. Mar. Biol. Ecol.
1002   300, 217-252. doi:10.1016/j.jembe.2003.12.027
1003
1004   Lesser, M. P. (2011). Coral bleaching: causes and mechanisms. In Coral Reefs: An Ecosystem
1005   in Transition (ed. Z. Dubinsky and N. Stambler). Dordrecht: Springer.
1006   doi:10.1007/978-94-007-0114-4_23
1007
1008   Loya, Y., Sakai, K., Yamazato, K., Nakano, Y., Sambali, H. and van Woesik, R. (2001). Coral
1009   bleaching: the winners and the losers. Ecol. Lett. 4, 122-131.
1010   doi:10.1046/j.14610248.2001.00203.x
1011
1012   Lueker, T. J., Dickson, A. G. and Keeling, C. D. (2000). Ocean pCO_2 calculated from dissolved
1013   inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory
1014   measurements of CO2 in gas and seawater at equilibrium. Mar. Chem. 70, 105-119.
1015   doi:10.1016/S0304-4203(00)00022-0
1016
1017   Magnusson, S. H., Fine, M. and Ku¨ hl, M. (2007). Light microclimate of endolithic phototrophs
1018   in the scleractinian corals Montipora monasteriata and Porites cylindrica. Mar. Ecol. Prog. Ser.
1019   332, 119-128. doi:10.3354/meps332119
1020

Marshall, B. and Biscoe, P. V. (1980). A model for C3 leaves describing the dependence of net
photosynthesis on irradiance. J. Exp. Bot. 31, 29-39. doi:10.1093/jxb/31.1.29

Matsuda, S., Huffmyer, A., Lenz, E. A., Davidson, J., Hancock, J., Przybylowski, A., Innis, T.,
Gates, R. D. and Barott, K. L. (2020). Coral bleaching susceptibility is predictive of subsequent
mortality within but not between coral species. Front. Ecol. Evol. 8, 178.
doi:10.3389/fevo.2020.00178

Matthews, J. L., Crowder, C. M., Oakley, C. A., Lutz, A., Roessner, U., Meyer, E., Grossman, A.
R., Weis, V. M. and Davy, S. K. (2017). Optimal nutrient exchange and immune responses
operate in partner specificity in the cnidarian–dinoflagellate symbiosis. Proc. Natl. Acad. Sci.
U.S.A. 114, E11058.

Matthews, J. L., Oakley, C. A., Lutz, A., Hillyer, K. E., Roessner, U., Grossman, A. R., Weis, V.
M. and Davy, S. K. (2018). Partner switching and metabolic flux in a model cnidarian–
dinoflagellate symbiosis. Proc. R. Soc. B 285, 20182336. doi:10.1098/rspb.2018.2336

McLachlan, R. H., Price, J. T., Solomon, S. L. and Grottoli, A. G. (2020). Thirty years of coral
heat-stress experiments: a review of methods. Coral Reefs 39, 885-902. doi:10.1007/s00338-
020-01931-9

McLachlan, R. H., Price, J. T., Mun˜oz-Garcia, A., Weisleder, N. L., Jury, C. P., Toonen, R. J.
and Grottoli, A. G. (2021). Environmental gradients drive physiological variation in Hawaiian
corals. Coral Reefs 40, 1505-1523. doi:10.1007/s00338-021-02140-8

Mollica, N. R., Guo, W., Cohen, A. L., Huang, K.-F., Foster, G. L., Donald, H. K. and Solow, A.
R. (2018). Ocean acidification affects coral growth by reducing skeletal density. Proc. Natl.
Acad. Sci. U.S.A. 115, 1754-1759. doi:10.1073/pnas.1712806115

Mydlarz, L. D., McGinty, E. S. and Drew Harvell, C. (2010). What are the physiological and
immunological responses of coral to climate warming and disease? J. Exp. Biol. 213, 934-945.
doi:10.1242/jeb.037580

Neal, B. P., Khen, A., Treibitz, T., Beijbom, O., O'Connor, G., Coffroth, M. A., Knowlton, N.,
Kriegman, D., Greg Mitchell, B. and Kline, D. I. (2017). Caribbean massive corals not recovering
from repeated thermal stress events during 2005–2013. Ecol. Evol. 7, 1339-1353.

Oakley, C. A. and Davy, S. K. (2018). Cell biology of coral bleaching. In Coral Bleaching.
Ecological Studies, vol. 233 (ed. M. van Oppen and J. Lough). Cham: Springer.
doi:10.1007/978-3-319-75393-5_8

Olito, C., White, C. R., Marshall, D. J. and Barneche, D. R. (2017). Estimating monotonic rates
from biological data using local linear regression. J. Exp. Biol. 220, 759-764.
doi:10.1242/jeb.148775

Oliver, T. A. and Palumbi, S. R. (2011). Many corals host thermally resistant symbionts in high-
temperature habitat. Coral Reefs 30, 241-250. doi:10.1007/s00338-010-0696-0

Oliver, E. C. J., Donat, M. G., Burrows, M. T., Moore, P. J., Smale, D. A., Alexander, L. V.,
Benthuysen, J. A., Feng, M., Sen Gupta, A., Hobday, A. J. et al. (2018). Longer and more

frequent marine heatwaves over the past century. Nat. Commun. 9, 1324. doi:10.1038/s41467-
018-03732-9

Oliver, E. C. J., Benthuysen, J. A., Darmaraki, S., Donat, M. G., Hobday, A. J., Holbrook, N. J.,
Schlegel, R. W. and Gupta, A. S. (2021). Marine heatwaves. Ann. Rev. Mar. Sci. 13, 313-342.
doi:10.1146/annurev-marine-032720-095144

Osborne, K., Thompson, A. A., Cheal, A. J., Emslie, M. J., Johns, K. A., Jonker, M. J., Logan,
M., Miller, I. R. and Sweatman, H. P. A. (2017). Delayed coral recovery in a warming ocean.
Glob. Change Biol. 23, 3869-3881. doi:10.1111/gcb.13707

Pendleton, L. H., Hoegh-Guldberg, O., Langdon, C. and Comte, A. (2016). Multiple stressors
and ecological complexity require a new approach to coral reef research. Front. Mar. Sci. 3, 36.
doi:10.3389/fmars.2016.00036

Perez, F. F. and Fraga, F. (1987). Association constant of fluoride and hydrogen ions in
seawater. Mar. Chem. 21, 161-168. doi:10.1016/0304-4203(87)90036-3

Putnam, H. M. (2021). Avenues of reef-building coral acclimatization in response to rapid
environmental change. J. Exp. Biol. 224, jeb239319. doi:10.1242/jeb.239319

Putnam, H. M. and Edmunds, P. J. (2011). The physiological response of reef corals to diel
fluctuations in seawater temperature. J. Exp. Mar. Biol. Ecol. 396, 216-223.
doi:10.1016/j.jembe.2010.10.026

Putnam, H. M., Mayfield, A. B., Fan, T. Y., Chen, C. S. and Gates, R. D. (2013). The
physiological and molecular responses of larvae from the reef-building coral Pocillopora
damicornis exposed to near-future increases in temperature and pCO_2. Mar. Biol. 160, 2157-
2173. doi:10.1007/s00227-012-2129-9

Putnam, H. M., Barott, K. L., Ainsworth, T. D. and Gates, R. D. (2017). The vulnerability and
resilience of reef-building corals. Curr. Biol. 27, R528-R540. doi:10.1016/j.cub.2017.04.047

Roach, T. N., Dilworth, J., Jones, A. D., Quinn, R. A. and Drury, C. (2021). Metabolomic
signatures of coral bleaching history. Nature Ecol. Evol. 54, 495-503.doi:10.1038/s41559-020-
01388-7

Rodrigues, L. J. and Grottoli, A. G. (2007). Energy reserves and metabolism as indicators of
coral recovery from bleaching. Limnol. Oceanogr. 52, 1874-1882.
doi:10.4319/lo.2007.52.5.1874

Rodrıdguez-Troncoso, A. P., Carpizo-Ituarte, E. and Cupul-Magan~a, A. L. (2016). Physiological
response to high temperature in the tropical Eastern Pacific coral Pocillopora verrucosa. Mar.
Ecol. 10, 1212717. doi:10.3389/fmars.2023.1212717

Rohwer, F., Seguritan, V., Azam, F. and Knowlton, N. (2002). Diversity and distribution of coral-
associated bacteria. Mar. Ecol. Prog. Ser. 243, 1-10. doi:10.3354/meps243001

Safaie, A., Silbiger, N. J., McClanahan, T. R., Pawlak, G., Barshis, D. J., Hench, J. L., Rogers,
J. S., Williams, G. J. and Davis, K. A. (2018). High frequency temperature variability reduces the
risk of coral bleaching. Nat. Commun. 9, 1671. doi:10.1038/s41467-018-04074-2

1122

1123 Scheufen, T., Kra¨mer, W. E., Iglesias-Prieto, R. and Enrı´quez, S. (2017). Seasonal variation
1124 modulates coral sensibility to heat-stress and explains annual changes in coral productivity. Sci.
1125 Rep. 7, 4937. doi:10.1038/s41598-017-04927-8
1126

1127 Schoepf, V., Grottoli, A. G., Levas, S. J., Aschaffenburg, M. D., Baumann, J. H., Matsui, Y. and
1128 Warner, M. E. (2015). Annual coral bleaching and the long-term recovery capacity of coral.
1129 Proc. R. Soc. B 282, 20151887. doi:10.1098/rspb.2015.1887
1130

1131 Schoepf, V., Sanderson, H. and Larcombe, E. (2022). Coral heat tolerance under variable
1132 temperatures: effects of different variability regimes and past environmental history vs. current
1133 exposure. Limnol.Oceanogr. 67, 404-418. doi:10.1002/lno.12000
1134

1135 Shaw, E. C., Hamylton, S. M. and Phinn, S. R. (2016). Incorporating benthic community
1136 changes into hydrochemical-based projections of coral reef calcium carbonate production under
1137 ocean acidification. Coral Reefs 35, 739-750. doi:10.1007/s00338-016-1407-2
1138

1139 Silverstein, R. N., Cunning, R. and Baker, A. C. (2015). Change in algal symbiont communities
1140 after bleaching, not prior heat exposure, increases heat tolerance of reef corals. Glob. Change
1141 Biol. 21, 236-249. doi:10.1111/gcb.12706
1142

1143 Stat, M., Morris, E. and Gates, R. D. (2008). Functional diversity in coral–dinoflagellate
1144 symbiosis. Proc. Natl Acad. Sci. USA 105, 9256-9261. doi:10.1073/pnas.0801328105
1145

1146 Sweet, M. J. and Brown, B. E. (2016). Coral responses to anthropogenic stress in the twenty-
1147 first century: an ecophysiological perspective. Oceanogr. Mar. Biol. Ann. Rev. 54, 271-314.
1148

1149 Therneau, T. M., and Grambsch, P. M. (2013). Modeling Survival Data: Extending the Cox
1150 Model. Springer Science & Business Media.
1151

1152 Thomas, L. and Palumbi, S. R. (2017). The genomics of recovery from coral bleaching. Proc.
1153 Biol. Sci. R. Soc. 284, 20171790. doi:10.1098/rspb.2017.1790
1154

1155 Thornhill, D. J., Rotjan, R. D., Todd, B. D., Chilcoat, G. C., Iglesias-Prieto, R., Kemp, D.W.,
1156 LaJeunesse, T. C., Reynolds, J. M. C., Schmidt, G.W., Shannon, T. et al. (2011). A connection
1157 between colony biomass and death in caribbean reef-building corals. PloS One 6, e29535.
1158 doi:10.1371/journal.pone.0029535
1159

Tresguerres, M., K. L. Barott, M. E. Barron, D. D. Deheyn, D. I. Kline, and Linsmayer, L. B.
(2017). Cell biology of reef-building corals: ion transport, acid/ base regulation, and energy
metabolism. In Acid-Base Balance and Nitrogen Excretion in Invertebrates: Mechanisms and
Strategies in Various Invertebrate Groups with Considerations of Challenges Caused by Ocean
Acidification (ed. Dirk Weihrauch and Michael O'Donnell), 193-218. Cham: Springer
International Publishing.
1166

1167 van Oppen, M. J. H. and Lough, J. M. (2018). Coral Bleaching: Patterns,
1168 Processes, Causes and Consequences. Springer. Veal, C. J., Carmi, M., Fine, M. and Hoegh-
1169 Guldberg, O. (2010). Increasing the accuracy of surface area estimation using single wax
1170 dipping of coral fragments. Coral Reefs 29, 893-897. doi:10.1007/s00338-010-0647-9
1171

1172   Wall, C. B., Ritson-Williams, R., Popp, B. N. and Gates, R. D. (2019). Spatial variation in the
1173   biochemical and isotopic composition of corals during bleaching and recovery. Limnol.
1174   Oceanogr. 64, 2011-2028. doi:10.1002/lno.11166
1175

1176   Wall, C. B., Kaluhiokalani, M., Popp, B. N., Donahue, M. J. and Gates, R. D. (2020). Divergent
1177   symbiont communities determine the physiology and nutrition of a reef coral across a light-
1178   availability gradient. ISME J. 14, 945-958. doi:10.1038/s41396-019-0570-1
1179

1180   Wall, C. B., Ricci, C. A., Wen, A. D., Ledbetter, B. E., Klinger, D. E., Mydlarz, L. D., Gates, R. D.
1181   and Putnam, H. M. (2021). Shifting baselines: physiological legacies contribute to the response
1182   of reef corals to frequent heatwaves. Funct. Ecol. 35, 1366-1378. doi:10.1111/1365-2435.13795
1183

1184   Weis, V. M. (2008). Cellular mechanisms of cnidarian bleaching: stress causes the collapse of
1185   symbiosis. J. Exp. Biol. 211, 3059-3066. doi:10.1242/jeb.009597
1186

1187   van Oppen, M. J., Oliver, J. K., Putnam, H. M. and Gates, R. D. (2015). Building coral reef
1188   resilience through assisted evolution. Proc. Natl. Acad. Sci. USA 112, 2307-2313.
1189   doi:10.1073/pnas.1422301112
1190

1191   van Woesik, R. and Kratochwill, C. (2022). A global coral-bleaching database, 19802020. Sci.
1192   Data 9, 20. doi:10.1038/s41597-022-01121-y
1193

1194   vanWoesik, R., Sakai, K., Ganase, A. and Loya, Y. (2011). Revisiting the winners and the losers
1195   a decade after coral bleaching. Mar. Ecol. Prog. Ser. 434, 67-76. doi:10.3354/meps09203
1196

1197   van Woesik, R., Shlesinger, T., Grottoli, A. G., Toonen, R. J., Thurber, R. V., Warner, M. E.,
1198   Hulver, A. M., Chapron, L., McLachlan, R. H., Albright, R. et al. (2022). Coral-bleaching
1199   responses to climate change across biological scales. Glob. Change Biol. 28, 4229-4250.
1200   doi:10.1111/gcb.16192
1201

1202   Ziegler, M., Anton, A., Klein, S. G., Ra¨decker, N., Geraldi, N. R., Schmidt-Roach, S., Saderne,
1203   V., Mumby, P. J., Cziesielski, M. J., Martin, C. et al. (2021). Integrating environmental variability
1204   to broaden the research on coral responses to future ocean conditions. Glob. Change Biol. 27,
1205   5532-5546. doi:10.1111/gcb.15840
1206

1207   Ziegler, M., Roder, C. M., Bu¨ chel, C. and Voolstra, C. R. (2015). Mesophotic coral depth
1208   acclimatization is a function of host-specific symbiont physiology. Front. Mar. Sci. 2, 4.
1209   doi:10.3389/fmars.2015.00004
1210

1211

1212

Fig. 1. Experimental design for an environmentally and seasonally realistic extended
heatwave scenario. (A) A map of O'ahu, Hawai'i, with a highlight box around the study
site. (B) A map of Kāne'ohe Bay with six circled collection sites. (C) Twelve tanks (n=3
per treatment) were assigned a treatment with the following color and acronym code:
ATAC (dark blue), ATHC (light blue), HTAC (light red), HTHC (dark red). (D) Sampling
timeline with an 'x' to indicate when sampling occurred for each response category
(n=6 physiology; n=3 molecular per species per treatment). Physiology includes
antioxidant capacity, soluble protein, ash-free dry weight (AFDW), chlorophyll a (*chl a*)
content and endosymbiont density. Metabolism includes photosynthetic and respiration
rates. (E) Discrete measurements for pH treatment conditions (pH total scale) for the
full experimental period. (F) Average daily pH treatment conditions modeled after
natural fluctuations that occur in Kāne'ohe Bay. Data come from continuous
measurements from an Apex System set up during the first 2 weeks of the exposure
period. (G) Discrete measurements for temperature treatment conditions (°C) for the
full experimental period. A timeline of physiological sampling dates is indicated above
the x-ais. (H) Average daily temperature treatment conditions modeled after natural
fluctuations that occur in Kāne'ohe Bay.

1230

1231
1232

Fig. 2. *Montipora capitata* and *Pocillopora acuta* baseline physiological measurements measured at week 1. (A) AFDW symbiont:host (S:H) ratio, (B) photosynthesis:respiration (P:R) ratio, (C) host total antioxidant capacity (CRE µmol mg−1), (D) symbiont cellular AFDW (×10−5 mg cell−1) and (E) cellular net photosynthetic rates (Pnet; ×10−5 µmol O2 cell−1 h−1).

1237

Fig. 3. Holobiont chlorophyll a content (μg cm−2) and endosymbiont cell density (×105 cells cm−2) levels during the exposure period (weeks 1–8). The x-axis of E–H is time in weeks with sampling at 1, 2, 4, 6 and 8 weeks. Color indicates the four treatments. Points represent means±s.e.m., and significant effects from Type III ANOVA of linear mixed models for each response are labeled with corresponding P-value. (A–D) Examples of (A,C) healthy and (B,D) bleached *M. capitata* and *P. acuta* fragments. (E,F) Cell density levels (×105 cells cm−2) of (E) *M. capitata* and (F) *P. acuta*. (G) *Chl a* content (μg cm−2) of (G) *M. capitata* and (H) *P. acuta*.

1246

1247

1248

Fig. 4. Principal component analyses of multivariate physiological state during the
exposure period (weeks 1–8) for both species. (A,B) *Pocillopora acuta* fragments.
(C,D) *Montipora capitata* fragments. Each point represents an individual coral
fragment's multivariate physiological state composed of 13 host and symbiont
response variables and colored by treatment. Significant effects from PERMANOVA
tests for each species are included within A and C (bottom right of each panel). PC
axis labels indicate the percent variance explained by that axis. (A,C) Centroid
multivariate state for each treatment at each time period is indicated by a bold,
treatment-colored point. All mean points are connected within each treatment to show
the shifts through time (week 1→8). (B,D) Biplots of the response variables are shown
on the multivariate plots. Each black arrow represents a physiological response
variable. The direction and length of each arrow indicates the relative strength of that
variable to the PC axes.

Fig. 5. Bleaching assessment and principal components analysis for multivariate physiological state of *M. capitata* fragments during the recovery period (weeks 8–16). (A) Endosymbiont densities (×105 cells cm−2) and (B) Holobiont *chl a* content (μg cm−2) for the recovery period. (C,D) Each point represents an individual fragment's multivariate physiological state composed of nine host and symbiont response variables (respiration and photosynthetic rates were not measured during the recovery period). (C) Centroid multivariate physiological state through time (week 8→16). (D) Biplot of individual physiological variable contributions to PCA separation. Each point (individual fragment) is colored by treatment. Significant effects from PERMANOVA tests for each species are indicated in gray text.

1276

1277

Fig. 6. Relative abundance of Symbiodiniaceae 'type profiles' identified by ITS2
sequencing for both species during exposure and recovery periods. The y-axis
includes each type profile, composed of a major sequence ID (in bold) and subsequent
full type profile ID. Relative abundance is indicated by 0–100% continuous color scale,
with the darkest green coloring indicating 100% abundance of that full type profile. The
exposure period includes 0, 6 and 12 h, and 1, 2, 4, 6 and 8 weeks. The recovery
period includes 12 and 16 weeks, but there was not a high enough sample size for *P.*
acuta fragments at 16 weeks.

1286

1287
Fig. S1. NOAA Moku o Loʻe Buoy data (CRIMP2) from September 2012 - June
2019 with treatment conditions overlayed. CRIMP2 buoy data (sea surface
temperature °C) was used to determine an environmentally realistic range of
temperatures for the experimental exposure period. The red dotted line represents the
average high temperature treatment condition. CRIMP2 information can be found at
https://www.ndbc.noaa.gov/station_page.php?station=mokh1.
1294

1295

1296

Fig. S2. Carbonate chemistry parameters during the acclimation, exposure, and
beginning of the recovery period for all treatments. Treatment is indicated by color
and vertical dotted lines mark the start and end of the treatment period. All x-axes
represent time by month. Parameters include the following: (A) Aragonite Saturation
State (Ω). (B) Carbon dioxide (CO_2; μmol kg^{-1}). (C) Carbonate (CO^{2-}_3; μmol kg^{-1}). (D)
Dissolved inorganic carbon (DIC; μmol kg^{-1}). (E) Bicarbonate (HCO_{3-}; μmol kg^{-1}). (F)
Partial pressure of CO_2 (pCO_2; ppm). (G) pH (total scale). (H) Total alkalinity (TA; mg
L^{-1} CaCO$_3$) (I). Temperature (°C).

1305

1306

1307
1308 **Fig. S3. Light data from the exposure period for all treatments.** Light reported in
1309 PAR (nm) measured with a LICOR-193 as described in methods. Color indicates
1310 treatment (dark blue = ATAC, light blue = ATHC, light red = HTAC, and dark red =
1311 HTHC).
1312

1313

1314

1315
Fig. S4. Photosynthetic Irradiance (PI) Curves for four time points during the
exposure period for both species. Photosynthetic rates (μmol O_2 cm^{-2} h^{-1}) measured
at a range of irradiance (μmol photons m^{-2} s^{-1}) levels plotted for four time points:
September 14 2018, October 17 2018, October 31 2018, and November 15 2018. PI
curves were conducted to identify saturating irradiance (I_k; solid vertical line in each
plot) and ensure coral fragments were not reaching photoinhibition during respiration
and photosynthesis measurements.
1323

1324

1325

Fig. S5. Survival probability calculated from a cox-mixed effects model for each
species. Treatment is indicated by color, significant effects are in black text, and non-
significant effects are in gray text. The dotted line is the point when all treatment
returns to ambient level conditions. Survival probability is on a scale of 0-1.00 for both
(A) *Montipora capitata* fragments. (B) *Pocillopora acuta* fragments across both
exposure (day 1 - week 8) and recovery (week 8-16) periods.

1332

1333

1334
Fig. S6. Mean ± s.e.m. weekly color score measurements through both exposure
(week 1-8) and recovery (week 9-16) periods for each species. As coral bleaches,
color score decreases into negative values. (A) *M. capitata* fragments. (B) *P. acuta*
fragments. Significant effects from the Type III ANOVA of linear mixed model outputs
are listed on each panel. Treatments are indicated by colored mean points for each
time point. The dotted line is when treatment conditions return to fully ambient values
(week 9).
1342

1343

1344
Fig. S7. Mean endosymbiont cell density levels (cells cm^{-2}) through both
exposure and recovery periods for (A) *M. capitata* and (B) *P. acuta*. Treatments
are indicated by color and significant effects for each species are labeled on the
corresponding panel. The dotted line marks when all treatments returned to ambient
level conditions at week 9. In *P. acuta*, high temperature treatments did not yield
enough sample sizes to complete physiological measurements for weeks 12 and 16.
1351

1352

Fig. S8. Mean chlorophyll-a-a content levels normalized to both surface area (µg cm^{-2})
and endosymbiont cell density (µg cell^{-1}) for (A, C) *M. capitata* and (B, D) *P. acuta*.
Treatments are indicated by color and significant effects for each species are labeled
on the corresponding panel. The dotted line marks when all treatments returned to
ambient level conditions at week 9. In all *P. actua* panels, high temperature treatments
did not yield enough sample sizes to complete physiological measurements for weeks
12 and 16.

1361

1362

Fig. S9. Mean host soluble protein levels normalized to surface area (mg cm⁻²) for
both (A) _M. capitata_ and (B) _P. acuta_ in exposure and recovery periods.
Treatments are indicated by color and significant effects for each species are labeled
on the corresponding panel. The dotted line marks when all treatments returned to
ambient level conditions at week 9. In _P. acuta_, high temperature treatments did not
yield enough sample sizes to complete physiological measurements for weeks 12 and
16.

1370

1371

1372

Fig. S10. Mean host total antioxidant capacity (TAC) levels normalized to host
soluble protein (CRE μmol mg⁻¹) for both (A) *M. capitata* **and (B)** *P. acuta* **through**
exposure and recovery periods. Treatments are indicated by color and significant
effects for each species are labeled on the corresponding panel. The dotted line marks
when all treatments returned to ambient level conditions at week 9. In *P. acuta*, high
temperature treatments did not yield enough sample sizes to complete physiological
measurements for weeks 12 and 16.

1380

1381

1382

Fig. S11. Mean tissue biomass levels measured in Ash-Free Dry Weight
normalized to surface area (AFDW mg cm⁻²) for host, symbiont, and the ratio of
symbiont:host for both (A, C, E) *M. capitata* and (B, D, F) *P. acuta*. Treatments are
indicated by color and significant effects are labeled on the corresponding panel. The
dotted line represents when treatments returned to ambient level conditions. (A,B)
Host AFDW (mg cm⁻²). (C,D) Symbiont AFDW (mg cm⁻²). (E,F) Symbiont:Host AFDW
Ratio. For all *P. acuta* panels, high temperature treatments did not yield enough
sample sizes to complete physiological measurements for weeks 12 and 16.

1391

1392

1393

Fig. S12. Mean symbiont cellular Ash-Free Dry Weight levels normalized to cell
densities (AFDW; mg cell⁻¹) for (A) *M. capitata* **and (B)** *P. acuta*. Treatments are
indicated by color and significant effects are labeled on the corresponding panel. The
dotted line represents when treatments returned to ambient level conditions. In *P.*
acuta, high temperature treatments did not yield enough sample sizes to complete
physiological measurements for weeks 12 and 16.

1400

1401

1402

Fig. S13. Mean dark respiration and net photosynthetic rates during the
exposure period for both (A, C, E) _M. capitata_ and (B, D, F) _P. acuta_. Treatments
are indicated by color and significant effects are labeled on the corresponding panel.
Both respiration (A,B) and photosynthetic rates (C,D) are reported in units of change in
oxygen levels (µmol cm^{-2} hr^{-1}). Respiration rates are the absolute value of the rate of
change so that a decrease in rates is equal to lower respiration happening within the
coral holobiont. Both respiration and photosynthetic rates were not measured for the
recovery period. (E,F) Net photosynthetic rates (µmol cm^{-2} hr^{-1}) to dark respiration rates
(µmol cm^{-2} hr^{-1}) ratio reported in unitless values.

1412

1413

1414

1415 **Fig. S14. Mean cellular net photosynthetic rates measured via change in oxygen**
1416 **levels μmol cell⁻¹ hr⁻¹) and normalized to cell densities for each species (A)** *M.*
1417 *capitata*. **(B)** *P. acuta*. Treatments are indicated by color and significant effects are
1418 labeled on the corresponding panel. Both respiration and photosynthetic rates were not
1419 measured for the recovery period.

1420

1421

1422

1423 **Fig. S15. Weekly mean holobiont growth rates measured in mg CaCO$_3$ cm^{-2} d^{-1} for**
1424 **both (A)** *M. capitata* **and (B)** *Pocilopora acuta*. Treatments are indicated by color
1425 and significant effects are labeled on the corresponding panel. The dotted line
1426 indicates when treatment returned to ambient levels. *P. acuta* fragments did not
1427 survive in enough quantities for sufficient sample sizes during the recovery period.
1428

1429

1430

Fig. S16. Correlation plot between all measured physiological variables (14
total). All treatments, species, and timepoints are collapsed into one figure to assess
which variables are highly correlated and redundant to include in multivariate analyses.
Dark blue indicates highly positively correlated (value of 1) and dark red indicates
highly negatively correlated (value of -1). The size of the dot in each cell indicates the
degree of correlation so that a larger dot is a higher correlation. Gross photosynthetic
rates and net photosynthetic rates were highly positively correlated and therefore only
net photosynthetic rates were included in multivariate analyses.

1439

1440

1441

Fig. S17. Relative abundance (%) of ITS2 Symbiodiniaceae Type Profiles. Each
individual square is the average percent relative abundance per treatment per time
point per species (n=3). The four column sections indicate treatments (ATAC, ATHC,
HTAC, and HTHC) and the two row sections indicate host species (*Montipora capitata*
and *Pocillopora acuta*). The following time points are included in the exposure period:
0 hour, 6 hour, 12 hour, 30 hour, 1, 2, 4, 6, and 8 weeks. The vertical dotted line
indicates the start of the recovery period, which includes 12 and 16 weeks. *P. acuta*
samples were only collected through week 12. Each square is colored by percentage
relative abundance, with 100% as the darkest green and 0% as white. In the ITS2
Type Profile name, the major sequence is bolded, followed by the rest of the full type
profile name.

1453

1454
1455
Fig S18. Baseline physiological measurements from week 1 ordered by most significant
differences to least. (A) Ash-free dry weight symbiont:host ratio, (B) Holobiont net photosynthetic
rates (μmol cm⁻² hr⁻¹), (C) Host total antioxidant capacity (CRE μmol mg⁻¹), (D)
Photosynthesis:Respiration ratio, (E) Cellular net photosynthetic rates (μmol cell⁻¹ hr⁻¹), (F)
Symbiont AFDW (mg cm⁻²), (G) Symbiont cellular AFDW (mg cell⁻¹), (H) Endosymbiont cell density
(cells cm⁻²), (I) Holobiont chlorophyll-a-a content (μg cm⁻²), (J) Symbiont cellular chlorophyll-a-a
content (μg cell⁻¹), (K) Host respiration rates (μmol cm⁻² hr⁻¹), (L) Host soluble protein (mg cm⁻²), (M)
Host AFDW (mg cm⁻²). *M. capitata* indicated by circles and *P. acuta* samples indicated by
triangles.
1465

Total Alkalinity: Stress Period

Treatment	N	Mean	St. Error
ATAC	48	2159.06	5.89
ATHC	48	2162.74	5.49
HTAC	48	2155.58	5.25
HTHC	48	2156.79	5.31

Timepoint * Treatment + (1 | Tank)

Effect	X^2	df	P
Timepoint	170.0467	15	**< 0.0001**
Treatment	1.1034	3	0.7762
Timepoint:Treatment	43.8277	45	0.5126

1466

1467

1468 **Table S1. Total Alkalinity sample size (N), mean, and standard error (St. Error)**
1469 **output from the exposure period for each treatment.** A linear mixed model was run
1470 to determine the effects of treatment and time point on tank conditions.
1471

aov: PAR ~ Position * Tank

Effect	df	Sum of Sq	Mean Sq	F	Pr(>F)
Position	5	23764	4753	0.155	0.978
Tank	11	498085	45280	1.475	0.143
Position:Tank	55	120265	2187	0.071	1.000

Light Summary Data: Stress Period

Treatment	N	Mean	St. Error
ATAC	212	179.70	9.97
ATHC	211	186.54	10.20
HTAC	210	176.64	10.84
HTHC	211	165.12	10.41

lmer: Treatment + (1 | Date) + (1 | Tank)

Effect	X^2	df	P
Treatment	2.0687	3	0.5583

1472

1473

1474 **Table S2. Light (PAR) output: sample size (N), mean, and standard error (St.**
1475 **Error) output from the exposure period for each treatment.** A linear mixed model
1476 was run to determine the effects of treatment on tank conditions.

1477

Stress Treatment Period

Montipora capitata

Dependent Variable	Effect	X^2	df	P
Respiration (μmol cm^{-2} h^{-1})	**Timepoint**	19.7897	4	**0.0005**
	Temperature	5.3829	1	**0.0203**
	CO2	0.2874	1	0.5919
	Timepoint:Temperature	3.5403	4	0.4718
	Timepoint:CO2	2.8987	4	0.5749
	Temperature:CO2	0.1348	1	0.7135
	Timepoint:Temperature:CO2	3.7495	4	0.4410
Net Photosynthesis (μmol cm^{-2} h^{-1})	**Timepoint**	14.4620	4	**0.0060**
	Temperature	18.1524	1	**< 0.0001**
	CO2	1.9896	1	0.1584
	Timepoint:Temperature	13.8077	4	**0.0079**
	Timepoint:CO2	9.1105	4	0.0584
	Temperature:CO2	1.5185	1	0.2178
	Timepoint:Temperature:CO2	9.0938	4	0.0588
Net Photosynthesis (μmol cell^{-1} h^{-1})	**Timepoint**	13.5702	4	**0.0088**
	Temperature	1.8312	1	0.1760
	CO2	0.3743	1	0.5407
	Timepoint:Temperature	1.6383	4	0.8019
	Timepoint:CO2	4.1280	4	0.3890
	Temperature:CO2	0.0064	1	0.9360
	Timepoint:Temperature:CO2	6.2204	4	0.1833
Gross Photosynthesis (μmol cm^{-2} h^{-1})	**Timepoint**	21.1984	4	**0.0003**
	Temperature	22.3058	1	**< 0.0001**
	CO2	0.3925	1	0.5310
	Timepoint:Temperature	15.5217	4	**0.0037**
	Timepoint:CO2	11.9872	4	**0.0174**
	Temperature:CO2	2.2441	1	0.1341
	Timepoint:Temperature:CO2	11.0932	4	**0.0255**
P:R Ratio	Timepoint	8.5378	4	0.0738
	Temperature	0.0187	1	0.8913
	CO2	3.4479	1	0.0633
	Timepoint:Temperature	2.2871	4	0.6831
	Timepoint:CO2	0.3761	4	0.9844
	Temperature:CO2	0.3484	1	0.5550
	Timepoint:Temperature:CO2	1.0685	4	0.8992

Pocillopora acuta

Dependent Variable	Effect	X^2	df	P
Respiration (μmol cm^{-2} h^{-1})	**Timepoint**	87.4169	4	**< 0.0001**
	Temperature	40.3078	1	**< 0.0001**
	CO2	0.3721	1	0.5419
	Timepoint:Temperature	9.9564	4	**0.0412**
	Timepoint:CO2	2.2023	4	0.6986
	Temperature:CO2	0.4268	1	0.5136
	Timepoint:Temperature:CO2	7.8998	4	0.0953
Net Photosynthesis (μmol cm^{-2} h^{-1})	**Timepoint**	51.8737	4	**< 0.0001**
	Temperature	61.5502	1	**< 0.0001**
	CO2	0.0146	1	0.9038
	Timepoint:Temperature	34.5005	4	**< 0.0001**
	Timepoint:CO2	2.3684	4	0.6684
	Temperature:CO2	0.7537	1	0.3853
	Timepoint:Temperature:CO2	7.6865	4	0.1038
Net Photosynthesis (μmol cell^{-1} h^{-1})	**Timepoint**	29.9940	4	**< 0.0001**
	Temperature	29.7790	1	**< 0.0001**
	CO2	2.8160	1	0.5957
	Timepoint:Temperature	18.3100	4	**0.001073**
	Timepoint:CO2	3.7644	4	0.4388
	Temperature:CO2	2.4520	1	0.1174
	Timepoint:Temperature:CO2	5.6688	4	0.2253
Gross Photosynthesis (μmol cm^{-2} h^{-1})	**Timepoint**	58.0206	4	**< 0.0001**
	Temperature	60.0036	1	**< 0.0001**
	CO2	0.3337	1	0.5635
	Timepoint:Temperature	24.9031	4	**< 0.0001**
	Timepoint:CO2	1.7910	4	0.7741
	Temperature:CO2	0.3442	1	0.5574
	Timepoint:Temperature:CO2	7.1393	4	0.1287
P:R Ratio	Timepoint	1.8573	4	0.7620
	Temperature	9.1988	1	**0.0024**
	CO2	0.8475	1	0.3573
	Timepoint:Temperature	12.5200	4	**0.0139**
	Timepoint:CO2	0.5815	4	0.9651
	Temperature:CO2	0.0370	1	0.8475
	Timepoint:Temperature:CO2	2.3578	4	0.6703

Table S3. Statistical output for metabolic (photosynthetic and respiration) rates normalized to both surface area (cm^{-2}) and cell density (cell^{-1}). Left: *M. capitata*. Right: *P. acuta*.

Stress Treatment Period

Montipora capitata

Dependent Variable	Effect	X^2	df	P
Host soluble protein mg cm⁻²	Timepoint	28.6767	4	< 0.0001
	Temperature	1.6554	1	0.1982
	CO2	1.4948	1	0.2215
	Timepoint:Temperature	9.6751	4	0.0463
	Timepoint:CO2	22.1155	4	0.0002
	Temperature:CO2	0.2089	1	0.6476
	Timepoint:Temperature:CO2	3.8747	4	0.4232
Host total antioxidant capacity (TAC) CRE μmol (mg prot)⁻¹	Timepoint	3.3971	4	0.4937
	Temperature	1.2026	1	0.2728
	CO2	0.2111	1	0.6459
	Timepoint:Temperature	5.7398	4	0.2194
	Timepoint:CO2	2.3971	4	0.6632
	Temperature:CO2	0.2418	1	0.6229
	Timepoint:Temperature:CO2	2.1796	4	0.7028
Chlorophyll-a ug cm⁻²	Timepoint	21.1924	4	0.0003
	Temperature	15.8058	1	< 0.0001
	CO2	0.7297	1	0.3930
	Timepoint:Temperature	13.5381	4	0.0089
	Timepoint:CO2	5.0723	4	0.2800
	Temperature:CO2	0.1355	1	0.7128
	Timepoint:Temperature:CO2	3.5923	4	0.4640
Chlorophyll-a ug cell⁻¹	Timepoint	7.2542	4	0.1231
	Temperature	3.0607	1	0.0802
	CO2	2.2806	1	0.1310
	Timepoint:Temperature	4.2774	4	0.3698
	Timepoint:CO2	1.9902	4	0.7376
	Temperature:CO2	0.2751	1	0.5999
	Timepoint:Temperature:CO2	4.2725	4	0.3704
Endosymbiont density cells cm⁻²	Timepoint	34.9996	4	< 0.0001
	Temperature	8.4654	1	0.0036
	CO2	0.0533	1	0.8173
	Timepoint:Temperature	13.2732	4	0.0100
	Timepoint:CO2	8.4822	4	0.0754
	Temperature:CO2	5.0336	1	0.0249
	Timepoint:Temperature:CO2	5.3255	4	0.2555

Pocillopora acuta

Dependent Variable	Effect	X^2	df	P
Host soluble protein mg cm⁻²	Timepoint	31.4707	4	< 0.0001
	Temperature	10.7168	1	0.0011
	CO2	0.1423	1	0.7060
	Timepoint:Temperature	9.5372	4	0.0490
	Timepoint:CO2	0.6262	4	0.9601
	Temperature:CO2	0.0761	1	0.7827
	Timepoint:Temperature:CO2	8.6149	4	0.0715
Host total antioxidant capacity (TAC) CRE μmol (mg prot)⁻¹	Timepoint	5.5116	4	0.2387
	Temperature	1.5631	1	0.2387
	CO2	0.0277	1	0.8679
	Timepoint:Temperature	11.2675	4	0.0237
	Timepoint:CO2	2.9275	4	0.5700
	Temperature:CO2	0.0066	1	0.9351
	Timepoint:Temperature:CO2	1.3579	4	0.8515
Chlorophyll-a ug cm⁻²	Timepoint	27.8694	4	< 0.0001
	Temperature	18.5191	1	< 0.0001
	CO2	0.6615	1	0.4160
	Timepoint:Temperature	25.1913	4	< 0.0001
	Timepoint:CO2	2.8095	4	0.5902
	Temperature:CO2	0.1432	1	0.7051
	Timepoint:Temperature:CO2	6.3085	4	0.1773
Chlorophyll-a ug cell⁻¹	Timepoint	9.3925	4	0.0520
	Temperature	5.3056	1	0.0213
	CO2	0.1696	1	0.6804
	Timepoint:Temperature	8.4497	4	0.0764
	Timepoint:CO2	1.9180	4	0.7508
	Temperature:CO2	0.0055	1	0.9408
	Timepoint:Temperature:CO2	1.8844	4	0.7570
Endosymbiont density cells cm⁻²	Timepoint	28.9709	4	< 0.0001
	Temperature	25.8864	1	< 0.0001
	CO2	0.0001	1	0.9905
	Timepoint:Temperature	36.2584	4	< 0.0001
	Timepoint:CO2	3.6063	4	0.4619
	Temperature:CO2	0.7820	1	0.3765
	Timepoint:Temperature:CO2	4.6170	4	0.3289

Recovery Period

Montipora capitata

Dependent Variable	Effect	X^2	df	P
Host soluble protein mg cm⁻²	Timepoint	1.2029	1	0.2727
	Temperature	1.2085	1	0.2716
	CO2	0.4967	1	0.4810
	Timepoint:Temperature	4.3028	1	0.0381
	Timepoint:CO2	0.1406	1	0.7077
	Temperature:CO2	0.2735	1	0.6010
	Timepoint:Temperature:CO2	0.4454	1	0.5045
Host total antioxidant capacity (TAC) CRE μmol (mg prot)⁻¹	Timepoint	0.446	1	0.5043
	Temperature	2.7182	1	0.0992
	CO2	0.4044	1	0.5248
	Timepoint:Temperature	4.8473	1	0.0277
	Timepoint:CO2	0.0354	1	0.8507
	Temperature:CO2	0.0317	1	0.8586
	Timepoint:Temperature:CO2	0.2589	1	0.6109
Chlorophyll-a ug cm⁻²	Timepoint	1.7639	1	0.1841
	Temperature	20.1042	1	< 0.0001
	CO2	0.0496	1	0.8237
	Timepoint:Temperature	0.458	1	0.4986
	Timepoint:CO2	0.2575	1	0.6118
	Temperature:CO2	0.0059	1	0.9389
	Timepoint:Temperature:CO2	2.2585	1	0.1329
Chlorophyll-a ug cell⁻¹	Timepoint	4.4612	1	0.0347
	Temperature	0.869	1	0.3512
	CO2	2.2783	1	0.1312
	Timepoint:Temperature	0.929	1	0.3351
	Timepoint:CO2	0.7022	1	0.4020
	Temperature:CO2	0.0001	1	0.9907
	Timepoint:Temperature:CO2	0.2732	1	0.6012
Endosymbiont density cells cm⁻²	Timepoint	1.5153	1	0.2183
	Temperature	7.2667	1	0.0070
	CO2	2.2871	1	0.1305
	Timepoint:Temperature	1.3407	1	0.2469
	Timepoint:CO2	0.1552	1	0.6936
	Temperature:CO2	0.1244	1	0.7244
	Timepoint:Temperature:CO2	0.9057	1	0.3413

Table S4. Statistical output for host soluble protein, host total antioxidant capacity, chlorophyll-a-a content, and endosymbiont densities during the exposure period for both *M. capitata* and *P. acuta* and the recovery period for *M. capitata.*

Stress Treatment Period

Montipora capitata

Dependent Variable	Effect	X^2	df	P
Host AFDW	**Timepoint**	**14.8616**	**4**	**0.0050**
mg cm^{-2}	Temperature	1.9343	1	0.1643
	CO2	0.6814	1	0.4091
	Timepoint:Temperature	1.0511	4	0.9020
	Timepoint:CO2	4.3846	4	0.3564
	Temperature:CO2	0.7694	1	0.3804
	Timepoint:Temperature:CO2	7.4577	4	0.1136
Symbiont AFDW	**Timepoint**	**17.8198**	**4**	**0.0013**
mg cm^{-2}	Temperature	1.2744	1	0.2589
	CO2	0.0635	1	0.8010
	Timepoint:Temperature	2.2496	4	0.6900
	Timepoint:CO2	2.105	4	0.7165
	Temperature:CO2	0.8867	1	0.3464
	Timepoint:Temperature:CO2	4.1668	4	0.3839
Symbiont AFDW	**Timepoint**	**23.2685**	**4**	**0.0001**
mg cell^{-1}	Temperature	2.064	1	0.1508
	CO2	0.6042	1	0.4370
	Timepoint:Temperature	8.7003	4	0.0690
	Timepoint:CO2	8.6447	4	0.0706
	Temperature:CO2	**6.1314**	**1**	**0.0133**
	Timepoint:Temperature:CO2	1.9128	4	0.7518
Symbiont:Host Ratio	Timepoint	9.1648	4	0.0571
	Temperature	0.3193	1	0.5720
	CO2	1.2031	1	0.2727
	Timepoint:Temperature	0.4943	4	0.9741
	Timepoint:CO2	6.4993	4	0.1648
	Temperature:CO2	0.2733	1	0.6011
	Timepoint:Temperature:CO2	3.1485	4	0.5333

Pocillopora acuta

Dependent Variable	Effect	X^2	df	P
Host AFDW	**Timepoint**	**11.8141**	**4**	**0.0188**
mg cm^{-2}	Temperature	3.3048	1	0.0691
	CO2	0.4392	1	0.5075
	Timepoint:Temperature	2.2453	4	0.6907
	Timepoint:CO2	1.8383	4	0.7655
	Temperature:CO2	1.6816	1	0.1947
	Timepoint:Temperature:CO2	2.5394	4	0.6376
Symbiont AFDW	**Timepoint**	**16.1781**	**4**	**0.0028**
mg cm^{-2}	Temperature	1.5581	1	0.2120
	CO2	1.6089	1	0.2046
	Timepoint:Temperature	6.0128	4	0.1982
	Timepoint:CO2	5.5813	4	0.2327
	Temperature:CO2	0.9101	1	0.3401
	Timepoint:Temperature:CO2	4.2046	4	0.3790
Symbiont AFDW	**Timepoint**	**12.4879**	**4**	**0.0141**
mg cell^{-1}	**Temperature**	**12.4936**	**1**	**0.0004**
	CO2	1.2175	1	0.2699
	Timepoint:Temperature	**54.2688**	**4**	**< 0.0001**
	Timepoint:CO2	**12.4029**	**4**	**0.0146**
	Temperature:CO2	0.0764	1	0.7822
	Timepoint:Temperature:CO2	1.0241	4	0.9061
Symbiont:Host Ratio	**Timepoint**	**13.5081**	**4**	**0.0090**
	Temperature	**5.5936**	**1**	**0.0180**
	CO2	0.3185	1	0.5725
	Timepoint:Temperature	4.2439	4	0.3740
	Timepoint:CO2	6.6815	4	0.1537
	Temperature:CO2	0.2966	1	0.5860
	Timepoint:Temperature:CO2	4.1574	4	0.3851

Recovery Period

Montipora capitata

Dependent Variable	Effect	X^2	df	P
Host AFDW	Timepoint	0.5857	4	0.4441
mg cm^{-2}	Temperature	0.3283	1	0.5666
	CO2	0.4115	1	0.5212
	Timepoint:Temperature	0.1518	4	0.6968
	Timepoint:CO2	0.0118	4	0.9136
	Temperature:CO2	0.0139	1	0.9061
	Timepoint:Temperature:CO2	0.9804	4	0.3221
Symbiont AFDW	Timepoint	3.0004	4	0.0832
mg cm^{-2}	Temperature	2.3528	1	0.1251
	CO2	1.0417	1	0.3074
	Timepoint:Temperature	1.8469	4	0.1741
	Timepoint:CO2	0.6433	4	0.4225
	Temperature:CO2	0.6127	1	0.4338
	Timepoint:Temperature:CO2	0.2205	4	0.6386
Symbiont AFDW	Timepoint	0.3926	4	0.5309
mg cell^{-1}	Temperature	2.8253	1	0.0928
	CO2	**4.6844**	**1**	**0.0304**
	Timepoint:Temperature	**8.2396**	**4**	**0.0041**
	Timepoint:CO2	1.0486	4	0.3058
	Temperature:CO2	0.0088	1	0.9254
	Timepoint:Temperature:CO2	0.1601	4	0.6891
Symbiont:Host Ratio	Timepoint	0.2829	4	0.5948
	Temperature	1.5012	1	0.2205
	CO2	0.0007	1	0.9794
	Timepoint:Temperature	1.7413	4	0.1870
	Timepoint:CO2	0.5863	4	0.4438
	Temperature:CO2	0.0001	1	0.9926
	Timepoint:Temperature:CO2	0.8463	4	0.3576

Table S5. Statistical output for tissue biomass measurements: host AFDW, symbiont AFDW normalized to both surface area (cm^{-2}) and cell density (cell^{-1}), and symbiont:host AFDW Ratio for both species during the exposure and the recovery period for *M. capitata*.

Buoyant Weight

Montipora capitata

Time Period	Effect	X^2	df	P
Stress	**Timepoint**	32.815	4	**< 0.0001**
	Temperature	0.2465	1	0.6195
	CO2	0.0469	1	0.8285
	Timepoint:Temperature	14.516	4	**0.0058**
	Timepoint:CO2	21.1175	4	**0.0003**
	Temperature:CO2	0.022	1	0.8820
	Timepoint:Temperature:CO2	32.3638	4	**< 0.0001**
Recovery	**Timepoint**	12.1538	4	**0.0069**
	Temperature	12.1538	1	**< 0.0001**
	CO2	1.1876	1	0.2758
	Timepoint:Temperature	28.2451	4	**< 0.0001**
	Timepoint:CO2	15.1806	4	**0.0017**
	Temperature:CO2	1.9284	1	0.1649
	Timepoint:Temperature:CO2	11.9167	4	**0.0077**

Pocillopora acuta

Time Period	Effect	X^2	df	P
Stress	**Timepoint**	47.5441	4	**< 0.0001**
	Temperature	3.2183	1	0.0728
	CO2	9.0017	1	**0.0027**
	Timepoint:Temperature	16.0271	4	**0.0030**
	Timepoint:CO2	16.2463	4	**0.0027**
	Temperature:CO2	2.3846	1	0.1225
	Timepoint:Temperature:CO2	4.5003	4	0.3425

Color Score

Montipora capitata

Time Period	Effect	X^2	df	P
Stress	Timepoint	3.3689	7	0.8489
	Temperature	0.0019	1	0.9655
	CO2	0.0001	1	0.9916
	Timepoint:Temperature	26.3114	7	**0.0004**
	Timepoint:CO2	1.8481	7	0.9678
	Temperature:CO2	0.0753	1	0.7837
	Timepoint:Temperature:CO2	26.5600	7	**0.0004**
Recovery	Timepoint	4.4963	3	0.7212
	Temperature	4.3216	1	**0.0376**
	CO2	0.0016	1	0.9680
	Timepoint:Temperature	87.9628	3	**< 0.0001**
	Timepoint:CO2	10.1081	3	0.1825
	Temperature:CO2	3.242	1	0.0718
	Timepoint:Temperature:CO2	17.8984	3	**0.0124**

Pocillopora acuta

Time Period	Effect	X^2	df	P
Stress	**Timepoint**	41.936	7	**< 0.0001**
	Temperature	0.0074	1	0.9312
	CO2	0.052	1	0.8196
	Timepoint:Temperature	57.1031	7	**< 0.0001**
	Timepoint:CO2	95.8169	7	**< 0.0001**
	Temperature:CO2	0.0231	1	0.8792
	Timepoint:Temperature:CO2	49.8236	7	**< 0.0001**
Recovery *(week 9-12)*	Timepoint	5.8895	3	0.1171
	Temperature	34.8274	1	**< 0.0001**
	CO2	3.9797	1	**0.0461**
	Timepoint:Temperature	1.2276	3	0.7464
	Timepoint:CO2	6.781	3	0.0792
	Temperature:CO2	0.0042	1	0.9485
	Timepoint:Temperature:CO2	0.509	3	0.9169

1497

1498 **Table S6.** Statistical output for calcification growth rates and color score for both species
1499 during the exposure and recovery period.
1500

Survivorship Probability

Montipora capitata

Type III Anova	Effect	df	X^2		P
	Temperature	1	21.004		**< 0.0001**
	CO2	1	11.4891		**0.0007**
	Temperature:CO2	1	0.2914		0.5893

Cox mixed-effects model fit by maximum likelihood	Effect	coef	exp(coef)	se(coef)	z	p
Surv(lifespan, status) ~ Temperature * CO2 + (1|PLUG.ID/Tank)	**High Temperature**	1.1421189	3.133401	0.2492068	4.58	**< 0. 0001**
	High CO2	0.7981529	2.221434	0.2354737	3.39	**0.0007**
	High Temp: High CO2	0.1732892	1.18921	0.3210069	0.54	0.5900
		NULL	*Integrated*		*Fitted*	
	Log-likelihood	-7359.299	-5183.304		-4472.52	

Pocillopora

Type III Anova	Effect	df	X^2		P
	Temperature	1	2.2173		0.1365
	CO2	1	40.5412		**< 0.0001**
	Temperature:CO2	1	79.9506		**< 0.0001**

Cox mixed-effects model fit by maximum likelihood	Effect	coef	exp(coef)	se(coef)	z	p
Surv(lifespan, status) ~ Temperature * CO2 + (1|PLUG.ID/Tank)	**High Temperature**	-0.9585059	0.383	0.6436949	-1.49	**0.1400**
	High CO2	-4.4442835	0.012	0.6979973	-6.37	**< 0.0001**
	High Temp: High CO2	7.8613358	2594.984	0.8791956	8.94	**< 0.0001**
		NULL	*Integrated*		*Fitted*	
	Log-likelihood	-24130.77	NaN		-18100.5	

1501

1502

1503 **Table S7.** Statistical output from survival probability models for both *M. capitata* and *P.*
1504 *acuta* throughout the entire experiment.

1505

Species Comparison (all samples at Week 1)

Dependent Variable	t	df	P
Host total antioxidant capacity (TAC) CRE μmol (mg prot)$^{-1}$	3.7566	27.413	**0.0008**
Host soluble protein mg cm^{-2}	-1.0843	37.811	0.2851
Respiration (μmol cm^{-2} h^{-1})	1.2234	44.266	0.2276
Gross Photosynthesis (μmol cm^{-2} h^{-1})	-3.4385	44.001	**0.0013**
Net Photosynthesis (μmol cm^{-2} h^{-1})	-4.0388	40.851	**0.0002**
Net Photosynthesis (μmol cell^{-1} h^{-1})	-3.0230	37.777	**0.0045**
P:R Ratio	-3.2361	45.631	**0.0023**
Endosymbiont Density cells cm^{-2}	1.8842	229.730	0.0608
Chlorophyll-a concentration ug cm^{-2}	0.4576	33.717	0.6502
Chlorophyll-a concentration ug cell^{-1}	-0.1252	37.549	0.9011
Host AFDW mg cm^{-2}	1.2561	36.851	0.2170
Symbiont AFDW mg cell^{-1}	2.5909	30.642	**0.0145**
Symbiont AFDW mg cm^{-2}	2.8515	28.989	**0.0079**
Symbiont:Host Ratio AFDW	5.0694	34.066	**< 0.0001**

Table S8. Statistical output for baseline physiological measurement comparison between *M. capitata* and *P. acuta* fragments at the initial week 1 measurement.

STRESS PERIOD

PERMANOVA: Timepoint * Temperature * CO2 * Species

Effect	df	Sum of Squares	R2	F	Pr(>F)
Timepoint	4	203.56	0.07022	5.4788	**0.001**
Temperature	1	130.16	0.04490	14.0127	**0.001**
CO2	1	13.74	0.00474	1.4789	0.166
Species	1	298.93	0.10311	32.1831	**0.001**
Timepoint:Temperature	4	115.98	0.04001	3.1218	**0.001**
Timepoint:CO2	4	69.68	0.02404	1.8755	**0.008**
Temperature:CO2	1	9.13	0.00315	0.9830	0.403
Timepoint:Species	4	145.09	0.05005	3.9051	**0.001**
Temperature:Species	1	21.56	0.00744	2.3213	**0.029**
CO2:Species	1	13.25	0.00457	1.4270	0.209
Timepoint:Temperature:CO2	4	62.88	0.02169	1.6926	**0.033**
Timepoint:Temperature:Species	4	32.22	0.01111	0.8672	0.652
Timepoint:CO2:Species	4	36.75	0.01268	0.9892	0.456
Temperature:CO2:Species	1	3.11	0.00107	0.3350	0.936
Timepoint:Temperature:CO2:Species	4	33.88	0.01169	0.9119	0.545

Montipora capitata PERMANOVA: Timepoint * Temperature * CO2

Effect	df	Sum of Squares	R2	F	Pr(>F)
Timepoint	4	184.65	0.12682	4.2291	**0.001**
Temperature	1	39.29	0.02699	3.5995	**0.005**
CO2	1	20.27	0.01393	1.8574	0.091
Timepoint:Temperature	4	60.73	0.04171	1.3909	0.099
Timepoint:CO2	4	71.81	0.04932	1.6447	**0.036**
Temperature:CO2	1	5.78	0.00397	0.5298	0.792
Timepoint:Temperature:CO2	4	58.28	0.04003	1.3349	0.135

Pocillopora acuta PERMANOVA: Timepoint * Temperature * CO2

Effect	df	Sum of Squares	R2	F	Pr(>F)
Timepoint	4	171.63	0.12002	4.3344	**0.001**
Temperature	1	139.81	0.09777	14.1231	**0.001**
CO2	1	4.67	0.00326	0.4716	0.824
Timepoint:Temperature	4	117.25	0.082	2.9611	**0.001**
Timepoint:CO2	4	42.69	0.02985	1.0781	0.377
Temperature:CO2	1	8.25	0.00577	0.8334	0.487
Timepoint:Temperature:CO2	4	44.86	0.03137	1.133	0.314

RECOVERY PERIOD

Montipora capitata PERMANOVA: Timepoint * Temperature * CO2

Effect	df	Sum of Squares	R2	F	Pr(>F)
Timepoint	4	40.95	0.0711	2.5587	**0.013**
Temperature	1	46.70	0.0811	5.8358	**0.002**
CO2	1	5.56	0.0097	0.6945	0.621
Timepoint:Temperature	4	29.98	0.0521	1.8732	0.060
Timepoint:CO2	4	14.18	0.0246	0.8861	0.507
Temperature:CO2	1	2.39	0.0042	0.2987	0.921
Timepoint:Temperature:CO2	4	12.08	0.0210	0.7550	0.661

1511

1512

1513 **Table S9.** Statistical output from a permutational multivariate ANOVA (PERMANOVA)
1514 on the multivariate physiology states for each species during the exposure and the
1515 recovery period for *M. capitata*.

1516

ITS2

PERMANOVA: Timepoint * Temperature * CO2 * Species

Effect	df	Sum of Squares	R2	F	Pr(>F)
Timepoint	10	119.18	0.04767	1.3438	**0.023**
Temperature	1	4.44	0.00178	0.5008	0.898
CO2	1	9.38	0.00375	1.0575	0.394
Species	1	329.71	0.13188	37.1762	**0.001**
Timepoint:Temperature	10	92.17	0.03687	1.0392	0.384
Timepoint:CO2	10	81.46	0.03258	0.9184	0.699
Temperature:CO2	1	13.62	0.00545	1.5361	0.121
Timepoint:Species	10	95.53	0.03821	1.1968	0.119
Temperature:Species	1	4.52	0.00181	0.5100	0.859
CO2:Species	1	9.32	0.00373	1.0505	0.407
Timepoint:Temperature:CO2	10	45.77	0.01831	0.5160	1.000
Timepoint:Temperature:Species	10	88.20	0.03528	1.1050	0.230
Timepoint:CO2:Species	10	69.92	0.02797	0.8760	0.776
Temperature:CO2:Species	1	13.35	0.00534	1.5057	0.138
Timepoint:Temperature:CO2:Species	10	42.32	0.01693	0.5302	1.000

Montipora capitata PERMANOVA: Timepoint * Temperature * CO2

Effect	df	Sum of Squares	R2	F	Pr(>F)
Timepoint	10	107.80	0.10286	1.3377	**0.030**
Temperature	1	7.88	0.00752	0.9782	0.457
CO2	1	6.37	0.00608	0.7903	0.593
Timepoint:Temperature	10	94.42	0.09010	1.1717	0.161
Timepoint:CO2	10	68.68	0.06554	0.8523	0.810
Temperature:CO2	1	10.94	0.01044	1.3573	0.215
Timepoint:Temperature:CO2	10	42.76	0.04080	0.5306	1.000

Pocillopora acuta PERMANOVA: Timepoint * Temperature * CO2

Effect	df	Sum of Squares	R2	F	Pr(>F)
Timepoint	10	44.00	0.07458	0.9719	0.501
Temperature	1	2.25	0.00381	0.4469	0.795
CO2	1	7.63	0.01292	1.5159	0.191
Timepoint:Temperature	10	55.42	0.09394	1.2242	0.177
Timepoint:CO2	10	40.50	0.06865	0.8946	0.630
Temperature:CO2	1	10.59	0.01795	2.1055	0.068
Timepoint:Temperature:CO2	10	32.21	0.05460	0.7115	0.916

Table S10. Statistical output from a permutational multivariate ANOVA (PERMANOVA) on the ITS2 full type profiles for each species for the entire time series.

Species	Metric	Sept 18 2018	Oct 17 2018	Oct 31 2018	Nov 15 2018
Montipora capitata	Am	0.95 ± 0.196	0.98 ± 0.178	0.81 ± 0.150	0.91 ± 0.328
	AQY	0.01 ± 0.001	0.01 ± 0.001	0.01 ± 0.001	0.01 ± 0.003
	Ic	40.40 ± 5.467	40.96 ± 5.092	38.02 ± 5.459	25.38 ± 2.962
	Ik	164.08 ± 14.970	163.35 ± 22.658	126.51 ± 6.308	83.19 ± 7.270
	Rd	0.22 ± 0.041	0.23 ± 0.028	0.22 ± 0.034	0.25 ± 0.066
Pocillopora acuta	Am	1.19 ± 0.257	0.89 ± 0.144	0.50 ± 0.079	0.36 ± 0.120
	AQY	0.005 ± 0.001	0.005 ± 0.001	0.004 ± 0.001	0.004 ± 0.001
	Ic	64.40 ± 8.240	58.28 ± 4.172	54.16 ± 4.208	50.12 ± 20.932
	Ik	246.29 ± 26.716	190.53 ± 9.757	121.90 ± 5.651	95.21 ± 7.303
	Rd	0.30 ± 0.069	0.26 ± 0.038	0.21 ± 0.040	0.12 ± 0.031

Table S11. Photosynthetic curve output (mean ± standard error, n=4) for maximal photosynthetic rate (Am; μmol O_2 cm^{-2} h^{-1}), apparent quantum yield (AQY, ф; mol O_2 mol photons^{-1}), Compensation irradiance (IC; PAR), saturating irradiance (I_k; PAR), and light enhanced dark respiration rate (Rd; μmol O_2 cm^{-2} h^{-1}).