ABC-HuMi: the Atlas of Biosynthetic Gene Clusters in the Human Microbiome

Pascal Hirsch [1,†]**, Azat Tagirdzhanov** [1,2,†]**, Aleksandra Kushnareva** [2,3]**, Ilia Olkhovskii** [2,3]**, Simon Graf** [4]**, Georges P. Schmartz** [1]**, Rolf Müller** [2,5,*]**, Andreas Keller** [1,2,*]**, Alexey Gurevich** [2,4,*]

[1]Chair for Clinical Bioinformatics, Saarland University, Saarbrücken 66123, Germany, [2]Helmholtz Institute for Pharmaceutical Research Saarland (HIPS), Helmholtz Centre for Infection Research, Saarbrücken 66123, Germany, [3]Saarbrücken Graduate School of Computer Science, Saarland University, Saarbrücken 66123, Germany, [4]Department of Computer Science, Saarland University, Saarbrücken 66123, Germany, [5]Department of Pharmacy, Saarland University, Saarbrücken 66123, Germany

ABSTRACT

The human microbiome has emerged as a rich source of diverse and bioactive natural products, harboring immense potential for therapeutic applications. To facilitate systematic exploration and analysis of its biosynthetic landscape, we present ABC-HuMi: the Atlas of Biosynthetic Gene Clusters (BGCs) in the Human Microbiome. ABC-HuMi integrates data from major human microbiome sequence databases and provides an expansive repository of BGCs compared to the limited coverage offered by existing resources. Employing state-of-the-art BGC prediction and analysis tools, our database ensures accurate annotation and enhanced prediction capabilities. ABC-HuMi empowers researchers with advanced browsing, filtering, and search functionality, enabling efficient exploration of the resource. At present, ABC-HuMi boasts a catalog of 14,821 representative BGCs derived from the human gut, oral, skin, respiratory and urogenital systems. By capturing the intricate biosynthetic potential across diverse human body sites, our database fosters profound insights into the molecular repertoire encoded within the human microbiome and offers a comprehensive resource for the discovery and characterization of novel bioactive compounds. The database is freely accessible at https://www.ccb.uni-saarland.de/abc_humi/.

GRAPHICAL ABSTRACT

INTRODUCTION

Bacteria, archaea, fungi, and protists residing on or within our bodies compose the human microbiota; the aggregate of their genomes is commonly referred to as the human microbiome (1). Many of these microorganisms produce bioactive compounds that play vital roles in human-microbe and microbe-microbe interactions and might greatly affect the health state of an individual (2, 3). The identification and thorough analysis of biosynthetic gene clusters (BGCs) encoding such compounds provide insights into the human microbiota capabilities and ultimately lead to the discovery of pharmaceutically relevant molecules (4). Community initiatives such as the Human Microbiome Project (5, 6) accumulated vast volumes of sequencing data that hold huge unlocked potential for the BGC exploration.

*To whom correspondence should be addressed. Emails: rolf.mueller@helmholtz-hips.de (R.M.), andreas.keller@ccb.uni-saarland.de (A.K.), alexey.gurevich@helmholtz-hips.de (A.G.) †These authors contributed equally to this work

The genome mining software paved the way for the creation of databases of BGCs computationally predicted from voluminous genomics datasets (7, 8, 9, 10). However, these resources either focus on general microbes, such as antiSMASH-DB (7) and IMG-ABC (8), or include BGCs only from a specific human body site, such as sBGC-hm (10) and BiG-FAM (9) collecting only human gut data. All databases applied the BGC mining to isolate genome sequences or high-quality metagenome-assembled genomes (MAGs), leaving the rich collections of unbinned metagenome assemblies unprocessed. Furthermore, most of these databases were created with already outdated software and do not benefit from the latest advances in genome mining techniques. Finally, only BiG-FAM enables custom BGC searches, crucial for prioritizing novel BGCs.

To address these limitations, we developed ABC-HuMi, the Atlas of BGCs in the Human Microbiome. Our resource accumulates data from three major human microbiome sequence databases and represents BGCs originating from five human body sites and systems. We employed the most advanced BGC mining and analysis software to populate the database and created a user-friendly interactive platform for exploring the collected BGCs and their associated functionalities.

DATA RETRIEVAL AND PROCESSING

Data sources

ABC-HuMi integrates data obtained from the Genomic Catalogue of Earth's Microbiomes (JGI GEM (11)), EMBL-EBI MGnify catalogs (12), and the Human Microbiome Project (HMP, (13)). The EMBL-EBI MGnify Unified Human Gastrointestinal Genome v2.0.1 and Human Oral v1.0 catalogs contain more than 290,000 isolates and MAGs from the human microbiome clustered into 4,744 and 452 species representatives, respectively. For further processing, we retrieved the representative genomes (https://www.ebi.ac.uk/metagenomics/browse/genomes). Selecting only the representative genomes as the input is a compromise between performance and sensitivity. This allowed us to keep the total number of BGCs small, but our experiments show that we still cover the vast majority of the metabolomic diversity present in the skipped genomes (75% at the gene cluster family (GCF) level and 98% at the gene cluster clan (GCC) level) (Supplementary Figure S1).

JGI GEM contains more than 52,000 MAGs from a wide range of environmental and host-associated microbiomes. We retrieved all human microbiome-associated MAGs, excluding the ones associated with the human gut, already covered by the EBI data (https://portal.nersc.gov/GEM/genomes/).

HMP contains metagenomes from more than 3,808 samples associated with different parts of the human body. From this dataset, we selected all metagenomes associated with respiratory, skin, and urogenital samples (https://portal.hmpdacc.org/).

Processing pipeline

As an initial processing step, we harmonized metadata obtained from the source databases and unified the body site categories. For EMBL-EBI MGnify catalogs, the body site was retrieved from the metadata of associated NCBI BioSamples. Metagenome assemblies were taxonomically annotated using Kraken2 (v2.1.2) (14).

The retrieved genomes were processed with antiSMASH (v7.0.0; command-line arguments (cla): --genefinding-tool=prodigal-m --asf) (15). A BGC was classified as fragmented if it is located on a contig edge and complete otherwise. All BGCs with the same body site and taxonomy were grouped and MMseqs2 (v14.7) (16) was used to compare BGCs inside each group. We then merged BGCs with a minimum sequence identity of 0.95 into a single database entry with combined metadata. Finally, BGCs were clustered with BiG-SCAPE (v1.1.5; cla: --mibig) (17) using Pfam (v35.0) (18).

Overview of the collected data

The resulting database comprises 14,821 BGCs grouped into 7,346 GCFs and 241 GCCs. Figure 1 shows the composition of the database with regard to the associated body site and product types of the BGCs.

Web server implementation

For the implementation of the ABC-HuMi web server, we set up a Django Python web framework (https://djangoproject.com/) and a PostgreSQL database (https://www.postgresql.org/) in docker containers (https://www.docker.com/) with the help of a Cookiecutter template (https://cookiecutter.readthedocs.io/). For the search job query, we are using the task queue manager Celery (http://docs.celeryproject.org) together with the in-memory data structure store Redis (https://redis.io/). The search jobs are handled with BLAST+ (19) using the Biopython wrapper (https://biopython.org/docs/dev/api/Bio.Blast.Applications.html) and cblaster (20) using a Snakemake pipeline (21). The database browsing table is using DataTables (https://datatables.net/) and the Cytoscape network visualization (22) is using cytoscape.js (https://js.cytoscape.org/). The front end of the website also uses Bootstraps (https://getbootstrap.com/) and Font Awesome (https://fontawesome.com/) for design purposes and jQuery (https://jquery.com/) as a utility library.

DATABASE FUNCTIONALITY

Versatile search options

ABC-HuMi provides the BLAST+ (19) search for identifying homologs of user-provided sequences in the database, for example, antibiotic resistance genes (Figure 2A, B). The function handles both nucleotide and protein sequences and can also search translated nucleotide sequences of the database BGCs using a protein query (the tBLASTn mode). Since the single gene search is rarely informative for identifying homologous gene clusters (23), we employ cblaster (20) for screening a custom BGC against the entire database (Supplementary Figure S2-3).

Figure 1. Overview of the ABC-HuMi content. The distributions of biosynthetic gene clusters (BGCs) by (**A**) data source, (**B**) human body site, and (**C**) product type. (**D**) The number of gene cluster clans (GCCs) that span across at least two different body sites.

Interactive tables and visualizations

All BGCs in ABC-HuMi are enriched with detailed metadata that augments antiSMASH-predicted properties with the source human microbiome sequence database and body site, the taxonomy of the producing organism, and the links to similar BGCs within our database and in MIBiG (24). The information is stored in interactive tables that allow filtering by metadata and seamless switching between BGCs and related gene cluster families (GCFs) and clans (GCCs) (Figure 2B, D). Each BGC is complemented with the fully-embedded antiSMASH report; GCF and GCC networks are visualized with Cytoscape (Figure 2C, E).

Application examples

Exploring individual BGCs. To explore the hidden bioactive potential of GCFs spanning across multiple human body sites, we employed the versatile ABC-HuMi search functionality. First, we used metadata filtration and Cytoscape visualizations to identify GCFs containing experimentally-validated MIBiG BGCs whose direct neighbors originated from distinct body sites. We further selected two out of 11 hits since the corresponding MIBiG BGCs are known for producing compounds with antimicrobial properties: bacteriocins gassericin T (BGC0000619) and gassericin E (BGC0001388). Both compounds were originally discovered in *Lactobacillus gasseri* (25), one of the main *Lactobacillus*

species identified in the vaginal, gastrointestinal, and oral microbiomes (26). The cblaster search of BGC0000619 and BGC0001388 revealed four BGCs in ABC-HuMi with identifiers HMBGC00006519-22 (Supplementary Figures S2-3). The body sites associated with these BGCs are gut, oral and urogenital system, thus perfectly matching the common localization of the known gassericin producer. Notably, three of the ABC-HuMi BGCs were predicted from *L. paragasseri*, never reported as a gassericin E producer before. Given the resemblance between these two bacterial species, such production is plausible and the ABC-HuMi BGCs might harbor a new variant of the bacteriocin. Though, this computational hypothesis requires experimental validation.

Large-scale comparison. To demonstrate the ABC-HuMi ability to prioritize the search for novel bioactive compounds, we explored BGCs in a recently published dataset of gut microbiota samples derived from patients with Parkinson's disease (27). First, we applied our data processing pipeline to the metagenomic data from (27). AntiSMASH identified 17,978 BGCs, which we further used as a query for a bulk comparison against the entire ABC-HuMi. Clustering with BiG-SCAPE revealed that 11,096 (62%) of the query BGCs were clustered together with the BGCs from ABC-HuMi on the GCF level and 16,819 (94%) on the GCC level, with 83% of matched ABC-HuMi BGCs attributed to the human gut.

Figure 2. ABC-HuMi search, browse, and visualization functionality. The BLAST search (**A**) identifies biosynthetic gene clusters (BGCs) harboring homologs of the user-provided sequence. The search result page (**B**, top) enables browsing of the BGC metadata (**B**, bottom) and the corresponding embedded antiSMASH reports (**C**). The metadata table links to the gene cluster family (GCF) or clan (GCC) of a BGC, thus allowing one to browse all similar clusters (**D**). GCC and GCF networks can also be visualized via interactive Cytoscape plots (**E**). Here all nodes of a GCC are titled with the BGC identifiers and colored according to the corresponding human body site: the selected black node is from the gut, green nodes are from the oral microbiome and red ones are from the respiratory system.

COMPARISON TO EXISTING RESOURCES

With over 2500 entries, MIBiG is by far the leading community resource for collecting experimentally-verified BGCs (24, 28). However, such BGCs are scarce. A more complete — though less accurate — view on the biosynthetic potential of environmental microbiomes relies on genome mining (4), which naturally leads to the emergence of resources storing computationally-predicted BGCs, such as our database. These resources visibly differ in data focus, processing techniques, and functionality provided to the end users. We compare the ABC-HuMi content and features to four state-of-the-art *in silico* BGC databases in Table 1.

ABC-HuMi and sBGC-hm are the only databases entirely focused on the human microbiome but the former covers five major body sites and systems while the latter contains only BGCs predicted in gut samples (10). BiG-FAM metadata allows the selection of BGCs originating from human gut metagenome-assembled genomes (MAGs) (9). Some of the remaining BiG-FAM BGCs as well as entries in antiSMASH-DB and IMG-ABC might also be relevant to the human microbiome but this is impossible to infer from the metadata (7, 8). Thus, a direct comparison of the total number of BGCs across the databases could be misleading for researchers interested in this specific environment. All databases but antiSMASH-DB contain BGCs predicted from both assemblies of isolate genomes and high-quality MAGs. On top of that, ABC-HuMi includes BGCs mined directly from unbinned metagenome assemblies.

While all five databases rely on antiSMASH for mining BGCs, the software version varies from v5 (BiG-FAM, antiSMASH-DB and IMG-ABC) to v6 (sBGC-hm) to the latest v7 (ABC-HuMi), which might greatly affect the accuracy and completeness of the predictions (15, 29, 30). ABC-HuMi, sBGC-hm, and BiG-FAM group identified BGCs into gene cluster families (GCFs) and clans (GCCs). The first two resources utilize BiG-SCAPE (17) while the enormous size of BiG-FAM requires the use of more resource-efficient but less accurate BiG-SLICE (31). The resulting BGCs could be downloaded in bulk only from the ABC-HuMi and BiG-FAM websites, but the further use of the BiG-FAM data might be complicated due to the non-standard data format.

All considered databases provide comprehensive filtering by metadata but the search for a custom sequence or BGC is available only in three of them (ABC-HuMi, BiG-FAM, and sBGC-hm) and only ABC-HuMi supports all search types. Though, antiSMASH-DB and IMG-ABC offer advanced data lookups based on a user-defined set of Pfam domains from a predefined list. The visualization facilities vary greatly among the databases but the most informative detailed view of each

Table 1. Comparison of *in silico* BGC databases. *# HM-related BGCs (GCFs)* stands for the total number of BGCs (GCFs) related to the human microbiome.

Category	ABC-HuMi	sBGC-hm	BiG-FAM	antiSMASH-DB	IMG-ABC
General information					
Data focus	Human microbiome	Human gut microbiome	Microbes in general	Microbes in general	Microbes in general
Genome types	isolates, MAGs, metagenomes	isolates, MAGs	isolates, MAGs	isolates	isolates, MAGs
Latest update	2023 (this work)	2023 (10)	2020 (9)	2020 (7)	2019 (8)
Data processing and availability					
BGC calling	antiSMASH v7 (15)	antiSMASH v6 (29)	antiSMASH v5 (30)	antiSMASH v5 (30)	antiSMASH v5 (30)
BGC clustering	BiG-SCAPE (17)	BiG-SCAPE (17)	BiG-SLICE (31)	✗	✗
# HM-related BGCs (GCFs)	14,821 (7346)	36,583 (8004)	>4791 (>811)[a]	unknown	unknown
Bulk data download	✓ (GenBank)	✗	✓ (custom)	✗	✗
Features					
Metadata search	✓	✓	✓	✓	✓
Sequence search	✓ (BLAST+ (19))	✓ (BLAST+ (19))	✗	✗	✗
Cluster search	✓ (cblaster (20))	✗	✓ (BiG-SLICE (31))	✗	✗
BGC detailed view	✓ (antiSMASH)	✗	✓ (custom[b])	✓ (antiSMASH)	✓ (antiSMASH[c])
Extra features	Cytoscape visualization of GCF/GCC networks with export to PNG and JSON	Matrices of gene co-occurrence in the HMP data Visualizations of differences in core gene coverage across HMP samples	Word cloud of Pfam features for each BGC	Cluster search based on NPRS/PKS module queries Taxonomy tree-based browsing	Pfam-based search with ClusterScout (32) Search by chemical attributes (for experimentally-verified products)

[a]The exact numbers are known for human gut MAGs only.
[b]BGC annotation into genes and Pfam domains.
[c]Unavailable for part of the database BGCs.

BGC (the antiSMASH report) is provided only in ABC-HuMi and antiSMASH-DB.

CONCLUSION

The unprecedented biosynthetic potential of the human microbiota remains underexplored. In response, we unveil ABC-HuMi, a repository of biosynthetic gene clusters (BGCs) computationally predicted in genomes of microbes populating major human body sites and systems. Despite its compact size, the database represents a huge biosynthetic diversity of the human microbiome. The streamlined nature of the resource facilitates easy and fast analysis queries, even for computationally demanding applications. ABC-HuMi empowers users with the ability to search not only for custom nucleotide and protein sequences but also entire BGCs and enhances the results with interactive browsing and visualization functionality. Unlike most existing BGC databases, the entire ABC-HuMi can be downloaded locally and used for bulk comparison and prioritization of novel BGCs in large-scale studies.

CODE AND DATA AVAILABILITY

The code used to collect and process the data can be found at https://github.com/gurevichlab/abc_humi. The resulting annotated BGC sequences in the GenBank format and all metadata are available from the database website. The resource is freely accessible at https://www.ccb.uni-saarland.de/abc_humi/.

SUPPLEMENTARY DATA

Supplementary Data are available online.

ACKNOWLEDGEMENTS

The graphical abstract was created with BioRender.com.

FUNDING

We are thankful for funding from Saarland University for the NextAID project through which the position of AT is supported.

Conflict of interest statement. None declared.

REFERENCES

1. Ursell, L. K., Metcalf, J. L., Parfrey, L. W., and Knight, R. (2012) Defining the human microbiome. *Nutrition reviews,* **70**(suppl_1), S38–S44.
2. Hou, K., Wu, Z.-X., Chen, X.-Y., Wang, J.-Q., Zhang, D., Xiao, C., Zhu, D., Koya, J. B., Wei, L., Li, J., et al. (2022) Microbiota in health and diseases. *Signal transduction and targeted therapy,* **7**(1), 135.
3. Donia, M. S., Cimermancic, P., Schulze, C. J., Brown, L. C. W., Martin, J., Mitreva, M., Clardy, J., Linington, R. G., and Fischbach, M. A. (2014) A systematic analysis of biosynthetic gene clusters in the human microbiome reveals a common family of antibiotics. *Cell,* **158**(6), 1402–1414.
4. Medema, M. H., de Rond, T., and Moore, B. S. (2021) Mining genomes to illuminate the specialized chemistry of life. *Nature Reviews Genetics,* **22**(9), 553–571.
5. Consortium, T. H. M. P. (2012) Structure, function and diversity of the healthy human microbiome. *nature,* **486**(7402), 207–214.
6. Consortium, T. H. M. P. (2012) A framework for human microbiome research. *Nature,* **486**(7402), 215–21.
7. Blin, K., Shaw, S., Kautsar, S. A., Medema, M. H., and Weber, T. (2021) The antiSMASH database version 3: increased taxonomic coverage and new query features for modular enzymes. *Nucleic acids research,* **49**(D1), D639–D643.
8. Palaniappan, K., Chen, I.-M. A., Chu, K., Ratner, A., Seshadri, R., Kyrpides, N. C., Ivanova, N. N., and Mouncey, N. J. (2020) IMG-ABC v. 5.0: an update to the IMG/Atlas of Biosynthetic Gene Clusters Knowledgebase. *Nucleic acids research,* **48**(D1), D422–D430.
9. Kautsar, S. A., Blin, K., Shaw, S., Weber, T., and Medema, M. H. (2021) BiG-FAM: the biosynthetic gene cluster families database. *Nucleic acids research,* **49**(D1), D490–D497.
10. Zou, H., Sun, T., Jin, B., and Wang, S. (2023) sBGC-hm: an atlas of secondary metabolite biosynthetic gene clusters from the human gut microbiome. *Bioinformatics,* **39**(3), btad131.
11. Nayfach, S., Roux, S., Seshadri, R., Udwary, D., Varghese, N., Schulz, F., Wu, D., Paez-Espino, D., Chen, I.-M., Huntemann, M., et al. (2021) A genomic catalog of Earth's microbiomes. *Nature biotechnology,* **39**(4), 499–509.
12. Richardson, L., Allen, B., Baldi, G., Beracochea, M., Bileschi, M. L., Burdett, T., Burgin, J., Caballero-Pérez, J., Cochrane, G., Colwell, L. J., et al. (2023) MGnify: the microbiome sequence data analysis resource in 2023. *Nucleic Acids Research,* **51**(D1), D753–D759.
13. Integrative HMP (iHMP) Research Network Consortium (2019) The integrative human microbiome project. *Nature,* **569**(7758), 641–648.
14. Wood, D. E., Lu, J., and Langmead, B. (2019) Improved metagenomic analysis with Kraken 2. *Genome biology,* **20**, 1–13.
15. Blin, K., Shaw, S., Augustijn, H. E., Reitz, Z. L., Biermann, F., Alanjary, M., Fetter, A., Terlouw, B. R., Metcalf, W. W., Helfrich, E. J., et al. (2023) antiSMASH 7.0: New and improved predictions for detection, regulation, chemical structures and visualisation. *Nucleic acids research,* p. gkad344.
16. Steinegger, M. and Söding, J. (2017) MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nature biotechnology,* **35**(11), 1026–1028.
17. Navarro-Muñoz, J. C., Selem-Mojica, N., Mullowney, M. W., Kautsar, S. A., Tryon, J. H., Parkinson, E. I., De Los Santos, E. L., Yeong, M., Cruz-Morales, P., Abubucker, S., et al. (2020) A computational framework to explore large-scale biosynthetic diversity. *Nature chemical biology,* **16**(1), 60–68.
18. Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E. L., Tosatto, S. C., Paladin, L., Raj, S., Richardson, L. J., et al. (2021) Pfam: The protein families database in 2021. *Nucleic acids research,* **49**(D1), D412–D419.
19. Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden, T. L. (2009) BLAST+: architecture and applications. *BMC bioinformatics,* **10**, 1–9.
20. Gilchrist, C. L., Booth, T. J., van Wersch, B., van Grieken, L., Medema, M. H., and Chooi, Y.-H. (2021) Cblaster: a remote search tool for rapid identification and visualization of homologous gene clusters. *Bioinformatics Advances,* **1**(1), vbab016.
21. Köster, J. and Rahmann, S. (2012) Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics,* **28**(19), 2520–2522.
22. Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T. (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome research,* **13**(11), 2498–2504.
23. Medema, M. H., Takano, E., and Breitling, R. (2013) Detecting sequence homology at the gene cluster level with MultiGeneBlast. *Molecular biology and evolution,* **30**(5), 1218–1223.
24. Terlouw, B. R., Blin, K., Navarro-Munoz, J. C., Avalon, N. E., Chevrette, M. G., Egbert, S., Lee, S., Meijer, D., Recchia, M. J., Reitz, Z. L., et al. (2023) MIBiG 3.0: a community-driven effort to annotate experimentally validated biosynthetic gene clusters. *Nucleic acids research,* **51**(D1), D603–D610.
25. Kasuga, G., Tanaka, M., Harada, Y., Nagashima, H., Yamato, T., Wakimoto, A., Arakawa, K., Kawai, Y., Kok, J., and Masuda, T. (2019) Homologous expression and characterization of gassericin T and gassericin S, a novel class IIb bacteriocin produced by Lactobacillus gasseri LA327. *Applied and environmental microbiology,* **85**(6), e02815–18.
26. Selle, K. and Klaenhammer, T. R. (2013) Genomic and phenotypic evidence for probiotic influences of Lactobacillus gasseri on human health. *FEMS microbiology reviews,* **37**(6), 915–935.
27. Becker, A., Schmartz, G. P., Gröger, L., Grammes, N., Galata, V., Philippeit, H., Weiland, J., Ludwig, N., Meese, E., Tierling, S., et al. (2022) Effects of resistant starch on symptoms, fecal markers, and gut microbiota in Parkinson's disease—the RESISTA-PD trial. *Genomics, Proteomics & Bioinformatics,* **20**(2), 274–287.
28. Medema, M. H., Kottmann, R., Yilmaz, P., Cummings, M., Biggins, J. B., Blin, K., De Bruijn, I., Chooi, Y. H., Claesen, J., Coates, R. C., et al. (2015) Minimum information about a biosynthetic gene cluster. *Nature chemical biology,* **11**(9), 625–631.
29. Blin, K., Shaw, S., Kloosterman, A. M., Charlop-Powers, Z., Van Wezel, G. P., Medema, M. H., and Weber, T. (2021) antiSMASH 6.0: improving cluster detection and comparison capabilities. *Nucleic acids research,* **49**(W1), W29–W35.
30. Blin, K., Shaw, S., Steinke, K., Villebro, R., Ziemert, N., Lee, S. Y., Medema, M. H., and Weber, T. (2019) antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic acids research,* **47**(W1), W81–W87.
31. Kautsar, S. A., van der Hooft, J. J., de Ridder, D., and Medema, M. H. (2021) BiG-SLiCE: A highly scalable tool maps the diversity of 1.2 million biosynthetic gene clusters. *Gigascience,* **10**(1), giaa154.
32. Hadjithomas, M., Chen, I.-M. A., Chu, K., Huang, J., Ratner, A., Palaniappan, K., Andersen, E., Markowitz, V., Kyrpides, N. C., and Ivanova, N. N. (2017) IMG-ABC: new features for bacterial secondary metabolism analysis and targeted biosynthetic gene cluster discovery in thousands of microbial genomes. *Nucleic acids research,* **45**(D1), D560–D565.