TITLE

Age-specific and compartment-dependent changes in mitochondrial homeostasis and cytoplasmic viscosity in mouse peripheral neurons

RUNNING TITLE

Aging peripheral neurons

James N. Sleigh[1,2]*, Francesca Mattedi[3,4]*, Sandy Richter[3,5]*, Emily Annuario[3]*, Kristal Ng[3]†, I. Emilie Steinmark[6]†, Iveta Ivanova[6]†, István L. Darabán[3], Parth P. Joshi[3,7], Elena R. Rhymes[1,2], Shirwa Awale[6], Gokhan Yahioglu[8], Jacqueline C. Mitchell[3], Klaus Suhling[6], Giampietro Schiavo[1,2], Alessio Vagnoni[3,9]‡

[1]Department of Neuromuscular Diseases and UCL Queen Square Motor Neuron Disease Centre, UCL Queen Square Institute of Neurology, University College London, London, UK.

[2]UK Dementia Research Institute, University College London, London, UK.

[3]Department of Basic and Clinical Neurosciences, Maurice Wohl Clinical Neuroscience Institute, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK.

[4]Current address: Department of Neuromuscular Diseases, UCL Queen Square Institute of Neurology, University College London, London, UK.

[5]Current address: Department of Biomedical Sciences, University of Padova, Padova, Italy.

[6]Department of Physics, King's College London, Strand, London, UK.

[7]Current address: Sunderland Medical School, University of Sunderland, UK.

[8]Antikor Biopharma Ltd, Stevenage Bioscience Catalyst, Stevenage, UK.

[9]MIA-Portugal – Multidisciplinary Institute of Ageing, University of Coimbra, Coimbra,

Portugal.

‡ Corresponding author: alessio.vagnoni@kcl.ac.uk

*These authors contributed equally

†These authors contributed equally

ABSTRACT

Mitochondria are dynamic bioenergetic hubs that become compromised with age. In neurons, declining mitochondrial axonal transport has been associated with reduced cellular health. However, it is still unclear to what extent the decline of mitochondrial transport and function observed during ageing are coupled, and if somal and axonal mitochondria display compartment-specific features that make them more susceptible to the ageing process. It is also not known whether the biophysical state of the cytoplasm, thought to affect many cellular functions, changes with age to impact mitochondrial trafficking and homeostasis. Focusing on the mouse peripheral nervous system, we show that age-dependent decline in mitochondrial trafficking is accompanied by reduction of mitochondrial membrane potential and intra-mitochondrial viscosity, but not calcium buffering, in both somal and axonal mitochondria. Intriguingly, we observe a specific increase in cytoplasmic viscosity in the neuronal cell body, where mitochondria are most polarised, which correlates with decreased cytoplasmic diffusiveness. Increasing cytoplasmic crowding in the somatic compartment of DRG neurons grown in microfluidic chambers reduces mitochondrial axonal trafficking, suggesting a mechanistic link between the regulation of cytoplasmic viscosity and mitochondrial dynamics. Our work provides a reference for studying the relationship between neuronal mitochondrial homeostasis and the viscoelasticity of the cytoplasm in a compartment-dependent manner during ageing.

INTRODUCTION

Ensuring precise distribution and functionality of mitochondria is essential for maintaining cellular homeostasis. The complex architecture and long distances between nerve terminals and soma of neurons cause these cells to be highly

dependent on mechanisms for long-range cytoskeletal transport of mitochondria and

on continuous mitochondrial supply and functionality along their extended cellular

architecture. This is a particularly critical challenge for neurons, which are post-mitotic

and can live throughout the adult life of an organism. It is therefore not surprising that

mitochondrial dysfunction is a recognised hallmark of ageing and neurodegeneration

(López-Otín et al., 2023).

We have previously shown that axonal transport of mitochondria decreases in the

peripheral nervous system of ageing *Drosophila melanogaster* (Vagnoni et al., 2016)

and that neuronal ageing phenotypes, such as oxidative stress and focal protein

accumulation, can be manipulated by modulation of mitochondrial transport (Vagnoni

& Bullock, 2018; Vagnoni et al., 2016). Mitochondrial motility was shown to be

downregulated during ageing to varying degrees also in *C. elegans* motor neurons

and in the mouse central nervous system neurons *in vivo* (Mattedi & Vagnoni, 2019;

Morsci et al., 2016; Takihara et al., 2015). Despite the consistent observation that

mitochondrial trafficking is slowed down during ageing, it remains unclear whether this

directly correlates with functional changes of the mitochondria.

The biophysical properties of the neuronal cytoplasm are under increasing scrutiny in

the context of neurodegeneration (Alberti & Hyman, 2016; Elbaum-Garfinkle &

Brangwynne, 2015), although the effect of age on these properties is less well studied.

Whether fundamental properties of the neuroplasm, such as viscosity and diffusion,

undergo significant changes that can affect intracellular cargo trafficking and

functionality during ageing is currently unknown.

A study that combines analyses of the motile behaviour of neuronal cargoes and their

functional state with the evaluation of the biophysical properties of the cytoplasm in

young and old neurons has been missing, partly due to lack of suitable experimental

systems to perform this type of work and the challenge of studying these processes in

live neurons throughout ageing. Interrogating cellular processes in animal models *in*

vivo is indispensable for ageing studies, although gaining mechanistic insight can be

difficult in this context (Sleigh et al., 2017a). For example, neuronal cells *in situ* often

have their soma, axons and dendrites embedded in different tissues and organs and

so studying functional compartmentalisation *in vivo* becomes challenging. In an ageing

context, being able to complement *in vivo* studies with a species-specific *in vitro* model

of neuronal ageing would be particularly advantageous.

In this study, by focusing on the mouse peripheral nervous system, we filled this gap

through comparative analyses of mitochondrial trafficking in dorsal root ganglion

(DRG) neurons *in vitro* and sciatic nerve axons *in vivo* from both young and aged mice.

Using the DRG model, we also monitored other important aspects of mitochondrial

biology within neurons, including mitochondrial membrane potential, calcium uptake

and intra-mitochondrial viscosity, and we revealed age- and compartment-dependent

alterations in these properties. We expanded on these data by quantifying intracellular

viscosity and diffusion in the neuronal cytoplasm of the somal and axonal

compartments and uncover a mechanistic link between increased crowding and

viscosity in the cell bodies and reduced mitochondrial axonal trafficking. With this work,

we propose a novel model that correlates mitochondrial homeostasis with the

viscoelasticity of the neuronal cytoplasm during ageing.

RESULTS

Mitochondrial trafficking declines with age in mouse peripheral neurons both *in*

vitro* and *in vivo

To better characterise the effect of ageing on mitochondrial trafficking in mouse

peripheral neurons, we cultured sensory neurons from DRG of young (3-4 months)

and old (19-22 months) mice (Fig. 1A) and analysed mitochondrial axonal transport

by live imaging with MitoTracker. DRG neurons grow robustly in culture after

dissection, retain their age-dependent transcriptional profile (Gumy et al., 2011) and

have previously been used to study axonal cargo trafficking during ageing (Mellone et

al., 2013; Stavoe et al., 2019). Time-lapse recording and quantification of

mitochondrial axonal transport showed abundant mitochondrial movement in both the

anterograde and retrograde directions, with a small proportion of mitochondria

displaying bidirectional motility (Fig. 1B-C, Fig. S1A-B) (Klinman & Holzbaur, 2015; Lin

et al., 2017; Mellone et al., 2013; Pandey et al., 2022). Neurons obtained from old

mice showed a strong reduction in the percentage of moving mitochondria (Fig. 1B-C,

Fig. S1A-B, Movie 1), but no significant change in the number of mitochondria present

in axons (Fig. 1D), which correlated with an increase in their overall size (Fig. 1E-G,

Fig. S1C-D). Moreover, the speed and displacement of these organelles were largely

unaffected by age, both in the anterograde (Fig. 1H) and retrograde directions (Fig.

1I).

To assess whether the age-dependent decline in mitochondrial transport observed in

cultured DRGs was preserved in peripheral neurons *in vivo*, we analysed

mitochondrial axonal transport in intact sciatic nerve axons of anaesthetised MitoMice

(Fig. 2A), in which neuronal mitochondria are marked by cyan fluorescent protein

(CFP) (Fig. 2B-C and Movie 2) (Misgeld et al., 2007). Old mice displayed a clear

reduction in mitochondrial motility (Fig. 2D and Movie 2), which was accompanied by

a reduction in the overall number of axonal mitochondria (Fig. 2E) and a stark increase

in mitochondrial area (Fig. 2F, Fig. S2A-B). When the proportion of mitochondrial

motility was assessed, no difference between young and old neurons was identified

(Fig. S2C) in either anterograde or retrograde directions (Fig. S2D). This indicates that

the decline in the number of motile mitochondria might not be due to a net change in

the proportion of moving mitochondria, rather a potential imbalance in mitochondrial

biogenesis and/or turnover that leads to altered mitochondrial content in axons. While

the motile properties of the retrograde-moving mitochondria remained unchanged with

age (Fig. 2H), the speeds and displacement of the fewer anterograde-moving

mitochondria were surprisingly upregulated in older mice (Fig. 2G), contrasting with

what observed in cultured DRG neurons. Interestingly, the distributions of

mitochondrial speed and displacement measured in the sciatic nerve and cultured

neurons were significantly different (Fig. 1H-I, 2G-H, and Table S1), potentially

indicating a different regulation of the molecular motor machinery responsible for the

movement of these organelles *in vivo* and *in vitro*. Overall, these data show that

mitochondrial axonal trafficking is affected by age in peripheral nerves both *in vitro*

and *in vivo*, although the underlying mechanisms may differ.

Mitochondrial membrane potential is compartment-dependent and decreases

with age and declining mitochondrial viscosity

To understand if the trafficking defects are associated with altered mitochondrial

functionality, we first asked whether the membrane potential ($\Delta\Psi_m$) of these organelles

was altered between young and old DRG neurons. We stained the neurons with a low

concentration of the live dye tetramethylrhodamine methyl ester (TMRM), which

accumulates in active mitochondria, as well as MitoTracker Green (MTG) to label all

mitochondria (Fig. 3A-D). Calculating the ratio of TMRM to MTG fluorescence has

previously been used to determine $\Delta\Psi_m$ in live cells, where a higher ratio indicates

more polarised mitochondria (Mitra & Lippincott-Schwartz, 2010). Using this method

here reveals that the membrane potential of neuronal mitochondria decreases with

age (Fig. 3E and Fig. S3A-B) and that the axonal compartment is characterised by

mitochondria with substantially lower membrane potential, independently of age (Fig.

3F-G). Using the ratiometric dye JC-1, we confirmed that axonal mitochondria are

more depolarised than somal mitochondria (Fig. S3C-D), further validating our

protocol to estimate $\Delta\Psi_m$, using the TMRM/MTG ratio.

It is possible that decreased trafficking and altered membrane potential of

mitochondria during ageing are mechanistically linked, although the two phenotypes

could also manifest independently. To test if decreased mitochondrial trafficking has

a direct effect on membrane potential in DRGs, we used an RNAi construct to reduce

the abundance of Trak1, which is essential to drive mitochondrial motility in axons (van

Spronsen et al., 2013). Trak1 knockdown in young DRG neurons reduced

mitochondrial transport (Fig. S3E) but had no impact on membrane potential (Fig. S3F-

G). This indicates that acute reduction of mitochondrial axonal transport does not

directly affect $\Delta\Psi_m$, suggesting that the two phenotypes in the aged neurons are not

mechanistically linked. Nevertheless, we cannot exclude that chronic reduction of

mitochondrial transport during ageing or a time-specific reduction in older neurons

could have a knock-on effect on the maintenance of $\Delta\Psi_m$.

Manipulations that decrease mitochondrial membrane potential have been shown to

affect the viscosity of these organelles in HeLa and COS7 cells (Chambers et al.,

2018; Chen et al., 2020; Guo et al., 2022; Steinmark et al., 2019), while altered

mitochondrial membrane viscosity has been observed in brains of people affected by

Alzheimer's disease (Mecocci et al., 1997) and in murine neurons during

neurodegeneration and ageing (Eckmann et al., 2014; Kuter et al., 2016). We

therefore asked if the age-dependent changes in $\Delta\Psi_m$ that we observe in DRGs might

be accompanied by changes in the viscosity of the mitochondria. To do so, we

performed fluorescence lifetime imaging microscopy (FLIM) of neurons incubated with

the BODIPY-based fluorescent molecular rotor FMR-1 targeted to the mitochondrial

matrix (Fig. 3H) (Steinmark et al., 2019). Fluorescence lifetime of FMR-1 responds to

changes in viscosity environments, where a longer fluorescence lifetime indicates

higher viscosity. Neuronal mitochondria display a heterogeneous viscosity profile both

in the cell bodies and axons (Figure 3H), in accordance with previous work carried out

in HeLa and COS7 cells (Chambers et al., 2018; Steinmark et al., 2019). Similar to

$\Delta\Psi_m$, average mitochondrial viscosity decreases during ageing as indicated by

decreased lifetime (Fig. 3I and Fig. S3H-I). In this case, however, the viscosity of the

mitochondria remains largely unchanged between the cell bodies and axons in both

young and old neurons (Fig. 3J-K). These results suggest that while the striking

difference in $\Delta\Psi_m$ between somal and axonal mitochondria is unlikely to be coupled to

organelle viscosity (as this remains constant across compartments), the decline of

$\Delta\Psi_m$ and mitochondrial viscosity during age may be linked.

Ca^{2+} buffering is unaffected in DRG neurons cultured from young and old mice

Loss of $\Delta\Psi_m$ has long been thought to negatively affect Ca^{2+} accumulation in

mitochondria (Brini et al., 1999; Duchen, 2000) and, in mouse sensory neurons,

$[Ca^{2+}]_m$ uptake causes a transient decrease in $\Delta\Psi_m$ (Duchen, 2000). Increased

mitochondrial viscosity was also suggested to correlate with augmented cytosolic

calcium in HeLa cells (Steinmark et al., 2019). We therefore asked whether the age-

dependent changes in the $\Delta\Psi_m$ and mitochondrial viscosity are accompanied by

alterations in $[Ca^{2+}]_m$ uptake. We depolarised DRG cultures by transient stimulation

with 50 mM KCl and measured the fluorescent signal of mitochondrial and cytosolic

Ca^{2+} using the Rhod-2-AM and Fluo4-AM dyes, respectively (De vos et al., 2012;

Duchen, 2000) (Fig. S4A-J). Combined measurements from both the somal and

axonal compartments showed that there was no overall difference in the timing and

abundance of mitochondrial Ca^{2+} uptake and release (Rhod-2-AM) between young

and old cells (Fig. S4K,M,O). Likewise, we did not detect any significant difference in

the dynamics of cytoplasmic Ca^{2+} (Fluo4-AM) between young and old neurons

following neuronal stimulation (Fig. S4L,N,P). Taken together, these data show that

age does not affect Ca^{2+} homeostasis in cultured DRGs.

Age-dependent increase in cytoplasmic viscosity is inversely correlated with

diffusiveness in neuronal cell bodies but not in axons

Previous studies have shown that hypertonic treatments, which acutely increase

molecular crowding and cytoplasmic viscosity, affect cytoskeletal dynamics in fission

yeast (Molines et al., 2022), alter ER-to-Golgi transport in COS7 cells (Lee & Linstedt,

1999) and abolish endosomal motility in mammalian immune cells (Nunes et al.,

2015). To test whether decreased mitochondrial transport in ageing DRG neurons

correlates with a change in fluid-phase viscosity of the cytoplasm, we performed time-

resolved fluorescence anisotropy imaging (TR-FAIM, Fig. 4A) (Clayton et al., 2002; Siegel et al., 2003). TR-FAIM measures the depolarisation of fluorescence emission of a fluorescein derivative (2′,7′-bis-(2-carboxyethyl)-carboxyfluorescein, BCECF) caused by rotational Brownian motion, and so it is used to obtain information about the viscosity of the cellular microenvironment (Dix & Verkman, 1990). Anisotropy decay measurements and fitting (Fig. S5A and see Methods) showed that the rotational correlation time of the BCECF is significantly reduced in older cells leading to slower depolarisation (Fig. 4B), which is indicative of higher cytoplasmic viscosity in old DRGs (Fig. 4C). Because of the specialised architecture of neurons, we hypothesised that the somal and axonal compartments might comprise different viscosity environments. Measurement of the anisotropy decay in cell bodies and axons revealed that, although the two compartments have similar viscosity values in young cells (Fig. 4D, F and Fig. S5B), the viscosity of the cell bodies undergoes a significant increase in aged cells (Fig. 4E-F and Fig. S5C-D), whereas the axonal viscosity remains unchanged during age (Fig. 4D-F and Fig. S5B-C, E). These results suggest that axonal viscosity alone is unlikely to explain the observed decrease in mitochondrial axonal transport.

Fluid-phase viscosity affects the diffusiveness of the cytoplasmic milieu, as determined by the Stokes-Einstein law (Miller & Walker, 1924). To gain more insight into cytoplasmic diffusiveness in our *in vitro* cell model of ageing, we performed rheological studies by measuring the diffusion rates of genetically-encoded multimeric nanoparticles (GEMs) (Delarue et al., 2018). GEMs can be visualised as bright spherical particles of uniform shape and size that are suitable for microrheological studies in different cellular contexts (Delarue et al., 2018; Molines et al., 2022; Shu et

al., 2022). We transduced DRGs with a lentiviral vector encoding sapphire-tagged

GEMs of 40 nm diameter (Fig. 5A, D) and tracked particle movement in cell bodies

and axons (Fig. 5B-C, E-F and Movie 3-4). Single particle tracking and MSD analysis

showed that GEM movement is consistent with sub-diffusive motion (Fig. 5G and Fig.

S6A), in accordance with previous reports in non-neuronal cell types (Delarue et al.,

2018; Molines et al., 2022). In both young and old neurons, the diffusion rate of axonal

GEMs is significantly lower than the rate of GEMs freely diffusing in cell bodies (Fig.

5G-I and Fig. S6), suggesting that the specialised axonal architecture poses a

significant physical constraint for particle diffusion. The diffusion of somal GEMs is

reduced in aged cultures (Fig. 5G-I), in accordance with the observed increase in

viscosity. Interestingly, although the axonal viscosity does not change during ageing

(Fig. 4D-F, Fig. S5B-C, E), GEMs display a small, albeit significant, reduction in

diffusiveness in aged culture (Fig. 5G-I), indicating that mechanisms other than

viscosity regulate axonal diffusion during ageing.

Based on these experiments, we conclude that the behaviour of cytoplasmic fluid-

phase viscosity during ageing is defined by the cellular compartment, with potentially

distinct effects on somal and axonal biology. Indeed, while increased viscosity likely

contributes to reduced cytoplasmic diffusion in the neuronal cell bodies, axonal

viscosity is not a driver of the decreased diffusiveness of the axonal cytoplasm nor of

the reduced active mitochondrial axonal transport during ageing.

Challenging DRG neuron cell bodies with crowding agents reduces axonal

mitochondrial transport.

It is conceivable that the stark increase in cytoplasmic viscosity observed in the soma

of aged DRGs is sufficient to impact mitochondrial dynamics, including mitochondrial

export into the axon, reducing axonal transport rates in old cells. To test this

hypothesis, we set out to manipulate the viscosity specifically within somal

compartments of young DRG neurons cultured in microfluidic chambers (MFCs)(Fig.

6A, Fig. S7A) to mimic the increased viscosity observed in the aged DRGs. We took

advantage of the physical compartmentalisation afforded by MFCs and devised a

pulse-chase experiment whereby the somal compartment was stained for 30 min with

MitoTracker Green (MTG) following incubation with low doses of either polyethylene

glycol (PEG) or sorbitol (Fig. 6B), two crowding agents routinely used to increase

intracellular viscosity (Chambers et al., 2018; Molines et al., 2022). The axonal

compartment was visualised by staining with CellMask Orange (CMask) to assess

axonal growth (Fig. 6B-E). After dyes wash-off, we immediately imaged mitochondrial

axonal transport in the chamber microgrooves. We reasoned that any MTG-positive

mitochondria observed in the axons would have originated from the somal

compartment after being stained during the 30-min pulse. Low-magnification images

consistently showed that while MTG-positive mitochondria can travel long distances

into the axons-filled microgrooves of untreated cells to reach the synaptic

compartments, PEG and sorbitol-treated cells only display faint MTG stain in the

proximal axons (Fig. 6C-E). High-resolution time-lapse imaging and quantification of

mitochondrial axonal transport reveals a drastic reduction of mitochondrial export from

the somal into the axonal compartment in both PEG and sorbitol-treated cells,

compared to controls, as indicated by transport fluxes in the initial segment of the

microgrooves (Fig. 6F-G, Movie 5). As a result, mitochondrial distribution and

occupancy of the axonal space is strongly reduced by these treatments (Fig. S7B,

Movie 5). Interestingly, the average velocity of mitochondria is also significantly

reduced after sorbitol treatment (Fig. 6H). This suggests the higher viscosity

environment of the cell soma induces a lasting effect on the molecular components

driving mitochondrial trafficking, which cannot be reversed by translocating into the

less crowded axonal environment.

The DRG cell body area was not noticeably affected by the treatments with these

crowding agents (Fig. S7C-D), at least at the concentrations and incubation times used

in this study. This is in line with the unchanged cell body area observed in young and

old DRGs (Fig. S6D) and indicates our relatively mild treatments do not trigger an

overt osmotic stress. In conclusion, we demonstrate that viscosity manipulations

restricted to the neuronal cell bodies, mimicking an aged state, are sufficient to affect

mitochondrial transport rates in the axonal compartment.

DISCUSSION

In this study, we report several new findings that define mitochondrial homeostasis

and cytoplasmic biophysics in aging neurons, spanning from mitochondrial axonal

trafficking to mitochondrial membrane potential and calcium buffering, and including

mitochondrial and cytoplasmic viscosity, and cytoplasmic diffusion. By providing a

holistic view of the changes occurring in the mouse peripheral nervous system as it

ages, we uncover significant differences that display age specificity and are

compartment dependent (Fig. 7).

<u>- Mitochondrial axonal trafficking</u>

Altered mitochondrial axonal trafficking is an established feature of ageing peripheral

neurons in *Drosophila* and *C. elegans in vivo* (Morsci et al., 2016; Vagnoni et al., 2016)

and of mouse sciatic nerve *ex vivo* (Milde et al., 2015). We now show that the number

of motile axonal mitochondria declines both in aged DRG neurons *in vitro* and in the

sciatic nerve *in situ* (Fig. 7), although the underpinning mechanisms are likely to be

different. Firstly, mitochondria travel at significantly different speeds between *in vitro*

and *in vivo* systems. Secondly, the total number of axonal mitochondria remains stable

in vitro showing that decreased transport may not be sufficient to determine the

steady-state distribution of mitochondria in axons, as previously suggested (Milde et

al., 2015; Vagnoni et al., 2016). On the other hand, decreased mitochondrial motility

in the sciatic nerve of older mice may be coupled to impairment in biogenesis or

excessive mitochondrial clearance that could lead to an overall reduced number of

axonal mitochondria. Our result that mitochondrial area increases in aged axons both

in vitro and *in vivo* lend support to the notion that altered mitochondrial biogenesis

and/or fission and fusion may be a conserved and key feature of neuronal ageing. At

the same time, the strength of our comparative study underlies that the DRG *in vitro*

system is not a like-for-like replacement of the sciatic nerve *in vivo*: specific

mechanisms can differentially affect mitochondrial dynamics *in vitro* and *in vivo* to

determine overall mitochondrial distribution.

Decreased mitochondrial motility is accompanied by a surprising increase in the

processivity of anterograde mitochondria in aged neurons *in vivo*. It is possible this is

a compensatory mechanism triggered by age, similar to upregulation of cargo

trafficking observed in response to peripheral nerve injury (Mar et al., 2014; Milde et

al., 2015). This result is interesting as it shows that the decline in mitochondrial motility

during ageing can be partly reversed in mammalian neurons, as demonstrated

previously in ageing *Drosophila* neurons (Vagnoni & Bullock, 2018). Understanding

the mechanisms behind the transport upregulation in older neurons might therefore

be of therapeutic interest for age-dependent neurodegeneration when decline in cargo

trafficking can play an important role in disease progression (Berth & Lloyd, 2023;

Sleigh et al., 2019).

<u>- Mitochondrial membrane potential and ion homeostasis</u>

Decline in mitochondrial functionality is an active focus of research in the ageing field

(Sharma et al., 2019; Sun et al., 2016) with mitochondrial oxidative stress (Stefanatos

& Sanz, 2018) and mitophagy (Caponio et al., 2022; Markaki et al., 2023; Rappe &

McWilliams, 2022) thought to play a potentially critical role during ageing. The

observation that the $\Delta\Psi_m$ decreases in aged DRG neurons (Fig. 7) implies that the

mitochondria become overall less functional and is consistent with reduced

mitochondrial bioenergetics observed in iNeurons derived from old human donors

(Kim et al., 2018). Upregulation of retrograde transport of dysfunctional mitochondria

with reduced $\Delta\Psi_m$ was observed in murine hippocampal (Lin et al., 2017) and chicken

DRG neurons (Miller & Sheetz, 2004) after acute mitochondrial stress, suggesting this

could be a mechanism for recycling of dysfunctional mitochondria in the cell soma. We

now show that this potential safeguarding mechanism may be impaired during ageing

due to a strong reduction in mitochondrial motility in both anterograde and retrograde

directions. Interestingly, we find that the mitochondria that reside in the cell body are

much more polarised than axonal mitochondria and this is observed independently of

age. These results suggest that single axonal mitochondria may be metabolically less

active than the those found in the interconnected mitochondrial network within the cell

body. Alternatively, axonal mitochondria might be more prone to damage, possibly

due to a reduced turnover in the axon, where basal mitophagy may occur at different

rates than in the cell body (Sung et al., 2016), or to impaired fission and fusion

dynamics as a result of decreased trafficking. Whatever the case, our observations

support the hypothesis of a mitochondrial functional gradient along the axons whereby

mitochondria become progressively less active as the distance from the cell body

increases (Baranov et al., 2019; Hagemann et al., 2022). In future studies, it would be

interesting to use microfluidics chambers, for example similar to the ones we

introduced in this work, to apply stressors to the proximal and distal compartments

and observe if mitochondrial transport is differently affected in young and old neurons.

The calcium content of mitochondria has been correlated with their membrane

potential (Brini et al., 1999; Duchen, 2000) and motile behaviour (Chang et al., 2011;

Niescier et al., 2018), and acute increase in cytoplasmic calcium levels can suppress

mitochondrial motility (MacAskill et al., 2009; Wang & Schwarz, 2009). Interestingly,

we find that even though mitochondria are overall less polarised in aged DRG cultures

when their motility declines, this does not affect their capacity to buffer calcium or the

wider neuronal excitability in this context (Fig. 7). Thus, it is conceivable that the

population of healthy mitochondria in the older DRG neurons is sufficient to

compensate for the dysfunctional organelles, although we cannot exclude that the

calcium uptake capability is retained in depolarised organelles.

- Mitochondrial viscosity

Reduced $\Delta\Psi_m$ is accompanied by an overall reduction in the intramitochondrial

viscosity in aged DRG (Fig. 7), which could conceivably reflect an adaptation to

metabolic activity (Persson et al., 2020; Puchkov, 2013) that becomes altered with

age. Alternatively, changes in mitochondrial membrane fluidity as a result of oxidative

damage (Eckmann et al., 2014; Mecocci et al., 1997) or intra-mitochondria molecular

dilution due to enlarged area (Fig. 1-2) could explain reduced viscosity. Interestingly,

mitochondria of human fibroblasts that are deficient for complex I activity display

enhanced protein diffusion (Koopman et al., 2008), likely a sign of reduced viscosity,

while mitochondria hyperpolarisation correlates with increased viscosity in HepG2

cells (Bulthuis et al., 2023). Thus, our data support the idea that the reduced $\Delta\Psi_m$ and

mitochondrial viscosity in aged DRGs are mechanistically linked.

Contrarily to $\Delta\Psi_m$, however, mitochondrial viscosity across compartments remained

uniform, which suggests independent mechanisms might regulate $\Delta\Psi_m$ and viscosity

in different compartments. Mitochondrial viscosity was shown to be selectively

unaffected by acute osmotic stress compared to other organelles (Chambers et al.,

2018). Therefore, it is possible that mitochondrial viscosity is less sensitive to stressors

that could determine a difference in $\Delta\Psi_m$ across compartments. Overall, our study

adds significantly to the notion of mitochondrial functional specialisation that has been

observed across different neuronal populations (Fecher et al., 2019) and within

subcellular compartments (Faitg et al., 2021). Future studies should be aimed at

further characterising the mitochondrial microviscosity environment to assess whether

sub-mitochondrial domains are differently affected by age in different cellular

compartments.

<u>- Cytoplasmic viscosity.</u>

Whether cytoplasmic viscosity and diffusion undergo significant changes during

ageing that can affect intracellular trafficking is not known. Previous studies showed

that increased viscosity reduces both endosomal trafficking and microtubule dynamics

(Molines et al., 2022; Nunes et al., 2015). Changes in cytoplasmic viscosity are

thought to depend on alterations of molecular crowding, with likely repercussions on

wider cellular functionality (Leduc et al., 2012; Metzner et al., 2022; Sabharwal &

Koushika, 2019; Sood et al., 2018). Using TR-FAIM, we find that axonal cytoplasmic

viscosity does not change with age and so it is unlikely to have a major impact on the

biology of the axon, including the decrease in mitochondrial motility that we observe

in older DRG neurons (Fig. 7). However, we find that, contrarily to axons, the viscosity

of the cell bodies increases in aged cultures. Why cell bodies selectively display

increased viscosity is not clear. This could result from an imbalance in protein

translation (Delarue et al., 2018), a potential temperature difference across

compartments as the cell ages (Persson et al., 2020), for example as a consequence

of reduced metabolic activity (Parry et al., 2014), or be the product of compartment-

dependent difference in plasma membrane stiffness (Kubánková et al., 2019;

Lamoureux et al., 2010; Sugawa et al., 1996). Whatever the case, we demonstrated

that acutely increasing cell body viscosity using molecular crowders is sufficient to

strongly reduce mitochondrial export into the axons and overall transport rates. We

are aware of the challenge posed by recapitulating age-dependent processes with

acute, albeit well controlled, experimental treatments. Nevertheless, our results are

strongly suggestive of a scenario in which the age-dependent increase in cell body

cytoplasmic viscosity contributes to reducing axonal transport rates, with a likely wider

impact on cellular functionality in aged cells. The mechanism behind this process is

not known. Our observation that mitochondria appeared more clustered and less

dynamics in the DRG cell bodies after PEG and sorbitol treatments (Movie 6),

suggests this might be a likely cause for the decreased mitochondrial export into the

axons. However, it would be valuable to establish in future studies if viscosity alters

the molecular composition of the AIS-like filter zone in cultured DRGs (Farías et al.,

2015; Gumy et al., 2017) to determine the abundance of selected cargoes entering

the axons in ageing cells.

The apparent reduction of mitochondrial axonal transport speed after sorbitol

treatment of the cell bodies is interesting: it suggests that, in the highly crowded

environment of the cell body, the mitochondria and/or the mitochondrial transport

machinery may undergo molecular changes that affect their long-term trafficking. At

first, this result would not seem consistent with the unchanged mitochondrial velocity

measured in the axons of aged, cultured neurons (Fig. 1H-I). However, we think that

the acute increase in viscosity reveals that reduced speeds may be the precursor to

overall reduced mitochondrial motility observed during ageing (Fig. 1C), when

exposure to increased viscosity is likely to occur more gradually.

These data further support the hypothesis that failed mitochondrial export from the

soma is a primary contributor of reduced axonal trafficking during ageing. In turn,

reduced mitochondrial supply may trigger an imbalance in mitochondrial fission and

fusion in axons leading to enlarged organelles. It will be intriguing to test the validity of

our model in future work by studying if manipulating cell body viscosity directly affects

a pool of predominantly axonal mitochondria in aging neurons.

- Molecular diffusion

We find that the molecular diffusion in the neuronal cytoplasm, ascertained using the

GEM probe, decreases with age in both soma and axons (Fig. 7). While this is

consistent with the increased viscosity of the cell soma, it seems apparently at odds

with the constant axonal viscosity measured by TR-FAIM. However, our data are in

accordance with the notion that macromolecular crowding in complex cellular

environments can cause deviation from the Stokes-Einstein law that relates diffusion

and viscosity (Zustiak et al., 2011). Moreover, TR-FAIM measures rotational diffusion

of a dye whereas GEMs explore lateral diffusion in the mesoscale. Therefore, we

speculate that older axons may develop proteinaceous aggregates and/or molecular

condensates that, while not sufficient to alter overall viscosity, can provide significant

spatial hindrance by increasing local crowding in small calibre DRG axons. This could

explain both the slower diffusion of the 40 nm GEM particles and the well documented

decrease in organelle motility observed during ageing and neurodegeneration. In

future studies, it would be valuable to determine whether local viscosity and diffusion

in proximal and distal axonal regions display different rates, and to establish if this is

a general feature of the nervous system or confined to a specific subset of neurons.

Ultimately, it would be relevant to understand whether age-dependent

neurodegenerative diseases are characterised by specific viscosity and diffusion

patterns and whether this could be exploited to model disease progression.

Overall, this work will provide an important reference for future studies aimed at better

understanding mitochondrial functionality and the role of the cytoplasmic milieu in

cellular homeostasis.

ACKNOWLEDGEMENTS

We thank personnel of the Denny Brown Laboratory for assistance with mouse

colonies, Gábor Mórotz for suggestions regarding calcium imaging experimental

setup, BFK Lab for assistance with MSD analyses, the Wohl Cellular Imaging Centre

at King's College London for help with light microscopy, Frank Hirth and members of

the Vagnoni lab for comments on the draft manuscript. Fig. 1A, 2A, 6A and 7 were

created using https://www.biorender.com. This project was funded by a NC3Rs David

Sainsbury Fellowship NC/N001753/2 and SKT grant NC/T001224/1 (to A.V.), a

BBSRC CTP award BB/T508597/1 (to A.V.), Royal Society Research Grant

RGS\R2\180450 (to A.V.), an Academy of Medical Sciences Springboard Award

SBF004/1088 (to A.V.), a van Geest Fellowship in Dementia and Neurodegeneration

and van Geest PhD studentship awards (to A.V.), a King's Together Multi and

Interdisciplinary Research Scheme Wellcome Trust Institutional Strategic Support

Fund 204823/Z/16/Z (to A.V.), an European Union's Horizon 2020 research and

innovation programme, grant agreement No. 857524 (to A.V.), and by the Alzheimer's

Research UK London Network Centre grant ARUK-NC2020-KCL, a BBSRC award

BB/R004803/1 (to K.S.), Medical Research Council awards MR/S006990/1 and

MR/Y010949/1 (to J.N.S.), the UCL Neurogenetic Therapies Programme funded by

The Sigrid Rausing Trust (to J.N.S., G.S.), a Wellcome Trust Sir Henry Wellcome

Postdoctoral Fellowship 103191/A/13/Z (to J.N.S.), Wellcome Trust awards

107116/Z/15/Z and 223022/Z/21/Z (to G.S.), and a UK Dementia Research Institute

Foundation award UKDRI-1005 (to G.S.). D.S. and I.E.S. were supported by a BSCB

Studentship and a LIDo iCASE Studentship, respectively.

The authors declare no conflict of interest.

MATERIALS AND METHODS

Study approval

Experimentation involving mice was performed under license from the UK Home

Office in accordance with the Animals (Scientific Procedures) Act (1986). Work

conducted at University College London was approved by the Queen Square Institute

of Neurology Ethical Review Committee and experiments performed at King's College

London were approved by King's College Denmark Hill Ethical Review Body.

Animals

Mice were maintained under a 12 h light/dark cycle at constant room temperature

(≈21°C) with free access to water and food. Cages were enriched with nesting

material, plastic/cardboard tubes and wooden chew sticks. MitoMice (B6.Cg-Tg(Thy1-

CFP/COX8A)S2Lich/J – RRID:IMSR_JAX:007967) were obtained from The Jackson

Laboratory and maintained as hemizygous × wild-type breeding pairs on a C57BL/6J

background. Presence of the CFP transgene was confirmed by a standard condition

polymerase chain reaction using forward primer 5'- GAC GTA AAC GGC CAC AAG

TT-3' and reverse primer 5'- GTC CTC CTT GAA GTC GAT GC-3'. C57BL/6J wild-

type mice were obtained from Charles River Laboratories UK Ltd. Mice of both sexes

were used in all experiments. DRGs obtained from MitoMice were used for live

imaging of mitochondrial transport in mass cultures and Ca^{2+} imaging experiments,

while DRGs from C57BL/6J mice were used for live imaging of mitochondrial

membrane potential and mitochondrial viscosity, TR-FAIM and microrheology

experiments, and mitochondrial transport in microfluidic chambers. Young and old

cells were imaged in the same or in contemporary experimental sessions, whenever

possible, rather than 20 months apart.

DRG dissection and mass culture

DRGs were dissected from the spinal cord of euthanised adult mice (Sleigh et al.,

2016). Approximately 10 DRG pairs (*i.e.*, 20 DRG) were collected for each mouse,

and kept on ice in Ca^{2+}/Mg^{2+}-free HBSS with 1% Pen/Strep until dissociation. DRGs

were cultured as previously described (Sleigh et al., 2017b); briefly, after removing the

HBSS, the DRGs were incubated in 4 mg/ml collagenase type II (Thermo Fischer

Scientific #17101015) and 4.7 mg/ml dispase type II (Sigma-Aldrich #D4693) in

Ca^{2+}/Mg^{2+}-free HBSS at 37°C for 10-15 min with periodic gentle agitation. The

collagenase/dispase solution was replaced with pre-warmed F12 medium (F12

(Gibco) containing 10% heat-inactivated-FBS (hi-FBS) and 1% Pen/Strep (both from

Thermo Fischer Scientific)). Upon centrifugation, cells were resuspended in F12

medium and triturated using fire-polished glass Pasteur pipettes. After centrifugation

at 200 *g* for 5 min, cells were resuspended in 320 μl of fresh DRG medium (F12, 10%

hi-FBS, 1% Pen/Strep, 20 ng/ml mouse GDNF (Peprotech #450-44), 50 ng/ml mouse

NGF (Peprotech #450-34), 20 ng/ml human NT3 (Peprotech #450-03)) and plated as

single 20 μl drops (1-2 drops/well or coverslip) at 37°C, 5% CO_2 for 1-2 h. 0.3-1 ml of

DRG medium was then gently added to each well and cells were returned to the

incubator. Cells were plated in μ-Slide 8-well glass bottom imaging chambers (Ibidi

#80827), in 8- and 4-well Nunc Lab-Tek chambered coverglass (Thermo Fischer

Scientific) or 18-mm round glass coverslips (Marienfeld), depending on the

experiments. 20 μg/ml laminin (Corning #354232) in Ca^{2+}/Mg^{2+}-free HBSS (Gibco)

was used to coat the culture surface for at least 2 h before plating.

Microfluidic chambers (MFCs) preparation and DRG cultures

A preparation of 10:1 polydimethylsiloxane (PDMS) and curing agent (Sylgard 184

Silicone Elastomer Kit) was vigorously mixed (approximately 3.5 ml per MFC mould),

then bubbles removed in a desiccator vacuum pump for 30 min. PDMS mix was

carefully poured into 500 µm MFCs moulds, then baked at 65˚C for 60-75 min. Baked

PDMS was removed from moulds and punched to create chips. Glass-bottomed

dishes were coated with poly-D-lysine for 3-16 h to create static attraction to the PDMS

chips, washed 5 times with sterile H_2O, then PDMS chips were stuck down. The

channels were treated with 0.8% BSA overnight, then coated with laminin as described

previously (Rhymes et al., 2022).

DRGs were isolated as described above and resuspended in 40 µl of DRG medium.

10 µl of cell suspension were added to one well of the MFC cell body compartment

(cell body well 1, Fig. S7A) held at a slightly tilted angle to aid distribution throughout

the cell body channel (Fig. S7A), while 20 µl of DRG medium were added to the

corresponding axonal well (axon well 1, Fig. S7A). Cell suspension was incubated for

approximately 1 h at 37°C before adding 140 µl of DRG medium dropwise across both

wells of the cell body compartment (cell body well 1 and 2, Fig. S7A), and 80 µl across

the wells of the axonal compartment (axon well 1 and 2, Fig. S7A).

Live imaging and analysis of mitochondrial transport and morphology in mass

cultures of DRG neurons

DRGs were imaged at 3-4 days *in vitro* (DIV). Before imaging, cells were incubated

with 200 nM of MitoTracker Deep Red for 30 min and the dye washed off prior to

imaging. Mitochondrial axonal transport was recorded with a Nikon spinning disk

system comprising of a CSU-X1 scanning head (Yokogawa) and a Nikon Eclipse Ti-E

inverted microscope equipped with a Du 897 iXon Ultra electron-multiplying charge coupled device camera (Andor) and a 60× CFI Apo/1.4 NA objective. Mitochondrial movements were typically recorded for 2 min with an acquisition rate of 0.5 frames/s, 640 nm excitation laser (5-10% power), 100-150 ms exposure, gain 1, EM gain 100. Temperature and CO_2 were maintained at 37°C and 5%, respectively, via a temperature unit with CO_2 control and a microscope enclosure (Okolab). Axonal transport was recorded approximately 100 μm away from the cell body and kymographs were produced by first straightening 100 μm of axonal tract (approximately in mid axon) using the Straighten plugin and followed by the Velocity Measurement Tool in ImageJ, as described previously (Annuario et al., 2022; Mattedi et al., 2023). Mitochondrial speed, run length and number of total mitochondria were analysed directly from the kymographs. Time-lapse movies were used to unambiguously confirm the mitochondrial tracks, when necessary. A mitochondrion was defined as motile if containing at least one continuous bout of net motion of at least 2 µm (a 'run') in the anterograde or retrograde direction. Because mitochondrial traces can be composed of multiple runs of variable speeds, velocity is defined as the average run velocities per single mitochondrion. Mitochondrial displacement is defined as the sum of the mitochondrial run lengths per single mitochondrion. Bidirectional mitochondria display at least one run in both the anterograde and retrograde directions.

Mitochondrial morphology was quantified by calculating single mitochondrial surface area, length, and elongation, using the NIS Elements Analysis software. Mitochondrial length was defined as the length of the medial axis of the mitochondria while elongation was defined as ratio between maximal and minimal Feret's diameter. The

outline of each mitochondrion was obtained by manually adjusting the intensity

threshold for each movie and quantification of morphological parameters was

performed on individual representative timeframes at which mitochondria were best

visible. The same movies were utilised to quantify mitochondrial transport and

morphology.

DRG treatments and live imaging in MFCs

DRGs were imaged at 3-4 DIV. Prior to imaging, the cell body compartment was

treated with either PEG (polyethylene glycol, MW 8,000, Sigma) for 1 h or with sorbitol

(Sigma) for 30 min. PEG and sorbitol were separately diluted in DRG medium to 20%

and 300 mM, respectively, and 100 µl of each treatment were added dropwise across

the cell body wells. 150 µl of DRG medium without treatments were added across the

axonal wells. Following the initial incubation with the crowding agents, the medium in

the cell body wells was replaced with 100 µl DRG medium with treatments

supplemented with 200 nM MitoTracker Green and 1X CellMask Orange (CMO,

ThermoFisher). The medium in the axon wells was replaced with 150 µl medium

containing 1X CMO only. After 30 min, the dyes were washed off and the neurons

imaged immediately, with the treatments kept on in the cell body compartments

throughout imaging.

Mitochondrial axonal transport was recorded in the initial portion of the microgrooves

containing axonal projections using the same Nikon spinning disk system used for the

mass cultures and a 60× CFI Apo/1.4 NA objective. Mitochondrial movements were

recorded for 1 min with an acquisition rate of 0.5 frames/s, 488 nm excitation laser

(10% power), 200 ms exposure, gain 1, EM gain 100. Mitochondrial flux analysis was

performed by counting how many times in 1 min the mitochondria crossed a line drawn

in the initial segment of the microgrooves, close to the somal compartment.

Mitochondrial mass was defined as the total mitochondrial surface area in a region of

interest of 54 μm adjacent to the somal compartment and quantified as described

above for DRG cultures.

Cell body area was calculated by drawing a line around the border of the cell body, as

defined by CMO staining, using the Fiji Polygon selection tool.

Low magnification images to capture the length of the MCF microgrooves were

acquired using a 20×/0.75 NA CFI Plan Apochromat VC air objective, and the 'Large

Image' function on the NIS Acquisition software.

Live imaging and analysis of mitochondrial transport and morphology in the

mouse sciatic nerve

Exposure and *in vivo* imaging of intact sciatic nerve axons in anaesthetised mice were

performed under terminal anaesthesia as previously described (Sleigh et al., 2020;

Tosolini et al., 2021). Time-lapse imaging of mitochondrial transport was executed

using an inverted LSM780 laser-scanning microscope (ZEISS) and 63× Plan

Apochromat oil immersion objective lens (ZEISS) at 100× digital zoom. Images were

acquired at a rate of 0.5 frames/s for 8 min, with laser power 2%, resolution 1024×1024

and line averaging of 4. Analysis of mitochondrial motility was carried out essentially

as previously described (Vagnoni & Bullock, 2018). Tracking of mitochondria was

performed manually in Fiji/ImageJ on the raw movies using MtrackJ by marking the

start and end of each run and tracking was performed blind to the age and sex of the

animals. Speed and displacement were calculated from MtrackJ-generated tracks of

runs ≥ 2 μm, as defined above. Total mitochondrial number was averaged from two

frames at the beginning and at the end of the time series. The StackReg plugin of

Fiji/ImageJ was used to correct for sample drifting, if necessary.

Mitochondrial morphology was quantified as described above for DRG cultures.

Live imaging of mitochondrial membrane potential and siRNA treatment

DRGs were cultured to DIV3-4 and incubated with 100 nM MitoTracker Green FM

(Thermo Fischer Scientific) in DRG medium for 30 min at 37°C. After washing off the

dye, 50 nM TMRM (Thermo Fischer Scientific) was added to fresh medium and

incubated for 15 min. After washing off the TMRM dye with fresh medium, imaging

was performed with a Nikon A1RHD inverted confocal microscope, equipped with

GaAsP and multi-alkali PMT detectors and a 60×/1.4NA Plan Apochromat oil

immersion objective. Cells were kept at a constant temperature of 37°C, 5% CO_2

throughout imaging. For both 488 nm (MitoTracker Green) and 561 nm (TMRM)

excitation (laser power of 1% and 0.3%, respectively), we set the pixel dwell time at

2.2 μs, image size at 512×512, averaging at 2× and pinhole size at 39.6 μm. Low dye

concentration and laser power were used to prevent TMRM quenching and FRET

between dyes (Mitra & Lippincott-Schwartz, 2010). Quantification of fluorescent signal

was performed using the raw intensity values in Fiji/ImageJ. TMRM fluorescence

intensity was measured and divided by the fluorescence signal of MitoTracker Green

to obtain the normalised value of the mitochondrial membrane potential. For

presentation purposes, the 'Smooth' filter in Fiji/ImageJ was used to reduce salt and

pepper noise.

Dharmacon Accell SMARTpool Trak1 siRNA (Cat#E-056081-00-0010) and non-

targeting control siRNA (Cat#D-001910-10-05) from Horizon Discovery were

reconstituted to 100 μM following the manufacturer's instruction. When cultured DRGs

reached DIV1, siRNAs were mixed with the Accell Delivery Media (ADM, Cat #B-

005000-100) and, after removing 200 μl of conditioned medium from each well of an

8-well imaging chamber, 200 μl of siRNA-ADM complexes were delivered dropwise

on the cells at 1 μM final concentration (300 μl total medium volume/well). After 72 h

(DIV4), neurons were incubated with 4 μM of the JC-1 dye (Thermo Fisher Scientific)

for 30 min and the medium was replaced before imaging. The aggregation state of the

dye is affected by $\Delta\Psi_m$ with the fluorescence emission shifting from green to red in

more polarised mitochondria. The fluorescence emission at ~ 590 nm (red) and ~ 530

nm (green) was captured with emission filters ET595/50m and ET525/50m (both from

Chroma Technology Corporation), respectively. The ~ 590/530 fluorescence ratio

gives a measurement of $\Delta\Psi_m$. For trafficking studies using MitoTracker DeepRed, 0.5

μl non-targeting Green/6-FAM siRNA (Cat#D-001950-01-05) was added to the siRNA-

ADM complexes to assess the efficiency of siRNA delivery.

Live cell calcium imaging

DRGs were imaged with a Nikon Ti-E inverted epifluorescence microscope equipped

with a Nikon 20×/0.8 NA Plan Apochromat air objective, a mercury lamp (Nikon

Intensilight C-HGFI) for illumination and a Dual Andor Neo sCMOS camera for

detection. Cells were plated on laminin-coated 18 mm coverslips and treated with 2

μm Rhod2-AM and 4 μm Fluo4-AM (both from Thermo Fisher Scientific) for 15 min at

37°C in external solution (145 mM NaCl, 2 mM KCl, 5 mM NaHCO$_3$, 1 mM MgCl$_2$, 2.5

mM CaCl$_2$, 10 mM glucose, 10 mM HEPES pH 7.4) (De vos et al., 2012) prior to

imaging. Cells were imaged at DIV4 in a Ludin imaging chamber (Type 1, Life Imaging

Services) in external solution at 1 frames/s for 3 min using GFP/mCherry filter set and

exposure time of 30 ms. Cells were imaged for 30 s before transiently perfusing the

imaging chamber with 50 mM KCl with a peristaltic pump (Ismatec). KCl was washed

off after 1 min and each coverslip was subjected to only one round of perfusion and

imaging.

To quantify KCl-induced Fluo4 and Rhod2 response, a region of interest (ROI) was

drawn to outline single cell bodies and axonal tracts from each field of view and the

fluorescence intensity of the dyes measured in Fiji/ImageJ from raw images displaying

non-saturated fluorescent intensity. Response curves were aligned at the base of the

response peak and the fluorescence intensity at every time point was normalised to

the value of the first frame. Fluorescence 'peak' was defined as the maximum fold-

change value reached during imaging and the time to reach the peak was calculated

as the Δt between the 'peak' timepoint and the time of first response to stimulation, as

reported previously (Mattedi et al., 2023). Time to decay was defined as the time

necessary for the fluorescence signal to decline by 80% of the peak value.

GEMs viral preparation and transduction

To obtain GEMs-encoding lentiviral particles, the pCMV-PfV-Sapphire-IRES-

tdTomato plasmid (Addgene, #116934) (Delarue et al., 2018) was transfected into

Lenti-X 293T cells (TaKaRa, #632180) using the polyethylenimine(PEI)-based

Transporter 5 transfection reagent (Polysciences, #26008) and a 2nd generation viral

packaging system comprising gag/pol/tat/rev (psPAX2, Addgene, #12260) and VSV-

G (pMD2.G, Addgene, #12259). Cells were seeded in a 15 cm dish in 22.5 ml medium

(DMEM with 4.5g/l glucose and pyruvate, 10% hi-FBS, 1x GlutaMAX) at 37°C with 5%

CO_2 until they reached 60-80% confluency. The medium was then replaced and cells

incubated for 2 h with fresh medium containing 2% FBS. DNA-PEI complexes (14.6

μg psPAX2, 7.9 μg pMD2.G, 22.5 μg GEM-transfer vector and 135 μl Transporter 5)

were prepared according to the manufacturer's instructions and added dropwise to the

cells. After overnight incubation, the medium was replaced with 17 ml fresh medium

with 2% FBS, the supernatant harvested 3 times every 8-12 h and kept at 4°C.

Typically, the three harvests were centrifuged at 500 g for 5 min at 4°C before filtering

through 0.22 μm filter units and further ultracentrifugation at 50,000 x g for 2 h at 4°C.

After gently discarding the supernatant, the viral pellet was let dry for 2-3 min before

resuspension in DMEM without FBS or Opti-MEM to obtain a 1000-fold concentration

and stored at -80°C in 10-20 μl aliquots. Between 1-10 μl of concentrated virus was

added to each well of a μ-Slide 8-well glass bottom imaging chambers.

GEMs imaging and single particle analysis

DRGs were plated in μ-Slide 8-well glass bottom imaging chambers and the

concentrated lentiviral particles added at DIV4. The medium was replaced after 24 h

with fresh medium without growth factors and the cells imaged after an additional 36

h (DIV7) by spinning disk confocal microscopy with a Nikon 100x/NA 1.40 CFI Plan

Apochromat VC oil immersion objective and 100 ms continuous acquisition for 30 s,

488 nm laser at 40-50% intensity, gain 1, EM gain 250. Cells expressing similar levels

of transduced construct, as assessed by the tdTomato signal, were selected for

imaging.

Single particle tracking (SPT) was performed on raw time series using TrackMate on

selected ROIs defining the cell body and axonal compartments using the DoG detector

for small spot detection (<5 pixels), with estimated particle diameter set at 0.4-0.5,

thresholds between 0.4-0.6. with median filter and sub-pixel localization. The Simple

LAP tracker was used to link the particles throughout the movie (linking max distance

= 1, gap-closing max distance = 1.6, gap-closing max frame gap = 2). Tracks were

then exported and analysed in MATLAB. Time-averaged mean square displacement

(MSD) was calculated via creation of a SPT Analyser using a custom script (available

on request). The mean diffusion coefficient (D_{eff}) was calculated by fitting the first 10

s of the time-averaged MSD with the equation $MSD\ (t) = 4Dt^{\alpha}$, where α is the exponent

describing the anomalous diffusion and D is the diffusion coefficient.

FLIM

Calibration and FLIM measurements were carried out essentially as in Steinmark et

al. (Steinmark et al., 2019). DRGs were plated in μ-Slide 8-well glass bottom imaging

chambers. On the day of imaging (DIV3-4), cells were washed in external solution and

incubated with 1 μM FMR-1 for 30 min at 37°C and the dye was washed off prior to

imaging. Cells were imaged on an inverted TCS-SP2 microscope (Leica

Microsystems) using two SPC-150 TCSPC boards (Becker & Hickl). FMR-1 was

excited using a picosecond-pulsed (90 ps optical pulse width, 20 MHz repetition rate)

467 nm diode laser (Hamamatsu, Japan) using a HCX PL APO CS 63X 1.2 NA water

objective, with a 485 nm dichroic mirror and a 514/30 nm band pass filter. All

measurements were carried out at 37°C and 5% CO_2. Analyses were performed at

single cell level by measuring the fluorescence lifetime separately in cell bodies and

proximal-mid axons regions.

TR-FAIM

DRG neurons were grown to DIV3-4, as for the FLIM experiments, and labelled with

1 µM BCECF (Themo Fisher Scientific #B1150) for 30 min at 37°C. The dye was

washed off prior to imaging in FluoroBrite DMEM. Time-resolved anisotropy

measurements were conducted on the same TCS-SP2 microscope system used for

FLIM, with the BCECF dye excited using a picosecond-pulsed laser. In this case, the

fluorescence emission was split into its parallel and perpendicular components using

a polarizing beam splitter (Edmund Optics).

Calibration curve

Rotational correlation times of BCECF in solutions of increasing viscosity (through

addition of glycerol) were measured and plotted. To avoid depolarisation effects from

the objective, a 5x NA 0.15 objective was used and measurements were conducted at

room temperature, but otherwise the setup was identical to that described above. The

anisotropy was calculated using the equation:

821
$$r(t) = \frac{I(t)_{par} - GI(t)_{perp}}{I(t)_{par} + zGI(t)_{perp}} \ (1)$$

822 Where z = 2 and G is a correction factor used to correct for the different sensitivity of

823 the two detectors. It is calculated by taking the tail value of:

824
$$G = I(t)_{par}/I(t)_{perp} \ (2)$$

G-values used were around 1 +/- 0.1. Here the G-factor was measured using a solution

of fluorescein in water and 10% glycerol. The calibration curve was calculated to avoid

assumptions about molecular shape and diffusion boundary conditions in a classic

Stokes-Einstein-Debye calculation.

Fluorescence Anisotropy decay

For each raw neuron image, two regions of interests (ROIs) were extracted: from the

axon and the soma. For each ROI, a single set of parallel and perpendicular decays

were generated by summing the photon counts in each time bin over all pixels within

the given ROI. Anisotropy pre-processing, curve fitting and graphing were performed

in OriginLab. The perpendicular (I_{perp}) and parallel (I_{par}) decays for each ROI of each

image were aligned and the anisotropy decay was calculated using equation (1) where

z, due to the high NA of the objective, was determined to be 1, and G was determined

as described above. Analyses were performed at single cell level by calculating

anisotropy separately on cell bodies and proximal-mid axon region of the same cell.

For calculating the anisotropy of an entire cell, the soma and axon photon counts were

summed up and analysis was performed as described. It was established that the

hindered rotation model (below) provides an optimum fit for most anisotropy decays,

accounting for a restricted model of rotation in all environments.

$$r_{model} = (r_0 - r_\infty)e^{\frac{-t}{\theta}} + r_\infty \ (3)$$

Where r_0 is the initial anisotropy, r_∞ is the limiting anisotropy and θ is the rotational

correlation time. The weighting factor v(r) for the anisotropy fits was the variance of r

(Lidke et al., 2005):

$$v(r) = \frac{(1-r)(1+2r)(1-r+G(1+2r))}{3G} \ (4)$$

Only the first 10 ns of the decay were used for the fit and only meaningful fits with an

acceptable adjusted R-squared value and standard errors of the parameter fits were

accepted for further analysis.

Statistical analyses

Details of the statistical analyses are reported in the figure legends. Shapiro-Wilk test

was performed to test data for normal distribution before determining the appropriate

statistical test. Sample sizes, which were predetermined using power calculations and

previous experience, are also reported in figure legends.

FIGURES

Figure 1

Figure 1. Mitochondrial transport is reduced in aged DRG neurons in culture

(A) DRGs were dissected from the spinal cord of 6 young (P72-107, 3 females and 3

males) and 6 old (P560-637, 3 females and 3 males) mice and dissociated neurons

were cultured *in vitro.* **(B)** Representative kymographs of mitochondria moving in the

axons of DRG neurons dissected from young and old animals. **(C-D)** The proportion

of motile mitochondria (C), but not the mitochondrial mass (D), is strongly reduced

during ageing in the DRG neuron axons. Individual data points representing the mice

used (bigger circles with line depicting the median) are overlayed on data points

(smaller circles) representing individual fields of view (FOV) imaged (1-3 axons/FOV).

(E-F) Representative frames of MitoTracker-stained mitochondria in the axons of cells

extracted from young (E) and old animals (F). Bottom panels show the thresholded

mitochondrial area (magenta) used for quantitative analysis of morphology. Scale bar:

5 μm. **(G)** Quantification of mitochondrial area in young and old cells. Individual data

points represent the fields of view (FOV) imaged (1-3 axons/FOV) and are shown as

mean ± SEM. **(H-I)** Violin plot distributions show that anterograde (H) and retrograde

(I) mitochondrial velocity and displacement remain largely stable during ageing, albeit

with a subtle but significant reduction in anterograde displacement in aged neurons.

In brackets, number of mitochondria. Unpaired student's t-test (C-D), Mann-Whitney

test (G-I). * $p < 0.05$; ** $p < 0.01$.

Figure 2

Figure 2. Mitochondrial transport decreases with age in the sciatic nerve *in vivo*

(A) 5 young (P72-107, 3 females and 2 males) and 5 old (P560-637, 2 females and 3

males) MitoMice were anaesthetised and the sciatic nerve exposed for time-lapse

imaging of mitochondrial axonal transport. **(B-C)** Still frames from movies showing

CFP-marked mitochondria (grey, top panels) in single sciatic nerve axons from young

(B) and old (C) MitoMice. The bottom panels depict the traces of mitochondria moving

in the anterograde and retrograde directions (cyan and magenta, respectively) in the

corresponding movies. Scale bars: 5 μm. **(D-E)** Both the number of motile

mitochondria and the total number of mitochondria present in the axons of the sciatic

nerve are strongly reduced during ageing. Individual data points representing the mice

used (bigger circles with line depicting median) are overlayed on data points (smaller

circles) representing individual sciatic nerve axons. **(F)** Quantification of mitochondrial

area in young and old sciatic nerves. Individual data points represent the sciatic nerve

axons and are shown as mean ± SEM. **(G-H)** Violin plot distributions show that

mitochondrial anterograde velocity and displacement are both upregulated in old

neurons (G), while the motile properties of the retrograde-moving mitochondria are not

affected by age (H). In brackets, number of mitochondria. Mann-Whitney test (D-H). *

$p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Figure 3

Figure 3. Mitochondrial membrane potential and viscosity decline during ageing

(A-D) Representative images of young (A-B) and aged (C-D) DRG neuron cell bodies

(top panels) and axons (bottom panels), stained with TMRM and MitoTracker Green

(MTG). Scale bars: 10 μm. **(E-G)** Quantification of the ratio between TMRM and MTG

in neuronal cell bodies and axons combined indicates that the $\Delta\Psi_m$ declines during

ageing at cellular level (E) with axonal mitochondria displaying consistently reduced

$\Delta\Psi_m$ in both young (F) and old (G) cells. **(H)** Representative images of a DRG neuron

cell body (top panel) and axon (bottom panel) stained with FMR-1 to capture

mitochondrial viscosity. Rainbow colour scale indicates fluorescence lifetime range

(2,000-5,500 ps) in which orange/red and cyan/blue hues correspond to areas of lower

and higher viscosity, respectively. Scale bars: 15 μm. **(I)** Overall mitochondrial

viscosity is decreased in old cells, as indicated by quantification in neuronal cell bodies

and axons combined. **(J-K)** There is no significant difference between somal and

axonal mitochondrial viscosity in young DRG neurons (J), while mitochondrial viscosity

is slightly higher in old DRG axons (K). In E-G and I-K, data points represent the cell

bodies and axons imaged. Each neuron analysed provides both cell body and axons

to the analysis. DRGs were dissected from 3 young (P49-67, 1 female and 2 males)

and 3 old (P672-679, 2 females and 1 male) mice for TMRM experiments (A-G), and

from 3 young (P81-84, 2 females and 1 male) and 2 old (P715, 1 female and 1 male)

mice for FLIM experiments (H-K). Data are shown as mean ± SEM. Mann-Whitney

test (E, I), paired student's t-test (F, K), Wilcoxon test (G, J). * $p < 0.05$; ** $p < 0.01$;

**** $p < 0.0001$.

Figure 4

Figure 4. The cytoplasmic viscosity of aged DRG neurons in culture increases

in cell bodies but not in axons

(A) Schematic of the fluorescence anisotropy optical set up. A 467 nm depolarised

pulsed laser beam (cyan) is linearly polarised before exciting BCECF molecules with

a transition dipole (dashed red line) parallel to the electric field vector of the light. The

510 nm emitted light (green) is passed into a polarising beam splitter to two detectors

for light with either parallel (I∥) or perpendicular (I⊥) emission polarisation. The

rotational correlation time (θ) of the BCECF molecule is defined as the time taken to

rotate one radian (during the fluorescence lifetime). **(B)** Fluorescence anisotropy

decay and fit (see Methods, equation 3) of BCECF fluorescence obtained from

perpendicular and parallel decays (according to equation 1, see Methods) from young

and old DRG neurons. Older cells display a slower decay, indicative of higher

cytoplasmic viscosity. Red lines show the exponential decay fit and circles the

individual time points. **(C)** Viscosity values obtained from anisotropy decay analysis

show increased viscosity in old DRG neurons. Circles show individual cellular

compartments. Relative to (B). **(D-E)** The somal and axonal compartments display

similar viscosity values in young DRG neurons (D, $p = 0.5065$), whereas the

cytoplasmic viscosity of old DRG neurons is significantly higher in the somal

compartment. Circles depict individual cellular compartments. **(F)** Rotational

correlation time (θ) values relative to somal and axonal anisotropy data. Note the

average θ is proportional to viscosity. Data are shown as mean ± SEM, Mann-Whitney

test. * $p < 0.05$; ** $p < 0.01$. DRGs for the anisotropy experiments were obtained from

2 young (P80) and 3 old (P709-712) male mice.

 Figure 5

Figure 5. The diffusiveness of the neuronal cytoplasm decreases in aged cultured neurons

(A-F) Representative images of a cell body (A-C) and axon (D-F) of a DRG neuron transduced with lentiviral vectors expressing GEMs. In (B) and (E), single particle tracking (SPT) markers are overlayed onto the cell body and axons, respectively. (C) and (F) show the colour-coded GEMs trajectories tracked during 30 s of imaging of (B) and (E), respectively. Scale bars: 5 μm. **(G)** Mean square displacement (MSD) of GEM tracks in cell bodies and axons of young and old cells. Data are plotted as mean ± SEM, Kolmogorov-Smirnov test. The motion of GEMs is captured by a sub-diffusion

964 model described by the equation $MSD\ (t) = 4Dt^{\alpha}$ (young cell bodies: R = 0.999, old

965 cell bodies: R = 1, young axons: R = 0.997, old axons: R = 1). In brackets, number of

966 tracks analysed from 61 cell bodies and 170 axons. **(H)** Cumulative distribution

967 function (CDF) of GEMs diffusion coefficient (D_{eff}) in young and old DRG axons and

968 cell bodies, relative to (G). **(I)** Mean diffusion coefficient in the cell body and axonal

969 compartments of young and old cells. Relative to (H). DRGs for the microrheology

970 experiments were obtained from 2 young (P49-70) male mice and 2 old (P726-757, 1

971 female and 1 male) mice. Data are presented as mean ± SEM, Kruskal-Wallis with

972 Dunn's multiple comparisons test. * $p < 0.05$; **** $p < 0.0001$.

Figure 6

Figure 6. Increasing crowding and viscosity in DRG cell bodies is sufficient to

reduce mitochondrial axonal transport

(A) Schematic depicting the microfluidic chamber (MFC) used and the treatments

specifically applied to the somal and axonal compartments. **(B)** Timeline of the

experimental setup, starting with sorbitol or PEG treatment, followed by a 'pulse'

consisting of staining with live dyes, and a 'chase' (i.e. microscopy observation). **(C-**

E) Representative images of MFCs in which the somal compartment was either left

untreated (C) or treated with sorbitol (D) or PEG (E). Top panels: MFC microgrooves

(Microgr.) show the extent of transport from the soma of MitoTracker (MTG)-stained

mitochondria during the 'pulse' period. Bottom panels: correspondent axonal

compartment (Ax. comp.) stained with CellMask (CMask). Scale bar: 100 μm. **(F)**

Representative higher magnification of an area of the microgrooves typically used for

quantification of mitochondrial motility, indicated by a white rectangle in (C). Scale bar:

5 μm. **(G)** Quantification of mitochondrial transport fluxes (see Methods) shows strong

decline in sorbitol and PEG-treated DRGs. Datapoints in the violin plots represent

individual microgrooves with each colour indicating a different animal (4 females, P45-

86), Kruskal-Wallis with Dunn's multiple comparisons test. **(H)** Combined axonal

mitochondrial velocities are reduced in sorbitol-treated neurons. Due to the very low

number of mitochondrial runs in PEG-treated axons, this sample has not been used

for statistical analysis. ** $p < 0.01$; **** $p < 0.0001$.

Figure 7

Figure 7. Model of mitochondrial homeostasis and cytoplasmic viscosity during ageing

Decreased axonal mitochondrial transport characterises the ageing process of mouse DRG neurons *in vivo* and *in vitro* ('Age-dependent', yellow). The overall decrease in mitochondrial membrane potential observed in aged DRG cultured neurons correlates with reduced intra-mitochondrial viscosity both in the cell bodies and axons ('Age-dependent', green) but not with their capacity to buffer calcium, which is unaffected in aged cultures ('Age-independent', green). Cytoplasmic viscosity is upregulated in the cell bodies ('Age-dependent', blue), but not in axons ('Age-independent', yellow). Compartment-dependent and age-specific changes may reflect functional specialisation of the mitochondria and specific regulation of viscosity across sub-cellular compartments and time.

1012 **Supplementary Figure 1**

1013

1014

Figure S1

Proportion of motile mitochondria displaying anterograde, retrograde and bidirectional

motility in young **(A)** and old **(B)** DRG neurons *in vitro* (relative to Figure 1C). A general

decrease of motility is observed during age. **(C-D)** Quantification of mitochondrial

length and elongation (see Methods) in young and old cells. Individual data points

represent fields of view (FOV) imaged (1-3 axons/FOV). Data are shown as mean ±

SEM. Kruskal-Wallis with Dunn's multiple comparisons test (A-B) and Mann-Whitney

test (C-D). * p < 0.05; ** p < 0.01; **** p < 0.0001.

1023

1024 **Supplementary Figure 2**

1025

1026

Figure S2

(A-B) Quantification of mitochondrial length and elongation (see Methods) in young

and old sciatic nerve axons *in vivo*. Individual data points represent sciatic nerve

axons. **(C)** There is no difference in the mitochondrial motility index between young

and old neurons (*i.e.*, total moving mitochondria/(total moving + stationary

mitochondria)). **(D)** The relative proportions of anterograde and retrograde

mitochondrial transport are not changed by age in the sciatic nerve, indicating that the

reduction of both types of transport contributes to the overall transport decline (Figure

2D). Individual data points represent fields of view (FOV) imaged (1-3 axons/FOV).

Data are shown as mean ± SEM. Mann-Whitney test (A-B), unpaired student's t-test

(C), two-way ANOVA with Tukey's multiple comparisons test (D). ** $p < 0.01$; **** $p <$

0.0001.

Supplementary Figure 3

Figure S3

 (A-B) Quantification of the ratio between TMRM and MTG in neuronal cell bodies (A)

 and axons (B) indicates that the $\Delta\Psi_m$ declines during ageing in both compartments.

 (C) DRG neurons stained with the JC-1 vital dye and pseudocoloured in cyan and

magenta (~ 530 and ~ 590 nm emission, respectively). Arrows, somal compartments;

arrowheads, axons. Note the higher polarisation of somal mitochondria compared to

axonal mitochondria, indicated by widespread magenta, corroborating the data in Fig.

3A-G. Scale bar: 20 μm. **(D)** Quantification of the 590/530 JC-1 ratio from young DRG

neurons indicates that the axonal mitochondria display reduced $\Delta\Psi_m$ compared with

cell body mitochondria. Data points represent the cell bodies and axons imaged, with

each neuron providing both cell body and axons for paired analysis. **(E)** Increased

number of DRG neuron axons displaying no motile mitochondria after Trak1 RNAi. A

process is scored as displaying motility if at least two mitochondria are motile within 1

min of imaging. Number of axons analysed are 58 (Ctrl siRNA) and 46 (Trak1 siRNA).

(F-G) Trak1 RNAi does not significantly change the 590/530 JC-1 ratio in the cell

bodies (F) or axons (G) of DRG neurons, suggesting the $\Delta\Psi_m$ is not affected by this

manipulation. Data points represent individual cell bodies and axons, respectively.

Each neuron analysed provides both cell body and axons to the analysis. DRGs for

JC-1 quantifications were obtained from 3 young (P102-105) male mice. **(H-I)**

Mitochondrial viscosity decreases during age both in the cell bodies (H) and axons (I)

of DRG neurons. Data points represent cell bodies and axons imaged. Data are shown

as mean ± SEM. Unpaired student's t-test (A, F, H), Mann-Whitney test (B, G, I), paired

student's t-test (D), Fisher's exact test (E). * $p < 0.05$; ** $p < 0.01$; **** $p < 0.0001$.

1065

Supplementary Figure 4

1067

Figure S4. Mitochondrial and cytosolic calcium uptake are not significantly

affected by age in cultured DRG neurons

(A-H) Representative images of DRGs cultured before (A,C,E,G) and after (B,D,F,H)

stimulation with 50 mM KCl. Cells are co-stained with Rhod2-AM (magenta) and

Fluo4-AM (cyan) for measuring the fluorescent signal of mitochondrial and cytosolic

Ca^{2+}, respectively. Neuronal depolarisation by KCl increases calcium uptake in cell

bodies (arrows) and axons (arrowheads). Panels A-D are shown with enhanced

brightness in E-H to highlight the response after treatment. Scale bars: 20 μm. **(I-J)**

Traces indicate the average Rhod2-AM (black, circles) and Fluo4-AM (red, triangles)

fluorescence intensity values in 18 young (I) and 20 old (J) neurons at individual time

points (circles and triangles, respectively) normalised to the average fluorescence

value before KCl stimulation (time 0). **(K-L)** Time to reach the peak, **(M-N)** normalised

response peak and **(O,P)** decay time for cytoplasmic (Fluo4-AM) and mitochondrial

(Rhod2-AM) calcium. Mann-Whitney tests (K,M-P) and an unpaired student's t-test

with Welch's correction (L) showed no difference between conditions. DRGs were

obtained from 3 young (P58-93, 3 males) and 4 old (P560-676, 1 female, 3 males)

mice.

1085

 Supplementary Figure 5

 Figure S5

 (A) Calibration curve of BCECF rotational correlation time (θ) calculated in accordance

 with the Stokes-Einstein-Debye equation. The experimental data were compared to

 historical data for BCECF θ as a function of viscosity (Verkman et al., 1991) and

 showed good agreement. **(B-C)** Fluorescence anisotropy decay and fit (see Methods,

 equation 3) of BCECF fluorescence obtained from perpendicular and parallel decays

 (according to equation 1, see Methods) from cell bodies and axons of young (B) and

 old (C) DRG neurons. Red and black lines depict exponential decay fit for cell bodies

 and axons, respectively. The anisotropy decay of young cells (B) is not significantly

 different between compartments, leading to similar viscosity values (Fig. 4D). The

 slower decay of cell bodies compared to axons in old DRGs is indicative of higher

1099   viscosity, as shown in Fig. 4E. Circles, individual time points. **(D-E)** Cytoplasmic

1100   viscosity increases during age in the cell bodies (D) but not axons (E) of DRG neurons.

1101   Circles depict cell bodies and axons imaged. Data are shown as mean ± SEM. Mann-

1102   Whitney test (D-E). ** $p < 0.01$.

1103

 Supplementary Figure 6

 Figure S6

 (A) Combined MSD of GEM tracks in cell bodies (CB, blues) and axons (Ax, orange).

 Data are plotted as mean ± SEM in a log-log format. Dashed lines are curve fit with

 anomalous exponent of 0.910 ± 0.011 (CB) and 0.724 ± 0.234 (Ax), respectively. **(B-**

 C) Positive correlation between total number of GEMs tracks analysed and axonal (B)

 and cell body (C) areas, suggesting that GEMs equally cover the full extent of the

 neuronal cytoplasm, independently of cell size. Circles represent individual axons (B)

 and cell bodies (C). Red lines, linear regression fit. Spearman rho test. **(D)** There is

 no significant difference between the mean area of the young and old cell bodies,

1116   indicating that cell size is unlikely to affect viscosity and diffusiveness. Circles

1117   represent individual cell bodies. Data are shown as mean ± SEM. Mann-Whitney test

1118   (p = 0.5148).

1119   **Supplementary Figure 7**

1120

Figure S7

A) Microfluidic chamber (MFC) showing reliable fluidic separation between the cell

body and axonal compartments. A blue dye was applied to the somal compartment at

a lower volume than the connecting axonal compartment and incubated at 37°C for

up to 5 h. Visual inspection confirmed that there is no dye leakage against the volume

gradient. The same outcome was achieved after staining the axonal compartment. **(B)**

Quantification of mitochondrial mass (see Methods) shows strong decline in sorbitol

and PEG-treated DRGs. Datapoints in the violin plots represent individual

microgrooves with each colour indicating a different animal (4 females, P45-86),

Kruskal-Wallis with Dunn's multiple comparisons test. **(C)** Representative images of

DRG cell bodies that have been left untreated (top panels) or treated with sorbitol

(middle panels) or PEG (bottom panels). The outline of the cell body defined by

CellMask staining was used as a reference for the quantification of cell body area in

(D). Scale bar: 5 μm. **(D)** Quantification of the DRG cell body area (see Methods)

shows no significant change across conditions, unpaired t-tests with Welch's

correction (untreated vs sorbitol) and Mann-Whitney test (untreated vs PEG).

Datapoints in the violin plots represent cell bodies from different MFCs, with each

colour indicating a different animal (3-4 females, P45-86). *** $p < 0.001$.

1139

Supplementary Table 1

A. Young

Velocity (μm/s)			
Anterograde		Retrograde	
In vivo	*In vitro*	*In vivo*	*In vitro*
0.63 (2.55), n = 1018	0.45 (2.94), n = 236 ****	0.58 (2.01), n = 428	0.44 (2.98), n = 141 ****
Displacement (μm)			
Anterograde		Retrograde	
In vivo	*In vitro*	*In vivo*	*In vitro*
11.86 (64.92), n = 1018	11.73 (83.80), n = 236	11.95 (64.26), n = 428	9.02 (72.47), n = 141 **

B. Old

Velocity (μm/s)			
Anterograde		Retrograde	
In vivo	*In vitro*	*In vivo*	*In vitro*
0.69 (1.7), n = 507	0.51 (4.41), n = 183 ****	0.67 (2.17), n = 233	0.43 (2.85), n = 137 ****
Displacement (μm)			
Anterograde		Retrograde	
In vivo	*In vitro*	*In vivo*	*In vitro*
13.31 (65), n = 507	8.62 (95.13), n = 183 ****	13.58 (65), n = 233	7.21 (85.50), n = 137 ****

Table S1

Median mitochondrial velocities and displacements are significantly higher in the sciatic nerve axons (*in vivo*) compared to DRG axons (*in vitro*) in all conditions except for anterograde displacement in young samples (A) where no difference was detected. Values are medians with, in brackets, maximum speeds and maximum displacements. n = number of mitochondria, Mann-Whitney test. ** $p < 0.01$; **** $p < 0.0001$. The data suggest that, while mitochondria are able to reach faster maximum velocities *in vitro*, on average they move more slowly *in vitro* than *in vivo*. Note that the displacement is calculated over 65 μm and 100 μm of axonal tract in the sciatic nerve and cultured DRGs, respectively, suggesting the calculated values may underestimate the significance between *in vivo* and *in vitro* displacements.

1156 **Movie 1**

1157 Representative time-lapse movie of mitochondria in axons of DRG neurons obtained

1158 from young (P92) and old (P676) mice and stained with MitoTracker Deep Red. Width:

1159 73 μm. Movie plays at 20 frames per seconds and represents 2 min of real-time. Scale

1160 bar: 5 μm.

1161

1162 **Movie 2**

1163 Representative time-lapse movie of mitochondria from young (P72) and old (P624)

1164 MitoMouse sciatic nerve axons. Width: 64 μm. Movie plays at 20 frames per seconds

1165 and represents 4:26 min of real-time. The 'Smooth' filter in Fiji/ImageJ was applied for

1166 presentation purposes. Scale bar: 5 μm.

1167

1168 **Movie 3**

1169 Representative time-lapse movie of GEM particles in the cell body of a cultured DRG

1170 neuron. Movie plays at 20 frames per seconds and represents 30 seconds of real-

1171 time.

1172

1173 **Movie 4**

1174 Representative time-lapse movie of GEM particles in the axon of a cultured DRG

1175 neuron. Width: 55 μm. Movie plays at 20 frames per seconds and represents 30

1176 seconds of real-time. The 'Smooth' filter in Fiji/ImageJ was applied for presentation

1177 purposes.

1178

Movie 5

Representative time-lapse movie of mitochondrial transport in MFCs acquired from

axons of neurons in which the somal compartment was either left untreated (top

movie) or was treated with sorbitol (middle movie) or PEG (bottom movie). Width: 80

µm. Movie plays at 20 frames per seconds and represents 1 min of real-time. The

'Smooth' filter in Fiji/ImageJ was applied for presentation purposes.

1185

Movie 6

Representative time-lapse movie of mitochondrial dynamics in MFCs acquired from

neuronal cell bodies. Top movie: untreated control; middle movie: cell body treated

with sorbitol; bottom movie: cell body treated with PEG. Movie plays at 20 frames per

seconds and represents 1 min of real-time. In the cell bodies treated with crowding

agents, the mitochondria appear more clustered and less dynamic.

1192

AUTHOR CONTRIBUTIONS

JNS, FM, SR, EA, KN, IES, AV performed experiments, collected and analysed data; II, ILD, PPJ, SA analysed and interpreted data; ERR, JCM critically supported the experimental work; GY generated and provided reagents; JNS, KS, GS, AV supervised the work; JNS, KS, GS, AV were responsible for funding acquisition. AV wrote the manuscript with critical inputs from JNS, KS, GS. AV conceived and led the study. All the authors read and approved the manuscript.

DATA AVAILABILITY STATEMENT

All relevant data that support the findings are within the paper and its supporting information files. All raw data are available from the corresponding author upon request.

BIBLIOGRAPHY

Alberti, S., & Hyman, A. A. (2016). Insights & Perspectives Are aberrant phase transitions a driver of cellular aging? *BioEssays*, *38*(10), 959-968. https://doi.org/10.1002/bies.201600042

Annuario, E., Ng, K., & Vagnoni, A. (2022). High-Resolution Imaging of Mitochondria and Mitochondrial Nucleoids in Differentiated SH-SY5Y Cells. *Methods in molecular biology (Clifton, N.J.)*, *2431*, 291-310. https://doi.org/10.1007/978-1-0716-1990-2_15

Baranov, S. V., Baranova, O. V., Yablonska, S., Suofu, Y., Vazquez, A. L., Kozai, T. D. Y., Tracy Cui, X., Ferrando, L. M., Larkin, T. M., Tyurina, Y. Y., Kagan, V. E., Carlisle, D. L., Kristal, B. S., & Friedlander, R. M. (2019). Mitochondria modulate programmed neuritic retraction. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(2), 650-659. https://doi.org/10.1073/pnas.1811021116

Berth, S. H., & Lloyd, T. E. (2023). Disruption of axonal transport in neurodegeneration. *The Journal of Clinical Investigation*, *133*(11), e168554-e168554. https://doi.org/10.1172/JCI168554

Brini, M., Pinton, P., King, M. P., Davidson, M., Schon, E. A., & Rizzuto, R. (1999). A calcium signaling defect in the pathogenesis of a mitochondrial DNA inherited oxidative phosphorylation deficiency. *Nature Medicine*, *5*(8), 951-954. https://doi.org/10.1038/11396

Bulthuis, E. P., Dieteren, C. E. J., Bergmans, J., Berkhout, J., Wagenaars, J. A., van de Westerlo, E. M. A., Podhumljak, E., Hink, M. A., Hesp, L. F. B., Rosa, H. S.,

1229   Malik, A. N., Lindert, M. K.-T., Willems, P. H. G. M., Gardeniers, H. J. G. E.,
1230   den Otter, W. K., Adjobo-Hermans, M. J. W., & Koopman, W. J. H. (2023).
1231   Stress-dependent macromolecular crowding in the mitochondrial matrix. *The*
1232   *EMBO Journal*, *42*(7), e108533-e108533.
1233   https://doi.org/10.15252/embj.2021108533

1234 Caponio, D., Veverová, K., Zhang, S.-q., Shi, L., Wong, G., Vyhnalek, M., & Fang, E.
1235   F. (2022). Compromised autophagy and mitophagy in brain ageing and
1236   Alzheimer's diseases. *Aging Brain*, *2*, 100056-100056.
1237   https://doi.org/https://doi.org/10.1016/j.nbas.2022.100056

1238 Chambers, J. E., Huber, R. G., Avezov, E., Bond, P. J., Marciniak, S. J., & Kuimova,
1239   M. K. (2018). An Optical Technique for Mapping Microviscosity Dynamics in
1240   Cellular Organelles. *ACS Nano*, *12*(5), 4398-4407.
1241   https://doi.org/10.1021/acsnano.8b00177

1242 Chang, K. T., Niescier, R. F., & Min, K.-T. (2011). Mitochondrial matrix Ca2+ as an
1243   intrinsic signal regulating mitochondrial motility in axons. *Proceedings of the*
1244   *National Academy of Sciences of the United States of America*, *108*(37),
1245   15456-15461. https://doi.org/10.1073/pnas.1106862108

1246 Chen, Q., Fang, H., Shao, X., Tian, Z., Geng, S., Zhang, Y., Fan, H., Xiang, P., Zhang,
1247   J., Tian, X., Zhang, K., He, W., Guo, Z., & Diao, J. (2020). A dual-labeling probe
1248   to track functional mitochondria–lysosome interactions in live cells. *Nature*
1249   *Communications 2020 11:1*, *11*(1), 6290-6290. https://doi.org/10.1038/s41467-
1250   020-20067-6

1251 Clayton, A. H. A., Hanley, Q. S., Arndt-Jovin, D. J., Subramaniam, V., & Jovin, T. M.
1252   (2002). Dynamic fluorescence anisotropy imaging microscopy in the frequency
1253   domain (rFLIM). *Biophysical Journal*, *83*(3), 1631-1649.
1254   https://doi.org/10.1016/S0006-3495(02)73932-5

1255 De vos, K. J., Mórotz, G. M., Stoica, R., Tudor, E. L., Lau, K. F., Ackerley, S., Warley,
1256   A., Shaw, C. E., & Miller, C. C. J. (2012). VAPB interacts with the mitochondrial
1257   protein PTPIP51 to regulate calcium homeostasis. *Human Molecular Genetics*,
1258   *21*(6), 1299-1311. https://doi.org/10.1093/HMG/DDR559

1259 Delarue, M., Brittingham, G. P., Pfeffer, S., Surovtsev, I. V., Pinglay, S., Kennedy, K.
1260   J., Schaffer, M., Gutierrez, J. I., Sang, D., Poterewicz, G., Chung, J. K., Plitzko,
1261   J. M., Groves, J. T., Jacobs-Wagner, C., Engel, B. D., & Holt, L. J. (2018).
1262   mTORC1 Controls Phase Separation and the Biophysical Properties of the
1263   Cytoplasm by Tuning Crowding. *Cell*, *174*(2), 338-349.e320.
1264   https://doi.org/10.1016/j.cell.2018.05.042

1265 Dix, J. A., & Verkman, A. S. (1990). Mapping of fluorescence anisotropy in living cells
1266   by ratio imaging. Application to cytoplasmic viscosity. *Biophysical Journal*,
1267   *57*(2), 231-240. https://doi.org/10.1016/S0006-3495(90)82526-1

1268 Duchen, M. R. (2000). Mitochondria and calcium: from cell signalling to cell death. *The*
1269   *Journal of Physiology*, *529*(Pt 1), 57-68. https://doi.org/10.1111/J.1469-
1270   7793.2000.00057.X

1271 Eckmann, J., Clemens, L. E., Eckert, S. H., Hagl, S., Yu-Taeger, L., Bordet, T., Pruss,
1272   R. M., Muller, W. E., Leuner, K., Nguyen, H. P., & Eckert, G. P. (2014).
1273   Mitochondrial Membrane Fluidity is Consistently Increased in Different Models
1274   of Huntington Disease: Restorative Effects of Olesoxime. *Molecular*
1275   *Neurobiology*, *50*(1), 107-118. https://doi.org/10.1007/s12035-014-8663-3

1276 Elbaum-Garfinkle, S., & Brangwynne, C. P. (2015). Liquids, Fibers, and Gels: The
1277   Many Phases of Neurodegeneration. *Developmental Cell*, *35*(5), 531-532.
1278   https://doi.org/https://doi.org/10.1016/j.devcel.2015.11.014

Faitg, J., Lacefield, C., Davey, T., White, K., Laws, R., Kosmidis, S., Reeve, A. K., Kandel, E. R., Vincent, A. E., & Picard, M. (2021). 3D neuronal mitochondrial morphology in axons, dendrites, and somata of the aging mouse hippocampus. *Cell Reports*, *36*(6), 109509-109509. https://doi.org/https://doi.org/10.1016/j.celrep.2021.109509

Farías, G. G., Guardia, C. M., Britt, D. J., Guo, X., & Bonifacino, J. S. (2015). Sorting of Dendritic and Axonal Vesicles at the Pre-axonal Exclusion Zone. *Cell Reports*, *13*(6), 1221-1232. https://doi.org/10.1016/j.celrep.2015.09.074

Fecher, C., Trovò, L., Müller, S. A., Snaidero, N., Wettmarshausen, J., Heink, S., Ortiz, O., Wagner, I., Kühn, R., Hartmann, J., Karl, R. M., Konnerth, A., Korn, T., Wurst, W., Merkler, D., Lichtenthaler, S. F., Perocchi, F., & Misgeld, T. (2019). Cell-type-specific profiling of brain mitochondria reveals functional and molecular diversity. *Nature Neuroscience*, *22*(10), 1731-1742. https://doi.org/10.1038/s41593-019-0479-z

Gumy, L. F., Katrukha, E. A., Grigoriev, I., Jaarsma, D., Kapitein, L. C., Akhmanova, A., & Hoogenraad, C. C. (2017). MAP2 Defines a Pre-axonal Filtering Zone to Regulate KIF1- versus KIF5-Dependent Cargo Transport in Sensory Neurons. *Neuron*, *94*(2), 347-362.e347. https://doi.org/10.1016/j.neuron.2017.03.046

Gumy, L. F., Yeo, G. S. H., Tung, Y. C. L., Zivraj, K. H., Willis, D., Coppola, G., Lam, B. Y. H., Twiss, J. L., Holt, C. E., & Fawcett, J. W. (2011). Transcriptome analysis of embryonic and adult sensory axons reveals changes in mRNA repertoire localization. *RNA*, *17*(1), 85-98. https://doi.org/10.1261/rna.2386111

Guo, S., Sun, Y., Wu, Z., Yang, R., Qu, L., & Li, Z. (2022). Simultaneous monitoring of mitochondrial viscosity and membrane potential based on fluorescence changing and location switching of carbon dots in living cells. *Carbon*, *195*, 112-122. https://doi.org/10.1016/J.CARBON.2022.04.006

Hagemann, C., Moreno Gonzalez, C., Guetta, L., Tyzack, G., Chiappini, C., Legati, A., Patani, R., & Serio, A. (2022). Axonal Length Determines Distinct Homeostatic Phenotypes in Human iPSC Derived Motor Neurons on a Bioengineered Platform. *Advanced Healthcare Materials*, *11*(10), e2101817-e2101817. https://doi.org/10.1002/ADHM.202101817

Kim, Y., Zheng, X., Ansari, Z., Bunnell, M. C., Herdy, J. R., Traxler, L., Lee, H., Paquola, A. C. M., Blithikioti, C., Ku, M., Schlachetzki, J. C. M., Winkler, J., Edenhofer, F., Glass, C. K., Paucar, A. A., Jaeger, B. N., Pham, S., Boyer, L., Campbell, B. C., . . . Gage, F. H. (2018). Mitochondrial Aging Defects Emerge in Directly Reprogrammed Human Neurons due to Their Metabolic Profile. *Cell Reports*, *23*(9), 2550-2558. https://doi.org/10.1016/j.celrep.2018.04.105

Klinman, E., & Holzbaur, E. L. F. (2015). Stress-Induced CDK5 Activation Disrupts Axonal Transport via Lis1/Ndel1/Dynein. *Cell Reports*, *12*(3), 462-473. https://doi.org/10.1016/j.celrep.2015.06.032

Koopman, W. J. H., Distelmaier, F., Hink, M. A., Verkaart, S., Wijers, M., Fransen, J., Smeitink, J. A. M., & Willems, P. H. G. M. (2008). Inherited complex I deficiency is associated with faster protein diffusion in the matrix of moving mitochondria. *American Journal of Physiology-Cell Physiology*, *294*(5), C1124-C1132. https://doi.org/10.1152/ajpcell.00079.2008

Kubánková, M., Summers, P. A., López-Duarte, I., Kiryushko, D., & Kuimova, M. K. (2019). Microscopic Viscosity of Neuronal Plasma Membranes Measured Using Fluorescent Molecular Rotors: Effects of Oxidative Stress and Neuroprotection. *ACS Applied Materials and Interfaces*, *11*(40), 36307-36315. https://doi.org/10.1021/acsami.9b10426

Kuter, K., Kratochwil, M., Berghauzen-Maciejewska, K., Głowacka, U., Sugawa, M. D., Ossowska, K., & Dencher, N. A. (2016). Adaptation within mitochondrial oxidative phosphorylation supercomplexes and membrane viscosity during degeneration of dopaminergic neurons in an animal model of early Parkinson's disease. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, *1862*(4), 741-753. https://doi.org/10.1016/J.BBADIS.2016.01.022

Lamoureux, P. L., O'Toole, M. R., Heidemann, S. R., & Miller, K. E. (2010). Slowing of axonal regeneration is correlated with increased axonal viscosity during aging. *BMC Neuroscience*, *11*(1), 140-140. https://doi.org/10.1186/1471-2202-11-140

Leduc, C., Padberg-Gehle, K., Varga, V., Helbing, D., Diez, S., & Howard, J. (2012). Molecular crowding creates traffic jams of kinesin motors on microtubules. *Proceedings of the National Academy of Sciences*, *109*(16), 6100-6105. https://doi.org/10.1073/pnas.1107281109

Lee, T. H., & Linstedt, A. D. (1999). Osmotically Induced Cell Volume Changes Alter Anterograde and Retrograde Transport, Golgi Structure, and COPI Dissociation. *Molecular Biology of the Cell*, *10*(5), 1445-1462. https://doi.org/10.1091/mbc.10.5.1445

Lidke, K. A., Rieger, B., Lidke, D. S., & Jovin, T. M. (2005). The role of photon statistics in fluorescence anisotropy imaging. *IEEE transactions on image processing*, *14*(9), 1237-1245. https://doi.org/10.1109/tip.2005.852458

Lin, M.-Y., Cheng, X.-T., Tammineni, P., Xie, Y., Zhou, B., Cai, Q., & Sheng, Z.-H. (2017). Releasing Syntaphilin Removes Stressed Mitochondria from Axons Independent of Mitophagy under Pathophysiological Conditions. *Neuron*, *94*(3), 595-610.e596. https://doi.org/10.1016/j.neuron.2017.04.004

López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2023). Hallmarks of aging: An expanding universe. *Cell*, *186*(2), 243-278. https://doi.org/10.1016/J.CELL.2022.11.001

MacAskill, A. F., Rinholm, J. E., Twelvetrees, A. E., Arancibia-Carcamo, I. L., Muir, J., Fransson, A., Aspenstrom, P., Attwell, D., & Kittler, J. T. (2009). Miro1 Is a Calcium Sensor for Glutamate Receptor-Dependent Localization of Mitochondria at Synapses. *Neuron*, *61*(4), 541-555. https://doi.org/10.1016/j.neuron.2009.01.030

Mar, F. M., Simoes, A. R., Leite, S., Morgado, M. M., Santos, T. E., Rodrigo, I. S., Teixeira, C. A., Misgeld, T., & Sousa, M. M. (2014). CNS Axons Globally Increase Axonal Transport after Peripheral Conditioning. *The Journal of Neuroscience*, *34*(17), 5965-5970. https://doi.org/10.1523/JNEUROSCI.4680-13.2014

Markaki, M., Tsagkari, D., & Tavernarakis, N. (2023). Mitophagy and long-term neuronal homeostasis. *Journal of Cell Science*, *136*(11). https://doi.org/10.1242/jcs.260638

Mattedi, F., Lloyd-Morris, E., Hirth, F., & Vagnoni, A. (2023). Optogenetic cleavage of the Miro GTPase reveals the direct consequences of real-time loss of function in Drosophila. *PLOS Biology*, *21*(8), e3002273-e3002273. https://doi.org/10.1371/journal.pbio.3002273

Mattedi, F., & Vagnoni, A. (2019). Temporal Control of Axonal Transport: The Extreme Case of Organismal Ageing. *Frontiers in Cellular Neuroscience*, *13*, 393-393. https://doi.org/10.3389/fncel.2019.00393

Mecocci, P., Beal, M. F., Cecchetti, R., Polidori, M. C., Cherubini, A., Chionne, F., Avellini, L., Romano, G., & Senin, U. (1997). Mitochondrial membrane fluidity

and oxidative damage to mitochondrial DNA in aged and AD human brain. *Molecular and Chemical Neuropathology*, *31*(1), 53-64. https://doi.org/10.1007/BF02815160

Mellone, M., Kestoras, D., Andrews, M. R., Dassie, E., Crowther, R. A., Stokin, G. B., Tinsley, J., Horne, G., Goedert, M., Tolkovsky, A. M., & Spillantini, M. G. (2013). Tau pathology is present in vivo and develops in vitro in sensory neurons from human P301S tau transgenic mice: a system for screening drugs against tauopathies. *The Journal of Neuroscience*, *33*(46), 18175-18189. https://doi.org/10.1523/JNEUROSCI.4933-12.2013

Metzner, K., Darawsha, O., Wang, M., Gaur, N., Cheng, Y., Rödiger, A., Frahm, C., Witte, O. W., Perocchi, F., Axer, H., Grosskreutz, J., & Brill, M. S. (2022). Age-dependent increase of cytoskeletal components in sensory axons in human skin. *Frontiers in Cell and Developmental Biology*, *10*. https://doi.org/10.3389/fcell.2022.965382

Milde, S., Adalbert, R., Elaman, M. H., & Coleman, M. P. (2015). Axonal transport declines with age in two distinct phases separated by a period of relative stability. *Neurobiology of Aging*, *36*(2), 971-981. https://doi.org/10.1016/j.neurobiolaging.2014.09.018

Miller, C. C., & Walker, J. (1924). The Stokes-Einstein law for diffusion in solution. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, *106*(740), 724-749. https://doi.org/10.1098/rspa.1924.0100

Miller, K. E., & Sheetz, M. P. (2004). Axonal mitochondrial transport and potential are correlated. *Journal of Cell Science*, *117*(13), 2791-2804. https://doi.org/10.1242/jcs.01130

Misgeld, T., Kerschensteiner, M., Bareyre, F. M., Burgess, R. W., & Lichtman, J. W. (2007). Imaging axonal transport of mitochondria in vivo. *Nature Methods*, *4*(7), 559-561. https://doi.org/10.1038/nmeth1055

Mitra, K., & Lippincott-Schwartz, J. (2010). Analysis of mitochondrial dynamics and functions using imaging approaches. *Current protocols in cell biology*, *4*(46), 4251-42521. https://doi.org/10.1002/0471143030.CB0425S46

Molines, A. T., Lemière, J., Gazzola, M., Steinmark, I. E., Edrington, C. H., Hsu, C. T., Real-Calderon, P., Suhling, K., Goshima, G., Holt, L. J., Thery, M., Brouhard, G. J., & Chang, F. (2022). Physical properties of the cytoplasm modulate the rates of microtubule polymerization and depolymerization. *Developmental Cell*, *57*(4), 466-479.e466. https://doi.org/10.1016/J.DEVCEL.2022.02.001

Morsci, N. S., Hall, D. H., Driscoll, M., & Sheng, Z.-H. (2016). Age-Related Phasic Patterns of Mitochondrial Maintenance in Adult Caenorhabditis elegans Neurons. *The Journal of Neuroscience*, *36*(4), 1373-1385. https://doi.org/10.1523/JNEUROSCI.2799-15.2016

Niescier, R. F., Hong, K., Park, D., & Min, K. T. (2018). MCU Interacts with Miro1 to Modulate Mitochondrial Functions in Neurons. *The Journal of Neuroscience*, *38*(20), 4666-4677. https://doi.org/10.1523/JNEUROSCI.0504-18.2018

Nunes, P., Roth, I., Meda, P., Féraille, E., Brown, D., & Hasler, U. (2015). Ionic imbalance, in addition to molecular crowding, abates cytoskeletal dynamics and vesicle motility during hypertonic stress. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(24), E3104-E3113. https://doi.org/10.1073/pnas.1421290112

Pandey, J. P., Shi, L., Brebion, R. A., & Smith, D. S. (2022). LIS1 and NDEL1 Regulate Axonal Trafficking of Mitochondria in Mature Neurons. *Frontiers in Molecular Neuroscience*, *15*, 841047-841047. https://doi.org/10.3389/fnmol.2022.841047

Parry, B. R., Surovtsev, I. V., Cabeen, M. T., Hern, C. S. O., & Dufresne, E. R. (2014). The Bacterial Cytoplasm Has Glass-like Properties and Is Fluidized by Metabolic Activity. *Cell*, *156*(1-2), 183-194. https://doi.org/10.1016/j.cell.2013.11.028

Persson, L. B., Ambati, V. S., & Brandman, O. (2020). Cellular Control of Viscosity Counters Changes in Temperature and Energy Availability. *Cell*, *183*(6), 1572-1585.e1516. https://doi.org/10.1016/j.cell.2020.10.017

Puchkov, E. O. (2013). Intracellular viscosity: Methods of measurement and role in metabolism. *Biochemistry (Moscow) Supplement Series A: Membrane and Cell Biology*, *7*(4), 270-279. https://doi.org/10.1134/S1990747813050140

Rappe, A., & McWilliams, T. G. (2022). Mitophagy in the aging nervous system. *Frontiers in Cell and Developmental Biology*, *10*, 978142-978142. https://doi.org/10.3389/fcell.2022.978142

Rhymes, E. R., Tosolini, A. P., Fellows, A. D., Mahy, W., McDonald, N. Q., & Schiavo, G. (2022). Bimodal regulation of axonal transport by the GDNF-RET signalling axis in healthy and diseased motor neurons. *Cell Death Dis*, *13*(7), 584. https://doi.org/10.1038/s41419-022-05031-0

Sabharwal, V., & Koushika, S. P. (2019). Crowd Control: Effects of Physical Crowding on Cargo Movement in Healthy and Diseased Neurons. *Frontiers in Cellular Neuroscience*, *13*. https://doi.org/10.3389/FNCEL.2019.00470

Sharma, A., Smith, H. J., Yao, P., & Mair, W. B. (2019). Causal roles of mitochondrial dynamics in longevity and healthy aging. *EMBO reports*, *20*(12), e48395-e48395. https://doi.org/10.15252/embr.201948395

Shu, T., Szórádi, T., Kidiyoor, G. R., Xie, Y., Herzog, N. L., Bazley, A., Bonucci, M., Keegan, S., Saxena, S., Ettefa, F., Brittingham, G., Lemiere, J., Fenyö, D., Chang, F., Delarue, M., & Holt, L. J. (2022). nucGEMs probe the biophysical properties of the nucleoplasm. *BioRxiv*, 2021.2011.2018.469159-462021.469111.469118.469159. https://doi.org/10.1101/2021.11.18.469159

Siegel, J., Suhling, K., Lévêque-Fort, S., Webb, S. E. D., Davis, D. M., Phillips, D., Sabharwal, Y., & French, P. M. W. (2003). Wide-field time-resolved fluorescence anisotropy imaging (TR-FAIM): Imaging the rotational mobility of a fluorophore. *Review of Scientific Instruments*, *74*(1), 182-192. https://doi.org/10.1063/1.1519934

Sleigh, J. N., Dawes, J. M., West, S. J., Wei, N., Spaulding, E. L., Gómez-Martín, A., Zhang, Q., Burgess, R. W., Cader, M. Z., Talbot, K., Yang, X.-L., Bennett, D. L., & Schiavo, G. (2017b). Trk receptor signaling and sensory neuron fate are perturbed in human neuropathy caused by Gars mutations. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(16), E3324-E3333. https://doi.org/10.1073/pnas.1614557114

Sleigh, J. N., Rossor, A. M., Fellows, A. D., Tosolini, A. P., & Schiavo, G. (2019). Axonal transport and neurological disease. *Nat Rev Neurol*, *15*(12), 691-703. https://doi.org/10.1038/s41582-019-0257-2

Sleigh, J. N., Tosolini, A. P., & Schiavo, G. (2020). In Vivo Imaging of Anterograde and Retrograde Axonal Transport in Rodent Peripheral Nerves. *Methods Mol Biol*, *2143*, 271-292. https://doi.org/10.1007/978-1-0716-0585-1_20

Sleigh, J. N., Vagnoni, A., Twelvetrees, A. E., & Schiavo, G. (2017a). Methodological advances in imaging intravital axonal transport. *F1000Research*, *6*, 200-200. https://doi.org/10.12688/f1000research.10433.1

Sleigh, J. N., Weir, G. A., & Schiavo, G. (2016). A simple , step - by - step dissection protocol for the rapid isolation of mouse dorsal root ganglia. *BMC Research Notes*, *9*, 82-82. https://doi.org/10.1186/s13104-016-1915-8

Sood, P., Murthy, K., Kumar, V., Nonet, M. L., Menon, G. I., & Koushika, S. P. (2018). Cargo crowding at actin-rich regions along axons causes local traffic jams. *Traffic*, *19*(3), 166-181. https://doi.org/https://doi.org/10.1111/tra.12544

Stavoe, A. K. H., Gopal, P. P., Gubas, A., Tooze, S. A., & Holzbaur, E. L. F. (2019). Expression of WIPI2B counteracts age-related decline in autophagosome biogenesis in neurons. *Elife*, *8*, e44219-e44219. https://doi.org/10.7554/eLife.44219

Stefanatos, R., & Sanz, A. (2018). The role of mitochondrial ROS in the aging brain. *FEBS Letters*, *592*(5), 743-758. https://doi.org/10.1002/1873-3468.12902

Steinmark, I. E., James, A. L., Chung, P.-H., Morton, P. E., Parsons, M., Dreiss, C. A., Lorenz, C. D., Yahioglu, G., & Suhling, K. (2019). Targeted fluorescence lifetime probes reveal responsive organelle viscosity and membrane fluidity. *PLoS ONE*, *14*(2), e0211165-e0211165. https://doi.org/10.1371/journal.pone.0211165

Sugawa, M., Coper, H., Schulze, G., Yamashina, I., Krause, R., & Dencher, A. (1996). Impaired Plasticity of Neurons in Aging. *Annals of the New York Academy of Sciences*, *786*(1), 274-282. https://doi.org/https://doi.org/10.1111/j.1749-6632.1996.tb39070.x

Sun, N., Youle, R. J., & Finkel, T. (2016). The Mitochondrial Basis of Aging. *Molecular Cell*, *61*(5), 654-666. https://doi.org/10.1016/j.molcel.2016.01.028

Sung, H., Tandarich, L. C., Nguyen, K., & Hollenbeck, P. J. (2016). Compartmentalized Regulation of Parkin-Mediated Mitochondrial Quality Control in the Drosophila Nervous System In Vivo. *The Journal of Neuroscience*, *36*(28), 7375-7391. https://doi.org/10.1523/JNEUROSCI.0633-16.2016

Takihara, Y., Inatani, M., Eto, K., Inoue, T., Kreymerman, A., Miyake, S., Ueno, S., Nagaya, M., Nakanishi, A., Iwao, K., Takamura, Y., Sakamoto, H., Satoh, K., Kondo, M., Sakamoto, T., Goldberg, J. L., Nabekura, J., & Tanihara, H. (2015). In vivo imaging of axonal transport of mitochondria in the diseased and aged mammalian CNS. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(33), 10515-10520. https://doi.org/10.1073/pnas.1509879112

Tosolini, A. P., Villarroel-Campos, D., Schiavo, G., & Sleigh, J. N. (2021). Expanding the Toolkit for In Vivo Imaging of Axonal Transport. *J Vis Exp*(178). https://doi.org/10.3791/63471

Vagnoni, A., & Bullock, S. L. (2018). A cAMP/PKA/Kinesin-1 Axis Promotes the Axonal Transport of Mitochondria in Aging Drosophila Neurons. *Current Biology*, *28*, 1265-1272. https://doi.org/10.1016/j.cub.2018.02.048

Vagnoni, A., Hoffmann, P. C., & Bullock, S. L. (2016). Reducing Lissencephaly-1 levels augments mitochondrial transport and has a protective effect in adult Drosophila neurons. *Journal of Cell Science*, *129*(1), 178-190. https://doi.org/10.1242/jcs.179184

van Spronsen, M., Mikhaylova, M., Lipka, J., Schlager, M. A., van den Heuvel, D. J., Kuijpers, M., Wulf, P. S., Keijzer, N., Demmers, J., Kapitein, L. C., Jaarsma, D.,

Gerritsen, H. C., Akhmanova, A., & Hoogenraad, C. C. (2013). TRAK/Milton
motor-adaptor proteins steer mitochondrial trafficking to axons and dendrites.
Neuron, *77*(3), 485-502. https://doi.org/10.1016/j.neuron.2012.11.027
Verkman, A. S., Armijo, M., & Fushimi, K. (1991). Construction and evaluation of a
frequency-domain epifluorescence microscope for lifetime and anisotropy
decay measurements in subcellular domains. *Biophysical chemistry*, *40*(1),
117-125. https://doi.org/10.1016/0301-4622(91)85036-p
Wang, X., & Schwarz, T. L. (2009). The Mechanism of Ca2+-Dependent Regulation
of Kinesin-Mediated Mitochondrial Motility. *Cell*, *136*(1), 163-174.
https://doi.org/10.1016/j.cell.2008.11.046
Zustiak, Silviya P., Nossal, R., & Sackett, D. L. (2011). Hindered Diffusion in Polymeric
Solutions Studied by Fluorescence Correlation Spectroscopy. *Biophysical
Journal*, *101*(1), 255-264.
https://doi.org/https://doi.org/10.1016/j.bpj.2011.05.035

1539
1540
1541
1542
1543