Higher-order combinatorial chromatin perturbations by engineered CRISPR-Cas12a for functional genomics

Hsiung CC[1,2,4,9,10], Wilson CM[2,3,4,9,10], Sambold NA[9], Dai R[2,4,5,9,10], Chen Q[7], Misiukiewicz S[2,4,5], Arab A[10], Teyssier N[6], O'Loughlin T[2,4,9], Cofsky JC[8], Shi J[7], Gilbert LA[#,2,4,9,10]

Author affiliations

[1]Dept. of Pathology, Stanford University School of Medicine, Stanford, CA 94305, USA
[2]Dept. of Urology, University of California, San Francisco, CA 94158, USA
[3]Tetrad Graduate Program, University of California, San Francisco, CA 94158, USA
[4]Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco, San Francisco, CA 94158
[5]Biomedical Sciences Graduate Program, University of California, San Francisco, San Francisco, CA 94158, USA
[6]Biological and Medical Informatics Graduate Program, University of California, San Francisco, San Francisco, CA 94158, USA
[7]Department of Cancer Biology, University of Pennsylvania, Philadelphia, PA, 19104, USA
[8]Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA, 02115, USA
[9]Innovative Genomics Institute, University of California, San Francisco, San Francisco, CA 94158, USA
[10]Arc Institute, Palo Alto, CA 94304, USA

[#]Correspondence to luke@arcinstitute.org

Abstract

Multiplexed genetic perturbations are critical for testing functional interactions among coding or non-coding genetic elements. Compared to double-stranded DNA cutting, repressive chromatin formation using CRISPR interference (CRISPRi) avoids genotoxicity and is more effective for perturbing non-coding regulatory elements in pooled assays. However, current CRISPRi pooled screening approaches are limited to targeting 1-3 genomic sites per cell. To develop a tool for higher-order ($\geq$3) combinatorial targeting of genomic sites with CRISPRi in functional genomics screens, we engineered an *Acidaminococcus* Cas12a variant -- referred to as multiplexed transcriptional interference AsCas12a (multiAsCas12a). multiAsCas12a incorporates a key mutation, R1226A, motivated by the hypothesis of nicking-induced stabilization of the ribonucleoprotein:DNA complex for improving CRISPRi activity. multiAsCas12a

significantly outperforms prior state-of-the-art Cas12a variants in combinatorial CRISPRi targeting using high-order multiplexed arrays of lentivirally transduced CRISPR RNAs (crRNA), including in high-throughput pooled screens using 6-plex crRNA array libraries. Using multiAsCas12a CRISPRi, we discover new enhancer elements and dissect the combinatorial function of cis-regulatory elements. These results instantiate a group testing framework for efficiently surveying potentially numerous combinations of chromatin perturbations for biological discovery and engineering.

Introduction

Functional interactions among combinations of genetic elements underlie many natural and engineered phenotypes [1–3], often involving higher-order ($\geq$3-plex) combinations of coding [4,5] or non-coding elements [6–9]. Experimentally testing higher-order combinations of genetic perturbations has been limited by throughput, with prior systematic analyses primarily performed in yeast [10–14]. In mammalian functional genomics, Cas9-based pooled CRISPR screens [15,16] using sequencing readouts have been limited in multiplexing capability, with only a few studies targeting more than 2 genomic sites per cell [17,18]. Further multiplexing in Cas9-based pooled screening is challenging due to 1) increasingly complex cloning schemes for large constructs encoding multiple guides, each expressed from a separate promoter [17–19], and 2) length-dependent high recombination frequencies of lentiviral guide libraries [20–22]. Conceptually, it also remains unclear how to tractably survey the potentially vast combinatorial spaces for $\geq$3-plex perturbations.

Cas12a, a member of the type V CRISPR-Cas family, has been proposed as an alternative to Cas9 for genetic perturbations due to enhanced multiplexing capabilities. Cas12a harbors RNase activity, separable from its DNase activity, that can process a compact primary transcript expressed from a single promoter into multiple CRISPR RNAs (crRNA) [23,24]. An array of multiple Cas12a crRNAs, each composed of 19nt direct repeat and 19-23nt spacer, can be encoded on a chemically synthesized oligo for single-step cloning into an expression vector [25–29]. Cas12a has been engineered for mammalian cell applications using its DNase activity to disrupt coding gene function using single or multiplexed crRNA constructs in individual well-based assays [24–27,30–32] and in pooled sequencing screens [28,29,32–36]. However, extended multiplexing with fully DNase-competent Cas12a is expected to be constrained by genotoxicity from double-stranded DNA breaks in many biological contexts [28,37,38,39–42]. In principle, avoiding genotoxicity can be achieved by using DNase-dead Cas fusion proteins to control chromatin state, such as by dCas9-based transcriptional repression (CRISPRi) and activation (CRISPRa)[43–45]. Moreover, CRISPRi is more efficient than DNA cutting at perturbing enhancers in pooled screens [46–48], likely due to CRISPRi's larger genomic

window of activity via formation of repressive chromatin [49]. Thus, a DNase-dead Cas12a
(dCas12a) functional genomics platform for multi-site CRISPRi targeting would be
highly desirable for testing the combinatorial functions of coding and non-coding genetic
elements. However, no dCas12a-based pooled CRISPRi screening platform has been
reported. Several studies have used dCas12a fusion proteins for CRISPRi in human
cells in individual well-based assays, reporting either successful [27,50–52] or unsuccessful
[53] repression of target genes. These dCas12a CRISPRi studies delivered crRNA
plasmids by transient transfection, rather than lentiviral transduction. Transient plasmid
transfections express synthetic components at 10 to 1000-fold higher than single-copy
lentiviral integration of crRNA constructs, which is required in pooled screens to attribute
cellular phenotypes to unique crRNA constructs by high-throughput sequencing [15,16].
Whether prior dCas12a CRISPRi constructs are sufficiently potent for pooled screens
remains unclear.
94
In this study, we show that existing dCas12a CRISPRi fusion constructs function poorly
when used with limiting doses of lentivirally delivered components, thus precluding their
application in pooled screens. We engineered a new Cas12a variant that incorporates a
key mutation, R1226A, which enhances stability of the ribonucleoprotein:DNA complex
in the form of a nicked DNA intermediate in vitro [54,55]. We show that in human cells,
R1226A significantly improves CRISPRi activity in the setting of lentivirally delivered
crRNA constructs, enabling use of 6-plex crRNA arrays in high-throughput pooled
screens using 6-plex crRNA arrays and up to 10-plex crRNA arrays in well-based
assays. We use this combinatorial CRISPRi platform to efficiently discover new
enhancer elements and to test higher-order combinatorial perturbations of cis-regulatory
elements. These results instantiate a group testing framework that enables efficient
searches of potentially large combinatorial spaces of chromatin perturbations.
107
108 **Results**
109
110 **CRISPRi using state-of-the-art dAsCas12a fusion proteins is dose-limited and**
111 **hypoactive in the setting of lentivirally delivered components**
112
We focused on building a CRISPRi functional genomics platform using
Acidaminococcus Cas12a (AsCas12a), the only Cas12a ortholog with demonstrated
success in pooled screens in mammalian cells [28,29,32–35,56,57]. A previous study reported
using dAsCas12a for CRISPRi by plasmid transient transfection delivery of dAsCas12a-
KRABx3 protein (harboring the E993A DNase-dead mutation) and crRNA in HEK 293T
cells (Campa et al. 2019). To test this construct in the setting of lentivirally delivered
crRNA, we introduced dAsCas12a-KRABx3 by piggyBac transposition in K562 cells,
followed by lentiviral transduction of single crRNA constructs targeting canonical TTTV

protospacer adjacent motifs (PAM) proximal to transcriptional start sites of 4 cell surface
genes, 3 of which (CD55, CD81 and B2M) have been successfully knocked down by
dCas9-KRAB CRISPRi [58]. Throughout this study we encoded crRNAs in a previously
optimized CROP-seq lentiviral vector [29,59]. We observed no expression change in any of
the targeted genes (Fig. 1A-B and Fig. S1-S2). We confirmed the expression of
dAsCas12a-KRABx3 by western blot (Fig. S3) and by flow cytometry monitoring of the
in-frame P2A-BFP (Fig. S4A). We also observed this lack of CRISPRi activity for
dAsCas12a-KRABx3 using lentivirally transduced crRNAs in C4-2B prostate cancer
cells (Fig. S4B). In contrast, transient co-transfection of dAsCas12a-KRABx3 and
CD55-targeting crRNA plasmids shows modest CRISPRi knockdown in HEK 293T cells
(Fig. S5), consistent with prior work [27]. These findings indicate that the requirements for
CRISPRi activity using dAsCas12a-KRABx3 with lentiviral crRNA constructs are distinct
from those of plasmid transient transfection in HEK 293T cells [27].
134
In an attempt to overcome this lack of CRISPRi activity, we tested combinations of
several Cas12a mutations representing state-of-the-art optimizations of Cas12a. These
include: 1) E174R/S542R/K548R (enhanced AsCas12a, enAsCas12a)[31]; 2)
M537R/F870L (AsCas12a ultra)[60]; 3) W382A, a mutation that reduces R-loop
dissociation in vitro for an orthologous enzyme (*Lachnospiraceae* Cas12a W355A)[61], but
which has not yet been tested in cells. We generated six dAsCas12a variants that each
harbor the DNase-inactivating D908A mutation, plus a select combination of the
aforementioned mutations. We delivered these variants in K562 cells by stable lentiviral
expression, followed by lentiviral transduction of crRNA construct targeting the TSS of
CD55 or CD81 (Fig. 1C and Fig. S6). Among this panel, denAsCas12a-KRAB
(E174R/S542R/K548R, plus D908A DNase-dead mutation) performed the best and
demonstrated strong repression of CD55. However, even for this best construct, we
observed weak repression of CD81, indicating inconsistent performance across crRNAs
(Fig. 1C and Fig. S6).
149
Dose-response and construct potency are key considerations for multiplexed
applications, as increased multiplexing effectively reduces the Cas protein available to
bind each individual crRNA. Focusing on denAsCas12a-KRAB as the top variant, we
tested the effect of separately altering the dosage of Cas12a protein and crRNAs. We
found that increasing the MOI of the denAsCas12a-KRAB construct from ~1 to ~5 can
improve CRISPRi knockdown by some crRNA constructs (Fig. 1D and Fig. S7A).
However, CRISPRi activity of denAsCas12a-KRAB is lost when the crRNA MOI is
reduced to <1 to mimic the low MOI required for pooled screens (Fig. 1E and Fig. S7B).
More problematically, CD81 knockdown by a 3-plex crRNA (CD81-1_CD151-3_CD55-4)
is extremely weak (~0%-25% median expression knockdown relative to non-targeting

control) across all doses of protein (Fig. 1D and Fig. S7A) and crRNA (Fig. 1E and Fig.
S7B) tested.
162
Given the inconsistent and deficient performance of denAsCas12a-KRAB, we tested an
alternative CRISPRi approach without mutating the RuvC DNase active site. In the
setting of transient plasmid transfection delivery in HEK 293T cells, wild-type AsCas12a
has been used for transcriptional control with truncated (15nt) crRNA spacers, which
enable DNA binding but not cleavage [26,27]. We tested this approach by fusing KRAB or
KRABx3 to opAsCas12a, a fully DNase-active Cas12a optimized for pooled screens [29].
We confirmed that 15nt spacers do not support DNA cleavage, while 23nt spacers do
(Fig. S8). However, using 15nt spacers, we observed weak or no CRISPRi activity in
two cell lines (Fig. S8). In total, we demonstrated 3 separate approaches to abolish the
DNase activity of AsCas12a (E993A in Fig. 1B and Fig. S4; D908A in Fig. 1C-E and Fig.
S7B; and truncated spacers in Fig. S8) result in poor CRISPRi activity using lentivirally
transduced crRNA constructs.
175
176 **multiAsCas12a-KRAB (R1226A/E174R/S542R/K548R), a variant that favors a**
177 **nicked DNA intermediate, substantially improves lentivirally delivered CRISPRi**
178
The mediocre performance of dAsCas12a for CRISPRi surprised us given the success
of AsCas12a in DNA-cutting pooled screens [28,29,35,36]. We wondered whether full
inactivation of DNA cutting in dAsCas12a may preclude strong CRISPRi activity by
reducing DNA affinity. Previous studies indicate that DNA cleavage strengthens the
Cas12a:DNA interaction [54,62,63]. In the Cas12a DNA cleavage process, the RuvC active
site first cuts the non-target strand, then the target strand [64]. While double-strand breaks
are undesired for CRISPRi applications, we wondered whether favoring the nicked DNA
intermediate might reduce the R-loop dissociation rate (Fig. 2A, see Discussion). In
support of this possibility, in vitro studies showed that dCas12a:DNA complexes are 20-
fold more stable when the non-target strand is pre-cleaved [54], and that non-target strand
nicking biases Cas12a:DNA complexes away from dissociation-prone conformations
[65,66].
191
To engineer nicking-induced stabilization of AsCas12a binding to DNA for CRISPRi
applications, we incorporated R1226A, a mutation that has not been tested in the
context of transcriptional control. Relative to wild-type AsCas12a, the R1226A mutant,
originally described as a nickase [55], is ~100-1,000 fold slower in cleaving the non-target
DNA strand and ~10,000-fold slower in cleaving the target DNA strand in vitro [54].
Consistent with nicking-induced stabilization, AsCas12a R1226A indeed binds DNA
more strongly in vitro than the fully DNase-inactivated D908A variant [54]. We expect the
R1226A mutation to both disfavor R-loop reversal and slow progression to double-

stranded breaks (Fig 2A; see Discussion). We hypothesized that, by trapping the
ribonucleoprotein:DNA in a nicked DNA intermediate, the R1226A mutation would
prolong chromatin occupancy and thus the time available for the KRAB domain to
recruit transcriptional repressive complexes.
204
To test the impact of R1226A on CRISPRi activity, we replaced the DNase-inactivating
D908A in denAsCas12a-KRAB with R1226A, and hereafter refer to this Cas12a variant
as multiAsCas12a (multiplexed transcriptional interference, i.e.
R1226A/E174R/S542R/K548R). To compare dose-sensitivity of their CRISPRi activities,
we stably expressed denAsCas12a-KRAB and multiAsCas12a-KRAB protein and
crRNA constructs by lentiviral transduction at high vs. low MOI's for protein and crRNA
constructs. Across a panel of single and 3-plex crRNA constructs, multiAsCas12a-
KRAB consistently exhibits robust CRISPRi with less sensitivity to low MOI of protein or
crRNA constructs (Fig. 2B-C, Fig. S9 and Fig. S10). Notably, multiAsCas12a-KRAB
substantially rescues the activities of several crRNA constructs that are virtually inactive
for denAsCas12a-KRAB even at high protein dose (Fig. 2C), including a non-canonical
GTTC PAM target (crCD151-3, Fig. 2E). Targeting by multiAsCas12a-KRAB results in
low indel frequencies at crCD81-1 (2.48%) and crCD55-4 (7.9%) target sites (Fig. 2F).
Simulations accounting for DNA copy number indicate that any possible gene
expression impact from these indels are far lower than the observed target gene
knockdown (Fig. S14A-B). multiAsCas12a-KRAB CRISPRi activity shows generally
minimal or no off-target transcriptomic effects as evaluated by bulk RNA-seq (Fig. S11).
222
223   **multiAsCas12a-KRAB enables multi-gene transcriptional repression using higher-**
224   **order arrayed crRNA lentiviral constructs**
225
We next tested performance of multiAsCas12a-KRAB in targeting >3 genomic sites per
cell for CRISPRi using lentiviral crRNA arrays [29]. To minimize the possibility of lentiviral
recombination, the expression construct (Fig. 3A) uses a unique direct repeat variant at
each position of the array, selected from a set of previously tested direct repeat variants
[28]. We assembled a panel of 13 distinct crRNA constructs (7 single-plex, two 3-plex, two
4-plex, two 5-plex, and two 6-plex) from individually active TSS-targeting spacers (Fig.
3B-C and Fig. S12). For these 13 crRNA constructs, we compared the CRISPRi
activities of denAsCas12a-KRAB, multiAsCas12a-KRAB, and multiAsCas12a (no
KRAB). For a subset of crRNA constructs we also added enAsCas12a-KRAB (DNase
fully active) as comparison. For these experiments and the remainder of this study we
use piggyBac transposition to constitutively express all fusion proteins at very similar
levels (Fig. S3 and Fig. S13B), which yields results similar to that obtained from high
MOI (~5) lentiviral fusion protein delivery and avoids day-to-day variations in lentiviral
titers. Across the entire crRNA panel, multiAsCas12a-KRAB substantially outperforms

denAsCas12a-KRAB in CRISPRi activity for 7 out of 7 constructs targeting CD81 (Fig.
3B); 4 out of 6 constructs targeting B2M (Fig. S12); and 6 out of 6 constructs targeting
KIT (Fig. 3C). For CD55 (Fig. S12), multiCas12a-KRAB substantially outperforms
denAsCas12a-KRAB using crCD55-5 (weaker spacer), and performs the same as or
marginally better than denAsCas12a-KRAB for 7 constructs containing crCD55-4
(stronger spacer). Similarly superior CRISPRi performance by multiAsCas12a-KRAB
was observed in C4-2B cells (Fig. S4).
247
For all crRNA constructs tested, multiAsCas12a alone shows much lower impact on
target gene expression than multiAsCas12a-KRAB (e.g. Fig. 3B for CD81),
demonstrating strong dependence of gene knockdown on the KRAB domain. For some
target genes, such as KIT, partial knockdown can be observed for multiAsCas12a alone
(Fig. 3C). Such gene knockdown may be due to 1) non-genetic perturbation of
transcription, or 2) alteration of DNA sequences crucial for transcription due to residual
DNA cutting. To distinguish these possibilities, we used short-read Illumina sequencing
of PCR amplicons (Fig. 3D, Fig. S14C and Fig. S15A), as well as long-read Nanopore
sequencing of native genomic DNA up to tens of kilobases in length (Fig. S16C), to
quantify indels generated by the panel of fusion proteins using single-site or dual-site
targeting in the KIT locus. For multiAsCas12a-KRAB we observed a maximum indel
frequency of 5% anywhere in the 340 bp PCR amplicon when simultaneously targeting
two sites spaced ~112 bp apart in the KIT locus (Fig. 3D). Based on this observed indel
frequency, we calculated an upper estimate of expected 1.3% median KIT expression
knockdown driven solely by indels, far less than the observed 90.4% median expression
knockdown, which is 44.4% in excess of the observed for denAsCas12a-KRAB (Fig.
S15B). Similar conclusions are supported by additional measurements at this and other
loci obtained for multiAsCas12a-KRAB and/or multiAsCas12a using short-read PCR
amplicon sequencing (Fig. S14) and/or long-read Nanopore sequencing (Fig. S16).
Altogether, our analyses of indel frequencies demonstrate that target gene knockdown
by multiAsCas12a-KRAB is largely attributable to non-genetic perturbation of
transcription via a combination of direct obstruction of transcription by the Cas protein
(as was observed for dCas9[43,44,67]) and KRAB-mediated repression.
271
At the single cell level, multiAsCas12a-KRAB consistently outperforms denAsCas12a-
KRAB in the fraction of cells with successful double knockdown (Fig. 3E-F and Fig. S17)
and triple knockdown (Fig. 3G and Fig. S17) of target genes using higher-order crRNA
arrays. To test the upper limit of multiplexing, we constructed 8-plex and 10-plex
constructs assembled using individually active spacers. In these 8-plex and 10-plex
arrays, spacers encoded in various positions within the array maintain robust CRISPRi
activity (i.e. for CD55, KIT and B2M, Fig 3H). However, for crCD81-1 encoded at the 3'
most position shows progressive diminishment in CRISPRi activity with further

multiplexing at 8-plex and 10-plex (Fig. 3H). This pattern suggests an intrinsic deficiency
of crCD81-1 that is unmasked by further multiplexing, perhaps related to the dose
sensitivity of this spacer (Fig. 2B-C). Nevertheless, these results indicate that 8-plex and
10-plex crRNA arrays can support robust CRISPRi activity for most spacers within these
arrays. We also observed that a specific 6-plex crRNA construct (crCD81-1_crB2M-
1_crB2M-3_crKIT-2_crKIT-3_crCD55-4, 6-plex #2 in Fig. 3H) fails to knockdown B2M,
despite robust CRISPRi of the other target genes. However, the same combination of
spacers in a slightly different 6-plex arrangement (crCD55-4_crB2M-1_crB2M-3_crKIT-
2_crKIT-3_crCD81-1) and also in 8-plex and 10-plex embodiments achieve ~50% B2M
knockdown (Fig. 3H). These results indicate the existence of still unpredictable pre-
crRNA sequence context influences on CRISPRi activity of specific spacers, unrelated
to genomic distance from the U6 promoter.
292
293   **multiAsCas12a-KRAB outperforms denCas12a-KRAB and performs similarly to**
294   **dCas9-KRAB in pooled single-guide CRISPRi screens**
295
Given the success of multiAsCas12a-KRAB in individual well-based assays using
lentivirally delivered crRNAs, we next evaluated its performance in the context of high-
throughput pooled screens. We designed a library, referred to as Library 1
(summarized in Fig. S18), aimed at extracting patterns for Cas12a CRISPRi activity with
respect to genomic position relative to the TSS using cell fitness as a readout. Library 1
contains 77,387 single crRNA lentiviral constructs tiling all predicted canonical TTTV
PAM sites and non-canonical PAM's (recognizable by enAsCas12a [31]) in the -50bp to
+300bp region around the TSS's of 559 common essential genes with K562 cell fitness
defects in prior genome-wide dCas9-KRAB screens [68].
305
Using K562 cells piggyBac-engineered to constitutively express multiAsCas12a-KRAB
or denAsCas12a-KRAB, we conducted a pooled cell fitness screen using this TSS tiling
crRNA library (MOI = 0.15). In this assay, CRISPRi knockdown of target essential genes
results in the relative depletion of cells harboring the corresponding crRNA over time,
quantified as a cell fitness score (Fig. 4A). Concordance between cell fitness scores of
screen replicates is high for multiAsCas12a-KRAB (R = 0.71) and much lower for
denAsCas12a-KRAB (R = 0.32), the latter due to much lower signal-to-background ratio
(Fig. S19). The cell fitness score distributions are virtually indistinguishable between the
intergenic targeting negative controls and the non-targeting negative controls (Fig. S20),
indicating no appreciable non-specific genotoxicity from multiAsCas12a-KRAB single-
site targeting. Among the 3,326 crRNA's targeting canonical TTTV PAM's, 24.5% vs.
17.5% showed a fitness defect in multiAsCas12a-KRAB vs. denAsCas12a-KRAB,
respectively (using the 5th percentile of intergenic negative controls as a threshold),

with the magnitude of effect for each crRNA overall stronger for multiAsCas12a-KRAB
(Fig. 4B).
321
Previous studies using dCas9-KRAB have identified a strong association between
CRISPRi activity and genomic proximity of the crRNA binding site to the TSS [44,52]. We
found that multiAsCas12a-KRAB generates a metagene profile of average CRISPRi
activity around the TSS that is remarkably similar in magnitude and bimodal genomic
distribution of CRISPRi activity obtained by dCas9-KRAB targeting NGG PAM's (Fig.
4C), consistent with nucleosomal hindrance near the +150bp region [52,69]. denCas12a-
KRAB is substantially weaker than both at all positions relative to the TSS (Fig. 4C).
multiAsCas12a-KRAB also outperforms denAsCas12a-KRAB in the average CRISPRi
activity of the top 3 best performing crRNAs/sgRNA for each TSS, and is similar to
dCas9-KRAB (Fig. 4D). Compared to the canonical TTTV PAM's, a smaller proportion
of crRNAs targeting non-canonical PAM's are active, in agreement with lower median
CRISPick on-target activity predictions (Fig. 4E) [28,70]. Within each PAM sequence,
individual crRNAs show significant variations in activity not accounted for by CRISPick
predictions (Fig. S21).
336
337 **multiAsCas12a-KRAB enables pooled sequencing screens using 6-plex crRNA**
338 **arrays**
339
To evaluate the performance of multiAsCas12a-KRAB in pooled sequencing screens
using multiplexed crRNA constructs, we constructed a library consisting of 6-plex
crRNAs. We refer to this 6-plex library as Library 2 (summarized in Fig. S18), which
includes Sublibrary A (described in this section) and Sublibrary B (described in the next
section). Sublibrary A was designed to contain 84,275 6-plex constructs for evaluating
CRISPRi activity at each of the 6 positions in the array in a K562 cell fitness screen
(Fig. 4F). Each 6-plex construct has one of the 6 positions designated as the "test"
position, which can encode either 1) a spacer targeting one of the top 50 essential gene
TSS's (ranked based on prior dCas9-KRAB screen data [52]), or 2) an intergenic negative
control (Fig. 4F). The remaining 5 positions in the array are designated as "context"
positions that encode negative control spacers drawn from a separate set of 30
negative control spacers (Fig. 4F). The motivation for this library design was to enable
sampling multiple sets of context spacers for a given test position.
353
The entirety of Library 2 was used in a cell fitness screen in K562 cells piggyBac-
engineered to stably express multiAsCas12a-KRAB. For a given test position spacer,
we calculated the average cell fitness scores from the top 3 context constructs ranked
by cell fitness score (Fig. 4F). The 123 spacers that previously showed strong cell
fitness scores as single crRNAs in the Library 1 screen also showed cell fitness score

distributions (average of top 3 contexts) that are clearly distinct from the corresponding
negative control distributions for the same test position (Fig. 4G). Each TSS-targeting
spacer encoded in the test position shows weaker cell fitness score than the same
spacer encoded singly in the Library 1 screen (Fig. S22). The recall of empirically active
single crRNA spacers from the Library 1 screen by the 6-plex crRNA constructs in this
Library 2 Sublibrary A screen ranges from 56%-92% across test positions (Fig. 4G). As
each position in the array is assigned a unique direct repeat variant held constant
across all constructs in this analysis, these apparent positional biases may reflect
contributions from differences among direct repeat variants. Together, these results
systematically demonstrate that the majority of individually active crRNAs retain
measurable activities when embedded within 6-plex crRNA arrays in the setting of
pooled screens using multiAsCas12a-KRAB.
371
372   **multiAsCas12a-KRAB enables discovery and higher-order combinatorial**
373   **perturbations of cis-regulatory elements**
374
The human genome contains ~500,000 predicted enhancers [71], a small minority of
which have been functionally tested by perturbations. To our knowledge, no study has
reported enhancer perturbation by CRISPRi using Cas12a. We confirmed that, similar
to dCas9-KRAB [72], multiAsCas12a-KRAB targeting using single crRNAs can effectively
perturb a known enhancer of the HBG gene, HS2 (Fig. 5A and Fig. S11). We next used
multiAsCas12a-KRAB to discover previously uncharacterized enhancers using the
CD55 locus in K562 cells as a myeloid cell model. CD55 encodes for decay-
accelerating factor, a cell surface protein that inhibits the activation of complement and
is expressed in most human cell types [73]. CD55 function in the myeloid lineage is
relevant in multiple disease states, including paroxysmal nocturnal hemoglobinuria [74]
and malaria [75,76]. To our knowledge, no known enhancers in myeloid cells have been
identified for CD55. In K562 cells, several DNase hypersensitive sites (DHS) bearing
histone 3 lysine 27 (H3K27Ac), a modification associated with active enhancers [71,77],
reside near CD55 (Fig. 5B). To conduct a well-based flow cytometry screen of the DHSs
within this general region for enhancers that regulate CD55, we designed 21 4-plex
crRNAs (encompassing 88 unique spacers) targeting 11 manually selected regions (R1-
R11) bearing varying levels of DNase hypersensitivity and H3K27Ac, plus a negative
control region (R12) devoid of DHS and H3K27Ac. R1-R4 are predicted by the activity-
by-contact (ABC) model [78] as candidate enhancers. Each region is independently
targeted by two completely distinct 4-plex crRNAs (except R10 and R12 are each
targeted by one 4-plex crRNA). We observed ~50%-75% reduction in CD55 expression
upon multiAsCas12a CRISPRi targeting of the ABC-predicted R1-R4, whereas no
decrease in CD55 expression is observed for R5-R12. For each region, the two distinct
4-plex crRNA arrays show quantitatively similar levels of CD55 knockdown (Fig. 5B),

indicating each array contains some 4-plex or lower-order combination of active
spacers. This consistency in the magnitude of CD55 expression knockdown likely
reflects the magnitude of true enhancer impact on gene transcription, rather than
technical peculiarities of individual spacer activities, which might be more unpredictably
variable and labor-intensive to test if encoded as single-plex perturbations. To our
knowledge, R1-R4 are the first functionally demonstrated enhancers for CD55 in a
myeloid cell type, in addition to another enhancer recently reported in a B-cell model [79].
Using opAsCas12a to target R1-R4 for DNA cutting using the same 4-plex crRNAs
elicits very little or no CD55 expression knockdown, despite potent knockdown by a
positive control crRNA targeting a coding exon (Fig. 5C).
409
To further test the utility of multiAsCas12a-KRAB in studies of enhancer function, we
used the MYC locus as a model. Prior studies using CRISPRi pooled screens in K562
cells have shown that MYC expression is proportional to cell fitness [80] and is regulated
by several enhancers identified by screens using cell fitness[80] and mRNA expression [81]
readouts. A recent study found that pairwise dCas9-KRAB perturbations of these
enhancers elicit stronger phenotypes than perturbing single enhancers [82]. Here we use
multiasCas12a-KRAB to dissect the phenotypic impact of $\geq$3-plex perturbations of MYC
cis-regulatory elements, which remained unknown. To avoid testing intractably
numerous higher-order combinations of crRNA spacers that are largely uninformative
due to the inclusion of weak or inactive spacers, we pre-screened for a small group of
active 3-plex crRNA combinations that can be subsequently assembled into higher-
order combinations. We used multiAsCas12a-KRAB to test four 3-plex crRNA
constructs targeting combinations of MYC cis-regulatory elements (3 crRNAs for
promoter and 3 crRNAs for each of 3 known enhancers, e1, e2 and e3) in a well-based
cell competition assay (Fig. 6A-B). We found that these four 3-plex crRNAs induce
varying degrees of cell fitness defect as a proxy of MYC expression knockdown,
indicating that each construct contains some spacer combination with CRISPRi activity.
For comparison, we included denAsCas12a-KRAB, multiAsCas12a, enAsCas12a-
KRAB and opAsCas12a, which showed relative activities that further demonstrate
multiAsCas12a-KRAB's superior gene knockdown is largely attributable to non-genetic
transcriptional perturbation (Fig. 6B).
431
We in silico assembled these 12 nominated spacers and 3 intergenic negative control
spacers into Library 2 Sublibrary B, consisting of 6,370 6-plex permutations encoded as
6-plex crRNA arrays (Fig. 6C and summarized in Fig. S18). These 6-plex arrays each
target up to 4 cis-regulatory elements (promoter + 3 enhancers) with up to 3 spacers per
element. Negative control spacers fill in the remaining array positions not filled by
targeting spacers. This Sublibrary B was included as part of the cell fitness screen for
the entirety of Library 2. Among 1,823 6-plex arrays with sufficient read coverage for

analysis, we grouped them into 16 categories, based on whether it encodes at least 1
spacer targeting the promoter, and/or at least 1 spacer targeting each of the 3
enhancers (Fig. 6D). We found that higher-order targeting of enhancers shows stronger
cell fitness defects beyond lower-order enhancer combinations (Fig. 6D, left panel). Co-
targeting the promoter together with any combination of enhancers showed greater cell
fitness defect over targeting the promoter alone while also exhibiting the cumulative
effects of multi-enhancer targeting (Fig. 6D, right panel). These results suggest that
when targeting subsets of cis-regulatory elements in a locus by CRISPRi, other cis-
regulatory elements can compete with CRISPRi to partially sustain gene transcription.
Such effects may reflect how cis-regulatory elements combinatorially respond to
endogenous repressive cues in the natural regulation of MYC gene transcription. These
results indicate that co-targeting multiple cis-regulatory elements may be more effective
for gene knockdown than targeting the promoter alone for some genes.
452
Discussion
In this study, we engineered multiAsCas12a-KRAB as a new platform for higher-order
combinatorial CRISPRi perturbations of gene transcription and enhancer function. The
enhanced CRISPRi potency of multiAsCas12a-KRAB is more robust to lower doses of
ribonucleoprotein (Fig. 2B-C, Fig. 2E), critically enabling higher-order multiplexing (Fig.
3) and high-throughput pooled screening applications conducted at single-copy
integrations of crRNA (Fig. 4 and Fig. 6C-D). We propose that the improved CRISPRi
activity of multiAsCas12a-KRAB emerges from prolonged chromatin occupancy due to
DNA nicking (Fig. 2A). This strategy is conceptually distinct from prior protein
engineering approaches to improving Cas12a function in mammalian cells, which
focused on substituting for positively charged amino acid residues near the protein:DNA
interface [31,50], using directed evolution to optimize DNA cleavage [60,83], or optimizing
transcriptional effector domain function [57]. We propose the following biophysical
explanation for improved multiAsCas12a function, grounded in prior in vitro literature. In
the absence of nicking, R-loop reversal occurs by invasion of the non-target strand into
the crRNA:target strand duplex, displacing the crRNA in a process analogous to
toehold-mediated nucleic acid strand displacement [84]. Severing the non-target strand
increases its conformational entropy and effectively destroys the toehold, decreasing
the rate at which the non-target strand can invade the crRNA:target strand duplex [84].
This model can also explain previous observations of cutting-dependent complex
stabilization [54,62,63] and suggests that engineering a nicking preference may improve the
efficacy of other Cas enzymes in chromatin targeting. Other potential explanations for
multiAsCas12a's enhanced CRISPRi activity include protein:DNA contacts formed after
non-target-strand nicking (Naqvi et al., 2022) and/or nicking-induced relaxation of DNA
supercoiling [85]. We have demonstrated that the effects of multiAsCas12a-KRAB's
residual DNase activity on DNA sequence contributes minimally to target gene

knockdown for typical functional genomics experimental conditions (Fig. 2F, Fig. 3B-G,
Fig. 5C, Fig. 6B, Fig. S12, and Fig. S14-S16). Nevertheless, for screens involving
strong positive selection, it may be possible for infrequent deletions to exert more
appreciable influence on screen outcome.
483
We propose that multiAsCas12a-KRAB provides new solutions to a major challenge in
combinatorial genetics: the infeasibility of surveying potentially enormous combinatorial
spaces of $\geq$3-plex genetic perturbations. Testing a single higher-order N-plex
combination also indirectly tests all or many of its constituent lower-order combinations,
for up to a total of 2^N combinations. Thus, increases in multiplexing capability potentially
yield exponential increases in search efficiency using group testing [86,87]. In group testing
(Fig. 7), a primary screen is conducted on grouped subjects (e.g. a multiplexed array of
crRNA constructs) to reduce the costs otherwise incurred by individually testing all
subjects (e.g. an individual crRNA). Our screen for CD55 enhancers instantiates this
approach by testing 22 4-plex crRNA arrays targeting 12 candidate regions, indirectly
testing 22 x 2^4 = 352 crRNA combinations in a cost-effective well-based experiment
(Fig. 5B). For this experimental objective, the grouped hits can be biologically
interpreted without further testing (Fig. 7). For other objectives, such as the analysis of
combinatorial cis-regulation at the MYC locus (Fig. 6), grouped hits can be followed by a
focused secondary screen as needed (Fig. 7). For pooled sequencing screens, the
ability to deterministically encode specific higher-order crRNA combinations in a single
array is crucial for group testing. In contrast, cloning combinatorial guide libraries by a
multiplicative and stochastic approach [17] requires testing all combinations at the onset,
and thus is incompatible with group testing. Group testing can significantly compress
the size of crRNA libraries to facilitate screens limited by assayable cell numbers. Group
testing may be combined with compressed sensing [88,89] to facilitate screens with
multidimensional phenotypic readouts [19,20,59,90–95].
506
A key parameter in group testing is the extent of potential signal dilution and/or
interference relative to individual testing. Signal dilution can arise from limiting doses of
ribonucleoprotein due to delivery format or reduction in the protein available to bind
each individual crRNA due to increased crRNA multiplexing. Signal interference can
arise if certain sequence features in one part of the crRNA array masks the activity of
individual constituent crRNAs in a dominant fashion. Despite some evidence of signal
dilution and/or interference for multiplexed crRNA arrays (Fig. 3H, Fig. 4G, and Fig.
S22), multiAsCas12a-KRAB demonstrates sufficient robustness for yielding new
biological insights into combinatorial cis-regulation at the MYC locus using 6-plex
crRNAs in high-throughput pooled screens (Fig. 6C-D). While we have focused our
optimizations to meet the stringent requirements of pooled screening formats,
multiAsCas12a also significantly lowers technical barriers to higher-order combinatorial

perturbations in array-based screening (Fig. 5B), which has recently improved in
throughput [96]. The assay format will likely influence the deliverable dose of synthetic
components and thus the upper limit of multiplexing for effective CRISPRi using
multiAsCas12a-KRAB. Improved prediction of crRNA array activities will likely further
support highly multiplexed and/or dose-limited applications, thus extending the
scalability of combinatorial genetic screens by group testing. A fully active 10-plex
crRNA array could indirectly screen up to $2^{10} = 1,024$ crRNA combinations. Another
area for improvement is the observation that a low proportion of crRNAs targeting non-
canonical PAMs show CRISPRi activity when targeted by multiAsCas12a-KRAB (Fig.
4E). Increasing the fraction of active crRNAs, especially for non-canonical PAM's, would
enable more reliable targeting with fewer crRNAs, especially in GC-rich TSS-proximal
regions.
531
While we have focused on CRISPRi applications using the KRAB domain, the discovery
and engineering of effector domains for chromatin perturbations by CRISPR-Cas is
rapidly evolving. Recent advances include new repressive effectors [97–100], activation
effectors [57,98,100,101], and combination effectors for epigenetic memory [52,102–105]. We
expect that multiAsCas12a can be flexibly combined with these and other effector
domains to support group testing for many chromatin perturbation objectives. We
envision multiAsCas12a and the group testing framework will enable elucidating and
engineering combinatorial genetic processes underlying broad areas of biology at
previously intractable scales.
541

Author Contributions
C.C.H. and L.A.G. conceived of and led the design of the overall study. C.C.H. wrote
the paper incorporating input from L.A.G., C.M.W., J.C.C., J.S., and agreement from all
authors. C.C.H. conceived of testing the R1226A mutation for CRISPRi applications,
proposed the conceptual link to group testing, and led the design, execution, and
computational analysis of all experiments, with contributions from others noted below.
L.A.G. proposed the dose of lentivirally delivered CRISPR components as a key
parameter for evaluation, interpreted results, and guided and obtained funding for the
overall study. C.M.W. designed, executed, and analyzed quantification of indel
frequencies by short-read sequencing, and contributed to cloning and flow cytometry-
based CRISPRi experiments. N.A.S. contributed to the execution of flow cytometry-
based CRISPRi experiments, including all experiments in C4-2B cells, testing of higher-
order crRNAs arrays, replicates of dose-response CRISPRi experiments, and enhancer
CRISPRi experiments at the CD55 and MYC loci. R.D. contributed to cell line
engineering and CRISPRi analysis of the DNase-dead mutant panel. Q.C. designed,
executed, and analyzed all experiments testing truncated crRNAs, with guidance and
funding obtained by J.S.. J.C.C. provided feedback on interpretations of the biophysics

literature and proposed the entropic barrier to R-loop reversal upon severance of the non-target strand. S.M. contributed to crRNA cloning and cell culture. T.O. prepared 3' RNA-seq Illumina sequencing libraries. A.A. contributed to screen data processing and analysis for 3' RNA-seq. N.T. wrote scripts in Rust for screen read mapping and counting.

Acknowledgements

We thank Brian Yu and Khushali Patel for assistance in Illumina and Nanopore sequencing; Michael Herschl for sharing a modified target-enrichment Nanopore sequencing protocol and Vince Tran for helpful discussions; Eric Chow for suggestions on Illumina sequencing of screen libraries; April Pawluk for feedback on manuscript; Gavin Knott, Ashir Borah, Garrett Wong, James Nunez, Jonathan Weissman, Howard Chang, Patrick Hsu, Silvana Konermann, and the Gilbert Lab for helpful discussions.

Funding

C.C.H. is supported by the Physician Scientist Incubator, Dept. of Pathology, Stanford University School of Medicine and NIH NHGRI (K01HG012789). Q.C. and J.S. were supported by the Mark Foundation for Cancer Research. J.C.C. is a fellow of the Helen Hay Whitney Foundation. This work was funded in part by NIH (R01HG012227) and in part by GSK through an award to the UCSF Laboratory for Genomics Research to L.A.G.. L.A.G. is funded by the Arc Institute, NIH (DP2CA239597, UM1HG012660), CRUK/NIH (OT2CA278665 and CGCATF-2021/100006), a Pew-Stewart Scholars for Cancer Research award and the Goldberg-Benioff Endowed Professorship in Prostate Cancer Translational Biology. Sequencing of CRISPR screens and 3' RNA-seq libraries was performed at the UCSF CAT, supported by UCSF PBBR, RRP IMIA, and NIH 1S10OD028511-01 grants.

Declaration of Interests

C.C.H., C.M.W., R.D. and L.A.G. have filed patent applications related to multiAsCas12a. J.S. is a scientific consultant for Treeline Biosciences. L.A.G has filed patents on CRISPRoff/on, CRISPR functional genomics and is a co-founder of Chroma Medicine.

Methods

Plasmid Design and Construction

A detailed table of constructs generated in this study will be provided as a Supplemental File with all sequences. Constructs will be made available on Addgene. Cloning was performed by Gibson Assembly of PCR amplified or commercially synthesized gene fragments (from Integrated DNA Technologies or Twist Bioscience) using NEBuilder Hifi

599   Master Mix (NEB Cat# E262), and final plasmids sequence-verified by Sanger
600   sequencing of the open reading frame and/or commercial whole-plasmid sequencing
601   service provided by Primordium.
602
603   Protein constructs components:
604
Fusion protein sequence maps are provided in Supplemental Files. To summarize,
denAsCas12a-KRAB, multiAsCas12a-KRAB, multiAsCas12a, and enAsCas12a-KRAB
open reading frames were embedded in the same fusion protein architecture consisting
of an N-terminal 6x Myc-NLS (Gier et al. 2020) and C-terminal XTEN80-KRAB-P2A-
BFP (Replogle et al. 2022). The denAsCas12a open reading frame was PCR amplified
from pCAG-denAsCas12a(E174R/S542R/K548R/D908A)-NLS(nuc)-3xHA-VPR
(RTW776) (Addgene plasmid # 107943, from [31]). AsCas12a variants described were
generated by using the denAsCas12a open reading frame as starting template and
introducing the specific mutations encoded in overhangs on PCR primers that serve as
junctions of Gibson assembly reactions. opAsCas12a[29] is available as Addgene plasmid
149723, pRG232. 6xMyc-NLS was PCR amplified from pRG232. KRAB domain
sequence from KOX1 was previously reported in [43]. The lentiviral backbone for
expressing Cas12a fusion protein constructs are expressed from an SFFV promoter
adjacent to UCOE and is a gift from Marco Jost and Jonathan Weissman, derived from
a plasmid available as Addgene 188765. XTEN80 linker sequence was taken from [52]
and was originally from [106]. For constructs used in piggyBac transposition, the open
reading frame was cloned into a piggyBac vector backbone (Addgene #133568) and
expressed from a CAG promoter. Super PiggyBac Transposase (PB210PA-1) was
purchased from System Biosciences.
624
dAsCas12a-KRABx3 open reading frame sequence is from [27], in that study encoded
within a construct referred to as SiT-ddCas12a-[Repr]. We generated SiT-ddCas12a-
[Repr] by introducing the DNase-inactivating E993A by PCR-based mutagenesis using
SiT-Cas12a-[Repr] (Addgene #133568) as template. Using Gibson Assembly of PCR
products, we inserted the resulting ddCas12a-[Repr] open reading frame in-frame with
P2A-BFP in a piggyBac vector (Addgene #133568) to enable direct comparison with
other fusion protein constructs cloned in the same vector backbone (crRNA's are
encoded on separate plasmids as described below).
633
Fusion protein constructs described in Fig. S8 were assembled by subcloning the
protein-coding sequences of AsCas12a and KRAB into a lentiviral expression vector
using the In-Fusion HD Cloning system (TBUSA). AsCas12a mutants were cloned by
mutagenesis PCR on the complete wildtype AsCas12a vector to generate the final
lentiviral expression vector.

crRNA expression constructs:

Unless otherwise specified, individual single and 3-plex crRNA constructs were cloned
into the human U6 promoter-driven expression vector pRG212 (Addgene 149722,
originally from [29]). Library1, Library2, some 3-plex and all 4-plex, 5-plex, and 6-plex *As.*
crRNA constructs were cloned into pCH67, which is derived from pRG212 by replacing
the 3' DR with the variant DR8 [28]. For constructs cloned into pCH67, the specific *As.* DR
variants were assigned to each position of the array as follows, in 5' to 3' order:
3-plex: WT DR, DR1, DR3, DR8
4-plex: WT DR, DR1, DR10, DR3, DR8
5-plex: WT DR, DR1, DR16, DR10, DR3, DR8
6-plex: WT DR, DR1, DR16, DR18, DR10, DR3, DR8
8-plex: WT DR, DR1, DR16, DR_NS1, DR17, DR18, DR10, DR3, DR8
10-plex: WT DR, DR1, DR16, DR_NS1, DR4, DR_NS2, DR17, DR18, DR10, DR3, DR8

Where the sequences of DR_NS1 and 2 were based on combining hits from the variant
DR screen from DeWeirdt et al., 2020. The sequences are DR_NS1:
aattcctcctcttggaggt, and DR_NS2: aattcctcctataggaggt.

1-plex,3-plex, 8-plex, and 10-plex crRNA constructs were cloned by annealing
complementary oligos, phosphorylation by T4 polynucleotide kinase (NEB M0201S),
and ligated with T4 DNA ligase (NEB M0202) into BsmbI site of vector backbones. 4-
plex, 5-plex and 6-plex crRNA arrays were ordered as double-stranded gene fragments
and cloned into the BsmbI site of vector backbones by Gibson Assembly.

Design of individual crRNAs
For cloning individual crRNA constructs targeting TSS's, CRISPick was used in the
enAsCas12a CRISPRi mode to design spacers targeting PAM's located within -50bp to
+300bp region around the targeted TSS. We manually selected spacers from the
CRISPick output by picking TTTV PAM-targeting spacers (except for crCD151-3, which
targets a non-canonical GTTC PAM) with the highest On-Target Efficacy Scores and
generally excluded any spacers with high off-target predictions. The same non-targeting
spacer was used throughout the individual well-based experiments and was randomly
generated and checked for absence of alignment to the human genome by BLAT [107].
The hg19 genomic coordinates for MYC enhancers are: e1 chr8:128910869-
128911521, e2 chr8:128972341-128973219, and e3 chr8:129057272-129057795. DNA
sequences from those regions were downloaded from the UCSC Genome Browser and
submitted to CRISPick. The top 3 spacers targeting TTTV PAM's for each enhancer
were picked based on CRISPick On-target Efficacy Score, having no Tier I or Tier II Bin

I predicted off-target sites, and proximity to the zenith of the ENCODE DNase
hypersensitivity signal in K562 cells.
681

Cell culture, lentiviral production, lentiviral transduction

All cell lines were cultured at 37deg. C with 5% CO2 in tissue culture incubators. K562
and C4-2B cells were maintained in RPMI-1640 (Gibco cat# 22400121) containing 25
mM HEPES, 2mM L-glutamine, and supplemented with 10% FBS (VWR), 100 units/mL
streptomycin, and 100 mg/mL penicillin. For pooled screens using K562 cells cultured in
flasks in a shaking incubator, the culture media was supplemented with 0.1% Pluronic
F-127 (Thermo Fisher P6866).
689

HEK 293T cells were cultured in media consisting of DMEM, high glucose (Gibco
11965084, containing 4.5g/mL glucose and 4mM L-glutamine) supplemented with 10%
FBS (VWR) and 100units/mL streptomycin, 100mg/mL penicillin. Adherent cells are
routinely passaged and harvested by incubation with 0.25% Trypsin-EDTA (Thermo
Fisher 25200056) at 37deg. C for 5-10min, followed by neutralization with media
containing 10% FBS.
696

Unless otherwise specified below, lentiviral particles were produced by transfecting
standard packaging vectors into HEK293T using TransIT-LT1 Transfection Reagent
(Mirus, MIR2306). At <24 hours post-transfection culture media with exchanged with
fresh media supplemented with ViralBoost (Alstem Bio, cat# VB100) at 1:500 dilution.
Viral supernatants were harvested ~48-72 hours after transfection and filtered through a
0.45 mm PVDF syringe filter and either stored in 4deg. C for use within <2 weeks or
stored in -80deg. C until use. Lentiviral infections included polybrene (8 mg/ml). MOI
was estimated from the fraction of transduced cells (based on fluorescence marker
positivity) by the following equation [108,109]: MOI = -ln(1-fraction of cells transduced).
706

For experiments described in Fig. S8, lentivirus was produced by transfecting HEK293T
cells with lentiviral vector, VSVG and psPAX2 helper plasmids using polyethylenimine.
Media was changed ~6–8 h post transfection. Viral supernatant was collected every 12
h for 5 times and passed through 0.45 μm PVDF filters. Lentivirus was added to target
cell lines with 8 μg/mL Polybrene and centrifuged at 650 × g for 25 min at room
temperature. Media was replaced 15 h post infection. Antibiotics (1 μg/mL puromycin)
was added 48 h post infection.
714

Antibody staining and flow cytometry

Antibodies used: CD55-APC (Biolegend 311312), CD81-PE (Biolegend 349506), B2M-
APC (Biolegend 316311), KIT-PE (Biolegend 313204), FOLH1-APC (Biolegend

342508), CD56(NCAM1)-APC (Invitrogen 17-0567-42). Cells were stained with
antibodies in 96-well plates, using 500g 5min at 4deg. C for centrifugation steps and
decanting in between each step. Cells were washed once with 200ul with FACS Buffer
(PBS with 1% BSA), then resuspended in 50ul of antibodies diluted at 1:100 in FACS
Buffer for 30min at 4deg. C. Then 150ul of FACS Buffer was added, followed by
centrifugation and supernatant, then washed one more time with 200ul FACS buffer,
followed by final resuspension in 200ul FACS Buffer for flow cytometry. For CRISPRi
experiments, all data points shown in figures are events first gated for single cells based
on FSC/SSC, then gated on GFP-positivity as a marker for cells successfully
transduced with crRNA construct. Flow cytometry was performed on the Attune NxT
instrument unless otherwise specified.
730
For cell fitness competition assays, the percentage of cells expressing the GFP marker
encoded on the crRNA expression vector is quantified by flow cytometry. log2 fold
change of % GFP-positive cells was calculated relative to day 2 (for experiments
targeting the Rpa3 locus in Fig. S8) or day 6 (for experiments targeting the MYC locus
in Fig. 6B). For experiments targeting the Rpa3 locus, flow cytometry was performed on
the Guava Easycyte 10 HT instrument.
737
Indel analysis by Illumina short-read sequencing
200,000 cells were collected on day 14 after crRNA transduction and genomic DNA was
isolated using NucleoSpin Blood (Macherey-Nagel, Catalog no. 740951.50). Briefly,
PCRs for loci of interest were run using Amplicon-EZ (Genewiz) partial IlluminaÒ
adapters and amplicons were processed using NucleoSpin Gel and PCR Clean-up Kit
(Macherey-Nagel, Catalog no. 740609.250). Paired end (2 x 250 bp) sequencing was
completed at GENEWIZ (Azenta Life Sciences). Raw fastq files were obtained from
GENEWIZ and aligned to reference sequences using *CRISPResso2* [110] with the
following modifications:
--quantification_window_size 12
--quantification_window_center -3
CRISPResso --fastq_r1 R1.fastq.gz --fastq_r2 R2.fastq.gz --amplicon_seq
acccgtcttgtttgtcccacccttggtgacgcagagccccagcccagaccccgcccaaagcactcatttaactggtattgcg
gagccacgaggcttctgcttactgcaactcgctccggccgctgggcgtagctgcgactcggcggagtcccggcggcgcg
tccttgttctaacccggcgcgccatgaccgtcgcgcggccgagcgtgcccgcggcgctgcccctcctcggggagctgccc
cggctgctgctgctggtgctgttgtgcctgccggccgtgtggggtgagtaggggcccggcggccggggaagcccctggg
ctgggtgggaggtccaagtcggtctctgaga -g actggtattgcggagccacgagg -wc -3 -w 12
755
For crRNA constructs in which the PAM is found on the opposite strand with respect to
the amplicon sequence (in this case, CD81) the following modifications were included:
--quantification_window_size 20

--quantification_window_center -18

CRISPResso --fastq_r1 pCH45H-CD81-array_S5_L001_R1_001.fastq.gz --fastq_r2

pCH45H-CD81-array_S5_L001_R2_001.fastq.gz --amplicon_seq

ctgcttcgcgggggacgaggggggggggctcgcgggcgggactcctggcgccccgcccccatgagctcatcaagagccgc

cgcccctggatggtgggggcgggggcgcacactttgccggaggttgggggcgatccgcctcactctttccccagcccagct

cactctccaatctgcggtcaccacccgagaccttcctgggggtcgcgcctaaaaggagcgcagactcccgccgggatgg

cccagaagctggggtgcgcgcaccctggccgtccctgcctgggagccgatctccctctcctcacccagacacgttccagc

ggaggcctcctcccagaagggctctggaggcctcgcaggagtggggatcccgcggttctgagttgg -p 3 -g

gagaccttcctgggggtcgcgcc -wc -18 -w 20

Quantification diagrams were generated in R.

For analysis of dual cutting at the *KIT* TSS, briefly, DNA was isolated using QuickExtract

DNA Solution (Lucigen) and amplicons were generated using 15 cycles of PCR to

introduce Illumina sequencing primer binding sites and 0-8 staggered bases to ensure

library diversity. After reaction clean-up using ExoSAP-IT kit (Thermo Fisher 78201), an

additional 15 cycles of PCR was used to introduce unique dual indices and Illumina P5

and P7 adaptors. Libraries were pooled and purified by SPRIselect magnetic beads

before paired-end sequencing using an Illumina MiSeq at the Arc Institute Multi-omics

Technology Center. Sequencing primer binding sites, unique dual indices (from Illumina

TruSeq kits), P5 and P7 adaptor sequences are from Illumina Adaptor Sequences

Document # 1000000002694 v16.

Reads were analyzed using CRISPRessoBatch from CRISPResso2 [110] with the following

modifications: wc -4 -w 15

CRISPRessoBatch --batch_settings batch2.batch --amplicon_seq

aagagcaggggccagacgCCGCCGGGAAGAAGCGAGACCCGGGCGGGCGCGAGGGAGG

GGAGGCGAGGAGGGGCGTGGCCGGCGCGCAGAGGGAGGGCGCTGGGAGGAGG

GGCTGCTGCTCGCCGCTCGCGGCTCTGGGGGCTCGGCTTTGCCGCGCTCGCTGCA

CTTGGGCGAGAGCTGGAACGTGGACCAGAGCTCGGATCCCATCGCAGCTACCGCG

ATGAGAGGCGCTCGCGGCGCCTGGGATTTTCTCTGCGTTCTGCTCCTACTGCTTCG

CGTCCAGACAGGTGGGACACCGCGGCTGGCACCCCGACCGTGcgactactcggcgaagcc

tgtg -p 3 -g TCTGCGTTCTGCTCCTACTGCTT -wc -4 -w 15

For dual gRNA cutting, both guides were included in the batch analysis.The total number

of insertions and deletions at each amplicon position were calculated and displayed

using the effect_vector_combined.txt output.

Frequencies of nucleotide substitutions with multiAsCas12a-KRAB targeting are

negligible and indistinguishable from sequencing error ($\leq$4.5%) observed in unmodified

K562 cells.

Simulations of indel impacts on gene expression

The fraction of reads containing indels within each specified region was subjected to

technical noise background subtraction by the fraction of reads containing indels

observed in K562 parental cells. This background-subtracted indel allelic frequency was

used to calculate per-copy deletion probabilities in Fig. S14 and Fig. S15B. For

denAsCas12a-KRAB this background-subtraction can result in a small negative value

and in those cases 0% is reported as per-copy deletion probability.

We simulated the impact of indels on gene expression under the assumption that gene

expression changes are entirely driven by indels generated by the given AsCas12a

fusion protein at the crRNA target site. A prior study reported at one Cas9 target site

that the frequency of larger (>250bp) indels is ~20% relative to the smaller (<250bp)

indels. This 20:80 ratio of unobserved-to-observed indels is very likely a high

overestimate in our case because our PCR amplicons are 340bp-382bp and thus are

expected to capture a large fraction of even the >250bp indels. Nevertheless we added

an additional 20% to the observed indel frequencies to arrive at our final estimates of

probability of the occurrence of any $\geq$1bp indels per DNA copy for a single target site.

Based on this indel probability we calculated the proportion of cell population expected

to harbor a given number of DNA copies with indels assuming indels occur

independently among DNA copies and using previously measured DNA copy number

for each genomic locus [111]. For dual targeting of the KIT locus by crKIT-2 and crKIT-3

we assume that the occurrence of a large (>250bp) unobserved deletion at one target

site precludes a deletion at the other target site. Starting from the single-cell expression

distributions obtained by flow cytometry from the non-targeting crRNA control, we

simulated the expected change in single-cell expression distribution under the

assumption that any $\geq$1bp indel in the PCR amplicon would generate a null allele

completely abolishing expression of the target gene in cis (i.e. indel in 1 out of 3 copies

would reduce expression by 33%). We refer to this as the "expected null" expression

change, which is expected to be a high overestimate of impact on expression. To better

more accurately estimate the true hypomorphic effects of indels we calculated a

"hypomorphic coefficient" defined as the ratio of the observed median expression
change vs. the expected null expression change for opAsCas12a. We multiply the
expected null expression change for all other fusion proteins by this hypomorphic
coefficient to derive an "expected hypomorph" expression change for each fusion
protein and crRNA construct combination.
837
Nanopore long-read sequencing analysis of deletion frequencies
Genomic DNA was harvested from 20 million cells using the Qiagen Genomic Tips Kit
(cat #10243). We used a custom protocol adapted from the Nanopore Cas9 Sequencing
Kit user's manual (SQK-CS9109, though this kit was not actually used) to enrich for
genomic DNA surrounding crRNA target sites for Nanopore sequencing using Kit 14
chemistry. Except where specified below, reagents are from the Nanopore Ligation
Sequencing Kit V14 (SQK-LSK114). To summarize in brief, synthetic spCas9 Alt-R
crRNAs targeting ~20kb regions surrounding target sites of crCD55-4, crCD81-1, crKIT-
2, crKIT-3, crB2M-1, and crB2M3 were designed according to instructions for the
Nanopore Cas9 Sequencing Kit and ordered from IDT. At least two guides were used to
target the (+) strand upstream of the Cas12a crRNA target sites, and at least two guides
were used to target the (-) strand downstream of the target sites. Synthetic spCas9
guides were pooled and annealed with spCas9 tracrRNA (IDT Cat # 1072532) by
heating to 95deg. C for 5min and cooling on the bench at room temperature, followed by
assembly into ribonucleoprotein complexes by incubating the following mixture at room
temperature for 30min: 10ul annealed crRNA:tracrRNA pool (100uM), 10ul 10x NEB
rCutSmart Buffer (NEB B6004S), 79.2ul nuclease-free water, and 0.8ul HiFi Cas9
(62uM, S. pyogenes HiFi Cas9 nuclease V3 IDT Cat#1081060). 5ug of genomic DNA
was dephosphorylated in a 30ul reaction volume using 15U of Quick calf intestinal
phosphatase (NEB cat #M0525) and 3ul of 10x rCutsmart Buffer (NEB B6004S),
incubated for 37deg. C for 10min, 80deg. C for 2min, and hold at 20deg. C. The entire
volume of dephosphorylated genomic DNA is then mixed with 10ul of Cas9
ribonucleoprotein, 1ul 10mM dATP, and 1ul Taq polymerase for a total of 42ul,
incubated at 37deg. C for 20min, 72 deg. C for 5min, and held at 4deg. C. The reaction
is then cleaned up using 21ul (0.5X) AMPure XP using two washes with 200ul 80%
ethanol, and eluted in 61ul of water. Adapters ligation was set up as follows: 60ul DNA
eluate, 25ul Ligation Buffer, 10ul NEBNExt Quick T4 DNA Ligase (NEB E6056S), and
5ul Ligation Adapter, and incubated for 40min at room temperature. The entire ligation
reaction was subjected to AMPure XP bead clean up using 40ul (0.4X) of AMPure
beads, two washes with Long Fragment Buffer, and eluted in 15ul Elution Buffer. Size
distribution and concentration of library was checked by Agilent Tapestation Genomic
DNA ScreenTape Analysis. ~75ng-560ng of each library per flow cell was each run on 1
or 2 flow cells on a Promethion P2 Solo, yielding ~100k reads per flow cell run over 40h-
72h.

fastq files generated by the MiniKNOW software were aligned to the ~20kb regions

(defined by the outermost Cas9 sgRNA protospacer sites flanking each targeted locus)

surrounding each crRNA target site in MiniKNOW to generate bam files. Bam files for

each sample were merged using *samtool merge* (samtools v1.6). Merged bam files

were filtered for alignments that overlap the start and end coordinates of the

protospacer region of the Cas12a crRNA using *bamtools filter -region* (bamtools v2.5.1).

Filtered bam files were loaded into the Integrative Genomics Viewer 2.17.0 for

visualization of individual read alignments. *pysamstats –fasta –type variation*

(pysamstats v1.1.2) was used to extract per base total read coverage and deletion

counts. The fraction of aligned reads harboring a deletion at each base was plotted

using custom scripts in R.

Pooled crRNA library design

For all crRNAs in Library 1 and Library 2: we excluded in the analysis spacers with the

following off-target prediction criteria using CRISPick run in the CRISPRi setting: 1) off-

target match = 'MAX' for any tier or bin, or 2) # Off-Target Tier I Match Bin I Matches >

1). The only crRNAs for which this filter was not applied are the non-targeting negative

control spacers, which do not have an associated CRISPick output. All crRNA

sequences were also filtered to exclude BsmbI sites used for cloning and >3

consecutive T's, which mimic RNA Pol III termination signal.

<u>Library 1 (single crRNA's)</u>

To design crRNA spacers targeting gene TSS's for Library 1, we used the -50bp to

+300bp regions of TSS annotations derived from capped analysis of gene expression

data and can include multiple TSS's per gene [68]. We targeted the TSS's of 559 common

essential genes from DepMap with the strongest cell fitness defects in K562 cells based

on prior dCas9-KRAB CRISPRi screen [68]. We used CRISPick with enAsCas12a

settings to target all possible PAM's (TTTV and non-canonical) in these TSS-proximal

regions. Except for the criteria mentioned in the previous paragraph, no other exclusion

criteria were applied. For the TSS-level analyses shown in Fig. 4D-E, each gene was

assigned to a single TSS targeted by the crRNA with the strongest fitness score for that

gene.

Negative controls in Library 1 fall into two categories: 1) 524 intergenic negative

controls, and 2) 445 non-targeting negative controls that do not map to the human

genome. Target sites for intergenic negative controls were picked by removing all

regions in the hg19 genome that are within 10kb of annotated ensembl genes (retrieved

from biomaRt from https://grch37.ensembl.org) or within 3kb of any ENCODE DNase
hypersensitive site (wgEncodeRegDnaseClusteredV3.bed from
http://hgdownload.cse.ucsc.edu/goldenpath/hg19/encodeDCC/wgEncodeRegDnaseClu
stered/). The remaining regions were divided into 1kb fragments. 90 such 1kb
fragments were sampled from each chromosome. Fragments containing >=20
consecutive N's were removed. The remaining sequences were submitted to CRISPick
run under CRISPRi settings. The CRISPick output was further filtered for spacers that
meet these criteria: 1) off-target prediction criteria described in the beginning of this
section, and 2) On-target Efficacy Score >= 0.5 (the rationale is to maximize
representation by likely active crRNAs to bias for revealing any potential cell fitness
effects from non-specific genotoxicity due to residual DNA cutting by multiCas12a-
KRAB), 3) mapping uniquely to the hg19 genome by Bowtie [112] using '-m 1' and
otherwise default parameters, 3) filtered once more against those whose uniquely
mapped site falls within 10kb of annotated ensembl genes or any ENCODE DNase
hypersensitive site.
927
Non-targeting negative control spacers were generated by combining 1) non-targeting
negative controls in the Humagne C and D libraries, 2) taking 20nt non-targeting
spacers from the dCas9-KRAB CRISPRi_v2 genome-wide library [68], removing the G in
the 1st position, and appending random 4-mers to the 3' end. This set of spacers were
then filtered for those that do not map to the hg19 genome using Bowtie with default
settings.
934
935 Library 2 (6-plex cRNA's)
936
Sublibrary A (84,275 constructs): Test position spacers were encoded at each position
of the 6-plex array, with remaining positions referred to as context positions and filled
with negative control spacers. Test positions encodes one of 506 intergenic negative
control spacers and 2,303 essential TSS-targeting spacers. The essential TSS-targeting
spacers were selected from among all spacers targeting PAM's within -50bp to +300bp
TSS-proximal regions of 50 common essential genes with the strongest K562 cell
fitness defect in prior dCas9-KRAB CRISPRi screen [68], and must have $\geq$0.7 CRISPick
On-target Efficacy Score. Negative control context spacers consist of 5 6-plex
combinations, 3 of these combinations consist entirely of non-targeting negative
controls and 2 of the combinations consist entirely of intergenic negative controls.
947
948 Sublibrary B (6,370 constructs): crRNA combinations targeting cis-regulatory elements
949 at the MYC locus were assembled from a subset of combinations possible from 15
950 starting spacers (3 targeting MYC TSS, 3 targeting each of 3 enhancers, and 3
951 intergenic negative control spacers). The 3 enhancer elements are described in the

subsection "Design of individual crRNAs." These 15 starting spacers were grouped into 5 3-plex combinations, each 3-plex combination exclusively targeting one of the 4 cis-regulatory elements, or consisting entirely of intergenic negative controls. Each 3-plex was then encoded in positions 1-3 of 6-plex arrays, and positions 4-6 were filled with all possible 3-plex combinations chosen from the starting 15 spacers. All 6-plex combinations were also encoded in the reverse order in the array.

All-negative control constructs (2000 constructs): 1500 6-plex combinations were randomly sampled from the intergenic negative control spacers described for Library 1. 500 6-plex combinations were randomly sampled from non-targeting negative control spacers described for Library 1.

Intergenic negative controls and non-targeting negative controls are defined the same as in Library 1.

As Library 2 was designed and cloned prior to the completion of the Library 1 screen, the majority of Library 2 contains constructs encoding for spacers in the test position that in hindsight do not produce strong phenotypes as single crRNAs in the Library 1 screen.

crRNA library construction

For Library 1, ~140 fmol of pooled oligo libraries from Twist were subjected to 10 cycles of PCR amplification using primers specific to adaptor sequences flanking the oligos and containing BsmbI sites. The PCR amplicons were cloned into a crRNA expression backbone (pCH67) by Golden Gate Assembly with ~1:1 insert:backbone ratio using ~500 fmol each. Golden Gate Assembly reaction was carried out in a 100ul reaction containing 2.5U Esp3I (Thermo ER0452) and 1000U T4 DNA Ligase in T4 DNA Ligase reaction buffer (NEB M0202L). The reaction mix was incubated for 31 cycles alternating between 37deg. C and 16deg. C for 20min at each temperature, then heat-inactivated at 65deg. C for 5min. Assembly reactions were column purified with Zymo DNA clean and concentrator-5 (Zymo D4004), eluted in 12ul of water and <7ul added to 70ul of MegaX DH10B T1R Electrocomp Cells (C640003) for electroporation using BioRad Gene Pulser Xcell Electroporator with settings 2.0kV, 200ohms, 25μF. Cells were recovered at 37deg. C for rotating for 1h in ~5ml recovery media from the MegaX DH10B T1R Electrocomp Cells kit and small volumes plated onto bacterial LB plates containing carbenicillin for quantification of colony forming units. The remaining recovery culture was inoculated directly into 200ml liquid LB media with carbenicillin and incubated in 37deg. C shaker for 12h-16h prior to harvesting for plasmid purification using ZymoPURE II Plasmid Midiprep kit (Zymo D4200). Based on the colony forming

units from the small volumes in the bacterial plates, the estimated coverage of the
library is 778x. 24 individual colonies were verified by Sanger sequencing and the
library subjected to deep sequencing as described in Illumina sequencing library
preparation. For Library 2, 915 fmol of pooled oligo libraries from Twist was subjected to
18 cycles of PCR amplification and agarose gel purification of the correctly sized band
before proceeding similarly with the remainder of the protocol as described above. The
estimated coverage of the library from colony forming units is ~60x.
999

Illumina sequencing library preparation
crRNA inserts were amplified from genomic DNA isolated from screens using 16 cycles
of first round PCR using pooled 0-8nt staggered forward and reverse primers, treated
with ExoSAP-IT (Thermo Fisher 78201.1.ML), followed by second round of PCR to
introduce Illumina unique dual indices and adaptors. Sequencing primer binding sites,
unique dual indices, P5 and P7 adaptor sequences are from Illumina Adaptor
Sequences Document # 1000000002694 v16. PCR amplicons were subject to size
selection by magnetic beads (SPRIselect, Beckman B23318) prior to sequencing on an
Illumina NovaSeq6000 using SP100 kit for Library 1 or SP500 kit for Library 2.
Sequencing of plasmid libraries were performed similarly, except 7 cycles of
amplification were each used for Round 1 and Round 2 PCR. The size distribution of
the final library was measured on an Agilent TapeStation system. We noted that even
after magnetic bead selection of Round 2 PCR-amplified Library 2 plasmid library
(colonies from which were Sanger sequencing verified) and genomic DNA from
screens, smaller sized fragments from non-specific PCR amplification during Illumina
sequencing library preparation persisted. This might contribute to the fraction of reads
that could not be mapped to our reference 6-plex array. Thus, these unmapped reads
do not necessarily reflect recombination of the crRNA library constructs, though the
latter could contribute as well.
1019
1020

Cell fitness screens
Library 1 screen: K562 cells engineered by piggyBac transposition to constitutively
express denAsCas12a-KRAB or multiAsCas12a-KRAB were transduced with lentivirally
packaged Library 1 constructs at MOI = ~0.15. Transduced cells were then selected
using 1ug/ml puromycin for 2 days, followed by washout of puromycin. On Day 6 after
transduction, initial (T0) time point was harvested, and the culture was split into 2
replicates that are separately cultured henceforth. 10 days later (T10), the final time
point was harvested (8.6 total doublings for multiAsCas12a-KRAB cells, 9.15 total
doublings for denasCas12a-KRAB cells). A cell coverage of >500x was maintained
throughout the screen. Library 2 screen: K562 cells engineered by piggyBac
transposition to constitutively express multiAsCas12a-KRAB were transduced with

lentivirally packaged Library 2 constructs at MOI = ~0.15. The screen was carried out
similarly as described for Library 1 screen, except the screen was carried out for 14
days (T14) or 13.5 total doublings and maintained at a cell coverage of >2000x
throughout. Genomic DNA was isolated using the NucleoSpin Blood XL Maxi kit
(Machery-Nagel 740950.50).
1037
1038 **Screen data processing and analysis**
1039
1040 Summary of library contents are in Fig. S18.
1041
1042 Library 1: Reads were mapped to crRNA constructs using sgcount
1043 (https://noamteyssier.github.io/sgcount/), requiring perfect match to the reference
1044 sequence.
1045
Library 2: First, reference construct sequences were created by interspersing provided
spacer and constant regions. Each construct is then given a unique construct id (CID).
Each CID is then split into R1 and R2 reference sequences, which are constructed by
taking the first three and last three spacer-construct pairs of the reference sequence
respectively. The R2 sequence is then reverse complemented for matching against the
R2 sequencing reads. Next, two hashmaps are created for the R1 and R2 spacer-
construct pairs respectively, which map the R1/R2 sequences to a set of corresponding
CIDs. Finally, for each R1/R2 sequencing pair, each k-mer (k = length of R1/R2
respective construct sequence) in the sequence is mapped against their respective
R1/R2 hashmap. If both sequencing pairs are able to be mapped to a CID set, then the
intersection of their sets is their original construct, and the total count of that CID is
incremented. We implemented the above algorithm as the `casmap constructs`
command in a package written in Rust, available at
https://github.com/noamteyssier/casmap.
1060
Starting from read counts, the remainder of analyses were performed using custom
scripts in R. Constructs that contained <1 read per million reads (RPM) aligned to the
reference library in either replicates at T0 were removed from analysis. From the
constructs that meet this read coverage threshold, a pseudocount of 1 was added for
each construct and the RPM re-calculated and used to obtain a fitness score [113] that
can be interpreted as the fractional defect in cell fitness per cell population doubling:
$\gamma = log2\left(\frac{(RPMfinal/negctrlmedianRPMfinal)}{(RPMinitial/negctrlmedianRPMinitial)}\right)/totaldoublings$,
where RPM = read count per million reads mapped to reference (initial = at T0, final = at
end of screen), negctrlmedian = median of RPM of intergenic negative control
constructs, totaldoublings = total cell population doublings in the screen. For Library 1,
data from a single T0 sample was used to calculate the fitness score for both replicates

due to an unexpected global loss of sequencing read counts for one of two originally
intended T0 replicate samples.
1074
1075 **3' RNA-seq experiment and data analysis**
1076
1077 *Experimental procedure*
1078
3' RNA-seq was performed as part of a batch processed using a QuantSeq-Pool
Sample-Barcoded 3′ mRNA-Seq Library Prep Kit for Illumina (Lexogen cat#139) in
accordance with the manufacturer's instructions. Briefly, 10 ng of each purified input
RNA was used for first strand cDNA synthesis with an oligo(dT) primer containing a
sample barcode and a unique molecular identifier. Subsequently, barcoded samples
were pooled and used for second strand synthesis and library amplification. Amplified
libraries were sequenced on an Illumina HiSeq4000 with 100 bp paired-end reads. The
QuantSeq Pool data was demultiplexed and preprocessed using an implementation of
pipeline originally provided by Lexogen (https://github.com/Lexogen-
Tools/quantseqpool_analysis). The final outputs of this step are gene level counts for all
samples (including samples from multiple projects multiplexed together).
1090
1091 *Gene level and differential expression analysis*
1092
1093 For generating scatter plots, normTransform function from DESeq2 [114] used to
1094 normalize the raw counts and then a pseudocount of 1 added, and log2-transformed.
1095 The output was plotted in R as scatter plots.
1096
For differential expression analysis, DESeq2 (version 1.34) default Wald-test was used
to compare each targeting construct (one replicate) with non-targeting samples (two
replicates). We calculated log2 transformed TPM counts and applied the threshold of
6.5 to eliminate genes with low expression. Using ggplot2, volcano plots visualized in R
are then displayed for genes with log2FoldChange above or below 2.055 and p-values
smaller than 0.01. The log2FoldChange cutoff was based on visually examining the
concordance between two replicates of untransduced controls and manually identifying
a threshold below which the log2FoldChange are poorly correlated between the
replicates of the untransduced control.
1106
1107 *Off-target analysis of spacers*
1108
To evaluate potential off-target effect of spacers, we used the crisprVerse (version
1.0.0) [115] and

crisprBowtie (version 1.2.0) together with other R packages including GenomicRanges
(version 1.50) [116] and tidyverse (version 1.3.2) [117]. First, we defined dictionary of
spacers as "TCCTCCAGCATCTTCCACATTCA":"HBG-2",
"TTCTTCATCCCTAGCCAGCCGCC":"HBG-3",
"CTTAGAAGGTTACACAGAACCAG":"HS2-1",
"TGTGTAACCTTCTAAGCAAACCT":"HS2-2",
"AGGTGGAGTTTTAGTCAGGTGGT":"HS2-3",
"ATTAACTGATGCGTAAGGAGATC":"NT-3". Then, `runCrisprBowtie` function used
with these parameters: `crisprNuclease` as `enAsCas12a`, `n_mismatches` equal 3,
`canonical` equal FALSE, and `bowtie_index` as a path to folder including pre-indexed
hg38 reference genome. Thus, the results from this step allows us to assess our
previously designed spacers and annotate potential off-target loci in the human
genome. To annotate results, we used reference annotation GENCODE (version 34) [118]
and we defined pam_site +/- 2500 bp for each predicted off-target to overlap them with
matched transcription start sites (TSS) +/- 1000 bp of all annotated genes. Results
shown as annotated tables.
1127
RT-qPCR
For the CRISPRi experiments targeting the HBG TSS or HS2 enhancer, K562 cells
engineered (by lentiviral transduction at MOI ~5) for constitutive expression of
multiAsCas12a-KRAB were transduced with crRNAs and sorted, followed by
resuspension of ~200k to 1 million cells in 300ul RNA Lysis Buffer from the Quick-RNA
Miniprep Kit (Zymo R1055) and stored in -70deg. C. RNA isolation was performed
following the kit's protocols, including on-column DNase I digestion. 500ng of RNA was
used as input for cDNA synthesis primed by random hexamers using the RevertAid RT
Reverse Transcription Kit (Thermo fisher K1691), as per manufacturer's instructions.
cDNA was diluted 1:4 with water and 2ul used as template for qPCR using 250nM
primers using the SsoFast EvaGreen Supermix (BioRad 1725200) on an Applied
Biosystems ViiA 7 Real Time PCR System. Data was analyzed using the ddCT method,
normalized to GAPDH and no crRNA sample as reference.
1141
Transient transfection experiments
For co-transfection experiments, transfections were performed similar to prior study [27].
Briefly, the day before transfection, 100,000 HEK293T cells were seeded into wells of a
24 well plate. The following day, we transiently transfected 0.6µg of each protein
construct and 0.3µg gRNA construct per well (in duplicate) in Mirus TransIT-LT1
transfection reagent according to manufacturer's instructions. Mixtures were incubated
at room temperature for 30 minutes and then added in dropwise fashion into each well.
24 hours after transfection, cells were replenished with fresh media. 48 hours after
transfection, BFP and GFP positive cells (indicative of successful delivery of protein and

crRNA constructs) were sorted (BD FACSAria Fusion) and carried out for subsequent
flow-cytometry experiments.
1153

1154   **Data access and availability**
1155   Tables of all sequence and read counts are included as Supplementary Files. Plasmids
1156   will be made available on Addgene. Engineered cell lines will be made available upon
1157   request.
1158
1159

References

1. Wong, A. S. L., Choi, G. C. G. & Lu, T. K. Deciphering Combinatorial Genetics. *Annu. Rev. Genet.* **50**, 515–538 (2016).

2. Costanzo, M. *et al.* Global Genetic Networks and the Genotype-to-Phenotype Relationship. *Cell* **177**, 85–100 (2019).

3. Domingo, J., Baeza-Centurion, P. & Lehner, B. The Causes and Consequences of Genetic Interactions (Epistasis). *Annu. Rev. Genomics Hum. Genet.* **20**, 433–460 (2019).

4. Takahashi, K. & Yamanaka, S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* **126**, 663–676 (2006).

5. Ewen-Campen, B., Mohr, S. E., Hu, Y. & Perrimon, N. Accessing the Phenotype Gap: Enabling Systematic Investigation of Paralog Functional Complexity with CRISPR. *Dev. Cell* **43**, 6–9 (2017).

6. Kvon, E. Z., Waymack, R., Gad, M. & Wunderlich, Z. Enhancer redundancy in development and disease. *Nat. Rev. Genet.* **22**, 324–336 (2021).

7. Blayney, J. *et al.* Super-enhancers require a combination of classical enhancers and novel facilitator elements to drive high levels of gene expression. *bioRxiv* 2022.06.20.496856 (2022) doi:10.1101/2022.06.20.496856.

8. Osterwalder, M. *et al.* Enhancer redundancy provides phenotypic robustness in mammalian development. *Nature* **554**, 239–243 (2018).

9. Blobel, G. A., Higgs, D. R., Mitchell, J. A., Notani, D. & Young, R. A. Testing the super-enhancer concept. *Nat. Rev. Genet.* **22**, 749–755 (2021).

10. Kuzmin, E. *et al.* Systematic analysis of complex genetic interactions. *Science* **360**,

(2018).

11. Domingo, J., Diss, G. & Lehner, B. Pairwise and higher-order genetic interactions

during the evolution of a tRNA. *Nature* **558**, 117–121 (2018).

12. Taylor, M. B. & Ehrenreich, I. M. Genetic interactions involving five or more genes

contribute to a complex trait in yeast. *PLoS Genet.* **10**, e1004324 (2014).

13. Taylor, M. B. & Ehrenreich, I. M. Transcriptional Derepression Uncovers Cryptic

Higher-Order Genetic Interactions. *PLoS Genet.* **11**, e1005606 (2015).

14. Celaj, A. *et al.* Highly Combinatorial Genetic Interaction Analysis Reveals a Multi-

Drug Transporter Influence Network. *Cell Syst* **10**, 25–38.e10 (2020).

15. Doench, J. G. Am I ready for CRISPR? A user's guide to genetic screens. *Nat.*

Rev. Genet. **19**, 67–80 (2018).

16. Przybyla, L. & Gilbert, L. A. A new era in functional genomics screens. *Nat. Rev.*

Genet. **23**, 89–103 (2022).

17. Zhou, P. *et al.* A Three-Way Combinatorial CRISPR Screen for Analyzing

Interactions among Druggable Targets. *Cell Rep.* **32**, 108020 (2020).

18. Wong, A. S. L., Choi, G. C. G., Cheng, A. A., Purcell, O. & Lu, T. K. Massively

parallel high-order combinatorial genetics in human cells. *Nat. Biotechnol.* **33**, 952–

961 (2015).

19. Adamson, B. *et al.* A Multiplexed Single-Cell CRISPR Screening Platform Enables

Systematic Dissection of the Unfolded Protein Response. *Cell* **167**, 1867–1882.e21

(2016).

20. Adamson, B., Norman, T. M., Jost, M. & Weissman, J. S. Approaches to maximize

sgRNA-barcode coupling in Perturb-seq screens. *bioRxiv* 298349 (2018)

doi:10.1101/298349.

21. Sack, L. M., Davoli, T., Xu, Q., Li, M. Z. & Elledge, S. J. Sources of Error in

Mammalian Genetic Screens. *G3* **6**, 2781–2790 (2016).

22. Basu, V. P. *et al.* Strand transfer events during HIV-1 reverse transcription. *Virus*

Res. **134**, 19–38 (2008).

23. Fonfara, I., Richter, H., Bratovič, M., Le Rhun, A. & Charpentier, E. The CRISPR-

associated DNA-cleaving enzyme Cpf1 also processes precursor CRISPR RNA.

Nature **532**, 517–521 (2016).

24. Zetsche, B. *et al.* Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-

Cas system. *Cell* **163**, 759–771 (2015).

25. Zetsche, B. *et al.* Multiplex gene editing by CRISPR-Cpf1 using a single crRNA

array. *Nat. Biotechnol.* **35**, 31–34 (2017).

26. Breinig, M. *et al.* Multiplexed orthogonal genome editing and transcriptional

activation by Cas12a. *Nat. Methods* **16**, 51–54 (2019).

27. Campa, C. C., Weisbach, N. R., Santinha, A. J., Incarnato, D. & Platt, R. J.

Multiplexed genome engineering by Cas12a and CRISPR arrays encoded on single

transcripts. *Nat. Methods* **16**, 887–893 (2019).

28. DeWeirdt, P. C. *et al.* Optimization of AsCas12a for combinatorial genetic screens

in human cells. *Nat. Biotechnol.* **39**, 94–104 (2021).

29. Gier, R. A. *et al.* High-performance CRISPR-Cas12a genome editing for

combinatorial genetic screening. *Nat. Commun.* **11**, 3455 (2020).

30. DeWeirdt, P. C. *et al.* Accounting for small variations in the tracrRNA sequence

improves sgRNA activity predictions for CRISPR screening. *bioRxiv*

2022.06.27.497780 (2022) doi:10.1101/2022.06.27.497780.

31. Kleinstiver, B. P. *et al.* Engineered CRISPR-Cas12a variants with increased

activities and improved targeting ranges for gene, epigenetic and base editing. *Nat.*

Biotechnol. **37**, 276–282 (2019).

32. Gonatopoulos-Pournatzis, T. *et al.* Genetic interaction mapping and exon-resolution

functional genomics with a hybrid Cas9-Cas12a platform. *Nat. Biotechnol.* **38**, 638–

648 (2020).

33. Chow, R. D. *et al.* In vivo profiling of metastatic double knockouts through CRISPR-

Cpf1 screens. *Nat. Methods* **16**, 405–408 (2019).

34. Liu, J. *et al.* Pooled library screening with multiplexed Cpf1 library. *Nat. Commun.*

10, 3144 (2019).

35. Dede, M., McLaughlin, M., Kim, E. & Hart, T. Multiplex enCas12a screens detect

functional buffering among paralogs otherwise masked in monogenic Cas9

knockout screens. *Genome Biol.* **21**, 262 (2020).

36. Anvar, N. E. *et al.* Efficient gene knockout and genetic interactions: the IN4MER

CRISPR/Cas12a multiplex knockout platform. *bioRxiv* 2023.01.03.522655 (2023)

doi:10.1101/2023.01.03.522655.

37. Meyers, R. M. *et al.* Computational correction of copy number effect improves

specificity of CRISPR-Cas9 essentiality screens in cancer cells. *Nat. Genet.* **49**,

1779–1784 (2017).

38. Aguirre, A. J. *et al.* Genomic Copy Number Dictates a Gene-Independent Cell

Response to CRISPR/Cas9 Targeting. *Cancer Discov.* **6**, 914–929 (2016).

39. Chen, Z. *et al.* In vivo CD8+ T cell CRISPR screening reveals control by Fli1 in

infection and cancer. *Cell* **184**, 1262–1280.e22 (2021).

40. Ihry, R. J. *et al.* p53 inhibits CRISPR-Cas9 engineering in human pluripotent stem

cells. *Nat. Med.* **24**, 939–946 (2018).

41. Bowden, A. R. *et al.* Parallel CRISPR-Cas9 screens clarify impacts of p53 on

screen performance. *Elife* **9**, (2020).

42. Haapaniemi, E., Botla, S., Persson, J., Schmierer, B. & Taipale, J. CRISPR-Cas9

genome editing induces a p53-mediated DNA damage response. *Nat. Med.* **24**,

927–930 (2018).

43. Gilbert, L. A. *et al.* CRISPR-mediated modular RNA-guided regulation of

transcription in eukaryotes. *Cell* **154**, 442–451 (2013).

44. Gilbert, L. A. *et al.* Genome-Scale CRISPR-Mediated Control of Gene Repression

and Activation. *Cell* **159**, 647–661 (2014).

45. Konermann, S. *et al.* Genome-scale transcriptional activation by an engineered

CRISPR-Cas9 complex. *Nature* **517**, 583–588 (2015).

46. Ren, X. *et al.* Parallel characterization of cis-regulatory elements for multiple genes

using CRISPRpath. *Sci Adv* **7**, eabi4360 (2021).

47. Tycko, J. *et al.* Mitigation of off-target toxicity in CRISPR-Cas9 screens for essential

non-coding elements. *Nat. Commun.* **10**, 4063 (2019).

48. Yao, D. *et al.* Multi-center integrated analysis of non-coding CRISPR screens.

bioRxiv 2022.12.21.520137 (2022) doi:10.1101/2022.12.21.520137.

49. Ecco, G., Imbeault, M. & Trono, D. KRAB zinc finger proteins. *Development* **144**,

2719–2729 (2017).

50. Guo, L. Y. *et al.* Multiplexed genome regulation in vivo with hyper-efficient Cas12a.

Nat. Cell Biol. **24**, 590–600 (2022).

51. Liu, Y. *et al.* Engineering cell signaling using tunable CRISPR-Cpf1-based

transcription factors. *Nat. Commun.* **8**, 2095 (2017).

52. Nuñez, J. K. *et al.* Genome-wide programmable transcriptional memory by

CRISPR-based epigenome editing. *Cell* **184**, 2503–2519.e17 (2021).

53. O'Geen, H. *et al.* dCas9-based epigenome editing suggests acquisition of histone

methylation is not sufficient for target gene repression. *Nucleic Acids Res.* **45**,

9901–9916 (2017).

54. Cofsky, J. C. *et al.* CRISPR-Cas12a exploits R-loop asymmetry to form double-

strand breaks. *Elife* **9**, (2020).

55. Yamano, T. *et al.* Crystal Structure of Cpf1 in Complex with Guide RNA and Target

DNA. *Cell* **165**, 949–962 (2016).

56. Esmaeili Anvar, N. *et al.* Combined genome-scale fitness and paralog synthetic

lethality screens with just 44k clones: the IN4MER CRISPR/Cas12a multiplex

knockout platform. *bioRxiv* 2023.01.03.522655 (2023).

57. Griffith, A. L. *et al.* Optimization of Cas12a for multiplexed genome-scale

transcriptional activation. *Cell Genomics* **0**, (2023).

58. Replogle, J. M. *et al.* Maximizing CRISPRi efficacy and accessibility with dual-

sgRNA libraries and optimal effectors. *bioRxiv* 2022.07.13.499814 (2022)

doi:10.1101/2022.07.13.499814.

59. Datlinger, P. *et al.* Pooled CRISPR screening with single-cell transcriptome

readout. *Nat. Methods* **14**, 297–301 (2017).

60. Zhang, L. *et al.* AsCas12a ultra nuclease facilitates the rapid generation of
therapeutic cell medicines. *Nat. Commun.* **12**, 3908 (2021).

61. Naqvi, M. M., Lee, L., Montaguth, O. E. T., Diffin, F. M. & Szczelkun, M. D.
CRISPR-Cas12a-mediated DNA clamping triggers target-strand cleavage. *Nat.*
Chem. Biol. **18**, 1014–1022 (2022).

62. Singh, D. *et al.* Real-time observation of DNA target interrogation and product
release by the RNA-guided endonuclease CRISPR Cpf1 (Cas12a). *Proc. Natl.*
Acad. Sci. U. S. A. **115**, 5444–5449 (2018).

63. Knott, G. J. *et al.* Broad-spectrum enzymatic inhibition of CRISPR-Cas12a. *Nat.*
Struct. Mol. Biol. **26**, 315–321 (2019).

64. Swarts, D. C. & Jinek, M. Mechanistic Insights into the cis- and trans-Acting DNase
Activities of Cas12a. *Mol. Cell* **73**, 589–600.e4 (2019).

65. Zhang, L. *et al.* Conformational Dynamics and Cleavage Sites of Cas12a Are
Modulated by Complementarity between crRNA and DNA. *iScience* **19**, 492–503
(2019).

66. Jeon, Y. *et al.* Direct observation of DNA target searching and cleavage by
CRISPR-Cas12a. *Nat. Commun.* **9**, 2777 (2018).

67. Qi, L. S. *et al.* Repurposing CRISPR as an RNA-guided platform for sequence-
specific control of gene expression. *Cell* **152**, 1173–1183 (2013).

68. Horlbeck, M. A. *et al.* Compact and highly active next-generation libraries for
CRISPR-mediated gene repression and activation. *Elife* **5**, (2016).

69. Horlbeck, M. A. *et al.* Nucleosomes impede Cas9 access to DNA in vivo and in
vitro. *Elife* **5**, (2016).

70. Kim, H. K. *et al.* Deep learning improves prediction of CRISPR-Cpf1 guide RNA

activity. *Nat. Biotechnol.* **36**, 239–241 (2018).

71. ENCODE Project Consortium *et al.* Expanded encyclopaedias of DNA elements in

the human and mouse genomes. *Nature* **583**, 699–710 (2020).

72. Li, K. *et al.* Interrogation of enhancer function by enhancer-targeting CRISPR

epigenetic editing. *Nat. Commun.* **11**, 485 (2020).

73. Dho, S. H., Lim, J. C. & Kim, L. K. Beyond the Role of CD55 as a Complement

Component. *Immune Netw.* **18**, e11 (2018).

74. Hillmen, P. *et al.* Effect of eculizumab on hemolysis and transfusion requirements in

patients with paroxysmal nocturnal hemoglobinuria. *N. Engl. J. Med.* **350**, 552–559

(2004).

75. Egan, E. S. *et al.* Malaria. A forward genetic screen identifies erythrocyte CD55 as

essential for Plasmodium falciparum invasion. *Science* **348**, 711–714 (2015).

76. Shakya, B., Patel, S. D., Tani, Y. & Egan, E. S. Erythrocyte CD55 mediates the

internalization of Plasmodium falciparum parasites. *Elife* **10**, (2021).

77. Creyghton, M. P. *et al.* Histone H3K27ac separates active from poised enhancers

and predicts developmental state. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 21931–

21936 (2010).

78. Fulco, C. P. *et al.* Activity-by-contact model of enhancer-promoter regulation from

thousands of CRISPR perturbations. *Nat. Genet.* **51**, 1664–1669 (2019).

79. Cheng, J. *et al.* Regulatory Architecture of the RCA Gene Cluster Captures an

Intragenic TAD Boundary, CTCF-Mediated Chromatin Looping and a Long-Range

Intergenic Enhancer. *Front. Immunol.* **13**, 901747 (2022).

80. Fulco, C. P. *et al.* Systematic mapping of functional enhancer-promoter

connections with CRISPR interference. *Science* **354**, 769–773 (2016).

81. Reilly, S. K. *et al.* Direct characterization of cis-regulatory elements and functional

dissection of complex genetic associations using HCR–FlowFISH. *Nat. Genet.* **53**,

1166–1176 (2021).

82. Lin, X. *et al.* Nested epistasis enhancer networks for robust genome regulation.

Science **377**, 1077–1085 (2022).

83. Ma, E. *et al.* Improved genome editing by an engineered CRISPR-Cas12a. *Nucleic*

Acids Res. **50**, 12689–12701 (2022).

84. Srinivas, N. *et al.* On the biophysics and kinetics of toehold-mediated DNA strand

displacement. *Nucleic Acids Res.* **41**, 10641–10658 (2013).

85. Baranello, L., Levens, D., Gupta, A. & Kouzine, F. The importance of being

supercoiled: how DNA mechanics regulate dynamic processes. *Biochim. Biophys.*

Acta **1819**, 632–638 (2012).

86. Dorfman, R. The Detection of Defective Members of Large Populations. *aoms* **14**,

436–440 (1943).

87. Du, D. *Combinatorial Group Testing and Its Applications*. (World Scientific, 1993).

88. Yao, D. *et al.* Compressed Perturb-seq: highly efficient screens for regulatory

circuits using random composite perturbations. *bioRxiv* 2023.01.23.525200 (2023).

89. Cleary, B. & Regev, A. The necessity and power of random, under-sampled

experiments in biology. *arXiv [q-bio.QM]* (2020).

90. Dixit, A. *et al.* Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell

RNA Profiling of Pooled Genetic Screens. *Cell* **167**, 1853–1866.e17 (2016).

91. Wessels, H.-H. *et al.* Efficient combinatorial targeting of RNA transcripts in single

cells with Cas13 RNA Perturb-seq. *Nat. Methods* **20**, 86–94 (2023).

92. Schraivogel, D. *et al.* Targeted Perturb-seq enables genome-scale genetic screens

in single cells. *Nat. Methods* **17**, 629–635 (2020).

93. Replogle, J. M. *et al.* Combinatorial single-cell CRISPR screens by direct guide

RNA capture and targeted sequencing. *Nat. Biotechnol.* **38**, 954–961 (2020).

94. Norman, T. M. *et al.* Exploring genetic interaction manifolds constructed from rich

single-cell phenotypes. *Science* **365**, 786–793 (2019).

95. Feldman, D. *et al.* Optical Pooled Screens in Human Cells. *Cell* **179**, 787–799.e17

(2019).

96. Yin, J.-A. *et al.* Robust and Versatile Arrayed Libraries for Human Genome-Wide

CRISPR Activation, Deletion and Silencing. *bioRxiv* 2022.05.25.493370 (2022)

doi:10.1101/2022.05.25.493370.

97. Alerasool, N., Segal, D., Lee, H. & Taipale, M. An efficient KRAB domain for

CRISPRi applications in human cells. *Nat. Methods* **17**, 1093–1096 (2020).

98. Mukund, A. X. *et al.* High-throughput functional characterization of combinations of

transcriptional activators and repressors. *Cell Syst* **0**, (2023).

99. Replogle, J. M. *et al.* Maximizing CRISPRi efficacy and accessibility with dual-

sgRNA libraries and optimal effectors. *Elife* **11**, (2022).

100.DelRosso, N. *et al.* Large-scale mapping and mutagenesis of human transcriptional

effector domains. *Nature* **616**, 365–372 (2023).

101.Alerasool, N., Leng, H., Lin, Z.-Y., Gingras, A.-C. & Taipale, M. Identification and

functional characterization of transcriptional activators in human cells. *Mol. Cell* **82**,

677–695.e7 (2022).

102. Van, M. V., Fujimori, T. & Bintu, L. Nanobody-mediated control of gene expression

and epigenetic memory. *Nat. Commun.* **12**, 537 (2021).

103. Nakamura, M., Ivec, A. E., Gao, Y. & Qi, L. S. Durable CRISPR-Based Epigenetic

Silencing. *BioDesign Research* **2021**, (2021).

104. Amabile, A. *et al.* Inheritable Silencing of Endogenous Genes by Hit-and-Run

Targeted Epigenetic Editing. *Cell* **167**, 219–232.e14 (2016).

105. O'Geen, H., Tomkova, M., Combs, J. A., Tilley, E. K. & Segal, D. J. Determinants of

heritable gene silencing for KRAB-dCas9 + DNMT3 and Ezh2-dCas9 + DNMT3 hit-

and-run epigenome editing. *Nucleic Acids Res.* **50**, 3239–3253 (2022).

106. Schellenberger, V. *et al.* A recombinant polypeptide extends the in vivo half-life of

peptides and proteins in a tunable manner. *Nat. Biotechnol.* **27**, 1186–1190 (2009).

107. Kent, W. J. BLAT--the BLAST-like alignment tool. *Genome Res.* **12**, 656–664

(2002).

108. Shabram, P. & Aguilar-Cordova, E. Multiplicity of infection/multiplicity of confusion.

Mol. Ther. **2**, 420–421 (2000).

109. Ellis, E. L. & Delbrück, M. THE GROWTH OF BACTERIOPHAGE. *J. Gen. Physiol.*

22, 365–384 (1939).

110. Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing

sequence analysis. *Nat. Biotechnol.* **37**, 224–226 (2019).

111. Zhou, B. *et al.* Comprehensive, integrated, and phased whole-genome analysis of

the primary ENCODE cell line K562. *Genome Res.* **29**, 472–484 (2019).

112. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-

efficient alignment of short DNA sequences to the human genome. *Genome Biol.*

10, R25 (2009).

113. Kampmann, M., Bassik, M. C. & Weissman, J. S. Integrated platform for genome-

wide screening and construction of high-density genetic interaction maps in

mammalian cells. *Proc. Natl. Acad. Sci. U. S. A.* **110**, E2317–26 (2013).

114. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

115. Hoberecht, L., Perampalam, P., Lun, A. & Fortin, J.-P. A comprehensive

Bioconductor ecosystem for the design of CRISPR guide RNAs across nucleases

and technologies. *Nat. Commun.* **13**, 6568 (2022).

116. Lawrence, M. *et al.* Software for computing and annotating genomic ranges. *PLoS*

Comput. Biol. **9**, e1003118 (2013).

117. Wickham, H. *et al.* Welcome to the tidyverse. *J. Open Source Softw.* **4**, 1686

(2019).

118. Frankish, A. *et al.* GENCODE 2021. *Nucleic Acids Res.* **49**, D916–D923 (2021).

Table of Contents for Main and Supplemental Figures

Figure 1 – dAsCas12a-KRAB variants are dose-limited and weak in CRISPRi activity when delivered lentivirally, despite incorporating state-of-the-art optimizations.

A) Schematic for assaying CRISPRi activity of Cas12a constructs using lentivirally transduced single-plex or 3-plex crRNAs targeting cell surface marker genes assayed by antibody staining and flow cytometry.

B) K562 cells constitutively expressing dAsCas12a-KRABx3 (Campa et al., 2019) were lentivirally transduced with single crRNAs targeting CD55, CD81, B2M, KIT, or a non-targeting crRNA, and assayed by flow cytometry 6 days after crRNA transduction.

C) A panel of Cas12a variants harboring combinations of mutations are tested using crCD55-4 and crCD81-1 using the fusion protein domain architecture shown. Both Cas12a fusion protein and crRNA constructs are delivered by lentiviral transduction. D908A is a mutation in the RuvC catalytic triad that renders Cas12a DNase-inactive (Yamano et al., 2016; Zetsche et al., 2015). Other mutations are described in detail in the main text. Shown are single-cell distributions of target gene expression assayed by flow cytometry 6 days after crRNA transduction for one of 3 independent replicates. One-sided Wilcoxon rank-sum test was performed comparing the single-cell distributions of the non-targeting control vs. the corresponding targeting crRNA; asterisk indicates p<0.01. Additional replicates and results for additional crRNA constructs (up to 3-plex crRNA constructs) are summarized in Fig. S6.

D) Analysis of CD81 knockdown in cells lentivirally transduced with denAsCas12a-KRAB protein construct at MOI ∼1 vs. MOI ∼5, while maintaining constant crRNA MOI (<0.74) for each crRNA construct. CD81 expression was assayed by flow cytometry 6 days after crRNA transduction. Shown are single-cell distributions of target gene expression knockdown as a percentage of non-targeting control for one of 3-6 biological replicates for each crRNA construct. Median and interquartile range are shown for each distribution. Percentage of cells below the 5th percentile (dashed line) of non-targeting crRNA are shown. One-sided Wilcoxon rank-sum test was performed on single-cell distributions for this replicate; asterisk indicates p<0.01. Summaries of all replicates shown in Fig. S7A.

E) Similar to D, but maintaining constant denAsCas12a-KRAB protein construct MOI at ∼5, while crRNA MOI is changed from high to low as indicated. CD81 expression was assayed by flow cytometry 10 days after crRNA transduction. Shown are single-cell distributions of CD81 knockdown for one of two biological replicates. One-sided Wilcoxon rank-sum test was performed on single-cell distributions for each replicate; asterisk indicates p<0.01. Additional replicate shown in Fig. S7B.

Figure 2 – multiAsCas12a-KRAB (R1226A/E174R/S542R/K548R), an engineered variant that favors a nicked DNA intermediate, substantially improves lentivirally delivered CRISPRi activity.

A) Model of Cas12a DNA binding and cleavage states for wildtype DNase vs. R1226A mutant based on prior in vitro studies as detailed in main text. Sizes of arrows qualitatively reflect reaction rates.

B) Comparison of denCas12a-KRAB (D908A/E174R/S542R/K548R) vs. multiAsCas12a-KRAB (R1226A/E174R/S542R/K548R) in CRISPRi knockdown of CD81 using crCD81-1. CD81 expression assayed by flow cytometry 10 days after crRNA transduction. Left panel: Holding crRNA MOI constant at ~3 while testing protein MOI ~1 vs. ~5. Right panel: Holding protein MOI constant at ~5 while testing crRNA MOI at ~3 vs. ~0.5. Asterisks indicate p <0.01 for one-sided Wilcoxon rank-sum test of single-cell distributions. Percentage of cells below the 5th percentile of non-targeting crRNA control (dashed line) is shown for each condition. One biological replicate is shown for each condition; additional replicates shown in Fig. S9.

C) Comparison of CD81 knockdown by lentivirally delivered denAsCas12a-KRAB vs. multiAsCas12a-KRAB at protein MOI ~1 vs. ~5 across a panel of single and 3-plex crRNA constructs, while holding constant crRNA MOI for each paired fusion protein comparison for each crRNA construct. Dashed gray line indicates 5th percentile of non-targeting crRNA control. crRNA MOI indicated by color scale. Lines connect paired replicates. One-sided Wilcoxon rank-sum tests were performed on single-cell distributions for each replicate, and asterisk denotes p<0.01 for all paired replicates within each condition. Dots indicate flow cytometry measurement 10 days after crRNA transduction; triangles indicate flow cytometry measurement 16 days after crRNA transduction.

D) Same as C but showing scatter plot of CD55-APC and CD81-PE antibody co-staining signals on flow cytometry performed 16 days after transduction of the indicated crRNA constructs in K562 cells lentivirally transduced with denAsCas12a-KRAB vs. multiAsCas12a-KRAB at protein MOI ~5. Quadrants drawn based on the 5th percentile of non-targeting controls and the percentage of cells in each quadrant denoted.

E) K562 cells piggyBac-engineered to constitutively express denAsCas12a-KRAB or multiAsCas12a-KRAB were transduced with the indicated crRNA constructs, followed by measurement of CD151 expression by antibody staining and flow cytometry 13 days after crRNA transduction. Median CD151 expression knockdown relative to non-targeting control is shown for each individual replicate. Dashed gray line indicates 5th percentile of non-targeting crRNA control. crRNA MOI indicated by color scale. One-sided Wilcoxon rank-sum test was performed for single-cell expression distributions comparing denAsCas12a-KRAB vs. multiAsCas12a-KRAB for each replicate; asterisk indicates p<0.001 for all replicate-level comparisons.

F) Indel quantification from PCR amplicons surrounding target sites of crCD81-1 and crCD55-4 in cells lentivirally transduced at protein MOI ~5 for denAsCas12a-KRAB and multiAsCas12a-KRAB. Cells lentivirally transduced with opAsCas12a (DNase fully active) are shown for comparison. Percent of reads containing indels at each base position within the amplicon is plotted, with labels indicating maximum indel frequency observed across all bases within the amplicon.

Figure 3 – multiAsCas12a-KRAB enables multi-gene CRISPRi perturbations using higher-order arrayed crRNA lentiviral constructs
A) Schematic for higher-order crRNA expression constructs. 23nt spacers are interspersed by 19nt direct repeat variants (DeWeirdt et al., 2020) uniquely assigned to each position within the array. **B)** Flow cytometry analysis of CD81 expression knockdown by antibody staining 6 days after transduction of the indicated lentiviral crRNA constructs in K562 cells engineered to constitutively express the specified fusion protein construct. Shown are averages of median single-cell expression knockdown from 2-5 biological replicates for each crRNA construct, with error bars indicating SEM. One-sided Wilcoxon rank-sum test was performed for differences in single-cell expression distributions for each fusion protein against multiAsCas12a-KRAB for each individual replicate. Asterisk indicates p < 0.01 for all replicates for a given pairwise comparison. **C)** Same as B, but shown for KIT expression knockdown. **D)** Indel quantification for the indicated fusion protein constructs using a 6-plex crRNA construct encoding crKIT-2 and crKIT-3 that target opposite strands at sites spaced 112 bp apart near the KIT TSS. Following crRNA transduction, cells were sorted on day 3 for GFP marker on the crRNA construct, and the 340bp genomic region surrounding both crRNA binding sites was PCR amplified from cell lysates harvested 15 days after crRNA transduction. The maximum percentages of reads containing indels overlapping any base position within each of the demarcated regions (region A, region B, region C) are shown. **E)** Comparison of the indicated fusion protein constructs in dual CD55 and CD81 CRISPRi knockdown 10 days after lentiviral transduction of a 6-plex crRNA construct by flow cytometry. Shown are log10 fluorescence intensity for each antibody stain and the percentages of cells in each quadrant, defined by the 5th percentile of non-targeting crRNA for each fluorescence signal, are indicated. **F)** Summary of the same experiment in E for a larger panel of crRNA constructs, showing the percentage of cells with successful double-knockdown of CD55 and CD81 (e.g. same gating strategy as bottom left quadrant in E). 2-6 biological replicates are shown as individual data points and summarized by the mean and SEM as error bars. Two-sample chi-square test was used compare the proportion of cells with double-knockdown between multiAsCas12a-KRAB and each of the other fusion protein constructs; asterisk indicates p<0.01 for all replicates. **G)** Analogous to F, except triple knockdown of CD55, KIT, and CD81 was quantified by the percentage of cells that are below the 5th percentile along all 3 dimensions on day 33 after transduction of crRNA constructs. 2 biological replicates are shown as individual data points. Two-sample chi-square test was used compare the proportion of cells with double-knockdown between multiAsCas12a-KRAB and each of the other fusion protein constructs; asterisk indicates p<0.01 for all replicates. **H)** Gene expression knockdown by multiAsCas12a-KRAB using 6-plex, 8-plex and 10-plex crRNA array constructs was measured by flow cytometry 10-11 days after lentiviral transduction of crRNA constructs. Shown are median gene expression knockdown averaged from 2-4 biological replicates, with error bars denoting SEM. CRISPRi activities of crFOLH1-1, crCD151-3 and crHBG-3 were not assayed in this experiment, but these spacers are active when encoded as individual crRNAs as shown in Fig. S4B, Fig. 2E, and Fig. 5A, respectively. All protein constructs shown in A-H were delivered by piggyBac transposition into K562 cells and sorted for the same expression level of the P2A-BFP marker (except opAsCas12a was delivered by lentiviral transduction and selected for by puromycin-resistance marker).

Figure 4 – multiAsCas12a-KRAB enables TSS-targeting pooled CRISPRi screens, including with 6-plex crRNA arrays.

A) Design of Library 1 consisting of single crRNAs tiling TSS-proximal regions of essential genes.

B) Library 1: Scatter plot of cell fitness scores in K562 cells for multiAsCas12a-KRAB vs. denCas12a-KRAB for 3,334 single crRNA constructs with sufficient read coverage for analysis and targeting canonical TTTV PAMs within -50bp to +300bp window of 584 essential gene TSS's. Marginal histograms show percentage of crRNA constructs with cell fitness scores exceeding the 5th percentile of negative control crRNAs. y = x line is shown.

C) Library 1: Moving average cell fitness score across all TTTV PAM-targeting crRNAs at each PAM position relative to the TSS (left), shown for the 240 essential gene TSS's for which analogous dCas9-KRAB NGG PAM tiling screen data (Nuñez et al., 2021) is available in K562 cells (right).

D) Library 1: Boxplots of average cell fitness scores of top 3 crRNAs for each essential TSS for multiAsCas12a-KRAB or denCas12a-KRAB, subtracted by the average cell fitness scores from top 3 sgRNAs for the same TSS for dCas9-KRAB (Nuñez et al., 2021). The top 3 sgRNA's per TSS for dCas9-KRAB are taken from a prior genome-wide screen that used 10 sgRNAs per TSS, which were pre-selected based on bioinformatic prediction of strong sgRNA activity (Horlbeck et al. 2016a). Boxplots show median, interquartile range, whiskers indicating 1.5x interquartile range, and are overlaid with individual data points.

E) Library 1: Bubble plot of the fraction of crRNAs for each PAM that exceed the 5th percentile of negative controls, versus the median CRISPick on-target efficacy score for that PAM. Size of the bubble indicates number of PAM sites per gene.

F) Design of Library 2 Sublibrary A, aimed at evaluating CRISPRi activity each position in the 6-plex array. For each 6-plex array, a specific position is defined as the test position (which can encode either a TSS-targeting spacer or a negative control spacer), and the remaining positions are referred to as context positions encoding one of 5 sets of negative control spacers designated only for context positions.

G) Library 2 Sublibrary A: Analysis of 2,987 6-plex constructs with sufficient read coverage and encodes in the test position one of 123 spacers that scored as strong hits as single crRNAs in the Library 1 screen, compared to constructs that encode an intergenic negative control spacer in the test position. Boxplots show cell fitness scores averaged from the top 3 context constructs of each test position spacer in the 6-plex array, grouped by negative control spacers vs. essential TSS-targeting spacers in each test position. Recall is calculated as the percentage of essential TSS-targeting spacers (that were empirically active in the single crRNA Library 1 screen) recovered by the Library 2 6-plex crRNA array screen for a given test position, using the 5th percentile of constructs containing negative control spacer in the same test position as a threshold for calling hits. Boxplots display median, interquartile range, whiskers indicating 1.5x interquartile range, and outliers. One-sided Wilcoxon rank-sum test was performed for the difference in the distributions of negative control spacers vs. TSS-targeting spacers at each position, with asterisks indicating p<0.0001.

Figure 5 – multiAsCas12a-KRAB CRISPRi enables enhancer perturbation and discovery.
A) K562 cells constitutively expressing multiAsCas12a-KRAB are lentivirally transduced with single crRNAs targeting the HBG TSS or its known enhancer, HS2. Shown are HBG mRNA levels measured by RT-qPCR, normalized to GAPDH levels and averaged across 6-7 technical replicates from 2 independent experiments. Error bars denote SEM. One-sided unpaired Student's t-test was performed to compare untransduced control vs. each individual crRNA; asterisk denotes p<0.05.
B) Genome browser view of the CD55 locus, including predicted enhancers using the activity-by-contact model and DNase-seq and H3K27Ac ChIP-seq tracks from ENCODE. K562 cells piggyBac-engineered to constitutively express multiAsCas12a-KRAB was transduced with 4-plex crRNA constructs targeting regions (R1-R11) in the CD55 locus, and R12 as a negative control region devoid of enhancer features. Each unique 4-plex crRNA construct is labeled as "a" or "b". For comparison, targeting the CD55 promoter using a 6-plex crRNA array (crCD55-4_crB2M-1_crKIT-2_crKIT-3_crCD81-1) is included. CD55 expression was assayed by flow cytometry between 9 and 11 days after crRNA transduction. One-sided Wilcoxon rank-sum test was performed on the median expression knockdown across 4-7 biological replicates for each crRNA construct, compared to the median expression knockdown of R12 (negative control region); asterisk indicates p < 0.01.
C) Comparison of CRISPRi targeting in K562 cells engineered to constitutively express multiAsCas12a-KRAB vs. opAsCas12a using a subset of lentivirally transduced crRNA constructs from B, plus a crRNA construct targeting a coding exon of CD55 as a positive control for knockdown by DNA cutting. CD55 expression was assayed by flow cytometry 11 days after crRNA transduction. One-sided Wilcoxon rank-sum test was performed to compare the median expression knockdown of multiAsCas12a-KRAB vs. opAsCas12a across 2-7 biological replicates; asterisk indicates p≤0.05.

Figure 6 – multiAsCas12a-KRAB enables combinatorial targeting of cis-regulatory elements in pooled CRISPRi screens.

A) Genome browser view of the MYC locus, including activity-by-contact model predictions, and DNase-seq and H3K27Ac ChIP-seq tracks from ENCODE. 3 of the known MYC enhancers (e1, e2, e3) in the body of the non-coding RNA, PVT1, are shown.

B) K562 cells piggyBac-engineered to constitutively express the indicated panel of fusion protein constructs were transduced with one of 4 3-plex crRNA constructs targeting the MYC promoter or co-targeting the 3 enhancers using one crRNA per enhancer. Cell fitness as a proxy of MYC expression is measured as log2 fold-change in percentage of cells expressing GFP marker on the crRNA construct, relative to day 6 after crRNA transduction. Shown are the average of 2 biological replicates, with error bars denoting the range.

C) 6,370 6-plex permutations of the 12 individual spacers from B, together with 3 intergenic negative control spacers, were designed and cloned as 6-plex crRNA arrays used in the design of Library 2 Sublibrary B.

D) Library 2 Sublibrary B: Analysis of 1,823 constructs with sufficient read coverage, categorized based on whether each contains at least one of 3 crRNAs that target the MYC promoter, and/or at least one crRNA that targets each of the MYC enhancers. Boxplots summarize cell fitness score distributions (as proxy of MYC expression) of all constructs that fall in each category. Boxplots show median, interquartile range, whiskers indicating 1.5x interquartile range, and are overlaid with individual data points each representing a 6-plex construct.

Figure 7 – Group testing framework for efficient exploration of combinatorial CRISPR perturbations.

Supplemental Figures

Figure S1 – Example of flow cytometry gating strategy for CRISPRi experiments. Example of the general gating strategy for CRISPRi experiments using flow cytometry readouts, shown for K562 cells as an example. Single live cells are gated by FSC vs. SSC, followed by gating for the fluorescent marker on the crRNA construct (typically GFP), which are subsequently analyzed for target gene expression in the respective fluorescence channels.

Figure S2 – Additional replicate testing dAsCas12a-3xKRAB CRISPRi activity.
Additional replicate for Fig. 1B testing CRISPRi activity of K562 cells piggyBac-engineered to constitutively express dAsCas12a-3xKRAB and using lentivirally delivered crRNA constructs. See Fig. 1B for details.

Well	Protein construct	Plasmid ID
1	multiAsCas12a-HA-XTEN80-KRAB-P2A-BFP	pCH61
2	denAsCas12a-HA-XTEN80-KRAB-P2A-BFP	pCH62
3	enAsCas12a-HA-XTEN80-KRAB-P2A-BFP	pNS49
4	multiAsCas12a-HA-P2A-BFP	pNS35
5	dAsCas12a-3xKRAB-3xHA-P2A-BFP	pCH68
6	No construct	No construct

Figure S3 – Western blot of fusion proteins. Western blot of whole-cell lysates prepared from K562 cells piggyBac engineered to constitutively express each of the fusion proteins in the panel. anti-HA tag was used for detection of the fusion protein and anti-GAPDH for detection of GAPDH as loading control.

Figure S4 – CRISPRi activity of multiCas12a-KRAB, denAsCas12a-KRAB, and dAsCas12a-KRABx3 in C42B cells.

A) C4-2B cells piggyBac-engineered to constitutively express each of the fusion protein constructs are lentivirally transduced with the indicated crRNA constructs. Cells were sorted based on P2A-BFP marker signal. Because some of these cell lines showed slightly different levels of BFP signal as a proxy of fusion protein expression, to account for fusion protein expression we performed propensity score matching to subset for populations of cells for each fusion protein construct with the same distributions in BFP signals after data acquisition for flow cytometry in CRISPRi experiments. The BFP signals before and after matching are shown as violin blots overlaid with boxplots showing median and interquartile range.

B) Target gene expression are measured by cells surface antibody staining and flow cytometry 13-14 days after crRNA transduction. Single-cell distributions of expression knockdown relative to non-targeting crRNA are shown with mean and interquartile range indicated using the cells after propensity score matching for BFP levels as described in A. CRISPRi knockdown results are indistinguishable with and without propensity score matching (not shown).

HEK 293T, dAsCas12a–KRABx3
Day6 co–transfection

Figure S5 – dAsCas12a-KRABx3 CRISPRi by transient transfection in HEK 293T cells. HEK 293T cells were co-transfected with a plasmid encoding for dAsCas12a-KRABx3 and plasmids encoding for the indicated crRNA constructs targeting CD55. Cells were sorted 2 days after transfection for successful co-transfection based on BFP and GFP markers on the plasmids and CD55 expression was measured by antibody staining on flow cytometry 6 days after transfection. Violin plots of single-cell distributions of CD55 expression knockdown as a percentage of the median of non-targeting control are shown. Median and interquartile range are shown in the plot. The percentage of cells below the 5th percentile of the non-targeting control are also shown.

Figure S6 – Comparisons of dAsCas12a variant fusion CRISPRi constructs using up to 3-plex crRNA constructs.
A) The same fusion protein schematic as shown in Fig. 1C, labeled with construct IDs for ease of reference.
B) CD55 expression knockdown measured by flow cytometry using the indicated crRNA constructs and the panel of fusion protein constructs in A. Shown are averages of the median single-cell expression knockdown relative to non-targeting crRNA for 3 biological replicates (including the replicate for crCD55-4 shown in Fig. 1C) for all comparisons, except the comparison for crCD81-1_crCD151-3_crCD55-4 contains 2 replicates. One-sided Wilcoxon rank-sum test comparing denAsCas12a-KRAB (pCH4) to each of the other fusion constructs in the panel was performed. Asterisk indicates p<0.01 for all replicate-level comparisons for a given construct comparison.
C) Analogous to B, but for CD81 knockdown. Summaries shown for 3 biological replicates (including the replicate for crCD81-1 shown in Fig. 1C) for all comparisons, except the comparison for crCD81-1_crCD151-3_crCD55-4 contains 2 replicates.

Figure S7 – Additional replicates testing effect of dose on denAsCas12a-KRAB CRISPRi activity
A) Summary of all replicates for experiment shown in Fig. 1D: shown are averages of median expression knockdown for each crRNA construct (N = 3-6 biological replicates for each crRNA construct, including the replicate shown in Fig. 1D). Error bars denote SEM. One-sided Wilcoxon rank-sum test was performed on the medians of single-cell expression knockdown of each replicate and p-values indicated where relevant.
B) Additional biological replicate for Fig. 1E testing denAsCas12a-KRAB CRISPRi activity with varying crRNA MOI while holding protein MOI constant; see Fig. 1E for details.

Figure S8 – Testing CRISPRi activity of lentivirally delivered truncated crRNAs. In all panels, the indicated cell line (RN2 or B16) was engineered for constitutive expression of the indicated fusion protein constructs by lentiviral transduction, followed by lentiviral transduction (at MOI between 0.3-0.4) of the indicated single-plex crRNA constructs containing spacers of the indicated lengths targeting Rpa3, an essential gene. The spacers target either the gene's coding exon, the TSS region, or the Rosa locus (negative control) as indicated in the legends. Cell fitness defect is used as a proxy of Rpa3 gene expression knockdown due to either DNA cutting or CRISPRi activity. Cell fitness is measured in a competition assay by quantifying log2 fold change in percent of cells expressing the GFP marker on the crRNA expression constructs (relative to day 2). Error bars indicate SEM for N = 3 biological replicates for all panels.

Figure S9 – CD81 knockdown denAsCas12a-KRAB vs. multiAsCas12a-KRAB at different protein and crRNA MOIs. Second biological replicate for testing denAsCas12a-KRAB vs. multiAsCas12a-KRAB CRISPRi activity in the setting of varying protein MOI or crRNA MOI, same as Fig. 2B. See Fig. 2B for further details.

Figure S10 – CD55 knockdown by denAsCas12a-KRAB vs. multiAsCas12a-KRAB at different protein MOIs. Comparison of CD55 knockdown by lentivirally delivered denAsCas12a-KRAB vs. multiAsCas12a-KRAB at protein MOI ~1 vs. ~5 across a panel of single and 3-plex crRNA constructs, while holding constant crRNA MOI for each paired fusion protein comparison for each crRNA construct. Dashed gray line indicates 5th percentile of non-targeting crRNA control. crRNA MOI indicated by color scale. Lines connect paired replicates. One-sided Wilcoxon rank-sum tests were performed on single-cell distributions for each replicate, and asterisk denotes p<0.01 for all paired replicates within each condition. Dots indicate flow cytometry measurement 10 days after crRNA transduction; triangles indicate flow cytometry measurement 16 days after crRNA transduction.

Figure S11 – RNA-seq analysis of crRNA specificity.
A) K562 cells lentivirally engineered (MOI ∼5) to constitutively express multiAsCas12a-KRAB were either transduced with the indicated crRNA's at MOI <0.3, followed by sorting for crRNA-transduced cells based on GFP marker, or received no crRNAs. RNA was isolated from the sorted cells 32 days of culture after crRNA transduction and subjected to 3' RNA-seq. Scatter plot of normalized mRNA expression levels for crRNA transduced (1 biological replicate each) vs. cells without crRNA (2 biological replicates), and Pearson correlation coefficient calculated for the transcriptome, excluding HBG. RT-qPCR quantifications are shown in Fig. 5A.
B) Volcano plots of p-values vs. log2FoldChange from differential expression analysis using DE-seq2 are shown. Genes that fall beyond p-value and log2FoldChange cutoffs (dashed lines) are highlighted.
C) Lists of differentially expressed genes (other than HBG) from the analysis in B for the crHBG-3 and crHS2-3 transduced cells are shown. For comparison, a list of all off-target predictions generated by crisprVerse are shown for all crRNAs in the panel in A and B.

Figure S12 – CRISPRi knockdown of CD55 and B2M using up to 6-plex crRNA arrays. Analogous to Fig. 3B-C, but shown for CD55 and B2M knockdown on day 6 after crRNA transduction, measured by antibody staining of those targets using flow cytometry. Shown are averages of median single-cell expression knockdown from 2-5 biological replicates for each crRNA construct, with error bars indicating SEM.

Figure S13 – Monitoring P2A-BFP reporter as proxy of fusion protein expression level.
A) K562 cells lentivirally engineered (at MOI ~1 or MOI ~5) to constitutively express the indicated fusion protein constructs were monitored for P2A-BFP expression levels by flow cytometry. Shown are representative biological replicates from routine monitoring.
B) Same as A for K562 cells piggyBac-engineered to constitutively express the indicated fusion protein constructs. opAsCas12a does not contain BFP reporter and is shown as fluorescence negative control. Shown are representative biological replicates from routine monitoring.

Figure S14 – Indel quantification and gene expression knockdown simulation for single-targeting.
A) To obtain an upper estimate on the impact on gene expression arising from indels induced by crCD81-1 at the target site, we added an additional 20% unobserved large deletions (upper estimate based on long-read sequencing in Fig. S16A) to the observed indel frequency from short-read sequencing of PCR amplicon (Fig. 2F) to arrive at an estimated per-copy deletion probability. Based on this estimate, the proportions of cells harboring deletions at varying DNA copy numbers are shown assuming independent probability of indel formation across the 3 DNA copies in this region of the K562 genome. Assuming that an indel of any size within the PCR amplicon results in a complete genetic null abolishing CD81 gene expression in cis, we simulate the expected effect of these indels on single-cell distributions of gene expression measured by flow cytometry ("expected null"), compared to the observed gene expression knockdown by crCD81-1 ("observed").
B) Analogous to A, but for crCD55-4, which targets a tetraploid region of the K562 genome. **C)** Indel quantification by short-read sequencing of PCR amplicon for a single-site targeting of the KIT TSS region using crKIT-2 encoded within the indicated 4-plex crRNA array.
D) Analogous to A-B, we simulated expected gene expression knockdown by crKIT-2 single-site targeting based on indel frequencies observed in C. Expected knockdown under this genetic null assumption exceeds that observed for opAsCas12a (fully active DNase), demonstrating the genetic null assumption is an overestimate of gene expression effects of indels in this region. To correct for this overestimate, we use the ratio of observed vs. expected null median expression knockdown by opAsCas12a as an estimate of the hypomorphic effect of deletions in this region ("hypomorphic coefficient"). We multiply the expected null median expression knockdown for all other fusion proteins by this hypomorphic coefficient to obtain an "expected hypormorph" median expression knockdown, which we propose as our final estimate of the effects arising from indels.

Construct	Per-copy deletion probability	No deletion	Deletion in 1 copy	Deletion in 2 copies	Deletion in 3 copies
			Proportion of cells with		
multiAsCas12a-KRAB	0.041	0.882	0.113	0.005	0.000
opAsCas12a	0.530	0.104	0.351	0.396	0.149
enAsCas12a-KRAB	0.996	0.000	0.000	0.012	0.988
multiAsCas12a	0.083	0.770	0.210	0.019	0.001
denAsCas12a-KRAB	0.000	1.000	0.000	0.000	0.000

Figure S15 – Indel quantification and gene expression knockdown simulation for dual-targeting of the KIT TSS region.

A) Related to Fig. 3D. Indel quantification of PCR amplicon near the KIT TSS region in K562 cells lentivirally engineered to constitutively express opAsCas12a 15 days after transduction of the indicated 6-plex crRNA array (sorted for crRNA transduced cells on 2 days after transduction). Note that opAsCas12a is encoded in a different expression backbone using a puromycin selectable marker and thus is not directly matched to other fusion constructs in Fig. 3D in transgenic expression level.

B) We estimated the per-copy deletion probability occuring any where within the PCR amplicon in the KIT locus in Fig. 3D and Fig. S15A based on the on the observed indel allelic frequencies from short-read sequencing, plus an additional 20% unobserved large deletions (upper estimate based on long-read sequencing in Fig. S16A). A single composite per-copy deletion probability is assigned to the locus assuming independent probabilities of small indels generated separately at the crKIT-2 and crKIT-3 target sites, plus assuming that large deletions at one crRNA target site precludes additional alterations in DNA sequence at the other target site. The proportions of cells that harbor a specified number of DNA copies containing indels of any size in this locus are calculated assuming indels occur independently across DNA copies within each cell. We simulated the expected distribution of single-cell gene expression levels under the assumption that indels of any size in this locus result in a genetic null abolishing KIT expression in cis ("expected null"). The expected knockdown under this genetic null assumption exceeds that observed for opAsCas12a (fully active DNase), demonstrating the genetic null assumption is an overestimate of gene expression effects of indels in this region. To correct for this overestimate, we use the ratio of observed vs. expected null median expression knockdown by opAsCas12a as an estimate of the hypomorphic effect of deletions in this region ("hypomorphic coefficient"). We multiply the expected null median expression knockdown for all other fusion proteins by this hypomorphic coefficient to obtain an "expected hypormorph" median expression knockdown, which we propose as our final estimate of the effects arising from indels.

Figure S16 – Long-read sequencing quantification of deletions surrounding single and dual crRNA target sites.
A) K562 cells piggyBAC-engineered to constitutively express the indicated fusion protein constructs were lentivirally transduced with the indicated 6-plex crRNA constructs at MOI < 0.13, sorted for transduced cells based on GFP reporter fluorescence, and the genomic DNA harvested 16 days later and subjected to long-read Nanopore sequencing. Left: Representative view of individual reads aligned to the region surrounding the crCD81-1 target site is shown, with deletions indicated by black horizontal lines. Right: Quantification of deletion frequency across the region surrounding the crCD81-1 target site.
B) Analogous to A), but shown for region surrounding the crCD55-4 target site.
C Analogous to A), but shown for region surrounding the crKIT-3 and crKIT-2 target sites located on opposing DNA strands spaced ~112bp apart.

Figure S17 – Double and triple gene knockdown by CRISPRi using higher-order crRNA arrays.
A) Single-cell view of CD81, KIT, and CD55 3-way knockdown using a 6-plex crRNA construct in K562 cells piggyBac-engineered to constitutively express each of the indicated fusion protein constructs, measured by multiplexed flow cytometry. Summary of percentage of cells with triple knockdown is shown in Fig. 3G.
B) Quantification of the fraction of cells showing double-knockdown of pairs of target genes in the experiment described in A for 4-plex and 6-plex crRNA arrays. Double knockdown is defined as the fraction of cells with expression below the 5th percentile for the non-targeting crRNA for the expression of a given pair of target genes.

A.

Library 1 (single-plex)	TSS-targeting constructs	Negctrl-intergenic	Negctrl-unmapped
# constructs	77387	524	445
# genes targeted	559		
For constructs with ≥1 RPM			
# constructs	76293	524	445
# constructs targeting TTTV PAM	3326		
# target genes with >0 crRNA exceeding fitness score of 1%tile of neg controls	450		
# target genes shared with prior dCas9 CRISPRi screens	240		

B.

Library 2 (6-plex)	Sublibrary A (positional shuffling)	Neg-ctrl only arrays
# constructs	84275	2000
For constructs with ≥1 RPM		
# constructs	67534	324
# constructs with TSS-targeting spacer in test position	44802	NA
# constructs with neg control spacer in test position	12029	NA
# constructs with TSS-targeting spacer in test position exceeding fitness score of 1%tile of neg controls in Library 1 screen	2987	NA
# unique TSS-targeting spacers in test position exceeding fitness score of 1%tile of neg controls in Library 1 screen	123	NA
# unique neg control spacers in test position	506	NA

C.

Library 2 (6-plex)	Sublibrary B (MYC locus)		Neg-ctrl only arrays
# constructs	6370		2000
For constructs with ≥1 RPM			
# constructs	1823		324
Promoter-targeting	**Enhancer-targeting (≥1 for each listed)**	**# constructs**	
≥1 promoter targeting spacer	none	48	
none	none	197	
≥1 promoter targeting spacer	e1	187	
none	e1	65	
≥1 promoter targeting spacer	e2	108	
none	e2	50	
≥1 promoter targeting spacer	e3	90	
none	e3	33	
≥1 promoter targeting spacer	e1e2	216	
none	e1e2	164	
≥1 promoter targeting spacer	e1e3	161	
none	e1e3	121	
≥1 promoter targeting spacer	e2e3	84	
none	e2e3	90	
≥1 promoter targeting spacer	e1e2e3	64	
none	e1e2e3	145	

Figure S18 – Summaries of Library 1 and Library 2 screens.
A) Summary of crRNA constructs in the Library 1 screen.
B) Summary of crRNA constructs in the Library 2, Sublibrary A screen.
C) Summary of crRNA constructs in the Library 2, Sublibrary B screen.

A.

B.

Figure S19 – Screen replicate concordance for Library 1 and Library 2. Shown are 2D density plots of cell fitness scores for individual crRNA constructs in **A)** Library 1 and **B)** Library 2 (Sublibrary A and Sublibrary B), with the Pearson correlation coefficients calculated for all constructs in each library shown.

Figure S20 – Cell fitness score distributions of intergenic vs. non-targeting negative control crRNAs. Boxplots of cell fitness scores for **A)** Library 1 single-crRNA constructs, and **B)** Library 2 6-plex constructs, categorized by whether the construct encodes exclusively intergenic vs. non-targeting negative control crRNAs. Boxplots display median, interquartile range, whiskers indicating 1.5x interquartile range, and outliers.

Figure S21 – Relationship between CRISPick predictions and empirical CRISPRi activity.
A) Library 1: 2D density plots of cell fitness scores of individual crRNAs for multiAsCas12a-KRAB vs. predicted crRNA on-targeting efficacy score from the CRISPick algorithm, grouped by TTTV PAM vs. non-canonical PAM's. The 5th percentiles of intergenic negative control crRNAs cell fitness scores are shown as a dashed horizontal line and the percentage of crRNAs below that threshold shown in the marginal histogram. Pearson correlation coefficients are shown.
B) Library 1: Pearson correlation coefficient between the cell fitness score and the CRISPick on-target efficacy score is shown for each individual PAM sequence for multiAsCas12a-KRAB.

Figure S22 – Difference in cell fitness scores for 6-plex vs. 1-plex crRNA constructs. Library 2 Sublibrary A (6-plex crRNA construct with one test position targeting essential TSS) screen cell fitness scores for a given test position spacer are compared to the cell fitness scores of the same spacer in the Library 1 (single-plex crRNA) screen. Difference in cell fitness scores (6-plex minus single-plex) are shown as boxplots, which display the median, interquartile range, and whiskers indicating 1.5x interquartile range.