Steroid hormones produced by coral-associated *Endozoicomonas* prime the coral immune response during thermal stress

Amin R. Mohamed[1,2$*], Michael A. Ochsenkühn[1$], Trent D. Haydon[3], Salah Abdelrazig[1], Lisa S.Y. Coe[1], Gabriel Mochales-Riaño[3], Woonkee S. Jo[4], Alan R. Healy[4], David Abrego[5,6], Jean-Baptiste Raina[7], John A. Burt[3,8,9], Shady A. Amin[1,3,9*]

[1]Marine Microbiomics Laboratory, New York University Abu Dhabi, Saadiyat Island, Abu Dhabi, United Arab Emirates

[2]Zoology Department, Faculty of Science, Benha University, Benha, Egypt

[3]Center for Genomics and Systems Biology, New York University Abu Dhabi, Saadiyat Island, Abu Dhabi, United Arab Emirates

[4]Chemistry Program, New York University Abu Dhabi (NYUAD), Saadiyat Island, United Arab Emirates

[5]Faculty of Science and Engineering, Southern Cross University, Coffs Harbor, NSW, Australia

[6]Lizard Island Research Station, Australian Museum, Sydney NSW, Australia

[7]Max Planck Institute for Marine Microbiology, Bremen, Germany

[8]Marine Biology Laboratory, New York University Abu Dhabi, Saadiyat Island, Abu Dhabi, United Arab Emirates

[9]Mubadala Arabian Center for Climate and Environmental Sciences, New York University Abu Dhabi, Saadiyat Island, Abu Dhabi, United Arab Emirates

[*]Corresponding Authors:

Shady A. Amin, Biology Program, New York University Abu Dhabi, Saadiyat Island, Abu Dhabi, United Arab Emirates. Email: samin@nyu.edu , Tel: (+971) 2 628 5743

Amin R. Mohamed, Biology Program, New York University Abu Dhabi, Saadiyat Island, Abu Dhabi, United Arab Emirates. Email: amin.mohamed@nyu.edu, Tel: (+971) 2 628 7863

[$]equally-contributed

Abstract

The coral microbiome is believed to play a critical role in sustaining corals and enabling adaptation to thermal stress. A ubiquitous group of coral-associated bacteria, known as *Endozoicomonas*, are hypothesized to provide essential metabolites to corals, though the exact roles these bacteria play during ambient or thermal stress conditions are largely unknown. We demonstrate that while *Endozoicomonas* synthesizes the iron-binding siderophore aerobactin *in vitro*, this trait may be suppressed *in hospite*. Integrated metagenomics, metabolomics, and lab validations reveal that within the coral holobiont *Endozoicomonas* is the only microbe capable of metabolizing coral-derived testosterone and converting ubiquitous cholesterol into testosterone and progesterone, important steroids that accumulate within the coral holobiont during thermal stress. Provision of testosterone and progesterone to corals during thermal stress induced transcriptional reprogramming of the coral innate immune response, promoting cellular survival and suggesting active microbial involvement in immune priming and stress resilience. These findings highlight a previously unrecognized mechanism of microbial contribution to coral resilience. Lastly, the capacity of bacteria to synthesize eukaryotic steroids implies deep evolutionary connections between microbial metabolism and the origin of steroid-mediated processes in early eukaryogenesis.

Keywords: coral symbiosis, *Endozoicomonas*, thermal stress, chemical diversity, steroids, siderophore.

Introduction

Corals form intricate associations with a diverse microbiome, collectively referred to as the coral holobiont[1]. This microbiome has been recognized as an essential contributor to coral health and resilience, especially during thermal stress[2–4]. While the symbiotic interactions between corals and their primary algal endosymbiont, Symbiodiniaceae, have been extensively studied[5–7], the functional roles of bacteria in the coral holobiont remain largely unexplored[2,8], despite their importance for coral health[3,9,10]. Among the myriad of bacteria associated with corals, the genus *Endozoicomonas* has received considerable attention due to its ubiquitous distribution across coral species and geographic locations[11–13] and the correlation between its abundance and coral health[11,14,15].

While *Endozoicomonas* are sometimes present in the coral mucus[16,17], they are primarily localized within coral tissues as cell-associated microbial aggregates (CAMAs)[18]. Based on genomic predictions, *Endozoicomonas* spp. are hypothesized to play a crucial role in coral holobiont metabolism through the recycling of essential nutrients and biosynthesis of B-vitamins, amino acids and organosulfur compounds[11,19–21]. Indeed, when exposed to coral tissue extracts, *E. marisrubri* activates vitamins B_1 and B_6 biosynthesis and glycolytic processes[22]. However, despite recent efforts[13,21,23–25], a definitive role for *Endozoicomonas* within the coral holobiont and the precise function underlying its symbiotic association is yet to be identified.

In this study, we combine multiomics data (16S rRNA gene amplicons, shotgun metagenomics, and untargeted metabolomics) to elucidate the functional roles of *Endozoicomonas* during coral response to extreme thermal conditions in the Arabian Gulf, the hottest sea on Earth with surface temperature exceeding 36°C[26]. To investigate symbiotic interactions in the coral holobiont, our sampling scheme included both healthy corals, representing stable host-microbiome associations, and corals infected with white syndrome, representing a dysbiotic state. Diseased samples, characterized by a marked reduction in the relative abundance of *Endozoicomonas*, provided a natural contrast to healthy corals that enabled us to determine the functional contribution of this taxon. By integrating metagenomic and metabolomic data, we uncover both shared and niche-specific metabolic pathways among members of the coral holobiont. Given the prominence of *Endozoicomonas* as a key member of the healthy coral microbiome, we leverage pangenomic analysis to delineate conserved and unique functions in *Endozoicomonas* lineages. We then perform additional experimental validations of these functional predictions.

We show that *Endozoicomonas* encodes genes for the biosynthesis and uptake of the iron-binding siderophore aerobactin, and confirm its ability to produce it *in vitro*, likely to access bioavailable

iron *ex hospite*. In addition, we show that during thermal stress, two terminal steroid hormones,

testosterone and progesterone, are abundant in corals despite the depletion of most of their

precursors. We reveal that *Endozoicomonas* is the only holobiont member to possess the gene

repertoire required for testosterone degradation and, surprisingly, can metabolize cholesterol to

produce progesterone and testosterone *in vitro*, a unique capability only present in select bacteria.

Finally, we show that progesterone and testosterone play a critical role in modulating the coral

immune response during thermal stress by promoting pro-survival signaling while suppressing

pro-inflammatory responses. These findings constitute a major breakthrough in our understanding

of bacterial contributions to coral resilience and host–microbe interactions.

Results

We monitored and sampled colonies of the coral *Acropora pharaonis*[27] for 16S rRNA gene

amplicons, shotgun metagenomics and untargeted metabolomics from Saadiyat reef on the

Arabian Gulf off the coast of Abu Dhabi. Sampling occurred across a temperature gradient starting

with early thermal stress showing preliminary signs of white syndrome disease on some colonies

at 32°C (June), severe thermal stress and further white syndrome progression at 34°C (August),

and recovery at 27°C (October) (Extended Data Table 1). Samples were collected from healthy

coral tissues at all three temperatures, and from dysbiotic controls (i.e., white-syndrome lesion

sites (L) and healthy appearing tissue adjacent to the lesion (AL)) at 34°C (Fig. 1a). All samples

were extracted for metabolites and genomic DNA to profile the coral holobiont metabolome and

microbiome composition and functional potential.

Divergent metabolite and bacterial profiles in corals under extreme temperature

The intracellular metabolites of the coral holobiont were extracted and analyzed with reverse

phase liquid chromatography-high resolution mass spectrometry (RP-LC/HRMS) in positive and

negative ionization modes and in tandem with MS^2 mass fragmentation. A total of 5878 putative

molecular features were predicted based on *in-silico* mass fragmentation network analysis

(SIRIUS) (*see* Methods; Supplementary Information Table 1). Principal component analysis (PCA)

patterns were significantly different (PERMANOVA; F-value = 7.58; R^2 = 0.62; adj. *p*-value=0.001)

and showed that early stress (32°C) and recovery (27°C) metabolomes clustered together away

from severe stress (34°C), suggesting that despite being separated by four months (June to

October) the holobiont metabolic state during early stress and recovery were similar (Fig. 2a;

Extended Data Fig. 1). The metabolomes of the diseased samples (L, AL) displayed inter-colony

variations and showed a significant separation from all other samples, suggesting a fundamentally different chemical composition compared to healthy tissue (Fig. 2a). These findings confirm that acroporid corals in the Arabian Gulf exhibit a notable tolerance to temperatures up to 32°C, lethal to corals elsewhere, while exposure to 34°C poses a critical challenge to their overall fitness, which aligns with previous reports[28].

The bacterial microbiome composition, based on 16S rRNA gene amplicons, revealed that disease samples (L, AL) differed significantly from all other samples (PERMANOVA; F-value = 1.98; R^2 = 0.0318; *adj.* p-value = 0.01), suggesting the bacterial microbiome is resilient and only undergoes a significant shift during disease progression (Fig. 2b). Thirty-two bacterial ASVs belonging to the family Endozoicomonadaceae dominated the microbiome, with an average relative abundance of 60% in healthy samples. In addition, Endozoicomonadaceae persisted under elevated (32°C) and even extreme temperatures (34°C), which was particularly remarkable as they usually disappear during coral exposure to thermal stress[29] (Fig. 2c; Extended Data Fig. 2a). At lesion sites, the relative abundance of Endozoicomonadaceae was only 1.3%, while Rhodobacteraceae and Saprospiraceae contributed 23% and 39% of the microbiome, respectively, indicating the proliferation of potentially opportunistic microbes at the expense of Endozoicomonadaceae during disease onset (Fig. 2c; Extended Data Fig. 2a)[30–32].

Integrated metagenome/metabolome analysis implicates the microbiome in steroid hormone metabolism

The metagenomes of coral holobionts produced a total of 589 gigabases (Gb), averaging 42 Gb per sample (Supplementary Information Table 3; Extended Data Fig. 3). We leveraged publicly available genomes for *Acropora* and *Cladocopium* to partition the holobiont metagenome before assembling coral and algal contigs (see Methods). A total of 257 million reads (~7% of the holobiont reads) were classified as microbial (i.e., non-coral, non-algal reads; Fig. 2d). These microbial reads were used for downstream read-based and assembly-based metagenomic analyses (Extended Data Fig. 2b and 3; Supplementary Information Table 3). Taxonomic profiling of putative metagenomic microbial reads revealed a diverse coral microbiome primarily dominated by bacteria, comprising 95% of the microbial community (Extended Data Fig. 2b). To explore the potential roles of the microbiome, we assigned all genes and putatively annotated metabolites in our metagenomic and metabolomic datasets to members of the coral holobiont. To do this, metagenomic reads assigned to host, algal symbiont or bacteria were assembled and functionally annotated using the KEGG database to a total of 11,333 KEGG Orthology terms[33] (KOs). While a third of the predicted KOs (3,065) were shared between different members of the holobiont, most

were uniquely assigned to the coral host (4,619), followed by microbes (3,029) and algal

symbionts (620) (Fig. 2e; Supplementary Information Table 4). Using these KO taxonomic

assignments, we mapped the putatively annotated metabolites associated with KEGG compounds

(284) (Supplementary Information Table 2). This analysis indicated 148 of these putative

metabolites were shared between different members of the holobiont, while 43 were assigned

uniquely to the host, 42 to the algal symbiont and 51 to other microbes (Fig. 2e, Extended Data

Table 2). Examining the KEGG pathways across metagenomes and metabolomes indicated that

the host dominates the biosynthesis of steroid hormones, retinol metabolism, and vitamin

digestion and absorption, while microbes dominate biosynthesis of cofactors, biosynthesis of

secondary metabolites, and steroid degradation (Fig. 2e).

To assign functions to specific bacterial taxa, we assembled genomes from our metagenomes

(MAGs). Coral samples devoid of mucus and seawater yielded 11 medium to high-quality MAGs

(completeness >75% and contamination <10%). Core microbiome members were defined as

MAGs consistently present at relatively high (40-60%) abundance in healthy tissues (at all

temperatures), but showing reduced abundance in dysbiotic samples (AL and L) (Fig. 3a). Out of

the 11 MAGs, two MAGs stood out as important members of the coral holobiont: MAG01 assigned

to the phylum Myxococcota and MAG10 assigned to the genus *Endozoicomonas* (Extended Data

Fig. 4; Supplementary Information Table 5). *Endozoicomonas* (MAG10; completeness: 91%,

contamination: 1.4%), exhibited a dynamic abundance profile as its relative abundance peaked at

32°C during early stress (65.4%), followed by a dramatic decrease at 34°C (35.8%) and then

bounced back with a modest abundance during recovery at 27°C (43%) (Fig. 3a). The relative

abundance of MAG10 dropped drastically in L samples (0.8%), coinciding with dysbiosis and

consistent with previous reports[30–32]. Given its dynamic abundance pattern and strong association

with healthy coral states, we next explored the functional potential of MAG10 to better understand

its role within the coral holobiont.

Pangenome analysis reveals conserved functions across *Endozoicomonas* genomes

Since culturing efforts of *Endozoicomonas* from *Acropora* in the Arabian Gulf were not successful,

we isolated *Endozoicomonas acroporae* AT381 from *A. kenti* from the Great Barrier Reef (GBR),

referred to hereafter as AT381 (Fig. 1b), and obtained a nearly complete genome (size = 6.02

Mbp; completeness = 99.1%) predicted to harbor 5001 coding sequences (CDS). AT381 and

MAG10 are closely related as they share an average nucleotide identity (ANI) of 94.3%[34].

Phylogenomic analysis between MAG10, AT381 and 27 publicly available *Endozoicomonas*

genomes and MAGs from diverse invertebrate hosts further confirmed the close relationship between MAG10, AT381 and other *E. acroporae* genomes (Extended Data Fig. 5). Functional conservation and specialization across all *Endozoicomonas* genomes were examined by constructing a pangenome of their 34,846 gene clusters (Fig. 3b, Supplementary Information Table 6)[35]. Because of the inability to culture a MAG10 relative from the Arabian Gulf, we used pangenomic analysis to confirm functional conservations within the *Endozoicomonas* genus and further functionally validated some of these conserved capabilities using the closely related *E. acroporae* AT381 from the GBR and two other *Endozoicomonas* isolates.

We hypothesized that *Endozoicomonas* genomes symbiotic with corals possess specialized, but shared functions enabling interactions with their coral hosts, compared to genomes isolated from other hosts. As reported for other genomes of coral-associated *Endozoicomonas* spp., MAG10 harbors genes related to symbiosis establishment, e.g., flagellar assembly, type IV pili[18], ankyrin and tetratricopeptide repeats[21,22], several secretion systems involved in host infection (T2SS, T3SS, T6SS)[18,21–23], complete biosynthesis pathways for the vitamins riboflavin (B$_2$), biotin (B$_7$) and pyridoxal-phosphate (B$_6$), and nine essential amino acids and their transporters (Extended Data Fig. 4b and c, Supplementary Information Table 7)[11]. MAG10 was also significantly enriched within the holobiont for antioxidant production[22], heme biosynthesis[25], glycoside hydrolases that cleave the most abundant marine polymers, starch and chitin[36], and diverse ABC transporters (Supplementary Information Tables 7 and 8). In addition to these previously known functions, MAG10 is capable of **(a)** catabolizing histidine to glutamate, an essential intermediate for ammonium assimilation and a common feature of most *Endozoicomonas* genomes, **(b)** degrading galactose (Leloir pathway), a Symbiodiniaceae photosynthate[37], which is a common feature in most *Endozoicomonas* genomes, **(c)** biosynthesizing the nucleoside antibiotic showdomycin, **(d)** biosynthesizing and exporting the polyamine spermidine, a reactive oxygen species scavenger, and export proteins for its precursor putrescine, **(e)** and biosynthesizing ectoine, a bacterial osmoprotectant (Fig. 3b, Extended Data Fig. 4c, Extended Data Tables 3 and 4, Supplementary Information Table 7). Cumulatively, these functions may enhance symbiosis and increase the fitness of the coral host and/or Symbiodiniaceae, especially during thermal stress.

Endozoicomonas acquires iron *in vitro* using the siderophore aerobactin

Siderophores are small molecules secreted by some bacteria and fungi to acquire iron in iron-limited environments[38] and can play important roles in iron cycling and bioavailability in the pelagic

ocean[39,40]. The genome of *Klebsiella pneumoniae,* a model bacterium for studying the
biosynthesis of the siderophore aerobactin, possesses the canonical, complete gene
cluster/operon containing aerobactin synthases (*iucA/iucC*) and the TonB-dependent uptake
receptor (*iutA*) used to produce apo-aerobactin and take up Fe-bound aerobactin, respectively[41].
In contrast, MAG10, all other *E. acroporae* and 8 additional coral- and invertebrate-associated
Endozoicomonas genomes possessed only the biosynthesis genes within the gene cluster, while
having either a distal *iutA* that is outside of the gene cluster in the case of AT381 or a homolog of
iutA in a completely separate region of the genome for all other *Endozoicomonas* that possess
the gene cluster (Fig. 4a, Extended Data Table 4). Aerobactin requires a functional *iutA* for Fe
acquisition to function[42] and a distal *iutA* suggests a potential decoupling from biosynthesis. To
confirm whether *Endozoicomonas* can use aerobactin to acquire iron, we grew multiple strains *in*
vitro under Fe-limiting conditions and tested their siderophore production using the chrome azurol
S assay (CAS)[43]. The tested strains include two isolates from *Acropora*: *E. acroporae* AT381 and
E. acroporae Acr14*,* and *E. montiporae* CL33 (isolated from *Montipora*), a negative control that
lacks aerobactin biosynthesis genes (Extended Data Table 4). The CAS assay showed that *E.*
acroporae AT381 and Acr14 were able to grow and utilize Fe from CAS (Fig. 4b) while *E.*
montiporae was not able to grow, consistent with its inability to produce aerobactin. These findings
suggest that select *Endozoicomonas* strains and species (12 out of 27 in total) can utilize
siderophores for Fe acquisition *in vitro* and likely *ex hospite*. We were not able to detect any m/z
signatures that match aerobactin, its Fe-complex or photoproducts[44] in the holobiont
metabolomes. Indeed, siderophore production *in hospite* is mainly associated with pathogenic
bacteria[45], and siderophore production may take place when *Endozoicomonas* are in seawater *ex*
hospite.

The fact that all examined *Endozoicomonas* genomes capable of biosynthesizing aerobactin
possess distal, not proximal, *iutA* homologs indicates a potential decoupling of expression
between biosynthesis and transport, which can make Fe acquisition less efficient and more
energetically costly for the bacteria. This pattern also suggests a gradual loss of function of the
transporter that would eventually abolish aerobactin biosynthesis over evolutionary timescale.
Indeed, a loss of function was concluded from the lack of *iutA* in the genomes of *Bacillus*
licheniformis and *B. halodurans* despite the presence of the aerobactin biosynthesis operon[46].

To assess the evolutionary pressures acting on the *iutA* gene, we calculated rates of non-
synonymous and synonymous substitutions (dN/dS) ratios across multiple *E. acroporae* strains
compared to the housekeeping gene DNA Polymerase III subunit delta as a conserved core gene

against *E. acroporae* AT381 as a reference (Supplementary Information Table 9). Several
Endozoicomonas isolates from *Acropora* displayed signatures of relaxed purifying selection
(significantly elevated dN/dS) in *iutA*, suggesting ecological redundancy or diminished reliance on
siderophore-mediated iron acquisition in certain niches. In contrast, *iutA* in *E. acroporae* Acr14
exhibited low dN/dS value (0.1), consistent with strong purifying selection and functional
conservation. These patterns suggest that while some *Endozoicomonas* strains may reduce
dependence on siderophore systems, others retain this capability, potentially as a survival strategy
during transitions to a free-living lifestyle *ex hospite*. MAG10 was not included in this analysis
because MAGs are typically a mosaic of sequences from multiple closely related strains, rather
than a single, clonal genome. This chimerism, along with assembly artifacts, can distort
substitution rate estimates and obscure true evolutionary signals[47], a concern that is particularly
relevant here given the high strain-level diversity we observed, with 32 distinct ASVs from
Endozoicomonadaceae identified by 16S rRNA gene amplicon sequencing. Thus, we propose
that coral-associated *Endozoicomonas* do not utilize siderophore-mediated iron uptake *in hospite*,
a function that could be used in the free-living stage, and that over evolutionary time aerobactin-
dependent Fe acquisition will be lost. Coral-associated *Endozoicomonas* can instead utilize other
pathways to acquire iron, such as heme transporters (Supplementary Information Table 7).

Steroid metabolism is a prominent function of *Endozoicomonas* among the coral
microbiome

MAG10 was the only holobiont member that possessed the complete pathway to utilize carbon
from the degradation of the steroid hormone testosterone via 3-oxosteroid 1-dehydrogenase
(*kstD*) gene (Fig. 5a,b; Supplementary Information Table 4, 7; Extended Data Fig. 3). This function
was shared with most *Endozoicomonas* genomes (Fig. 3b; Extended Data Table 4), consistent
with their broader capacity to break down this important steroid molecule[48]. The relative
abundance of most steroids gradually and significantly decreased as temperature increased, with
abundance at 34°C, AL, and L near background levels (Fig. 5a, Extended Data Fig. 6,
Supplementary Information Table 9), consistent with previous reports[49,50] showing suppression of
steroid hormone production under stress and during infection. Surprisingly, a notable exception to
this pattern were select key terminal steroids and steroid degradation products. Specifically, the
putative eukaryotic hormones, progesterone, testosterone and estradiol-17β, in addition to
Endozoicomonas-derived androsta-1,4-diene-3,17-dione and HIP showed the highest relative
abundance at 34°C, with most other samples at near background levels (Fig. 5a, Supplementary

Information Table 10). Since the sampling timepoints fell outside the spawning season in the

Arabian Gulf (March to May)[51], and most precursors of progesterone and testosterone showed

low relative abundance near background levels at 34°C, the elevated levels of these putative

steroid hormones during severe stress at 34°C were unexpected. Since these precursors are

required for the synthesis of testosterone and progesterone, their absence suggests they are

either depleted by the host or a source other than the host, is providing testosterone and

progesterone. Notably, MAG10 was the only microbiome member that possessed a 17-β-

hydroxysteroid dehydrogenase (K05296; E1.1.1.51) that synthesizes testosterone from

androstenedione, the reverse reaction of KstD (Supplementary Information Table 7)[52]. Therefore,

we hypothesized that *Endozoicomonas* may provide corals with some of these steroids during

severe stress.

MAG10 and other *Endozoicomonas* genomes (e.g., *E. montiporae*) encode genes associated with

cholesterol binding, such as those containing the steroidogenic acute regulatory protein-related

lipid transfer (START) and sterol carrier protein-2 (SCP2) domains, as well as

cholesterol/phospholipid transport, including MlaD, MlaE, and ABC transporters (Supplementary

Information Table 7). Therefore, we investigated whether *Endozoicomonas* can catabolise

testosterone, convert androstenedione to testosterone and if it synthesizes the terminal steroids

progesterone, testosterone, or estradiol-17β from cholesterol, given that cholesterol is the

precursor of steroid hormones and can be one of the abundant steroid molecules in the coral

holobiont[48,53]. Cultures of three *Endozoicomonas* strains, *E. acroporae* AT381, *E. acroporae* Acr14

and *E. montiporae* were grown at 26°C and 34°C in minimal media with or without cholesterol,

testosterone or androstenedione (Fig. 1b). After 24 hours, all three strains catabolized

testosterone into androstenedione and downstream metabolites leading to the TCA cycle, with *E.*

acroporae Acr14 only carrying out this transformation at 34°C, while *E. acroporae* AT381 and *E.*

montiporae doing so at both temperatures (Fig. 5c, Supplementary Information Table 11),

suggesting thermal regulation of these transformations in *E. acroporae* Acr14. All three strains

were also able to convert androstenedione to testosterone at both temperatures likely via 17-β-

hydroxysteroid dehydrogenase (Fig. 5d, Supplementary Information Table 11). Surprisingly, all

three strains were capable of transforming cholesterol into terminal steroids. *E. acroporae* AT381

converted cholesterol into the terminal products progesterone, testosterone, androstenedione and

other intermediates at both temperatures (Fig. 5e, Extended Data Fig. 6c). Conversely, *E.*

acroporae Acr14 converted cholesterol to cholesterol derivatives and progesterone only at 34°C

with no transformations observed at 26°C. Finally, *E. montiporae* converted cholesterol to

androstenedione, testosterone and related metabolites only at 26°C and switched to converting

cholesterol to cholesterol derivatives and progesterone at 34°C, indicating a strong temperature

influence on the regulation of these transformations (Fig. 5e). These findings suggest that

depending on the species/strain of *Endozoicomonas*, cholesterol conversion towards testosterone

or progesterone is controlled by temperature, which may explain the high relative abundance of

progesterone and testosterone in the holobiont at 34°C (Fig. 5 and Extended Data Fig. 6a,b).

Using high-resolution mass spectrometry and fragmentation information, we propose a metabolic

pathway that bifurcates cholesterol into **(a)** progesterone via pregnenolone and **(b)** testosterone

via oxo-cholestenoate and androstenedione (Fig. 5f). Although the *Endozoicomonas* pangenome

and auto-annotation highlight important enzymes for steroid degradation (e.g. *kstD*), further mining

of the *E. montiporae* genome yielded steroid delta isomerases of the CyP450 family, which

putatively convert pregnenolone into progesterone (Fig. 5f; Supplementary Information Table 7)[48].

MAG10 also encodes a member of the 3-beta hydroxysteroid dehydrogenase/isomerase protein

family [PF01073.22] (Supplementary Information Table 7). Orthologs of this protein are present in

all three *Endozoicomonas* strains we used for steroid incubations. This enzyme family includes

3β-hydroxysteroid dehydrogenase/Δ^5-Δ^4 isomerases (3β-HSDs), which plays a central role in

steroid biosynthesis by catalyzing the oxidation and isomerization of steroid precursors into active

steroid hormones. Although mining *Endozoicomonas* genomes did not reveal any specific

exporters for testosterone or progesterone, both hormones are lipophilic and can diffuse freely

through the cytoplasmic membrane[54–56]. Collectively, these findings indicate that *Endozoicomonas*

possess the genetic repertoire not only for steroid degradation, but for their conversion too,

positioning them as active contributors to steroid metabolism in the coral holobiont and suggesting

that these steroids play an integral role in the coral response to thermal stress. Because

testosterone is an immunosuppressive substrate under stress and progesterone is an

immunomodulatory and anti-inflammatory substrate in various animal models, we hypothesized

that the production of these hormones by *Endozoicomonas* likely modulates how the coral immune

system responds to thermal stress.

Steroid hormones prime the coral immune response during thermal stress

To determine whether corals can take up exogenous steroid hormones and their effective

concentrations, we incubated *Acropora* nubbins with three different concentrations of [13]C-labeled

testosterone (10 nM, 100 nM, 1 *μ*M) for 48 hours and examined its relative abundance after 24

and 48 hours (Fig. 6a). While corals incubated with 1*μ*M accumulated the most testosterone after

24 hours, the steroid relative abundance was highly reduced (3-fold) at 48 hours, suggesting saturation. Similar results were obtained with incubations with 100 nM testosterone. At 10 nM, *Acropora* accumulated testosterone progressively at 24 and 48 hours, resulting in an 11-fold increase indicative of active uptake and limited metabolic removal, consistent with a potential biological role; therefore, we used this concentration for subsequent experiments for both testosterone and progesterone. To test the effect of each molecule on the immune response of *Acropora,* coral nubbins were incubated with each molecule for 3 hours at an ambient temperature of 28°C (T1), followed by an increase in temperature to 33°C (T2), simulating thermal stress and at which point a second dose of progesterone or testosterone were added. To map the coral immune system response to these steroids during thermal stress, samples were collected at T1, T2 and at 24 hours post T1 (T3) (Fig. 6b) for RNA sequencing. Coral fragments were monitored by measuring algal symbiont's photosynthetic efficiency (*Fv/Fm)* to ensure corals were stable and exposure to both steroid hormones did not lead to physiological toxicity (Fig. 6c).

To examine differential expression of immune-related genes upon coral exposure to testosterone or progesterone during thermal stress relative to untreated controls (Supplementary Information Table 12, 13, 14), we compiled a list of all known innate immunity-related genes from our assembled transcriptome (Fig. 6d). While many immune-related genes were activated due to heat stress, exposure to testosterone or progesterone further increased their transcription (Fig. 6d,e). For example during T2, both testosterone and progesterone treatments caused on average an increase in the fold-change expression of master regulators of innate immunity, including tumor necrosis factor receptor (TNFR; +129%), nuclear factor kappa B (NF-κB; +80%), and activator protein 1 (AP-1; +10%) relative to controls, indicating increased immune signal transduction and immune priming. Concomitantly, upstream immune-sensing and adaptor pathways (e.g., RIG1 and *TRAF2*; +131%, +1.9% respectively), as well as downstream genes linked to immune regulation and anti-apoptosis (e.g., *TNFAIP3*, *BIRC2/3, BCL2*; +18.5%, +64%, +104% respectively) were activated. This suggests that both testosterone and progesterone prime the coral innate immune system during thermal stress by enhancing immune signaling and activating both upstream sensing and downstream protective genes. Expression of pro-inflammatory signaling components decreased (*TRAF6*; -32% and the JAK/STAT receptor *DOME*; -84%), indicating immune-dampening that limits immune overreaction. Despite these shared effects in the presence of testosterone and progesterone, progesterone uniquely enhanced transcriptional responses associated with cell survival and anti-apoptotic signaling (*MCL1, BIRC4, PIM1*; +25%, +161%, +245% respectively) and preferentially promoted negative feedback regulation of cytokine-activated JAK/STAT pathway (*CISH/SOCS2*; +190%), consistent with suppression of

prolonged or excessive immune activation (Fig. 6d,e). Coincidentally, progesterone was
preferentially produced by some *Endozoicomonas* from cholesterol only during temperatures
resembling thermal stress (Fig. 5e), suggesting *Endozoicomonas* modulates the coral immune
system to suppress prolonged inflammatory responses to thermal stress.

Discussion

Despite the central role of microbial associates to coral health and fitness, the functional
mechanisms bacteria utilize to influence host physiology remain largely unresolved. Here, we
provide the first evidence that ubiquitous *Endozoicomonas* possesses an unexpected capacity to
synthesize terminal eukaryotic steroid hormones, a novel metabolic capability in the coral
microbiome, and that these hormones modulate the host immune response during thermal stress,
suggesting a critical functional role for this genus within the coral holobiont. Particularly, the ability
of *Endozoicomonas* to synthesize testosterone and progesterone in a temperature-dependent
manner confirms that their accumulation in the holobiont when seawater temperature is high may
be partially or wholly driven by *Endozoicomonas* symbionts. While steroid degradation is
widespread in many alpha- and gamma-proteobacteria[48,57,58], the ability to synthesize terminal,
host-active steroid hormones from cholesterol is rare and positions *Endozoicomonas* as a
metabolic outlier among coral symbionts. Specific human gut microbiota, such as *Bacteroides*
acidifaciens and *Ruminococcus gnavus*, synthesize steroid hormones and play a major role in
steroid hormone homeostasis in mammals[59]. Moreover, *Mycobacterium smegmatis* was shown to
synthesize testosterone and estrogens *in vitro* from steroid precursors[60]. While the mycobacterial
conversion of cholesterol to testosterone has been observed under fermentative conditions,
Endozoicomonas differs in its ability to carry these transformations under aerobic conditions,
highlighting a distinct biochemical strategy that can influence major functions in their host.

While steroid hormones play critical roles in many physiological functions, such as growth,
reproduction, and development in vertebrates[48], their role and origin in invertebrates have been
subject to recent debate[61]. While definitive CYP19A1 and HSD17B3 orthologs (the vertebrate
enzymes that catalyze the final steps in estrogen and androgen synthesis, respectively) have not
been identified in invertebrate genomes, several studies have demonstrated functional aromatase
activity in corals (e.g., *Euphyllia ancora*, *Pocillopora damicornis*) using substrate conversion
assays and inhibition by vertebrate aromatase inhibitors[62,63] that suggest invertebrates are capable
of synthesizing these molecules. We also identified several coral genes involved in steroid

hormone biosynthesis (Supplementary Information Table 7), including 17β-hydroxysteroid dehydrogenase and several members of the cytochrome P450 enzyme family that can biosynthesize testosterone and progesterone, respectively, as reported in previous coral genomes[64]. This suggests that although the exact gene homologs that produce steroid hormones may differ, the enzyme function is present.

In corals, steroid hormones play important roles in gametogenesis and spawning and exogenous additions of steroid hormones influence the coral microbiome composition[61,65,66]. Our findings indicate that, in addition to these functions, steroid hormones modulate the immune response to thermal stress within hours of progesterone or testosterone addition. Specifically, hormone exposure under thermal stress preferentially engaged TRAF2-associated TNF/NFKappaB signaling and upregulated immune negative regulators, suggesting a survival-oriented, immune-modulating state rather than a canonical inflammation response. TRAF signaling can diverge toward either pro-survival or pro-inflammatory outcomes, with TRAF2 typically implicated in cell survival and anti-apoptotic signaling[67] and TRAF6 commonly associated with inflammatory pathways[68]. Indeed, exposure to either progesterone or testosterone under thermal stress induced TRAF2 expression while suppressing TRAF6, suggesting a shift from inflammatory to anti-apoptotic response relative to the coral's response under thermal stress alone (Fig. 6e). In contrast to the combined steroid hormone response, progesterone uniquely attenuated the JAK/STAT signaling pathway by downregulating DOME receptors and inducing the negative regulators of JAK/STAT, CIS and SOCS (Fig. 6e). These patterns may fine-tune the immune response by limiting immune overactivation while preserving basal immune competence[69,70]. By priming innate immunity while constraining excessive inflammation, progesterone may help balance pathogen defense with tissue protection, enhancing resistance to opportunistic microbes without exacerbating heat-induced cellular damage or algal symbiont loss[71,72]. This balance may be further fine-tuned by the temperature-sensitive regulation of steroidogenic biosynthesis pathways in *Endozoicomonas* (Fig. 5), which may exert a disproportionate influence on holobiont physiology relative to its abundance during thermal stress. Given the deep evolutionary conservation of innate immune pathways in cnidarians[73,74], bacterially-derived steroid-based signaling may represent an underappreciated metabolic interface between bacteria and early diverging metazoa that contributes to the evolution of metazoa-microbe associations.

Conclusion

Recent evidence suggests the origin of protosteroids occurred up to 1.6 billion years ago[75],

coinciding with the proposed evolution of unicellular eukaryotes. This correlation indicates that the

origin of these molecules is most likely eukaryotic, especially given their role in sexual

reproduction, which is absent in prokaryotes. Through interkingdom interactions, some bacteria

might have inherited select genes from steroid hormone biosynthesis that enabled them to hijack

their host immune system and thus pave the way for establishing symbiotic interactions. The

limited distribution of this function across bacterial taxa associating with higher organisms

suggests these microbes occupy a unique niche that distinguishes them from other microbiome

members. Because of this selectivity, it may be that bacteria, such as *Endozoicomonas* that can

synthesize steroid hormones, may exert a disproportionate influence on holobiont physiology

compared to other bacteria. In the absence of *Endozoicomonas* during temperature-driven

disease[76], depletion of steroid precursors or reduced hormone availability may compromise host

immune regulation, creating favorable conditions for opportunistic pathogens to proliferate. Our

work reveals the unusual metabolic capabilities of coral-associated *Endozoicomonas* and the

functional role of this bacterial genus in the coral holobiont during heat stress, as they prime the

immune system of their hosts to withstand stressful events. We therefore propose a broader role

for steroids in the coral holobiont and highlight the significance of understanding coral symbiosis

in efforts to protect endangered coral reef habitats in a changing climate.

Data Availability

Metagenomic reads of the coral holobiont community consortium are deposited in NCBI under the

BioProject PRJNA1001615. Metagenomically assembled genomes and their annotations are

available on Zenodo (https://zenodo.org/records/15276884). The untargeted mass spectrometry

data is deposited on the Global Natural Products Social Molecular Network server

(https://gnps.ucsd.edu/) as raw data and can be accessed under the accession numbers

MSV000092488 and MSV000097641. Raw 16S rRNA gene amplicon libraries are deposited

under the BioProject PRJNA1262123 in NCBI. Raw sequences and reference genome for *E.*

acroporae AT381 are deposited under the accession number PRJNA1267713. Raw

transcriptomic sequences of the coral are deposited under the accession number PRJNA1414119.

De novo-assembled transcriptome, annotations, raw and normalised expression matrices are

provided under the Zenodo record number 18379496.

Acknowledgements

This work was supported by a Tamkeen award to S.A.A. (AD179) and by the NYUAD Research

Institute grants ADHPG-CGSB and CG009 to the Center for Genomics and Systems Biology

(CGSB) and Mubadala ACCESS, respectively. The authors acknowledge the NYU Abu Dhabi

Core Technology Platform facilities for access to mass spectrometry facilities and boats for

sampling. This research was carried out on the High-Performance Computing resources at New

York University Abu Dhabi. Figure 1 and 6e were created using https://www.biorender.com.

Author contributions

490 MAO, DA and SAA conceived the project; MAO, TDH and DA conducted the sampling; MAO,

491 TDH, LSYC, WJ and ARH processed the samples; TDH, SA, MAO, and JBR conducted lab

492 experiments; AM, SA, MAO, WSJ, ARH and GMR carried out data processing and analysis; AM,

493 MAO and SAA drafted the manuscript; AM, TDH, WSJ, SAA and MAO plotted all the figures; all

494 authors contributed to manuscript preparation; SAA and JAB acquired funding for the project.

Competing Interest Declaration

496 The authors declare no competing interests.

Additional information

498 This paper is accompanied by extended data and Supplementary Information.

References

1.   Rohwer, F., Seguritan, V., Azam, F. & Knowlton, N. Diversity and distribution of coral-associated bacteria. *Mar. Ecol. Prog. Ser.* **243**, 1–10 (2002).

2.   Bourne, D. G., Morrow, K. M. & Webster, N. S. Insights into the Coral Microbiome: Underpinning the Health and Resilience of Reef Ecosystems. *Annu. Rev. Microbiol.* **70**, 317–340 (2016).

3.   Ziegler, M. *et al.* Coral bacterial community structure responds to environmental change in a host-specific manner. *Nat. Commun.* **10**, 3092 (2019).

4.   Li, J. *et al.* Cultured Bacteria Provide Insight into the Functional Potential of the Coral-Associated Microbiome. *mSystems* **7**, e00327-22 (2022).

5.   Davy, S. K., Allemand, D. & Weis, V. M. Cell Biology of Cnidarian-Dinoflagellate Symbiosis. *Microbiol. Mol. Biol. Rev.* **76**, 229–261 (2012).

6.   LaJeunesse, T. C. *et al.* Systematic Revision of Symbiodiniaceae Highlights the Antiquity and Diversity of Coral Endosymbionts. *Curr. Biol.* **28**, 2570-2580.e6 (2018).

7.   Weis, V. M. Cell Biology of Coral Symbiosis: Foundational Study Can Inform Solutions to the Coral Reef Crisis. *Integr. Comp. Biol.* **59**, 845–855 (2019).

8.   Mohamed, A. R., Ochsenkühn, M. A., Kazlak, A. M., Moustafa, A. & Amin, S. A. The coral microbiome: towards an understanding of the molecular mechanisms of coral–microbiota interactions. *FEMS Microbiol. Rev.* **47**, fuad005 (2023).

9.   Meunier, V. *et al.* Microbes support enhanced nitrogen requirements of coral holobionts in a high CO_2 environment. *Mol. Ecol.* **30**, 5888–5899 (2021).

10. Voolstra, C. R. & Ziegler, M. Adapting with Microbial Help: Microbiome Flexibility Facilitates Rapid Responses to Environmental Change. *BioEssays* **42**, 2000004 (2020).

11. Neave, M. J., Michell, C. T., Apprill, A. & Voolstra, C. R. Endozoicomonas genomes reveal functional adaptation and plasticity in bacterial strains symbiotically associated with diverse marine hosts. *Sci. Rep.* **7**, 40579 (2017).

12. Ziegler, M., Seneca, F. O., Yum, L. K., Palumbi, S. R. & Voolstra, C. R. Bacterial community dynamics are linked to patterns of coral heat tolerance. *Nat. Commun.* **8**, 1–8 (2017).

13. Pogoreutz, C. *et al.* Dominance of *Endozoicomonas* bacteria throughout coral bleaching and mortality suggests structural inflexibility of the *Pocillopora verrucosa* microbiome. *Ecol. Evol.* **8**, 2240–2252

528   (2018).

14. Bayer, T. *et al.* The Microbiome of the Red Sea Coral Stylophora pistillata Is Dominated by Tissue-
530   Associated Endozoicomonas Bacteria. *Appl. Environ. Microbiol.* **79**, 4759–4762 (2013).

15. Roder, C., Bayer, T., Aranda, M., Kruse, M. & Voolstra, C. R. Microbiome structure of the fungid coral
532   *C tenactis echinata* aligns with environmental differences. *Mol. Ecol.* **24**, 3501–3511 (2015).

16. Pollock, F. J. *et al.* Coral-associated bacteria demonstrate phylosymbiosis and cophylogeny. *Nat.*
534   *Commun.* **9**, 4921 (2018).

17. Apprill, A., Weber, L. G. & Santoro, A. E. Distinguishing between Microbial Habitats Unravels
536   Ecological Complexity in Coral Microbiomes. *mSystems* **1**, e00143-16 (2016).

18. Maire, J. *et al.* Colocalization and potential interactions of *Endozoicomonas* and chlamydiae in
538   microbial aggregates of the coral *Pocillopora acuta*. *Sci. Adv.* **9**, eadg0773 (2023).

19. Raina, J.-B. *et al.* DMSP biosynthesis by an animal and its role in coral thermal stress response.
540   *Nature* **502**, 677–80 (2013).

20. Robbins, S. J. *et al.* A genomic view of the reef-building coral Porites lutea and its microbial
542   symbionts. *Nat. Microbiol.* **4**, 2090–2100 (2019).

21. Tandon, K. *et al.* Comparative genomics: Dominant coral-bacterium Endozoicomonas acroporae
544   metabolizes dimethylsulfoniopropionate (DMSP). *ISME J.* **14**, 1290–1303 (2020).

22. Pogoreutz, C. *et al.* Coral holobiont cues prime Endozoicomonas for a symbiotic lifestyle. *ISME J.* **16**,
546   1883–1895 (2022).

23. Wada, N. *et al.* High-resolution spatial and genomic characterization of coral-associated microbial
548   aggregates in the coral *Stylophora pistillata*. *Sci. Adv.* **8**, eabo2431 (2022).

24. Maire, J. & Van Oppen, M. J. H. A role for bacterial experimental evolution in coral bleaching
550   mitigation? *Trends Microbiol.* **30**, 217–228 (2022).

25. Hochart, C. *et al.* Ecology of Endozoicomonadaceae in three coral genera across the Pacific Ocean.
552   *Nat. Commun.* **14**, 3037 (2023).

26. Nesterov, O. *et al.* Validation and statistical analysis of the Group for High Resolution Sea Surface
554   Temperature data in the Arabian Gulf. *Oceanologia* **63**, 497–515 (2021).

27. Riegl, B. M., Benzoni, F., Samimi-Namin, K. & Sheppard, C. The Hermatypic Scleractinian (Hard)
Coral Fauna of the Gulf. in *Coral Reefs of the Gulf* (eds. Riegl, B. M. & Purkis, S. J.) vol. 3 187–224

(Springer Netherlands, Dordrecht, 2012).

28. Howells, E. J., Vaughan, G. O., Work, T. M., Burt, J. A. & Abrego, D. Annual outbreaks of coral

disease coincide with extreme seasonal warming. *Coral Reefs* **39**, 771–781 (2020).

29. Marangon, E. *et al.* Destabilization of mutualistic interactions shapes the early heat stress response of

the coral holobiont. *Microbiome* **13**, 31 (2025).

30. Rosales, S. M. *et al.* A meta-analysis of the stony coral tissue loss disease microbiome finds key

bacteria in unaffected and lesion tissue in diseased colonies. *ISME Commun.* **3**, 19 (2023).

31. Becker, C. C., Brandt, M., Miller, C. A. & Apprill, A. Microbial bioindicators of Stony Coral Tissue Loss

Disease identified in corals and overlying waters using a rapid field-based sequencing approach.

Environ. Microbiol. **24**, 1166–1182 (2022).

32. Pollock, F. J., Wada, N., Torda, G., Willis, B. L. & Bourne, D. G. White Syndrome-Affected Corals

Have a Distinct Microbiome at Disease Lesion Fronts. *Appl. Environ. Microbiol.* **83**, e02799-16 (2017).

33. Kanehisa, M. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res.* **28**, 27–30

(2000).

34. Lee, I., Ouk Kim, Y., Park, S.-C. & Chun, J. OrthoANI: An improved algorithm and software for

calculating average nucleotide identity. *Int. J. Syst. Evol. Microbiol.* **66**, 1100–1103 (2016).

35. Eren, A. M. *et al.* Community-led, integrated, reproducible multi-omics with anvi'o. *Nat. Microbiol.* **6**,

3–6 (2020).

36. Keller-Costa, T. *et al.* Metagenomics-resolved genomics provides novel insights into chitin turnover,

metabolic specialization, and niche partitioning in the octocoral microbiome. *Microbiome* **10**, 151

(2022).

37. Ochsenkühn, M. A., Röthig, T., D'Angelo, C., Wiedenmann, J. & Voolstra, C. R. The role of floridoside

in osmoadaptation of coral-associated algal endosymbionts to high-salinity conditions. *Sci. Adv.* **3**,

e1602047 (2017).

38. Vraspir, J. M. & Butler, A. Chemistry of Marine Ligands and Siderophores. *Annu. Rev. Mar. Sci.* **1**,

43–63 (2009).

39. Barbeau, K., Rue, E. L., Bruland, K. W. & Butler, A. Photochemical cycling of iron in the surface

ocean mediated by microbial iron(iii)-binding ligands. *Nature* **413**, 409–413 (2001).

40. Amin, S. A. *et al.* Photolysis of iron–siderophore chelates promotes bacterial–algal mutualism. *Proc.*

Natl. Acad. Sci. **106**, 17071–17076 (2009).

41. Carbonetti, N. H. & Williams, P. H. A cluster of five genes specifying the aerobactin iron uptake

system of plasmid ColV-K30. *Infect. Immun.* **46**, 7–12 (1984).

42. Li, C. *et al.* Aerobactin-Mediated Iron Acquisition Enhances Biofilm Formation, Oxidative Stress

Resistance, and Virulence of Yersinia pseudotuberculosis. *Front. Microbiol.* **12**, 699913 (2021).

43. Schwyn, B. & Neilands, J. B. Universal chemical assay for the detection and determination of

siderophores. *Anal. Biochem.* **160**, 47–56 (1987).

44. Küpper, F. C., Carrano, C. J., Kuhn, J.-U. & Butler, A. Photoreactivity of Iron(III)−Aerobactin:

Photoproduct Structure and Iron(III) Coordination. *Inorg. Chem.* **45**, 6028–6033 (2006).

45. Miethke, M. & Marahiel, M. A. Siderophore-Based Iron Acquisition and Pathogen Control. *Microbiol.*

Mol. Biol. Rev. **71**, 413–451 (2007).

46. Rey, M. W. *et al.* Complete genome sequence of the industrial bacterium Bacillus licheniformis and

comparisons with closely related Bacillusspecies. *Genome Biol.* **5**, r77 (2004).

47. Meziti, A. *et al.* The Reliability of Metagenome-Assembled Genomes (MAGs) in Representing Natural

Populations: Insights from Comparing MAGs against Isolate Genomes Derived from the Same Fecal

Sample. *Appl. Environ. Microbiol.* **87**, e02593-20 (2021).

48. Ding, J.-Y., Shiu, J.-H., Chen, W.-M., Chiang, Y.-R. & Tang, S.-L. Genomic Insight into the Host–

Endosymbiont Relationship of Endozoicomonas montiporae CL-33T with its Coral Host. *Front.*

Microbiol. **7**, (2016).

49. Sun, F., Yang, H., Zhang, X. & Shi, Q. Metabolic and metatranscriptional characteristics of corals

bleaching induced by the most severe marine heatwaves in the South China Sea. *Sci. Total Environ.*

858, 160019 (2023).

50. Williams, A. *et al.* Multi-omic characterization of the thermal stress phenome in the stony coral

Montipora capitata. *PeerJ* **9**, e12335 (2021).

51. Howells, E. J., Abrego, D., Vaughan, G. O. & Burt, J. A. Coral spawning in the Gulf of Oman and

relationship to latitudinal variation in spawning season in the northwest Indian Ocean. *Sci. Rep.* **4**,

7484 (2014).

52. Sood, U., Singh, Y., Shakarad, M. & Lal, R. Highlight on Engineering *Mycobacterium smegmatis* for

testosterone production. *Microb. Biotechnol.* **10**, 73–75 (2017).

53. Hambleton, E. A. *et al.* Sterol transfer by atypical cholesterol-binding NPC2 proteins in coral-algal

symbiosis. *eLife* **8**, e43923 (2019).

54. Oren, A. Thermodynamic limits to microbial life at high salt concentrations. *Environ. Microbiol.* **13**,

1908–1923 (2011).

55. Walker, W. H. Testosterone signaling and the regulation of spermatogenesis. *Spermatogenesis* **1**,

116–120 (2011).

56. Carlson, J. C., Gruber, M. Y. & Thompson, J. E. A Study of the Interaction between Progesterone and

Membrane Lipids[*]. *Endocrinology* **113**, 190–194 (1983).

57. Bergstrand, L. H., Cardenas, E., Holert, J., Van Hamme, J. D. & Mohn, W. W. Delineation of Steroid-

Degrading Microorganisms through Comparative Genomic Analysis. *mBio* **7**, e00166-16 (2016).

58. Chiang, Y., Wei, S. T., Wang, P., Wu, P. & Yu, C. Microbial degradation of steroid sex hormones:

implications for environmental and ecological studies. *Microb. Biotechnol.* **13**, 926–949 (2020).

59. Arp, G. *et al.* Gut Bacteria Encode Reductases that Biotransform Steroid Hormones. Preprint at

https://doi.org/10.1101/2024.10.04.616736 (2024).

60. Dlugovitzky, D. G., Fontela, M. S., Martinel Lamas, D. J., Valdez, R. A. & Romano, M. C.

Mycobacterium smegmatis synthesizes in vitro androgens and estrogens from different steroid

precursors. *Can. J. Microbiol.* **61**, 451–455 (2015).

61. Tan, E. S. *et al.* Does estrogen regulate vitellogenin synthesis in corals? *Comp. Biochem. Physiol. A.*

Mol. Integr. Physiol. **255**, 110910 (2021).

62. Twan, W.-H., Hwang, J.-S. & Chang, C.-F. Sex Steroids in Scleractinian Coral, Euphyllia ancora:

Implication in Mass Spawning. *Biol. Reprod.* **2260**, 2255–2260 (2017).

63. Rougée, L. R. A., Richmond, R. H. & Collier, A. C. Molecular reproductive characteristics of the reef

coral Pocillopora damicornis. *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* **189**, 38–44 (2015).

64. Chiu, Y.-L., Shikina, S., Yoshioka, Y., Shinzato, C. & Chang, C.-F. De novo transcriptome assembly

from the gonads of a scleractinian coral, Euphyllia ancora: molecular mechanisms underlying

scleractinian gametogenesis. *BMC Genomics* **21**, 732 (2020).

65. Vilela, C. L. S. *et al.* Estrogen induces shift in abundances of specific groups of the coral microbiome.

Sci. Rep. **11**, 2767 (2021).

66. Twan, W.-H. *et al.* Hormones and reproduction in scleractinian corals. *Comp. Biochem. Physiol. A.*

Mol. Integr. Physiol. **144**, 247–253 (2006).

67. Petersen, S., Chen, T., Lawrence, D. et al. TRAF2 is a biologically important necroptosis suppressor. Cell Death Differ 22, 1846–1857 (2015).

68. Mukundan L, Bishop GA, Head KZ, Zhang L, Wahl LM, Suttles J. TNF receptor-associated factor 6 is an essential mediator of CD40-activated proinflammatory pathways in monocytes and macrophages. J Immunol. 15;174(2):1081-90 (2005).

69. Amatya N, Garg AV, Gaffen SL. IL-17 Signaling: The Yin and the Yang. Trends Immunol. 38(5):310-322.(2017).

70. Kingsolver, MB & Hardy, R.W. Making connections in insect innate immunity, Proc. Natl. Acad. Sci. U.S.A. 109 (46) 18639-18640 (2012).

71. Palmer CV. Immunity and the coral crisis. Commun Biol. 9;1:91. doi: 10.1038/s42003-018-0097-4 2018.

72. Virginia M Weis, Cell Biology of Coral Symbiosis: Foundational Study Can Inform Solutions to the Coral Reef Crisis, Integrative and Comparative Biology, Volume 59, Issue 4, Pages 845–855 (2019).

73. Miller, D.J., Hemmrich, G., Ball, E.E. *et al*. The innate immune repertoire in Cnidaria - ancestral complexity and stochastic gene loss. Genome Biol 8, R59 (2007).

74. Augustin R, Bosch TC. Cnidarian immunity: a tale of two barriers. Adv Exp Med Biol. 2010;708:1-16.

 Main Figures and Figure Legends

Figure 1. Sampling, analysis, and validation experiments schemes. **a**, *Acropora pharaonis* colonies were sampled across a temperature gradient before gDNA and metabolites were extracted for microbiome and metabolome analyses. Samples were collected during thermal stress at 32°C and 34°C (June/August) and recovery at 27°C (October). Additional samples showing white syndrome disease phenotypes at 34°C were also collected (L: lesion, AL: adjacent to the lesion). Samples were processed for 16S rRNA gene amplicon, shotgun metagenome, and untargeted metabolome analyses. **b**, To test the hypothesis derived from the multiomics data, experiments were conducted on three coral coral-associated *Endozoicomonas* to examine siderophore-mediated iron acquisition and steroid production. **c**, Transcriptomics was then used to examine the effect of steroid hormones on the coral immune response during thermal stress.

Figure 2. Integrated microbiome/metabolome analysis of the coral holobiont. a, Principal
component analysis (PCA) of the metabolomics data (n=5878 putative molecular features).
Samples collected at 27°C and 32°C cluster together and are significantly different from samples
collected at 34°C. **b**, Principal coordinate analysis (PCoA) of microbial communities using Bray-
Curtis distance shows significant differences between the diseased samples (i.e., AL and L) and
all other samples. **c**, ASV-based bacterial community structure across temperatures and disease
samples at the family level. Each bar plot represents the bacterial abundance of a coral colony.

692   The top 20 bacterial families are shown. **d**, Distribution of metagenomic reads from coral

693   holobiont metagenomes across coral, *Cladocopium spp*. (algal symbiont) and putative microbial

694   members. **e**, Distribution of host-algal symbiont-bacterial genes (left) and metabolites (right)

695   across the top-most abundant KEGG metabolic pathways. Numbers in parentheses refer to the

696   number of genes and metabolites shown in each pathway, respectively.

Figure 3. Abundance of *Endozoicomonas* MAG10 and the putative functional importance
of *Endozoicomonas*. a, Relative abundance of tissue-associated MAGs inferred by read
recruitment from metagenomes across all samples. Taxonomic classifications of MAGs are
shown in SI Table 5. **b**, *Endozoicomonas* pangenome and biosynthetic gene clusters (BGCs) of
29 genomes and MAGs, including MAG10 and *E. acroporae* AT381. *Spartinivicinus marina*
SM1973 was included to represent a free-living member of the Oceanospirillales as a close free-
living isolate to *Endozoicomonas.* Pangenomic analyses based on the occurrence of gene
clusters were implemented in Anvi'o[34]. The central dendrogram represents hierarchical
clustering based on the presence (colored) or absence (opaque) of 34,846 gene clusters.
Branches were colored based on core, accessory or singleton/doubleton genes. Each track in
the pangenome represents a single genome colored according to its isolation source. Genomes
were clustered based on their pairwise average nucleotide identity (ANI) values shown in the
heatmap, which reveals high similarity between MAG10 and *E. acroporae* genomes. Distribution
of BGCs across genomes is shown below the heatmap. Key genes related to aerobactin
biosynthesis (*iucC*) and degradation of steroid hormones (*kstD*), galactose (*galM*), and histidine
(*hutH*) are labeled on the periphery of the pangenome.

Figure 4. Utilization potential of the siderophore aerobactin for iron acquisition in
***Endozoicomonas*. a**, Genomic neighborhood organization of the aerobactin BGC in genomes
of MAG10, *Endozoicomonas* spp. symbiotic with *Acropora* corals, soft corals, their closest free-
living isolate, *S. marinus* SM1973, and the aerobactin-producer *Klebsiella pneumoniae*. Shading
across genes represents gene synteny of aerobactin biosynthesis (blue), aerobactin uptake
(orange), or other genes (grey). Aerobactin is biosynthesized by 4 genes, one of which is
duplicated (*iucA/iucC*). Although the aerobactin biosynthesis operon is partially truncated in the
MAG10 scaffold, we confirmed that the only partially truncated gene is the duplicated *iucA/iucC*.
The TonB-dependent receptor refers to the aerobactin transporter gene, *iutA,* which is proximal
to the BGC of SM1973 and K. *pneumoniae.* Among all coral-associated *Endozoicomonas,* only
E. acroporae AT381 has *iutA* ~11kb distal to the BGC; all other *Endozoicomonas* genomes
display a distal ortholog/paralog of *iutA* elsewhere in the genome (>11kb away from the
biosynthesis BGC) **b**, CAS assay of cultures of two *E. acroporae* strains (AT381 and Acr14),
encoding the aerobactin cluster, grown in iron-limited media on agar. Siderophore activity was
then assessed through the formation of halos (black arrow) around bacterial colonies. * indicates
isolation from hard coras from the genus *Acropora.*

Figure 5. Putative steroid hormone biosynthesis/degradation in the holobiont across the temperature gradient and an abbreviated steroid hormone production pathway by *Endozoicomonas*. a, Steroid biosynthesis and degradation pathways based on coral and Endozoicomonas MAG10 contigs. Arrows indicate genes, and arrowheads correspond to the number of genes carrying a transformation. The relative abundance of putative steroid hormones is presented along with adjusted p-values, calculated using the Benjamini-Hochberg method, based on ANOVA results that assess the impact of temperature on changes in their abundance. Middle: Circos plot depicting all detected steroid-related metabolites and their connection to genes of *Endozoicomonas* (Endozoic.), the coral host, or *Cladocopium* spp. (Cl). The plot shows that steroid degradation is restricted to *Endozoicomonas*, steroid biosynthesis is mostly dominated by *Cladocopium* and steroid hormone biosynthesis is mostly dominated by the coral. Colors indicate different pathways (brown=steroid biosynthesis, red=steroid hormone biosynthesis, blue=steroid degradation, black=metabolites). **b**, Presence/absence of steroid degradation genes in the holobiont. Note that microbial contigs excluded contigs used to construct the bacterial MAGs. cholesterol oxidase; choD, beta-hydroxysteroid dehydrogenase; βHSD, 3-oxo-5alpha-steroid 4-dehydrogenase; tesI, steroid Delta-isomerase; ksi, 3-oxosteroid 1-dehydrogenase; kstD, 3-ketosteroid 9alpha-monooxygenase subunit A; kshA, 3-ketosteroid 9alpha-monooxygenase subunit B; kshB, 3-hydroxy-9,10-secoandrosta-(10)1,3,5-triene-9,17-dione monooxygenase; hsaA, 3,4-dihydroxy-9,10-secoandrosta-(10)1,3,5-triene-9,17-dione 4,5-dioxygenase; hsaC, 4,5:9,10-diseco-3-hydroxy-5,9,17-trioxoandrosta-2,(10)1-diene-4-oate hydrolase; hsaD, HIP-CoA ligase; fadD3. **c, d and e,** Steroid hormone production by three *Endozoicomonas* strains. Two *E. acroporae* (AT381 and Acr 14) and one *E. montiporae* strains were incubated with or without testosterone (**c**) , Androstenedione (**d**), or cholesterol (**e**), at 26°C and 34°C. Heatmaps show the normalized mean relative abundance of each metabolite in each condition, with * indicating statistical significance of their treatment relative to controls (T-test, adjusted p-value <0.05). Statistical significance was not calculated for

testosterone/androstenedione in **c, d** because both were added to the media. All detected

metabolites were annotated based onaccurate mass within 5 ppm and high-resolution MS/MS

fragmentation of standards. Additionally, a testosterone standard was used to further confirm its

production using the aforementioned criteria and retention time. **c,** Novel pathway for the

production of progesterone and testosterone inferred from metabolomics analysis. Dotted lines

indicate the absence of genes involved in these specific metabolic reactions in the

Endozoicomonas genomes.

Figure 6. Activation of the coral innate immune response by progesterone and testosterone during heat stress. a, Uptake and accumulation of [13]C-labeled testosterone in coral tissues over a 48-h exposure period. Peak area was normalised to dry extract weight. **b**, Experimental design. *Acropora* colonies were exposed to acute thermal stress with or without the addition of progesterone or testosterone. Hormones were administered at 10 nM at the start of the experiment at 28°C. Three hours later, the first sampling point (T1) was collected for RNA-seq analysis, followed by a second hormone dose at 6 h. The temperature was ramped up from 28°C to 33°C over 3 h after T1 collection, and samples were collected at T2 along with untreated controls. Final sampling (T3) was conducted after 24 h of exposure at 33°C. **c**, Maximum quantum yield of photosystem II (F_v/F_m) was measured daily as an indicator of photophysiological stress and a proxy for the coral holobiont health. **d**, Heatmap showing normalized expression levels of selected immunity-related genes differentially regulated across treatments and time points. Significantly differentially expressed genes at 6 hours compared to controls (FDR < 0.05) were highlighted by asterisks. All differential expressions were based on biological triplicates. **e**, Pathway diagram based on KEGG annotations highlighting genes associated with innate immune pathways, including immune receptors, intracellular signaling cascades, and downstream effector responses. Tumor necrosis factor receptor 1 (TNFR1), TNF receptor–associated factor 2 (TRAF2), TNF receptor–associated factor 6 (TRAF6), TAK1-binding protein 2 (TAB2), Mitogen-activated protein kinase kinase 3 (MKK3), Mitogen-activated protein kinase kinase 4/7 (MKK4/7), Nuclear factor kappa-B subunit 1/2 (NFKB1/2), NF-κB inhibitor alpha (IκBα / NFKBIA), Inhibitor of nuclear factor κB kinase subunit alpha/beta

(IKBKA/B), NF-κB essential modulator (NEMO / IKBKG), Activator protein 1 (AP-1), C-type lectin
domain family 4 member L (CLEC4L), Retinoic acid–inducible gene I (RIG-I), Interleukin-17
receptor A (IL17RA), Signal transducer and activator of transcription 5A (STAT5A), Cytokine
receptor domeless (DOME), Spleen tyrosine kinase (SYK), B-cell lymphoma 2 (*BCL2*), B-cell
lymphoma extra-large (*BCL-XL*), baculoviral IAP repeat-containing proteins 2 and 3 (*BIRC2/3*),
tumor necrosis factor alpha–induced protein 3 (*TNFAIP3*), cytokine-inducible SH2-containing
protein (*CISH*), suppressor of cytokine signaling 2 (*SOCS2*), heat shock protein 90 alpha
(*HSP90A*).