Elevation, drainage, and spatially heterogeneous vulnerability of tidal marshes to sea level rise

Man Qi[*1,2](manqistar@gmail.com), Keryn Gedan[2] (kgedan@gwu.edu)

1 Key laboratory of Geographical Process Analysis & Simulation of Hubei Province/College of Urban and Environmental Science, Central China Normal University, Wuhan 430079, China

2 Department of Biological Sciences, George Washington University, 800 22[nd] St NW, Washington, DC 20052, USA

Abstract:

Marsh plants regulate marsh surface elevation through organic matter accumulation and sediment capturing, thus feedback loops and relationships between vegetation and marsh elevation form the basis of numerical models that estimate and predict marsh vulnerability to Sea Level Rise (SLR). However, with interior portions of shallow-sloping microtidal marshes suffering much more extensive inundation relative to edge marshes, it is necessary to identify factors in addition to elevation that create spatial heterogeneity in flood vulnerability. Here we studied the influence of drainage on vulnerability of the marsh interior and the marsh edge. Specifically, we compared the deviation of *in situ* observed hydrological metrics for surface (hydroperiod, daily inundation duration) and subsurface (soil saturation index, or SSI) soil levels of interior and edge marsh from expected values estimated using elevation and tidal regime alone. We further analyzed the bias that hydrological deviation introduced to marsh vulnerability estimation using growth response curves of typical marsh macrophytes to hydroperiod. We found that the water table fluctuation was attenuated from marsh edge to interior, which resulted in a higher sensitivity of interior marsh to elevation deficiency. At the brim of marsh collapse in the marsh interior, hydroperiod and SSI increased dramatically; they were 20-50% higher than at the same elevation at the marsh edge, which greatly reduces the window of opportunity for interior marsh plant communities to adapt to SLR. Our findings

demonstrate the importance of drainage in explaining spatial patterns in vulnerability

across the marsh platform, which should be considered in future empirical studies and

numerical simulations to develop practical tools for coastal management.

Key words:

Marsh stability; Drainage; Inundation; Groundwater; Plant dieback

Introduction

Marsh macrophytes regulate marsh elevation both through organic matter

accumulation and sediment capture (Mudd et al. 2010, Baustian et al. 2012, Kirwan

and Mudd 2012, Bouma et al. 2013). The long-term stability of marshes can be

explained by interactions among sea level, marsh surface elevation, plant productivity,

and sediment accretion, which interact to regulate marsh elevation in equilibrium with

mean sea level (Baustian et al. 2012, Morris et al. 2013, Kirwan et al. 2016, Cahoon et

al. 2019, FitzGerald and Hughes 2019, Cahoon et al. 2021).

The persistence of marsh ecosystems is thought to depend on the production of

aboveground and belowground biomass by marsh plants, which can raise marsh

surface elevation and avoid plant inundation caused by sea level rise (Morris et al.

2002, Kirwan and Guntenspergen 2012, Beckett et al. 2016, Gonneea et al. 2019).

Specifically, marsh ecosystems will be stable in the face of sea level rise when marsh

surface elevation is greater than the optimum for primary production, as an increase in

sea level will stimulate primary production, leading to increased sedimentation and

organic matter accretion. Otherwise, where surface elevation relative to sea level is

lower than the optimum, the system will be unstable, as rising sea levels reduce

primary production, leading to declines in mineral sedimentation and organic matter

accretion, and further elevation loss (Morris et al. 2002). Therefore, the growth curve

of marsh plants to varying elevation has been widely applied across entire marsh

platforms in empirical studies (Morris et al. 2002, Baustian et al. 2012, Kirwan and

Guntenspergen 2012, Voss et al. 2013, Kirwan and Guntenspergen 2015) and

eco-geomorphologic models (Kirwan and Murray 2007, L. et al. 2010, Marani et al.

2013, Mariotti and Carr 2014, D'Alpaos and Marani 2016, Mariotti 2016, Duran

Vinent et al. 2021) to estimate the stability of marsh platforms colonized by various

macrophytes.

However, whether the hydrological stress experienced by marsh plant can be

represented by elevation and tidal regime alone is still questionable, as hydrological

conditions, especially groundwater dynamics can also be greatly influenced by

distance from outlets such as tidal creeks and drainage conditions (Ursino et al. 2004,

Van Putte et al. 2022, Xin et al. 2022). For example, in field studies, wave attenuation

by marsh vegetation was observed as reduction in wave height over distance traveled

(Möller et al. 2014, Stark et al. 2015). In addition, field observations and numerical

simulations demonstrated that as the tide propagates toward the marsh interior, marsh

subsurface hydrology is less responsive to tidal height, especially in interior marsh

that is distant from tidal channels, which resulted in tide-induced water table

fluctuations which attenuated from creek to marsh interior (Ursino et al. 2004, Cao et

al. 2012a, Xin et al. 2013b, Byers and Chmura 2014, Xin et al. 2022). In addition,

compared with creek banks, soil condition and vegetation in interior marsh is

regulated more by rainfall and evapotranspiration than by tide (Xin et al. 2013a, Xin

et al. 2017, Walker et al. 2019). These suggest the influence of elevation- and tidally-

derived hydrological regimes weaken with increasing distance from tidal creeks, as

the immediate effects of drainage diminish.

Identifying the deviation of *in situ* hydrological condition from elevation- and

tidally- derived estimates is important, especially for interior marsh, as poorly-drained

interior marsh in microtidal marshes has experienced the greatest habitat losses

resulting from sea level rise, for example in the Mississippi Delta, Chesapeake Bay,

and Venice Lagoon (D'Alpaos and Marani 2016, Kearney and Turner 2016, Smith et

al. 2016, Mitchell et al. 2017, Ortiz et al. 2017, Schepers et al. 2017, Wang et al.

2021). Besides, an increasing body of empirical studies are applying the results of

mesocosm experiments conducted in tidal creeks designed to test the effects of

elevation relative to sea level (i.e. marsh organ experiments) without considering the

spatial variation in drainage conditions that occur across a marsh platform (Kirwan

and Guntenspergen 2012, Adam Langley et al. 2013, Kirwan and Guntenspergen 2015,

Janousek et al. 2016, Janousek et al. 2020, McLain et al. 2020, Payne et al. 2021).

Furthermore, numerical models of marsh vulnerability to accelerated SLR seldom

incorporate drainage conditions or subsurface hydrology into the current framework

of a dynamic marsh surface elevation that is driven by the feedback between elevation,

tide, plant growth and sediment deposition (Kirwan and Murray 2007, L. et al. 2010,

Marani et al. 2013, Mariotti and Carr 2014, D'Alpaos and Marani 2016, Mariotti 2016,

Duran Vinent et al. 2021).

Although some prior studies have noted the deficiency of overlooking

heterogeneity in drainage conditions (Morris et al. 2013, Kirwan and Guntenspergen

2015, McLain et al. 2020, Van Putte et al. 2022), to our knowledge, no field studies

have been conducted to quantify the deviation from elevation- and tidally- based

estimations that can be attributed to spatial variation in drainage conditions. To

address this, we selected two deteriorating marsh platforms in the Chesapeake Bay,

United States. At each marsh, we monitored water levels and soil moisture in three

different zones -- vegetated, plant dieback, and ponded -- interior marsh. For each

zone, we estimated the difference between elevation- and tidally- derived hydrological

regimes and *in situ* observations of hydroperiod, daily inundation duration, and soil

aeration. We also analyzed the influence of hydroperiod deviation between *in situ*

observations and elevation- and tidally- derived estimations on marsh stability using

empirical growth response curves of typical marsh macrophytes from the region. By

recognizing the effect of drainage on spatially variation in marsh stability, we hope to

provide a resource for future studies and models of marsh stability, which are

important and necessary tools for coastal management.

Study area

The Chesapeake Bay is one of the largest microtidal estuaries in the world, and

has experienced increasing deterioration in recent decades (Kearney et al. 1988,

Beckett et al. 2016). Within the Bay, we selected Deal Island (DI) and Farm Creek

Marsh (FCM), MD as study sites. These two marshes are in the early stages of transition from vegetated marsh to open pond (Walker et al. 2019, Qi et al. 2021). Areas in various states of deterioration can be found at both sites, but in general, deterioration at both sites is more apparent in marsh interiors, where drainage is poor and marsh accretion rates are lower compared with marsh edge (Walker et al. 2019, Qi et al. 2021).

Both DI and FCM are brackish marshes, but DI has a higher salinity range (7-15 ppt) than FCM (2-5 ppt) based on measurements in summer of 2019. Emergent vegetation in both marshes is typical of brackish marshes of the region, and includes grasses, sedges, and rushes of the species *Spartina patens*, *Distichlis spicata, Spartina alterniflora, Schoenoplectus americanus* and *Juncus roemarianus.* The dominant species is *S. alterniflora* in DI, and *S. americanus* in FCM, reflecting differences in salinity levels and the salinity tolerances of these two species. In DI, inundation stressed *S. alterniflora* colonize a spatially homogenous substrate of weak soil strength as a sparse meadow, whereas in deteriorating areas of FCM, *S. americanus* forms a heterogenous surface of hollows and hummocks.

At the marsh interior of each site, we selected three zones: vegetated, dieback, and pond, to capture the hydrological alteration along the trajectory of marsh deterioration. In DI, we monitored a tidal creek bank located 800 m from interior marsh as a reference point where the water table was likely to resemble an unrestricted tidal fluctuation. We did not instrument such a reference point at FCM because we found that the tidal creek water levels at FCM and DI were highly correlated. Therefore, we took hourly tidal height readings at Bishops Head station, MD from NOAA tide and current website, (Station ID: 8571421, https://tidesandcurrents.noaa.gov/stationhome.html?id=8571421) to represent tidal height at the local scale from FCM to DI. This Bishops Head station is 11 km southeast of FCM, and 12 km Northwest of DI.

<Fig. 1>

Methodology

In situ groundwater observation

At each site, three groundwater wells were installed in the vegetated zone,

dieback zone, and pond, respectively, in April 2019 to monitor water table dynamics.

In DI, a groundwater well was installed at a nearby tidal creeks (as shown in Fig. 1) in

comparison with the interior part of the marsh.

Groundwater level data was collected at 15 min interval from May 2019 to

March 2020 with HOBO water level loggers (U20L -04). Marsh surface elevation of

each zone was measured within a 0.5 m radius from each groundwater well in May

2019 using a Trimble R8 Real Time Kinematic Global Positioning System (RTK-GPS)

to reference water table data to the NAVD88 datum and to allow for comparison

within and between sites. The flooding regime was estimated using tidal height data

from the Bishops Head and *in situ* marsh elevation collected at each zone, as

described above.

Hydrological metrics

We selected three metrics to represent surface inundation, daily inundation stress

and soil aeration: 1) hydroperiod, 2) daily inundation duration, and 3) soil saturation

index (SSI). Hydroperiod is the fraction of time   marsh was flooded over the

observation period. SSI is the fraction of the tidal period over which the soil remains

fully saturated at a given depth, and SSI at depth=0.0 m is equivalent to the

hydroperiod (Xin et al. 2010).

In situ hydrological metrics were calculated from observations of groundwater

levels and marsh surface elevation. As a counterpart, estimated hydrological metrics

was calculated from tidal height and marsh surface elevation. Comparison between in

situ metrics and estimated metrics was conducted to analyze the hydrological bias

caused by spatially dependent drainage.

Influence of hydrological deviation on marsh vulnerability

To understand the deviation of *in situ* observations from estimations based solely

on elevation and tidal fluctuations, and how these affect forecasts of marsh

vulnerability to deterioration, we extracted growth response curves of typical

macrophytes to predicted and observed hydroperiods. Growth response curves were

extracted from the literature, from studies using marsh organ experiments in sites on

the Atlantic Coast of the United States. Growth response curves, including the ones

used in this study, have been widely used to estimate and forecast marsh vulnerability

to sea level rise.

We only adopted growth response curves that used plant biomass (aboveground

or belowground) or biomass change relative to inundation duration. Some other

response indicators were also used in relevant studies e.g. shoot/stem density, stem

length, but these were less comparable across studies and could not be related to the

hydrological metrics of interest in this study (Kirwan and Guntenspergen 2012, Adam

Langley et al. 2013, Janousek et al. 2020).

Based on these criteria, three empirical studies were identified with the variables

of interest for four out of five of the typical macrophytes (except *D. spicata*) at our

study sites. Cumulatively, these studies contained 246 measurements of aboveground

biomass and 178 measurements of belowground biomass. To synthesize, we

normalized aboveground biomass and belowground biomass of a given species from

each study separately with the equation of $x_n=[x_i - \min(x_i)]/[\max(x_i)-\min(x_i)]$. Then

we merged the normalized data (with a range of [0~1]) from different studies to form

a unified growth response curve. After a preliminary analysis, we found aboveground

and belowground biomass response curves had similar shapes (Appendix S1: Fig. S1),

therefore we merged the standardized aboveground biomass and belowground

biomass into a uniform biomass response curve to inundation using quadratic

regression.

204   <Table 1>

Results

Marsh elevation and hydrological series

In both sites, there was a difference in elevation of 0.22 m between healthy zones

and ponds (dashed lines in Fig. 2). In DI, the elevation difference was similar between

vegetated zone to dieback zone and to pond, while elevation in FCM did not vary

greatly between vegetated zone and dieback zone. However, there was a large

elevation difference between the transition and dieback zones with ponds.

The water table at DI, which was monitored at the bank of tidal creek, was

highly consistent with the tidal level measured at Bishops Head station (Fig, 2), and

exhibited a tidal range of approximately 60 cm (Fig. 3). In the interior marsh, the

water table levels at DI had a similar range of variability (20~30cm) as at FCM, but

the daily variation at DI was muted compared to FCM (Fig. 3).

Within each site, interior marsh had smaller water table fluctuations but higher

average water table height than nearby tidal creeks (Fig. 2~3), though there is slight

difference in the distribution of water table   across interior zones (Fig. 3).

221   <Fig. 2>

222   <Fig. 3>

Hydrological metrics deviation

The difference between in situ observed and estimated hydroperiod and SSI was

small at the tidal creek but increased within interior marsh (Fig. 4). Estimated

hydroperiod and SSI increased linearly with decreasing marsh surface elevation,

while observed hydroperiod and SSI increased nonlinearly with increasing elevation

and reached a maximum of 100% in several zones. Within any given zone, soil

saturation index at depth was generally higher than hydroperiod other than where both

reach a maximum of 100%. The deviation between estimated and in situ monitored

hydroperiod reached a maximum of 50% at dieback zone, and is 22-33% at the pond

and vegetated zones. In situ observed SSI at 0.1 m depth presented the greatest

deviation (50%) from estimations from the dieback zone, and observed SSI at 0.2 m

depth exhibited the greatest deviation from estimations of 40% in the vegetated zone.

236   <Fig. 4>

237   The distribution of daily inundation duration differed greatly between in situ

238   observations and estimations. Estimated daily inundation durations averaged between

239   4 and 8 hrs in all zones for both sites, and increased slightly with the degree of interior

marsh deterioration (Fig. 5). Whereas measured observations showed that in the majority of observed days (65% in DI, and 30% in FCM), vegetated marsh zones were continuously inundated for the entire day, which almost never occurred in the estimation. For dieback zones and ponds, continuous inundation occurred on 100% of days in DI, and 75% and 100% of days, respectively, in FCM.

<Fig. 5>

Bias in marsh vulnerability estimation

Among the four macrophytes, only *S. americanus* showed a unimodal growth curve that peaked at intermediate daily durations of inundation; the other three species' performance decreased gradually with increasing hydroperiod. For vegetated zones of the marsh interior, the estimated hydroperiod (dashed green lines in Fig. 6) is 22%~26% lower than in situ observations (solid green lines in Fig. 6). Therefore, if species performance is estimated on the basis of elevation and tidal fluctuations alone, this results in an overestimation of plant performance by 10%-25%. This result in general overestimation of marsh plant resilience to sea level rise, especially for *S. americanus*, the dominant species at FCM. At FCM, for example, whereas the vegetated zone's elevation was estimated to be at the species' optimal performance level, based on these hydrologic corrections, it was actually within a suboptimal range.

In the dieback and pond zones of DI, the estimated hydroperiod indicated a suboptimal but non-lethal status for all species, but in situ hydrologic observations suggests a hydroperiod that no species could survive, which more accurately reflected the vegetation condition. Similar bias also happened in ponds of FCM, but not in dieback zones of FCM, because there was a smaller elevation difference between healthy and dieback areas.

Therefore, plant performance and marsh resilience in the vegetated zones of interior marsh was overestimated for all species when elevation and tidally-derived hydrologic regimes were used

<Fig. 6>

Discussion

Tide station data was a good match to tidal creek water levels. However, within

the marsh platform, as the tide propagates over and through the peat matrix into marsh

interior, tide-induced water table fluctuations are greatly attenuated. Due to the

narrow fluctuation of the water table in interior marsh zones, even minor elevation

change can greatly shift the hydroperiod and period of soil saturation. These dynamics

are poorly captured when estimated using tidal height and marsh elevation. As a result

of the underestimation of hydroperiod and soil saturation in deteriorating microtidal

marshes, we have also underestimated the vulnerability of interior marsh to sea level

rise by applying elevation and tide-derived hydrological regimes to the whole marsh

platform.

Effect of elevation and drainage on hydrological regime

A decrease in the water table fluctuations in response to reduced drainage in

interior marshes has been previously observed in field investigations and numerical

simulations (Ursino et al. 2004, Cao et al. 2012a, Cao et al. 2012b, Byers and Chmura

2014, O'Connor and Moffett 2015). However, most observations were only conducted

for days and focus on transient asynchronization in groundwater table at different

distance from tidal creek, and seldom were further interpreted or upscaled to inform

difference in vegetation performance and marsh vulnerability between marsh edge

and marsh interior. With a year-long observation, we found estimates by elevation and

tidal height understimate hydroperiod in interior marsh by 25% ~ 50% (Fig. 4a).

Rather, this reduction in the tidal fluctuation in interior marsh is also greatly

influenced by dampening (by friction or water spreading) as tide propagates across the

marsh (Stark et al. 2015, Xin et al. 2022).

The difference in the fluctuation of the water table between marsh exterior and

interior marsh creates a divergent response in hydroperiod and SSI to elevation

variation (Fig. 4). At Deal Island, for example, there was little fluctuation at all in the

groundwater table within interior marsh (Fig. 3). Let's take an ideally occasion at

Deal Island that we can find marsh points with elevation ranging from 0 to 0.7 m in

marsh exterior zone and marsh interior zone We then assume marsh points within

each zone shared the same groundwater level as but differed in elevation. Finally we

can calculate the hydroperiod and SSI under different elevation levels at each section

(Fig. 7). We can found that when elevation drops from 0.36 m to 0.15 m, there is a

sharp increase in the hydroperiod in marsh interior (88% increase) relative to the

marsh creek bank (30% increase) (Fig.7). Interestingly, the hydroperiod and the SSI in

interior marsh is underestimated at the left side of the intersections of the curves,

whereas it is overestimated, albeit to a lesser degree, at the right side of the

intersections, indicating that at very high elevations of marsh, soils may be more

drained and less frequently inundated than expected (Fig. 7). Given that we observed

that deteriorated marsh stages are lower in elevation and peat collapse can be develop

fast following vegetation death (Day et al. 2011), we expect that elevation loss during

marsh deterioration means that the estimation of hydroperiod based on elevation can

quickly become out of date, and distort the true experience of plants in these

deteriorating microtidal marshes. Therefore, it is important to recognize the range of

water table fluctuations in interior marsh in order to better predict the plant response

within interior marshes and its feedbacks to marsh elevation gain and loss. Elevation

loss in interior marshes leads to pond expansion and habitat loss.

317   <Fig. 7>

Spatially heterogeneous marsh vulnerability

In many studies (D'Alpaos and Marani 2016, Ortiz et al. 2017, Schepers et al.

2017), the higher vulnerability of interior marsh is attributed to a lower accretion rate

in marsh interiors than at marsh edges (Kirwan et al. 2016). Our study suggests that

the higher inundation at the surface and saturation of the subsurface layer in interior

marshes, which is caused by poor drainage, should be a more direct factor. The high

frequency of saturation and inundation reduces belowground biomass production,

often before aboveground biomass reductions are observed (Turner et al. (2004),

meaning that these effects can be easily missed in monitoring efforts. This might

explain the shallow subsidence rates observed in the Chesapeake Bay, where accretion

rates in interior marshes can be much higher than the rate of sea level rise, as high as

9-15 mm/yr, yet marsh surface elevation has decreased 1.8 ± 2.7 mm/yr on average,

and this occurred before any evidence of plant dieback was observed (Beckett et al.

2016).

This deficiency of using elevation as the sole indicator of marsh stability, and the importance of drainage in understanding spatial heterogeneity in marsh vulnerability has been reported in prior studies. For example, Kirwan and Guntenspergen (2015) reported that *S. americanus* produced 2-4 times greater aboveground biomass in marsh organs than at the equivalent elevations in its natural habitat of interior marsh. Similarly, drainage associated with creek banks facilitated seedling survival and marsh stability in microtidal or macro marshes (Liu et al. 2020, Cao 2021). Therefore, it is necessary to consider spatial heterogeneity in drainage in future studies of marsh vulnerability and risk.

Besides overall higher vulnerability of interior marsh than marsh edge, the higher sensitivity of hydroperiod to a narrow range of elevation in interior marsh makes deterioration occur more quickly and with less warning than at more exterior locations. It may also leave interior marshes less resilient to minor disturbances because only a trivial loss of relative elevation, of about 15 cm in this study, can create a disproportional increase in hydroperiod that can greatly change the growing conditions and cause plant mortality or prevent recovery (Raposa et al. 2017, Watson et al. 2017, Qi et al. 2021). From this aspect, microtidal systems are much more vulnerable to sea level rise effects than meso- and macrotidal systems (Kirwan and Guntenspergen 2010, D'Alpaos et al. 2012).

At Deal Island and many other sites, plant displacement in interior marshes has been observed prior to pond formation and expansion (Raposa et al. 2017, Watson et al. 2017, Qi et al. 2021). Additionally, mortality of flood intolerant species, such as *S. patens* and *D. spicata,* occurred before colonization and replacement by flood tolerant species, such as *S. americanus* and *J. roemerianus* (Fig. 8). The elevation range of *S. americanus* is only a scant 4 to 5 cm lower than *S. patens* (Holmquist et al. 2021), despite a large difference in flood tolerance (Fig. 7, and Qi et al. 2021). It is likely that the steep response curve of hydroperiod and elevation in interior marshes didn't given sufficient window of opportunity for displacement to be accomplished, i.e. after preexisting plant die off and elevation loss, the hydroperiod has already shifted

outside the appropriate range of subsequent plant species before the latter can

establish. Therefore, understanding the elevation building capacity and flood

tolerance of different plant assemblages could improve predictions of the elevation

trajectory in deteriorating marshes (Kelleway et al. 2017).

<Fig. 8>

Achieving more accurate estimation of interior marsh

The deteriorating marshes selected for this study were located within the range of

elevation where elevation and tidally-based estimates severely underestimate

hydroperiod and SSI (Fig. 7). This is consistent with observations of rapid vegetation

displacement and pond expansion at these same sites over the past decade (Taylor et

al. 2020, Qi et al. 2021). Based on the relationship between observed hydroperiod and

elevation, we predict that marsh vulnerability may be overestimated for interior marsh

areas at higher-elevation or with greater elevation capital. This provides a hypothesis

that can be tested in future studies.

Current numeric models that simulate marsh vulnerability to sea level rise are

based on geomorphological drivers such as elevation, tidal range, and distance from

surface water features. In general, subsurface flow and its interaction with surface

flow are not incorporated in these models (Kirwan and Murray 2007, L. et al. 2010,

Marani et al. 2013, Mariotti and Carr 2014, D'Alpaos and Marani 2016, Mariotti 2016,

Duran Vinent et al. 2021). We urge future modeling studies to consider more than

surface elevation of the marsh platform, but also to incorporate subsurface

hydrological processes in order to more accurately reflect the hydrological stress on

marsh plants and the feedbacks to marsh platform integrity. In this way, the estimates

of marsh vulnerability and tools for marsh conservation and restoration can improve.

Empirical studies can also better incorporate additional measurements to

represent the factors influencing the drainage condition of a marsh. We recommend

considering the distance from tidal creeks and soil permeability in study designs and

monitoring efforts. These factors can be used to establish regression functions

between tidal and groundwater fluctuations to improve data interpretation and

interpret spatial variability in marsh plant performance, ponding, and ecosystem

deterioration in microtidal marshes.

Acknowledgement

This study was supported by Washington Biologist's Field Club (awarded to M

Qi), … We thank Jessica MacGregor for helping with the field work.

References

Adam Langley, J., T. J. Mozdzer, K. A. Shepard, S. B. Hagerty, and J. Patrick Megonigal. 2013. Tidal

marsh plant responses to elevated CO2, nitrogen fertilization, and sea level rise. Global

Change Biology **19**:1495-1503.

Baustian, J. J., I. A. Mendelssohn, and M. W. Hester. 2012. Vegetation's importance in regulating

surface elevation in a coastal salt marsh facing elevated rates of sea level rise. Global Change

Biology **18**:3377-3382.

Beckett, L. H., A. H. Baldwin, and M. S. Kearney. 2016. Tidal marshes across a Chesapeake Bay

subestuary are not keeping up with Sea-Level Rise. PLOS ONE **11**:e0159753.

Bouma, T. J., S. Temmerman, L. A. van Duren, E. Martini, W. Vandenbruwaene, D. P. Callaghan, T.

Balke, G. Biermans, P. C. Klaassen, P. van Steeg, F. Dekker, J. van de Koppel, M. B. de Vries,

and P. M. J. Herman. 2013. Organism traits determine the strength of scale-dependent

bio-geomorphic feedbacks: A flume study on three intertidal plant species. Geomorphology

180-181:57-65.

Byers, S. E., and G. L. Chmura. 2014. Observations on Shallow Subsurface Hydrology at Bay of

Fundy Macrotidal Salt Marshes. Journal of Coastal Research **30**:1006-1016.

Cahoon, D. R., J. C. Lynch, C. T. Roman, J. P. Schmit, and D. E. Skidds. 2019. Evaluating the

Relationship Among Wetland Vertical Development, Elevation Capital, Sea-Level Rise, and

Tidal Marsh Sustainability. Estuaries and Coasts **42**:1-15.

Cahoon, D. R., K. L. McKee, and J. T. Morris. 2021. How Plants Influence Resilience of Salt Marsh

and Mangrove Wetlands to Sea-Level Rise. Estuaries and Coasts **44**:883-898.

Cao, H. 2021. Tidal marsh dynamics in a changing climate: A study on the mechanisms of pioneer

establishment and erosion. Utrecht University.

Cao, M., P. Xin, G. Jin, and L. Li. 2012a. A field study on groundwater dynamics in a salt marsh -

Chongming Dongtan wetland. Pages 61-69   Ecological Engineering.

Cao, M., P. Xin, G. Q. Jin, and L. Li. 2012b. A field study on groundwater dynamics in a salt marsh -

Chongming Dongtan wetland. Ecological Engineering **40**:61-69.

D'Alpaos, A., C. Da Lio, and M. Marani. 2012. Biogeomorphology of tidal landforms: physical and

biological processes shaping the tidal landscape. Ecohydrology **5**:550-562.

D'Alpaos, A., and M. Marani. 2016. Reading the signatures of biologic–geomorphic feedbacks in

salt-marsh landscapes. Advances in Water Resources **93**:265-275.

Day, J. W., G. P. Kemp, D. J. Reed, D. R. Cahoon, R. M. Boumans, J. M. Suhayda, and R. Gambrell.

2011. Vegetation death and rapid loss of surface elevation in two contrasting Mississippi delta

salt marshes: The role of sedimentation, autocompaction and sea-level rise. Ecological

Engineering **37**:229-240.

Duran Vinent, O., E. R. Herbert, D. J. Coleman, J. D. Himmelstein, and M. L. Kirwan. 2021. Onset of

runaway fragmentation of salt marshes. One Earth **4**:506-516.

FitzGerald, D. M., and Z. Hughes. 2019. Marsh Processes and Their Response to Climate Change and

Sea-Level Rise. Annual Review of Earth and Planetary Sciences **47**:481-517.

Gonneea, M. E., C. V. Maio, K. D. Kroeger, A. D. Hawkes, J. Mora, R. Sullivan, S. Madsen, R. M.

Buzard, N. Cahill, and J. P. Donnelly. 2019. Salt marsh ecosystem restructuring enhances

elevation resilience and carbon storage during accelerating relative sea-level rise. Estuarine,

Coastal and Shelf Science **217**:56-68.

Holmquist, J. R., L. Schile-Beers, K. Buffington, M. Lu, T. J. Mozdzer, J. Riera, D. E. Weller, M.

Williams, and J. P. Megonigal. 2021. Scalability and performance tradeoffs in quantifying

relationships between elevation and tidal wetland plant communities. Marine Ecology

Progress Series **666**:57-72.

Janousek, C. N., K. J. Buffington, K. M. Thorne, G. R. Guntenspergen, J. Y. Takekawa, and B. D.

Dugger. 2016. Potential effects of sea-level rise on plant productivity: species-specific

responses in northeast Pacific tidal marshes. Marine Ecology Progress Series **548**:111-125.

Janousek, C. N., B. D. Dugger, B. M. Drucker, and K. M. Thorne. 2020. Salinity and inundation effects

on productivity of brackish tidal marsh plants in the San Francisco Bay-Delta Estuary.

Hydrobiologia **847**:4311-4323.

Kearney, M. S., R. E. Grace, and J. C. Stevenson. 1988. Marsh Loss in Nanticoke Estuary, Chesapeake

Bay. Geographical Review **78**:205-220.

Kearney, M. S., and R. E. Turner. 2016. Microtidal Marshes: Can These Widespread and Fragile

Marshes Survive Increasing Climate–Sea Level Variability and Human Action? Journal of

Coastal Research:686-699.

Kelleway, J. J., N. Saintilan, P. I. Macreadie, J. A. Baldock, and P. J. Ralph. 2017. Sediment and carbon

deposition vary among vegetation assemblages in a coastal salt marsh. Biogeosciences

14:3763-3779.

Kirwan, M. L., and G. R. Guntenspergen. 2010. Influence of tidal range on the stability of coastal

marshland. Journal of Geophysical Research: Earth Surface **115**.

Kirwan, M. L., and G. R. Guntenspergen. 2012. Feedbacks between inundation, root production, and

shoot growth in a rapidly submerging brackish marsh. Journal of Ecology **100**:764-770.

Kirwan, M. L., and G. R. Guntenspergen. 2015. Response of plant productivity to experimental

flooding in a stable and a submerging marsh. Ecosystems **18**:903-913.

Kirwan, M. L., J. A. Langley, G. R. Guntenspergen, and J. P. Megonigal. 2013. The impact of sea-level

rise on organic matter decay rates in Chesapeake Bay brackish tidal marshes. Biogeosciences

10:1869-1876.

Kirwan, M. L., and S. M. Mudd. 2012. Response of salt-marsh carbon accumulation to climate change.

Nature **489**:550-+.

Kirwan, M. L., and A. B. Murray. 2007. A coupled geomorphic and ecological model of tidal marsh

evolution. Proceedings of the National Academy of Sciences **104**:6118-6122.

Kirwan, M. L., S. Temmerman, E. E. Skeehan, G. R. Guntenspergen, and S. Fagherazzi. 2016.

Overestimation of marsh vulnerability to sea level rise. Nature Climate Change **6**:253.

L., K. M., G. G. R., D. A. Andrea, M. J. T., M. S. M., and T. Stijn. 2010. Limits on the adaptability of

coastal marshes to rising sea level. Geophysical Research Letters **37**.

Liu, Z., S. Fagherazzi, X. She, X. Ma, C. Xie, and B. Cui. 2020. Efficient tidal channel networks

alleviate the drought-induced die-off of salt marshes: Implications for coastal restoration and

management. Science of The Total Environment **749**:141493.

Marani, M., C. Da Lio, and A. D'Alpaos. 2013. Vegetation engineers marsh morphology through

multiple competing stable states. Proceedings of the National Academy of Sciences of the

United States of America **110**:3259-3263.

Mariotti, G. 2016. Revisiting salt marsh resilience to sea level rise: Are ponds responsible for

permanent land loss? Journal of Geophysical Research: Earth Surface **121**:1391-1407.

Mariotti, G., and J. Carr. 2014. Dual role of salt marsh retreat: Long-term loss and short-term resilience.

Water Resources Research **50**:2963-2974.

McLain, N., L. Camargo, C. R. Whitcraft, and J. G. Dillon. 2020. Metrics for Evaluating Inundation

Impacts on the Decomposer Communities in a Southern California Coastal Salt Marsh.

Wetlands **40**:2443-2459.

Mitchell, M., J. Herman, D. M. Bilkovic, and C. Hershner. 2017. Marsh persistence under sea-level rise

is controlled by multiple, geologically variable stressors. Ecosystem Health and Sustainability

3:1379888.

Möller, I., M. Kudella, F. Rupprecht, T. Spencer, M. Paul, B. K. van Wesenbeeck, G. Wolters, K.

Jensen, T. J. Bouma, M. Miranda-Lange, and S. Schimmels. 2014. Wave attenuation over

coastal salt marshes under storm surge conditions. Nature Geoscience **7**:727.

Morris, J., K. Sundberg, and C. Hopkinson. 2013. Salt Marsh Primary Production and Its Responses to

Relative Sea Level and Nutrients in Estuaries at Plum Island, Massachusetts, and North Inlet,

South Carolina, USA. Oceanography **26**:78-84.

Morris, J. T., P. V. Sundareshwar, C. T. Nietch, B. Kjerfve, and D. R. Cahoon. 2002. Response of

coastal wetlands to rising sea level. Ecology **83**:2869-2877.

Mudd, S. M., A. D'Alpaos, and J. T. Morris. 2010. How does vegetation affect sedimentation on tidal

marshes? Investigating particle capture and hydrodynamic controls on biologically mediated

sedimentation. Journal of Geophysical Research: Earth Surface **115**.

O'Connor, M. T., and K. B. Moffett. 2015. Groundwater dynamics and surface water-groundwater

interactions in a prograding delta island, Louisiana, USA. JOURNAL OF HYDROLOGY

524:15-29.

Ortiz, A. C., S. Roy, and D. A. Edmonds. 2017. Land loss by pond expansion on the Mississippi River

Delta Plain. Geophysical Research Letters **44**:3635-3642.

Payne, A. R., D. M. Burdick, G. E. Moore, and C. Wigand. 2021. Short-Term Effects of Thin-Layer

Sand Placement on Salt Marsh Grasses: A Marsh Organ Field Experiment. Journal of Coastal

Research **37**:771-778.

Qi, M., J. MacGregor, and K. Gedan. 2021. Biogeomorphic patterns emerge with pond expansion in

deteriorating marshes affected by relative sea level rise. Limnology and Oceanography

66:1036-1049.

Raposa, K. B., R. L. J. Weber, M. C. Ekberg, and W. Ferguson. 2017. Vegetation Dynamics in Rhode

Island Salt Marshes During a Period of Accelerating Sea Level Rise and Extreme Sea Level

Events. Estuaries and Coasts **40**:640-650.

Schepers, L., M. Kirwan, G. Guntenspergen, and S. Temmerman. 2017. Spatio-temporal development

of vegetation die-off in a submerging coastal marsh. Limnology and Oceanography **62**:137-150.

Smith, S. M., M. Tyrrell, K. Medeiros, H. Bayley, S. Fox, M. Adams, C. Mejia, A. Dijkstra, S. Janson, and M. Tanis. 2016. Hypsometry of Cape Cod Salt Marshes (Massachusetts, U.S.A.) and Predictions of Marsh Vegetation Responses to Sea-Level Rise. Journal of Coastal Research **33**:537-547.

Snedden, G. A., K. Cretini, and B. Patton. 2015. Inundation and salinity impacts to above- and belowground productivity in Spartina patens and Spartina alterniflora in the Mississippi River deltaic plain: Implications for using river diversions as restoration tools. Ecological Engineering **81**:133-139.

Stark, J., T. Oyen, P. Meire, and S. Temmerman. 2015. Observations of tidal and storm surge attenuation in a large tidal marsh. Limnology and Oceanography **60**:1371-1381.

Taylor, L., D. Curson, G. M. Verutes, and C. Wilsey. 2020. Mapping sea level rise impacts to identify climate change adaptation opportunities in the Chesapeake and Delaware Bays, USA. Wetlands Ecology and Management **28**:527-541.

Turner, R. E., E. M. Swenson, C. S. Milan, J. M. Lee, and T. A. Oswald. 2004. Below-ground biomass in healthy and impaired salt marshes. Ecological Research **19**:29-35.

Ursino, N., S. Silvestri, and M. Marani. 2004. Subsurface flow and vegetation patterns in tidal environments. Water Resources Research **40**.

Van Putte, N., P. Meire, P. Seuntjens, I. Joris, G. Verreydt, L. Hambsch, and S. Temmerman. 2022. Solving hindered groundwater dynamics in restored tidal marshes by creek excavation and soil amendments: A model study. Ecological Engineering **178**.

Voss, C. M., R. R. Christian, and J. T. Morris. 2013. Marsh macrophyte responses to inundation anticipate impacts of sea-level rise and indicate ongoing drowning of North Carolina marshes. Marine Biology **160**:181-194.

Walker, C. W., T. R. Lester, and C. W. Nealen. 2019. Hydrologic study at Farm Creek Marsh, Dorchester County, Maryland, from April 2015 to April 2016 U.S. Geological Survey Scientific Investigations Report **2019-5032, 12p**.

Wang, C., L. Schepers, M. L. Kirwan, E. Belluco, A. D'Alpaos, Q. Wang, S. Yin, and S. Temmerman. 2021. Different coastal marsh sites reflect similar topographic conditions under which bare

patches and vegetation recovery occur. Earth Surf. Dynam. **9**:71-88.

Watson, E. B., K. B. Raposa, J. C. Carey, C. Wigand, and R. S. Warren. 2017. Anthropocene Survival

of Southern New England's Salt Marshes. Estuaries and Coasts **40**:617-625.

Xin, P., B. Gibbes, L. Li, Z. Song, and D. Lockington. 2010. Soil saturation index of salt marshes

subjected to spring-neap tides: a new variable for describing marsh soil aeration condition.

Hydrological Processes **24**:2564-2577.

Xin, P., J. Kong, L. Li, and D. A. Barry. 2013a. Modelling of groundwater-vegetation interactions in a

tidal marsh. Advances in Water Resources **57**:52-68.

Xin, P., L. Li, and D. A. Barry. 2013b. Tidal influence on soil conditions in an intertidal creek-marsh

system. Water Resources Research **49**:137-150.

Xin, P., A. Wilson, C. J. Shen, Z. M. Ge, K. B. Moffett, I. R. Santos, X. G. Chen, X. H. Xu, Y. Y. Y. Yau,

W. Moore, L. Li, and D. A. Barry. 2022. Surface Water and Groundwater Interactions in Salt

Marshes and Their Impact on Plant Ecology and Coastal Biogeochemistry. REVIEWS OF

GEOPHYSICS **60**.

Xin, P., T. Z. Zhou, C. H. Lu, C. J. Shen, C. M. Zhang, A. D'Alpaos, and L. Li. 2017. Combined effects

of tides, evaporation and rainfall on the soil conditions in an intertidal creek-marsh system.

Advances in Water Resources **103**:1-15.

Table 1 Data sources of synthesized growth response curves of typical macrophytes to inundation duration.

Species	Growth response indicators
S. americanus	live aboveground biomass (AGB), live belowground biomass (BGB), root to shoot ratio[a]
S. patens	live AGB, live BGB, root to shoot ratio[a]; AGB, BGB[b]
S. alterniflora	AGB , BGB [b]; Total AGB, Live AGB, Total shoot density, End-of-season BGB[c]
J. roemerianus	total AGB, Live AGB, total shoot density, end-of-season BGB[c]

[a]Kirwan and Guntenspergen (2015)- Blackwater and Transquaking River, Maryland.

[b]Snedden et al. (2015)- Middle Breton Sound and lower Breton Sound, Louisiana.

[c]Voss et al. (2013)- Pine Knoll Shores and Lola, North Carolina.

Figure legends

Fig. 1 a) Location of the two study sites within the Chesapeake Bay. Shallow groundwater wells were located in the vegetated, dieback, and pond zones at each site: b) Farm Creek Marsh (FCM) and c) Deal Island (DI).

Fig. 2 Groundwater table (solid lines) and surface elevation (dashed lines) of different zones (color coded) of deteriorating marshes at DI (a) and FCM (b) (Datum: NAVD 1988). At DI, 'tidal creek' means nearest tidal creek to interior deteriorating marshes of DI (Fig. 1c), and at FCM, it represents tidal height at Bishops Head station.

Fig. 3 Groundwater level at marsh interiors (vegetated zone, dieback zone, pond) was distributed within a narrow height range compared to tidal creeks at both marsh sites (a-DI; b-FCM). Similar to Fig. 2, tidal creek in a) means nearby tidal creek in DI, and tidal creek in b) means Bishops Head station.

Fig. 4 Comparison of soil saturation index (SSI) at soil depth of 0 m (a, $SSI_{0.0}$, equivalent to the hydroperiod), 0.1 m (b, $SSI_{0.1}$) and 0.2 m (c, $SSI_{0.2}$) between in situ observations (blue) and estimations (red) derived from elevation and tidal height. TC-tidal creek; PD-pond; DB-dieback zone, VG-vegetated zone. The response curve of hydroperiod and SSI to elevation in interior marsh was fitted with logistic

regression, and results of tidal creek in DI was marked separately with *.

Fig. 5 Comparison of the distribution of daily inundation duration between

observations and estimations from elevation and tidal height.

Fig. 6 Growth response curves of typical macrophytes to hydroperiod was generated

from empirical data (Voss et al. 2013, Kirwan and Guntenspergen 2015, Snedden et al.

2015). The difference in hydroperiod estimated using elevation and tidally-based

method (dashed lines) and in situ observations (solid lines) are also displayed for each

marsh zone. Colors indicate the vegetated zone (green), the dieback zone (orange),

and the pond (blue) for Deal Island (DI, left) and Farm Creek Marsh (FCM, right

column).

Fig. 7 There is a divergent response of hydroperiod (a) and soil saturation index at 0.1

m depth (b) to elevation change in tidal creeks (red) and interior marshes (blue) at

Deal Island. Asterisks indicate the observed result in interior marsh, these levels were

used to validate the simulated result (filled circles).

Fig. 8 Dieback of *S. patens* and *D. spicata* before being displaced by more flood

tolerant species, such as *S. americanus* (a) or *J. roemerianus* (b) in interior marsh of

Deal Island.

609   Fig. 1

612   Fig. 2

614
615   Fig. 3
616

618   Fig. 4

622   Fig. 5

625 Fig. 6

628   Fig. 7

630
631   Fig. 8
632