Cancer immunotherapy responses persist after lymph node resection

Hengbo Zhou[1,2], James W. Baish[3], Meghan J. O'Melia[1], Laurel B. Darragh[4], Emma Specht[5], Juliane Czapla[5], Pin-ji Lei[1], Lutz Menzel[1], Johanna J. Rajotte[1], Mohammad R. Nikmaneshi[1], Mohammad S. Razavi[1], Matthew G. Vander Heiden[2,6], Jessalyn M. Ubellacker[7], Lance L. Munn[1], Genevieve M. Boland[5], Sonia Cohen[5], Sana D. Karam[4] and Timothy P. Padera[1]

1. Edwin Steele Laboratories, Department of Radiation Oncology, Massachusetts General Hospital

2. Koch Institute for Cancer Research and Department of Biology, Massachusetts Institute of Technology

3. Department of Biomedical Engineering, Bucknell University

4. Department of Radiation Oncology, University of Colorado Anschutz Medical Campus

5. Department of Surgery, Massachusetts General Hospital

6. Department of Medical Oncology, Dana-Farber Cancer Institute

7. Department of Molecular Metabolism, Harvard T.H. Chan School of Public Health

Abstract:

Due to the critical role of lymph nodes (LNs) in the initiation and maintenance of adaptive immune responses, it is unclear whether surgical removal or ablative radiation therapy of LNs should be performed in patients with metastatic LNs that will receive immunotherapy. Surgical removal of LNs to prevent metastatic recurrence, including sentinel lymph node biopsy (SLNB) and completion lymph node dissection (CLND), are performed in routine practice. However, removing LNs eliminates opportunities for generating anti-cancer immune responses that are enhanced with immune checkpoint blockade (ICB). Balancing the potential risks and benefits of LN surgery is necessary to maximize outcomes for patients. In contrast to mouse studies using ectopic tumor implantation[1,2], the phase III clinical trial NCT00636168 found patients with completely resected stage III melanoma (primary tumor, sentinel LNs and disease-related LNs all removed) still benefited from anti-CTLA4 inhibition. Our retrospective analysis demonstrated that stage III melanoma patients with SLNB or CLND have similar response to anti-PD1 inhibition. Using orthotopic murine mammary carcinoma and melanoma that have spontaneous LN metastases, we show that responses to ICB persist in mice after resection of TDLNs. Mechanistically, after disruption of normal lymphatic drainage, antigen can access adjacent lymphatic basins (lymphosomes) and be transported to distant LNs, which extends the responsiveness to ICB. By evaluating head and neck cancer patients treated by neoadjuvant durvalumab and irradiation, we show that distant LNs remain reactive in the absence of metastatic disease in ICB responder patients after tumor and disease related-LN resection, indicating anti-cancer immune reactions occur in distant LNs after TDLN removal in patients. Additionally, after TDLN dissection in mouse models, ICB delivered to distant LNs generated greater survival benefit, compared to systemic administration. In complete responders, anti-tumor immune memory induced by ICB was systemic rather than confined within lymphoid organs. Based on these findings, we constructed a computational model to predict free antigen trafficking in patients that will undergo LN dissection. Collectively, we propose that ICB efficacy persists in tumors of the trunk and extremities after removing original TDLNs because antigen can be transported through interstitial tissues to adjacent lymphosomes to gain access to alternative LNs, where anti-cancer immune responses can be generated and maintained.

Introduction:

Recent advances in understanding of the mechanisms of T cell activation and exhaustion led to the identification of several immune checkpoints, such as programmed death 1 (PD-1)[3,4] and cytotoxic T lymphocytes antigen 4 (CTLA-4)[5,6], which have been used as therapeutic targets[7,8]. Immune checkpoint blockade (ICB) has been approved to treat many cancers including

melanoma[9,10], non-small cell lung cancer[11,12], renal cell carcinoma[13], Hodgkin lymphoma[14],
hepatocellular carcinoma[15,16], and gastric carcinoma[17]. However, an improved understanding of
ICB mechanisms *in vivo* is necessary for immunotherapy to benefit more patients.

Lymph nodes (LNs) are secondary lymphoid organs that surveil tissue fluids and initiate
adaptive immunity[18]. Due to the direct drainage of interstitial fluid from tumor tissue in the
lymphatic basin (lymphosome), metastatic dissemination to tumor-draining lymph nodes
(TDLNs) occurs in the early stage of cancer growth[19]. Colonization in LNs can block nodal entry
of naïve lymphocytes[20], expand regulatory T cells (Tregs)[21,22] and induce systemic tolerance to
favor distant metastasis of cancer cells[22]. LN metastases can also further disseminate to
additional metastatic sites[23,24]. Therefore, LN metastatic burden is a strong prognostic factor for
patient outcome. Sentinel lymph node biopsy (SLNB) and completion lymph node dissection
(CLND) are commonly performed[25-29] to help stage disease and reduce disease burden. However,
in an era of effective systemic therapies, the morbidity of these surgical LN procedures and the
controversy regarding their role in preventing recurrence/metastatic disease in certain cancers,
including early-stage breast cancer[30], ovarian cancer[31], cervical cancer[32], and melanoma[33], has
shifted clinical practice toward less aggressive LN surgery for the initial management of
microscopic metastatic disease.

Recent work has demonstrated that tumor-specific T cells and non-exhausted tumor-specific
memory cells are enriched in TDLNs, but not in non-TDLNs[34,35]. In addition, PD-1/PD-L1
interaction between conventional dendritic cells (DCs) (PD-L1+) and tumor-specific T cells (PD-
1+) is abundant in TDLNs[34]. Intrapleural delivery of low-dose PD-L1 antibody to TDLNs can
induce anti-tumor immunity against AC29 tumors (mesothelioma)[34]. Additional studies, using
B16F10 (melanoma), 4T1 (mammary carcinoma) and E0771(mammary carcinoma) tumors,
found selective administration of ICB to TDLNs can improve ICB potency, compared to
systemic administration[36,37]. These data suggest that removal of TDLNs could potentially impair
ICB outcome. Indeed, anti-PD-1 or anti-4-1BB efficacy can be abolished by TDLN dissection in
MC38 (colon adenocarcinoma) or CT26 (colon carcinoma) tumors ectopically grafted in the
skin[1,2]. Similarly, responses can be ablated by TDLN lymphadenectomy at any point during anti-
PD-L1 treatment in subcutaneously grafted B16F10 (melanoma) tumors expressing lymphocytic
choriomeningitis virus glycoprotein[35]. However, in the phase III clinical trial NCT00636168,
completely resected stage III melanoma patients, from whom sentinel LNs and metastatic LNs
were removed, still responded and benefited from ipilimumab (targeting CTLA-4), indicating
that persistent presence of TDLNs may not be required for ICB efficacy[38]. Further, in the phase I
trial NCT04274816, lymphatic basin administration of tremelimumab (targeting CTLA4) after
tumor resection led to a systemic reduction of Tregs and activation of effector T cells. This
response persisted up to 3 months after subsequent SLNB, suggesting that the ICB-mediated
immune response was not confined only within sentinel LNs[39]. Due to the important role of LNs
in the adaptive immune response, it is imperative to evaluate the effects of SLNB and CLND on
ICB efficacy.

Here, through retrospective analysis, we found that regardless of the extent of LN surgery, stage
III melanoma patients benefited similarly from adjuvant anti-PD1 inhibition. By using orthotopic
breast cancer and melanoma models in mice, we demonstrated that neither SLNB nor CLND
eliminates ICB efficacy, despite the purported important role of TDLNs in anti-tumor responses.
Mechanistically, after TDLN dissection, antigen can be transported to distant LNs through
redirected lymph drainage, which contributes to the residual responsiveness to ICB. This
phenomenon is corroborated in head and neck cancer patients—treated by neoadjuvant
durvalumab and irradiation therapy followed by adjuvant durvalumab—by the presence of
reactive distant LNs after primary tumor resection and LN dissection. Thus, consistent with
clinical observations, ICB efficacy is not reliant solely on the presence of the original TDLNs,
and ICB can be effective against locoregional recurrence even after TDLN dissection.

Results:

The extent of lymph node surgery does not impact melanoma response to ICB clinically

Prior to the availability of effective systemic therapies for patients with melanoma, LN surgery played a major role in the treatment of patients with locoregional disease. Any patient with clinically evident LN metastases or a positive SLNB underwent CLND for disease control. In 2017 MSLT-II[33], a randomized trial of CLND vs observation in patients with positive SLNB, demonstrated that CLND provided no melanoma-specific survival (MSS) benefit to patients with melanoma identified within the SLN. Given the morbidity of CLND, this trial changed clinical practice to favor observation after a positive SLNB, with CLND reserved for clinical disease recurrence in the draining LN basin. Approximately 20% of patients would ultimately recur in the draining nodal basin, sparing 80% of patients with a positive SLNB from the unnecessary morbidity of CLND[33].

The MSLT-II trial reflected the era of melanoma treatment prior to the introduction of ICB. With the approval of effective systemic therapies, the role of LN surgery—other than for staging—has been called into question. Given the importance of TDLNs for ICB response in mouse models[1,2,35], we asked whether the extent of LN surgery would impact the effectiveness of ICB treatment. To assess this, we analyzed a cohort of 66 patients who underwent surgery for stage III melanoma followed by adjuvant treatment with at least one line of anti-PD1 therapy (nivolumab or pembrolizumab) at Massachusetts General Hospital. Within this adjuvant ICB cohort, 50 patients underwent SLNB alone while 16 patients underwent CLND. There was no statistically significant difference in MSS (**Figure 1 A**) or recurrence-free survival (RFS) (**Figure 1 B**) between patients who underwent SLNB alone versus those who underwent CLND. After further stratification, no difference was detected in local (**Figure 1 C**) or distant (**Figure 1 D**) RFS either. These data demonstrate that patients who have undergone LN surgery still benefit from ICB treatment, and that the extent of LN surgery within the draining lymph node basin does not impact outcomes in the setting of ICB.

ICB response persists after TDLN dissection in mice

To experimentally test the impact of TDLN dissection on ICB efficacy in murine models, we used orthotopic, syngeneic mouse melanoma (B16F10) and mammary carcinoma (E0771) models. SLNB was modeled by removing the inguinal (IN) LN in C57BL6 mice, which drains the upper hindleg and 4th mammary fat pad of mice. Of note, as surgery can cause multiple effects in cancer patients, including the potential for causing metastatic spread after primary tumor removal and trauma-induced immunosuppression[40,41], sham LN removal surgery was performed on all animals in control groups. After 10 days of recovery, B16F10 or E0771 cells were introduced to the dermis on the lower flank or 4th mammary fat pad, respectively. 7 days later, 3 doses of PD-1 blocking antibody (10mg/kg) were administered intraperitoneally (every 2 days for B16F10 tumors or every 3 days for E0771 tumors). Compared to IgG control treated mice, PD-1 inhibition increased overall survival in both the sham and SLN removed (SLNB) mice. Interestingly, in groups treated with anti-PD-1, there was no difference in overall survival between the sham surgery mice and the SLNB counterparts (**Figure 2A**). To mimic CLND, a radical procedure that is increasingly less common in the clinic[30-33], ipsilateral IN, axillary (AX), and brachial (BR) LNs were surgically removed, as each is in the region can receive lymphatic drainage from the tumor site[2,42]. Similar to SLNB, there was residual ICB response after CLND, and no statistically significant difference in overall survival was found between sham and CLND conditions when animals were treated with PD-1 blocking antibody (**Figure 2B**).

Next, we measured whether anti-tumor immunity is retained solely in TDLNs after tumor establishment. Thus, we tested the effects of SLNB and CLND after primary tumor implantation. B16F10 or E0771 cells were orthotopically implanted 6 days prior to surgery. Sham, SLNB or CLND was performed at day 6 and 3 doses of anti-PD-1 treatment was initiated at day 7 (every 2 days for B16F10 tumors or every 3 days for E0771 tumors). Similar to pre-emptive surgery, neither SLNB nor CLND significantly affected overall survival in anti-PD-1 treated mice compared to sham surgery (**Figure 2C**). Although there was no significant decrease in survival

after LN surgery, fewer mice remained tumor-free in the LN dissection conditions (**Figure 2D**).
These findings are not consistent with recent investigations using MC38 and CT26 tumor models
grown ectopically in the subcutaneous setting[1,2]. To determine whether implantation site can
affect ICB outcome, E0771 cells were introduced to the dermis and LN dissection was conducted
either pre-emptively or after tumor establishment. Intriguingly, aggressive CLND can impair
anti-PD-1 efficacy in the ectopic setting (**Supp Figure 1 A&B**), indicating altered immune
response can occur when tumor cells are introduced to an ectopic site.

Other LNs are required for ICB response after TDLN dissection

Because of the surprising finding that neither SLNB nor CLND affects the overall survival
benefit of ICB treatment in mice, we sought to examine whether LNs in general are essential for
ICB response. To that end, lymphotoxin alpha (Lta) knock out (KO) mice —Lta^{tm1Dch} (C57bl/6
background), referred to throughout as *Lta* KO mice were used. These mice lack Peyer's patches
and multiple LNs including popliteal (PO), IN, BR and AX LNs, among other missing lymphoid
tissues[43]. Of note, the bone marrow is functional in these animals and the lack of LNs is likely
caused by development defects[43-45]. Although the splenic structure is disorganized by *Lta*
deletion, the number of lymphocytes, the composition of the types of lymphocytes and their
homing and effector abilities are normal in both the spleen and in peripheral blood[43-46]. B16F10
or E0771 cells were orthotopically implanted in *Lta KO* mice and 3 doses of PD-1 blocking
antibody were administered starting 7 days after tumor implantation (every 3 days). Strikingly,
compared to responses in wildtype (WT) mice (sham animal survival curves from SLNB study,
Figure 2A, were plotted for comparison), slowed tumor growth and improved survival from PD-
1 inhibition were abolished in these mice lacking LNs (**Figure 2E**), suggesting adaptive immune
responses generated in LNs contribute to ICB response. Further, these results suggest that LN-
dependent compensatory mechanisms occur after TDLN dissection to maintain anti-PD-1
responses in wild-type mice.

**LN dissection alters lymph drainage and promotes antigen access to distant lymphoid
organs**

To interrogate the mechanism by which animals that have undergone LN surgery generate ICB-
induced immune responses, we first measured the local transport of surrogate antigens to
alternative draining lymph nodes (DLNs) after SLNB or CLND. 2 million Dalton (MD) FITC-
dextran and 4.4 kilodalton (kD) TRITC-dextran were used as surrogate biomolecules with
different sizes. The dextran mixture was introduced to the 4th mammary fat pad in animals that
had undergone sham surgery or dissection of the IN LN (SLNB) or the IN/AX/BR LNs (CLND).
After 6 hours, ipsilateral IN, AX, BR LNs remaining in the relevant groups as well as all
contralateral IN LNs were collected for microscopic examination. In LNs, the majority of 2 MD
dextran was found to be aggregated in the subcapsular sinus (SCS) region—showing transport by
the lymphatic system—while 4.4kD dextran was sparsely distributed throughout the LN (**Figure
3A**). Based on these data, it is not clear whether the 4.4kD dextran arrived in the LN from the
lymph or the blood. Small biomolecules can enter LNs via the lymph or blood circulation[47].
However, our data show that the large biomolecules rely heavily on the lymphatic system to
arrive in the LN. Both types of dextran can be taken up by phagocytes resident in cortical,
paracortical and inter-follicular spaces (**Figure 3A**). 2 MD dextran abundance in AX LNs from
SLNB animals is comparable to counterparts in IN or AX LNs from sham animals (**Figure 3B**),
indicating resection of IN LN does not disrupt the drainage to the downstream AX LN.
Interestingly, 2 MD dextran was also detected entering the SCS of contralateral IN LNs in
certain animals (**Figure 3C**). Compared to sham animals, there was an increased incidence of 2
MD dextran drained to contralateral IN LNs of CLND animals (**Figure 3B**). Inconsistent with a
previous report[2], there was limited drainage from mammary fat pad to BR LNs across conditions,
suggesting BR LNs are not main DLNs (**Figure 3B**). Nonetheless, because BR LNs are in axilla
region, removing BR LN along with IN LN and AX LN is a relevant approach to model CLND
(basin dissection) which also removes non-DLNs[30,33]. Collectively, these data show that CLND

disrupts the normal patterns of lymph drainage, which results in lymph (and surrogate antigen)
being transported to alternative LNs.

Next, we determined whether interstitial transport from a disrupted lymphosome could cross the
body midline and enter adjacent lymphosomes. To that end, an incision was made to the midline
skin (from lower abdomen to chest) in CLND mice, and 2MD dextran was injected to the
ipsilateral 4th mammary fat pad. Strikingly, 2MD dextran transport to contralateral LNs was only
detected in mice with intact midline skin, but not in mice with midline injury (**Figure 3D**). Thus,
these data show that interstitial fluid with solutes can move across the midline through the
dermis and reach an adjacent lymphosome draining to distant LNs after CLND. These results
suggest that when primary LN(s) are removed, lymph can be driven to the next accessible
lymphosome, even if the DLNs are anatomically distant (**Figure 3E**).

To further test whether—beyond the mammary fat pad region—antigen access to distant LNs
can be enhanced when local lymphatic drainage is ablated, we performed sham surgery or
ipsilateral popliteal (PO), IN, AX and BR LN dissection on mice (ILND condition).
Subsequently, a mixture of 2MD and 4.4kD dextran was injected to the foot pad of those
animals. Due to this anatomic site, injection-induced antigen convection to distant sites was
minimized, and all LNs along the direct drainage path were absent. Similar to findings described
in **Figure 3B**, we discovered that 6 hours after injection, 2MD dextran was mainly located in
ipsilateral PO, IN and AX LNs when lymphatic drainage was intact, whereas ILND re-routed
2MD dextran transport to contralateral IN and AX LNs (**Supp Figure 2**). The lack of 2MD
dextran in contralateral PO LNs suggests that, once taken up by lymphatic vessels of an adjacent
lymphosome, the lymph still follows the normal drainage patterns, and does not undergo
retrograde flow or circulate significantly through the blood in ILND mice (**Supp Figure 2**). In
contrast, 4.4kD dextran, with its smaller size was found in all remaining LNs (**Supp Figure 2**),
suggesting transport through the blood. The compensatory drainage phenomenon in the lower
limb suggested by our data has also been reported clinically. A recent indocyanine green
fluorescence lymphography study found that in patients with cancer treatment related lower limb
lymphoedema, lower limb interstitial fluid can be transported across the lower abdomen to the
contralateral inguinal region[48].

Collectively, our data show that lymph is drained to downstream LNs after SLNB, while CLND
causes disruption of lymphatic vessel transport, which leads to a reliance on interstitial transport
to carry the surrogate antigens to adjacent lymphosomes (drainage basins), where they are taken
up by local lymphatics and transported to distant LNs. These findings are consistent with the
clinical observation that breast cancer patients who have undergone AX LN dissections are more
likely to develop altered lymph drainage to contralateral AX LNs[49]. Similarly, radiation therapy
can cause lymphatic drainage aberration in breast cancer patients, which decreases flow to the
ipsilateral axilla while increasing flow to contralateral axilla[50].

In addition to distant LNs, we found that 2MD dextran was also detectable in spleens from both
sham and ILND mice. 2MD dextran was more abundant in spleens of mice in the ILND
condition (**Figure 3F**). These large dextran molecules were mainly detected in the B cell follicles
and adjacent marginal zone macrophages, instead of residing in macrophage-rich red pulp[51]
(**Figure 3G**), indicating the potential for the generation of immune responses in the spleen
against antigen arising from distal tissue, compensating for the loss of DLNs.

TDLN dissection stimulates immune responses in distant LNs

Since distant LNs of CLND mice can be exposed to interstitial fluid from the primary tumor site,
we next examined whether tumor-specific immunity can be generated in these distant LNs.
B16F10 cells expressing ovalbumin (257-264: SIINFEKL) were implanted to sham surgery or
CLND mice and anti-PD-1 treatment started after 7 days. Subsequently, the spleen as well as any
ipsi- and contra-lateral LNs were collected for immunofluorescence and flow cytometry
analyses. CD8 T cell expansion was detected in ipsilateral LNs in sham surgery mice and

contralateral LNs in CLND mice (**Figure 4A**). Moreover, spleens in CLND animals contained an
equal percentage of CD8 T cells compared to the sham surgery counterparts (**Figure 4B**),
indicating that the immune reaction to the tumor was not confined within TDLNs. We also
measured the abundance of SIINFEKL-specific CD8 T cells (referred to throughout as OVA-T
cells) to assess tumor-specific immune responses (**Figure 4C**). Consistent with previous
investigations[34,36], OVA-T cells were found to be predominantly enriched in sham surgery
ipsilateral LNs, whereas certain CLND contralateral LNs, but not cervical LNs, also harbored
high numbers of OVA-T cells (**Figure 4D**), indicating a directional spread of antigen consistent
with adjacent lymphosomes, instead of systemic spread . Additionally, the abundance of OVA-T
cells was comparable in spleens and blood of both sham and CLND mice (**Figure 4D**),
demonstrating that the absence of TDLNs does not ablate circulating tumor-specific T cells.

To better recapitulate the clinical situation, ICB-induced tumor-specific T cell response was
tested in an adjuvant setting. Two weeks after tumor implantation in intact mice, primary tumor
removal was performed with or without CLND. After 3 days of recovery, mice were given anti-
PD-1 treatment. No difference was found in OVA-T cell abundance between sham ipsilateral
LNs and CLND contralateral LNs (**Figure 4E**), indicating, that even without the primary tumor
antigen source, distant LNs can generate tumor-specific immune response after the removal of
the initial TDLNs. To test whether tumor-specific T cells from distant LNs can infiltrate
recurrent tumors, B16F10-OVA tumors were implanted into Kaede mice[52], and the tumors were
resected with and without CLND; the mice were then treated with adjuvant anti-PD1 inhibition.
After 14 days, these mice were implanted with B16F10-OVA tumors in the same primary site—
to model recurrence—followed by photoconversion of contralateral IN LNs. Strikingly, Kaede
red+ OVA-T cells were present in recurrent tumors, and the abundance was greater in CLND
mice, compared to mice that did not undergo LN surgery (**Figure 4F**), suggesting that immune
cells from distant LNs can be activated and traffic to recurrent tumors.

Antigen can disseminate by advection with interstitial fluid, but it can also be carried by
antigen-presenting cells (APCs), which may access distant LNs after primary DLN resection. To
assess the contribution of APCs, sham surgery or dissection of PO, IN, AX and BR LNs (ILND
condition) was conducted in cohorts of mice. After recovery from surgery, bone marrow-derived
DCs (BMDCs) from donor mice were loaded with OVA, stimulated with lipopolysaccharide
(LPS) and subsequently injected in the foot pad of sham or ILND mice. Even though it has been
demonstrated that BMDCs contain heterogeneous macrophage and DC populations[53], this model
is able to test APC-mediated T cell priming in LNs. Intriguingly, we found that OVA-T cells
were significantly enriched in ipsilateral LNs of sham mice and contralateral LNs of ILND mice,
but not contralateral LNs of sham animals (**Figure 4G**). These results suggest that antigen
carrying-APCs also have greater access to distant LNs when lymphatic drainage is disrupted.

As our data show that large dextran molecules are enriched in the spleen after CLND (**Figure 3
E&F**) and accompanied by CD8 T cell expansion (**Figure 4B**) and a tumor-specific T cell
response (**Figure 4D**), we next determined whether the spleen is essential for residual ICB
response in mice after CLND. To that end, sham surgery (SP+ condition) or splenectomy (SP-
condition) was performed 2 weeks prior to CLND, and E0771 tumors were implanted and treated
as described before. Unexpectedly, splenectomy did not significantly affect the overall survival
in anti-PD-1 treated mice. Complete responders were less frequent, but still present, in SP- mice
(**Supp Figure 3A**). Further, all complete responders (5/5) from the SP+ condition and 2 out of 3
of SP- counterparts could completely reject tumor rechallenge, (**Supp Figure 3B**). Altogether,
these data suggest that the spleen contributes, but only marginally, to the ICB response and long-
term immune memory formation in CLND mice.

Mathematical modeling of antigen access to adjacent lymphosomes via dermal interstitial
transport after TDLN dissection

Because our data show that antigen can access adjacent lymphosomes and activate distant LNs
(**Figure 3&4**), we next used mathematical modeling to help understand this process from a

pharmacokinetic perspective. To that end, we developed a multi-scale computational model of
transport of free antigen and imaging tracers **(Figure 5A&B)** that recapitulates our primary
experimental findings in mice and predicts how these results may translate to clinical practice in
humans (details of the model provided in **Supplemental Materials**).

When all LNs and lymphosomes are functioning normally, interstitial fluid is absorbed by nearby
initial lymphatic capillaries that drain locally into collecting lymphatic vessels that pump the
lymph to the DLN and eventually return it to the blood circulation. Normally, a local equilibrium
exists in which interstitial fluid pressure remains low—near or just below atmospheric
pressure—and little lateral movement of fluid occurs in the superficial tissue layers. In contrast,
Figure 5C-F shows how disabling various combinations of the IN, AX, BR, and PO nodes in the
mouse diverts flow to adjacent lymphosomes due to the lack of local pressure minima provided
by lymphatic drainage. In general, disabled collecting lymphatic vessels lead to the accumulation
of interstitial fluid and then its redirection to the next adjacent lymphosome – even across the
midline. This transport can occur through the tissue interstitial space or through the non-valved,
highly interconnected lymphatic capillary network. However, the DLN(s) for the adjacent
lymphosome might be relatively distal, helping to explain our in vivo data in models of CLND.
This computational model predicts that antigen can activate immune responses in specific distal
LNs in the body after LN resections.

Our results in a human model (**Figure 5G-I**) were qualitatively similar to those in mice, but
important differences arise because of the relative size of the lymphosomes, anatomical
connectivity of the vessels, and the degree of redundancy in the lymphatic networks. The
pressures in the skin obtained from the full, 3-D human model were compared to an analytical
result for a planar, circular lymphosome with the same surface area (**Figure 5J**) in **Figure 5K**.
Close agreement is seen despite the planar model not accounting for the spatial curvature of the
human chest. Dimensional analysis of the governing equations (see **Supplemental Materials**)
revealed that the lateral distance over which interstitial fluid must flow in the skin before being
absorbed by functional lymphatics increases with the fluid mobility in the initial, valveless
lymphatic vessels (tissue permeability κ or hydraulic conductivity), but decreases with the fluid
conductance of the blood vessel and lymph vessel walls.

To examine the size dependence of the lymphosome, we performed a parametric study of the
simplified model **(Figure 5L)**. Here, we found that larger lymphosomes, such as those present in
humans, generate more total fluid that must be drained and result in higher interstitial pressures
than expected in mice. This size difference alone may account for much of the relative difficulty
in creating experimental models of lymphedema in laboratory mice. An additional result of the
simplified model **(Figure 5M)** was that the mobility of the fluid (κ) in the network of initial
lymphatic vessels strongly affects the interstitial fluid pressure, but has little influence on the
total quantity of antigen-carrying fluid diverted elsewhere or its destination. Increasing κ by a
factor of 10,000 resulted in an increase in the velocity of interstitial flow at the perimeter of the
disabled lymphosome by only a factor of about 15 because changes in pressure gradient were
offset by changes in κ. For a wide range of parameter values, fluid was diverted to the same
nearest neighboring drainage basins.

However, anatomical differences between mouse and human alter the path of flow in the
collecting vessels. For example, the direct connection from the inguinal region to the axillary
region that exists in the mouse is absent in human. Disabling an AX LN in a human did not
affect drainage to the inguinal basin, but like the mouse diverted fluid to contralateral LNs
(Figure 5I). Moreover, disabling a single AX LN in a human did not produce the same extensive
re-routing of flow seen in the mouse because the human has numerous nodes in the axillary
region that form a redundant network. Note that anatomical variations in the network can exist
between individuals, and this will affect the flow redundancy.

Administration of ICB to distant LNs enhances efficacy after TDLN dissection

Previous preclinical investigations demonstrated that directly delivering a low dose of ICB
agents to TDLNs can achieve superior outcome compared to systemic delivery[36,37]. After tumor
resection, local intradermal injection of low-dose anti-CTLA4 antibody into the lymphatic
drainage basin to the SLNs can activate CD8 effector T cells while decreasing Treg frequency
prior to SLNB in patients[39]. These findings highlight the potential of LNs to be used as an ICB
target site. Because distant LNs are more likely to be exposed to tumor antigen after LN
dissection (**Figure 3 & Supp Figure 2**), we assessed the importance of contralateral LNs for
low-dose ICB treatment in mice that have undergone CLND. Since anti-CTLA4 and anti-PD-1
combination has shown efficacy in clinical settings[54], we delivered this ICB combination at low
dose (5mg/kg) by intradermal administration in a lymphatic basin that is drained by ipsilateral
LNs (tumor draining IN+AX) in sham LN surgery mice or contralateral LNs in CLND mice.
Compared to isotype IgG, we found ICB drastically increases the volume of ipsilateral LNs in
sham LN surgery mice and contralateral LNs in CLND mice (**Figure 6A**). Additionally, these
LNs share similar T cell composition (**Figure 6B**) and possess comparable abundance of stem-
like effector memory T cells (CD44+, PD-1+, TCF1+, CD39-, CD62L-) which are key
contributors to ICB response[55] (**Figure 6C**). Altogether, similar to TDLNs, distant LNs can be
activated by ICB in mice after CLND.

To directly test the potency of distant LNs during ICB treatment, we treated sham LN surgery
and CLND mice bearing E0771 tumors with anti-CTLA4 and anti-PD-1 antibodies (5mg/kg) by
intraperitoneal (IP) administration or intradermal administration to contralateral LNs (C.LNs).
Strikingly, compared to systemic delivery, combination ICB given to C.LNs provides greater
survival benefit and response rate in CLND mice, but not in sham LN surgery counterparts
(**Figure 6D**). These results provocatively suggest that delivering ICB therapeutics to downstream
or distant LNs may benefit patients that have undergone LN surgery. Notably, CLND decreased
ICB efficacy in the systemic anti-CTLA4/PD-1 treatment setting, which may be attributed to the
robust response rate (~75%) in this experiment, which is not normally observed clinically[38,56-59].
In trial NCT01844505, compared to single agent treatment, the combination of nivolumab and
ipilimumab generated a superior response in patients with untreated advanced melanoma[54].
However, in trial NCT03068455, compared to nivolumab alone, this combination given in the
adjuvant setting did not improve outcome in patients with completely resected melanomas[60].
Thus, primary tumor and/or TDLNs may be required for CTLA4 inhibition to provide additional
survival benefit when combined with PD-1 inhibition.

Compared to adjuvant-only regimens, neoadjuvant-adjuvant ICB can provide better outcomes for
patients with advanced melanoma[61]. To examine whether LN surgery compromises neoadjuvant-
adjuvant ICB efficacy, B16F10 tumors expressing Gaussia luciferase (B16F10-Gluc) were
treated with ICB delivered to the TDLN, and curative primary tumor resection with or without
LN surgery (SLNB or CLND) was performed (**Figure 6E**). Mice were monitored for tumor
recurrence and metastases (Gluc activity) for 100 days. After 100 days tumor-free mice were
rechallenged with B16F10-Gluc cells, and mice that developed tumors were treated with
adjuvant ICB delivered to the TDLN (IN LN in no LN surgery condition, AX LN in SLNB
condition and contralateral IN LN in CLND condition) (**Figure 6E**). Interestingly, after surgery,
ICB, but not IgG, induced vitiligo, especially, at the surgery sites of both primary tumor
resection and LN surgery (**Figure 6F**). The appearance of vitiligo indicates that the ICB-induced
immune response to antigens involved in melanin production is not abolished by removal of
TDLNs and can disseminate beyond the primary tumor location[62]. After neoadjuvant treatment,
only one mouse in the No LN surgery + ICB condition had a spontaneous recurrent tumor and
one mouse in the SLNB + ICB condition died due to a cause unrelated to cancer (**Figure 6G**),
demonstrating the potency of curative surgery. Upon rechallenge, compared to naïve mice, all
surgically treated mice exhibited immune memory against the original tumors, while ICB-treated
mice possessed greater immune memory compared to IgG-treated counterparts (**Figure 6H**). At
this sample size (~15 mice), neither SLNB nor CLND significantly diminished the ICB-induced
immune memory or survival benefit (**Figure 6H&I**). Surprisingly, these results suggest that, in a

TDLN targeted neoadjuvant-adjuvant ICB treatment regimen, LN surgery along with curative primary tumor resection does not ablate the formation of immune memory against cancer cells and improvement of overall survival.

LN-egressing immune cells are expendable for long-term anti-tumor immune memory

Complete responders emerged after ICB treatment in sham LN surgery and TDLN resected mice (**Figure 2&6**). To evaluate the long-term immune memory, we re-introduced E0771 cells to the original tumor sites of anti-PD1 treatment complete responders (**Figure 2**). Survivors in both sham LN surgery and TDLNs removed conditions rejected tumor rechallenge (**Supp Figure 4A**). To further test whether this immune memory is systemic, E0771 cells were introduced to both ipsi- and contra-lateral sides of animals. Similarly, all survivors, regardless of the presence of TDLNs, remained tumor-free (**Supp Figure 4B**). Finally, we determined that this immune memory is tumor specific, as E0771-immune mice cannot reject B16F10 tumors implanted in either mammary fat pad or dermis (**Supp Figure 4C**).

Lymphocytes use sphingosine 1-phosphate (S1P) signaling to exit lymphoid organs including bone marrow, LNs and Peyer's patches[63-65]. We aimed to test whether central memory T cells from LNs enter the circulation and greatly contribute to long-term immune memory in tumor-free survivors. To that end, tumor naïve control mice or complete responders generated by ICB combo treatment (**Figure 6**) were fed with water with or without FTY720 (2.5μg/ml), a potent inhibitor against S1PR[66]. Subsequently, E0771 tumors were implanted to both the primary tumor location and the contralateral site for re-challenge. FTY720 treatment drastically reduced CD4, CD8 and central memory (CD44+/CD62L+) T cells in the circulation ~90% (**Supp Figure 4D)**, suggesting this treatment can block T cells egress from lymphoid organs including LNs. Intriguingly, the vast majority of complete responders remained tumor-free, except two mice from the conditions-one from sham LN surgery and one from CLND-treated with FTY720 (**Supp Figure 4E**). Like previous observations (**Supp Figure 4 A&B**), LN surgery status did not affect the immune memory (**Supp Figure 4E**). Strikingly, this anti-tumor immune memory did not only reside within primary tumor tissues, as subsequent E0771 tumors implanted ectopically in the dermis of these tumor-free survivors from the orthotopic rechallenge study were also rejected (**Supp Figure 4F)**. These findings suggest that ICB-induced anti-tumor immune memory cells spread systemically in complete responders, and the contribution of immune cells egressing from lymphoid organs is dispensable after memory responses have formed.

Distant LNs are reactive after resection of tumor and disease-related LNs in head and neck cancer patients treated by ICB

To test whether, after LN dissection, distant LNs contribute to ICB efficacy on suppressing cancer recurrence in patients, we investigated head and neck cancer patients treated by ICB that tend to have a high local recurrence rate[67]. We examined a recent phase I/Ib trial (NCT03635164) in which HPV negative head and neck cancer patients that received neoadjuvant treatment to manage gross disease (primary and grossly involved nodes) consisting of combination hypofractionated radiotherapy and durvalumab (targeting PD-L1). Following a recovery time ranging from 5-6 weeks in the extension cohort, patients underwent surgical resection of the primary tumor and bilateral neck dissection per standard of care. Those with major pathological response or complete pathological response to neoadjuvant radiotherapy and durvalumab were classified as responders (response>95%). Under permission of multi-disciplinary tumor board, with the exception of four cycles of durvalumab, all other adjuvant therapy in the form of radiation or chemotherapy was omitted in responders[68]. To date, no recurrent disease was detected in these responders during post-operative follow-up visits (>2 years). We compared CT and PET images at diagnosis to post-surgery follow-up examinations in the responder cohort. Strikingly, after removal of the tumor and initial TDLNs, enlargement of distant LNs was found in most responders, but not in non-responders (**Figure 7A**). Of note, no nodal disease was detected by radiology-guided biopsy, indicating these enlarged LNs were activated and immune-reactive. The reactive LNs were higher echelon LNs in areas that were

now draining the original tumor site. For instance, in patients 01-013 and 01-015, lower neck
LNs (posterior cervical) and upper neck LNs (submandibular and submental) were enlarged,
respectively, at follow-up, which was not observed at diagnosis (**Figure 7B**). Similarly, in
patient 01-009, neck draining mediastinal LNs above the aortic arch were inflamed after tumor
and LN resection (**Figure 7C**). These data suggest that surgical removal of the primary tumor
and TDLNs can result in reactive distant LNs and extend ICB response in patients.

Discussion:

TDLNs are exposed to high antigen burden, allowing them to readily generate adaptive immune
response against cancer cells[19]. Recent reports using mouse tumor models advocate that TDLNs
are indispensable for ICB response[1,2,35,69], likely due to unique tumor-specific T cell memory
populations[35]. A phase II trial (NCT03698019) of patients with resectable stage III or stage IV
melanoma, neoadjuvant-adjuvant pembrolizumab treatment provided improved event-free
survival compared to adjuvant-only setting[61], indicating the presence of the primary tumor and/or
related LNs is important for optimal ICB response. However, clinical trials involving patients
with completely resected stage IIIb/c or stage IV melanoma, demonstrate that patients still
benefit from ICB treatment after sentinel LNs and bulky disease-related LNs removal[59,60,70],
arguing TDLNs are not necessary for ICB response. Further, retrospective analysis of stage III
melanoma patients demonstrates that patients benefited similarly from adjuvant ICB, regardless
of the extent of LN surgery. Recent work from our lab and others has shown that metastatic
TDLNs are immune suppressed and can impair systemic anti-cancer immune responses[20-22].
Ultimately, the metastatic burden in TDLNs may determine whether SLNB or CLND has a
positive, negative or neutral effect on disease recurrence and survival. Because human LN chains
are complex with multiple echelons[71], it is key to determine immune activation or suppression
status of LNs downstream of sentinel LNs. Identifying LNs with sufficient exposure to antigen
but not suppressive factors from tumor, may identify optimal target sites for immunotherapy
delivery.

In this study, we emphasize the dynamic and adaptive aspect of antigen transportation.
Disruption of normal lymphatic drainage drives the antigen-carrying fluid to flow to adjacent
lymphosomes and induces immune response in the LNs of that lymphosome. Often these LNs
could be quite distant (and thus non-intuitive) from the original TDLNs, which is corroborated
by clinical observations[49,50]. However, the maximal transport distance of interstitial antigens,
without being substantially diluted and degraded, remains to be determined. Further, limits of
active transport of APCs from the tumor site are also indeterminate. Understanding these spatial
limits can explain the anatomic distribution of LNs for regional immune surveillance.
Intriguingly, we found CLND induces tumor-specific T cells enrichment in distant LNs, when
performed along with primary tumor resection (adjuvant setting). The antigen source fueling the
tumor-specific T cell response in distant LNs is unknown. Conceivably, antigens were vastly
spread by the surgery, from residual cancer cells at the primary tumor site, or in occult metastatic
sites. Despite the importance of antigen trafficking to distant LNs, direct T cell priming in the
local tumor microenvironment[72] and tertiary lymphoid structures[73] may also contribute to
residual ICB response.

The finding that long-term anti-tumor immune memory is not predominantly reliant on the T cell
egress from LNs (**Supp Figure 4**), leads to critical unanswered questions. Do anti-tumor
immune memory cells disseminate from TDLNs immediately and accumulate systemically
during ICB? Or is systemic immune memory mainly formed after primary tumor clearance? Our
neoadjuvant-adjuvant ICB study (**Figure 6**) demonstrated that mice that underwent curative
surgery had systemic immune memory development, indicating immune reactions are present at
the time of tumor resection. This also suggests that adjuvant ICB may act through existing anti-
tumor tissue-resident immune memory cells at the time of treatment. One caveat of the FTY720
experiment to block T cell egress, including memory T cell egress, from LNs is that tissue-
resident memory cells can enter the circulation and reject tumor cells at distant sites[74]. S1P

inhibition mainly blocks lymphocytes egress from LNs and bone marrow[63] not other tissues, and
circulating memory cells are still present in FTY720 treated mice. It is possible a small number
of cancer-specific memory clones are sufficient to suppress recurrence. Additionally, S1P
metabolism promotes metastasis via activating cell-autonomous signaling pathways such as the
C3 complement pathway[75] or PAK1/LIMK1/Cofilin1 pathway[76] in cancer cells, which adds
another caveat of using systemic S1P inhibition.

Additional limitations are present in our preclinical model when used to address clinical
questions[77]. In this work, surgery of skin LNs in mice was used to recapitulate SLNB and CLND
procedures. However, large scale and anatomical differences are present between humans and
mice. On one hand, the size of a mouse is comparable to a human axilla, thus, the murine version
of CLND used in this work may be similar to a SLNB scenario in patients. On the other hand,
compared to simple chains of a few LNs in mice, humans have more complex and interconnected
lymphatic networks as well as multiple LNs draining one common region[71]; hence, removing a
single LN in mice may equate to CLND in human. Encouragingly, consistent with our preclinical
data, compensatory lymphatic drainage[49,50], even in the presence of lymphedema[48,78], and distant
LN activation are observed in cancer patients (**Figure 7**). Further, it is important to investigate
lymphatic drainage patterns after non-skin LN dissection as ICB treatment increases survival in
non-small cell lung cancer[79] and colorectal cancer[80] patients. Notably, regional specification is
found in immunity generated in DLNs[81]. Therefore, after CLND, activated T cells in distant LNs
may not be able to infiltrate the cancer lesion, which is dictated by anatomic location[81]. In
contrast to preclinical studies, clinical trials are more statistically powered[82]. Thus, it is
conceivable that LN dissection can cause a statistically significant difference in ICB-mediated
survival in trials, whereas the biological impact of this difference is expected to be marginal.
Emergence of more powerful biomarkers for patient stratification can enhance ICB response in
specific patient cohorts and may exacerbate the effects of LN dissection. Finally, dissection is
performed on SLNs and metastatic LNs which are immunosuppressed[83] with disrupted CD8 T
cell responses[84]. Consequently, the net benefit of removing these immune suppressive LNs is
likely to outweigh the marginal decrease in systemic response to ICB.

Method:

Cell lines

Parental E0771 (ATCC, CRL-3461), mammary carcinoma line, and parental B16F10 (ATCC,
CRL-6475), melanoma line, were obtained from ATCC. All cells were cultured using
Dulbecco's modified Eagle's medium (DMEM) (Corning, Cat#10-014-CV) supplemented with
10%(v/v) fetal bovine serum (GeminiBio, Cat#100-106) in Forma Steri-cycle incubator (Thermo
Scientific, Cat#381), where the CO_2 concentration was set at 5% and temperature was set at
37 °C. Cells were trypsinized (Gibco, Cat#25-050-CI) and passaged at a 1:5 ratio when reaching
80-90% confluency. All cells were authenticated before use and checked for mycoplasma
monthly using PCR Mycoplasma Test Kit I/C (Promokine, Cat # PK-CA91-102).

Transduction

HEK293 cells were transfected with 1μg of plasmids of gene of interest as well as psPAX2 and
VSVG, using 5% Fugene transfection reagent (Promega, Cat#E2311) diluted in Opt-MEM
(Gibco, Cat#11058-021). After overnight incubation, media containing transfection reagents was
replaced with fresh culture media. After another 2 days, lentiviral supernatant was collected,
filtered and applied with polybrene (8μg/ml) to target cells.

pLV-Ef1a-ZSgreen-SIINFEKL-IRES-puro plasmid (a gift from Stefanie Spranger lab at MIT)
was used to generate ovalbumin (257-264) overexpressing cells. CSCW-mbGluc-biotin-IG
plasmid (generated by MGH Vector Core)[85] was used to generate Gaussia luciferase expressing
cells.

Animal models

In addition to wild-type C57Bl6 mice, lymphotoxin alpha knockout mice B6.129s2-Lta^{tm1Dch}/J
(the Jackson Laboratory, Strain# 002258) were used for tumor immunology and dextran
trafficking study. Animal experiment protocols were approved by the Institutional Animal Care
and Use Committee of the Massachusetts General Hospital, Boston, MA.

Tumor study and ICB administration

200,000 E0771 cells or 500,000 B16F10 cells were trypsinized and resuspended in DMEM base
media before implantation. E0771 cells were injected to 4th mammary fat pad of female C57BL6
mice while B16F10 cells were injected to the dermis (atop bottom flank) of male or female
C57BL6 mice. After 7 days, anti-PD1 blocking antibody (Bio X cell, Cat# BP0146) at 10mg/kg
was given intraperitoneally or anti-PD1 and anti-CTLA4 (Bio X cell, Cat# BP0131) blocking
antibodies each at 5mg/kg were given intraperitoneally or intradermally. Additional ICB was
administered on day 10 and 13 for E0771 tumors or day 9 and 11 for B16F10 tumors. In
neoadjuvant-adjuvant treatment regimen, the adjuvant ICB was given when secondary tumors
reach the size greater than 100mm^3. Rat (Jackson ImmunoResearch Lb, Cat#012-000-003) and
Syrian Hamster (Leinco Technologies, Cat# I-444) IgGs were used as control.

Tumor volume (ellipsoid shape is assumed) was monitored at least twice a week and mice were
sacrificed when tumor volume reached 1000mm^3 or any tumor dimension was greater than
15mm, or when ulceration is greater than 4mm in any dimension. Tumor rechallenge was
performed on complete responders, which were defined as mice that remained tumor free for at
least 60 days after initial inoculation. In certain studies, LNs were harvested and volume was
measured using caliper (ellipsoid shape is assumed).

Surgical procedures

<u>Primary tumor resection:</u> After anesthesia, the fur growing over the tumor site was removed by
hair clippers and hair removal cream and the skin sterilized. A small skin incision was made to
begin separating the tumor from the underlying fascia and overlying skin through blunt
dissection. The tumor was then excised where necessary. The skin was closed with wound clips
that were subsequently removed after 2 weeks when the wound is completely closed.

<u>LN dissection:</u> Mice skin was shaved and sterilized by wiping ethanol prior to surgery. A 1cm
incision was made to skin atop the inguinal or axillary region, which allowed the surgical
removal of either inguinal or axillary and brachial LNs, while a 0.5cm incision was made to
popliteal skin to remove the popliteal LN. The skin was closed after LN dissection with wound
clips. Mice were let recover for at least 10 days before tumor implantation or tracer injection.

<u>Splenectomy:</u> A 1 cm incision was introduced to the right of ventral midline (below diaphragm)
to open peritoneal cavity. Splenic arteries and veins were ligated at the splenic hilum with a 3–0
silk suture and the spleen was removed. The peritoneum was closed after splenectomy with
polydioxanone suture, while skin was closed with wound clips. Mice were let recover for at least
14 days before tumor implantation.

Dextran tracing

50mg/ml TRITC-dextran with 4400 Dalton molecular weight (Sigma Aldrich, Cat# T1037-
50MG), mixed with 25mg/ml FITC-dextran with 2 million Dalton molecular weight (Sigma
Aldrich, Cat# FD2000S-100MG) in a total volume of 20μl, were injected to mammary fat pad,
dorsal foot or dermis of mice (specified in text). After 6 hours, ipsilateral and contralateral
popliteal, inguinal, axillary and brachial LNs as well as spleens were harvested and stained for
microscopy.

Immune cells tracing using Kaede mice

500,000 B16F10-OVA cells were implanted to the lower flank dermis of Kaede mice B6.Cg-Tg
(CAG-tdKaede) 5Utr (Riken Bioresource Center, RBRC05737). After 2 weeks, primary tumors

were resected, and two doses of 10mg/kg anti-PD1 antibodies were administered
intraperitoneally (two days and four days after tumor resection). After another 2 weeks,
2,500,000 B16F10-OVA cells were implanted to the primary tumor site as rechallenge.
Contralateral inguinal LNs were photoconverted (405nm UV light at 12V/0.09A for 5mins),
when recurrent tumors exceeded 100mm^3. After 1 day, recurrent tumors were harvested for flow
analysis.

Antigen-presenting cells-based vaccination

Bone marrow was flushed out from hind limbs of 7-12 week old C57Bl6 mice. After ACK buffer
lysis, cells were cultured in RPMI media supplemented with 10% FBS, 1% Pen/Strep, 50μM β-
mercaptoethanol, 20ng/ml GM-CSF (PeproTech, Cat# 315-03) and 10ng/ml IL-4 (PeproTech,
Cat# 214-14). Media was refreshed every three days to generate BMDCs. At day 9, BMDCs
were pulsed with 20μg/ml OVA$_{257-264}$ (Invitrogen, Cat# vac-sin) for 3 hours, then 2μg/ml LPS
(Sigma, Cat# L-2630-10MG) was added as adjuvant. The next day, 500,000 activated BMDCs
(in 20μl DMEM base media) were collected and injected to the foot of C57Bl6 mice that
underwent sham surgery or LN surgery. After 7 days, LNs were harvested from immunized mice
for further analysis.

Immunofluorescence staining and imaging

Fresh tissue was embedded in OCT compound (Tissue-Tek, Cat# FIS 23730571) and frozen at -
80 °C for overnight. Subsequently, frozen tissue was cut into serial sections with 10μm interval
using cryostat station. 3-5 tissue sections representing different tissue depths were used for
staining. Tissue sections were fixed using 4% paraformaldehyde for 15mins at room temperature.
After fixation, slides were rinsed in PBS-T and blocked in 5% normal donkey serum for 1 hour.
Then slides were stained with primary antibody in humidity chamber for overnight at 4 degree.
Next day, primary antibody was washed away, and slides were incubated with secondary
antibody for 1 hour at room temperature. At last, DAPI was added to label DNA and slides were
mounted. Slides were imaged using Olympus IX81 fluorescence microscope with FV10-ASW
4.2 software or ZEISS Axio scan Z.1 with ZEN 2.3 slidescan software.

Flow cytometry

Fresh tissue was dissociated and filtered using 70μm mesh. Red blood cells were lysed by
incubation with ACK lysis buffer for 2 minutes, followed by neutralization using PBS with 10%
FBS. Next, samples were stained with viability dye: DAPI or Zombie Violet (1:200) (Biolegend,
Cat# 423113) for 15 mins, then blocked by Fc receptor (CD16/32) antibody (1:500) (Biolegend,
Cat# 101302) for 15 mins. After blocking, samples were stained with primary antibody (or
tetramer) for 1 hour at 4 °C in the dark, then washed with PBS and fixed using BD fixative. At
last, samples were analyzed by Fortessa X20 cytometer within two weeks of fixation.

Image processing

All image processing of fluorescence images was done in ImageJ (https://imagej.nih.gov/ij/) or
QuPath (https://qupath.github.io) to adjust brightness and contrast. Constant adjustment setting
was done to all images from individual experiment. For signal quantification, Cellprofiler
(https://cellprofiler.org) and Qupath were used. To be specific, the tissue border was identified
from each image manually. Specific staining fluorescence signal was identified by adaptive
thresholding-minimum cross-entropy-algorithm. Positive signal objects were de-clumped by
intensity (Cellprofiler), or positive cell detection command with manual threshold (QuPath).
Another object size-based filter was additionally applied to cell identification (1-10μm) to
remove non-specific staining.

Retrospective analysis of melanoma patient survival

We performed a retrospective analysis of patients with stage III cutaneous melanoma according to
the 8th American Joint Committee on Cancer criteria. The patients were selected from a large

melanoma cohort study and tumor biobank maintained at Massachusetts General Hospital for over
13 years (2009-2022) and approved by the Institutional Review Board. Informed consent was
obtained from all patients. All patients included in this analysis received at least one line of PD1
immune checkpoint blockade adjuvant therapy. For each patient, we extracted general
demographic information, histopathological reports, extent of surgery, adjuvant treatments,
recurrence and survival outcomes. The data extraction was followed by statistical analysis using
R (Core Team 2021) package[86] (survminer) to generate Kaplan-Meier survival curves.
Significance was determined using a log-rank test and the p-value is reported on each plot. P-value
significance was considered to be p<0.05.

Clinical trial data and patient records

In phase I/Ib trial NCT03635164, patient response rate (Laurel Darragh & Sana Karam):
Pathological response was evaluated independently by two board-certified pathologists who were
blinded to treatment outcome. Specimens collected at time of surgery were evaluated for
microscopic residual tumor (only scattered residual foci) or macroscopic residual tumor and
scored based on these findings.

In phase I/Ib trial NCT03635164, radiology images (Laurel Darragh & Sana Karam): At
baseline, patients received a PET/CT of skull base to mid-thigh and an MRI with and without
contrast of the neck no more than 28 days before radiation therapy. Long-term follow up imaging
assessment was done with either PET/CT and/or MRI at the discretion of the treating physician.
Enlarged lymph nodes at the follow-up timepoints were noted on imaging by a radiologist and
concerning nodes were biopsied to rule out recurrence.

Statistics

Statistical significance was determined with either two-tailed unpaired Student's t test (for two
conditions) or one-way ANOVA (for conditions more than 2) with Tukey post-hoc adjustment.
Log-rank (Mantel-Cox) test was used for overall survival. Data are presented as mean ± SEM.
Fisher's exact test was used to analyze contingency tables including responders vs. non-
responders and survivors vs. non-survivors. Statistical significance threshold was set at $p<0.05$.
All calculation was performed using GraphPad Prism 9 software.

Mathematical modeling

The mathematical model predicts interstitial fluid pressures, flow rates, and the patterns of fluid
redistribution after selective disabling of LNs. Details of the model formulation, validations and
parameter dependence are available in the **Supplemental Materials**. In summary, our
mathematical model focuses on transport of antigen, antigen-presenting cells and tracer by fluid
flow rather than molecular diffusion. Relatively small molecules such as the 4k MW dextran take
only minutes to diffuse along concentration gradients over the distances between blood vessels
or within lymph nodes (~100 μm) so that they can quickly reach the bloodstream for distribution
throughout the body. In contrast, larger materials such as 2M MW dextran might take days or
longer to diffuse similar distances which means that the larger tracer, free antigens and cells
move predominantly by interstitial flow.

Interstitial fluid in the skin is assumed to arise from the semipermeable walls of the capillaries,
and then flow locally through the interstitial space to the nearest initial lymphatic capillaries. The
initial lymphatic vessels in the skin, which lack intraluminal valves and contractile muscle, form
a highly redundant meshwork that carry fluid passively down gradients in the interstitial pressure
to the nearest collecting lymphatic vessels which then actively pump lymph toward the sentinel
lymph node for that lymphosome of the skin[18]. The boundaries of the drainage basins or
lymphosomes are determined by the locations of the collecting vessels and which lymph nodes
are downstream.

Antibodies and tetramer

Information (source and concentration) of antibodies and tetramer used in immunofluorescence and flow cytometry are listed in **Supplemental Materials**.

Acknowledgement

This work was financially supported by grants from NIH (K00CA234940 to H.Z.; R21AI097745, R01CA214913, R01HL128168, and R01CA284372 to T.P.P; R01CA284603 to T.P.P., L.L.M. and J.W.B.; 1R01DE028529-01, 1R01DE028282-01, 1R01DE028529-01 and 1P50CA261605-01 to S.D.K.; R35CA242379 to M.G.V.H.), the Rullo Family MGH Research Scholar Award (T.P.P.; G.M.B.), the Patricia K. Donahoe Award from the Huiying Foundation (G.M.B.), and the Adelson Medical Research Foundation (G.M.B.), the Ludwig Center at MIT (M.G.V.H.) and the MIT Center for Precision Cancer Medicine (M.G.V.H.), the Breast Cancer Alliance (J.M.U.), and the Melanoma Research Foundation (J.M.U.). The Phase I/Ib clinical trial NCT03635164 was funded by AstraZeneca, S.D.K also received clinical trial funding from Genentech and preclinical funding from Roche.

Competing Interests

M.G.V.H. acknowledges that he is a scientific advisor for Agios Pharmaceuticals, iTeos Therapeutics, Sage Therapeutics, Pretzel Therapeutics, Lime Therapeutics, Auron Therapeutics, and Droia Ventures. G.M.B has sponsored research agreements through her institution with: Olink Proteomics, Teiko Bio, InterVenn Biosciences, Palleon Pharmaceuticals. She served on advisory boards for: Iovance, Merck, Nektar Therapeutics, Novartis, and Ankyra Therapeutics. She consults for: Merck, InterVenn Biosciences, Iovance, and Ankyra Therapeutics. She holds equity in Ankyra Therapeutics.

724
725 **Figure 1. Adjuvant ICB response is not affected by the extent of LN surgery in stage III melanoma patients.**
726 Retrospective analysis was performed on a cohort of cutaneous stage III melanoma patients who underwent SLNB
727 or CLND, and were treated by adjuvant nivolumab or pembrolizumab at Massachusetts General Hospital. (**A**)
728 Melanoma-specific survival, (**B**) recurrence-free survival, (**C**) local recurrence-free survival and (**D**) distant
729 recurrence-free survival are plotted.

Figure 2. ICB response persists after LN dissection. (**A**) Sham surgery or SLNB was performed on C57Bl6 mice (day -10) prior to orthotopic tumor implantation (day 1). Mice were treated by rat IgG control or anti-PD1 antibody (day 7) three times (every 2 days for B16F10 or every 3 days for E0771, 10mg/kg). Tumor volume (*left & middle*) and overall survival (*right*) were plotted for mice bearing orthotopic B16F10 (*top*) or E0771 (*bottom*) tumors, respectively. (**B**) To evaluate the impact of more aggressive surgery, sham surgery or CLND was performed followed by tumor implantation and subsequent anti-PD1 inhibition using the same regimen described above. Tumor volume (*left*) and overall survival (*right*) were plotted for mice bearing B16F10 (*top*) or E0771 (*bottom*) tumors, respectively. (**C**) In contrast with pre-emptive approach, impact of post-tumor implantation LN dissection was also assessed. After tumor implantation (day 1), sham surgery, SLNB or CLND was performed on C57Bl6 mice (day 6) right before anti-PD1 treatment (day 7). Tumor volume (*left*) and overall survival (*right*) were plotted for mice bearing B16F10 (*top*) and E0771 (*bottom*) tumors. (**D**) Frequency of complete responders from pre-emptive and post-tumor TDLN removal studies were plotted. Fisher's exact test was used for statistical analysis. (**E**) B16F10 (*top*) or E0771 (*bottom*) tumor response to anti-PD1 treatment was tested in *Lta^{tm1Dc}* mice. ICB response of wildtype (WT) sham mice from (**A**) SLNB study was plotted in dashed lines for comparison. Tumor volume (*left*) and overall survival (*right*) were plotted. In all animal studies, "(n)" indicates number of mice used. Log-rank test with Bonferroni test was used for statistical analysis of survival data. *$p<0.05$, ****$p<0.0001$.

748
Figure 3. DLN dissection alters lymph drainage and promotes antigen access to distant lymphoid organs.
FITC-conjugated 2 million Dalton (FITC-2MD) and TRITC-conjugated 4.4 kilodalton (TRITC-4.4kD) dextran were
injected to the mammary fat pad of mice that underwent sham surgery, SLNB or CLND. (**A**) Representative
immunofluorescence images of mouse with sham surgery demonstrate intranodal dextran distribution. (**B**) Dextran
distribution in ipsilateral inguinal (IN), axillary (AX), brachial (BR) LNs as well as contralateral IN LNs in animals
that underwent sham, SLNB or CLND. (**C**) Greater magnification image shows that large dextran molecules enter
subcapsular sinus of contralateral IN LN in CLND mice, suggesting transport through lymphatic vessels. (**D**) FITC-
2MD dextran distribution in contralateral inguinal LN from CLND mice with or without acute midline skin incision,
prior to dextran injection to mammary fat pad. (**E**) Diagram shows lymph flow pattern in various surgery settings.
(**F**) FITC-2MD dextran distribution in spleens from mice that underwent sham surgery or ipsilateral LN dissection
(ILND). (**G**) Representative immunofluorescence images demonstrate FITC-2MD dextran location in spleen from
ILND mice relative to B cells (B220) and macrophages (F4/80). In this figure, fluorescence signal quantification
was calculated by the occupied area, while DAPI was used to normalize LN area. Two tailed student t test (two
conditions) and one way ANOVA with Tukey post analysis (more than two conditions) were used for statistical
analysis. *$p<0.05$, **$p<0.01$, ****$p<0.0001$.
764

Figure 4. TDLN dissection stimulates immune responses in distant LNs. To assess the tumor-specific T cell response, B16F10 tumors overexpressing Ovabulmin$_{257-264}$ (OVA) were implanted to sham and CLND mice followed by anti-PD1 inhibition. (**A**) In ipsilateral and contralateral LNs as well as (**B**) spleens, CD8 T cell expansion is shown by immunofluorescence images. (**C**) Additionally, OVA-specific CD8 T cells in ipsilateral, contralateral and cervical LNs, spleens and blood were determined by tetramer staining using flow cytometry analysis. Gating approach and representative density plots from ipsilateral and contralateral LNs are shown. (**D**) Abundance of OVA-specific CD8 T cells (CD8+, Tetramer+) is plotted for various LNs, blood and spleens collected from sham and CLND mice; each dot represents data from a single mouse. (**E**) OVA-specific CD8 T cell response in ipsilateral and contralateral LNs is shown in an adjuvant setting where primary tumor resection was performed with or without CLND before anti-PD1 inhibition. (**F**) Similar to **E**, Kaede mice bearing B16F10-OVA tumors were treated by adjuvant anti-PD1 inhibition, followed by tumor rechallenge to primary site. Once recurrent tumor formed, contralateral inguinal LNs were photoconverted prior to tissue harvesting for flow analysis. Abundance of Kaede red+ tumor infiltrating OVA-specific CD8 T cells is plotted. (**G**) To assess antigen-presenting cell-mediated T cell response after LN dissection, BMDCs were pulsed with OVA and LPS for 1 day, and injected to the footpad of mice that underwent sham surgery or ILND. OVA-specific CD8 T cells in ipsilateral and contralateral LNs are plotted. In this figure, fluorescence signal quantification is calculated by the occupied area, while DAPI is used to normalize. Two tailed student t test (two conditions) and one way ANOVA with Tukey post analysis (more than two conditions) were used for statistical analysis. *$p<0.05$, **$p<0.01$, ****$p<0.0001$.

Modeling of interstitial fluid transport in mouse after LN dissection

Modeling of interstitial fluid transport in human after LN dissection

Simplified model for lymphosome drainage

Active lymphosomes
Disabled lymphosome

Figure 5. Mathematical modeling of antigen access to adjacent lymphosomes via dermal network after node dissection. (**A**) Mouse model with network of collecting lymphatic vessels (black), collecting points (blue) and lymph nodes (red). (**B**) Computational mesh for network of initial lymphatic vessels in the skin. (**C-F**) Drainage basins (red) with disabled nodes (black *), four most active compensatory nodes (black o) for (**C**) inguinal node resection, (**D**) inguinal and axillary node resection, (**E**) popliteal, inguinal, axillary and brachial node resection, and (**F**) inguinal and axillary node resection with severed midline model (intact **C-E**). (**G**) Human male model with network of collecting lymphatic vessels (black), collecting points (blue) and lymph nodes (red). (**H**) Computational mesh for network of initial lymphatic vessels in the skin. (**I**) Drainage basin (red) with disabled axillary node (black *), four most active compensatory nodes (black o). (**J**) Planar geometry for analytical model with a circular disabled lymphosome (red) surrounded by tissue with active lymph clearance (blue). (**K**) Interstitial pressure inside and outside of the disabled lymphosome from the full, 3D human model compared to an analytical result for the planar

797   geometry in (**J**) with the same surface area. (**L**) Dependence of interstitial pressure on radius of disabled
798   lymphosome (arrows) showing how small lymphosomes produce little increase in interstitial pressure. (**M**)
799   Dependence of interstitial pressure and fluid velocity u at the periphery of the disabled lymphosome on tissue
800   permeability (radius 50 mm) showing how changes in tissue permeability can alter the interstitial pressure, but
801   barely affects the rate of fluid/antigen flow to compensatory lymphosomes (u).
802

803
Figure 6. Distant LNs can be targeted to improve ICB response. Sham surgery or CLNB was performed on
C57Bl6 mice (day -10) prior to orthotopic E0771 tumor implantation (day 1). IgG control or combination of anti-
PD1 and anti-CTLA4 antibodies were intradermally delivered to ipsilateral TDLNs or contralateral LNs in sham and
CLND conditions, respectively (day 7) for three doses at 5mg/kg. LNs exposed to treatment were harvested for flow
cytometry analysis measuring CD45, TCR-beta, CD4, CD8, CD44, PD-1, TCF-1, CD39 and CD62L. Non-tumor
bearing naïve mice are used as control. (**A**) LN Volume of naïve mice or mice treated by IgG or ICB combination is
plotted. (**B**) Principle component analysis is performed on naïve or treatment-exposed LNs, using the abundance of
antigen-experienced T cells (CD44+, PD-1+), memory T cells (CD44+, PD-1+, TCF1+), central memory T cells
(CD44+, PD-1+, TCF1+, CD62L+), effector memory T cells (CD44+, PD-1+, TCF1+, CD62L-), stem-like effector
memory T cells (CD44+, PD-1+, TCF1+, CD62L-, CD39-) and their CD4/CD8 subsets. (**C**) The percentage of
stem-like effector memory T cells to total T cells is plotted. (**D**) Using the same surgery, tumor implantation and
treatment procedure, ICB was given intraperitoneally or intradermally to the lymphatic basin of contralateral
inguinal LN (day 7) for three times at 5mg/kg in sham and CLND mice. Tumor volume (*left*) and overall survival
(*right*) were plotted. (**E**) Timeline for neoadjuvant-adjuvant treatment and tumor rechallenge. (**F**) Representative
images of vitiligo (arrows) that developed in mice after neoadjuvant treatment (*left*). Percentage of mice with local,
distant or local & distant vitiligo across conditions (*right*). (**G**) Overall survival of mice treated by neoadjuvant ICB
with various LN surgery status. (**H**) Tumor-free survival of mice previously treated by neoadjuvant ICB after tumor
rechallenge. Naïve mice were used as control. (**I**) Overall survival of mice previously treated by neoadjuvant-
adjuvant ICB after tumor rechallenge. Of note, only mice that developed tumors after rechallenge were given
adjuvant treatment. In all studies, "(n)" indicates the number of mice used. One way ANOVA with Tukey post
analysis (comparing more than two conditions) and Log-rank test with Bonferroni test (survival) were used for
statistical analysis. *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.
826

827
Figure 7. Distant LNs are inflamed after resection of tumor and related LNs in cancer patients. (**A**) *Left*, from
phase I trial NCT03635164, response rate and reactive status of remaining LNs after neck dissection are shown.
Right, percentage of patients that have enlarged or reactive LNs in more distal LN basins are plotted in non-
responder and responder cohorts. Fisher's exact test 2x2 is used to evaluate statistical significance, *p<0.05. (**B**) CT
images of patient 01-013's and 01-015's neck at diagnosis and 2-year post-surgery follow-up scan are demonstrated.
Arrows indicate enlarged neck LNs (posterior cervical LNs in patient 01-013; submandibular and submental LNs in
patient 01-015). (**C**) CT and PET images of patient 01-009 at diagnosis and 7-month post-surgery follow-up scan.
Arrows indicate reactive cervical (diagnosis) and apex mediastinal LNs (follow up).
836

E0771 ectopic implantation to dermis

837
Supplementary Figure 1. Aggressive LN dissection damages ICB response in ectopic setting. (**A**) For pre-
emptive setting, sham surgery or CLND was performed on C57Bl6 mice (day -10) prior to E0771 tumor
implantation to dermis (day 1). Mice were treated by anti-PD1 inhibition (day 7) for three times at 10mg/kg. Tumor
volume (*left*) and overall survival (*right*) were plotted. (**B**) For post-tumor setting, after E0771 tumor implantation to
dermis (day 1), sham surgery, SLNB or CLND was performed on C57Bl6 mice (day 6) right before anti-PD1
treatment. Tumor volume (*left*) and overall survival (*right*) were plotted. In all animal studies, "(n)" indicates the
number of mice used. Log-rank test with Bonferroni test was used for statistical analysis of survival data. *$p<0.05$.
845

2M Dextran 4.4k Dextran DAPI

846
Supplementary Figure 2. Lack of primary drainage promotes antigen to access distant LNs. FITC-conjugated
2 million Dalton (FITC-2MD dextran) and TRITC-conjugated 4.4 kilo Dalton dextran (TRITC-4.4kD dextran) were
injected to the footpad of mice have undergone sham surgery or ipsilateral LN dissection (ILND). Representative
immunofluorescence images demonstrate dextran distribution in ipsilateral popliteal, inguinal, axillary, brachial LNs
as well as contralateral counterparts. In this figure, fluorescence signal quantification is calculated by the occupied
area, while DAPI is used to normalize the LN area. One way ANOVA with Tukey post analysis was used for
statistical analysis, all conditions were compared to Sham ipsilateral popliteal LN, *$p<0.05$.
854

855
Supplementary Figure 3. Spleen contributes marginally to residual ICB response after CLND. (**A**) Sham
surgery (SP+ condition) or splenectomy (SP- condition) was performed on mice 14 days prior to CLND. E0771
tumor implantation and IgG or anti-PD1 treatment were conducted using same procedure as before. Tumor volume
(*left*) and overall survival (*right*) were plotted. (**B**) E0771 tumor rechallenge was performed using complete ICB
responders. Tumor volume was plotted when tumor was palpable (after day 7) and when the study was ended. In all
animal studies, "(n)" indicates the number of mice used. Log-rank test with Bonferroni test was used for statistical
analysis of survival data. *$p<0.05$.
863

864
Supplementary Figure 4. LN-egressing immune cells are not necessary for long-term anti-tumor immune
memory. (**A-C**) In these rechallenge studies, tumor volume was plotted when tumor was palpable (after day 7) and
when the mouse was euthanized. (**A**) E0771 tumors were implanted orthotopically to the primary tumor site of naïve
mice or complete ICB responders. (**B**) E0771 tumors were orthotopically implanted to both ipsilateral (primary
tumor site) and contralateral sides of naïve mice or complete ICB responders. (**C**) B16F10 tumors were implanted to
the mammary fat pad (*left*) or dermis (*right*) of naïve mice or E0771-immune mice. (**D**) To test the potency of
FTY720 in depleting circulating lymphocytes, mice were given water containing or FTY720 (2.5ug/ml) for 5 days
or maintained on regular water (control). Blood was then collected for characterization. *Left*, representative density
plot demonstrates the abundance of circulating B cells (CD45+, B220+), T cells (CD45+, TCR-beta+) and central
memory T cells (CD45+, TCR-beta+, CD8+, CD44+, CD62L+). *Right,* quantification of CD4, CD8 and central
memory T cells, relative to live cell gate, was graphed. To test the contribution of lymphocytes egressing from
lymphoid organs to anti-tumor immune memory, E0771 tumor rechallenge to 4th mammary fat pad (**E**) and dermis
(**F**) were performed sequentially on both sides in complete ICB responders fed with vehicle or FTY720 water.
Tumor naïve mice were used as controls. Tumor volume (*left*) and overall survival (*right*) were plotted. Of note,
dermal tumors were often ulcerated leading to euthanasia. In all animal studies, "(n)" indicates the number of mice
used. Log-rank test with Bonferroni test was used for statistical analysis of survival data. ****$p<0.0001$.
881
882

References:

Fransen, M. F. *et al.* Tumor-draining lymph nodes are pivotal in PD-1/PD-L1 checkpoint therapy. *JCI Insight* **3** (2018). https://doi.org:10.1172/jci.insight.124507

Zhao, X., Kassaye, B., Wangmo, D., Lou, E. & Subramanian, S. Chemotherapy but Not the Tumor Draining Lymph Nodes Determine the Immunotherapy Response in Secondary Tumors. *iScience* **23**, 101056 (2020). https://doi.org:10.1016/j.isci.2020.101056

Ishida, Y., Agata, Y., Shibahara, K. & Honjo, T. Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J* **11**, 3887-3895 (1992). https://doi.org:10.1002/j.1460-2075.1992.tb05481.x

Iwai, Y. *et al.* Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *Proc Natl Acad Sci U S A* **99**, 12293-12297 (2002). https://doi.org:10.1073/pnas.192461099

Brunet, J. F. *et al.* A new member of the immunoglobulin superfamily--CTLA-4. *Nature* **328**, 267-270 (1987). https://doi.org:10.1038/328267a0

Leach, D. R., Krummel, M. F. & Allison, J. P. Enhancement of antitumor immunity by CTLA-4 blockade. *Science* **271**, 1734-1736 (1996). https://doi.org:10.1126/science.271.5256.1734

Hargadon, K. M., Johnson, C. E. & Williams, C. J. Immune checkpoint blockade therapy for cancer: An overview of FDA-approved immune checkpoint inhibitors. *Int Immunopharmacol* **62**, 29-39 (2018). https://doi.org:10.1016/j.intimp.2018.06.001

Wei, S. C., Duffy, C. R. & Allison, J. P. Fundamental Mechanisms of Immune Checkpoint Blockade Therapy. *Cancer Discov* **8**, 1069-1086 (2018). https://doi.org:10.1158/2159-8290.CD-18-0367

Hodi, F. S. *et al.* Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* **363**, 711-723 (2010). https://doi.org:10.1056/NEJMoa1003466

Robert, C. *et al.* Nivolumab in previously untreated melanoma without BRAF mutation. *N Engl J Med* **372**, 320-330 (2015). https://doi.org:10.1056/NEJMoa1412082

Borghaei, H. *et al.* Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer. *N Engl J Med* **373**, 1627-1639 (2015). https://doi.org:10.1056/NEJMoa1507643

Brahmer, J. *et al.* Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer. *N Engl J Med* **373**, 123-135 (2015). https://doi.org:10.1056/NEJMoa1504627

Motzer, R. J. *et al.* Nivolumab plus Ipilimumab versus Sunitinib in Advanced Renal-Cell Carcinoma. *N Engl J Med* **378**, 1277-1290 (2018). https://doi.org:10.1056/NEJMoa1712126

Chen, R. *et al.* Phase II Study of the Efficacy and Safety of Pembrolizumab for Relapsed/Refractory Classic Hodgkin Lymphoma. *J Clin Oncol* **35**, 2125-2132 (2017). https://doi.org:10.1200/JCO.2016.72.1316

El-Khoueiry, A. B. *et al.* Nivolumab in patients with advanced hepatocellular carcinoma (CheckMate 040): an open-label, non-comparative, phase 1/2 dose escalation and expansion trial. *Lancet* **389**, 2492-2502 (2017). https://doi.org:10.1016/S0140-6736(17)31046-2

16   Duffy, A. G. *et al.* Tremelimumab in combination with ablation in patients with advanced
hepatocellular carcinoma. *J Hepatol* **66**, 545-551 (2017).
https://doi.org:10.1016/j.jhep.2016.10.029

17   Janjigian, Y. Y. *et al.* First-line nivolumab plus chemotherapy versus chemotherapy alone
for advanced gastric, gastro-oesophageal junction, and oesophageal adenocarcinoma
(CheckMate 649): a randomised, open-label, phase 3 trial. *Lancet* **398**, 27-40 (2021).
https://doi.org:10.1016/S0140-6736(21)00797-2

18   Padera, T. P., Meijer, E. F. & Munn, L. L. The Lymphatic System in Disease Processes and
Cancer Progression. *Annu Rev Biomed Eng* **18**, 125-158 (2016).
https://doi.org:10.1146/annurev-bioeng-112315-031200

19   Zhou, H., Lei, P. J. & Padera, T. P. Progression of Metastasis through Lymphatic System.
Cells **10** (2021). https://doi.org:10.3390/cells10030627

20   Jones, D. *et al.* Solid stress impairs lymphocyte infiltration into lymph-node metastases.
Nat Biomed Eng **5**, 1426-1436 (2021). https://doi.org:10.1038/s41551-021-00766-1

21   Lei, P. J. *et al.* Cancer cell plasticity and MHC-II-mediated immune tolerance promote
breast cancer metastasis to lymph nodes. *J Exp Med* **220** (2023).
https://doi.org:10.1084/jem.20221847

22   Reticker-Flynn, N. E. *et al.* Lymph node colonization induces tumor-immune tolerance to
promote distant metastasis. *Cell* **185**, 1924-1942 e1923 (2022).
https://doi.org:10.1016/j.cell.2022.04.019

23   Brown, M. *et al.* Lymph node blood vessels provide exit routes for metastatic tumor cell
dissemination in mice. *Science* **359**, 1408-1411 (2018).
https://doi.org:10.1126/science.aal3662

24   Pereira, E. R. *et al.* Lymph node metastases can invade local blood vessels, exit the node,
and colonize distant organs in mice. *Science* **359**, 1403-1407 (2018).
https://doi.org:10.1126/science.aal3622

25   Alatengbaolide *et al.* Lymph node ratio is an independent prognostic factor in gastric
cancer after curative resection (R0) regardless of the examined number of lymph nodes.
Am J Clin Oncol **36**, 325-330 (2013). https://doi.org:10.1097/COC.0b013e318246b4e9

26   Aoyama, T. *et al.* The Lymph Node Ratio Is an Independent Prognostic Factor in
Pancreatic Cancer Patients Who Receive Curative Resection Followed by Adjuvant
Chemotherapy. *Anticancer Res* **38**, 4877-4882 (2018).
https://doi.org:10.21873/anticanres.12801

27   Chen, C. C., Lin, J. C. & Chen, K. W. Lymph node ratio as a prognostic factor in head and
neck cancer patients. *Radiat Oncol* **10**, 181 (2015). https://doi.org:10.1186/s13014-015-
0490-9

28   Scoggins, C. R. *et al.* Prognostic information from sentinel lymph node biopsy in patients
with thick melanoma. *Arch Surg* **145**, 622-627 (2010).
https://doi.org:10.1001/archsurg.2010.115

29   Voordeckers, M., Vinh-Hung, V., Van de Steene, J., Lamote, J. & Storme, G. The lymph
node ratio as prognostic factor in node-positive breast cancer. *Radiother Oncol* **70**, 225-
230 (2004). https://doi.org:10.1016/j.radonc.2003.10.015

30   Veronesi, U. *et al.* A randomized comparison of sentinel-node biopsy with routine
axillary dissection in breast cancer. *N Engl J Med* **349**, 546-553 (2003).
https://doi.org:10.1056/NEJMoa012782
31   Cheng, A. & Lang, J. Survival Analysis of Lymph Node Resection in Ovarian Cancer: A
Population-Based Study. *Front Oncol* **10**, 355 (2020).
https://doi.org:10.3389/fonc.2020.00355
32   Gortzak-Uzan, L. *et al.* Sentinel lymph node biopsy vs. pelvic lymphadenectomy in early
stage cervical cancer: is it time to change the gold standard? *Gynecol Oncol* **116**, 28-32
(2010). https://doi.org:10.1016/j.ygyno.2009.10.049
33   Faries, M. B. *et al.* Completion Dissection or Observation for Sentinel-Node Metastasis in
Melanoma. *N Engl J Med* **376**, 2211-2222 (2017).
https://doi.org:10.1056/NEJMoa1613210
34   Dammeijer, F. *et al.* The PD-1/PD-L1-Checkpoint Restrains T cell Immunity in Tumor-
Draining Lymph Nodes. *Cancer Cell* **38**, 685-700 e688 (2020).
https://doi.org:10.1016/j.ccell.2020.09.001
35   Huang, Q. *et al.* The primordial differentiation of tumor-specific memory CD8(+) T cells
as bona fide responders to PD-1/PD-L1 blockade in draining lymph nodes. *Cell* (2022).
https://doi.org:10.1016/j.cell.2022.09.020
36   Francis, D. M. *et al.* Blockade of immune checkpoints in lymph nodes through
locoregional delivery augments cancer immunotherapy. *Sci Transl Med* **12** (2020).
https://doi.org:10.1126/scitranslmed.aay3575
37   O'Melia, M. J., Manspeaker, M. P. & Thomas, S. N. Tumor-draining lymph nodes are
survival niches that support T cell priming against lymphatic transported tumor antigen
and effects of immune checkpoint blockade in TNBC. *Cancer Immunol Immunother* **70**,
2179-2195 (2021). https://doi.org:10.1007/s00262-020-02792-5
38   Eggermont, A. M. *et al.* Adjuvant ipilimumab versus placebo after complete resection of
high-risk stage III melanoma (EORTC 18071): a randomised, double-blind, phase 3 trial.
Lancet Oncol **16**, 522-530 (2015). https://doi.org:10.1016/S1470-2045(15)70122-1
39   van Pul, K. M. *et al.* Local delivery of low-dose anti-CTLA-4 to the melanoma lymphatic
basin leads to systemic Treg reduction and effector T cell activation. *Sci Immunol* **7**,
eabn8097 (2022). https://doi.org:10.1126/sciimmunol.abn8097
40   Tohme, S., Simmons, R. L. & Tsung, A. Surgery for Cancer: A Trigger for Metastases.
Cancer Res **77**, 1548-1552 (2017). https://doi.org:10.1158/0008-5472.CAN-16-1536
41   Tang, F., Tie, Y., Tu, C. & Wei, X. Surgical trauma-induced immunosuppression in cancer:
Recent advances and the potential therapies. *Clin Transl Med* **10**, 199-223 (2020).
https://doi.org:10.1002/ctm2.24
42   Nakajima, Y. *et al.* Near-Infrared Fluorescence Imaging Directly Visualizes Lymphatic
Drainage Pathways and Connections between Superficial and Deep Lymphatic Systems
in the Mouse Hindlimb. *Sci Rep* **8**, 7078 (2018). https://doi.org:10.1038/s41598-018-
25383-y
43   De Togni, P. *et al.* Abnormal development of peripheral lymphoid organs in mice
deficient in lymphotoxin. *Science* **264**, 703-707 (1994).
https://doi.org:10.1126/science.8171322

44 Mariathasan, S. *et al.* Absence of lymph nodes in lymphotoxin-alpha(LT alpha)-deficient
mice is due to abnormal organ development, not defective lymphocyte migration. *J*
Inflamm **45**, 72-78 (1995).
45 Davis, I. A. & Rouse, B. T. Immune responsiveness of lymphotoxin-alpha-deficient mice:
two reconstitution models. *Cell Immunol* **189**, 116-124 (1998).
https://doi.org:10.1006/cimm.1998.1371
46 Banks, T. A. *et al.* Lymphotoxin-alpha-deficient mice. Effects on secondary lymphoid
organ development and humoral immune responsiveness. *J Immunol* **155**, 1685-1693
(1995).
47 Meijer, E. F. J. *et al.* Lymph node effective vascular permeability and chemotherapy
uptake. *Microcirculation* **24** (2017). https://doi.org:10.1111/micc.12381
48 Suami, H. *et al.* A new indocyanine green fluorescence lymphography protocol for
diagnostic assessment of lower limb lymphoedema. *J Plast Reconstr Aesthet Surg*
(2022). https://doi.org:10.1016/j.bjps.2022.08.017
49 Maaskant-Braat, A. J. *et al.* Lymphatic mapping after previous breast surgery. *Breast* **21**,
444-448 (2012). https://doi.org:10.1016/j.breast.2011.10.007
50 Sato, A. *et al.* Altered lymphatic drainage patterns in re-operative sentinel lymph node
biopsy for ipsilateral breast tumor recurrence. *Radiat Oncol* **14**, 159 (2019).
https://doi.org:10.1186/s13014-019-1367-0
51 Bronte, V. & Pittet, M. J. The spleen in local and systemic regulation of immunity.
Immunity **39**, 806-818 (2013). https://doi.org:10.1016/j.immuni.2013.10.010
52 Tomura, M. *et al.* Monitoring cellular movement in vivo with photoconvertible
fluorescence protein "Kaede" transgenic mice. *Proc Natl Acad Sci U S A* **105**, 10871-
10876 (2008). https://doi.org:10.1073/pnas.0802278105
53 Helft, J. *et al.* GM-CSF Mouse Bone Marrow Cultures Comprise a Heterogeneous
Population of CD11c(+)MHCII(+) Macrophages and Dendritic Cells. *Immunity* **42**, 1197-
1211 (2015). https://doi.org:10.1016/j.immuni.2015.05.018
54 Wolchok, J. D. *et al.* Long-Term Outcomes With Nivolumab Plus Ipilimumab or
Nivolumab Alone Versus Ipilimumab in Patients With Advanced Melanoma. *J Clin Oncol*
40, 127-137 (2022). https://doi.org:10.1200/JCO.21.02229
55 Siddiqui, I. *et al.* Intratumoral Tcf1(+)PD-1(+)CD8(+) T Cells with Stem-like Properties
Promote Tumor Control in Response to Vaccination and Checkpoint Blockade
Immunotherapy. *Immunity* **50**, 195-211 e110 (2019).
https://doi.org:10.1016/j.immuni.2018.12.021
56 Paz-Ares, L. *et al.* Pembrolizumab plus Chemotherapy for Squamous Non-Small-Cell Lung
Cancer. *N Engl J Med* **379**, 2040-2051 (2018). https://doi.org:10.1056/NEJMoa1810865
57 Reck, M. *et al.* Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-Cell
Lung Cancer. *N Engl J Med* **375**, 1823-1833 (2016).
https://doi.org:10.1056/NEJMoa1606774
58 Swoboda, A. & Nanda, R. Immune Checkpoint Blockade for Breast Cancer. *Cancer Treat*
Res **173**, 155-165 (2018). https://doi.org:10.1007/978-3-319-70197-4_10
59 Weber, J. *et al.* Adjuvant Nivolumab versus Ipilimumab in Resected Stage III or IV
Melanoma. *N Engl J Med* **377**, 1824-1835 (2017).
https://doi.org:10.1056/NEJMoa1709030

Weber, J. S. *et al.* Adjuvant Therapy of Nivolumab Combined With Ipilimumab Versus Nivolumab Alone in Patients With Resected Stage IIIB-D or Stage IV Melanoma (CheckMate 915). *J Clin Oncol*, JCO2200533 (2022). https://doi.org:10.1200/JCO.22.00533

Patel, S. P. *et al.* Neoadjuvant-Adjuvant or Adjuvant-Only Pembrolizumab in Advanced Melanoma. *N Engl J Med* **388**, 813-823 (2023). https://doi.org:10.1056/NEJMoa2211437

Lo, J. A. *et al.* Epitope spreading toward wild-type melanocyte-lineage antigens rescues suboptimal immune checkpoint blockade responses. *Sci Transl Med* **13** (2021). https://doi.org:10.1126/scitranslmed.abd8636

Baeyens, A., Fang, V., Chen, C. & Schwab, S. R. Exit Strategies: S1P Signaling and T Cell Migration. *Trends Immunol* **36**, 778-787 (2015). https://doi.org:10.1016/j.it.2015.10.005

Cyster, J. G. & Schwab, S. R. Sphingosine-1-phosphate and lymphocyte egress from lymphoid organs. *Annu Rev Immunol* **30**, 69-94 (2012). https://doi.org:10.1146/annurev-immunol-020711-075011

Girard, J. P., Moussion, C. & Forster, R. HEVs, lymphatics and homeostatic immune cell trafficking in lymph nodes. *Nat Rev Immunol* **12**, 762-773 (2012). https://doi.org:10.1038/nri3298

Brinkmann, V., Cyster, J. G. & Hla, T. FTY720: sphingosine 1-phosphate receptor-1 in the control of lymphocyte egress and endothelial barrier function. *Am J Transplant* **4**, 1019-1025 (2004). https://doi.org:10.1111/j.1600-6143.2004.00476.x

Borsetto, D. *et al.* The risk of recurrence in surgically treated head and neck squamous cell carcinomas: a conditional probability approach. *Acta Oncol* **60**, 942-947 (2021). https://doi.org:10.1080/0284186X.2021.1925343

Darragh, L. B. *et al.* A phase I/Ib trial and biological correlate analysis of neoadjuvant SBRT with single-dose durvalumab in HPV-unrelated locally advanced HNSCC. *Nat Cancer* **3**, 1300-1317 (2022). https://doi.org:10.1038/s43018-022-00450-6

Saddawi-Konefka, R. *et al.* Lymphatic-preserving treatment sequencing with immune checkpoint inhibition unleashes cDC1-dependent antitumor immunity in HNSCC. *Nat Commun* **13**, 4298 (2022). https://doi.org:10.1038/s41467-022-31941-w

Ascierto, P. A. *et al.* Adjuvant nivolumab versus ipilimumab in resected stage IIIB-C and stage IV melanoma (CheckMate 238): 4-year results from a multicentre, double-blind, randomised, controlled, phase 3 trial. *Lancet Oncol* **21**, 1465-1477 (2020). https://doi.org:10.1016/S1470-2045(20)30494-0

Haley, P. J. The lymphoid system: a review of species differences. *J Toxicol Pathol* **30**, 111-123 (2017). https://doi.org:10.1293/tox.2016-0075

Fu, C. & Jiang, A. Dendritic Cells and CD8 T Cell Immunity in Tumor Microenvironment. *Front Immunol* **9**, 3059 (2018). https://doi.org:10.3389/fimmu.2018.03059

Bery, A. I. *et al.* Role of tertiary lymphoid organs in the regulation of immune responses in the periphery. *Cellular and molecular life sciences : CMLS* **79**, 359 (2022). https://doi.org:10.1007/s00018-022-04388-x

Wijeyesinghe, S. *et al.* Expansible residence decentralizes immune homeostasis. *Nature* **592**, 457-462 (2021). https://doi.org:10.1038/s41586-021-03351-3

75 Janneh, A. H. *et al.* Crosstalk between pro-survival sphingolipid metabolism and
complement signaling induces inflammasome-mediated tumor metastasis. *Cell Rep* **41**,
111742 (2022). https://doi.org:10.1016/j.celrep.2022.111742
76 Shi, W. *et al.* SphK2/S1P Promotes Metastasis of Triple-Negative Breast Cancer Through
the PAK1/LIMK1/Cofilin1 Signaling Pathway. *Front Mol Biosci* **8**, 598218 (2021).
https://doi.org:10.3389/fmolb.2021.598218
77 Perlman, R. L. Mouse models of human disease: An evolutionary perspective. *Evol Med*
Public Health **2016**, 170-176 (2016). https://doi.org:10.1093/emph/eow014
78 Heydon-White, A., Suami, H., Boyages, J., Koelmeyer, L. & Peebles, K. C. Assessing breast
lymphoedema following breast cancer treatment using indocyanine green
lymphography. *Breast Cancer Res Treat* **181**, 635-644 (2020).
https://doi.org:10.1007/s10549-020-05661-y
79 Onoi, K. *et al.* Immune Checkpoint Inhibitors for Lung Cancer Treatment: A Review. *J Clin*
Med **9** (2020). https://doi.org:10.3390/jcm9051362
80 Boukouris, A. E. *et al.* Latest evidence on immune checkpoint inhibitors in metastatic
colorectal cancer: A 2022 update. *Crit Rev Oncol Hematol* **173**, 103663 (2022).
https://doi.org:10.1016/j.critrevonc.2022.103663
81 Barreto de Albuquerque, J. *et al.* Microbial uptake in oral mucosa-draining lymph nodes
leads to rapid release of cytotoxic CD8(+) T cells lacking a gut-homing phenotype. *Sci*
Immunol **7**, eabf1861 (2022). https://doi.org:10.1126/sciimmunol.abf1861
82 Li, A. & Bergan, R. C. Clinical trial design: Past, present, and future in the context of big
data and precision medicine. *Cancer* **126**, 4838-4846 (2020).
https://doi.org:10.1002/cncr.33205
83 Jones, D., Pereira, E. R. & Padera, T. P. Growth and Immune Evasion of Lymph Node
Metastasis. *Front Oncol* **8**, 36 (2018). https://doi.org:10.3389/fonc.2018.00036
84 Rahim, M. K. *et al.* Dynamic CD8(+) T cell responses to cancer immunotherapy in human
regional lymph nodes are disrupted in metastatic lymph nodes. *Cell* **186**, 1127-1143
e1118 (2023). https://doi.org:10.1016/j.cell.2023.02.021
85 Wurdinger, T. *et al.* A secreted luciferase for ex vivo monitoring of in vivo processes. *Nat*
Methods **5**, 171-173 (2008). https://doi.org:10.1038/nmeth.1177
86 CoreTeam, R. R: A language and environment for statistical computing. *R Foundation for*
Statistical Computing, Vienna, Austria. (2021).
1129