Title:

Capture of circulating metastatic cancer cell clusters from a lung cancer patient can

reveal a unique genomic profile and potential anti-metastatic molecular targets: A proof

of concept study.

Kourosh Kouhmareh [1], Erika Martin [1], Darren Finlay [2], Anukriti Bhadada [3], Hector Hernandez-

Vargas [4], Francisco Downey [3], Jeffrey K. Allen [3] and Peter Teriete [5±]

1) PhenoVista Biosciences, 6195 Cornerstone Ct E STE 114, San Diego, CA 92121; 2) NCI Cancer Center Sanford

Burnham Prebys Medical Discovery Institute, 10901 N. Torrey Pines Rd., La Jolla, CA 92037; 3) TumorGen Inc.,

6197 Cornerstone Ct E STE #101, San Diego, CA 92121; 4) Centre Léon Bérard, 28 Prom. Léa et Napoléon

Bullukian, 69008 Lyon, France; 5) IDEAYA Biosciences, 7000 Shoreline Ct STE #350, South San Francisco, CA

94080; ±) corresponding author

Short title:

Metastatic cancer cell clusters reveal a unique genomic profile identifying anti-metastatic

targets.

Abstract

Metastasis remains the leading cause of cancer deaths worldwide and lung cancer, known for its highly metastatic progression, remains among the most lethal of malignancies. The heterogeneous genomic profile of lung cancer metastases is often unknown. Since different metastatic events can selectively spread to multiple organs, strongly suggests more studies are needed to understand and target these different pathways. Unfortunately, access to the primary driver of metastases, the metastatic cancer cell clusters (MCCCs), remains difficult and limited. These metastatic clusters have been shown to be 100-fold more tumorigenic than individual cancer cells. Capturing and characterizing MCCCs is a key limiting factor in efforts to help treat and ultimately prevent cancer metastasis. Elucidating differentially regulated biological pathways in MCCCs will help uncover new therapeutic drug targets to help combat cancer metastases. We demonstrate a novel, proof of principle technology, to capture MCCCs directly from patients' whole blood. Our platform can be readily tuned for different solid tumor types by combining a biomimicry-based margination effect coupled with immunoaffinity to isolate MCCCs. Adopting a selective capture approach based on overexpressed CD44 in MCCCs provides a methodology that preferentially isolates them from whole blood. Furthermore, we demonstrate a high capture efficiency of more than 90% when spiking MCCC-like model cell clusters into whole blood. Characterization of the captured MCCCs from lung cancer patients by immunofluorescence staining and genomic analyses, suggests highly differential morphologies and genomic profiles., This study lays the foundation to identify potential drug targets thus unlocking a new area of anti-metastatic therapeutics.

Introduction

Significance of metastatic cancer cell clusters

Metastasis remains the primary cause of cancer-related deaths. A recently published clinical analysis identifies metastasis as the cause of death in 50.2% to 90.4% of cases, depending on the origin of the primary tumor. The resulting high metastatic death rates indicate that increased focus on anti-metastatic therapy is urgently needed. Many cancer types have a demonstrated propensity to establish distant metastases among a defined set of organs. Non-small cell lung cancer (NSCLC), a major malignancy with high mortality, typically metastasizes to bone, followed by brain, liver and adrenal glands.[1] It is a prime example of mortality driven by distant metastases to many vital organs.[2, 3] However, the mechanism driving the spread, in particular cancer cell transportation and immune evasion is not well understood. The inability to isolate and profile circulating metastatic cells constitutes a major challenge in understanding these pathways.

Over the past decade, tremendous strides in developing better therapies via molecularly targeted and immune-checkpoint inhibitor treatments, have improved patient outcomes.[4] However, similar significant gains have not been observed for metastatic disease.[5] A key factor in this discrepancy is the complexity of cancer metastasis. Evidence has shown that signaling pathways driving metastatic lesions are often different than found within the primary tumor.[6] A recent publication from Anderson *et al.*, representing the Cancer Research UK and Australia Metastasis Working Group stated: "The standard cancer drug discovery and development pathway, including that for molecularly targeted and immunotherapies, generally ignores the ability of experimental medicines to inhibit metastasis." The authors went on to highlight: "To treat metastasis effectively, we must inhibit fundamental metastatic processes and

develop specific preclinical and clinical strategies that do not rely on primary tumor

responses."[7]

Extensive research into individual circulating tumor cells (CTCs), during this past decade, has

already begun to shed light on tumor proliferation but the underlying mechanism of metastasis

remains largely unknown.[8] Adding further complexity in understanding these mechanisms, the

majority of individual cancer cells in circulation undergo anoikis or are eliminated by the immune

system.[9] However, an important smaller subset of CTCs circulates in the bloodstream as a

cluster of two or more cells. Compelling evidence has demonstrated that CTC clusters have a

significantly higher tumorigenic potential once they enter circulation as opposed to individual

CTCs.[10-12]

Regarding circulating clusters or metastatic clusters, publications have referred to them with

different nomenclature such as: circulating tumor cell clusters (CTC clusters),[13] circulating

tumor micro-emboli (CTMs),[14] clustered circulating cancer stem cells (cCSCs),[15] metastatic

cancer cell clusters (MCCCs),[16] or by using a more general term called collective cell

migration,[17] yet all share common characteristics such as a being composed of 2-50 cells with

up to 100-fold higher metastatic potential than individual CTCs.[13, 18] [19] Here we will use the

terms MCCCs and CTC clusters interchangeably to refer to these circulating, highly metastatic

entities.

A number of recent studies have begun to elucidate the phenotypic composition of these

MCCCs and have shown them as either homotypic, composed of all cancer cells, or heterotypic,

containing multiple cell types including non-cancerous cells.[20-22] The MCCC phenotype

possesses an adaptive mechanism that enhances their survival in the harsh bloodstream

environment contributing to their metastatic potential.[23-26] Specifically, heterotypic MCCCs

demonstrate an increased ability to evade immune surveillance which is postulated to be a

result of cancer cell and immune cell communication within the cluster.[27-29] The metastatic

cancer cell clustering phenomenon is also thought to mitigate an immune response from natural

killer (NK) cells, due to surface receptors in addition to intercellular communication present

within CTC clusters.[30, 31]

New drug therapies, focused on MCCCs, will have the potential to dramatically reduce

metastasis and improve patient survival. A landmark paper published in 2019 by Aceto *et al.*

demonstrated that targeting MCCCs significantly reduces metastasis in an orthotopic breast

cancer model in mice.[32] Genomic analysis of breast cancer clusters from patients revealed 10

differentially regulated pathways unique to metastatic clusters.[32] Cardiac glycosides, an

existing class of drugs targeting one of those pathways, was evaluated in an *in vivo* animal

model. Treatment with one of the drugs (Ouabain) reduced the number of CTC clusters by 60%

and decreased metastasis to the lungs and other organs by 80-fold. An intriguing aspect from

this work is that the Ouabain dosage levels utilized are not cytotoxic and only disrupt the

MCCCs leading to a reduced metastatic index *in vivo.* This study strongly supports the

necessity to characterize the MCCCs from patients, to identify previously unseen therapeutic

targets and ultimately develop novel drugs thus opening a completely new paradigm in anti-

metastatic therapy.

Current Tumor Cell Isolation Methods from Whole Blood

Insights into how solid tumors migrate to specific pre-metastatic sites is necessary to develop

focused therapeutic approaches that can prevent metastasis. While the importance of MCCCs

has been well established, technologies to capture and characterize the MCCCs remain

problematic and has limited research into this field.

Existing technologies used to capture CTC clusters or MCCCs, rely on systems designed to

capture single CTCs. Many technologies are based on a size filtration mechanism for capture,

relying upon the premise that individual CTCs are larger than most leukocytes.[33, 34] Using

size filtration, patient blood samples are processed and then sorted based on the individual cell

diameter. Whole blood consists of three main components: platelets, red blood cells, and white

blood cells. White blood cells, which are the largest of the three components, are generally 12 -

15µm in size. The larger MCCCs, when present in whole blood, would be isolated and available

for further profiling. Although the main benefit of size-based technology is that its label-free and

requires little pre-processing, it has major limitations when it comes to successfully identifying

and isolating several types of individual CTCs and MCCCs. It is possible for smaller MCCC's

and many CTC's to be falsely binned as white blood cells, since the average cell diameters of

CTC's fall within the 10 - 15µm range.[35-37] A recent review also stated that microfluidic

channel size can induce critical flow velocities resulting in shearing and disaggregation of CTC

clusters.[18, 38] Another review by Amintas *et al.* concluded that individual CTC isolation

techniques are typically not compatible with CTC cluster isolation. They identified three key

requirements stating: "Future developments should aim towards methods that allow phenotypic,

molecular, and even functional analyses".[39]

New Metastatic Cancer Cell Cluster Isolation Methods

New approaches are needed to capture MCCC's. We have developed a highly selective capture

technology with a positive capture bias toward MCCCs by specifically exploiting the surface

molecular signature found on most solid tumor cancer cells. Our selective capture approach is

based on a biomimetic CD44-induced cell cluster margination effect combined with immobilized

capture antibodies coated within the inner walls of a microfluidic chip. The CD44 surface

expression is an abundant marker of MCCCs, upregulation of which closely correlates to their

metastatic potential.[40-44]

We are reporting on the success of our highly selective capture platform, focused on lung

cancer, that preferentially isolates cell clusters predisposed to metastasis. Our technology

allows routine and efficient capture of MCCCs providing insights into their biology as well as

elucidation of new anti-metastatic therapeutic targets. Ultimately, this will expedite the

development of new drugs capable of disrupting clusters and/or cluster formation.

Materials and Methods

Microfluidic Chip Manufacturing and Coating Procedure

Design, manufacturing, and coating of the microfluidic chips has been published previously.[45]

Production of the microfluidic chips used for this study were outsourced to a commercial

partner, µFluidix (Scarborough Ontario, Canada). Each chip was produced from two separate

layers of polydimethylsiloxane (PDMS) which were then precision aligned to create our open

channel microfluidic chips. The design parameters include 32 individual channels (100µm H x

200µm W each) along with a 'ceiling' feature composed on alternating patterns of recessed

chevrons, 50µm high which induces chaotic flow pattern.[45]

Ultrapure pharmaceutical grade alginate (Novamatrix; Sandvika, Norway) used for the hydrogel

substrate of the Smart-Coating™, was derivatized with streptavidin (AAT Bioquest, Cat.#16885)

molecules using carbodiimide chemistry via the carboxylic groups of the sugar residues.[46]

Coating of the microfluidic chips with a streptavidin derivatized, calcium (Ca^{+2}) cross-linked

alginate hydrogel was performed as described previously[45] A secondary coating consisting of

a mixture of biotinylated hyaluronic acid and monoclonal antibodies (mAbs) was added to the

alginate hydrogel. The secondary coating was prepared using cosmetic grade hyaluronic acid

(HA) powder (Resurrection Beauty; Cat.# High-MW-1800kDA) that was biotinylated and Cy5-

labeled (1:1:1 molar ratios) using carbodiimide chemistry via the carboxylic groups of the

glucuronic acid moiety using Biotin-Hydrazide, Thermo Scientific, Cat.# 21340; and Cy5-

Hydrazide from ApexBio, Cat.# A8145. The labeled HA plus a combination of biotinylated

monoclonal antibodies was mixed at a 1:16 ratio into phosphate buffered saline (PBS, 1X

Caisson Labs PBL01-500ML) along with 0.1% Tween-20.. The 4.5 mL mixture was infused onto

the alginate coated chip at 200 uL/min and the chip was subsequently washed with 2.5 mL of

PBS at 2.5 mL/min. After the second coating was complete chips were stored in a sealed

conical tube containing PBS at room temperature until needed. Figure S1 shows a successful

coating of our chip with the fluorescently labeled biotinylated hyaluronic acid-Cy5 (bHA-Cy5)

bound to the streptavidin derivatized, cross-linked alginate hydrogel.

Three surface epitopes, EGFR (ERBB1), MET and HER3 (ERBB3) were selected based on

published reports that they are typically expressed on the surface of lung cells.[47-49] Three

commercially available antibodies that bind to these epitopes were selected: biotinylated anti-

EGFR (Santa Cruz Biotechnology, Cat.# SC-120B) biotinylated anti-MET (Cell Signaling

Technology, Cat.#64526BC) and biotinylated anti-HER3 (LS Bio Cat.#LS-C87995). All

purchased antibodies were tested to confirm cell surface binding on established NSCLC cell

lines, HCC827 (ATCC, CRL-2868) and A549 (Angio-Proteomie cAP-0097GFP) by

immunofluorescence. Cell line authentication was based provided in vendor information. See

Supporting Information.

Experiments to assess the microfluidic chip capture efficiency were conducted with multi-cellular

co-cultured spheroids, generated using established cell lines mixed with the non-cancerous

human foreskin fibroblasts, HFF cells, as cancer associated fibroblast surrogates. A known

number of model NSCLC MCCCs using co-cultured spheroids (1:1 ratio of NSCLC HCC827 and

non-cancerous human foreskin fibroblasts, HFF cells) were spiked into normal donor whole

blood samples and processed in triplicate. Results comparing zero antibodies, (Smart-

Coating™ biotinylated hyaluronic acid / alginate hydrogel only) to 1 antibody (anti-EGFR) and 3

antibodies, (anti-EGFR, anti-MET and anti-HER3 combined) were performed. See the

Supporting Information for additional details.

NSCLC Patient Whole Blood Processing

After IRB approval, a total of eight patient samples (mean age male [3]: 70.9, ssd 1.1; female

[5]: 57.1, ssd 1.2) were collected from the University of California, San Diego (UCSD) Moores

Cancer Center, an NCI-Designated Comprehensive Cancer Center (UCSD Moores Cancer

Center, Project ID# INMC-044, IRB# 181755). Patients previously diagnosed with non-small cell

lung cancer (NSCLC), at stages III-A, III-B, III-C, IV-A, IV-B and agreeing to participate in a

study, graciously consented to provide whole blood samples. Patient information was compiled

for each blood sample, including age/gender, confirmed diagnosis, and cancer stage, at the

time of blood draw. Patient's sex and demographic information was not considered as inclusion

or exclusion criteria for this proof-of-concept study. Future studies involving a higher number of

patients will consider these factors to help understanding existing disparities. Sample

processing was blinded to patient information.

Collection tubes were treated with EDTA and Tirofiban to prevent coagulation and platelet

aggregation respectively. Samples were transported on the same day at ambient temperature

and processed through our microfluidic system within 8 hours. In an effort to preserve sample

integrity to the fullest extent, no red blood cell (RBC) lysis or whole blood pre-processing was

performed, since reports of CTC loses have been noted due to pre-treatment methods.[50, 51]

The entirety of the patient collected sample was aspirated into a 10mL syringe, attached to a

syringe pump and processed through our coated microfluidic chip at a flow rate of

100µL/minute. See Supplementary Methods for additional details.

Flow was not disturbed until the entire sample had been infused through the microfluidic chip.

After processing our patient blood sample, the chip was flushed with PBS at a rate of

100μL/minute for 5 minutes to remove any remaining unbound blood or white blood cells. A 4%

formaldehyde, methanol-free (PFA) solution was then pumped through at 100μL/minute

followed by another PBS wash for 5 minutes. Five of the patients' samples were processed for

immunofluorescence imaging and analysis. Cells were stained with a 10 minute flush of our

staining palette consisting of Hoechst 33342 live nuclear stain, (Thermo Scientific Cat.# H3570),

anti-CD44-FITC (Abeomic Cat.# 10-7516-F) and anti-EGFR-AF594 (Santa Cruz Biotechnology

Cat.# SC-120-AF594) all diluted in PBS, followed by a 5 minute PBS wash, all at the

100μL/minute flow rate.

Our processed chip was imaged across all flow channels using the Yokogawa CQ1 scanner by

stitching together 936 fields captured using a 10x objective. Chips were scanned for Brightfield,

Hoechst staining (405nm), CD44 (488nm), EGFR (594nm), and our bHA-Cy5 chip coating

(647nm). A relative increase of CD44 and EGFR fluorescent signal against background,

including well-defined nuclei morphology, were essential gating parameters used to identify

potential MCCCs. Using our 10X stitched composite image as a map for locating potential

MCCCs, we re-imaged at pre-selected points using a 40x objective to confirm identification.

Using a relatively strict inclusion criteria, any irregular nuclei illuminated via the Hoechst signal

even if coupled with positive CD44 and EGFR signal were rejected as MCCC candidates.

Furthermore, at least 3 individual nuclei had to be discernable in order to be classified as a

candidate MCCC.

NSCLC Whole Blood Processing for RNA-seq Analysis

Three additional patient LC-6, LC-7, LC-8 (selected at random, see Table 1) whole blood

samples were processed for RNA-sequencing analysis. Following whole blood infusion, a

modified staining protocol was used to reduce imaging time and enhance RNA stability. The

chips were flushed with Hoechst 33342 stain (without adding the anti-EGFR and anti-CD44

mAbs), at 100uL/min. for 15 minutes. The Hoechst stain was flushed out using a DMEMs-

phenol red media (Thermo Fisher, Cat.# 11965084) at 100uL/min. for 5 minutes..

The Inlet and Outlet ports were sealed and imaged using fluorescence microscopy. Brightfield

and 405nm images were collected to allow visualization of the captured MCCC nuclei. Post

imaging, RNA lysis / stabilization reagent (TaKaRa, Cat. # STO948) was flushed through the

microfluidic chip and incubated at zero flow for 15 minutes. Following lysis, fractions were

collected at the Outlet port every 15 seconds providing a total of 8 fractions, each with an

approximate volume of 25µL. Fractions were immediately frozen at -20°C and sent for RNA

sequencing analysis.

One of the patient samples, LC-6 was chosen for RNA-seq analysis. This sample corresponded

to a 64-year-old female patient, who was diagnosed by transthoracic biopsy with a TTF-1

positive Stage IV lung adenocarcinoma that had metastasized to the brain. Three RNA lysis

aliquots of the LC6 sample were processed and library preparation completed on the 3 aliquots

with SMART-Seq® HT Kit (Takara) and Illumina Nextera reagents for tagmentation. RNA

sequencing was performed at The Scripps Research Institute NGS core facility. The resulting

FASTQ files were downloaded and imported into R. All subsequent analyses were performed

using R/Bioconductor packages. A MultiQC v1.7 bioinformatics analysis was generated,

revealing the following QC summary data: Reads: 45.1M, % Alignment 94.7% and Phred

Quality score ranging from: 31.64 - 33.39.

Results

Capture Optimization

Our strategy was to develop a more 'selective' technology within a microfluidic system that would preferentially isolate and capture specific metastatic cell clusters rather than utilize a filtration-based or other size-based methodology. The premise was based on the extreme rarity of MCCCs among billions of red and white blood cells thus promoting the need to use an existing molecular biology process to enhance our cell selectivity. Leukocyte extravasation is a natural process involved with tissue damage, inflammation or infection.[52] Within the circulatory system, binding of CD44-presenting cells to endothelial hyaluronic acid (HA) is the first and essential step initiating margination, rolling, and extravasation.[53] Taking advantage of this existing biological principle, we designed and manufactured a microfluidic chip with a Smart-Coating™ technology that combines a tunable biomimetic and immuno-based dual capture approach.[45] An advantage of using our Smart-Coating™ is the ability to vary the number of monoclonal antibodies and biomimetic tethers which are integrated into the coating of the inner wall channels within the microfluidic chip. Immobilized streptavidin was derivatized on the cross-linked alginate hydrogel as previously described.[45] Antibodies with a biotin conjugate are easily bound to the streptavidin within the coating. Combinations of antibodies as well as biotinylated hyaluronic acid, which serves as a biomimetic tether, make up the second layer of our Smart-Coating. Since our platform is tunable, incorporating different biomimetic tethers and monoclonal antibodies enables the system to be adapted to other solid tumor cancer types.

Capture Antibody Selection for NSCLC

Identifying biomarkers on the surface of lung cells, that are relatively specific for lung tissue, is an important element of our platform. Multiple epitopes were utilized in our immuno-capture method, since this would likely isolate a larger diversity of MCCCs in circulation from different lung cancer patients. Experiments to assess the capture efficiency were conducted using multi-

cellular co-cultured spheroids which were counted and spiked into normal whole blood samples.

The co-cultured spheroids generated using established NSCLC cell lines resemble the MCCCs

found in patients' whole blood. As shown in Fig. 1, using 3 capture antibodies resulted in a 90%

capture efficiency, thus illustrating an advantage using multiple antibodies. Therefore 3 capture

antibodies were used for subsequent NSCLC patient samples. The ability to capture co-cultured

spheroids using the Smart-Coating alone, without any immobilized antibody, is likely due to the

CD44 expression found on the co-cultured spheroids that have an avidity for the hyaluronic acid

in our second coating.[54, 55]

Figure 1: Capture efficiency comparison using varying numbers of tissue specific

capture antibodies. The NSCLC co-cultured spheroids were spiked into freshly collected whole

blood samples from normal donors. The initial spike count varied for each sample run but

consistently fell within the range of 71-91 spheroids. Post processing, NSCLC co-cultured

spheroids were quantified via IF. Samples were processed in triplicate.

Patient Samples

A total of 8 patient samples were collected from the UCSD Moores Cancer Center and

processed through our Smart-Coated™ microfluidic chips that incorporated the 3 monoclonal

antibodies (mAbs) which demonstrated the highest capture efficiency. Two additional normal,

healthy donor, patient whole blood samples were processed from a 54-year-old female and 68-

year-old male, both of which revealed no confirmed MCCCs.

Five of the patient samples, following processing on the chip, included immuno-fluorescence

imaging for confirmation of the captured MCCCs using conjugated monoclonal antibodies as

listed in Fig. 2. Imaging revealed MCCCs were identified and confirmed in all 5 patient samples,

see Table 1.

Table 1: Basic patient information and the number of MCCCs confirmed.

Patient	Age/Gender	Diagnosis	Stage	# Clusters Detected	Detection Method
LC1	82 / M	lung adenocarcinoma	IVb	2	MCCC detection by bright field microscopy and confirmation by immunofluorescence staining for CD44, EGFR, Hoechst
LC2	53 / F	squamous cell lung carcinoma	IV	1	
LC3	59 / F	metastatic lung adenocarcinoma	IV	2	
LC4	65 / M	squamous cell lung carcinoma	IV	1	
LC5	67 / M	non-small cell lung carcinoma	IIIa	3	
LC6	64 / F	lung adenocarcinoma	IV	MCCC were not IF stained to preserve RNA quality	MCCC detection by bright field microscopy, samples processed for genomic analysis by RNA-seq.
LC7	46 / F	non-small cell lung cancer	IV		
LC8	66 / F	non-small cell lung cancer	IV		
Normal Blood Donors	54 / F	N/A	N/A	0	Neg. control
	68 / M			0	

Basic patient information including diagnosis and disease staging and detection method are

listed along with the number of confirmed MCCCs each consisting of a minimum of 3

morphologically distinct nuclei.

The rationale for the selected surface antigens to confirm MCCCs is that the overexpression of

both CD44 and EGFR is associated with cancerous lung cells.[15, 40, 56] Elevated fluorescent

signal of anti-CD44 mAb conjugate and anti-EGFR(528) mAb combined with at least 3 distinct

nuclei with well-defined nuclear morphology (Hoechst 33342) were essential gating parameters

used to identify potential MCCCs. Observations of the captured MCCCs appear to show

clusters with multiple nuclei that were relatively compact and not significantly larger than some

single cell, non-cancerous monocytes. See Fig. 2.[57, 58] The remaining three NSCLC patient

samples were processed with a modified protocol without immunofluorescence to help ensure

RNA stability was maintained for subsequent ultra-low concentration bulk RNA sequencing

analysis.

Figure 2: Representative 40X IF images of microfluidic chip captured NSCLC MCCCs

from the whole blood of two patients. Nuclei stained with Hoechst 33342 (blue) plus surface

antigens anti-CD44-FITC (488nm green) and anti-EGFR-AF594 (594nm red). The Smart-

Coating™ is visualized via HA-Cy5 (647nm purple). The overlapping of both CD44 and EGFR

fluorescence (yellow) in the composite image indicates cells expressing both markers. Scale bar

= 20μm for both LC2 and LC3 images.

Imaged cell clusters were binned into three groups:

1. Hoechst (+), EGFR (-) and CD44 (-)   Disregarded

2. Hoechst (+), CD44 (+) and EGFR (-)   Disregarded

3. Hoechst (+), CD44 (+) and EGFR (+)   Confirmed MCCC

Cells positive for only Hoechst and CD44 were presumed to be white bloods cells.[59] Any

clusters with green or red positive staining without clear nuclear morphology via Hoechst

staining were disregarded as potential MCCCs. Only clusters with all three stains, including well

defined nuclei were considered MCCCs. Examples of the confirmed NSCLC MCCCs captured

from whole blood are shown in Fig. 2. Note the heterogeneous shapes and sizes plus the multi-

cellular composition as seen by the variation in fluorescence signal distribution among the cells

in the MCCCs. A single MCCC was observed in patient LC2 which appears to have 5 nuclei.

Also note the apparent compact cellular morphology of the clusters. Two separate, multi-cellular

MCCCs were observed in patient LC3. The upper LC3 MCCC appears to have 3 separate

nuclei and the lower MCCC has 4 nuclei present.

MCCC Characterization via RNA-seq Analysis

Determining the unique gene expression profiles represented within the MCCCs via RNA-seq

analysis is crucial to learning more about the mechanisms of metastasis. Specifically, the ability

to characterize the metastatic cancer cell clusters, captured while in transit, directly from a

whole blood specimen will begin to shed light on the numerous signaling pathways involved in

the metastatic cascade. Three additional patient samples with confirmed NSCLC diagnoses

were processed through our microfluidic chips, using an abbreviated protocol, for the purpose of

identifying gene expression profiles represented by the MCCCs. Following a modified

processing method, MCCC capture was confirmed with Hoechst staining. After imaging, the

microfluidic chip was infused with RNA lysis/stabilization buffer, incubated and then aliquoted

fractions were collected at the outlet port.

Understanding the unique gene expression profile emanating from MCCCs found in NSCLC

patient's whole blood is essential to begin discovering new anti-metastatic targets for

therapeutic development. The 6[th] patient, designated LC6, is a 64-year-old female with TTF-1

positive Stage IV lung adenocarcinoma. A 10mL whole blood sample was processed using our

Smart-Coating™ microfluidic system. Post analysis imaging revealed multi-nucleated cell

clusters (each with 3 or more nuclei based on Hoechst staining). RNA-seq analysis was

performed on 3 separate lysate aliquots eluted from the chip processed from patient LC6.

Analysis revealed that 138 genes were highly expressed (over 1000 reads) among the MCCCs.

Bioinformatic analysis of the RNA-seq data was used to confirm that the MCCCs captured

originated from lung tissue. Expression levels were based on 6 well known genes found in lung

cells from previously published datasets of human lung tissue, GSM18949 and GSM18950.[60,

61] The boxplots shown in Fig.3 also included normalized gene expression levels found in lung

CTCs and lung primary tumor cells published elsewhere.[61] Our MCCC data was integrated

with the GSE74639 dataset. This RNA-seq dataset is comprised of lung tumor circulating single

cells, designated (SCs) in the original manuscript, but also referred to here as CTCs as well as

lung primary tumor cells (PTs).

A   **B**

Boxplot of lung-specific genes.

Figure 3: RNA-seq gene expression comparisons between lung circulating tumor cells,

primary tumor cells and MCCCs.

A) Normalized RNA-seq gene expression on a logarithmic y axis of 6 published genetic markers

correlated to human lung tissue. Also shown are the comparisons of the MCCCs (patient LC6)

to lung individual cells (CTCs) and lung primary tumor cells (PT) for those selected markers.

The plots from 5 of the 6 markers selected confirm the lung cell origin of the MCCCs. **B)** Boxplot

showing the normalized gene expression of *NKX2.1* (aka *TTF-1*) of the MCCCs compared to

both lung CTCs and lung primary tumor samples (PT). The detection of *TTF-1* from lung biopsy

specimens is typically used to aid in the diagnosis of NSCLC.

Expression of five out of the six genes assessed were found within the MCCCs from patient

LC6: *SFTPC*, *SCGB1A1*, *SLC34A2*, *SFTPB* and *CLDN18* thus confirming the tissue of origin

for the captured MCCCs was lung. Another gene was identified within the captured MCCCs,

TTF-1 (aka *NKX2.1*), which is commonly used to aid in the diagnosis of primary pulmonary

adenocarcinoma. The *NKX2.1* expression is found in 70-90% of non-mucinous adenocarcinoma

subtypes.[62] The boxplot shown in Fig. 3 reveals the expression of *TTF-1*(*NKX2.1*) in the

MCCCs from patient LC6, which is consistent with the patient diagnosis obtained from the

UCSD Moores Cancer Center, see Table 1. Data supports the finding that the MCCCs are from

lung tissue and are adenocarcinoma cells.

An important objective of the RNA-seq analysis of NSCLC patient LC6 was to determine if a

unique gene expression profile could be discerned for the captured MCCCs in comparison to

circulating lung CTCs and lung primary tumor cells. Additional statistical analysis of the

aliquoted samples revealed a unique profile of differentially expressed genes when compared to

other individual lung cancer circulating tumor cells (CTCs) and primary tumor (PT) lung biopsy

samples. Figure 4 is a Principal Component Analysis (PCA) of the integrated data sets

comparing lung CTCs, lung PTs and the 3 MCCC aliquoted samples (LC6). The PCA plot

shows a tight grouping of the LC6 aliquots that are significantly different than the PT and CTC

lung samples, supporting the postulation that a unique gene expression profile exists for the

MCCCs from NSCLC patient LC6.

Principal component analysis (PCA).

Figure 4: **Principal Component Analysis (PCA) of the integrated data sets**. PCA comparing

lung circulating tumor cells (CTC), lung primary tumor cells (PT) and the 3 MCCC aliquoted

samples (LC6). The PCA plot of the patient's MCCCs sequence data are grouped separately

compared to the other lung cancer cell types.

In order to better visualize a unique gene expression profile, based on differentially expressed

genes (DEGs) for the MCCCs, a heatmap was generated showing lung MCCCs relative to

individual lung CTCs and lung primary tumor cells. The MCCCs from patient LC6 display an

intermediate expression profile between lung CTCs and PTs, suggesting some common gene

expression among all 3 groups but also showing significant differences as shown in Fig. 5.

Gene expression mapping of MCCCs suggests a unique profile exists that can reveal pathways

driving cancer cell clusters migration to distant metastatic tumor microenvironments.

Figure 5: An RNA-seq heatmap was generated using differentially expressed gene

(DEGs) between PTs, CTCs and MCCCs. Although the MCCCs share some similarities

between lung primary tumor cells PTs and lung CTCs, the DEG grouping of the MCCCs

displays a unique profile.

Using additional bioinformatic data analysis, actionable targets were identified among the

MCCCs with corresponding therapeutics available or in development. (Fig.6) Based on the

sampling of 9 actionable targets, 3 genes from the MCCCs had noticeably higher expression

levels: *EGF*, *ALK* and *ERBB4* (aka *HER4*) compared with lung CTCs and primary tumor cells.

Additional research is necessary to demonstrate that inhibiting the identified targets will interfere

with metastatic cluster formation thus ultimately preventing distant metastasis.

A   **B**

Figure 6:   A sampling of actionable therapeutic targets

A) Box plots comparing normalized gene expression levels on a logarithmic y axis from a

sampling of genes among the MCCCs from patient LC6. The selected actionable genes may be

targets since existing drug interventions or new therapeutics are in development. Note the

increased expression of *EGF*, *ERBB4* (*HER4*) and *ALK* compared to lung CTCs and lung

primary tumor cells. **B)** An Illustration of several genes within the *Ras/Raf* pathway. Note the

relatively high expression of *BRAF* among the MCCCs suggesting a potential unseen

therapeutic target not seen in lung CTCs or primary tumor cells.

Another potential therapy strategy to disrupt MCCCs is to interfere with intercellular interaction

and communication. We reviewed the MCCC RNA-seq data to identify elevated expression

levels for adhesion and epithelial to mesenchymal (EMT) molecules which may be associated

with preparing metastatic clusters for intravasation. Two genes in particular, *COL4A1* and

FGFR1 showed higher expression levels than other lung cell types (See Fig. 7).

Figure 7: Gene expression correlated to intercellular adhesion and EMT.

Two box plots representing *COL4A1* and *FGFR1* normalized expression on a logarithmic scale.

MCCCs from patient LC6 appear to show relatively high expression levels compared to

individual circulating lung CTCs and lung primary tumor cells.

Discussion

Understanding the molecular signaling pathways of metastasis is essential for developing new

therapeutics designed to prevent cancer's spread. Isolating and characterizing the cancer cell

clusters which are more tumorigenic versus individual cancer cells is an important precursor for

new anti-metastatic drug development. [32, 63] Facilitating the isolation and characterization of

MCCCs will increase the number of patients analyzed for these metastatic clusters in real time,

as well as during various stages of their disease. Technology that allows for frequent sampling

at multiple time points may improve our understanding of the metastatic cascade during

cancer's progression. Additionally, easier access to MCCCs will increase research focus on the

mechanisms responsible for cancer cluster intravasation and ultimately aid the development of

therapeutics to stop their genesis.

Our proof of concept study utilized a completely novel dual capture approach based on a highly

selective, biomimetic-immunocapture 'Smart Coating' that fast tracks isolation and

characterization of extremely rare MCCCs. The Smart Coating™ technology provides a positive

selection bias for capturing extremely rare MCCCs directly from unprocessed whole blood.

Furthermore, we have shown that a differentially expressed gene profile (DEG) unique to the

MCCCs provides evidence that these specific vectors of metastasis require closer observation

to better understand the mechanisms responsible for distant metastatic lesions.

The RNA-seq heatmap, which illustrates a clear DEG profile, (Fig.5) further supports our proof-

of-concept study, demonstrating the ability to capture MCCCs in circulation and characterize

them to identify potentially relevant therapeutic targets. Also, identifying a unique DEG profile for

MCCCs supports the assertion that different signaling pathways are in effect during the

metastatic cascade.[8, 64]

We acknowledge that our study has limitations. Specifically, the number of patient samples

processed using fluorescently labeled mAbs to characterize the MCCCs and the corresponding

RNA-seq analyses was relatively low. However, the results demonstrating that one or more

MCCCs were found in all patient samples processed are highly promising. Our genomic

analysis of MCCCs provides evidence that our dual capture technology can be utilized to reveal

previously unseen metastatic pathways. Rather than being conclusive, our goal is to showcase

the potential of this type of information and prepare a roadmap for future MCCC studies.

The RNA-seq analyses from our captured MCCCs have shown the expression of several genes

that are considered actionable relative to targeted therapies in the treatment of cancer. An

example of a potential, previously unseen therapeutic target among the MCCCs is suggested by

the elevated *ERBB4* expression as seen in Fig. 6A. The *ERBB4* is a tyrosine kinase receptor

and a member of the epidermal growth factor receptor (EGFR) family, also known as HER4.

The ERBB4 receptor is activated via ligand binding which subsequently impacts MAPKs and

PI3K/AKT pathways.[65] Much research continues to elucidate the role of *ERBB4* mutations and

expression among many cancer types but it may be associated with proliferative and migratory

abilities of NSCLC.[66] While, *ERBB4* mutations and/or high expression levels observed among

NSCLC patients have been identified, their clinical significance remains unclear.[67, 68]

Elevated levels of ERBB4 among MCCCs, confirmed when additional patient samples are

processed, suggests a possible role in the metastatic cascade thus warranting additional study.

Another gene with high expression levels within the MCCCs relative to lung CTCs and primary

tumor is *BRAF* (See Fig.6B). Detection of mutated *BRAF* has been low as reported in only 2-7%

of advanced NSCLC patients.[69, 70] The *BRAF* gene is part of the MAPK/ERK signaling

pathway, which plays a crucial role in regulating cell growth, differentiation, and survival.

Mutated *BRAF* is a well-known driver of various cancers but not usually observed with lung

cancer.[71] However, a recent publication identified elevated wild type (WT) *BRAF* expression

levels at all stages (I-IV) from lung adenocarcinoma patients and was significantly associated

with decreased overall survival.[72] Identifying elevated *BRAF* expression among these rare

MCCCs reinforces the need to characterize the metastatic clusters to reveal previously unseen

therapeutic targets.

An important strategy for disrupting MCCCs (aka CTC clusters) was described by Nicola

Aceto's group where they identified molecules which interfered with intercellular adhesion.[32]

Results from an animal model study using patient derived xenografts demonstrated an 80-fold

reduction in metastatic index with the adhesion molecule inhibitor treatment group.

Two genes associated with EMT and adhesion, *COL4A1* and *FGFR1,* showed elevated gene

expression (see Fig. 7). The *COL4A1* (Collagen, type IV, alpha 1) is known to promote the

growth and metastasis of hepatocellular carcinoma cells.[73] Collagen, a major component of

the extracellular matrix (ECM), plays a significant role in cancer metastasis. Cancer cells can

alter the composition and organization of the ECM within the tumor microenvironment. Tumor

cells often produce enzymes such as matrix metalloproteinases (MMPs) that degrade and

remodel the ECM, allowing cancer cells to invade surrounding tissues more easily.[74] A recent

publication discusses targeting specific focal adhesion and ECM receptor pathways associated

with *COL4A1* for liver and lung metastases.[75]

Also shown is the upregulation of *FGFR1* (Fibroblast growth factor receptor 1) which has been

associated with the lung cancer progression pathway in some studies.[76] *FGFR1* is a receptor

tyrosine kinase also linked to the *MAPK/ERK* pathway and overexpression of this gene was

found to promote epithelial–mesenchymal transition (EMT) plus metastasis.[77] An interesting

note regarding elevated expression of *FGFR1* was its link to decreased efficacy of EGFR

tyrosine kinase inhibitor (TKI) treatments.[78]

Conclusion

Understanding the molecular signaling pathways of metastasis is essential for developing new

therapeutics designed to prevent cancers' spread. Isolating and characterizing the metastatic

cancer cell clusters, which are more tumorigenic versus individual cancer cells, directly from

cancer patients, is essential for finding unseen targets for new anti-metastatic drug

development. Expanding research efforts have highlighted the highly promising therapeutic

value in targeting and disrupting MCCCs directly in the fight against cancer metastasis.

Identification of *ERBB4*, *BRAF*, *COL4A1* and *FGFR1* as previously unseen therapeutic targets,

observed in highly tumorigenic MCCCs, may serve as focal points for new anti-metastatic drug

development. Although the gene expression data is based on replicates from a single patient

sample, the results demonstrate a proof of concept for our approach. The simplicity and utility of

our new platform has the potential to greatly expand the study of metastatic cancer cell clusters

for many different solid tumor cancers.

Author's Disclosures

Jeffrey K. Allen is the President/Founder of TumorGen and holds shares of the company. Peter

Teriete is a Co-Founder/Scientific Advisor of TumorGen and holds shares of the company.

Author's Contributions

Conceptualization   Peter Teriete, Jeffrey K. Allen

Formal Analysis   Hector Hernandez-Vargas

Funding Acquisition   Jeffrey K. Allen, Peter Teriete

Investigation   Kourosh Kouhmareh, Anukriti Bhadada, Francisco Downey

Methodology   Darren Finlay

Project Administration   Jeffrey K. Allen

Resources   Jeffrey K. Allen

Supervision   Erika Martin

Writing – Original Draft Prep.   Jeffrey K. Allen, Kourosh Kouhmareh

Writing – Review & Editing   Jeffrey K. Allen, Peter Teriete, Anukriti Bhadada

Acknowledgments

We are truly grateful for the participation of the patients that consented to provide blood

samples for our study. Also, we very much appreciate the support of Dr. Sandip Patel and Dr.

Greg Botta from the UCSD Moores Cancer Center. Additionally, thanks go out to Steve Head,

Director of the NGS Core facility at Scripps Research Translational Institute for aiding us with

the RNA-seq sample preparation and analysis. Further, we appreciate the support provided by

Dr. James Evans at PhenoVista Biosciences. The study was supported with an SBIR grant from

the National Cancer Institute, 1R43CA261362-01. https://reporter.nih.gov/project-

details/10255872

Note

Supporting Information

<u>References</u>

1.   Tamura T, Kurishima K, Nakazawa K, Kagohashi K, Ishikawa H, Satoh H, Hizawa N.

Specific organ metastases and survival in metastatic non-small-cell lung cancer.

Molecular and Clinical Oncology. 2015;3(1):217-21.

2.   Disibio G, French SW. Metastatic patterns of cancers: results from a large autopsy

study. Arch Pathol Lab Med. 2008;132(6):931-9.

3.   Riihimaki M, Hemminki A, Fallah M, Thomsen H, Sundquist K, Sundquist J, Hemminki K.

Metastatic sites and survival in lung cancer. Lung Cancer. 2014;86(1):78-84.

4.   Thallinger C, Füreder T, Preusser M, Heller G, Müllauer L, Höller C, et al. Review of

cancer treatment with immune checkpoint inhibitors : Current concepts, expectations,

limitations and pitfalls. Wien Klin Wochenschr. 2018;130(3-4):85-91.

5.   Ganesh K, Massagué J. Targeting metastatic cancer. Nat Med. 2021;27(1):34-44.

6.   Caswell DR, Swanton C. The role of tumour heterogeneity and clonal cooperativity in

metastasis, immune evasion and clinical outcome. BMC Med. 2017;15(1):133.

7.   Anderson RL, Balasas T, Callaghan J, Coombes RC, Evans J, Hall JA, et al. A

framework for the development of effective anti-metastatic agents. Nat Rev Clin Oncol.

2019;16(3):185-204.

8.   Riggi N, Aguet M, Stamenkovic I. Cancer Metastasis: A Reappraisal of Its Underlying

Mechanisms and Their Relevance to Treatment. Annu Rev Pathol. 2018;13:117-40.

9.   Wang WC, Zhang XF, Peng J, Li XF, Wang AL, Bie YQ, et al. Survival Mechanisms and

Influence Factors of Circulating Tumor Cells. Biomed Res Int. 2018;2018:6304701.

10.   Aceto N, Bardia A, Miyamoto DT, Donaldson MC, Wittner BS, Spencer JA, et al.

Circulating tumor cell clusters are oligoclonal precursors of breast cancer metastasis.

Cell. 2014;158(5):1110-22.

11.   Cheung KJ, Padmanaban V, Silvestri V, Schipper K, Cohen JD, Fairchild AN, et al.

Polyclonal breast cancer metastases arise from collective dissemination of keratin 14-

expressing tumor cell clusters. Proc Natl Acad Sci U S A. 2016;113(7):E854-63.

12.   Wrenn ED, Yamamoto A, Moore BM, Huang Y, McBirney M, Thomas AJ, et al.

Regulation of Collective Metastasis by Nanolumenal Signaling. Cell. 2020;183(2):395-

410.e19.

13.   Schuster E, Taftaf R, Reduzzi C, Albert MK, Romero-Calvo I, Liu H. Better together:

circulating tumor cell clustering in metastatic cancer. Trends Cancer. 2021;7(11):1020-

32.

14.   Wan S, Kim TH, Smith KJ, Delaney R, Park GS, Guo H, et al. New Labyrinth Microfluidic

Device Detects Circulating Tumor Cells Expressing Cancer Stem Cell Marker and

Circulating Tumor Microemboli in Hepatocellular Carcinoma. Sci Rep. 2019;9(1):18575.

15.   Liu X, Adorno-Cruz V, Chang YF, Jia Y, Kawaguchi M, Dashzeveg NK, et al. EGFR

inhibition blocks cancer stem cell clustering and lung metastasis of triple negative breast

cancer. Theranostics. 2021;11(13):6632-43.

16.   Xu K, Wang R, Xie H, Hu L, Wang C, Xu J, et al. Single-cell RNA sequencing reveals

cell heterogeneity and transcriptome profile of breast cancer lymph node metastasis.

Oncogenesis. 2021;10(10):66.

17.   Marchant CL, Malmi-Kakkada AN, Espina JA, Barriga EH. Cell clusters softening

triggers collective cell migration in vivo. Nat Mater. 2022;21(11):1314-23.

18.   Ring A, Nguyen-Sträuli BD, Wicki A, Aceto N. Biology, vulnerabilities and clinical

applications of circulating tumour cells. Nat Rev Cancer. 2023;23(2):95-111.

19.   Wrenn E, Huang Y, Cheung K. Collective metastasis: coordinating the multicellular

voyage. Clin Exp Metastasis. 2021;38(4):373-99.

20.   Piñeiro R, Martínez-Pena I, López-López R. Relevance of CTC Clusters in Breast

Cancer Metastasis. Adv Exp Med Biol. 2020;1220:93-115.

21.   Castro-Giner F, Aceto N. Tracking cancer progression: from circulating tumor cells to

metastasis. Genome Med. 2020;12(1):31.

22.   Herath S, Razavi Bazaz S, Monkman J, Ebrahimi Warkiani M, Richard D, O'Byrne K,

Kulasinghe A. Circulating tumor cell clusters: Insights into tumour dissemination and

metastasis. Expert Rev Mol Diagn. 2020;20(11):1139-47.

23.   Arnoletti JP, Fanaian N, Reza J, Sause R, Almodovar AJ, Srivastava M, et al. Pancreatic

and bile duct cancer circulating tumor cells (CTC) form immune-resistant multi-cell type

clusters in the portal venous circulation. Cancer Biol Ther. 2018;19(10):887-97.

24.   Jia D, Li X, Bocci F, Tripathi S, Deng Y, Jolly MK, et al. Quantifying Cancer Epithelial-

Mesenchymal Plasticity and its Association with Stemness and Immune Response. J

Clin Med. 2019;8(5).

25.   Uchôa Guimarães CT, Ferreira Martins NN, Cristina da Silva Oliveira K, Almeida CM,

Pinheiro TM, Gigek CO, et al. Liquid biopsy provides new insights into gastric cancer.

Oncotarget. 2018;9(19):15144-56.

26.   Zajac O, Raingeaud J, Libanje F, Lefebvre C, Sabino D, Martins I, et al. Tumour spheres

with inverted polarity drive the formation of peritoneal metastases in patients with

hypermethylated colorectal carcinomas. Nat Cell Biol. 2018;20(3):296-306.

27.   Sharma U, Medina-Saenz K, Miller PC, Troness B, Spartz A, Sandoval-Leon A, et al.

Heterotypic clustering of circulating tumor cells and circulating cancer-associated

fibroblasts facilitates breast cancer metastasis. Breast Cancer Res Treat.

2021;189(1):63-80.

28.   Szczerba BM, Castro-Giner F, Vetter M, Krol I, Gkountela S, Landin J, et al. Neutrophils

escort circulating tumour cells to enable cell cycle progression. Nature.

2019;566(7745):553-7.

29.   Heeke S, Mograbi B, Alix-Panabières C, Hofman P. Never Travel Alone: The Crosstalk

of Circulating Tumor Cells and the Blood Microenvironment. Cells. 2019;8(7).

30. Nakamura K, Smyth MJ. Immunoediting of cancer metastasis by NK cells. Nat Cancer.

2020;1(7):670-1.

31. Chan IS, Knútsdóttir H, Ramakrishnan G, Padmanaban V, Warrier M, Ramirez JC, et al.

Cancer cells educate natural killer cells to a metastasis-promoting cell state. J Cell Biol.

2020;219(9).

32. Gkountela S, Castro-Giner F, Szczerba BM, Vetter M, Landin J, Scherrer R, et al.

Circulating Tumor Cell Clustering Shapes DNA Methylation to Enable Metastasis

Seeding. Cell. 2019;176(1-2):98-112.e14.

33. Sharma S, Zhuang R, Long M, Pavlovic M, Kang Y, Ilyas A, Asghar W. Circulating tumor

cell isolation, culture, and downstream molecular analysis. Biotechnol Adv.

2018;36(4):1063-78.

34. Pugia M, Magbanua MJM, Park JW. Enrichment and Detection of Circulating Tumor

Cells and Other Rare Cell Populations by Microfluidic Filtration. Adv Exp Med Biol.

2017;994:119-31.

35. Kojić N, Milošević M, Petrović D, Isailović V, Sarioglu AF, Haber DA, et al. A

computational study of circulating large tumor cells traversing microvessels. Comput Biol

Med. 2015;63:187-95.

36. Takagi H, Dong L, Kuczler MD, Lombardo K, Hirai M, Amend SR, Pienta KJ. Analysis of

the Circulating Tumor Cell Capture Ability of a Slit Filter-Based Method in Comparison to

a Selection-Free Method in Multiple Cancer Types. Int J Mol Sci. 2020;21(23).

37. Bergman R, Afifi A, Heidger P. Atlas of Microscopic Anatomy. Michael P. D'Alessandro

MD, editor: WB Saunders; 1996.

38. Yamamoto A, Doak AE, Cheung KJ. Orchestration of Collective Migration and

Metastasis by Tumor Cell Clusters. Annu Rev Pathol. 2023;18:231-56.

39. Amintas S, Bedel A, Moreau-Gaudry F, Boutin J, Buscail L, Merlio JP, et al. Circulating

Tumor Cell Clusters: United We Stand Divided We Fall. Int J Mol Sci. 2020;21(7).

40. Liu X, Taftaf R, Kawaguchi M, Chang YF, Chen W, Entenberg D, et al. Homophilic CD44

Interactions Mediate Tumor Cell Aggregation and Polyclonal Metastasis in Patient-

Derived Breast Cancer Models. Cancer Discov. 2019;9(1):96-113.

41. Kawaguchi M, Dashzeveg N, Cao Y, Jia Y, Liu X, Shen Y, Liu H. Extracellular Domains I

and II of cell-surface glycoprotein CD44 mediate its trans-homophilic dimerization and

tumor cluster aggregation. J Biol Chem. 2020;295(9):2640-9.

42. San Juan BP, Garcia-Leon MJ, Rangel L, Goetz JG, Chaffer CL. The Complexities of

Metastasis. Cancers (Basel). 2019;11(10).

43. Kapeleris J, Zou H, Qi Y, Gu Y, Li J, Schoning J, et al. Cancer stemness contributes to

cluster formation of colon cancer cells and high metastatic potentials. Clin Exp

Pharmacol Physiol. 2020;47(5):838-47.

44. Rodrigues P, Vanharanta S. Circulating Tumor Cells: Come Together, Right Now, Over

Metastasis. Cancer Discov. 2019;9(1):22-4.

45. Shaner SW, Allen JK, Felderman M, Pasko ET, Wimer CD, Cosford NDP, et al. Design

and production of a novel microfluidic device for the capture and isolation of circulating

tumor cell clusters. AIP Advances. 2019;9(6):065313.

46. Lee KY, Mooney DJ. Alginate: properties and biomedical applications. Prog Polym Sci.

2012;37(1):106-26.

47. Giannopoulou E, Nikolakopoulos A, Kotsirilou D, Lampropoulou A, Raftopoulou S,

Papadimitriou E, et al. Epidermal growth factor receptor status and Notch inhibition in

non-small cell lung cancer cells. J Biomed Sci. 2015;22:98.

48. Salgia R. MET in Lung Cancer: Biomarker Selection Based on Scientific Rationale. Mol

Cancer Ther. 2017;16(4):555-65.

49. Scharpenseel H, Hanssen A, Loges S, Mohme M, Bernreuther C, Peine S, et al. EGFR

and HER3 expression in circulating tumor cells and tumor tissue from non-small cell lung

cancer patients. Sci Rep. 2019;9(1):7406.

50. Andree KC, Mentink A, Zeune LL, Terstappen L, Stoecklein NH, Neves RP, et al.

Toward a real liquid biopsy in metastatic breast and prostate cancer: Diagnostic

LeukApheresis increases CTC yields in a European prospective multicenter study

(CTCTrap). Int J Cancer. 2018;143(10):2584-91.

51. Joosse SA, Gorges TM, Pantel K. Biology, detection, and clinical implications of

circulating tumor cells. EMBO Mol Med. 2015;7(1):1-11.

52. Newton K, Dixit VM. Signaling in innate immunity and inflammation. Cold Spring Harb

Perspect Biol. 2012;4(3).

53. Zarbock A, Ley K, McEver RP, Hidalgo A. Leukocyte ligands for endothelial selectins:

specialized glycoconjugates that mediate rolling and signaling under flow. Blood.

2011;118(26):6743-51.

54. Christophis C, Taubert I, Meseck GR, Schubert M, Grunze M, Ho AD, Rosenhahn A.

Shear stress regulates adhesion and rolling of CD44+ leukemic and hematopoietic

progenitor cells on hyaluronan. Biophys J. 2011;101(3):585-93.

55. Hümmer J, Koc J, Rosenhahn A, Lee-Thedieck C. Microfluidic Shear Force Assay to

Determine Cell Adhesion Forces. Methods Mol Biol. 2019;2017:71-84.

56. Bethune G, Bethune D, Ridgway N, Xu Z. Epidermal growth factor receptor (EGFR) in

lung cancer: an overview and update. J Thorac Dis. 2010;2(1):48-51.

57. Au SH, Storey BD, Moore JC, Tang Q, Chen YL, Javaid S, et al. Clusters of circulating

tumor cells traverse capillary-sized vessels. Proc Natl Acad Sci U S A.

2016;113(18):4947-52.

58. Hao SJ, Wan Y, Xia YQ, Zou X, Zheng SY. Size-based separation methods of

circulating tumor cells. Adv Drug Deliv Rev. 2018;125:3-20.

59. Johnson P, Ruffell B. CD44 and its role in inflammation and inflammatory diseases.

Inflamm Allergy Drug Targets. 2009;8(3):208-20.

60.   Song Y, Ahn J, Suh Y, Davis ME, Lee K. Identification of novel tissue-specific genes by

767   analysis of microarray databases: a human and mouse model. PLoS One.

768   2013;8(5):e64483.

769   61.   Zheng Y, Miyamoto DT, Wittner BS, Sullivan JP, Aceto N, Jordan NV, et al. Expression

770   of β-globin by cancer cells promotes cell survival during blood-borne dissemination. Nat

771   Commun. 2017;8:14344.

772   62.   Moisés J, Navarro A, Santasusagna S, Viñolas N, Molins L, Ramirez J, et al. NKX2-1

773   expression as a prognostic marker in early-stage non-small-cell lung cancer. BMC Pulm

774   Med. 2017;17(1):197.

775   63.   Scheidmann MC, Castro-Giner F, Strittmatter K, Krol I, Paasinen-Sohns A, Scherrer R,

776   et al. An In Vivo CRISPR Screen Identifies Stepwise Genetic Dependencies of

777   Metastatic Progression. Cancer Res. 2022;82(4):681-94.

778   64.   Fabisiewicz A, Grzybowska E. CTC clusters in cancer progression and metastasis. Med

779   Oncol. 2017;34(1):12.

780   65.   Wang Z. ErbB Receptors and Cancer. Methods Mol Biol. 2017;1652:3-35.

781   66.   El-Gamal MI, Mewafi NH, Abdelmotteleb NE, Emara MA, Tarazi H, Sbenati RM, et al. A

782   Review of HER4 (ErbB4) Kinase, Its Impact on Cancer, and Its Inhibitors. Molecules.

783   2021;26(23).

784   67.   Tomizawa K, Suda K, Onozato R, Kuwano H, Yatabe Y, Mitsudomi T. Analysis of

785   ERBB4 mutations and expression in japanese patients with lung cancer. J Thorac Oncol.

786   2010;5(11):1859-61.

787   68.   Kurppa KJ, Denessiouk K, Johnson MS, Elenius K. Activating ERBB4 mutations in non-

788   small cell lung cancer. Oncogene. 2016;35(10):1283-91.

789   69.   Skoulidis F, Heymach JV. Co-occurring genomic alterations in non-small-cell lung

790   cancer biology and therapy. Nat Rev Cancer. 2019;19(9):495-509.

70.   Tissot C, Couraud S, Tanguy R, Bringuier PP, Girard N, Souquet PJ. Clinical

characteristics and outcome of patients with lung cancer harboring BRAF mutations.

Lung Cancer. 2016;91:23-8.

71.   Sumimoto H, Imabayashi F, Iwata T, Kawakami Y. The BRAF-MAPK signaling pathway

is essential for cancer-immune evasion in human melanoma cells. J Exp Med.

2006;203(7):1651-6.

72.   Dora D, Vörös I, Varga ZV, Takacs P, Teglasi V, Moldvay J, Lohinai Z. BRAF RNA is

prognostic and widely expressed in lung adenocarcinoma. Transl Lung Cancer Res.

2023;12(1):27-41.

73.   Wang T, Jin H, Hu J, Li X, Ruan H, Xu H, et al. COL4A1 promotes the growth and

metastasis of hepatocellular carcinoma cells by activating FAK-Src signaling. J Exp Clin

Cancer Res. 2020;39(1):148.

74.   Kleiner DE, Stetler-Stevenson WG. Matrix metalloproteinases and metastasis. Cancer

Chemother Pharmacol. 1999;43 Suppl:S42-51.

75.   DaSilva NA, Barlock BJ, Guha P, Ghosh CC, Trebino CE, Camberg JL, et al. Proteomic

signatures of myeloid derived suppressor cells from liver and lung metastases reveal

functional divergence and potential therapeutic targets. Cell Death Discov.

2021;7(1):232.

76.   Katoh M. FGFR inhibitors: Effects on cancer cells, tumor microenvironment and whole-

body homeostasis (Review). Int J Mol Med. 2016;38(1):3-15.

77.   Wang K, Ji W, Yu Y, Li Z, Niu X, Xia W, Lu S. FGFR1-ERK1/2-SOX2 axis promotes cell

proliferation, epithelial-mesenchymal transition, and metastasis in FGFR1-amplified lung

cancer. Oncogene. 2018;37(39):5340-54.

78.   Gammelgaard KR, Vad-Nielsen J, Clement MS, Weiss S, Daugaard TF, Dagnæs-

Hansen F, et al. Up-Regulated FGFR1 Expression as a Mediator of Intrinsic TKI

Resistance in EGFR-Mutated NSCLC. Transl Oncol. 2019;12(3):432-40.