Authors: Hélène Pidon[1,2,†*], Brigitte Ruge-Wehling[3,†], Torsten Will[4], Antje Habekuß[4], Neele Wendler[5],
Klaus Oldach[6], Anja Maasberg-Prelle[6], Viktor Korzun[5], Nils Stein[1,7]

Title: High-resolution mapping of *Ryd4^{Hb}*, a major resistance gene to *Barley yellow dwarf virus* from *Hordeum bulbosum*

Authors affiliations:
[1] Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Gatersleben, Seeland, Germany

[2] IPSiM, Univ Montpellier, CNRS, INRAE, Institut Agro, Montpellier, France

[3] Julius Kühn Institute (JKI) – Federal Research Centre for Cultivated Plants, Institute for Breeding Research
on Agricultural Crops, Sanitz, Germany

[4] Julius Kühn Institute (JKI) – Federal Research Centre for Cultivated Plants, Institute for Resistance
Research and Stress Tolerance, Quedlinburg, Germany

[5] KWS SAAT SE & Co. KGaA, Einbeck, Germany

[6] KWS LOCHOW GMBH, Bergen, Germany

[7] Center for Integrated Breeding Research (CiBreed), Georg-August University, Göttingen, Germany

[†]These authors contributed equally to the work.

*Corresponding authors: helene.pidon@inrae.fr, stein@ipk-gatersleben.de

Authors ORCID:
Pidon H.: 0000-0002-9802-1787
Will T.: 0000-0001-7985-4292
Habekuβ A.: 0000-0001-9627-0280
Oldach K.: 0000-0002-7114-2650
Korzun V.: 0000-0003-0760-8249
Stein N.: 0000-0003-3011-8731

Abstract
Virus diseases are causing high yield losses in crops worldwide. The *Barley yellow dwarf virus* (BYDV)
complex is responsible for one of the most widespread and economically important viral diseases of cereals.
While no complete resistance gene has been uncovered in the primary genepool of barley, sources of
resistance were identified in the wild relative *Hordeum bulbosum*, representing the secondary genepool of
barley. One such locus, *Ryd4^{Hb}*, has been previously introgressed into barley, and was allocated to
chromosome 3H, but is tightly linked to a sublethality factor that prevents the incorporation and utilization
of *Ryd4^{Hb}* in barley varieties. To solve this problem, we fine-mapped *Ryd4^{Hb}* and separated it from this
negative factor. We narrowed the *Ryd4^{Hb}* locus to a 66.5 kbp physical interval in the barley 'Morex'
reference genome. The region comprises one complete and one partial gene from the nucleotide-binding
and leucine-rich repeat immune receptor family, typical of dominant virus resistance genes. The closest
homolog to these two *Ryd4^{Hb}* candidate genes is the wheat *Sr35* stem rust resistance gene. In addition to the
fine mapping, we reduced the sublethality factor interval to 600 kbp in barley. Aphid feeding experiments
demonstrated that *Ryd4^{Hb}* provides a direct resistance to BYDV rather than a resistance to its vector. The
presented results, including the high-throughput molecular markers, will permit a more targeted selection
of the resistance in breeding, enabling the use of *Ryd4^{Hb}* in barley varieties.

Keywords

High-resolution mapping, *Barley yellow dwarf virus*, disease resistance, barley, *Hordeum bulbosum*, *Ryd4^{Hb}*

Declarations

Funding   This work was supported as part of the collaborative projects "Dwarfbulb" (grant 2814501910 from the German Federal Ministry of Food and Agriculture (BMEL)) and "BulbOmics" (grant 2818201615 from the German Federal Ministry of Food and Agriculture (BMEL)). The exome capture data production was part of the collaborative project "TRANSBULB" (grant 0315966 from the German Federal Ministry of Education and Research (BMBF)).

Conflicts of interest/Competing interests   Neele Wedler, Viktor Korzun, Klaus Oldach, and Anja Maasberg-Prelle are employed at KWS SAAT SE & Co and KWS LOCHOW. The other authors declare no conflict of interest.

Availability of data and material   The Exome capture sequencing datasets generated and/or analyzed in this study are deposited at EMBL-ENA under the project IDs PRJEB7909 and PRJEB65283.

Authors' contributions   Brigitte Ruge-Wehling, Nils Stein, Neele Wedler, and Viktor Korzun conceived the project and acquired the funding. Brigitte Ruge-Wehling, Neele Wedler, Klaus Oldach, Anja Maasberg-Prelle, and Viktor Korzun designed and built the mapping populations. Brigitte Ruge-Wehling and Kristin Fischer performed the initial linkage mapping and broke the linkage between the resistance and the sublethality loci. Brigitte Ruge-Wehling and Antje Habekuß performed the phenotyping experiments. Neele Wedler did the exome capture experiment. Torsten Will performed and analyzed the aphid feeding experiments. Hélène Pidon performed the high-resolution mapping and the pangenome analysis and drafted the manuscript. Nils Stein supervised the project. All authors provided critical feedback and helped shape the manuscript.

Acknowledgments

We gratefully acknowledge the excellent technical support by Manuela Kretschmann in DNA extraction and KASP genotyping, Dörte Grau in BaMMV resistance phenotyping, and Evelyn Betke, Katharina Stein and Kerstin Welzel in aphid rearing. We thank Anne Fiebig for raw data submission and Kristin Kutter for marker analysis.

Key message (<30 words)

We mapped *Ryd4^{Hb}* in a 66.5 kpb interval in barley and dissociated it from a sublethality factor. These results will enable a targeted selection of the resistance in barley breeding.

Introduction

Virus diseases cause significant yield losses and represent an increasing threat to agricultural crop production worldwide (Oerke, 2006). Among them, *Barley yellow dwarf virus* (BYDV) complex is responsible for one of the most widespread and economically important viral diseases of cereals. Transmitted in a persistent and circulative manner by several species of aphids, BYDV causes dwarfing and leaf discoloration, leading to significant yield loss in major cereal crops, in particular barley, wheat, maize, and oats (Ali et al., 2018). In recent years, it has become increasingly important in winter barley with an incidence that could reach 70% and yield loss of up to 80% (Beoni et al., 2016; Dedryver et al., 2010; Ordon et al., 2009). As climate change scenarios predict longer and warmer autumns, which favor aphid infestations of winter crop fields, BYDV could become one of the most threatening diseases of cereal crops (Roos et al., 2011; Trebicki, 2020). Reduction of yield losses by insecticide-based vector control is possible in principle, but undesirable for ecological reasons. To ensure sustainable barley cultivation in the expanding infestation areas and thus secure yields and quality in the long term, the cultivation of virus resistant varieties would provide the best solution.

So far, three genes and some QTLs have been described as providing partial resistance or tolerance to BYDV in barley. The gene *ryd1*, providing recessive intermediate tolerance was identified by Suneson (1955) but is still not cloned (Niks et al., 2004). Its effectiveness is low and it is rarely used in breeding. *Ryd2* was identified from an Ethiopian barley landrace (Schaller et al., 1964). It provides field tolerance to the virus serotypes BYDV-PAV, BYDV-MAV, and BYDV-SGV (Baltenberger et al., 1987). Mapped close to the centromere of chromosome 3H (Collins et al., 1996), *Ryd2* is used in several breeding lines (Kosova et al., 2008) where it can reduce significantly the yield loss caused by BYDV (Beoni et al., 2016). The third gene, *Ryd3* was also identified from an Ethiopian barley landrace (Niks et al., 2004). The gene was mapped in the centromeric region of chromosome 6H but, despite fine mapping on more than 3,000 F_2 plants (Lüpken et al., 2014), the mapping interval is still large. *Ryd3* has been transferred to commercial varieties where it provides a quantitative resistance, improved when in combination with *Ryd2* (Riedel et al., 2011). QTLs on chromosomes 1H, 2H, 4H, 5H, and 7H have been reported, however, providing only a limited level of tolerance (Toojinda et al., 2000; Riedel et al., 2011; Hu et al., 2019). No complete resistance to BYDV or its aphid vectors is known in barley, and broadening the genetic basis of resistance is therefore needed to ensure a durable and stable production of winter barley fields.

The secondary genepool of barley, consisting of the species *Hordeum bulbosum*, has not yet been used to improve resistance to the BYDV complex. Michel (1996) identified resistance to BYDV in the tetraploid (2n=4x=28) *Hordeum bulbosum* accession A17 (Bu10/2) from the Botanical Garden of Montevideo, Uruguay. Plants of this accession remained ELISA-negative for BYDV after several inoculations with aphids charged with the virus isolates BYDV-PAV1 Aschersleben, BYDV-MAV1 Aschersleben, and CYDV (*Cereal yellow dwarf virus*)-RPV Dittersbach (Habekuß et al., 2004). A17 was used as a parent in interspecific crosses and backcrosses with *H. vulgare* cv. Igri to generate an *H. bulbosum* introgression to barley. Its resistance was described as complete, dominant, and monogenic, and the locus, assigned to chromosome 3H, was named *Ryd4^{Hb}* (Scholz et al., 2009). Adversely, a recessive sublethality factor was cosegregating with *Ryd4^{Hb}* in the respective introgression. A study revealed low aphid feeding on the *H. bulbosum* A17 accession, suggesting that resistance may not be acting directly on the virus but rather indirectly against the aphid vector (Schliephake et al., 2013).

The present study reports the fine mapping of the *Ryd4^{Hb}* locus, the identification of candidate genes, and the description of aphids feeding behavior on susceptible and resistant introgression lines.

Material and methods

Plant material

BC_2F_5 and BC_2F_6 selected from BC_2F_4 plants from the Scholz et al. (2009) population were used for the low-resolution linkage mapping and development of an introgression line lacking the sublethality factor. This population is later named LM_Pop.

Two additional populations were generated to map $Ryd4^{Hb}$ at a higher resolution. FM_Pop1 was derived from a BC_2F_7 plant from LM_Pop crossed successively with three different barley elite varieties. The pedigree of the 15 lineages that constitute FM_Pop1 is presented in supplementary table 1. The donor of resistance in the FM_Pop2 was a BC_2F_8 derived from the BC_2F_6 JKI-5215 homozygous for the *H. vulgare* (Hv)-allele in the sublethality factor locus. As for FM_Pop1, it was crossed successively with two different barley elite varieties. The pedigree of the lines of four lineages that constitute FM_Pop2 is presented in supplementary table 2. The F_1 plants from the successive crosses were checked with markers to ensure the presence of the *H. bulbosum* (Hb)-allele at the $Ryd4^{Hb}$ locus and were further selfed to obtain the F_2 lineages forming FM_Pop1 and FM_Pop2.

Test of resistance to BYDV

Five to ten *Rhopalosiphum padi* aphids of the biotype R07, that were reared on BYDV-PAV1 Aschersleben (PAV1-ASL) infected plants as described by Kern et al. (2022), were placed on each one-week-old seedlings to be phenotyped in an air-conditioned greenhouse (20 °C, 16 h photoperiod, 10 klx). The aphids were killed after two days using the insecticide Confidor®WG 70 (Bayer CropScience AG, Germany). Further cultivation of the plants was carried out. Five to six weeks after inoculation, leaf samples of 50 mg from two leaves were taken and tested by a double-antibody sandwich-enzyme-linked immunosorbent assay (DAS-ELISA) according to Clark & Adams (1977) using custom-made antibodies. The viral content was evaluated by measuring extinction at 405 nm on a microtitre plate reader (Opsys MR, ThermoLabsystems or Tecan Sunrise, Tecan) one hour after the addition of the enzyme substrate. Based on negative controls, a extinction threshold was set in each experiment, usually at 0.1, under which a plant was classified as resistant. Phenotyping tests for gene mapping were performed on ten to 15 seeds of progenies of each genotype.

DNA extraction

Genomic DNA from the LM-Pop was either isolated following a slightly modified protocol after Stein et al. (2001) or using the Biosprint 96 DNA Plant Kit (Qiagen) and the Biosprint 96 working station (Qiagen) following the manufacturer's instructions. DNA was dissolved in TE buffer, quantified via photometric approaches (NanoQuant, Tecan, Austria) and diluted to a working concentration of 10 ng/μl. DNA extractions of plants from FM_Pop1 and FM_Pop2 were carried out according to the protocol described in Milner et al. (2019).

Marker development for low-resolution linkage mapping

The EST-derived SSR anchor markers *GBM1050* and *GBM1059* (Scholz et al., 2009; Stein et al., 2007; Thiel et al., 2003; Varshney et al., 2007) were kindly provided by Prof. Andreas Graner (Leibniz Institute of Plant Genetics and Crop Plant Research, Gatersleben). The STS markers *GLMWG883* and *GLABC161* were derived from the sequence of RFLP probe *MWG883* (Rostoks et al., 2005; Szűcs et al., 2009) and barley anchor marker *ABC161* (Close et al., 2009), respectively. Orthology of the interval on the rice chromosome 1 using the mapping of the anchor marker *ABC161* at 40.43 Mbp on said chromosome allowed

for the development of 18 tentative consensus (TC) markers polymorphic between Hv and Hb on
chromosome 3HL.

Additional polymorphisms between the *H. bulbosum* and *H. vulgare* genomes for marker development were
identified by RNASeq and Massive analysis of cDNA-ends (MACE). To this end, 1,000 plants from BC_2F_5
of LM_Pop were screened with the TC marker *TC173485*. Among them, 200 plants homozygous for the *H.
bulbosum* allele, considered resistant, and 200 plants homozygous for the *H. vulgare* allele, considered
susceptible, were selected. In total, 100 plants of each category were inoculated by aphids carrying the
isolate BYDV-PAV1, and an equal amount of plants were infested with control aphids without virus titer in
a separate climate chamber. 1h, 4h, 8h, and 24h after inoculation plant material of 25 genotypes per category
and treatment was harvested and sent to GenXPro (Frankfurt am Main, Germany) for RNA isolation and
sequencing. RNASeq and MACE were performed as described in Santos et al. (2018) and Braun et al.
(2019), respectively. In short, the raw data was cleaned off adapter sequences using the software TagDust
(Lassmann et al., 2009). All RNAseq datasets were combined to create a reference library. Assembly was
performed using the software Trinity (Grabherr et al., 2011). The reads of the individual libraries were
hereafter mapped to the reference library and single nucleotide polymorphisms (SNPs) were identified using
the software SNVMix (Goya et al., 2010). SNPs between the *H. bulbosum* and *H. vulgare* genomes were
identified and sequences 100 bps up- and downstream of each SNP were determined. Annotation of the
SNP-containing sequences was done by using the database Swiss-Prot (Boeckmann et al., 2003).

Additional polymorphisms were retrieved from exome capture sequencing, performed according to Wendler
et al. (2014) on the *H. bulbosum* parent A17 (Wendler et al., 2015) and the BC_2F_4 plant 5194/5, homozygous
for the 3HL-*H. bulbosum* introgression and mapped on the first barley genome assembly (International
Barley Genome Sequencing Consortium, 2012). Single nucleotide variants between *H. vulgare* and *H.
bulbosum* were called with samtools (SNP call score <200). Variants located within 200 bp of the end of a
reference sequence contig or supported by less than fivefold sequence read coverage were excluded from
further evaluation. The flanking sequences (50–60 bp) of variant positions were used for marker assay
development.

The primer design for PCR markers was carried out using Primer 3 (Untergasser et al., 2012). Conversion
of SNPs in CAPS markers was done by using SNP2CAPS (Thiel et al., 2004). All markers used for the low-
resolution linkage mapping are described in supplementary table 3.

Marker development for high-resolution linkage mapping
Exome capture data of the *H. bulbosum* A17 parent and of the BC_2F_4 plant 5194/5 was remapped onto the
barley reference genome MorexV3 (Mascher et al., 2021) together with the exome capture data of 13 barley
varieties from (Russell et al., 2016) (cultivars 'Barke', 'Bonus', 'Borwina', 'Bowman', 'Foma', 'Gull',
'Harrington', 'Haruna-Nijo', 'Igri', 'Kindred', 'Morex', 'Steptoe', and 'Vogelsanger Gold'). Reads
mapping and variant calling were performed as described in Milner et al., (2019). The SNP matrix was
filtered on the following criteria: heterozygous and homozygous calls had to have a minimum mapping
quality score of three and five, respectively, and be supported by a minimum of ten reads. SNP sites were
retained if they had less than 20% missing data and less than 20% heterozygous calls. SNPs that were within
the $Ryd4^{Hb}$ 20 Mbp interval defined by the low-resolution linkage mapping, homozygous for one allele in
all barley varieties and for the other allele in A17 and the BC_2F_4 introgression line, were retained.

For six SNPs, a 100 bp sequence containing the SNP in its center was provided to LGC genomics (Berlin,
Germany) for KASPar marker production (Supplementary table 4). Within the sublethality factor interval,

ten more SNPs were retrieved and sequences of 100 bp around each one were sent to 3CR Bioscience
(Welwyn Garden City, UK) for PACE assay design (Supplementary table 5). Primers were ordered from
Metabion (Germany) and mixed according to 3CR Bioscience (Welwyn Garden City, UK)
recommendations.

Thirteen CAPS markers (Supplementary table 6) were developed using NEBcutter (Vincze et al., 2003) to
identify the cutting enzyme and Primer 3 (Untergasser et al., 2012) to design the PCR primers.

Genotyping assays
For PCR of SSR, STS, and CAPS markers were carried out in a volume of 10µL containing 50-100ng of
DNA, 1X PCR buffer (Qiagen), 0.5 mM of each primer, 0.5 U of Taq DNA polymerase (Qiagen), and 0.2
mM of dNTPs. PCR amplification was carried out with an initial 10 min step at 95°C, followed by a
touchdown profile of ten cycles at 95°C for 30 seconds, 60°C for 30 seconds with a 0.5°C reduction per
cycle, and 72°C for 1 minute, followed by 35 cycles at 95°C for 30 seconds, 55°C for 30 seconds then 72°C
for 1 minute, and a last step of 7 minutes at 72°C. For CAPS markers, a 5 µl aliquot of the PCR product was
digested in 10 µL with 1 U of restriction enzyme and 1x of the appropriate digestion buffer at the temperature
recommended by the manufacturer. Pre- and post-digestion PCR products were separated on 2.5 % agarose
gels followed by ethidium bromide staining or in 10% polyacrylamide gels followed by silver nitrate
staining according to Budowle et al. (1991).

Detection of SNPs as genetic markers were performed by high-resolution melt analysis (HRM) by using the
Rotor Gene Technology (Qiagen). PCR was carried out in 20 µl volume containing 20 ng template DNA, 1
x buffer (Promega), 2.5 mM $MgCl_2$, 0.8 mM dNTP mix, 0.5 µM of each primer, 1 x EvaGreen Dye
(Biotium, Inc.) and 0.3 U *Taq* DNA polymerase (Promega). A touchdown PCR protocol was conducted
with a temperature gradient from 60–50°C. The melt curve analysis was conducted by ramping from 65 C
to 95 C with a 0.1 C decrease per capture.

Genotyping assays with KASPar and PACE markers were carried out as described in Pidon et al. (2020).

Genetic linkage analysis
220 LM_Pop was genotyped with 4 EST-derived SSR anchor markers, 18 TC-markers, 6 MACE-SNP, 13 STS-
221 markers derived from MACE (denoted as MACE_b), 9 markers derived from RNA-seq experiment, and 3
222 markers derived from exome capture. Linkage analysis was performed using the JoinMap® 4.1 software
223 (Van Ooijen, 2006). Genetic maps were displayed and edited in MapChart2.2 (Voorrips, 2002).

Pangenome comparison
The flanking markers and the *Ryd4^{Hb}* interval in the MorexV3 genome were searched on the 19 other
assemblies of the barley pangenome (Jayakodi et al., 2020) using BLAST+ (Camacho et al., 2009). The
resulting intervals were extracted and reannotated through a combination of alignments of the Morex
candidate genes to them, the search for the presence of conserved domains using NCBI *conserved domains*
(S. Lu et al., 2019), and NLRs annotation with NLR-Annotator (Steuernagel et al., 2020). The interval
structures were compared using Easyfig with blastn (Sullivan et al., 2011).

Aphids feeding experiment
In order to test if *Ryd4^{Hb}* provides resistance to aphids, a resistant and a susceptible progeny of the
heterozygous lines at *Ryd4Hb* locus EP_16-271_3_206 and EP_16-460_5_228 were selected. The
susceptible progenies EP_16-271_3_206_2 and EP_16-460_5_228_2 were carrying Hv alleles at both
Ryd4_CAPS19 and Ryd4_CAPS24, while the resistant lines EP_16-271_3_206_4 and EP_16-

460_5_228_6 were recombining in the interval and carrying a Hb allele at Ryd4_CAPS24 or
Ryd4_CAPS19, respectively. The feeding behavior of 12 to 16 adult apterouse non-viruliferous *R. padi*
(clone R07) of random age was observed on an individual healthy plant of the respective genotypes per
aphid by using the electrical penetration graph (EPG) technique (Tjallingii, 1978). Plants for EPG
experiments were reared in a greenhouse and were used at a 3-4 leaf stage where aphids were placed on the
lower side of the second leaf. Aphids were reared as described before (Kern et al., 2022). Aphids starved
for 1 hour before they were placed on the leaf. The recording started when all aphids were prepared. The
observation period was set to 8 hours and recording was started after all aphids were placed. For data
acquisition, the GIGA-8 EPG amplifier and EPG stylet software (EPG Systems, Wageningen, The
Netherlands) were used and data were analyzed with the EPG stylet analysis module. Waveforms were
annotated according to Tjallingii (1978) and Tjallingii & Esch (1993). Subsequently, selected parameters
were analyzed by using an Excel workbook (Alvarez et al., 2021). Recordings of aphids that fell from the
leaf or escaped during the experiment were not used.

Results

Low-resolution linkage mapping of *Ryd4^{Hb}*
From LM_Pop, 1,125 BC_2F_5 and BC_2F_6 were used for the low-resolution linkage mapping of *Ryd4^{Hb}*.
Phenotyping of progenies revealed 570 resistant, 276 susceptible plants, and 279 plants that died at early
stages. This high mortality rate was expected due to the recessively inherited sublethality factor very closely
linked to *Ryd4^{Hb}*. This locus prevents normal plant development, resulting in plant death (Figure 1). Because
of the close linkage between the two loci, we considered that all plants that died in the BC_2F_5 and BC_2F_6
families could be defined as homozygous resistant, while the resistant plants that survived would be
heterozygous at *Ryd4^{Hb}*. The phenotype distribution would indeed fit the expected 1:2:1 ratio of homozygous
resistant, heterozygous resistant, and homozygous susceptible genotypes for a dominant monogenic
inheritance of *Ryd4^{Hb}* resistance (Chi-Square Goodness of Fit Test: χ^2= 0.216, *p*=0.90), confirming the
previous observation for that locus on the BC_2F_4 generation (Scholz et al., 2009).

Fig. 1 Five months-old homozygous resistant plants either carrying a sublethality factor (left) or vital (right). The growth of plants homozygous at the sublethality factor is greatly reduced compared to plants carrying the resistance locus without the sublethality factor. They died before reaching heading.

The population was genotyped with 53 co-dominant markers and the linkage mapping was performed on the 1,014 individuals with a low amount of missing data. The introgression size was estimated to be 18.7 cM limited distally by the marker EXCAP_16 and proximally by the marker MACE_b_79 (Figure 2). The resistance locus $Ryd4^{Hb}$ is flanked by the MACE marker Mace_b_53 and the TC marker TC262452 with a genetic distance of 0.3 cM proximally and 0.5 cM distally, respectively. Moreover, the 200 BC_2F_5 plants screened with TC262452 and phenotyped by DAS-ELISA showed a perfect cosegregation of the TC262452 marker data with the resistance: the 100 plants carrying at least one Hb allele at the TC173485 marker had an extinction <0.1 whereas the 100 plants homozygous for the Hv allele had an extinction >0.5.

Fig. 2 Linkage map of chromosome 3HL carrying *Ryd4^Hb^*. The extent of the *Ryd4^Hb^* locus is represented in blue (1), marker bracket around *Ryd4^Hb^* displaying the flanking markers (2)

Development of an introgression line without sublethality factor

The exploitation of *Ryd4^Hb^* for breeding programs requires the selection of homozygous resistant and vital (by opposition with sublethal) plants. To select recombinants lacking the sublethality factor, 12,133 BC_2F_6 plants of LM_Pop were screened with markers TC262452 and Mace_b_53. Among those, 3,103 plants were homozygous for Hb-alleles, 6,020 heterozygous (Hb/Hv), and 3,010 homozygous for *H. vulgare* (Hv) at *Ryd4^Hb^* locus, confirming the 1:2:1 segregation (Chi-Square Goodness of Fit Test: χ^2= 2.14, *p*=0.34). The 3,103 plants homozygous for Hb-alleles were propagated and the progenies were checked for resistance. A single progeny, denoted as JKI-5215, was both completely resistant and vital. Marker analysis revealed that this population was segregating for Hb- and Hv-alleles distally from *Ryd4^Hb^*.

The JKI-5215 population, made of 43 BC$_2$F$_7$ plants from JKI-5215 progeny, was genotyped with all 16
markers that mapped distally from *Ryd4^{Hb}*. Markers TC262452 to RNASeq_b_1 were homozygous for Hb-
alleles whereas Mace_b_30 and all the remaining markers distally located were segregating in a 1:2:1
fashion (Figure 3). The recombination occurred within the initial 3HL introgression and resulted in a
reduced Hb-segment of 3.4 cM and all progenies are completely vital. A BC$_2$F$_8$ from the family JKI-5215
homozygous for the Hv-segment in the sublethality factor interval was selfed and used as the resistant donor
for the FM_Pop2.

Fig. 3 Characterization of 43 plants from the family JKI-5215 characterized by reduced introgressed Hb-
fragment without sublethality factor. Each vertical bar represents one offspring from JKI-5215. White,
black, and grey fragments represent Hb fragments, Hv fragments, and heterozygous genotypes at the
markers indicated on the right side, respectively. The light grey fragments at the dominant marker
Mace_b_69 are either Hv or heterozygous genotypes. The phenotype is indicated below the figure as 'nL'
for vital plants and 'L' for sublethal ones.

High-resolution mapping of *Ryd4^{Hb}* in two F$_2$ populations

To precisely map *Ryd4^{Hb}*, 5,589 F$_2$ plants from FM_Pop1 and 10,155 F$_2$ plants from FM_Pop2 were
genotyped with two KASPar markers to identify recombination at the locus. Plants from FM_Pop1 were
screened with Ryd4_KASP1 and Ryd4_KASP4, and plants from FM_Pop2 with Ryd4_KASP1 and
Ryd4_KASP5. Indeed, Ryd4_KASP4 is located in the sublethality factor interval and was segregating in
FM_Pop1, but not in FM_Pop2 where it was fixed Hv in its resistant donor. We identified 46 and 68
recombinant plants in FM_Pop1 and FM_Pop2, corresponding to 0.82% and 0.67% of recombination,
respectively. The positions of the SNPs of the markers Ryd4_KASP1, Ryd4_KASP4 and Ryd4_KASP5
were identified in the GBS data from 92 recombinant inbred lines of the cross 'Barke' x 'Morex' (Mascher
et al., 2013). A distance of 15 cM was observed between the markers Ryd4_KASP1 and Ryd4_KASP4, and
of 12.6 cM between Ryd4_KASP1 and Ryd4_KASP5, indicating by definition a 15% and 12.6% probability
of recombination in these intervals, respectively. The observed rate of recombination at the *Ryd4^{Hb}* locus is

therefore about 20-times lower than expected for the same genetic interval in a pure intraspecific barley
cross.

Recombinants were phenotyped on 15 offsprings and genotyped with the 13 CAPS markers. The resulting
interval for $Ryd4^{Hb}$ was 66.5 kbp-long in MorexV3 genome between the coordinates 592,685,940 and
592,752,329 flanked by CAPS19_2 and CAPS24, describing an interval comprising six recombination
events (Figure 4, Supplementary table 7). Four genes are annotated with high confidence on the MorexV3
genome in that interval: HORVU.MOREX.r3.3HG0318400, an S-formylglutathione hydrolase,
HORVU.MOREX.r3.3HG0318420, a partial nucleotide-binding and leucine-rich repeat immune receptors
(NLR) with a coiled-coil domain (CNL) lacking LRR domain which is likely a pseudogene,
HORVU.MOREX.r3.3HG0318450, a complete CNL, and HORVU.MOREX.r3.3HG0318470, an ankyrin-
repeat-containing gene. Those genes will later be referred to as *Ryd4_SFGH*, *Ryd4_pCNL1*, *Ryd4_CNL2*,
and *Ryd4_ANK*, respectively. The genes' homology revealed that locus $Ryd4^{Hb}$ is syntenic to the *Triticum*
monococcum locus *Sr35,* conferring resistance to wheat stem rust (Saintenac et al., 2013). The candidate
genes present a high similarity to one of the *Sr35* candidate genes. In particular, the translation of the
complete *Ryd4_CNL2* gene sequence from the MorexV3 reference genome shows 83% identity with the
SR35 protein while the respective coding sequence shows 88.7% nucleotide identity. The $Ryd4^{Hb}$ interval
also overlaps almost completely the ones of the *Rph13* leaf rust resistance gene from the *H. vulgare* ssp.
s*pontaneum* Hs2986 (Jost et al., 2020) and of the *Jmv2* resistance gene to the *Japanese soil-borne wheat*
mosaic virus from the barley cultivar 'Sukai Golden' (Okada et al., 2022). *Rph13* is located on chr3H
between the coordinates 592,658,337 and 592,786,929 on MovexV3 (128.6 kbp). Comparing the number
of recombinants, the size of the interval in which they occurred, and the size of the mapping population for
Rph13 and $Ryd4^{Hb}$ (4 recombinant in 128.6 kbp out of 719 plants and 6 recombinants plants in 66.5 kbp out
of 15,774, respectively), the recombination rate observed in the $Ryd4^{hb}$ populations is 7.5 times lower than
observed in the intraspecific cross used to map *Rph13*.

Fig. 4 High-resolution mapping of the *Ryd4*^{Hb} locus. (a) High-resolution mapping in the 5,589 F$_2$ plants from FM_Pop1 and 10,155 plants of FM_Pop2. The numbers in brackets show the sum of individual recombinants between the resistance locus and the corresponding marker in FM_Pop1 and FM_Pop2, respectively. (b) Marker saturation of the 18 recombinant plants at *Ryd4*^{Hb} locus. Recombinants from FM_Pop1 that died are excluded. (c) Candidate genes in the 66.5 kbp final *Ryd4*^{Hb} interval. (d) Comparison of *H. vulgare* 'Morex' and *T. monococcum* DV92 orthologous genes (only fragments with more than 80% nucleotide identity are shown).

Ryd4^{Hb} locus diversity in the barley pangenome

The orthologous intervals of the MorexV3 *Ryd4*^{Hb} region were retrieved from 19 additional diverse genome assemblies of the barley pangenome (Jayakodi et al., 2020) (supplementary table 8). We then annotated them using a combination of methods: mapping the 'Morex' genes, searching for NLR genes with NLR-Annotator (Steuernagel et al., 2020), and confirmation of the absence of additional conserved domains with NCBI conserved domains (S. Lu et al., 2019). The analysis revealed a very large divergence of the *Ryd4*^{Hb} interval in the different genotypes (Figure 5, supplementary figure 1). The shortest orthologous interval is the one of MorexV3. The largest is the one of the accession 'HOR 21599', 406 kbp-long and containing 10 NLRs, of which five are complete. The interval in the cultivars 'Akashinriki' and 'Du-Li Huang' is affected

by a large inversion of around 600 kbp. The observed diversity between haplotypes is mainly explained by
the presence of different repetitive elements and duplications. The degree of divergence in the *H. vulgare*
genepool in this interval suggests that an even greater diversity and divergence may be anticipated for the
corresponding region in the *H. bulbosum* genome.

Fig. 5 Graphical representation of the haplotype size and gene composition in the barley pangenome. Grey
horizontal bars represent the dimension of the interval; forward slashes represent the breakpoint due to the
large inversion in Akashinriki and Du-Li Huang. The interval between them is around 400 kbp-long and is
not homologous to 'Morex' interval. Arrows represent genes; yellow genes are S-formylglutathione
hydrolase, orange genes are partial NLRs, green genes are complete NLRs, and blue genes are ankyrin-
repeats-containing genes.

Mapping of the sublethality factor
To better understand why the *H. bulbosum* locus is causing the sublethality, we used recombinants identified
in the frame of *Ryd4^{Hb}* mapping to pinpoint the responsible factor more precisely. The low-resolution linkage
mapping located it distally from the marker *RNASeq_b_1*, which corresponds to position 594,019,595 on
chromosome 3H of the MorexV3 genome. To precise its interval, ten PACE markers were designed between
Ryd4_KASP18 and Ryd4_KASP22 (supplementary table 5), and used to genotype plants recombining in
the interval. Six recombination events were available: the one of JKI-5215 that we mapped using 24 non-
recombinant F$_2$ plants from lineage EP_16-460 of FM_Pop2, and the ones of five F$_2$ plants from FM_Pop1
recombining between Ryd4_KASP5 and Ryd4_C22, of which four vital plant and the sublethal plant
KW15_231/340_48 which died before heading (figure 5). The genotyping of 24 non-recombinant FM_Pop2
F$_2$ plants confirmed that the JKI-5215 recombination event occurred between Ryd4_KASP5 and
Ryd4_CAPS22 (Figure 5), and more precisely between markers Ryd4_leth2 and Ryd4_leth3 (594,034,042
to 594,290,776 bp). The genotyping and phenotyping of thirty-two F$_3$ plants from each of the four vital
plants recombining between Ryd4_KASP5 and Ryd4_C22 placed the sublethality factor proximally of
marker Ryd4_leth7. This was confirmed by the genotyping of the sublethal plant KW15_231/340_48 from
FM_Pop1 which was identified as recombinant between Ryd4_leth6 and Ryd4_leth7 (Figure 6). The
sublethality factor could therefore be assigned to a 483 kbp-interval between markers Ryd4_leth3 and
Ryd4_leth7 (594,290,776-594,773,972 bp on chromosome 3H of MorexV3). This interval is annotated with
15 high-confidence genes described in table 1. Among those genes, one or several could be essential genes
for plant development with no orthologs in the corresponding region of the *H. bulbosum* genome.

Fig. 6 Graphical representation of the genotype of lines recombining in the sublethality interval. Genotypes
of KW15-233/457_13, KW15-168/41_106, KW15-266/59_101, and KW15-231/344_20 are inferred from
those of 32 of their offsprings. The genotype of JKI-5215 is reconstructed from those of 24 F_2 plants of
FM_Pop2 from the lineage EP_16-460. Markers are depicted as vertical black lines and genotypes as
horizontal bars. White, black, and grey segments represent Hv, Hb, and heterozygotes genotypes,
respectively. Phenotypes are described on the right as sublethal (L), sublethality segregating in the progeny
(HL), and vital (nL). For better readability, marker positions are not to scale.

Gene name	Start	Stop	Annotation
HORVU.MOREX.r3.3HG0319200	594,288,404	594,295,050	B3 domain-containing protein
HORVU.MOREX.r3.3HG0319210	594,376,497	594,377,229	B3 domain-containing protein
HORVU.MOREX.r3.3HG0319220	594,404,110	594,405,348	Werner Syndrome-like exonuclease
HORVU.MOREX.r3.3HG0319240	594,551,247	594,553,562	B3 domain-containing protein
HORVU.MOREX.r3.3HG0319250	594,556,937	594,557,242	Ultraviolet-B-repressible protein
HORVU.MOREX.r3.3HG0319270	594,615,615	594,620,408	B3 domain-containing protein
HORVU.MOREX.r3.3HG0319280	594,622,773	594,626,954	Beta-1,3-glucanase
HORVU.MOREX.r3.3HG0319300	594,699,354	594,701,563	Beta-1,3-glucanase
HORVU.MOREX.r3.3HG0319310	594,712,025	594,712,888	F-box protein
HORVU.MOREX.r3.3HG0319320	594,721,495	594,725,844	RecA
HORVU.MOREX.r3.3HG0319330	594,731,099	594,738,322	Sentrin-specific protease
HORVU.MOREX.r3.3HG0319340	594,744,042	594,747,817	voltage-dependent L-type calcium channel subunit
HORVU.MOREX.r3.3HG0319350	594,748,418	594,749,859	Annexin
HORVU.MOREX.r3.3HG0319360	594,764,653	594,766,566	Amino acid permease
HORVU.MOREX.r3.3HG0319370	594,772,628	594,783,366	Acetyl-CoA carboxylase

387 **Table 1** High-confidence genes annotated in the sublethality interval on MorexV3 reference genome.
388 Coordinates refer to chromosome 3H.

Ryd4^{Hb} does not prevent aphid feeding

Resistance to insect-transmitted viruses can either be a direct resistance to the virus or an indirect one,
through resistance to the vector. To test if *Ryd4^{Hb}* provides resistance to its aphid vector, we monitored the
feeding of 12 to 16 aphids by EPG on five lines: two susceptible F_4 lines (EP_16-271_3_206_2 and
EP_16_460_5_228_2), their two resistant sister lines (EP_16-271_3_206_4 and EP_16_460_5_228_6) and
the susceptible barley cultivar 'Igri' which was the susceptible parent of LM-Pop and JKI-5215 (Figure 7a).
As none of the selected EPG parameters showed a normal distribution according to a Shapiro-Wilk test with
a p-value threshold of 0.05, we selected the Kruskal-Wallis test for multiple comparison. No significant
differences between the lines were observed for the selected parameters s_Np (χ^2=6.78, df=4, p=0.148),
s_C (χ^2=2.35, df=4, p=0.671), s_F (χ^2=4.64, df=4, p=0.327), s_G (χ^2=2.96, df=4, p=0.565), s_E1 (χ^2=1.35,
df=4, p=0.854), s_E2 (χ^2=3.16, df=4, p=0.534) and s_sE2 (χ^2=3.52, df=4, p=0.474) (Figure 7b,

supplementary table 9). The most divergent line was Igri, with an increased median duration for s_Np and decreased median durations for s_E2 and s_sE2, probably due to differences in the genetic background with the other lines.

Fig. 7 Aphid feeding behaviour on resistant and susceptible lines **a.** Graphical representation of the lines used to test the aphid feeding behaviour on resistant and susceptible lines. Loci are depicted as vertical black lines and genotypes as horizontal bars. White and black, segments represent Hv and Hb genotypes, respectively. **b.** Duration of the different feeding events on the five lines summed up by type. s_Np: total duration of all non-probing events. s_C: total duration of pathway phase. s_F: total duration of penetration problems. s_G: total duration of xylem drinking. s_E1: total duration of secretion of watery saliva. s_E2: total duration of phloem sap ingestion. s_sE2: total duration of sustained phloem sap ingestion. **c.** Duration ... Lines in the box plots indicate the median, and whiskers show the upper and lower 1,5xIQR (interquartile range) with dots indicating ouliers.

In addition to these general parameters, parameters associated with epidermis (t>1Pr) and mesophyll located (t>1E) but also sieve element located (t>1E2, t>1sE2) defense responses were inspected. Igri showed the

415 longest median durations for reaching the sieve elements (t>1E) and to reach ingestion (t>1E2 and t>1sE2).
416 However, none of the parameters t>1Pr (χ^2=3.83, df=4, p=0.43), t>1E (χ^2=6.32, df=4, p=0.176), t>1E2
417 (χ^2=3.43, df=4, p=0.489) and t>sE2 (χ^2=5.42, df=4, p=0.247) differs significantly between the tested lines
418 (Figure 7c, supplementary table 9).

Discussion

BYDV is a major threat to barley cultivation that is expected to increase in the following years, as autumns
become longer and warmer in Northern Europe (Roos et al., 2011; Trebicki, 2020), thus breeding for BYDV
resistance will become an even higher priority. So far only partial resistance has been discovered in the *H.*
vulgare primary genepool (Baltenberger et al., 1987; Collins et al., 1996; Lüpken et al., 2014; Niks et al.,
2004; Schaller et al., 1964; Suneson, 1955). Here, we report the high-resolution mapping of *Ryd4^Hb^*, the first
resistance gene to BYDV in barley, originating from the wild relative and secondary genepool species *H.*
bulbosum. We mapped the gene in a 66.5 kbp-interval of the MorexV3 barley reference genome despite the
linkage of the resistance locus to a sublethality factor and the reduced recombination between *H. vulgare*
and *H. bulbosum* genomes.

At the *Ryd4^Hb^* locus, four genes are annotated with high confidence on the MorexV3 genome, including two
genes from the CNL family, one pseudogene, and one likely to be functional. CNL genes are part of the
larger NLR family which are the most common class of resistance genes to biotic stress. They are coding
for intracellular proteins that form complexes (Wang et al., 2019) recognizing, directly or indirectly,
pathogen effector molecules and inducing local cell death in response. More than 30 NLR genes conferring
resistance to viruses have been cloned so far (Boualem et al., 2016; Sett et al., 2022) and more are candidates.
An *H. bulbosum* homolog of *Ryd4_pCNL1* or *Ryd4_CNL2* is therefore a very likely candidate for *Ryd4^Hb^*.
The *Ryd4^Hb^* locus is orthologous to the *Sr35* resistance locus from the wheat wild relative *Triticum*
monococcum (Saintenac et al., 2013). *Sr35* is coding for a CNL protein that shares 83% identity with the
translated *H. vulgare* sequence of *Ryd4_CNL2* and provides resistance to the fungal pathogen *Puccinia*
graminis f. sp. *tritici* causing wheat stem rust. This interval was also identified as candidate for the *Rph13*
resistance gene to leaf rust in the *H. vulgare* spp. *spontaneum* accession 'PI 531849' (Jost et al., 2020), and,
overlaps with the large interval of the *Jmv2* resistance gene to the *Japanese soil-borne wheat mosaic virus*
from the barley cultivar 'Sukai Golden' (Okada et al., 2022). Interestingly, the best homolog of *Sr35* in rice
is LOC_Os11g43700, which was identified as a resistance gene to the *Rice yellow mottle virus* in the African
rice species *Oryza glaberrima* (Pidon et al., 2017; Bonnamy et al., 2023). It is not rare that closely related
NLRs provide resistance to different classes of pathogens. A good example is the potato NLRs genes GPA2
and RX1 which provide resistance against the nematode *Globodera pallida* and potato virus X, respectively,
and share 88% of their amino acid sequence (Van Der Vossen et al., 2000). The comparison of the *Ryd4^Hb^*
interval in the barley pangenome (Jayakodi et al., 2020) demonstrated a very large diversity at this locus,
including NLR duplications. NLRs genes are indeed frequently under diversifying selection and tend to
evolve and duplicate by interallelic recombination between orthologs and by unequal crossing-over (Baggs
et al., 2017; Chen et al., 2010; Ding et al., 2007; Guo et al., 2011; Li et al., 2010; Michelmore & Meyers,
1998; Zhou et al., 2004). The *Sr35/Ryd4^Hb^* locus is one of those very diverse and dynamic loci that could be
described as R gene factories. Together with the homology with other resistance loci, this locus' NLR
diversity pleads for *Ryd4^Hb^* to be a CNL. However, in addition to the CNL genes, an *H. bulbosum* ortholog
of the *Ryd4_ANK* could also be a good candidate. Indeed, the structure of the encoded protein is close to the
one of *Arabidopsis* ACCELERATED CELL DEATH 6 (ACD6) protein. ACD6 confers enhanced resistance
to bacterial pathogens, including *Pseudomonas syringae*, by increasing the level of salicylic acid and

inducing spontaneous cell death (Rate et al., 1999; Dong, 2004; H. Lu et al., 2003, 2005). We also cannot
exclude that $Ryd4^{Hb}$ resistance is due to presence/absence variation of a gene in the primary and the
secondary genepool of barley, thus the resistance gene from *H. bulbosum* may have no ortholog in the *H.*
vulgare interval. Cloning of $Ryd4^{Hb}$ would therefore most likely require a *de novo* genome assembly of the
$Ryd4^{Hb}$ interval in a resistant genotype (introgression line of *H. vulgare* or resistance donor genotype of
H.bulbosum).

Resistance to insect-transmitted viruses, like the one provided by $Ryd4^{Hb}$, can either be a direct resistance
to the virus or a resistance to the vector, which would in effect prevent infection and therefore provide
indirect virus resistance. The melon NLR *Vat* resistance gene is the model of this indirect resistance. VAT
provides resistance to *Aphis gossypii* and to all the viruses it transmits tested so far, including the *Cucumber*
mosaic virus (Boissot et al., 2016). It recognizes an effector from *A. gossypii* and triggers the hypersensitive
response, stopping at the same time any viral infection that may have occurred. BYDV cannot be inoculated
to barley mechanically, so only resistance to the aphid vector was tested. A previous study showed that *R.*
padi aphids were feeding less and having a shorter salivation time on the $Ryd4^{Hb}$ *H. bulbosum* resistance
donor A17 compared to the BYDV-susceptible *H. bulbosum* line A21, suggesting that this could be the
reason for A17 BYDV resistance (Schliephake et al., 2013). However, our study showed no differences in
aphid feeding patterns on closely related resistant and susceptible lines, accompanied by an absence of
BYDV infection in the resistant lines, suggesting that the preliminary observation on the *H. bulbosum* donor
was probably due to A17 genetic background rather than $Ryd4^{Hb}$. $Ryd4^{Hb}$, therefore, provides direct
resistance to BYDV.

$Ryd4^{Hb}$ is a prime example of the importance of crop wild relatives serving as genetic resources and gene
donors in breeding schemes to achieve efficient and durable disease resistance. The advantage of using a
crop wild relative in prebreeding schemes as a unique source of resistance, however, comes at a cost:
reduced frequency of recombination and/or hybrid incompatibility leading to fertility or lethality problems.
In the case of the $Ryd4^{Hb}$ locus, recombination is 7.5 times less than in the intraspecific barley cross used to
map *Rph13*. To fine map the gene despite this handicap, we screened very large mapping populations with
high throughput genotyping technologies. At the $Ryd4^{Hb}$ locus, this negative linkage drag was strongly
materialized by a sublethality factor characterized by the reduced growth and early death of introgression
lines carrying the Hb allele at homozygous state at this locus. By screening a large number of plants for
recombination, we managed to break the linkage, producing a resistance donor without the sublethality
factor, that could be included in breeding schemes. We mapped the sublethality factor to a 600 kbp interval
on MorexV3 genome. The observed phenotype suggested that sublethal plants are possibly lacking an
essential gene, or a few genes, for development, therefore that one of the genes of the *H. vulgare* interval
has no ortholog in the donor *H. bulbosum* genome. Among the 15 genes annotated with high confidence in
the interval, B3 domain-containing proteins are part of a large transcription factor superfamily whose
members are playing key roles in various stages of plant development, from embryogenesis to seed
maturation (Swaminathan et al., 2008). F-box containing proteins are central part of the ubiquitin–26S
proteasome system and are thus key for different processes like phytohormone signaling, plant development,
cell cycle, or self-incompatibility (Stefanowicz et al., 2015). RecA proteins are maintaining DNA integrity
during meiosis by initiating double-strand break repair (Emmenecker et al., 2023). Annexins are widely
involved in regulating plant processes, from growth and development to responses to stresses (Wu et al.,
2022). One of the corresponding genes in the MorexV3 interval could be missing in the A17 haplotype and
explain the observed phenotype.

The results of this study would be helpful to breed barley varieties with an effective resistance to BYDV. We identified recombinants with a strongly reduced *H. bulbosum* fragment that can be used in breeding schemes, removing almost completely the negative linkage drag. The markers closely linked to the resistance can then be used in marker-assisted and genomic selection, postponing the tedious resistance evaluation to the last breeding step. Knowing that $Ryd4^{Hb}$ is a direct resistance gene to BYDV would also make it possible to establish the best strategy to avoid resistance breaking. Such a strategy could be pyramiding it with partial resistance or tolerance sources like *Ryd2* and *Ryd3*. Such resistant varieties would make a major contribution to sustainable barley cultivation.

Supplementary figure 1 Comparison of $Ryd4^{Hb}$ in *H. vulgare* pangenome modify from Easyfig 2.2.5 output. Only fragments with more than 80% nucleotide identity are shown.

References

Ali, M., Anwar, S., Shuja, M. N., Tripathi, R. K., & Singh, J. (2018). The genus Luteovirus from infection to disease. *European Journal of Plant Pathology*, *151*(4), 841-860. https://doi.org/10.1007/s10658-018-1425-8

Baggs, E., Dagdas, G., & Krasileva, K. (2017). NLR diversity, helpers and integrated domains : Making sense of the NLR IDentity. *Current Opinion in Plant Biology*, *38*, 59-67. https://doi.org/10.1016/j.pbi.2017.04.012

Baltenberger, D. E., Ohm, H. W., & Foster, J. E. (1987). Reactions of Oat, Barley, and Wheat to Infection with Barley Yellow Dwarf Virus Isolates1. *Crop Science*, *27*(2), cropsci1987.0011183X002700020010x. https://doi.org/10.2135/cropsci1987.0011183X002700020010x

Beoni, E., Chrpová, J., Jarošová, J., & Kundu, J. K. (2016). Survey of Barley yellow dwarf virus incidence in winter cereal crops, and assessment of wheat and barley resistance to the virus. *Crop and Pasture Science*, *67*(10), 1054-1063.

Boeckmann, B., Bairoch, A., Apweiler, R., Blatter, M.-C., Estreicher, A., Gasteiger, E., Martin, M. J., Michoud, K., O'Donovan, C., Phan, I., Pilbout, S., & Schneider, M. (2003). The SWISS-PROT protein knowledgebase and its supplement TrEMBL in 2003. *Nucleic Acids Research*, *31*(1), 365-370. https://doi.org/10.1093/nar/gkg095

Boissot, N., Thomas, S., Chovelon, V., & Lecoq, H. (2016). NBS-LRR-mediated resistance triggered by

aphids : Viruses do not adapt; aphids adapt via different mechanisms. *BMC Plant Biology*, *16*(1),

25. https://doi.org/10.1186/s12870-016-0708-5

Bonnamy, M., Pinel-Galzi, A., Gorgues, L., Chalvon, V., Hébrard, E., Chéron, S., Nguyen, T. H., Poulicard,

N., Sabot, F., Pidon, H., Champion, A., Césari, S., Kroj, T., & Albar, L. (2023). Rapid evolution of

an RNA virus to escape recognition by a rice nucleotide-binding and leucine-rich repeat domain

immune receptor. *New Phytologist*, *237*(3), 900-913. https://doi.org/10.1111/nph.18532

Boualem, A., Dogimont, C., & Bendahmane, A. (2016). The battle for survival between viruses and their

host plants. *Current Opinion in Virology*, *17*, 32-38. https://doi.org/10.1016/j.coviro.2015.12.001

Braun, E.-M., Tsvetkova, N., Rotter, B., Siekmann, D., Schwefel, K., Krezdorn, N., Plieske, J., Winter, P.,

Melz, G., Voylokov, A. V., & Hackauf, B. (2019). Gene Expression Profiling and Fine Mapping

Identifies a Gibberellin 2-Oxidase Gene Co-segregating With the Dominant Dwarfing Gene Ddw1

in Rye (Secale cereale L.). *Frontiers in Plant Science*, *10*.

https://www.frontiersin.org/articles/10.3389/fpls.2019.00857

Budowle, B., Chakraborty, R., Giusti, A. M., Eisenberg, A. J., & Allen, R. C. (1991). Analysis of the VNTR

locus D1S80 by the PCR followed by high-resolution PAGE. *American Journal of Human Genetics*,

48(1), 137-144.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009).

BLAST+ : Architecture and applications. *BMC Bioinformatics*, *10*(1), 421.

https://doi.org/10.1186/1471-2105-10-421

Chen, Q., Han, Z., Jiang, H., Tian, D., & Yang, S. (2010). Strong Positive Selection Drives Rapid

Diversification of R-Genes in Arabidopsis Relatives. *Journal of Molecular Evolution*, *70*(2),

137-148. https://doi.org/10.1007/s00239-009-9316-4

Clark, M. F., & Adams, A. N. (1977). Characteristics of the Microplate Method of Enzyme-Linked

Immunosorbent Assay for the Detection of Plant Viruses. *Journal of General Virology*, *34*(3),

475-483. https://doi.org/10.1099/0022-1317-34-3-475

Close, T. J., Bhat, P. R., Lonardi, S., Wu, Y., Rostoks, N., Ramsay, L., Druka, A., Stein, N., Svensson, J.

T., Wanamaker, S., Bozdag, S., Roose, M. L., Moscou, M. J., Chao, S., Varshney, R. K., Szűcs, P.,

Sato, K., Hayes, P. M., Matthews, D. E., … Waugh, R. (2009). Development and implementation

of high-throughput SNP genotyping in barley. *BMC Genomics*, *10*(1), 582.

https://doi.org/10.1186/1471-2164-10-582

Collins, N. C., Paltridge, N. G., Ford, C. M., & Symons, R. H. (1996). The Yd2 gene for barley yellow

dwarf virus resistance maps close to the centromere on the long arm of barley chromosome 3.

Theoretical and Applied Genetics, *92*(7), 858-864. https://doi.org/10.1007/BF00221898

Dedryver, C.-A., Le Ralec, A., & Fabre, F. (2010). The conflicting relationships between aphids and men :

A review of aphid damage and control strategies. *Comptes Rendus Biologies*, *333*(6), 539-553.

https://doi.org/10.1016/j.crvi.2010.03.009

Ding, J., Cheng, H., Jin, X., Araki, H., Yang, Y., & Tian, D. (2007). Contrasting patterns of evolution

between allelic groups at a single locus in Arabidopsis. *Genetica*, *129*(3), 235-242.

https://doi.org/10.1007/s10709-006-0002-9

Dong, X. (2004). The Role of Membrane-Bound Ankyrin-Repeat Protein ACD6 in Programmed Cell Death

and Plant Defense. *Science's STKE*, *2004*(221), pe6-pe6. https://doi.org/10.1126/stke.2212004pe6

Emmenecker, C., Mézard, C., & Kumar, R. (2023). Repair of DNA double-strand breaks in plant meiosis :

Role of eukaryotic RecA recombinases and their modulators. *Plant Reproduction*, *36*(1), 17-41.

https://doi.org/10.1007/s00497-022-00443-6

Goya, R., Sun, M. G. F., Morin, R. D., Leung, G., Ha, G., Wiegand, K. C., Senz, J., Crisan, A., Marra, M.

A., Hirst, M., Huntsman, D., Murphy, K. P., Aparicio, S., & Shah, S. P. (2010). SNVMix :

Predicting single nucleotide variants from next-generation sequencing of tumors. *Bioinformatics*,

26(6), 730-736. https://doi.org/10.1093/bioinformatics/btq040

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L.,

Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma,

F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., … Regev, A. (2011). Trinity : Reconstructing a

full-length transcriptome without a genome from RNA-Seq data. *Nature biotechnology*, *29*(7), 644-652. https://doi.org/10.1038/nbt.1883

Guo, Y.-L., Fitz, J., Schneeberger, K., Ossowski, S., Cao, J., & Weigel, D. (2011). Genome-Wide Comparison of Nucleotide-Binding Site-Leucine-Rich Repeat-Encoding Genes in Arabidopsis. *Plant Physiology*, *157*(2), 757-769. https://doi.org/10.1104/pp.111.181990

Habekuß, A., Schliephake, E., & Ehrig, F. (2004). Hordeum bulbosum—A source for BYDV resistance. *Proceedings of the 9th international barley genetics symposium*, 787-791.

Hu, H., Choudhury, S., Shabala, S., Gupta, S., & Zhou, M. (2019). Genomic regions on chromosome 5H containing a novel QTL conferring barley yellow dwarf virus-PAV (BYDV-PAV) tolerance in barley. *Scientific Reports*, *9*(1), Article 1. https://doi.org/10.1038/s41598-019-47820-2

International Barley Genome Sequencing Consortium. (2012). A physical, genetic and functional sequence assembly of the barley genome. *Nature*, *491*(7426), 711-716. https://doi.org/10.1038/nature11543

Jayakodi, M., Padmarasu, S., Haberer, G., Bonthala, V. S., Gundlach, H., Monat, C., Lux, T., Kamal, N., Lang, D., Himmelbach, A., Ens, J., Zhang, X.-Q., Angessa, T. T., Zhou, G., Tan, C., Hill, C., Wang, P., Schreiber, M., Boston, L. B., … Stein, N. (2020). The barley pan-genome reveals the hidden legacy of mutation breeding. *Nature*, *588*(7837), 284-289. https://doi.org/10.1038/s41586-020-2947-8

Jost, M., Singh, D., Lagudah, E., Park, R. F., & Dracatos, P. (2020). Fine mapping of leaf rust resistance gene Rph13 from wild barley. *Theoretical and Applied Genetics*, *133*(6), 1887-1895. https://doi.org/10.1007/s00122-020-03564-6

Kern, M., Meiners, T., Schliephake, · Edgar, Habekuss, A., Ordon, · Frank, & Will, T. (2022). Infection of susceptible/tolerant barley genotypes with Barley yellow dwarf virus alters the host plant preference of Rhopalosiphum padi clones depending upon their ability to transmit BYDV. *Journal of Pest Science*, *95*, 215-229. https://doi.org/10.1007/s10340-021-01367-2

Kosova, K., Chrpová, J., & Šíp, V. (2008). Recent advances in breeding of cereals for resistance to barley yellow dwarf virus. *Czech Journal of Genetics and Plant Breeding*, *44*(1), 1-10.

Lassmann, T., Hayashizaki, Y., & Daub, C. O. (2009). TagDust—A program to eliminate artifacts from
next generation sequencing data. *Bioinformatics*, *25*(21), 2839-2840.
https://doi.org/10.1093/bioinformatics/btp527

Li, J., Ding, J., Zhang, W., Zhang, Y., Tang, P., Chen, J.-Q., Tian, D., & Yang, S. (2010). Unique
evolutionary pattern of numbers of gramineous NBS–LRR genes. *Molecular Genetics and*
Genomics, *283*(5), 427-438. https://doi.org/10.1007/s00438-010-0527-6

Lu, H., Liu, Y., & Greenberg, J. T. (2005). Structure–function analysis of the plasma membrane- localized
Arabidopsis defense component ACD6. *The Plant Journal*, *44*(5), 798-809.
https://doi.org/10.1111/j.1365-313X.2005.02567.x

Lu, H., Rate, D. N., Song, J. T., & Greenberg, J. T. (2003). ACD6, a Novel Ankyrin Protein, Is a Regulator
and an Effector of Salicylic Acid Signaling in the Arabidopsis Defense Response. *The Plant Cell*,
15(10), 2408-2420. https://doi.org/10.1105/tpc.015412

Lu, S., Wang, J., Chitsaz, F., Derbyshire, M. K., Geer, R. C., Gonzales, N. R., Gwadz, M., Hurwitz, D. I.,
Marchler, G. H., Song, J. S., Thanki, N., Yamashita, R. A., Yang, M., Zhang, D., Zheng, C.,
Lanczycki, C. J., & Marchler-Bauer, A. (2019). CDD/SPARCLE: the conserved domain database
in 2020. *Nucleic Acids Research*, *48*(D1), D265-D268. https://doi.org/10.1093/nar/gkz991

Lüpken, T., Stein, N., Perovic, D., Habekuß, A., Serfling, A., Krämer, I., Hähnel, U., Steuernagel, B.,
Scholz, U., Ariyadasa, R., Martis, M., Mayer, K., Niks, R. E., Collins, N. C., Friedt, W., & Ordon,
F. (2014). High-resolution mapping of the barley Ryd3 locus controlling tolerance to BYDV.
Molecular Breeding, *33*(2), 477-488. https://doi.org/10.1007/s11032-013-9966-1

Mascher, M., Muehlbauer, G. J., Rokhsar, D. S., Chapman, J., Schmutz, J., Barry, K., Muñoz-Amatriaín,
M., Close, T. J., Wise, R. P., Schulman, A. H., Himmelbach, A., Mayer, K. F. X., Scholz, U.,
Poland, J. A., Stein, N., & Waugh, R. (2013). Anchoring and ordering NGS contig assemblies by
population sequencing (POPSEQ). *The Plant Journal*, *76*(4), 718-727.
https://doi.org/10.1111/tpj.12319

Mascher, M., Wicker, T., Jenkins, J., Plott, C., Lux, T., Koh, C. S., Ens, J., Gundlach, H., Boston, L. B.,
Tulpová, Z., Holden, S., Hernández-Pinzón, I., Scholz, U., Mayer, K. F. X., Spannagl, M., Pozniak,
C. J., Sharpe, A. G., Šimková, H., Moscou, M. J., … Stein, N. (2021). Long-read sequence
assembly : A technical evaluation in barley. *The Plant cell*, *accepted*.

Michel, M. (1996). *Untersuchungen zur Übertragung von Resistenzgenen aus der Wildart Hordeam*
bulbosum L. in die Kulturgerste Hordeum vulgare L. Diss., TU München, Munich, Germany.

Michelmore, R. W., & Meyers, B. C. (1998). Clusters of Resistance Genes in Plants Evolve by Divergent
Selection and a Birth-and-Death Process. *Genome Research*, *8*(11), 1113-1130.
https://doi.org/10.1101/gr.8.11.1113

Milner, S. G., Jost, M., Taketa, S., Mazón, E. R., Himmelbach, A., Oppermann, M., Weise, S., Knüpffer,
H., Basterrechea, M., König, P., Schüler, D., Sharma, R., Pasam, R. K., Rutten, T., Guo, G., Xu,
D., Zhang, J., Herren, G., Müller, T., … Stein, N. (2019). Genebank genomics highlights the
diversity of a global barley collection. *Nature Genetics*, *51*(2), 319-326.
https://doi.org/10.1038/s41588-018-0266-x

Niks, R. E., Habekuß, A., Bekele, B., & Ordon, F. (2004). A novel major gene on chromosome 6H for
resistance of barley against the barley yellow dwarf virus. *Theoretical and Applied Genetics*, *109*(7),
1536-1543. https://doi.org/10.1007/s00122-004-1777-7

Oerke, E.-C. (2006). Crop losses to pests. *The Journal of Agricultural Science*, *144*(1), 31-43.
https://doi.org/DOI: 10.1017/S0021859605005708

Okada, K., Tanaka, T., Fukuoka, S., Oono, Y., Mishina, K., Oikawa, T., Sato, K., Kato, T., Komatsuda, T.,
& Namai, K. (2022). Two dominant genes in barley (Hordeum vulgare L.) complementarily encode
perfect resistance to Japanese soil-borne wheat mosaic virus. *Breeding Science*, *72*(5), 372-382.
https://doi.org/10.1270/jsbbs.22046

Ordon, F., Habekuss, A., Kastirr, U., Rabenstein, F., & Kühne, T. (2009). Virus resistance in cereals :
Sources of resistance, genetics and breeding. *Journal of Phytopathology*, *157*(9), 535-545.
https://doi.org/10.1111/j.1439-0434.2009.01540.x

Pidon, H., Ghesquière, A., Chéron, S., Issaka, S., Hébrard, E., Sabot, F., Kolade, O., Silué, D., & Albar, L.

(2017). Fine mapping of RYMV3 : A new resistance gene to Rice yellow mottle virus from Oryza

glaberrima. *Theoretical and Applied Genetics*, *130*(4), 807-818. https://doi.org/10.1007/s00122-

017-2853-0

Pidon, H., Wendler, N., Habekuβ, A., Maasberg, A., Ruge-Wehling, B., Perovic, D., Ordon, F., & Stein, N.

(2020). High-resolution mapping of Rym14Hb, a wild relative resistance gene to barley yellow

mosaic disease. *Theoretical and Applied Genetics*. https://doi.org/10.1007/s00122-020-03733-7

Rate, D. N., Cuenca, J. V., Bowman, G. R., Guttman, D. S., & Greenberg, J. T. (1999). The Gain-of-

Function Arabidopsis acd6 Mutant Reveals Novel Regulation and Function of the Salicylic Acid

Signaling Pathway in Controlling Cell Death, Defenses, and Cell Growth. *The Plant Cell*, *11*(9),

1695-1708. https://doi.org/10.1105/tpc.11.9.1695

Riedel, C., Habekuß, A., Schliephake, E., Niks, R., Broer, I., & Ordon, F. (2011). Pyramiding of Ryd2 and

Ryd3 conferring tolerance to a German isolate of Barley yellow dwarf virus-PAV (BYDV-PAV-

ASL-1) leads to quantitative resistance against this isolate. *Theoretical and Applied Genetics*,

123(1), 69. https://doi.org/10.1007/s00122-011-1567-y

Roos, J., Hopkins, R., Kvarnheden, A., & Dixelius, C. (2011). The impact of global warming on plant

diseases and insect vectors in Sweden. *European Journal of Plant Pathology*, *129*(1), 9-19.

https://doi.org/10.1007/s10658-010-9692-z

Rostoks, N., Borevitz, J. O., Hedley, P. E., Russell, J., Mudie, S., Morris, J., Cardle, L., Marshall, D. F., &

Waugh, R. (2005). Single-feature polymorphism discovery in the barley transcriptome. *Genome*

Biology, *6*(6), R54. https://doi.org/10.1186/gb-2005-6-6-r54

Russell, J., Mascher, M., Dawson, I. K., Kyriakidis, S., Calixto, C., Freund, F., Bayer, M., Milne, I.,

Marshall-Griffiths, T., Heinen, S., Hofstad, A., Sharma, R., Himmelbach, A., Knauft, M.,

van Zonneveld, M., Brown, J. W. S., Schmid, K., Kilian, B., Muehlbauer, G. J., … Waugh, R.

(2016). Exome sequencing of geographically diverse barley landraces and wild relatives gives

insights into environmental adaptation. *Nature Genetics*, *48*(9), 1024-1030. https://doi.org/10.1038/ng.3612

Saintenac, C., Zhang, W., Salcedo, A., Rouse, M. N., Trick, H. N., Akhunov, E., & Dubcovsky, J. (2013). Identification of Wheat Gene Sr35 That Confers Resistance to Ug99 Stem Rust Race Group. *Science*, *341*(6147), 783-786. https://doi.org/10.1126/science.1239022

Santos, E., Benito, C., Silva-Navas, J., Gallego, F. J., Figueiras, A. M., Pinto-Carnide, O., & Matos, M. (2018). Characterization, genetic diversity, phylogenetic relationships, and expression of the aluminum tolerance MATE1 gene in Secale species. *Biologia Plantarum*, *62*(1), 109-120. https://doi.org/10.1007/s10535-017-0749-0

Schaller, C. W., Qualset, C. O., & Rutger, J. N. (1964). Inheritance and Linkage of the Yd2 Gene Conditioning Resistance to the Barley Yellow Dwarf Virus Disease in Barley1. *Crop Science*, *4*(5), cropsci1964.0011183X000400050034x. https://doi.org/10.2135/cropsci1964.0011183X000400050034x

Schliephake, E., Habekuss, A., Scholz, M., & Ordon, F. (2013). Barley yellow dwarf virus transmission and feeding behaviour of Rhopalosiphum padi on Hordeum bulbosum clones. *Entomologia Experimentalis et Applicata*, *146*(3), 347-356. https://doi.org/10.1111/eea.12033

Scholz, M., Ruge-Wehling, B., Habekuß, A., Schrader, O., Pendinen, G., Fischer, K., & Wehling, P. (2009). Ryd4Hb : A novel resistance gene introgressed from Hordeumbulbosum into barley and conferring complete and dominant resistance to the barley yellow dwarf virus. *Theoretical and Applied Genetics*, *119*(5), 837-849. https://doi.org/10.1007/s00122-009-1093-3

Sett, S., Prasad, A., & Prasad, M. (2022). Resistance genes on the verge of plant–virus interaction. *Trends in Plant Science*, *27*(12), 1242-1252. https://doi.org/10.1016/j.tplants.2022.07.003

Stefanowicz, K., Lannoo, N., & Van Damme, E. J. M. (2015). Plant F-box Proteins – Judges between Life and Death. *Critical Reviews in Plant Sciences*, *34*(6), 523-552. https://doi.org/10.1080/07352689.2015.1024566

Stein, N., Herren, G., & Keller, B. (2001). A new DNA extraction method for high-throughput marker

analysis in a large-genome species such as Triticum aestivum. *Plant Breeding*, *120*(4), 354-356.

https://doi.org/10.1046/j.1439-0523.2001.00615.x

Stein, N., Prasad, M., Scholz, U., Thiel, T., Zhang, H., Wolf, M., Kota, R., Varshney, R. K., Perovic, D.,

Grosse, I., & Graner, A. (2007). A 1,000-loci transcript map of the barley genome : New anchoring

points for integrative grass genomics. *Theoretical and Applied Genetics*, *114*(5), 823-839.

https://doi.org/10.1007/s00122-006-0480-2

Steuernagel, B., Witek, K., Krattinger, S. G., Ramirez-Gonzalez, R. H., Schoonbeek, H., Yu, G., Baggs, E.,

Witek, A. I., Yadav, I., Krasileva, K. V., Jones, J. D. G., Uauy, C., Keller, B., Ridout, C. J., & Wulff,

B. B. H. (2020). The NLR-Annotator Tool Enables Annotation of the Intracellular Immune

Receptor   Repertoire1   [OPEN].   *Plant*   *Physiology*,   *183*(2),   468-482.

https://doi.org/10.1104/pp.19.01273

Sullivan, M. J., Petty, N. K., & Beatson, S. A. (2011). Easyfig : A genome comparison visualizer.

Bioinformatics (Oxford, England), *27*(7), 1009-1010. https://doi.org/10.1093/bioinformatics/btr039

Suneson, C. A. (1955). Breeding for Resistance to Yellow Dwarf Virus in Barley. *Agronomy Journal*, *47*(6),

283-283. https://doi.org/10.2134/agronj1955.00021962004700060014x

Swaminathan, K., Peterson, K., & Jack, T. (2008). The plant B3 superfamily. *Trends in Plant Science*,

13(12), 647-655. https://doi.org/10.1016/j.tplants.2008.09.006

Szűcs, P., Blake, V. C., Bhat, P. R., Chao, S., Close, T. J., Cuesta-Marcos, A., Muehlbauer, G. J., Ramsay,

L., Waugh, R., & Hayes, P. M. (2009). An Integrated Resource for Barley Linkage Map and Malting

Quality   QTL   Alignment.   *The*   *Plant*   *Genome*,   *2*(2).

https://doi.org/10.3835/plantgenome2008.01.0005

Thiel, T., Kota, R., Grosse, I., Stein, N., & Graner, A. (2004). SNP2CAPS: a SNP and INDEL analysis tool

for   CAPS   marker   development.   *Nucleic*   *Acids*   *Research*,   *32*(1),   e5-e5.

https://doi.org/10.1093/nar/gnh006

Thiel, T., Michalek, W., Varshney, R., & Graner, A. (2003). Exploiting EST databases for the development

and characterization of gene-derived SSR-markers in barley (Hordeum vulgare L.). *Theoretical and*

Applied Genetics, *106*(3), 411-422. https://doi.org/10.1007/s00122-002-1031-0

Tjallingii, W. F. (1978). Electronic Recording of Penetration Behaviour by Aphids. *Entomologia*

Experimentalis et Applicata, *24*(3), 721-730. https://doi.org/10.1111/j.1570-7458.1978.tb02836.x

Tjallingii, W. F., & Esch, Th. H. (1993). Fine structure of aphid stylet routes in plant tissues in correlation

with EPG signals. *Physiological Entomology*, *18*(3), 317-328. https://doi.org/10.1111/j.1365-

3032.1993.tb00604.x

Toojinda, T., Broers, L. H., Chen, X. M., Hayes, P. M., Kleinhofs, A., Korte, J., Kudrna, D., Leung, H.,

Line, R. F., Powell, W., Ramsay, L., Vivar, H., & Waugh, R. (2000). Mapping quantitative and

qualitative disease resistance genes in a doubled haploid population of barley (Hordeum vulgare).

Theoretical and Applied Genetics, *101*(4), 580-589. https://doi.org/10.1007/s001220051519

Trebicki, P. (2020). Climate change and plant virus epidemiology. *Virus Research*, *286*, 198059.

https://doi.org/10.1016/j.virusres.2020.198059

Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B. C., Remm, M., & Rozen, S. G. (2012).

Primer3—New capabilities and interfaces. *Nucleic Acids Research*, *40*(15), e115.

https://doi.org/10.1093/nar/gks596

Van Der Vossen, E. A. G., Van Der Voort, J. N. A. M. R., Kanyuka, K., Bendahmane, A., Sandbrink, H.,

Baulcombe, D. C., Bakker, J., Stiekema, W. J., & Klein-Lankhorst, R. M. (2000). Homologues of

a single resistance-gene cluster in potato confer resistance to distinct pathogens : A virus and a

nematode. *The Plant Journal*, *23*(5), 567-576. https://doi.org/10.1046/j.1365-313x.2000.00814.x

Van Ooijen, J. W. (2006). JoinMap 4 : Software for the calculation of genetic linkage maps in experimental

populations of diploid species. *Wageningen, Netherlands: Plant Research International BV and*

Kayazma BV.

Varshney, R. K., Marcel, T. C., Ramsay, L., Russell, J., Röder, M. S., Stein, N., Waugh, R., Langridge, P.,

Niks, R. E., & Graner, A. (2007). A high density barley microsatellite consensus map with 775 SSR

761   loci. *Theoretical and Applied Genetics*, *114*(6), 1091-1103. https://doi.org/10.1007/s00122-007-

762   0503-7

763 Vincze, T., Posfai, J., & Roberts, R. J. (2003). NEBcutter : A program to cleave DNA with restriction

764   enzymes. *Nucleic Acids Research*, *31*(13), 3688-3691. https://doi.org/10.1093/nar/gkg526

765 Voorrips, R. E. (2002). MapChart : Software for the Graphical Presentation of Linkage Maps and QTLs.

766   *Journal of Heredity*, *93*(1), 77-78. https://doi.org/10.1093/jhered/93.1.77

767 Wang, J., Hu, M., Wang, J., Qi, J., Han, Z., Wang, G., Qi, Y., Wang, H.-W., Zhou, J.-M., & Chai, J. (2019).

768   Reconstitution and structure of a plant NLR resistosome conferring immunity. *Science*, *364*(6435),

769   eaav5870. https://doi.org/10.1126/science.aav5870

770 Wendler, N., Mascher, M., Himmelbach, A., Johnston, P., Pickering, R., & Stein, N. (2015). Bulbosum to

771   Go : A Toolbox to Utilize Hordeum vulgare/bulbosum Introgressions for Breeding and Beyond.

772   *Molecular Plant*, *8*(10), 1507-1519. https://doi.org/10.1016/j.molp.2015.05.004

773 Wendler, N., Mascher, M., Nöh, C., Himmelbach, A., Scholz, U., Ruge-Wehling, B., & Stein, N. (2014).

774   Unlocking the secondary gene-pool of barley with next-generation sequencing. *Plant Biotechnology*

775   *Journal*, *12*(8), 1122-1131. https://doi.org/10.1111/pbi.12219

776 Wu, X., Wang, Y., Bian, Y., Ren, Y., Xu, X., Zhou, F., & Ding, H. (2022). A critical review on plant

777   annexin : Structure, function, and mechanism. *Plant Physiology and Biochemistry*, *190*, 81-89.

778   https://doi.org/10.1016/j.plaphy.2022.08.019

779 Zhou, T., Wang, Y., Chen, J.-Q., Araki, H., Jing, Z., Jiang, K., Shen, J., & Tian, D. (2004). Genome-wide

780   identification of NBS genes in japonica rice reveals significant expansion of divergent non-TIR

781   NBS-LRR genes. *Molecular Genetics and Genomics*, *271*(4), 402-415.

782   https://doi.org/10.1007/s00438-004-0990-z