For *eLife*

The larva and adult of *Helicoverpa armigera* use differential gustatory receptors

to sense sugars

Shuai-Shuai Zhang[a, b, 1], Pei-Chao Wang[a, b, 1], Chao Ning[a, b], Ke Yang[a, b], Guo-Cheng

Li[a, b], Lin-Lin Cao[a, b], Ling-Qiao Huang[a], Chen-Zhu Wang[a, b, *]

[a] State Key Laboratory of Integrated Management of Pest Insects and Rodents,

Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, P.R. China

[b] Chinese Academy of Sciences Center for Excellence in Biotic Interactions,

University of Chinese Academy of Sciences, Beijing 100049, P. R. China

[1] These authors contributed equally to this work.

[*]Corresponding author

Correspondence to:

Chen-Zhu Wang

Institute of Zoology, Chinese Academy of Sciences, 1 Beichen West Road, Chaoyang

District, Beijing 100101, P. R. China

E-mail: czwang@ioz.ac.cn

Tel: +86-10-64807115

Fax: +86-10-64807099

Abstract

Almost all herbivorous insects feed on plants and use sucrose as a feeding stimulant,

but the molecular basis of their sucrose reception remains unclear. *Helicoverpa*

armigera as a notorious crop pest worldwide mainly feeds on reproductive organs of

many plant species in the larval stage, and its adult draws nectar. In this study, we

determined that the sucrose sensory neurons located in the contact chemosensilla of

larval maxillary galeae were 100–1000 times more sensitive to sucrose than those of

adult antennae, tarsi and proboscis. Using the *Xenopus* expression system, we

discovered that Gr10 highly expressed in the larval sensilla was specifically tuned to

sucrose, while Gr6 highly expressed in the adult sensilla responded to fucose, sucrose

and fructose. Moreover, using CRISPR/Cas9, we revealed that Gr10 was mainly used

by larvae to detect lower sucrose, while Gr6 was primarily used by adults to detect

higher sucrose and other saccharides, which results in differences in selectivity and

sensitivity between larval and adult sugar sensory neurons. Our results demonstrate

the sugar receptors in this moth are evolved to adapt towards the larval and adult

foods with different types and amounts of sugar, and fill in a gap in sweet taste of

animals.

Keywords: sucrose, gustatory receptor, cotton bollworm, CRISPR/Cas9, *Xenopus*

oocyte expression and two-electrode voltage-clamp

Introduction

Over 400,000 species of phytophagous insects live on approximately 300,000 species

of vascular plants worldwide (Schoonhoven et al. 2005). Carbohydrates produced by

plant photosynthesis are important nutrients for phytophagous insects. The

disaccharide sucrose, the major sugar in plants, is the principal transport form of

photoassimilates in higher plants. Sucrose varies considerably in concentration in

green leaves, fruits, and flower nectar where it occurs together with its constituent

monosaccharides fructose and glucose (Bernays and Chapman 1994). Sucrose,

fructose and glucose are the powerful phagostimulants for most phytophagous insects

studied (Bernays and Chapman 1994, Blaney and Simmonds 1988, Chapman 2003,

Schoonhoven et al. 2005).

The gustatory sensory neurons (GSNs) in one or more mouthpart sensilla of

phytophagous insects respond to sugars, usually including sucrose (Chapman 2003,

Dethier 1973). In Lepidoptera, caterpillars often have sugar sensitive GSNs in lateral

or medial sensilla styloconica in the maxillary galea (Glendinning et al. 2009, Martin

and Shields 2012, Schoonhoven and van Loon 2002), while adults also have sugar

sensitive GSNs in contact chemosensilla of the antennae (Jørgensen et al. 2007,

Popescu et al. 2013), tarsi (Schnuch and Seebauer 1998, Zhang et al. 2010), and

proboscis (Blaney and Simmonds 1988, Blaney and Simmonds 1990, Faucheux 2013).

The molecular basis of sugar reception in Lepidopteran insects remains unclear

although the sweet taste receptors in mammals and *Drosophila melanogaster* have

been extensively studied (Liman et al. 2014, Yarmolinsky et al. 2009). In mammals,

the combination of two taste receptors T1R2 and T1R3 recognizes sugars including

sucrose (Damak et al. 2003, Li et al. 2002, Nelson et al. 2001, Zhao et al. 2003). In

Drosophila, multiple gustatory receptors (GRs), including DmGr5a, DmGr61a, and

DmGr64a–f clusters, are required for the response of adults to multiple sugars

(Dahanukar et al. 2001, Dahanukar et al. 2007, Jiao et al. 2007, Jiao et al. 2008, Slone

et al. 2007); DmGr43a specifically responding to fructose is the major larval sugar

receptor, but it is only expressed in internal organs and some parts of the brain

(Mishra et al. 2013, Miyamoto et al. 2012). No external sugar-sensing mechanism in

Drosophila larvae has yet been characterized.

The cotton bollworm *Helicoverpa armigera* (Noctuidae, Lepidoptera) is a worldwide

agricultural pest, it causes over $3 billion in annual global economic loss (Pearce et al.

2017). The larvae of *H. armigera* mainly feeds on young leaves, flower buds, and

fruits of host plants; the adults mainly drink nectar of plants. Larval and adult diets

vary dramatically in the variety and concentration of sugars. Sucrose is the major

sugar in plant leaves and reproductive organs, whereas flower nectar mainly contains

sucrose, fructose and glucose (Nicolson 1998). Previous studies have shown that one

sugar sensitive GSN in *H. armigera* is located in each of two lateral sensilla

styloconica in the larva (Tang et al. 2000, Zhang et al. 2013), and many sugar

sensitive GSNs are distributed in the numerous contact chemosensilla of antennae

(Jiang et al. 2015), tarsi (Blaney and Simmonds 1990, Zhang et al. 2010), and

proboscis (Blaney and Simmonds 1988) in the adult.

To understand the molecular basis of sugar reception in these neurons, we first

compared electrophysiological and behavioral responses of *H. armigera* larvae and

adults. Second, we analyzed the expression patterns of nine candidate sugar GR genes

in larval and adult taste organs. Third, we carried out functional analysis of these GR

genes using the ectopic expression system, and revealed the response profiles of

Xenopus oocytes expressing Gr10, Gr6, and both of them. Finally, we individually

knocked out the two GR genes and obtained two homozygous mutants of *H. armigera*

by CRISPR/Cas9, and then detected changes in the electrophysiological and

behavioral responses of the mutant larvae and adults to sugars. Taking all the results

together, we demonstrate that larval and adult *H. armigera* used different GRs to

detect sucrose in food.

Results

Electrophysiological responses of contact chemosensilla to sugars

We first investigated the electrophysiological responses of the lateral and medial

sensilla styloconica in the larval maxillary galea to eight sugars. These sugars were

chosen because they are mostly found in host-plants of *H. armigera* or are

representative in the structure and source of sugars. The lateral sensilla styloconica

showed significant unicellular responses to 10 mM sucrose and fucose, with the firing

rates of 169 ± 8.57 and 43 ± 9.78 spikes/s, respectively (**Figure 1A, Figure 1—figure**

supplement 1A). The threshold concentrations for their responses to sucrose and

fucose were 0.1 mM and 10 mM, respectively (**Figure 1B, Figure 1—figure**

supplement 1B). The medial sensilla styloconica had a weak response to xylose

(**Figure 1—figure supplement 1C**). We also compared the responses of the lateral

sensilla styloconica to 0.05 mM sucrose, 5 mM fucose, and their mixture with the

same concentrations, and found that the mixture induced higher responses with only

one amplitude spikes and a firing rate of 71 ± 12.36 spikes/s, which was higher than

the firing rates induced by 0.05 mM sucrose (39 ± 6.37 spikes/s) and 5 mM fucose (36

± 8.87 spikes/s) (**Figure 1—figure supplement 1D**), confirming that the same cell

responded to sucrose and fucose. We conclude that there is only one GSN sensitive to

sucrose and fucose in each maxillary galea of larvae.

We also studied the responses of the contact chemosensilla in the tip of the antennae,

foreleg tarsi, and proboscis of female adults to eight sugar compounds at 10 mM and

the related dose responses (**Figure 1C–H**). The antennal sensilla responded to sucrose,

fucose, and fructose positively (**Figure 1C**), the tarsal sensilla only responded to

sucrose and fucose (**Figure 1E**), and the proboscis sensilla responded to sucrose,

fucose, and fructose (**Figure 1G**), but their firing rates were between 9-22 spikes/s.

All the evoked spike traces had one amplitude, indicating that they were single cell

responses (**Figure 1—figure supplement 1A**). The dose–response curves showed

that the threshold concentrations of sucrose, fucose and fructose were all 10 mM for

the antennal sensilla (**Figure 1D, Figure 1—figure supplement 1B**), 100 mM, 10

mM and 100 mM for the tarsal sensilla (**Figure 1F**, **Figure 1—figure supplement**

1B), and 10 mM, 100 mM and 10 mM for the proboscis sensilla, respectively (**Figure**

1H, **Figure 1—figure supplement 1B**).

In short, both larvae and adults of *H. armigera* had sugar GSNs, but their response

profiles, intensity and sensitivity differed distinctively. The sugar GSNs in lateral

sensilla styloconica of larvae responded strongly to sucrose with high sensitivity, and

weakly to fucose; while the sugar GSNs of adults responded to sucrose, fucose, and

fructose (except for those on tarsi) with lower intensity and sensitivity. The GSNs of

larvae were 100–1000 times more sensitive to sucrose than those of adults.

Behavioral responses of larvae and adults to sugars

To determine how the taste inputs of various sugars cause behavioral responses in

insects, the two-choice test was used to examine the feeding behavior of the 5[th] instar

larvae of *H. armigera* to eight sugars at 10 mM. Sucrose and fructose stimulated

larval feeding, and their threshold concentrations were 10 mM (**Figure 2A–B, Figure**

2—figure supplement 1); other tested sugars had no effect on larval feeding (**Figure**

2A).

We also studied the proboscis extension reflex (PER) induced by sugar stimulation in

the antennal and tarsal tip sensilla of virgin female adults. In consideration of the

sugar content in nectar and the response intensity of the contact chemosensilla in

adults, the test concentration of sugars was set at 100 mM. The results showed that the

PER percentage reached $64 \pm 4.35\%$, $44 \pm 5.96\%$ and $20 \pm 1.92\%$ when the antennal

sensilla were stimulated with sucrose, fucose and fructose (**Figure 2C**), while the

PER percentage reached $53 \pm 14.08\%$, $45 \pm 4.44\%$ and $36 \pm 10.19\%$ when the tarsal

sensilla were stimulated, respectively (**Figure 2D**). Stimulation of the antennal or

tarsal sensilla by other sugar compounds did not induce changes in the PER

percentage compared with stimulation of water (**Figure 2C–D**). The dose–response

curves showed that the threshold concentrations of sucrose, fucose, and fructose for

the antennal sensilla were 100 mM, 100 mM, and 1000 mM (**Figure 2E**), those for

the tarsal sensilla were 10 mM, 10 mM, and 100 mM, respectively (**Figure 2F**).

Phylogenetic tree and expression patterns of putative sugar gustatory receptors

On the basis of understanding the electrophysiological and behavioral responses of larvae and adults of *H. armigera* to various sugars, we aimed to reveal the molecular mechanisms of sugar reception in the contact chemosensilla of *H. armigera*. We first analyzed the putative sugar gustatory receptor genes based on the genome data of *H. armigera* (Pearce et al. 2017), the reported gene sequences of sugar GRs in *H. armigera* and their phylogenetic relationship of *D. melanogaster* sugar gustatory receptors (Jiang et al. 2015, Xu et al. 2012). All nine putative sugar GR genes in *H. armigera*, *Gr4–12* were validated, and their full-length cDNA sequences were cloned (The GenBank accession number is provided in **Appendix—Table 1**). The lengths of the amino acid sequences of the sugar GRs were consistent with the lengths of the corresponding reported sequences (the similarity was 97%-99%), except for Gr8, Gr9, and Gr10. Gr8 had two more amino acids at the C-terminal than the reported by Xu et al., 2017 with 99% sequence similarity (Xu et al. 2017). Gr9 had 9 more amino acids at the N-terminal than the reported in the *H. armigera* genome study with 99% sequence similarity (Pearce et al. 2017), while the sequence length of Gr9 was exactly the same as Gr4 in Jiang et al., 2015 with 99% sequence similarity (Jiang et al. 2015). Gr10 had 20 more amino acids than the sequence reported in the *H. armigera* genome study (Pearce et al. 2017) and 186 more amino acids than the sequence reported by Xu et al., 2017 (Xu et al. 2017), with 97% and 99% sequence similarities, respectively.

We established a phylogenetic tree using sugar GR genes of nine insect species, including *D. melanogaster*, *Aedes aegypti*, and *Anopheles gambiae* in Diptera; *Apis*

mellifera in Hymenoptera; and *Danaus plexippus*, *Heliconius melpomene*, *Bombyx*

mori, *Plutella xylostella*, and *H. armigera* in Lepidoptera (**Figure 3A**). The

phylogenetic analysis showed that candidate sugar GRs of these nine insect species

grouped within two clades, the sugar GR clade and the fructose GR clade. All the

putative sugar GRs of *H. armigera* were in the sugar GR clade except for HarmGr9,

which was in the fructose GR clade (**Figure 3A**).

To determine the expression patterns of nine putative sugar GR genes in larval and

adult taste organs, we measured their expression levels in the larval maxillary galea,

and the antennae, foreleg tarsi, and proboscis in virgin female and male adults by

real-time quantitative PCR (qRT-PCR) (**Figure 3B–3E, Figure 3—figure**

supplement 1). The results showed that the expression patterns of sugar GRs in the

same organ of male and female adults were nearly the same (**Figure 3C–3E**, **Figure**

3—figure supplement 1). The average expression level of all sugar GRs in the larval

maxillary galea was lower than that in the adult organs. In each taste organ, the

expression levels of 1–2 sugar GRs were much higher than those of other sugar GRs,

and these highly expressed GRs were significantly different between larvae and adults

(**Figure 3B–3E, Figure 3—figure supplement 1**). Among all sugar GRs, *Gr10* was

highly expressed in the larval maxillary galea, whereas *Gr6* was highly expressed in

the antennae, tarsi, and proboscis of both male and female adults. *Gr5* expression

levels greatly varied; they were highest in the proboscis followed by the tarsi, but very

low in the antennae and the maxillary galea of the larvae (**Figure 3B–E**, **Figure**

3—figure supplement 1).

Functional analysis of putative sugar gustatory receptors by ectopic expression

Of all the nine putative sugar GRs in *H. armigera* mentioned above, only Gr9,

originally named as Gr4, has been previous identified as a fructose receptor (Jiang et

al. 2015). To explore the functions of the other eight GRs, we first investigated the

inward current changes in *Xenopus* oocytes expressing sugar GRs individually in

response to the stimulation of 11 sugar compounds at 100 mM using two-electrode

voltage-clamp. The results showed that the oocytes expressing Gr10 only responded

to sucrose with a current value of 4.10 ± 0.53 μA (**Figure 4A–B**); the oocytes

expressing Gr6 showed responses to fucose and sucrose with a current value of $1.59 \pm$

0.08 μA and 0.24 ± 0.03 μA, respectively (**Figure 4A–B**). The oocytes expressing

remaining six sugar GR genes (Gr4, Gr5, Gr7, Gr8, Gr11, and Gr12) singly had no

response to the tested sugar compounds (**Figure 4—figure supplement 1A–B**). The

dose–response curves showed that the threshold concentration of sucrose for the

oocytes expressing Gr10 was 100 mM (**Figure 4C–D**); whereas the threshold

concentrations of sucrose and fucose for the oocytes expressing Gr6 were 250 mM

and 100 mM, respectively, and 500 mM fructose could also induce an inward current

(**Figure 4C–D**).

Given that there was only one sugar GSN responding to sucrose and fucose in each

larval maxillary galea, we speculated that Gr10 should be expressed together with Gr6

in this cell although their expressing levels were different. Therefore, we examined

the response profile of the oocytes expressing Gr10 together with Gr6. The results

showed that the oocytes co-expressing Gr10 and Gr6 significantly responded to

sucrose with a current value of 1.89 ± 0.58 μA at 100 mM, a threshold concentration,

and had a weak response to fucose with a threshold concentration of 250 mM (**Figure**

4).

Since Gr5 and Gr6 were highly expressed in the proboscis and tarsi (**Figure 3D–E,**

Figure 3—figure supplement 1), we suspected that Gr5 and Gr6 might be expressed

in the same cells, and then tested the response profile of their co-expression in oocytes.

Similar to the oocytes expressing Gr6, the oocytes expressing Gr5 and Gr6 had

responses to 100 mM fucose with a current value of 0.11 ± 0.015 μA and to 100 mM

sucrose with a current value of 0.03 ± 0.002 μA (**Figure 4—figure supplement**

1B–C).

In summary, we find that oocytes expressing single Gr10 or Gr6 confer sensitivity to

one or more sugars, indicating their direct roles in ligand recognition. Gr10 is tuned to

sucrose specifically, Gr6 is responsive to fucose, sucrose and fructose in the order of

sensitivity, and the combination of Gr10 and Gr6 is sensitive to sucrose and then

fucose, which is in line with the characteristics of the sugar GSNs in larval and adult

taste organs where the two GRs are highly expressed respectively.

Establishment of *Gr10* and *Gr6* homozygous mutants

In the above study, we discovered ligands of Gr10 and Gr6 using the ectopic

expression system. To validate the function of *Gr10* and *Gr6 in vivo*, we further

constructed *Gr10* and *Gr6* homozygous mutants (*Gr10$^{-/-}$* and *Gr6$^{-/-}$*, respectively)

using CRISPR/Cas9 techniques (**Figure 5**). *Gr10$^{-/-}$* and *Gr6$^{-/-}$* generated 4-bp and

17-bp deletions at the sgRNA targeting site **(Figure 5)**, respectively, which also

confirmed the deletion at the mRNA level in *Gr10$^{-/-}$* and *Gr6$^{-/-}$* (**Figure 5—figure**

supplement 1). The deletion resulted in truncated Gr10 and Gr6 proteins. The Gr10

sequence was reduced from 460 to 62 amino acids and contained no transmembrane

domains, whereas the Gr6 sequence was reduced from 452 to 115 amino acids and

contained only two transmembrane domains (**Figure 5**).

In order to predict the potential off-target sites of sgRNA of *Gr6* and *Gr10*, we used

online software Cas-OFFinder (http://www.rgenome.net/cas-offinder/) to blast the

genome of *H. armigera*, and the mismatch number was set to less than or equal to 3.

According to the predicted results, the *Gr10* sgRNA had no potential off-target region

but *Gr6* sgRNA had one. Therefore, we amplified and sequenced the potential

off-target region of *Gr6*$^{-/-}$ and found there was no frameshift or premature stop codon

in the region compared to WT (**Figure 5—figure supplement 2**). It is worth

mentioning that there was no potential off-target region of *Gr6* sgRNA or *Gr10*

sgRNA in all the other eight sugar receptor genes of *H. armigera*. We further found

there was no difference in the response to xylose of the medial sensilla styloconica

among WT, *Gr10*$^{-/-}$ and *Gr6*$^{-/-}$ (**Figure 5—figure supplement 2**). Furthermore, WT,

Gr10$^{-/-}$ and *Gr6*$^{-/-}$ did not show differences in the larval body weight, adult lifespan,

and number of eggs laid per female (**Figure 5—figure supplement 2**). All these

results suggest that no off-target effects occurred in the study.

Effects of *Gr10* or *Gr6* knockout on sugar reception

After obtaining the two homozygous mutants, we compared electrophysiological

responses of the sugar GSNs in larval lateral sensilla styloconica of the WT, *Gr10*$^{-/-}$,

and *Gr6*$^{-/-}$ to sucrose and fucose using 10 mM sinigrin as a positive control, which

induced the response of a bitter cell in the same sensilla. To 1 mM, 10 mM and 100

mM sucrose, the firing rates of the larval sugar GSNs in $Gr10^{-/-}$ were only 40%, 20%

and 28% of those in the WT, respectively, whereas those in $Gr6^{-/-}$ larvae did not differ

from those in the WT larvae (**Figure 6A–B**). To 10 mM and 100 mM fucose, the

firing rates of the larval sugar GSNs in $Gr6^{-/-}$ were only 20% and 5.8% of those in the

WT, respectively, whereas those of $Gr10^{-/-}$ did not differ from the WT (**Figure 6A–B**).

The electrophysiological responses of the bitter GSNs to sinigrin in the larval lateral

sensilla styloconica of the WT, $Gr10^{-/-}$, and $Gr6^{-/-}$ were all strong and did not differ

(**Figure 6A–B**). It is clear that $Gr10$ plays a key role in sucrose reception whereas

$Gr6$ plays a key role in fucose reception of the sugar GSNs in the lateral sensilla

styloconica of larvae.

Similarly, we compared the electrophysiological responses of the sugar GSNs in

contact chemosensilla of the adult antennae, foreleg tarsi, and proboscis of the WT,

$Gr10^{-/-}$, and $Gr6^{-/-}$ to 10 mM and 100 mM of sucrose and fucose, and 100 mM

fructose. Additionally, 10 mM nicotine was used as a positive control, that induced

electrophysiological responses of bitter GSNs in the tip sensilla of antennae and tarsi.

The firing rates of the sugar GSNs in the antennal sensilla of $Gr6^{-/-}$ to 10 mM and 100

mM sucrose, 10 mM and 100 mM fucose, and 100 mM fructose were only 32% and

9%, 22% and 11%, and 29% of those in the WT, respectively (**Figure 6C–D**). The

firing rates of the sugar GSNs in antennal sensilla of $Gr10^{-/-}$ to sucrose, fucose, or

fructose did not differ from those of the WT adults. The antennal sensilla of the WT,

$Gr10^{-/-}$, and $Gr6^{-/-}$ all showed strong responses to nicotine with no differences (**Figure

6C–D**). The response patterns of the sugar GSNs in the foreleg tarsi of the WT,

$Gr10^{-/-}$, and $Gr6^{-/-}$ to 10 mM and 100 mM sucrose, 10 mM and 100 mM fucose, and

100 mM fructose were similar to those of the sugar GSNs in antennae (**Figure 6E–F**).

The firing rates of the sugar GSNs in proboscis sensilla of $Gr6^{-/-}$ to sucrose, fucose,

and fructose were much lower and even close to zero (**Figure 6G–H**). The firing rates

of the sugar GSNs in tarsal and proboscis sensilla of $Gr10^{-/-}$ to sucrose, fucose, and

fructose did not differ from those of the WT adults (**Figure 6E–H**). The tarsal sensilla

of the WT, $Gr10^{-/-}$, and $Gr6^{-/-}$ responded to nicotine with no difference (**Figure**

6E–F).

Effects of *Gr10* and *Gr6* knockout on behavioral responses

To determine the effect of *Gr10* and *Gr6* knockout on feeding responses of *H.*

armigera larvae, we compared the larval feeding preferences of the WT, $Gr10^{-/-}$, and

$Gr6^{-/-}$ to sucrose by two-choice tests. The $Gr10^{-/-}$ larvae showed no preference for the

leaf discs painted with 10 mM sucrose ($P = 0.4258$). However, they still had

preference for the discs with 100 mM sucrose ($P = 0.0141$) (**Figure 7A–B**) though

their PI value was decreased than those of the WT (**Figure 7B**). The $Gr6^{-/-}$ larvae

were the same as the WT larvae and retained a significant preference for 10 mM or

100 mM sucrose diets (**Figure 7A–B**). The WT, $Gr10^{-/-}$, and $Gr6^{-/-}$ larvae all showed

similar preference for 10 mM and 100 mM fructose (**Figure 7—figure supplement**

1A–B).

We also compared the feeding amount in 1 hour of the larvae of WT, $Gr10^{-/-}$, and

$Gr6^{-/-}$ on cabbage leaves, corn kernels, pea seeds, pepper fruits, and tomato fruits by

no-choice tests. On each of these plant tissues, the $Gr10^{-/-}$ larvae fed significantly less

than the WT larvae, but the $Gr6^{-/-}$ larvae fed the same amount as the WT larvae

(**Figure 7C**).

Moreover, we examined the PER of the WT, $Gr10^{-/-}$, and $Gr6^{-/-}$ female adult antennae

and tarsi in response to stimulation of sucrose, fucose, and fructose at different

concentrations. When antennae or tarsi were stimulated with 10 mM, 100 mM, or

1000 mM sucrose, fucose, or fructose, $Gr6^{-/-}$ females showed a considerably lower

PER percentage than the WT females and even had little or no PER in response to

sucrose and fucose, whereas the PER percentage of $Gr10^{-/-}$ females was not different

from that of the WT females (**Figure 7D–I**).

In conclusion, $Gr10$ plays a key role in sucrose reception by the sugar GSNs in the

larval lateral sensilla styloconica, and mediates the feeding preference of larvae to

sucrose, whereas $Gr6$ plays a key role in sensing sucrose, fucose, and fructose by the

sugar GSNs in contact chemosensilla of the adult antennae, foreleg tarsi and proboscis,

and mediates the related PER behavior of adults.

Discussion

In this study, we reveal the different molecular bases of sucrose reception between *H.*

armigera larvae and adults through ectopic expression and genetic editing of *GR*s.

Gr10, which was highly expressed in the larval maxillary galea, is specifically tuned

to sucrose with high sensitivity, whereas Gr6, which was highly expressed in the adult

antennae, foreleg tarsi and proboscis, has a broader response spectrum but lower

sensitivity to sucrose, indicating that larvae mainly use Gr10 to detect low sucrose in

plant tissues, while adults primarily use Gr6 to detect a variety of sugars with high

content, including sucrose in nectar **(Figure 8)**.

Our findings indicate that the *H. armigera* larvae and adults adopt different

sugar-sensing systems to satisfy their requirements for food selection. The larvae

mainly feed on young leaves, flower buds, and fruits of host plants. As a universal

phagostimulant, sucrose is the major sugar in plants and considerably varies in

content in different tissues and developmental stages. Tomato is a principal host plant

of *H. armigera*. The sucrose content in its leaves generally ranges from 13.72 to 42.06

mM, but that in its fruits generally ranges from 35.2 to 988.22 mM (Schauer et al.

2005). The adults mainly feed on plant nectar. Sucrose and some monosaccharides,

especially fructose and glucose, are the main components in nectar, where the sugar

content is much higher than that in leaves and fruits. The sugar content in the nectar

of cotton plants is 10%–34% (which roughly corresponds to 0.5 M–1.7 M) (Vansell

1944, Waller et al. 1981), and that of soybean plants is even higher, around 23%–60%

(which roughly corresponds to 1.2–3 M) (Free 1993). It is clear that, in their life cycle,

a single sugar-sensing mechanism cannot meet the requirements for food selection by

this phytophagous insect species. From the perspective of insect and plant

co-evolution, using value-based and target-specific sugar-sensing systems is a

favorable choice for phytophagous insects to ensure their growth and reproduction on

plants.

Although Gr10 plays a major role in sucrose reception in the larval stage, the response

of the sugar GSNs in larval lateral sensilla styloconica to sucrose did not completely

disappear in *Gr10*$^{-/-}$ (**Figure 6A–B**), and the larvae still showed a behavioral

preference for the high concentration of sucrose (100 mM) (**Figure 7B**). It is most

likely that low-level expressed Gr6 is functional in the GSNs. The PER percentage of

$Gr6^{-/-}$ adults was not disappeared when the antennae and tarsi were stimulated by high

fructose concentrations (**Figure 7F and 7I**); this indicates that fructose-tuned GRs

other than *Gr6* were involved. Previous studies have shown that *Gr9* (*Gr4* in Jiang et

al., 2015), as an ortholog of DmelGr43a and BmorGr9, is a fructose-tuned GR (Ai et

al. 2022, Jiang et al. 2015, Sato et al. 2011, Xu et al. 2012). From our results,

knocking out *Gr10* or *Gr6* is unlikely to be compensated by overexpression of other

sugar GRs. One of our recent studies showed that *Orco* knockout had no significant

effect on the expression of most OR, IR and GR genes in adult antennae of *H.*

armigera although some genes were up- or down-regulated to a certain extent (Fan et

al. 2022). To elucidate the precise molecular mechanisms of sugar reception in *H.*

armigera is necessary to compare a series of single, double and even multiple Gr

knock-out lines and investigate the downstream effects of the GRs.

The larval maxillary galea with only two sensilla styloconica is an ideal system for

studying the mechanisms of sugar sensing. In each maxillary galea of *H. armigera*,

the only sugar GSN sensitive to sucrose and fucose is located in the lateral sensillum

styloconicum. Based on our results, we speculate that Gr10 and Gr6 are expressed in

this GSN with different levels, and mainly determine the response profile of the cell.

The firing rate of the cell in $Gr10^{-/-}$ to 1mM, 10 mM and 100 mM sucrose was

significantly lower than that in WT, but there was no difference between WT and

$Gr6^{-/-}$, indicating that Gr10 mainly determine the sensitivity of this cell to sucrose

(**Figure 6A–B**). In contrast, the firing rate of the cell in $Gr6^{-/-}$ to 10 mM and 100 mM

fucose was significantly lower than that in the WT, but there was no difference

between WT and *Gr10^{-/-}*, indicating that Gr6 was also functional in larvae (**Figure**

6A–B). This can be well explained by the response profiles of the oocytes expressing

Gr10, Gr6, and their combination **(Figure 4)**.

Sugars are common phagostimulants in animals, but the molecular basis of sugar

perception differs among animals. In mammals, the taste receptors T1R2 and T1R3

combine to complete the perception of all carbohydrate compounds, including sucrose,

and both receptors are indispensable (Damak et al. 2003, Li et al. 2002, Nelson et al.

2001, Zhao et al. 2003). In *D. melanogaster* adults, multiple sugar GRs are involved

in sugar reception, different GRs expressed in sweet taste neurons may function as

multimers (Dahanukar et al. 2001, Dahanukar et al. 2007, Fujii et al. 2015, Jiao et al.

2007, Jiao et al. 2008, Slone et al. 2007). Gr5a and Gr64a are widely used for

responses to complementary subsets of sugars in labellar sweet taste neurons

(Dahanukar et al. 2007), and Gr64f is also required broadly as a coreceptor for the

detection of sugars (Jiao et al. 2008). The recent successful ectopic expression of

single sugar GRs in olfactory neurons confirmed that each sugar GR is directly

involved in ligand recognition (Freeman et al. 2014). Sweet sensing neurons or

receptors have not been identified in the larval peripheral taste system (Kwon et al.

2011). Sugar perception in *Drosophila* larvae has been assigned to the fructose

receptor *Gr43a*, which, however, is expressed in pharyngeal sensory organs, foregut,

and in the brain, but absent from external taste neurons (Mishra et al. 2013). This is

most likely due to the fact that fruit fly larvae prefer to feed on rotting fruits at the

optimum stage of fermentation, which contain fructose as the main sugar. Anyway,

the sucrose sensing neurons were recently identified in the larval primary taste center

(Maier et al. 2021). In honeybees, AmGr1 acts as a GR for certain sugars other than

fructose, and AmGr2 acts as a coreceptor with AmGr1 (Jung et al. 2015). AmGr3 is

tuned to fructose and is phylogenetically closely related to the DmGr43a (Değirmenci

et al. 2020, Takada et al. 2018). The whitefly *Bemisia tabaci* consistently choose diets

containing higher sucrose concentrations. It has been shown that BtabGr1 displayed

significant sucrose specificity when expressed in *Xenopus* oocytes, and silencing of

this GR significantly interfered with the ability of *B. tabaci* adults to detect sucrose in

the phloem sap (Harari et al. 2023).

The most studied phytophagous insects for sugar perception are Lepidopterans.

BmGr9 in *B. mori* (Morinaga et al. 2022, Sato et al. 2011), HarmGr9 in *H. armigera*

(named as HarmGr4 by Jiang et al., 2015) (Jiang et al. 2015, Xu et al. 2012),

PxGr43a-1 in *P. xylostella* (Liu et al. 2020), and SlGr8 in *Spodoptera litura* (Liu et al.

2019) were all the orthologs of DmGr43a in *D. melanogaster* (Miyamoto et al. 2012),

and tuned to fructose without complexing with other GRs. BmGr4 was recently

identified as a receptor for sucrose and glucose and may function as a chemosensor

for nutritional compounds in both midgut and brain (Mang et al. 2022). In this study,

we find *Gr10* and *Gr6* were highly expressed in the larval and adult taste organs,

respectively. The experiments on ectopic expression in *Xenopus* oocytes shows that

Gr10 specifically responded to sucrose, Gr6 responded to fucose and sucrose, and

weakly to fructose, and the combination of Gr10 and Gr6 responded to sucrose and

weakly to fucose (**Figure 4A–B**). These results indicate that Gr10 and Gr6 are

capable of forming functional receptors singly, and the function of the two

co-expressed receptors is not equal to the sum of the functions of the two receptors

(Figure 4). When further tracing the genomic locations of the two GR genes, we

confirm that *Gr10* and *Gr6* together with other 6 sugar GR genes (except for *Gr9*) are

tandemly duplicated on the same scaffold of *H. armigera* (Pearce et al. 2017). The

sugar GR gene cluster and interactions between GRs could provide a degree of

evolutionary and regulatory flexibility that may meet the needs of the insect to sense

subtle sugar components and concentrations in food sources in a changing

environment (Delventhal and Carlson 2016). In this study, the expression patterns of 9

sugar GRs in three taste organs of adult *H. armigera* show that there is a disparity in

GRs, specifically GR5 and GR6, between the female antenna, tarsi and proboscis,

which may be an evolutionary adaptation reflecting subtle differentiation in the

function of these taste organs in adult foraging. Antennae and tarsi play a role in the

exploration of potential sugar sources, while the proboscis plays a more precise role in

the final decision to feed.

In this work, we report for the first time the molecular basis of sucrose reception in

external taste neurons of a Lepidopteran insect, and discover that different taste

receptors are used for food selection in *H. armigera* larval and adult stages. These

findings greatly improve our understanding of "sweet" taste of phytophagous insects.

According to the phylogenetic tree of sugar GRs in insects, *Gr10* is clustered in the

same branch with BmorGr5 of *Bombyx mori*, whereas *Gr6* is clustered in another

branch with BmorGr6 (**Figure 3A**). GRs closely associated with *Gr10* and *Gr6* are

also found in other Lepidopteran species such as *D. plexippus*, *H. melpomene*, and *P.*

xylostella (**Figure 3A**). Therefore, we speculate that similar sugar-sensing

mechanisms may also exist in other Lepidopteran species, which is worth verifying in

these species in the future. Knockout of *Gr10* or *Gr6* led to a significant decrease in

sugar sensitivity and food preference of the larvae and adults of *H. armigera*,

respectively, which is bound to bring adverse consequences to the survival and

reproduction of the insects. Therefore, studying the molecular mechanisms underlying

sugar perception in phytophagous insects would provide new insights into the

behavioral ecology of this highly diverse and important group of insects, and may also

provide new means for agricultural pest control by blocking or disrupting sugar

receptors.

Materials and Methods

Insects

Helicoverpa armigera were collected from Xuchang (Henan Province, China) and

raised in an indoor environment (27 ± 1°C, 70%–80% humidity) with a 16 : 8 h light :

dark cycle. The larvae fed on artificial diet (mainly including wheat bran, wheat germ,

and tomato paste) in a glass tube (I.D., 2.5 cm; length, 8 cm). When the larvae

reached the 3rd instar, they were transferred to a new glass tube and raised individually

until pupation. Female and male moths were raised separately until they became

sexually mature and then moved to a cage for mating. Since the taste system of the

female is similar to that of the male and the nutrient requirements are more critical for

females (Jiang et al. 2015, Zhang et al. 2010), the virgin female adults were used in

the related experiments.

Xenopus laevis

Xenopus laevis were fostered at 20 ± 1°C in the Animal Laboratory Center, Institute

of Genetics and Developmental Biology, Chinese Academy of Sciences. They fed on

artificial frog food (mainly including fish meal, yeast extract powder, high-gluten

flour). All experiments were approved by the Animal Care and Use Committee of the

Institute of Zoology, Chinese Academy of Sciences and followed the Guide for the

Care and Use of Laboratory Animals (IOZ17090-A).

Scanning electron microscopy

Just as our previous study on tarsal contact chemosensilla in fore legs of female moths

(Zhang et al. 2010), the samples of antennae or proboscis in adult female *H. armigera*

were mounted directly on stainless steel sample buds and sputter-coated with a 10 nm

thick layer of gold. Photomicrographs were obtained with a scanning electron

microscope (HITACHI SU8010) in the Institute of Microbiology, Chinese Academy

of Sciences.

Tip-recording

The tip-recording technique based on a previously described methods (Hodgson et al.

1955, van Loon 1990, Zhang et al. 2010) was used to record the electrophysiological

responses of lateral and medial sensilla styloconica in larval maxillary galea, sensilla

chaetica in adult antennae, sensilla trichodea in adult tarsi, and sensilla styloconica in

adult proboscis of *H. armigera*. A glass capillary filled with the test compound

solution, into which a silver wire was inserted, was placed in contact with the contact

chemosensillum to be recorded as the recording electrode. The tip diameter of the

electrode was about 50 μm, suitable for recording from a single sensillum. Action

potentials (spikes) generated during the first second after stimulus onset were

amplified by a preamplifier and was sampled with a computer equipped with a

Metrabyte DAS16 A/D conversion board. The amplifier used an AD 515-K (Analog

Devices) integrated circuit in the first stage, yielding < 1 pA input bias current, 1015

Ohm and 0.8 pF input impedance. An interface (GO-box) was used for signal

conditioning. This involved a second-order band pass filter (-3 dB frequencies: 180

and 1700 Hz). Stimuli were dissolved in 2 mM KCl, which was used as the control.

Two pentoses (arabinose and xylose), four hexoses (fructose, fucose, galactose and

glucose), and two disaccharides (sucrose and trehalose) (**Key resources table**) were

used as sugar stimuli. Each sugar first tested at 10 mM to screen active compounds,

which were then tested in a range of concentrations in dose-response assays (0.01 mM,

0.1 mM, 1 mM, 10 mM and 100 mM). To avoid possible adaptation of the contact

chemosensilla tested, the interval between two successive stimulations was at least 3

min. Digitized traces were analyzed by means of SAPID Tools software version 3.5

(Smith et al. 1990) and Autospike 3.7 (Syntech, Buchenbach, Germany). The spikes

were sorted on the basis of typical biphasic waveforms and spike amplitudes, and the

frequency of the same type of spikes was counted from the first second after

stimulation.

In recording of electrophysiological responses of larvae, the first day of larvae in the

5^{th} instar were used. To guarantee all test caterpillars were in the same stage, we first

selected 4^{th} instar larvae which were preparing to undergo ecdysis, and then

transferred them to a glass tube with green pepper. The test larva was starved for

approximately 2 h before recording. The head was cut and fixed in a silver holder

which was connected to a preamplifier with a copper miniconnector. The lateral and

medial sensilla styloconica in larval maxillary galea were exposed for testing

electrophysiological responses. About 12–13 replicates were set for each treatment.

The virgin female moths on the second day after eclosion were used for recording

electrophysiological responses (Jiang et al. 2015, Zhang et al. 2010). They were only

fed with ddH$_2$O. When testing sensilla in a tarsus, the foreleg was cut between coxa

and femur, and the silver holder connected to the amplifier was inserted into the femur.

When testing sensilla in the antennae and proboscis, the head and thorax without all

legs were reserved. The silver holder was inserted into the thorax. The orientation of

the taste organ was gently adjusted to allow the sensilla to be touched by the electrode.

To guarantee the sensilla was steadily exposed, the distal part of antenna or proboscis

was fixed on a small platform with double sided adhesive tape. For each individual,

three sensilla on the top area of antennae, foreleg tarsi, or proboscis were chosen for

testing electrophysiological responses, and about 15–21 replicates were used for each

treatment.

Two-choice tests

We determined the behavioral responses to sugars of the *H. armigera* larvae using the

two-choice test as previously described with small modifications (Chen et al. 2022).

We used the leaf of cabbage (*Brassica oleraceaas*, Shenglv-7) as the substrate since it

is not a favorite food for larvae. Circular cabbage leaf discs of 1 cm in diameter were

prepared. The upper surface of the treated leaf disc was painted with the 35.2 μL sugar

solution with 0.01% Tween 80 (CAS: 9005-55-6, Sigma-Aldrich, St Louis, USA)

using a paintbrush, and the control leaf disc was painted with 35.2 μL ddH$_2$O with

0.01% Tween 80. The 0.01% Tween 80 was added to reduce the surface tension of the

cabbage leaves and increase the malleability of the test sugar solution (Blaney and

Simmonds 1988). The feeding preference of larvae to eight sugars including arabinose,

fructose, fucose, galactose, glucose, sucrose, trehalose and xylose (**Key resources**

table) was tested at 10 mM. 0.01 mM, 0.1 mM, 1 mM, 10 mM, and 100 mM sucrose

and fructose were used in a dose-response assay. The intact and the remaining leaf

discs were scanned by DR-F120 scanner (Canon, Tokyo, Japan), and the area of intact

and remaining leaf discs were calculated by ImageJ softwere (NIH) (Abràmoff et al.

2004, Chen et al. 2022). The feeding area = the area of intact leaf disc – the area of

remaining leaf disc. The preference index (PI) was calculated according to the

following formula: PI = (the feeding area of test leaf discs – the feeding area of

control leaf discs) / (the feeding area of test leaf discs + the feeding area of control

leaf discs). 20 replicates were run.

Adult proboscis extension reflex (PER) tests

We determined the behavioral responses to sugars of the *H. armigera* female adults

using the proboscis extension reflex (PER) test as previously described with some

modifications (Jiang et al. 2015). Arabinose, fructose, fucose, galactose, glucose,

sucrose, trehalose, and xylose (**Key resources table**) were first tested at 100 mM to

screen active compounds, which were then tested in a range of concentrations in

dose-response assays (0.01 mM, 0.1 mM, 1 mM, 10 mM, 100 mM, and 1000 mM). In

brief, the virgin females on the first day after eclosion were used. Each moth was

harnessed in a 1 mL pipette with a tip removed, from which its head protruded, and its

antennae, proboscis and forelegs were permitted to move freely. They were allowed to

adapt to the environment for about 12 h, and satiated with ddH_2O until the trial. The

testing moth was stimulated to the distal part of one antenna with a given test solution

in an absorbent cotton ball, and the PER in response to the stimulation was recorded.

The interval between two successive stimulations was 5 min, during which the distal

part of the antenna was washed, and the insect was fed with ddH$_2$O if needed. Four

hours later, two foreleg tarsi of the moth were simultaneously stimulated with the test

solutions in the same way as the antenna was stimulated, and the PER was recorded.

20 to 30 individual moths were used for each set of tests. The experiments were

repeated three times with groups of insects from three different rearing batches.

RNA extraction and cDNA synthesis

The galea of larvae, and the antennae, foreleg tarsi, proboscis, and abdomen of both

male and female adults were removed and directly placed into 1.5 mL Eppendorf

tubes (Axygen, Tewksbury, MA, USA) containing 1 mL QIAzol Lysis Reagent

(QIAGEN, Hilden, Germany); then, the extraction of total RNA was carried out

following the manufacturer's protocols. After extraction, the RNA quality and

concentration were determined with a NanoDrop 2000 spectrophotometer (Thermo

Fisher Scientific, Waltham, MA, USA). First, for cDNA synthesis 2 μg total RNA

was mixed with 1 μL oligo-dT (Promega, Madison, WI, USA); then, the mixture was

heated at 70°C 5 min and followed by an ice bath for 5 min. M-MLV reverse

transcriptase (Promega, Madison, WI, USA) was added; subsequently, the mixture

was incubated at 42°C for 90 min. The products were dissolved in ddH$_2$O and stored

at −20°C for later use.

Gene cloning of sugar GRs

We designed specific primers (**Appendix—Table 2**) according to the reported sugar

GR sequence of *H. armigera* (the GenBank accession number is provided in

Appendix—Table 1) for PCR cloning (Pearce et al. 2017, Xu et al. 2017). The

GenBank accession number of *Gr9* (named as *Gr4* in Jiang et al., 2015) nucleic acid

sequence and protein sequence were JX982536.2 and AKG90011 in Jiang et al., 2015,

respectively (Jiang et al. 2015). The Q5 High-fidelity DNA Polymerase (New

England Biolabs, Beverly, MA, USA) was used to clone sugar sequences except *Gr10*.

The PCR instrument was Applied Biosystems (Thermo Fisher Scientific, Woodlands,

Singapore) and the sequence amplification procedure was as follows: 98°C for 10 s,

40 cycles (60°C for 30 s, 72°C for 1 min), and final extension at 72°C for 2 min.

TransStart FastPfu DNA Polymerase (TransGen Biotech, Beijing, China) was used to

clone the *Gr10* sequence. The amplification procedure was as follows: 95°C for 1 min,

40 cycles (95°C for 20 s, 50°C for 20 s, 72°C for 45 s), and 72°C for 5 min. PCR

products were purified, subcloned into pGEM-T vector according to the

manufacturer's instructions (Promega), and sequenced. *Gr4* (402 amino acids) and

Gr7 (428 amino acids) were obtained from the tissues of male and female *H.*

armigera tarsi, respectively. *Gr5* (482 amino acids)*, Gr6* (452 amino acids)*,* and *Gr10*

(460 amino acids) were all obtained from the tissues of female *H. armigera* proboscis.

Gr8 (427 amino acids) and *Gr9* (474 amino acids) was obtained from the tissues of

female *H. armigera* antennae. Finally, *Gr11* (435 amino acids) and *Gr12* (434 amino

acids) were obtained from the tissues of female and male *H. armigera* abdomens,

respectively. No alternatively spliced transcript variants were found in the gene

cloning of all the sugar GRs studied.

Xu et al., 2017 (Xu et al. 2017) reported that *Gr10* was a pseudogene because the

nucleotide sequence changed at positions 521–524 compared with the sequence

reported in Pearce et al., 2017 (Pearce et al. 2017), and it terminated early during

translation. The *Gr10* sequence that we obtained had no termination on the same

position. Aligning the *Gr10* sequences of Pearce et al., 2017 and Xu et al., 2017

revealed that *Gr10* in Xu et al., 2017 had 22 more amino acids at the C-terminus than

that in Pearce et al., 2017. Therefore, we designed upstream and downstream primers

of *Gr10* from the N-terminus in Pearce et al., 2017 and the C-terminus in Xu et al.,

2017. Finally, we obtained the full-length sequence of *Gr10* with 460 amino acids.

Phylogenetic analysis of sugar GRs

The phylogenetic tree was built with *H. armigera* and sugar GRs of eight other insect

species: Diptera, *Drosophila melanogaster* (Robertson et al. 2003), *Aedes aegypti*

(Kent et al. 2008), and *Anopheles gambiae* (Hill et al. 2002); Hymenoptera, *Apis*

mellifera (Robertson and Wanner 2006); and Lepidoptera, *Danaus plexippus* (Zhan et

al. 2011), *Heliconius melpomene* (Briscoe et al. 2013), *Bombyx mori* (Guo et al. 2017),

and *Plutella xylostella* (Engsontia et al. 2014). Amino acid sequences were aligned

with MAFFT version 7.455 (Rozewicki et al. 2019), and gap sites were removed with

trimAI version 1.4 (Capella-Gutierrez et al. 2009). The phylogenetic tree was

constructed using the maximum likelihood method in IQ-tree version 6.8 (Nguyen et

al. 2015). The bootstrap algorithm was based on the Jones–Taylor–Thornton (JTT) +

F + G4 model and run 5000 times. Visualization and later modification of

phylogenetic trees were performed in FigTree version 1.4.4

(http://tree.bio.ed.ac.uk/software/figtree/).

Quantitative real-time PCR

The relative expression levels of sugar GRs in *H. armigera* were analyzed by

quantitative real-time PCR (qRT-PCR) (Livak and Schmittgen 2001). The specific

primers **(Appendix—Table 3**) were designed by Primer3plus

(http://www.primer3plus.com/cgi-bin/dev/primer3plus.cgi). qRT-PCR was conducted

with a QuantStudio 3 Real-Time PCR System (Thermo Fisher Scientific, Waltham,

MA, USA). All reactions included 10 μL SYBR Premix Ex *Taq* (TaKaRa, Shiga,

Japan); 0.4 μL (10 mM) primer (forward and reverse), 2 μL template DNA, 0.4 μL

ROX Reference Dye II Dye, 6.8 μL ddH$_2$O (20 μL total). The procedure was as

follows: 95°C for 30 s; 40 cycles of 95°C for 5 s and 60°C for 34 s; 95°C for 15 s;

60°C for 1 min; and 95°C for 15 s. *18S* ribosomal RNA (GenBank number:

KT343378.1) was used as the standard gene because the Ct value of this gene was not

different between different tissues (Sharath Chandra et al. 2014). The relative

expression levels of the specific genes were quantified by using the $2^{-\Delta Ct}$ method,

where ΔCt is the Ct value of GR genes subtracted from that of *18S*. Three biological

replicates were run for all experiments, and there were 30–100 insects in each

replicate.

Ectopic expression and functional analysis of sugar GRs

The sugar GRs sequences which were used for ectopic expression and functional

identification need to contain restriction enzyme sites and Kozak sequences (the

primer sequences of *H. armigera* sugar GRs for functional identification are provided

in **Appendix—Table 4**). The full-length sequence of the sugar receptor was cloned

from cDNA and transferred into pGEM-T vector (Promega, Madison, WI, USA).

After cleavage by appropriate endonucleases, it was linked to pCS2$^+$ vector. pCS2$^+$

vector was linearized and cRNA was synthesized according to the protocol of the

mMESSAGE mMACHINE SP6 Kit (Ambion, Austin, TX, USA). The products were

dissolved in ddH$_2$O, diluted to 3000 ng/μL, and then stored at −80°C until use.

Each of Gr4, Gr5, Gr6, Gr7, Gr8, Gr10, Gr11, Gr12, Gr5 + Gr6, and Gr10 + Gr6 was

expressed in oocytes. The acquisition of *X. laevis* oocytes was performed following a

previously described protocol (Nakagawa and Touhara 2013). *X. laevis* were

anesthetized by a mixture of ice and water for 30 min; then, the abdomen was cut to

obtain oocytes. The *Xenopus* oocytes were treated with 2 mg/mL collagenase type I

(Sigma-Aldrich, St Louis, MO, USA) in a calcium-free saline buffer (82.5 mM NaCl,

2 mM KCl, 1 mM MgCl$_2$, 5 mM HEPES, pH = 7.5; the reagents are provided in **Key**

resource table) at room temperature for 25 min. The oocytes were microinjected with

27.6 nL (80 ng) cRNA of each *GR* by SMART Touch^{TM} (World Precision

Instruments, Sarasota, FL, USA). For Gr10 + Gr6 or Gr5 + Gr6, a cRNA mixture with

a ratio of 1:1 was used. The oocytes microinjected with RNAse-free water were used

as the control. Then, the oocytes were incubated for 3–4 days at 16°C in the bath

solution (96 mM NaCl, 2 mM KCl, 1 mM MgCl$_2$, 1.8 mM CaCl$_2$, 5 mM HEPES, pH

7.5; reagents are provided in **Key resource table**).

The whole-cell current was recorded with a two-electrode voltage clamp, the

procedures as previously described (Jiang et al. 2015). The glass electrode, filled with

3M KCl, was inserted into the oocytes and maintained at 0.2–2.0 MΩ. The signal was

amplified with an OC-725C amplifier (Warner Instruments, Hamden, CT, USA) at a

holding potential at –80 mV all the time, low-pass filtered at 50 Hz, and digitized at 1

kHz. The sugar compounds (**Key resource table**) were dissolved in $1 \times$ Ringer

solution (96 mM NaCl, 2 mM KCl, 1 mM $MgCl_2$, 1.8 mM $CaCl_2$, 5 mM HEPES, pH

7.5) as the test solution. The oocytes were stimulated by test solution via the perfusion

system for 20 s each time, and then washed by $1 \times$ Ringer solution (Nakagawa and

Touhara 2013). The concentration of 100 mM for each chemical was randomly used

at first, and then concentration gradients (ranging from 0.01 mM, 0.1 mM, 1 mM, 10

mM, 50 mM, 100 mM, 250 mM, and 500 mM) were recorded later when a clear

current response was detected. Data were obtained and analyzed using Digidata

1322A and pCLAMP software version 10.4.2.0 (Axon Instruments Inc., Foster City,

CA, USA).

In vitro **synthesis of single-guide RNA**

Gr10 and *Gr6* single-guide RNA (sgRNA) position screening was conducted at

http://www.rgenome.net/cas-designer/. The sgRNA target site of *Gr10* was designed

on exon 4 (**Figure 5B**), and the sgRNA target site of *Gr6* was designed on exon 2

(**Figure 5E**). The *Gr10* and *Gr6* sgRNA primers (**Appendix—Table 5**) were designed

based on the principles provided by the GeneArt^{TM} Precision gRNA Synthesis Kit

manual (Invitrogen, Vilnius, Lithuania). The sgRNAs were synthesized by the

GeneArt^{TM} gRNA Prep Kit (Invitrogen, Vilnius, Lithuania), and purified by the

GeneArt^{TM} gRNA Clean Up Kit (Invitrogen, Vilnius, Lithuania). The products were

dissolved in RNAse-free water and then stored at –80°C until use. TrueCut^{TM} Cas9

protein 2 (Invitrogen, Vilnius, Lithuania) was diluted to 500 ng/mL and stored at

–20°C.

Embryo microinjection

The fresh laid eggs of *H. armigera* were placed on microscope slides and fixed on

double-sided adhesive tape. These operations were finished within 2 h after

oviposition. Microinjection into these pretreated eggs was performed with a mixture

of *Gr10* (or *Gr6*) sgRNA and Cas9 protein, with final concentrations at 250 ng/μL and

100 ng/μL, respectively. Microinjection was finished using a PLI-100A

PICO-INJECTOR (Warner Instruments, Hamden, Connecticut, USA). The injected

eggs were placed in a 10-cm Petri dish, covered with wet filter paper, and incubated at

25°C until hatching.

DNA extraction and mutagenesis detection

To screen *Gr10* (or *Gr6*) homozygous mutants, the tarsi on hindlegs were used for

DNA extraction with the Animal Tissue PCR Kit (TransGen Biotech, Beijing, China).

Then, we used PCR to amplify the sequences that covered *Gr10* or *Gr6* sgRNA

targeting sites. The reactions were 50 μL total and included: 2 μL template, 1 μL

forward primer, 1 μL reverse primer (mutation detection primer sequences are

provided in **Appendix—Table 5**), 25 μL Premix *Taq*^{TM} (Takara, Dalian, China), and

19 μL ddH$_2$O. The procedure was as follows: 94°C for 3 min, 40 cycles [94°C for 30

s, 55°C (*Gr6*), or 60°C (*Gr10*) for 30 s], 72°C for 1 min, and finally 72°C for 5 min.

The PCR products were sent to Beijing Genomics Institute (BGI) or the

SinoGenoMax for Sanger sequencing, and the sequence results were analyzed by

SnapGene® software version 4.3.8 (from Insightful Science; available at

snapgene.com) and SeqMan software version 7.1 (DNASTAR, Madison, WI, USA).

Screening homozygous mutants

Gr10 and *Gr6* homozygous mutant lines (*Gr10*$^{-/-}$ and *Gr6*$^{-/-}$, respectively) were

established. A mixture of Gr10 sgRNA and Cas9 protein was co-injected into each of

732 eggs that were laid within two hours. After hatching, the larvae were raised on

artificial diet, and finally 125 G0 adults were obtained. Among 125 G0 adults, 15 G0

harbored multiple targeted mutations. Each mutated G0 was backcrossed with the

wild type to obtain G1 offspring. From 57 G1 adults, one DNA strands containing 4

bp-deleted modifications, were in-crossed to establish stable lines; for the expanding

mutant population, 4 bp-deleted homozygotes of G2 moths were in-crossed to

generate homozygotes G3 moths (**Figure 5A).** Similarly, a mixture of Gr6 sgRNA and

Cas9 protein was co-injected into each of 695 eggs. After hatching, the larvae were

raised on artificial diet, and finally 132 G0 adults were obtained. Among 132 G0

adults, 12 G0 harbored multiple targeted mutations. From 66 G1 adults, one DNA

strands containing 17 bp-deleted modifications, were in-crossed to establish stable

lines; for the expanding mutant population, 17 bp-deleted homozygotes of G2 moths

were in-crossed to generate homozygotes G3 moths (**Figure 5D**). The related

mutation rates are provided in **Appendix—Table 6.**

To confirm the deletion in *Gr10*$^{-/-}$ and *Gr6*$^{-/-}$ at mRNA level, the total RNA of the

female adult proboscis of WT, *Gr10*$^{-/-}$ and *Gr6*$^{-/-}$ were extracted, and the cDNA was

synthesized following the previous gene cloning protocol. The specific primers

(**Appendix—Table 5**) were designed based on the DNA exon sequences of *Gr10* and

Gr6, which covered the DNA-confirmed sgRNA targeting sites. During mRNA

sequence cloning, the PCR reaction conditions are largely the same as those used for

DNA sequence confirmation, with little modification (the annealing temperature

reduced to 50°C). The nucleic acid sequences of the identified *Gr10* and *Gr6* were

aligned manually by tools at the T-Coffee web server (Notredame et al. 2000).

Off-target effects detection

The potential off-target sites were predicted as previously described (Chen et al. 2020).

The online software Cas-OFFinder (http://www.rgenome.net/cas-offinder/) was used

to blast the *Gr10* and *Gr6* sgRNAs against the *H. armigera* genome (Pearce et al.

2017), with mismatch number ≤ 3. According to the blast results, *Gr10* has no

potential off-target site, and one sequence (GenBank accession number:

LOC126054697) in the presence of *Gr6* sgRNA showed a potential off-target region.

The specific primers (**Appendix—Table 5**) were designed based on the DNA

sequence of LOC126054697 covering the potential off-target sites. The DNA

extraction and mutagenesis detection methods were described previously. Five

individual replicates were performed.

To further illustrate that the mutation strains did not exhibit off-target effects, the

electrophysiological responses of larvae and the biological parameters were compared

among WT, $Gr10^{-/-}$, and $Gr6^{-/-}$. The larval medial sensilla styloconica to xylose, the

body weight of the fifth instar larvae on the first day, the adult lifespan, and the

number of eggs laid were recorded. Five to sixteen individuals were recorded.

Effects of *Gr10* or *Gr6* knockout on sugar reception in larvae and adults

We recorded the electrophysiological responses of contact chemosensilla in the wild

type (WT), $Gr10^{-/-}$, and $Gr6^{-/-}$ by the tip-recording technique as described earlier.

Concentration of 1 mM, 10 mM, and 100 mM for sucrose and fucose were used as

stimuli for lateral sensilla styloconica in larvae. As the response thresholds of sugar

GSNs in adults were higher than those in larvae, concentrations of 10 mM and 100

mM for sucrose and fucose and 100 mM for fructose were used for adult contact

chemosensilla. 2 mM KCl was used as the control for both larval and adult contact

chemosensilla. 10 mM sinigrin and 10 mM nicotine were used as a positive control

(induced the response of a bitter cell in the same sensilla) for larval and adult contact

chemosensilla, respectively. All stimuli were tested randomly, and 12–20 replicates

were set for each experiment.

Effects of *Gr10* or *Gr6* knockout on behavioral responses of larvae to sugars and

plant tissues

The feeding preference of WT, $Gr10^{-/-}$, and $Gr6^{-/-}$ larvae to 10 mM and 100 mM

sucrose was investigated in two-choice tests as described earlier. 20 replicates were

set for each experiment. We also performed the no-choice tests to measure the feeding

amount of WT, $Gr10^{-/-}$, and $Gr6^{-/-}$ larvae in 1 h on leaves of cabbage (*Brassica*

oleraceaas, Shenglv-7), corn kernels of maize (*Zea mays,* Jingke-968), seeds of pea

(*Pisum sativum*, Changshouren), fruits of pepper (*Capsicum frutescens*, Changjian),

and fruits of tomato (*Solanum lycopersicum*, Ailsa Craig-LA2838A). Fifth instar

larvae on the first day after molting were starved for 2.5 h before testing. Each plant

tissue was weighed and put into a Petri dish (10 cm diameter $\times$ 2 cm depth; Corning,

Wujiang, China), and then one larva was introduced. The larva was removed after 1 h,

and the rest of the plant tissue was weighed. Petri dishes with plant tissues but no

caterpillar was used to measure water evaporation of the plant tissues. The feeding

amount of larvae is calculated based on the weight changes of the plant tissue before

and after being fed. 20 replicates were set for each experiment.

Effects of *Gr10* or *Gr6* knockout on behavioral responses of adults to sugars

The behavioral responses of the female adults of WT, *Gr10*$^{-/-}$, and *Gr6*$^{-/-}$ to sugars

were determined by using the PER test as described earlier. sucrose, fucose, and

fructose at the concentrations of 10 mM, 100 mM, and 1000 mM were tested

randomly to antennae and tarsi. H_2O was used as the control. All the experiments

were repeated three times, and 20 individual moths were used for each set of tests.

Statistical analysis

Statistical analyses were performed by SPSS 20 (IBM, Chicago, IL, USA) and

GraphPad Prism 8.2.1 (Dotmatics, San Diego, CA, USA). For the tip-recording data

and the PER data, the comparison between test compounds and the control was

performed using two-tailed independent-samples *t* test. For the two-choice feeding

test, paired-samples *t* test was used. The data of dose-response curves in tip-recording,

PER experiments and two-electrode voltage-clamp recording, and gene relative

expression levels and feeding amount in the no-choice test were analyzed by one-way

ANOVA with Tukey's HSD tests. The comparison among mutation strains and WT in

electrophysiological and PER experiments were analyzed by two-way ANOVA with

post hoc Tukey's multiple comparison. All analytical methods were tested at $P < 0.05$

for significant differences.

Acknowledgments

We thank our colleagues Zi-Lin Li, Qing Zhang, Yan Chen, Jun Yang, Xin-Lin Gong,

Yan-Yan Jia for assistances in PER experiments, mutant line screening and scanning

electron microscopy. This work was supported by National Key R&D Program of

China (Grant No. 2022YFD1400800), the National Natural Science Foundation of

China (Grant No. 32130090), and the Beijing Natural Science Foundation (Grant No.

6224062).

Author Contributions

Conceptualization, C.Z.W. and S.S.Z.; Methodology, S.S.Z., P.C.W., L.Q.H., C.N.,

G.C.L., and C.Z.W.; Software, S.S.Z., P.C.W., C.N., G.C.L., and C.Z.W.; Validation,

S.S.Z. and C.Z.W.; Formal Analysis, S.S.Z., C.N., G.C.L., and C.Z.W.; Investigation,

S.S.Z., P.C.W., K.Y., L.L.C., L.Q.H., and C.Z.W.; Resources, S.S.Z., P.C.W., L.Q.H.,

and C.Z.W.; Data curation, S.S.Z., C.N., P.C.W., G.C.L., and C.Z.W.; Writing –

original draft, C.Z.W. and S.S.Z.; Writing – review & editing, C.Z.W., S.S.Z.;

Visualization, S.S.Z. and C.Z.W.; Supervision, C.Z.W.; Project administration, C.Z.W.

and L.Q.H.; Funding acquisition, C.Z.W. and K.Y.

Competing interests

826   The authors declare no competing interests.

Data availability

829 All data and materials generated in this study are available in the manuscript,

830 supplementary materials, Appendix, and source data. Sequence data that support the

831 finding of this study have been deposited in GenBank with the accession numbers

832 (see **Appendix—Table 1**).

References

Abràmoff MD, Magalhães PJ, Ram SJ. 2004. Image processing with ImageJ.

Biophotonics international 11:36-42

Ai D, Dong C, Yang B, Yu C, Wang G. 2022. A fructose receptor gene influences

development and feed intake in *Helicoverpa armigera*. *Insect Science* **29**:993-1005.

DOI: https://doi.org/10.1111/1744-7917.12984

Bernays EA, Chapman RF. 1994. *Host-Plant Selection by Phytophagous Insects*.

Chapman & Hall.

Blaney WM, Simmonds MSJ. 1988. Food selection in adults and larvae of three

species of Lepidoptera: a behavioral and electrophysiological study. *Entomologia*

Experimentalis Et Applicata **49**:111-121. DOI:

https://doi.org/10.1111/j.1570-7458.1988.tb02482.x

Blaney WM, Simmonds MSJ. 1990. A behavioural and electrophysiological study of

the role of tarsal chemoreceptors in feeding by adults of Spodoptera, *Heliothis*

virescens and *Helicoverpa armigera*. *Journal of Insect Physiology* **36**:743-756.

DOI: https://doi.org/10.1016/0022-1910(90)90048-K

Briscoe AD, et al. 2013. Female behaviour drives expression and evolution of

gustatory receptors in butterflies. *PLoS Genetics* **9**:e1003620. DOI:

https://doi.org/10.1371/journal.pgen.1003620

Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. 2009. trimAl: a tool for

automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*

25:1972-1973. DOI: https://doi.org/10.1093/bioinformatics/btp348

Chapman RF. 2003. Contact chemoreception in feeding by phytophagous insects.

Annual Review of Entomology **48**:455-484. DOI:

https://doi.org/10.1146/annurev.ento.48.091801.112629

Chen W, Dong YH, Saqib HSA, Vasseur L, Zhou WW, Zheng L, Lai YF, Ma XL, Lin

LY, Xu XJ. 2020. Functions of duplicated glucosinolate sulfatases in the

development and host adaptation of *Plutella xylostella*. *Insect Biochemistry and*

Molecular Biology 119. DOI: https://doi.org/ 10.1016/j.ibmb.2020.103316

Chen Y, Wang PC, Zhang SS, Yang J, Li GC, Huang LQ, Wang CZ. 2022. Functional

analysis of a bitter gustatory receptor highly expressed in the larval maxillary galea

of *Helicoverpa armigera*. *PLoS Genetics* 18:e1010455. DOI:

https://doi.org/10.1371/journal.pgen.

Dahanukar A, Foster K, van der Goes van Naters WM, Carlson JR. 2001. A Gr

receptor is required for response to the sugar trehalose in taste neurons of

Drosophila. *Nat Neurosci* **4**:1182-1186. DOI: https://doi.org/10.1038/nn765

Dahanukar A, Lei YT, Kwon JY, Carlson JR. 2007. Two Gr genes underlie sugar

reception in *Drosophila*. *Neuron* **56**:503-516. DOI:

https://doi.org/10.1016/j.neuron.2007.10.024

Damak S, Rong M, Yasumatsu K, Kokrashvili Z, Varadarajan V, Zou S, Jiang P,

Ninomiya Y, Margolskee RF. 2003. Detection of sweet and umami taste in the

absence of taste receptor T1r3. *Science* **301**:850-853. DOI:

https://doi.org/10.1126/science.1087155

Değirmenci L, Geiger D, Rogé Ferreira FL, Keller A, Krischke B, Beye M,

Steffan-Dewenter I, Scheiner R. 2020. CRISPR/Cas 9-mediated mutations as a new

tool for studying taste in Honeybees. *Chemical Senses* **45**:655-666. DOI:

https://doi.org/10.1093/chemse/bjaa063

Delventhal R, Carlson JR. 2016. Bitter taste receptors confer diverse functions to

neurons. *Elife* **5**:e11181. DOI: https://doi.org/10.7554/eLife.11181

Dethier VG. 1973. Electrophysiological studies of gustation in Lepidopterous larvae.

Journal of comparative physiology **82**:103-134. DOI:

https://doi.org/10.1007/BF00696148

Engsontia P, Sangket U, Chotigeat W, Satasook C. 2014. Molecular evolution of the

odorant and gustatory receptor genes in Lepidopteran insects: implications for their

adaptation and speciation. *Journal of Molecular Evolution* **79**:21-39. DOI:

https://doi.org/10.1007/s00239-014-9633-0

Fan XB, Mo BT, Li GC, Huang LQ, Guo H, Gong XL, Wang CZ. 2022. Mutagenesis

of the odorant receptor co-receptor (Orco) reveals severe olfactory defects in the

crop pest moth *Helicoverpa armigera. BMC Biology* 20:214. DOI:

https://doi.org/10.1186/s12915-022-01411-2

Faucheux MJ. 2013. Sensillum types on the proboscis of the Lepidoptera: a review.

Annales De La Societe Entomologique De France **49**:73-90. DOI:

https://doi.org/10.1080/00379271.2013.767514

Free JB. 1993. *Insect Pollination of Crops*. Academic Press.

Freeman EG, Wisotsky Z, Dahanukar A. 2014. Detection of sweet tastants by a

conserved group of insect gustatory receptors. *PNAS* **111**:1598-1603. DOI:

https://doi.org/10.1073/pnas.1311724111

Fujii S, Yavuz A, Slone J, Jagge C, Song X, Amrein H. 2015. *Drosophila* sugar

receptors in sweet taste perception, olfaction, and internal nutrient sensing. *Current*

Biology **25**:621-627. DOI: https://doi.org/10.1016/j.cub.2014.12.058

Glendinning JI, Foley C, Loncar I, Rai M. 2009. Induced preference for host plant

chemicals in the tobacco hornworm: contribution of olfaction and taste. *Journal of*

Comparative Physiology a-Neuroethology Sensory Neural and Behavioral

Physiology **195**:591-601. DOI: https://doi.org/10.1007/s00359-009-0434-7

Guo H, et al. 2017. Expression map of a complete set of gustatory receptor genes in

chemosensory organs of *Bombyx mori*. *Insect Biochemistry and Molecular Biology*

82:74-82. DOI: https://doi.org/10.1016/j.ibmb.2017.02.001

Harari OA, Dekel A, Wintraube D, Vainer Y, Mozes-Koch R, Yakir E, Malka O,

Morin S, Bohbot JD. 2023. A sucrose-specific receptor in *Bemisia tabaci* and its

putative role in phloem feeding. *iScience* 26. DOI:

https://doi.org/10.1016/j.isci.2023.106752

Hill CA, Fox AN, Pitts RJ, Kent LB, Tan PL, Chrystal MA, Cravchik A, Collins FH,

Robertson HM, Zwiebel LJ. 2002. G protein-coupled receptors in *Anopheles*

gambiae. *Science* **298**:176-178. DOI: https://doi.org/10.1126/science.1076196

Hodgson ES, Lettvin JY, Roeder KD. 1955. Physiology of a primary chemoreceptor

unit. *Science* **122**:417-418. DOI: https://doi.org/ 10.1126/science.122.3166.417-a

Jiang XJ, Ning C, Guo H, Jia YY, Huang LQ, Qu MJ, Wang CZ. 2015. A gustatory

receptor tuned to D-fructose in antennal sensilla chaetica of *Helicoverpa armigera*.

Insect Biochemistry and Molecular Biology **60**:39-46. DOI:

https://doi.org/10.1016/j.ibmb.2015.03.002

Jiao Y, Moon SJ, Montell C. 2007. A *Drosophila* gustatory receptor required for the

responses to sucrose, glucose, and maltose identified by mRNA tagging. *PNAS*

104:14110-14115. DOI: https://doi.org/10.1073/pnas.0702421104

Jiao YC, Moon SJ, Wang XY, Ren QT, Montell C. 2008. Gr64f is required in

combination with other gustatory receptors for sugar detection in *Drosophila*.

Current Biology **18**:1797-1801. DOI: https://doi.org/10.1016/j.cub.2008.10.009

Jørgensen K, Almaas TJ, Marion-Poll F, Mustaparta H. 2007. Electrophysiological

characterization of responses from gustatory receptor neurons of sensilla chaetica in

the moth *Heliothis virescens*. *Chemical Senses* **32**:863-879. DOI:

https://doi.org/10.1093/chemse/bjm057

Jung JW, Park KW, Ahn YJ, Kwon HW. 2015. Functional characterization of sugar

receptors in the western honeybee, *Apis mellifera*. *Journal of Asia-Pacific*

Entomology **18**:19-26. DOI: https://doi.org/10.1016/j.aspen.2014.10.011

Kent LB, Walden KK, Robertson HM. 2008. The Gr family of candidate gustatory

and olfactory receptors in the yellow-fever mosquito *Aedes aegypti*. *Chemical*

Senses **33**:79-93. DOI: https://doi.org/10.1093/chemse/bjm067

Kwon JY, Dahanukar A, Weiss LA, Carlson JR. 2011. Molecular and cellular

organization of the taste system in the *Drosophila* larva. *Journal of Neuroscience*

31:15300-15309. DOI: https://doi.org/10.1523/jneurosci.3363-11.2011

Li X, Staszewski L, Xu H, Durick K, Zoller M, Adler E. 2002. Human receptors for

sweet and umami taste. *PNAS* **99**:4692-4696. DOI:

https://doi.org/10.1073/pnas.072090199

Liman ER, Zhang YV, Montell C. 2014. Peripheral coding of taste. *Neuron*

81:984-1000. DOI: https://doi.org/10.1016/j.neuron.2014.02.022

Liu XL, Sun SJ, Hou W, Zhang J, Yan Q, Dong SL. 2020. Functional characterization

of two spliced variants of fructose gustatory receptor in the diamondback moth,

Plutella xylostella. *Pesticide biochemistry and physiology* **164**:7-13. DOI:

https://doi.org/10.1016/j.pestbp.2019.12.002

Liu XL, Yan Q, Yang YL, Hou W, Miao CL, Peng YC, Dong SL. 2019. A gustatory

receptor GR8 tunes specifically to D-fructose in the common cutworm *Spodoptera*

litura. *Insects* **10**:272. DOI: https://doi.org/10.3390/insects10090272

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using

real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods* **25**:402-408. DOI:

https://doi.org/10.1006/meth.2001.1262

Maier GL, Komarov N, Meyenhofer F, Kwon JY, Sprecher SG. 2021. Taste sensing

and sugar detection mechanisms in *Drosophila* larval primary taste center. *Elife*

959 DOI: https://doi.org/10.7554/eLife.67844

960 Mang D, Mayu K, Toyama T, Yamagishi T, Sato R. 2022. BmGr4 responds to sucrose

961  and glucose and expresses in tachykinin-related peptide-secreting enteroendocrine

962  cells. *Insect Biochemistry and Molecular Biology* **150**:103858. DOI:

963  https://doi.org/10.1016/j.ibmb.2022.103858

964 Martin TL, Shields VDC. 2012. Detection of alkaloids and carbohydrates by taste

965  receptor cells of the galea of gypsy moth larvae, *Lymantria dispar* (L.).

966  *Arthropod-Plant Interactions* **6**:519-529. DOI:

967  https://doi.org/10.1007/s11829-012-9209-0

968 Mishra D, Miyamoto T, Rezenom YH, Broussard A, Yavuz A, Slone J, Russell DH,

969  Amrein H. 2013. The molecular basis of sugar sensing in *Drosophila* larvae.

970  *Current Biology* **23**:1466-1471. DOI: https://doi.org/10.1016/j.cub.2013.06.028

971 Miyamoto T, Slone J, Song XY, Amrein H. 2012. A fructose receptor functions as a

972  nutrient sensor in the *Drosophila* brain. *Cell* **151**:1113-1125. DOI:

973  https://doi.org/10.1016/j.cell.2012.10.024

974 Morinaga S, Nagata K, Ihara S, Yumita T, Niimura Y, Sato K, Touhara K. 2022.

975  Structural model for ligand binding and channel opening of an insect gustatory

976  receptor. *Journal of Biological Chemistry* **298**:102573. DOI:

977  https://doi.org/10.1016/j.jbc.2022.102573

978 Nakagawa T, Touhara K. 2013. Functional assays for insect olfactory receptors in

979  *Xenopus* oocytes. Pages 107-119 in Touhara K, ed. Pheromone Signaling Totowa,

980  NJ: Humana Press.

981 Nelson G, Hoon MA, Chandrashekar J, Zhang Y, Ryba NJ, Zuker CS. 2001.

982  Mammalian sweet taste receptors. *Cell* **106**:381-390. DOI:

983  https://doi.org/10.1016/s0092-8674(01)00451-2

984 Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A fast and

985  effective stochastic algorithm for estimating maximum-likelihood phylogenies.

Molecular Biology **32**:268-274. DOI: https://doi.org/10.1093/molbev/msu300

Nicolson SW. 1998. The importance of osmosis in nectar secretion and its

consumption by insects. *American Zoologist* **38**:418-425. DOI:

https://doi.org/10.1093/icb/38.3.418

Notredame C, Higgins DG, Heringa J. 2000. T-Coffee: A novel method for fast and

accurate multiple sequence alignment. *Journal Of Molecular Biology* 302:205-217.

DOI: https://doi.org/10.1006/jmbi.2000.4042

Pearce SL, et al. 2017. Genomic innovations, transcriptional plasticity and gene loss

underlying the evolution and divergence of two highly polyphagous and invasive

Helicoverpa pest species. *BMC biology* **15**:63. DOI:

https://doi.org/10.1186/s12915-017-0402-6

Popescu A, Couton L, Almaas TJ, Rospars JP, Wright GA, Marion-Poll F, Anton S.

2013. Function and central projections of gustatory receptor neurons on the antenna

of the noctuid moth *Spodoptera littoralis*. *Journal of Comparative Physiology*

a-Neuroethology Sensory Neural and Behavioral Physiology **199**:403-416. DOI:

https://doi.org/10.1007/s00359-013-0803-0

Robertson HM, Wanner KW. 2006. The chemoreceptor superfamily in the honey bee,

Apis mellifera: expansion of the odorant, but not gustatory, receptor family.

Genome Research **16**:1395-1403. DOI: https://doi.org/10.1101/gr.5057506

Robertson HM, Warr CG, Carlson JR. 2003. Molecular evolution of the insect

chemoreceptor gene superfamily in *Drosophila melanogaster*. *PNAS* **100**

14537-14542. DOI: https://doi.org/10.1073/pnas.2335847100

Rozewicki J, Li S, Amada KM, Standley DM, Katoh K. 2019. MAFFT-DASH:

integrated protein sequence and structural alignment. *Nucleic Acids Research*

47:W5-W10. DOI: https://doi.org/10.1093/nar/gkz342

Sato K, Tanaka K, Touhara K. 2011. Sugar-regulated cation channel formed by an

insect gustatory receptor. *PNAS* **108**:11680-11685. DOI:

https://doi.org/10.1073/pnas.1019622108

Schauer N, Zamir D, Fernie AR. 2005. Metabolic profiling of leaves and fruit of wild species tomato: a survey of the *Solanum lycopersicum* complex. *Journal Of Experimental Botany* **56**:297-307. DOI: https://doi.org/10.1093/jxb/eri057

Schnuch M, Seebauer H. 1998. Sugar cell responses to lactose and sucrose in labellar and tarsal taste hairs of *Musca domestica*. *Journal of Comparative Physiology a-Neuroethology Sensory Neural and Behavioral Physiology* **182**:767-775. DOI: https://doi.org/10.1007/s003590050221

Schoonhoven LM, van Loon JJA. 2002. An inventory of taste in caterpillars: each species its own key. *Acta Zoologica Academiae Scientiarum Hungaricae* **48**:215-263

Schoonhoven LM, van Loon JJA, Dicke M. 2005. *Insect-Plant Biology*. Oxford University Press.

Sharath Chandra G, Asokan R, Manamohan M, Krishna Kumar NK, Sita T. 2014. Evaluation of reference genes for quantitative real-time PCR normalization in cotton bollworm *Helicoverpa armigera*. *Molecular Biology (Mosk)* 48:813-822

Slone J, Daniels J, Amrein H. 2007. Sugar receptors in *Drosophila*. *Current Biology* **17**:1809-1816. DOI: https://doi.org/10.1016/j.cub.2007.09.027

Smith JJB, Mitchell BK, Rolseth BM, Whitehead AT, Albert PJ. 1990. SAPID tools - microcomputer programs for analysis of multiunit nerve recordings. *Chemical Senses* **15**:253-270. DOI: https://doi.org/10.1093/chemse/15.3.253

Takada T, Sasaki T, Sato R, Kikuta S, Inoue MN. 2018. Differential expression of a fructose receptor gene in honey bee workers according to age and behavioral role. *Archives of Insect Biochemistry and Physiology* **97**:1-9. DOI: https://doi.org/10.1002/arch.21437

Tang DL, Wang CZ, Luo L, Qin JD. 2000. Comparative study on the responses of maxillary sensilla styloconica of cotton bollworm *Helicoverpa armigera* and

Oriental tobacco budworm *H. assulta* larvae to phytochemicals. *Science China Life*

Sciences **43**:606-612. DOI: https://doi.org/10.1007/Bf02882281

van Loon JJA. 1990. Chemoreception of phenolic acids and flavonoids in larvae of

two species of *Pieris*. *Journal of Comparative Physiology a-Neuroethology Sensory*

Neural and Behavioral Physiology **166**:889-899. DOI:

https://doi.org/10.1007/bf00187336

Vansell GH. 1944. Cotton nectar in relation to bee activity and honey production.

Journal of Economic Entomology **37**:528-530. DOI:

https://doi.org/10.1093/jee/37.4.528

Waller GD, Wilson FD, Martin JH. 1981. Influence of phenotype, season, and

time-of-day on nectar production in cotton. *Crop Science* **21**:507-511. DOI:

https://doi.org/10.2135/cropsci1981.0011183X002100040007x

Xu W, Liu N, Liao Y, Anderson A. 2017. Molecular characterization of sugar taste

receptors in the cotton bollworm *Helicoverpa armigera*. *Genome* **60**:1037-1044.

DOI: https://doi.org/10.1139/gen-2017-0086

Xu W, Zhang HJ, Anderson A. 2012. A sugar gustatory receptor identified from the

foregut of cotton bollworm *Helicoverpa armigera*. *Journal of Chemical Ecology*

38:1513-1520. DOI: https://doi.org/10.1007/s10886-012-0221-8

Yarmolinsky DA, Zuker CS, Ryba NJP. 2009. Common sense about taste: from

mammals to insects. *Cell* **139**:234-244. DOI:

https://doi.org/10.1016/j.cell.2009.10.001

Zhan S, Merlin C, Boore JL, Reppert SM. 2011. The monarch butterfly genome yields

insights into long-distance migration. *Cell* **147**:1171-1185. DOI:

https://doi.org/10.1016/j.cell.2011.09.052

Zhang HJ, Faucher CP, Anderson A, Berna AZ, Trowell S, Chen QM, Xia QY, Chyb S.

2013. Comparisons of contact chemoreception and food acceptance by larvae of

polyphagous *Helicoverpa armigera* and oligophagous *Bombyx mori*. *Journal of*

Chemical Ecology **39**:1070-1080. DOI: https://doi.org/10.1007/s10886-013-0303-2

Zhang YF, van Loon JJA, Wang CZ. 2010. Tarsal taste neuron activity and proboscis

extension reflex in response to sugars and amino acids in *Helicoverpa armigera*

(Hübner). *Journal of Experimental Biology* **213**:2889-2895. DOI:

https://doi.org/10.1242/jeb.042705

Zhao GQ, Zhang Y, Hoon MA, Chandrashekar J, Erlenbach I, Ryba NJ, Zuker CS.

2003. The receptors for mammalian sweet and umami taste. *Cell* **115**:255-266. DOI:

https://doi.org/10.1016/s0092-8674(03)00844-4

1075

Figure legends

Figure 1. Electrophysiological responses of larval and adult contact

chemosensilla in *Helicoverpa armigera* to sugars. **(A)** The maxilla morphology

(left), the representative spike traces of the responses of lateral sensilla styloconica in

larval maxillary galea to eight sugars at 10 mM (middle), and quantifications of the

firing rates (right) ($n = 13$). The red arrow marks the recorded lateral sensillum

styloconicum. Scale bar represents 20 μm. **(B)** Dose–responses of lateral sensilla

styloconica in larval maxillary galea to sucrose and fucose (sucrose: $n = 8$; fucose: n

$= 6$). **(C)** The antenna tip morphology (left), the representative spike traces of the

responses of sensilla chaetica in adult antennae to eight sugars at 10 mM (middle),

and quantifications of the firing rates (right) ($n = 21$). The red arrows mark the

recorded sensilla chaetica. Scale bar represents 100 μm. **(D)** Dose–responses of

sensilla chaetica in adult antennae to sucrose, fucose, and fructose ($n = 21$). **(E)** The

fifth tarsomere morphology (left) (Zhang et al. 2010), the representative spike traces

of the responses of sensilla trichodea in adult tarsi to eight sugars at 10 mM (middle),

and quantifications of the firing rates (right) ($n = 15$). The red arrows mark the

recorded sensilla trichodea. Scale bar represents 100 μm. **(F)** Dose–responses of

sensilla trichodea in adult tarsi to sucrose, fucose, and fructose ($n = 21$). **(G)** The

proboscis tip morphology (left), the representative spike traces of the responses of

sensilla styloconica in adult proboscis to eight sugars at 10 mM (middle), and

quantifications of the firing rates (right) ($n = 21$). The red arrows mark the recorded

sensilla styoloconica. Scale bar represents 200 μm. **(H)** Dose–responses to sucrose (n

$= 21$), fucose ($n = 21$), and fructose (fructose 0 mM, $n = 18$; fructose 0.01 mM-100

mM, $n = 21$) of sensilla styloconica in adult proboscis. **(A to H)** Data are mean ±

SEM; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. **(A, C, E, G)** Data were analyzed by

independent-samples t test (compared with control). **(B, D, F, H)** Data were analyzed

by one-way ANOVA with Tukey's HSD test.

Figure 2. Behavioral responses of *Helicoverpa armigera* larvae and adults to

sugars. (A) Feeding responses and the preference index (PI) value of 5^{th} instar larvae

to eight sugars painted on the cabbage leaf discs at 10 mM in two-choice tests. ** $P <$

0.01; ns indicates no significance, $P \geq 0.05$ (paired t test, $n = 20$). **(B)** Feeding

responses and the PI value of 5^{th} instar larvae to different concentrations of sucrose

painted on the cabbage leaf discs in two-choice tests. ** $P < 0.01$, *** $P < 0.001$; ns

indicates no significance, $P \geq 0.05$ (paired t test, $n = 20$). **(C)** Proboscis extension

reflex (PER) in adult females upon antennal stimulation by eight sugars at 100 mM.

** $P < 0.01$; *** $P < 0.001$ (independent-samples t test compared with control, $n = 3$).

(D) PER in adult females upon tarsal stimulation by eight sugars at 100 mM. * $P <$

0.05; ** $P < 0.01$ (independent-samples t test, $n = 3$). **(E)** PER in adult females upon

antennal stimulation by different concentrations of sucrose, fucose, and fructose. * P

< 0.05 (one-way ANOVA with Tukey's HSD test, $n = 3$). **(F)** PER in adult females

upon tarsal stimulation by different concentrations of sucrose, fucose, and fructose (n

$= 3$). * $P < 0.05$ (one-way ANOVA with Tukey's HSD test, $n = 3$). **(A to F)** Data are

mean ± SEM. The red arrow indicates the stimulating site.

Figure 3. The phylogenetic relationship and the expression level of sugar GRs in

larval maxilla and female adult antennae, tarsi and proboscis of *Helicoverpa*

***armigera*. (A)** The phylogenetic tree of insect sugar GRs. Diptera (orange): Aaeg,

Aedes aegypti; Agam, A*nopheles gambiae*; Dmel, *Drosophila melanogaster*.

Hymenoptera (blue): Am, *Apis mellifera*. Lepidoptera (black): Bmor, *Bombyx mori*;

Dple, *Danaus plexippus*; Hmel, *Heliconius melpomene*; Pxyl, *Plutella xylostella*.

Harm (red), *Helicoverpa armigera*. Numbers above branches indicate ultrafast

bootstrap approximation (UFBoot). **(B)** Relative expression levels of sugar GRs in the

maxillary galea of 5^{th} instar larvae of *H. armigera* determined by qRT-PCR. **(C)**

Relative expression levels of sugar GRs in the female adult antennae. **(D)** Relative

expression levels of sugar GRs in female adult tarsi. **(E)** Relative expression levels of

sugar GRs in female adult proboscis. **(B to E)** Data are mean ± SEM. One-way

ANOVA was used, and different letters labeled indicate significant difference

(Tukey's HSD test, $P < 0.05$, $n = 3$).

Figure 4. The inward current responses of the *Xenopus* oocytes expressing sugar

GRs of *Helicoverpa armigera* to sugar compounds. **(A)** The representative traces of

the oocytes expressing Gr10, Gr6, and Gr10 + Gr6 to 11 sugar compounds at 100 mM.

(B) The responses of the oocytes expressing Gr10, Gr6, and Gr10 + Gr6 to sugars at

100 mM (Gr10, $n = 14$. Gr6, $n = 20$. Gr10 + Gr6, $n = 7$). **(C)** The representative trace

of the oocytes expressing Gr10 to sucrose (the upper), the representative trace of the

oocytes expressing Gr6 to sucrose, fucose, and fructose (the middle), the

representative trace of the oocytes expressing Gr10 + Gr6 to sucrose and fucose (the

lower). **(D)** The dose-responses of the oocytes expressing Gr10 to sucrose (the upper,

$n = 13$); the dose-responses of the oocytes expressing Gr6 to sucrose, fucose, and

fructose (the middle, sucrose: $n = 6$; fucose: $n = 6$; fructose: $n = 3$); the

dose-responses of the oocytes expressing Gr10 + Gr6 to sucrose and fucose (the lower,

sucrose: $n = 4$; fucose: $n = 3$). **(B and D)** Data are mean ± SEM, and were analyzed

by one-way ANOVA with Tukey's HSD test ($P < 0.05$). Different letters labeled

indicate significant differences, $* P < 0.05$. **(A to D)** Ara, arabinose; Fru, fructose;

Fuc, fucose; Gal, galactose; Glu, glucose; Lac, lactose; Mal, maltose; Man, mannose;

Suc, sucrose; Tre, trehalose; Xyl, xylose; Rin, ringer solution.

Figure 5. Establishment of *Gr10* and *Gr6* homozygous mutants (*Gr10*^{-/-} and *Gr6*^{-/-})

of *Helicoverpa armigera* via CRISPR/Cas9. (A) The cross process of obtaining

Gr10^{-/-}. **(B)** The genomic structure of *Gr10*, the single-guide RNA (sgRNA) targeting

sequence (in green), and representative chromatograms of direct sequencing of the

PCR products obtained from wild types (WT) and *Gr10*^{-/-}, in which 4-bp of the *Gr10*

sequence were deleted. **(C)** The predicted secondary structures of the Gr10 protein in

WT and the truncated Gr10 protein in *Gr10*^{-/-}. **(D)** The cross process of obtaining

Gr6^{-/-}. **(E)** The genomic structure of *Gr6*, the single-guide RNA (sgRNA) targeting

sequence (in green), and representative chromatograms of direct sequencing of the

PCR products obtained from WT and *Gr6*^{-/-}, in which 17-bp of the *Gr6* sequence were

deleted. **(F)** The predicted secondary structures of the Gr6 protein in WT and the

truncated Gr6 protein in *Gr6*^{-/-}. **(B and D)** Boxes represent exons, black lines

represent introns, the green arrowhead indicates the direction of sgRNA; the

protospacer adjacent motif (PAM) is in blue. **(C and F)** The secondary structure of

Gr10 and *Gr6* in WT, *Gr10*^{-/-} and *Gr6*^{-/-} was predicted by

https://dtu.biolib.com/DeepTMHMM, and the image was constructed by TOPO2

software (http://www.sacs.ucsf.edu/TOPO2).

Figure 6. Electrophysiological responses of larval and adult contact

chemosensilla in WT, *Gr10*^{-/-} and *Gr6*^{-/-} of *Helicoverpa armigera* to sucrose and

other compounds. (A) The representative spike traces of lateral sensilla styloconica

in larval maxillary galea. **(B)** Quantifications of the firing rates of lateral sensilla

styloconica in larval maxillary galea (mean ± SEM; WT, $n = 12$; $Gr6^{-/-}$, $n = 13$;

$Gr10^{-/-}$, $n = 13$). **(C)** The representative spike traces of sensilla chaetica in female

antennae. **(D)** Quantifications of the firing rates of sensilla chaetica in female

antennae (mean ± SEM; WT, $n = 20$. $Gr6^{-/-}$: nicotine, $n = 17$; other compounds, $n =$

18. $Gr10^{-/-}$, $n = 20$). **(E)** The representative spike traces of sensilla trichodea in female

tarsi. **(F)** Quantifications of the firing rates of sensilla trichodea in female tarsi (mean

± SEM; WT, $n = 18$; $Gr6^{-/-}$: nicotine, $n = 14$. other compounds, $n = 18$. $Gr10^{-/-}$, $n =$

18). **(G)** The representative spike traces of sensilla styloconica in female proboscis.

(H) Quantifications of the firing rates of sensilla styloconica in female proboscis

(mean ± SEM; WT: sucrose 100 mM, $n = 17$; other compounds, $n = 18$. $Gr6^{-/-}$:

sucrose 100 mM, $n = 17$; other compounds, $n = 18$. $Gr10^{-/-}$, $n = 18$). **(B, D, F, and H)**

Two-way ANOVA with post hoc Tukey's multiple comparison was used separately

for sucrose and fucose, and one-way ANOVA with Tukey's HSD test was used for

KCl, sinigrin, fructose, and nicotine. Different letters labeled indicate significant

differences ($P < 0.05$). Suc: sucrose; Fuc: fucose; Sin: sinigrin; Fru: fructose; Nic:

nicotine.

Figure 7. Behavioral responses of WT, $Gr10^{-/-}$ and $Gr6^{-/-}$ larvae and adults of

***Helicoverpa armigera* to sugars and plant tissues. (A)** Feeding area of 5th instar

larvae in two-choice tests and the PI value to 10 mM sucrose ($n = 20$). ** $P < 0.01$; ns

indicates no significance, $P \geq 0.05$ (paired t test). Suc, sucrose. **(B)** Feeding area of

5th instar larvae in two-choice tests and the PI value to 100 mM sucrose ($n = 20$). * P

< 0.05; ** $P < 0.01$; *** $P < 0.001$ (paired t test). Suc, sucrose. **(C)** Feeding amount

of 5th instar larvae on cabbage leaves, corn kernels, pea seeds, pepper fruits, and

tomato fruits in no-choice tests ($n = 20$). Data were analyzed by one-way ANOVA

with Tukey's HSD test, and different letters labeled on the data of WT, *Gr10^{-/-}* and

Gr6^{-/-} for each plant tissue indicate significant differences ($P < 0.05$). Proboscis

extension reflex (PER) in adult females upon **(D)** antennal stimulation by sucrose

concentrations ($n = 3$), **(E)** antennal stimulation by fucose concentrations ($n = 3$), **(F)**

antennal stimulation by fructose concentrations ($n = 3$), **(G)** tarsal stimulation by

sucrose concentrations ($n = 3$), **(H)** tarsal stimulation by fucose concentrations ($n = 3$),

and **(I)** tarsal stimulation by fructose concentrations ($n = 3$). **(A to I)** Data are mean ±

SEM. **(D to I)** Data were analyzed by two-way ANOVA with post hoc Tukey's

multiple comparison. * $P < 0.05$. The red arrow indicates the stimulating site.

Figure 8. Two gustatory receptors in the cotton bollworm, *Helicoverpa armigera*

mainly mediate taste sensation of sugars in the larval and adult foods

1206

A

Sugar	Control	PI value
Arabinose ns		-0.0089 ± 0.017
Fructose **		0.0605 ± 0.016
Fucose **		-0.0010 ± 0.031
Galactose ns		-0.0173 ± 0.012
Glucose ns		-0.0002 ± 0.062
Sucrose **		0.1162 ± 0.031
Trehalose ns		-0.0024 ± 0.019
Xylose ns		-0.0088 ± 0.019

B

Sucrose	Control	PI value
0.01 mM ns		-0.0681 ± 0.047
0.1 mM ns		-0.0107 ± 0.069
1 mM ns		0.0451 ± 0.031
10 mM ***		0.1188 ± 0.052
100 mM ****		0.1712 ± 0.036

C

D

E

F

A

G0   Gr10 chimeras × wild types

G1   wild types   Gr10⁺ᐟ⁻   other heterozygous

G2   wild types   Gr10⁺ᐟ⁻   in-cross   Gr10⁺ᐟ⁻

                    in-cross

G3   Gr10⁻ᐟ⁻

B

Exon1   sgRNA

Gr10   5'   3'

PAM

TACACGGCTTTCCAAAAGGGATGAAGGTTCT

WT

Gr10⁻ᐟ⁻   -4bp
TACACGGCTTTCCAAAA-----TGAAGGTTCT

C

Extracellular

Cytoplasm   WT   Gr10⁻ᐟ⁻

D

G0   Gr6 chimeras × wild types

G1   wild types   Gr6ᵀᴮᐟ⁻   other heterozygous

G2   wild types   Gr6⁺ᐟ⁻   in-cross   Gr6ᵀᴮᐟ⁻

                    in-cross

G3   Gr6⁻ᐟ⁻

E

Exon1   sgRNA

Gr6   5'   3'

PAM

ACCAACTGCGGAGATGATGCTGTTCTCCA

WT

Gr6⁻ᐟ⁻   -17bp
ACCACTGCGG---------------TTCCA

F

Extracellular

Cytoplasm   WT   Gr6⁻ᐟ⁻

A 10 mM Suc | Control | PI value
WT *** 0.120 ± 0.059
Gr10⁻ ns 0.013 ± 0.015
Gr6⁻ ** 0.117 ± 0.035

B 100 mM Suc | Control | PI value
WT *** 0.192 ± 0.044
Gr10⁻ *** 0.100 ± 0.042
Gr6⁻ *** 0.171 ± 0.046

C Feeding amount (g): Cabbage leaf, Corn, Pea, Pepper, Tomato

D Sucrose (mM) — WT, Gr10⁻, Gr6⁻

E Fucose (mM)

F Fructose (mM)

G Sucrose (mM)

H Fucose (mM)

I Fructose (mM)

Percentage of PER

Larval gustatory system

Maxilla

Extracellular

N — — C

Cytoplasm

Sugar cell highly expressing Gr10

detecting

Sucrose

Sucrose in plant tissues

Adult gustatory system

Antenna
Proboscis
Tarsus

Extracellular

N — — C

Cytoplasm

Sugar cell highly expressing Gr6

detecting

Sucrose

Fucose   Fructose

Sugars in nectar