Threatened and Priority listed *Melaleuca* species from Western Australia display high susceptibility to

Austropuccinia psidii in controlled inoculations

A. M. Martino[1], R. F. Park[2], P. A. Tobias[1]

[1]School of Life and Environmental Sciences, The University of Sydney, Camperdown NSW 2006,

Australia, [2]Plant Breeding Institute, University of Sydney, Narellan NSW 2567, Australia

Alyssa M Martino[1] alyssa.martino@sydney.edu.au ORCID: 0000-0002-5062-3578

Robert F Park[2] robert.park@sydney.edu.au ORCID: 0000-0002-9145-5371

Peri A Tobias[1] peri.tobias@sydney.edu.au ORCID: 0000-0002-5645-0161

Threatened and Priority listed *Melaleuca* species from Western Australia display high susceptibility to

Austropuccinia psidii in controlled inoculations

Abstract

Austropuccinia psidii causes rust disease on species within the family Myrtaceae and was first

detected in Australia in 2010, with the first detection in Western Australia in 2022. While species

within the genus *Melaleuca* from Eastern Australia show variable responses to the pathogen, little is

known of the response of species from Western Australia. This study established that 13 previously

unscreened species of *Melaleuca*, including Threatened and Priority species that were grown from

seeds sourced from Western Australian populations, were susceptible to the pandemic strain of the

pathogen. The proportion of highly susceptible plants within a single species ranged from 2% – 94%,

with several species displaying highly variable levels of resistance to *A. psidii*. These results highlight

the importance of disease screening and may direct conservation efforts.

Keywords

Austropuccinia psidii - myrtle rust – Western Australia - *Melaleuca*

Introduction

Austropuccinia psidii, formerly *Puccinia psidii* (Beenken 2017), is a rust fungus and the causal agent of

the disease myrtle rust which impacts species within the family Myrtaceae. Originating in Brazil, the

first detection of the pathogen in Australia was in 2010 and it has since spread to all states and

territories except South Australia (Carnegie et al. 2010; Carnegie and Lidbetter 2012; Westaway

2016; Department of Natural Resources and Environment Tasmania 2020; Agriculture Victoria 2022;

The Department of Primary Industries and Regional Development 2022a). The most recent detection

within Australia was in the Kimberley region of Western Australia (WA), where infection was

observed on two *Melaleuca* species near the Northern Territory border (The Department of Primary

Industries and Regional Development 2022b).

Austropuccinia psidii infects the young and expanding tissues of susceptible hosts, including the

leaves, stems, petioles, and reproductive and seed-bearing structures. In susceptible species, yellow

urediniospores appear on the infected surfaces, which may be followed by other symptoms such as

leaf distortion and defoliation (Pegg et al. 2014). In species with no resistance to *A. psidii*, repeated

infections may lead to tree death as a result of defoliation, and impact reproduction through

infection of reproductive and seed-bearing structures (Carnegie et al. 2016). In Australia, *A. psidii* has

caused the near extinction of several rainforest understory species including *Rhodamnia rubescens*

and *Rhodomyrtus psidioides* (Pegg et al. 2014; Carnegie et al. 2016; Environment Protection and

Biodiversity Conservation Act, 1999), and could be potentially devastating for other keystone species

including *Melaleuca quinquenervia* (Pegg et al. 2018).

Melaleuca is the third largest genus within the family Myrtaceae, comprising over 200 species (Ryan

2016) that are adapted to a range of habitats (Naidu et al. 2000). Although well adapted, changing

conditions as a result of climate change are contributing to the decline of *Melaleuca* species in

Australia (Saintilan et al. 2019). An increased threat is placed on these species by *A. psidii*, with

several *Melaleuca* species found to be highly susceptible to the pathogen under field conditions and

in controlled inoculations (Carnegie and Lidbetter 2012; Morin et al. 2012; Pegg et al. 2014, 2018;

Berthon et al. 2019; Martino et al. 2022). Further, climatic modelling predicts changes in climatic

suitability for the pathogen as a consequence of climate change, with increased suitability in areas of

NSW, TAS, VIC, and WA (Berthon et al. 2018).

WA is rich in *Melaleuca* species, with the greatest diversity and highest level of endemism located

within the South-West region of the state, with up to 72 *Melaleuca* species per 100km^2 and

endemism scores of up to 9.9 (Brophy et al. 2013). Many of these species are valued for their

important ecological, cultural, and economic roles (Brophy et al. 2013). In the absence of the

pathogen in the many parts of WA, the vulnerability of many *Melaleuca* species remains unknown.

With the arrival of *A. psidii* into WA and high susceptibility of several *Melaleuca* species, there is an

urgent need to expand current disease screening of WA species to aid pre-emptive conservation and

monitoring efforts. Here, we investigated the response of 13 previously untested *Melaleuca* species

to controlled inoculation with *A. psidii*. Using seed sourced from populations in areas climatically

suited to *A. psidii* (Berthon et al. 2018), we aimed to determine the risk the pathogen may pose in

the natural environment.

Materials and methods

Species selection

To determine the response of selected *Melaleuca* species from WA to *A. psidii*, seed was obtained

from the Department of Biodiversity, Conservation and Attractions (DBCA) Kings Park and Kensington

seed banks. Seed was obtained for species listed under the Biodiversity Conservation (BC) Act

(2016) as Threatened, including critically endangered, endangered, or vulnerable species

(*Biodiversity Conservation Act* 2016 (WA) s 19), and species listed as Priority on DBCA's priority flora

list (Department of Biodiversity, Conservation and Attractions 2017). While not designated under the

BC Act, Priority listed species may be threatened but lack sufficient survey data to list under the Act.

Seed from Priority listed species for this work include; *Melaleuca dempta*, *Melaleuca incana* ssp.

gingilup, *Melaleuca penicula*, *Melaleuca similis*, and *Melaleuca sophisma*. Seed was also obtained for

the Threatened (endangered) listed species *Melaleuca* sp. *Wanneroo*. Seed was also obtained from

species listed as Not Threatened (*Biodiversity Conservation Act* 2016 (WA) s 19) and included

Melaleuca acutifolia, *Melaleuca argentea*, *Melaleuca cajuputi* ssp. *cajuputi*, *Melaleuca fulgens* ssp.

fulgens, *Melaleuca lanceolata*, *Melaleuca lateralis*, and *Melaleuca viminea* ssp. *appressa*. For each

species, seed was collected from multiple trees with co-ordinates obtained and mapped (Figure 1).

88 Figure 1. Seed collection sites by coordinate or nearest town for (A) *Melaleuca cajuputi* ssp*. cajuputi*, (B)

89 *Melaleuca argentea* (red), *Melaleuca acutifolia* (plum), (C) *Melaleuca fulgens* ssp. *fulgens* (mustard), *Melaleuca*

90 sp. *Wanneroo* (light green), *Melaleuca incana* ssp. *gingilup* (yellow), *Melaleuca penicula* (dark green),

91 *Melaleuca sophisma* (light blue), *Melaleuca lateralis* (green), *Melaleuca similis* (purple), *Melaleuca viminea* ssp.

92 *appressa* (dark blue), *Melaleuca dempta* (orange), and *Melaleuca lanceolata* (olive). Seed was collected from

93 multiple parents at each site. Image generated in Google My Maps and interactive map is viewable at

94 https://tinyurl.com/zvffxccd.

96 Seed germination and plant growth

97 Seeds were sown into perforated trays containing a mix of 2:1:1 peat, coconut coir, and perlite

98 supplemented with Osmocote® Native Controlled Release Fertiliser then covered with a fine coating

99 of vermiculite. Perforated trays were placed into solid trays filled with 1 cm of water, every 3-4 days

100 allowing for periods of drying to promote root growth. Seeds were germinated under natural light in

101 a climate-controlled greenhouse set at 24°C/20°C day-time/night-time temperature on a 12 hour

102 cycle. Germinated seedlings were transplanted into 85 mL pots (5 cm diameter and depth)

103 containing a mix of 2:1:1 Osmocote® Native Premium Potting Mix, peat, and perlite supplemented

104 with Osmocote® Native Controlled Release Fertiliser then placed on capillary mats. Seedlings were

105 grown under the same light and temperature conditions as for germination.

107 Seedling inoculations

108 For all species, we inoculated seedlings approximately 4 months post germination at the Plant

109 Breeding Institute at the University of Sydney (Cobbitty, NSW) alongside four highly susceptible

110 *Syzygium jambos* plants as positive controls. Approximately 50 mg of *A. psidii* urediniospores from a

111 greenhouse increased single pustule isolate (accession 622) (Sandhu and Park 2013) was added to 50

112 mL of Isopar® for a final concentration of 1 mg spores/mL. Seedlings were inoculated with the

113 suspension using an aerosol sprayer and relocated to a humid incubation chamber for 24 hours at

114 20°C. After incubation, seedlings were transferred to a greenhouse with the temperature set to

115 24°C/20°C day-time/night-time temperature on a 12-hour cycle under natural light.

117 Disease susceptibility scoring

118 Host response to *A. psidii* inoculation was scored using a 1 – 5 scoring system based on Morin et al.

119 (2012) and adapted for disease scoring on *Melaleuca* species (this study) where 1 indicates

120 completely resistant or no visible response and 5 indicates highly susceptible (Table 1). *Syzygium*

jambos was scored as score 5 for each round of inoculation indicating successful inoculation. As

inoculations were carried out in winter under shorter day-length conditions, disease symptoms were

slower to develop than in previous screening (Martino et al. 2022). Plants were left for 16 days prior

to scoring to allow for complete development of plant disease symptoms.

Table 1. Disease scoring scale adapted from Morin et al. (2012) to score *Melaleuca* species for their response to
Austropuccinia psidii in controlled inoculations. Scoring was based on the disease symptoms on *Melaleuca*
quinquenervia scored at 14-days post inoculation with greenhouse increased single pustule isolate (accession
622) *Austropuccinia psidii* urediniospores (Sandhu and Park 2013)

Infection Score	Disease Rating	Infection symptoms based on Morin et al., (2012)	Infection symptoms adapted from *Melaleuca quinquenervia* for other *Melaleuca* sp.	Representative leaf image
1	CR	No visible symptoms attributable to rust infection	No visible symptoms attributable to rust infection	
2	HR	Chlorotic, purplish, or necrotic spots or blotches	Chlorotic or necrotic spots or blotches	
3	MS	Purplish or necrotic flecks with underdeveloped uredinia. Pin sized uredinia, limited sporulation	Necrotic flecks with limited sporulation	
4	S	Fully developed uredinia with or without purplish halos that cover less than 25% of the leaf and abundant sporulation	Abundant sporulation with necrotic halos. Spores may appear on leaves, stems, and/or petioles	
5	HS	Fully developed uredinia with or without purplish halos that cover more than 25% of the leaf and abundant sporulation	Abundant sporulation with no visible necrosis. Spores may appear on leaves, stems, and/or petioles	

CR = Completely resistant, HR = Hypersensitive response, MS = Moderately susceptible, S = Susceptible, HS = Highly susceptible

Imaging

All images were captured using an Olympus OM-5 fitted with an Olympus M. Zuiko Premium 60mm

f/2.8 Macro Lens. Raw images were processed using Adobe Photoshop 2023.

Results

Within 16 days post inoculation, symptoms had developed on the highly susceptible *S. jambos*

control plants (Figure 2 A-B). Using the scoring system adapted for *Melaleuca* species (this study,

Table 1), disease scores across all plants ranged from completely resistant (score 1) to highly

susceptible (score 5) (Supplementary Figures 1 – 13). The proportion of highly susceptible plants

within a single species ranged from 2% for *M. sophisma,* to 94% for *M. lateralis*. Eight of the species

had individuals that were either completely resistant (score 1) or highly susceptible (score 4 or 5) to

A. psidii (Table 2). This binary response was observed for *M. lateralis*, with 94% of plants highly

susceptible and 6% with no observable symptoms and for *M. sophisma*, with 2% of plants highly

susceptible and the remaining 98% with no observable symptoms (Table 2). Only three of the 13

species tested; *M. argentea*, *M. cajupti* spp. *cajupti*, and *M. incana* ssp. *gingilup,* had representative

plants from each disease score (Table 2). Urediniospores were observed on the leaves of all highly

susceptible plants (Figure 2 C-O), as well as infection on stems and petioles on all species except for

M. sophisma and *M. incana* ssp. *gingilup*.

Figure 2. Representative highly susceptible disease symptoms on (A – B) *Syzygium jambos* positive control, (C) *Melaleuca acutifolia*, (D) *Melaleuca argentea*, (E) *Melaleuca cajuputi* ssp. *cajuputi*, (F) *Melaleuca dempta*, (G) *Melaleuca fulgens* ssp. *fulgens*, (H) *Melaleuca incana* ssp. *gingilup*, (I) *Melaleuca lanceolata*, (J) *Melaleuca lateralis*, (K) *Melaleuca penicula*, (L) *Melaleuca similis*, (M) *Melaleuca sophisma*, (N) *Melaleuca* sp. *Wanneroo*, and (O) *Melaleuca viminea* ssp. *appressa*. Scale bar = 0.5 cm.

156   Table 2. Disease scoring, based on Morin et al. (2012) and adapted for *Melaleuca* species (this study), of

157   controlled inoculation of *Austropuccinia psidii* of Threatened (*Biodiversity Conservation Act* 2016 (WA) s 19)

158   and Priority (Department of Biodiversity, Conservation and Attractions 2017) listed *Melaleuca* species included

159   total number of plants scored and the percentage of plants observed in each disease scoring category

Species name	Listing under Biodiversity Conservation Act 2016 or DBCA Priority Flora List	Total Number of Plants Scored	Disease Score (% of Total Plants)				
			1	2	3	4	5
Melaleuca acutifolia	NT	20	10	0	0	20	70
Melaleuca argentea	NT	52	44	23	4	4	25
Melaleuca cajuputi ssp. cajuputi	NT	54	28	15	13	35	9
Melaleuca dempta	P3	11	9	0	0	0	91
Melaleuca fulgens ssp. *fulgens*	NT	27	18	0	18	7	57
Melaleuca incana ssp. *gingilup*	P2	41	32	5	5	19	39
Melaleuca lanceolata	NT	27	15	0	18	26	41
Melaleuca lateralis	NT	31	6	0	0	0	94
Melaleuca penicula	P4	32	66	0	0	6	28
Melaleuca similis	P1	45	24	0	0	0	76
Melaleuca sophisma	P1	55	98	0	0	0	2
Melaleuca sp. Wanneroo	EN	80	14	0	0	0	86
Melaleuca viminea ssp. *appressa*	P2	25	28	0	0	4	68

EN = Endangered (Threatened species considered to be "facing a very high risk of extinction in the wild in the

near future, as determined in accordance with criteria set out in the ministerial guidelines"), NT = Not

Threatened, P1 - 3 = Priority 1 – 3 (Poorly-known species with conservation threat highest for Priority 1), P4 =

Priority 4 (Rare, Near Threatened and other species in need of monitoring).

Discussion

Here, we investigated host response to *A. psidii* in 13 previously unscreened *Melaleuca* species from

a range of geographic locations in Western Australia, revealing varying proportions of highly

susceptible plants within and between species. The broad-leaved paperbark species included in this

study, *M. cajuputi* ssp. *cajuputi* and *M. argentea*, both displayed variability in response to *A. psidii*

with plants displaying symptoms in each disease scoring category. This has previously been shown

for other broad-leaved species including *M. quinquenervia*, *M. viridiflora*, and *M. leucadendra* (Pegg

et al. 2018; Martino et al. 2022). Pegg et al. (2018) assessed the proportion of resistant *M. viridiflora*

from two provenances in WA determining 22-23 % of seedlings were resistant to *A. psidii*. The same

study assessed *M. leucadendra* seedlings from three provenances in WA determining 1- 53%

resistant seedlings, while a separate study assessing a population from the Wunaamin Conservation

Park in WA determined 30% of plants to be resistant to *A. psidii* (Martino et al. 2022). These results

indicate variability in host response to the pathogen between populations of broad-leaved

paperbarks. As the *M. cajuputi* ssp. *cajuputi* and *M. argentea* screened in this study were from grown

from seed collected from a single provenance, further studies should be conducted to determine

variation in host response between populations. Such information may be informative to shed light

on the forces driving differences in disease resistance between populations and indicating that they

may be useful for differential pathotype trials going forward.

Unlike the broad-leaved paperbarks, most species tested in this study displayed little variability in

response to the pathogen. This difference may be explained by the geographic distribution

differences of these species. For the broad-leaved paperbark species screened in this and in previous

studies, populations are numerous, and broadly distributed across large geographic regions of

Australia (Brophy et al. 2013). This distribution pattern is also true for *M. fulgens* ssp. *fulgens* and *M.*

lanceolata (Western Australian Herbarium) which both display similar variability in response to the

pathogen as the broad-leaved species. Conversely for *M*. *dempta*, *M*. *penicula*, *M*. *similis*, *M*.

sophisma, *M*. sp. *Wanneroo,* and *M*. *viminea* ssp. *appressa* where populations are geographically

sparse (Western Australian Herbarium), all display low variability in pathogen response. As

Melaleuca species are predominantly outcrossing (Quang Tan 2008; Baskorowati et al. 2010; Brophy

et al. 2013; Kartikawati et al. 2021), these differences may be explained by reductions in gene flow

within small, isolated populations, resulting in reduced genetic diversity within populations.

Of particular interest is the high proportion of resistant *M. sophisma* plants observed, with only 2%

of total plants susceptible to *A. psidii*. The remaining 98% of plants displayed no observable

symptoms or hypersensitive response, potentially indicating preformed resistance mechanisms. The

lack of a hypersensitive response has been observed in other Myrtaceae species inoculated with *A.*

psidii, including several *Eucalyptus* species (Dos Santos et al. 2019). In species with no observable

symptoms post inoculation, *A. psidii* was not detected within leaf tissues as determined by qPCR (Dos

Santos et al. 2019). The results indicated that the leaves were not colonised by the pathogen, with

the tested hypothesis that chemical compounds within cuticular waxes provide preformed resistance

in these species (Dos Santos et al. 2019). Leaf epidermal appendages have also been implicated in

contributing to responses to the pathogen with studies correlating rust susceptibility with increased

trichome density (Wang et al. 2020; Varma et al. 2023). Here, the suggestion is that trichomes

facilitate increased adherence of spores to the leaf surface. The lack of a hypersensitive response in

98% of unaffected *M. sophisma* plants may indicate reduced urediniospores adherence to the leaf

surface owing to the absence of trichomes, or the inability of *A. psidii* to penetrate or colonise the

leaves of this species owing to cuticular waxes. As many of these species remain poorly

characterised, histological analyses during rust infection may shed light on preformed resistance

mechanisms on these species.

Correlating disease responses in our greenhouse seedling tests with field responses will be important

in defining potential risks to these species. We were encouraged by the presence of some resistant

individuals within some of the listed Priority species. The results highlight the importance of

continued disease screening to determine the vulnerability of individual Myrtaceae species to *A.*

psidii. The identification of species with high susceptibility to the pathogen will be useful to inform

disease surveillance in the natural environment and to direct conservation efforts such as seed

collection.

Acknowledgements

This work was funded by the Australian Research Council under linkage project LP190100093 and

AMM by the Australian Government Research Training Program. We thank the Department of

Biodiversity, Conservation and Attractions (Kings Park and Kensington) for suppling the seed for this

work and Bob Makinson for the introductions that facilitated this work.

Conflict of interest

The authors declare no conflict of interest in the reporting of these results.

References

Agriculture Victoria (2022) About myrtle rust. https://agriculture.vic.gov.au/biosecurity/plant-
233   diseases/shrub-and-tree-diseases/myrtle-rust/about-myrtle-rust. Accessed 14 Jul 2022

234   Baskorowati L, Moncur MW, Cunningham SA, et al (2010) Reproductive biology of *Melaleuca*
235   *alternifolia* (Myrtaceae) 2. Incompatibility and pollen transfer in relation to the breeding
236   system. Aust J Bot 58:384–391. https://doi.org/10.1071/BT10036

237   Beenken L (2017) *Austropuccinia*: a new genus name for the myrtle rust *Puccinia psidii* placed within
238   the redefined family Sphaerophragmiaceae (Pucciniales). Phytotaxa 297:53-61-53–61.
239   https://doi.org/10.11646/PHYTOTAXA.297.1.5

240   Berthon K, Esperon-Rodrigueza M, Beaumonta LJ, et al (2018) Assessment and prioritisation of plant
241   species at risk from myrtle rust (*Austropuccinia psidii*) under current and future climates in
242   Australia. Biological Conservation journal 7:154–162.
243   https://doi.org/10.1371/journal.pone.0035434

244   Berthon KA, Fernandez Winzer L, Sandhu K, et al (2019) Endangered species face an extra threat:
245   susceptibility to the invasive pathogen *Austropuccinia psidii* (myrtle rust) in Australia.
246   Australasian Plant Pathology 48:385–393. https://doi.org/10.1007/s13313-019-00640-4

247   Brophy JJ, Craven LA, Doran JC (2013) *Melaleucas*: their botany, essential oils and uses. Australian
248   Centre for International Agricultural Research, Canberra

249   Carnegie AJ, Kathuria A, Pegg GS, et al (2016) Impact of the invasive rust *Puccinia psidii* (myrtle rust)
250   on native Myrtaceae in natural ecosystems in Australia. Biological Invasions 18:127–144.
251   https://doi.org/10.1007/s10530-015-0996-y

252   Carnegie AJ, Lidbetter JR (2012) Rapidly expanding host range for *Puccinia psidii* sensu lato in
253   Australia. Australasian Plant Pathology 41:13–29. https://doi.org/10.1007/s13313-011-0082-
254   6

255   Carnegie AJ, Lidbetter JR, Walker J, et al (2010) *Uredo rangelii*, a taxon in the guava rust complex,
256   newly recorded on Myrtaceae in Australia. Australasian Plant Pathology 39:463–466.
257   https://doi.org/10.1071/AP10102

258   Department of Biodiversity, Conservation and Attractions (2017) Threatened and Priority Flora
259   (DBCA-036). Government Printer for the State of Western Australia.
260   https://www.dbca.wa.gov.au/sites/default/files/2023-
261   10/Government%20Gazette%20135%20of%202023.pdf. Accessed 10 June 2023

262   Biodiversity Conservation Act 2016 (WA). Department of Biodiversity, Conservation and Attractions.
263   https://www.legislation.wa.gov.au/legislation/statutes.nsf/law_a147120.html

264   Department of Natural Resources and Environment Tasmania (2020) Plant species affected by myrtle
265   rust in Tasmania. https://nre.tas.gov.au/biosecurity-tasmania/plant-biosecurity/pests-and-
266   diseases/myrtle-rust/plant-species-affected-by-myrtle-rust-in-tasmania. Accessed 14 Jul
267   2022

268   Dos Santos IB, Lopes M da S, Bini AP, et al (2019) The *Eucalyptus* cuticular Waxes contribute in
269   preformed defense against *Austropuccinia psidii*. Frontiers in Plant Science 9:.
270   https://doi.org/10.3389/fpls.2018.01978

Environment Protection and Biodiversity Conservation Act 1999. Department of Climate Change,
Energy, the Environment and Water. https://www.legislation.gov.au/Details/C2016C00777

Kartikawati NK, Rimbawanto A, Na'iem M, et al (2021) Pollen dispersal and genetic structure in a
cajuput (*Melaleuca cajuputi* subsp. *cajuputi*) seed orchard in Yogyakarta, Indonesia.
Australian Forestry 84:82–90. https://doi.org/10.1080/00049158.2021.1911079

Martino AM, Park RF, Tobias PA (2022) Three species of *Melaleuca* from Western Australia are highly
susceptible to *Austropuccinia psidii* in controlled inoculations. Australasian Plant Disease
Notes 17:1–4. https://doi.org/10.1007/S13314-022-00476-W/FIGURES/2

Morin L, Aveyard R, Lidbetter JR, Wilson PG (2012) Investigating the host-range of the rust fungus
Puccinia psidii sensu lato across tribes of the family Myrtaceae present in Australia. PLoS
ONE 7:1–7. https://doi.org/10.1371/journal.pone.0035434

Naidu BP, Paleg LG, Jones GP (2000) Accumulation of proline analogues and adaptation of *Melaleuca*
species to diverse environments in Australia. Aust J Bot 48:611–620.
https://doi.org/10.1071/bt99059

Pegg GS, Giblin FR, McTaggart AR, et al (2014) *Puccinia psidii* in Queensland, Australia: disease
symptoms, distribution and impact. Plant Pathology 63:1005–1021.
https://doi.org/10.1007/s10530-015-0996-y

Pegg GS, Lee DJ, Carnegie AJ (2018) Predicting impact of *Austropuccinia psidii* on populations of
broad leaved *Melaleuca* species in Australia. Australasian Plant Pathology 47:421–430.
https://doi.org/10.1007/s13313-018-0574-8

Quang Tan N (2008) Pollination ecology of *Melaleuca cajuputi*, *Nypa fruticans* and their flower
visitors. Journal of Apicultural Research 47:10–16.
https://doi.org/10.1080/00218839.2008.11101417

Ryan M (2016) Australian forest profiles: *Melaleuca.* Australian Bureau of Agricultural and Resource
Economics and Sciences.
https://www.agriculture.gov.au/sites/default/files/abares/forestsaustralia/publishingimages
/forest%20profiles%202019/melaleuca/AusForProf_2019_Melaleuca_v.1.0.0.pdf. Accessed
12 Jan 2023_

Saintilan N, Rogers K, Kelleway JJ, et al (2019) Climate Change Impacts on the Coastal Wetlands of
Australia. Wetlands 39:1145–1154. https://doi.org/10.1007/s13157-018-1016-7

Sandhu KS, Park RF (2013) Genetic basis of pathogenicity in *Uredo rangelii. 10.13140/2.1.1965.6000_*

The Department of Primary Industries and Regional Development (2022a) Myrtle rust: Biosecurity
alert. https://www.agric.wa.gov.au/plant-biosecurity/myrtle-rust-threat-western-
australia?page=0%2C1. Accessed 14 Jul 2022

The Department of Primary Industries and Regional Development (2022b) Myrtle rust confirmed in
the Kimberley. https://www.agric.wa.gov.au/news/media-releases/myrtle-rust-confirmed-
kimberley. Accessed 20 Jul 2023

Varma PK, Chandrasekhar V, Charumati M, et al (2023) Correlation of leaf trichome density and
stomatal parameters of some commercial sugarcane genotypes to orange rust incidence and
severity. Indian Phytopathology 76:641–646. https://doi.org/10.1007/s42360-023-00624-x

Wang Y, Zeng J, Xia X, et al (2020) Comparative analysis of leaf trichomes, epidermal wax and
312   defense enzymes activities in response to *Puccinia horiana* in *Chrysanthemum* and
313   <i>Ajania<///i> species. Horticultural Plant Journal 6:191–198.
314   https://doi.org/10.1016/j.hpj.2020.03.006

315   Westaway JO (2016) The pathogen Myrtle Rust (*Puccinia psidii*) in the Northern Territory: First
316   detection, new host and potential impacts. Northern Territory Naturalist 27:13–28.
317   https://doi.org/10.3316/INFORMIT.426544422268311

318   Western Australian Herbarium (2023). Florabase-the Western Australian Flora. Department of
319   Biodiversity, Conservation and Attractions.
320   https://florabase.dbca.wa.gov.au/browse/profile/5921. Accessed 15 Nov 2023

321