Screening of Threatened and Priority listed *Melaleuca* species from Western Australia reveals high
susceptibility to *Austropuccinia psidii* in controlled inoculations

A. M. Martino[1], R. F. Park[2], P. A. Tobias[1]

[1]School of Life and Environmental Sciences, The University of Sydney, Camperdown NSW 2006,
Australia, [2]Plant Breeding Institute, University of Sydney, Narellan NSW 2567, Australia

Alyssa M Martino[1] alyssa.martino@sydney.edu.au ORCID: 0000-0002-5062-3578
Robert F Park[2] robert.park@sydney.edu.au ORCID: 0000-0002-9145-5371
Peri A Tobias[1] peri.tobias@sydney.edu.au ORCID: 0000-0002-5645-0161

Screening of Threatened and Priority listed *Melaleuca* species from Western Australia reveals high
susceptibility to *Austropuccinia psidii* in controlled inoculations

Abstract

Austropuccinia psidii causes rust disease on species within the family Myrtaceae and was first

detected in Northern Western Australia in 2022. While species within the genus *Melaleuca* from

Eastern Australia show variable responses to the pathogen, little is known of the response of species

from Western Australia. This study established that 13 previously unscreened species of *Melaleuca*,

including Threatened and Priority species that were grown from seeds sourced from Western

Australian populations were susceptible to the pandemic strain of the pathogen.

Keywords

Austropuccinia psidii - myrtle rust – Western Australia - *Melaleuca* species

Austropuccinia psidii, formerly *Puccinia psidii* (Beenken 2017), is the causal agent of the disease

myrtle rust on many species within the family Myrtaceae. Since the first detection of the pathogen in

Australia in 2010, *A. psidii* has spread to all states and territories aside from South Australia (Carnegie

et al. 2010; Carnegie and Lidbetter 2012; Westaway 2016; Department of Natural Resources and

Environment Tasmania 2020; Agriculture Victoria 2022; The Department of Primary Industries and

Regional Development 2022a). The most recent detection within Australia was in the Kimberley

region of Western Australia (WA), where infection was observed on two *Melaleuca* species near the

Northern Territory border (The Department of Primary Industries and Regional Development 2022b).

Melaleuca is the third largest genus within the Myrtaceae, comprising over 200 species (Ryan 2016).

Although adapted to a range of habitats, changing conditions as a result of climate change are

contributing to the decline of *Melaleuca* species in Australia (Saintilan et al. 2019). An increased

threat is placed on these species by *A. psidii*, with several *Melaleuca* species found to be highly

susceptible to the pathogen under field conditions and in controlled inoculations (Carnegie and

Lidbetter 2012; Morin et al. 2012; Pegg et al. 2014, 2018; Berthon et al. 2019; Martino et al. 2022).

WA is rich in *Melaleuca* species, with the greatest diversity and highest level of endemism located

within the South-West region of the state (Brophy et al. 2013). Many of these species are valued for

their important ecological, cultural, and economic roles (Brophy et al. 2013). In the absence of the

pathogen in the many parts of WA, the vulnerability of many *Melaleuca* species remains unknown.

With the arrival of *A. psidii* into WA and high susceptibility of several *Melaleuca* species, there is an

urgent need to expand current disease screening of WA species to aid pre-emptive conservation and

monitoring efforts.

To determine the susceptibility of selected *Melaleuca* species from WA in response to *A. psidii*, seed

was obtained from the Department of Biodiversity, Conservation and Attractions (DBCA) Kings Park

and Kensington seed banks. Seed was obtained for species listed under the Biodiversity Conservation

(BC) Act (2016) as Threatened, including critically endangered, endangered, or vulnerable species

(*Biodiversity Conservation Act* 2016 (WA) s 19), and species listed as Priority on DBCA's priority flora

list (Department of Biodiversity, Conservation and Attractions 2017). While not designated under the

BC Act, Priority listed species may be threatened but lack sufficient survey data to list under the Act.

Seed from Priority listed species for this work include; *Melaleuca dempta*, *Melaleuca incana* ssp.

gingilup, *Melaleuca penicula*, *Melaleuca similis*, and *Melaleuca sophisma*. Seed was also obtained for

the Threatened (endangered) listed species *Melaleuca* sp. *Wanneroo*. Seed was also obtained from

species listed as Not Threatened (*Biodiversity Conservation Act* 2016 (WA) s 19) and included

Melaleuca acutifolia, *Melaleuca argentea*, *Melaleuca cajuputi* ssp. *cajuputi*, *Melaleuca fulgens* ssp.

fulgens, *Melaleuca lanceolata*, *Melaleuca lateralis*, and *Melaleuca viminea* ssp. *appressa*. For each

species, seed was collected from multiple trees with co-ordinates obtained and mapped (Figure 1).

Figure 1. Seed collection sites by coordinate or nearest town for (A) *Melaleuca cajuputi* ssp*. cajuputi*, (B)
Melaleuca cardiophylla (brown), *Melaleuca argentea* (red), *Melaleuca acutifolia* (plum), (C) *Melaleuca fulgens*
ssp. *fulgens* (mustard), *Melaleuca* sp. *Wanneroo* (light green), *Melaleuca incana* ssp. *gingilup* (yellow),
Melaleuca penicula (dark green), *Melaleuca sophisma* (light blue), *Melaleuca lateralis* (green), *Melaleuca*
similis (purple), *Melaleuca viminea* ssp. *appressa* (dark blue), *Melaleuca dempta* (orange), and *Melaleuca*
lanceolata (olive). Seed was collected from multiple parents at each site. Image generated in Google My Maps
and viewable at https://tinyurl.com/zvffxccd.

Seeds were sown into perforated trays containing a mix of 2:1:1 peat, coconut coir, and perlite
supplemented with Osmocote® Native Controlled Release Fertiliser then covered with a fine coating
of vermiculite. Perforated trays were placed into solid trays filled with 1 cm of water, every 3-4 days
allowing for periods of drying to promote root growth. Seeds were germinated under natural light in

a climate-controlled greenhouse set at 24°C/20°C day-time/night-time temperature on a 12 hour

cycle. Germinated seedlings were transplanted into 85 mL pots (5 cm diameter and depth)

containing a mix of 2:1:1 Osmocote® Native Premium Potting Mix, peat, and perlite supplemented

with Osmocote® Native Controlled Release Fertiliser then placed on capillary mats. Seedlings were

grown under the same light and temperature conditions as for germination.

For all species, we inoculated seedlings approximately four months post germination at the Plant

Breeding Institute at the University of Sydney (Cobbitty, NSW) alongside four highly susceptible

Syzygium jambos plants as positive controls. Approximately 50 mg of *A. psidii* urediniospores from a

glasshouse increased single pustule isolate (accession 622) (Sandhu and Park 2013) was added to 50

mL of Isopar® for a final concentration of 1 mg spores/mL. Seedlings were inoculated with the

suspension using an aerosol sprayer and relocated to a humid incubation chamber for 24 hours at

20°C. After incubation, seedlings were transferred to a greenhouse with the temperature set to

24°C/20°C day-time/night-time temperature on a 12-hour cycle under natural light. As inoculations

were carried out in winter under shorter day-length conditions, plants were left for 16 days prior to

scoring to allow for development of plant disease symptoms.

Within 16 days post inoculation, symptoms had developed on the highly susceptible control plants

(Figure 2 A-B). Plants were scored using the system developed by Morin et al. (2012), ranging from

completely resistant (score 1) to highly susceptible (score 5). The proportion of highly susceptible

plants within a single species ranged from 2% – 94% with over half of the species displaying over 50%

of total plants highly susceptible to *A. psidii* (Table 1). The highest level of susceptibility was observed

in *M. lateralis* with 94% of plants scored as highly susceptible and 6% presenting with no observable

symptoms (Table 1). The lowest level of susceptibility was recorded in *M. sophisma*, with 2% of

plants scored as highly susceptible and the remaining 98% presenting with no observable symptoms

(Table 1). Urediniospores were observed on the leaves of all highly susceptible plants (Figure 2 C-O),

as well as infection on stems and petioles on all species aside from *M. sophisma* and *M. incana* ssp.

gingilup.

Figure 2. Representative highly susceptible disease symptoms on (A – B) *Syzygium jambos* positive control, (C) *Melaleuca acutifolia*, (D) *Melaleuca argentea*, (E) *Melaleuca cajuputi* ssp. *cajuputi*, (F) *Melaleuca dempta*, (G) *Melaleuca fulgens* ssp. *fulgens*, (H) *Melaleuca incana* ssp. *gingilup*, (I) *Melaleuca lanceolata*, (J) *Melaleuca lateralis*, (K) *Melaleuca penicula*, (L) *Melaleuca similis*, (M) *Melaleuca sophisma*, (N) *Melaleuca* sp. *Wanneroo*, and (O) *Melaleuca viminea* ssp. *appressa*. Scale = 0.5 cm.S

Table 1. Disease scoring, based on Morin et al. (2012), of controlled inoculation of *Austropuccinia psidii* of Threatened (*Biodiversity Conservation Act* 2016 (WA) s 19) and Priority (Department of Biodiversity, Conservation and Attractions 2017) listed *Melaleuca* species included total number of plants scored and the percentage of plants observed in each disease scoring category.

Species name	Listing under Biodiversity Conservation Act 2016 or DBCA Priority Flora List	Total Number of Plants Scored	Disease Score (% of Total Plants)				
			1	2	3	4	5
Melaleuca acutifolia	NT	20	10	0	0	20	70
Melaleuca argentea	NT	52	44	23	4	4	25
Melaleuca cajuputi ssp. cajuputi	NT	54	28	15	13	35	9
Melaleuca dempta	P3	11	9	0	0	0	91
Melaleuca fulgens ssp. *fulgens*	NT	27	18	0	18	7	57
Melaleuca incana ssp. *gingilup*	P2	41	32	5	5	19	39
Melaleuca lanceolata	NT	27	15	0	18	26	41
Melaleuca lateralis	NT	31	6	0	0	0	94
Melaleuca penicula	P4	32	66	0	0	6	28
Melaleuca similis	P1	45	24	0	0	0	76
Melaleuca sophisma	P1	55	98	0	0	0	2
Melaleuca sp. Wanneroo	EN	80	14	0	0	0	86
Melaleuca viminea ssp. *appressa*	P2	25	28	0	0	4	68

EN = Endangered (Threatened species considered to be "facing a very high risk of extinction in the wild in the near future, as determined in accordance with criteria set out in the ministerial guidelines"), NT = Not Threatened, P1 - 3 = Priority 1 – 3 (Poorly-known species with conservation threat highest for Priority 1), P4 = Priority 4 (Rare, Near Threatened and other species in need of monitoring).

Here, we investigated the extent of rust susceptibility in 13 previously unscreened *Melaleuca* species from a range of geographic locations in Western Australia in response to *A. psidii*, revealing varying levels of susceptibility between species. Correlating disease responses in our greenhouse seedling tests with field response will be important in refining risk to these species. We were encouraged by

the presence of some resistant individuals within some vulnerable species. These results highlight

the importance of continued disease screening to determine the vulnerability of individual

Myrtaceae species to *A. psidii*. The identification of species with high susceptibility to the pathogen

may inform disease surveillance in the natural environment and direct conservation efforts such as

seed collection.

Acknowledgements

This work was funded by the Australian Research Council under Linkage Project LP190100093 and

AMM by the Australian Government Research Training Program. We thank the Department of

Biodiversity, Conservation and Attractions (Kings Park and Kensington) for suppling the seed and Bob

Makinson for the introductions that facilitated this work.

Conflict of interest

The authors declare no conflict of interest in the reporting of these results.

References

Agriculture Victoria (2022) About myrtle rust. https://agriculture.vic.gov.au/biosecurity/plant-
diseases/shrub-and-tree-diseases/myrtle-rust/about-myrtle-rust. Accessed 14 Jul 2022

Beenken L (2017) *Austropuccinia*: a new genus name for the myrtle rust *Puccinia psidii* placed within
the redefined family Sphaerophragmiaceae (Pucciniales). Phytotaxa 297:53-61-53–61.
https://doi.org/10.11646/PHYTOTAXA.297.1.5

Berthon KA, Fernandez Winzer L, Sandhu K, et al (2019) Endangered species face an extra threat:
susceptibility to the invasive pathogen *Austropuccinia psidii* (myrtle rust) in Australia.
Australasian Plant Pathology 48:385–393. https://doi.org/10.1007/s13313-019-00640-4

Brophy JJ, Craven LA, Doran JC (2013) *Melaleucas*: their botany, essential oils and uses

Carnegie AJ, Lidbetter JR (2012) Rapidly expanding host range for *Puccinia psidii* sensu lato in
Australia. Australasian Plant Pathology 41:13–29. https://doi.org/10.1007/s13313-011-0082-
6

Carnegie AJ, Lidbetter JR, Walker J, et al (2010) *Uredo rangelii*, a taxon in the guava rust complex,
newly recorded on Myrtaceae in Australia. Australasian Plant Pathology 39:463–466.
https://doi.org/10.1071/AP10102

Department of Biodiversity, Conservation and Attractions (2017) Threatened and Priority Flora
(DBCA-036)

Biodiversity Conservation Act 2016 (WA)

Department of Natural Resources and Environment Tasmania (2020) Plant species affected by myrtle
rust in Tasmania. https://nre.tas.gov.au/biosecurity-tasmania/plant-biosecurity/pests-and-
diseases/myrtle-rust/plant-species-affected-by-myrtle-rust-in-tasmania. Accessed 14 Jul
2022

Martino AM, Park RF, Tobias PA (2022) Three species of *Melaleuca* from Western Australia are highly
susceptible to *Austropuccinia psidii* in controlled inoculations. Australasian Plant Disease
Notes 17:1–4. https://doi.org/10.1007/S13314-022-00476-W/FIGURES/2

Morin L, Aveyard R, Lidbetter JR, Wilson PG (2012) Investigating the host-range of the rust fungus
Puccinia psidii sensu lato across tribes of the family Myrtaceae present in Australia. PLoS
ONE 7:1–7. https://doi.org/10.1371/journal.pone.0035434

Pegg GS, Giblin FR, McTaggart AR, et al (2014) *Puccinia psidii* in Queensland, Australia: disease
symptoms, distribution and impact. Plant Pathology 63:1005–1021.
https://doi.org/10.1007/s10530-015-0996-y

Pegg GS, Lee DJ, Carnegie AJ (2018) Predicting impact of *Austropuccinia psidii* on populations of
broad leaved *Melaleuca* species in Australia. Australasian Plant Pathology 47:421–430.
https://doi.org/10.1007/s13313-018-0574-8

Ryan M (2016) Australian forest profiles: *Melaleuca*

Saintilan N, Rogers K, Kelleway JJ, et al (2019) Climate Change Impacts on the Coastal Wetlands of
Australia. Wetlands 39:1145–1154. https://doi.org/10.1007/s13157-018-1016-7

Sandhu KS, Park RF (2013) Genetic basis of pathogenicity in *Uredo rangelii*

The Department of Primary Industries and Regional Development (2022a) Myrtle rust: Biosecurity
alert. https://www.agric.wa.gov.au/plant-biosecurity/myrtle-rust-threat-western-
australia?page=0%2C1. Accessed 14 Jul 2022

The Department of Primary Industries and Regional Development (2022b) Myrtle rust confirmed in
the Kimberley. https://www.agric.wa.gov.au/news/media-releases/myrtle-rust-confirmed-
kimberley. Accessed 20 Jul 2023

Westaway JO (2016) The pathogen Myrtle Rust (*Puccinia psidii*) in the Northern Territory: First
detection, new host and potential impacts. Northern Territory Naturalist 27:13–28.
https://doi.org/10.3316/INFORMIT.426544422268311

186