Resting state aperiodic slope predicts Research Domain Criteria Social Processes in young typically developing adults

Talitha C. Ford*[1,2], Aron T. Hill[1], Nina-Francesca Parrella[1], Melissa Kirkovski[1,3], Peter Donaldson[1], Peter G. Enticott[1]

1. Cognitive Neuroscience Unit, School of Psychology, Deakin University, Geelong, Victoria, Australia
2. Faculty of Health, Arts and Design, Swinburne University of Technology, Hawthorn, Victoria, Australia
3. College of Sport, Health and Engineering, Victoria University, Melbourne, Victoria, Australia

*Corresponding author
Dr. Talitha Ford
Cognitive Neuroscience Unit, School of Psychology, Deakin University
221 Burwood Hwy, Burwood
Victoria 3125, Australia

Abstract

Background: The aperiodic slope has been utilized to demonstrate differences in brain excitation-inhibition (E/I) in ageing, cognition, and neuro- and psycho-pathology. Furthermore, E/I imbalance has been associated with social communication difficulties in clinical and non-clinical groups. No work to date has explored the association between aperiodic activity and social communication difficulties.

Methods: A total of 40 typically developing young adults aged 18-24 (24 female; age mean=21.6, SD=1.9) each underwent 5 minutes each of eyes open and eyes closed resting state EEG using a 64-channel HydroCel Geodesic Sensor Net. Participants also completed Research Domain Criteria Social Processes measures: Social Responsiveness Scale (SRS), Reading the Mind in the Eyes task (RME), and Penn Emotional Recognition Task (PER).

Results: Sex differences in aperiodic slope and offset were observed, with steeper slope and greater offset observed in males, suggestive of increased inhibitory tone in this subgroup ($ps<.05$). Whole-brain aperiodic activity was not associated with SRS or RME scores, however better PER performance was differentially associated with left social brain slope and right social brain offset, with left slope decreasing (i.e., increasing E/I) and right offset increasing (i.e., decreasing E/I) with better performance ($ps<.05$).

Conclusions: These data show promise for supporting the notion that brain E/I is associated with social communication and processing difficulties, and this may extend to clinical groups (e.g., autism).

1. Introduction

Maintaining the equilibrium of excitatory and inhibitory synaptic membrane currents is essential for neural development, information processing, and memory storage (1). The brain's primary inhibitory neurotransmitter, gamma-aminobutyric acid (GABA), is responsible for regulating levels of excitatory glutamate, with disturbance to inhibitory tone leading to hyper-excitation. Likewise, disruption to excitatory glutamatergic processes can result in hyper-excitation depending on the location and nature of the disturbance (2). An imbalance in brain excitation/inhibition (E/I) ratios biased toward hyper-excitation has been associated with several psychopathologies and neurodevelopmental conditions, such as autism spectrum disorder (3–8), schizophrenia (9,10), and epilepsy (11,12).

Both animal and human studies point firmly toward E/I imbalance as a putative marker of social communication difficulties (13–16). Human neuroscientific studies support the role of brain E/I in social communication (17,18), with increased glutamate and/or reduced GABA levels associated with social communication difficulties in clinical (14,19–24) and non-clinical samples (22,25,26). Furthermore, psychopharmacological interventions that enhance brain inhibition, such as cannabidiol (CBD), riluzole, and R-Baclofen (STX209), have been shown to improve social functioning in children and adults with autism (27–33). In mice, optogenetic enhancement of excitation led to impairment in social behaviors, while systematically increasing inhibitory tone thereafter partially rescued these impairments (16). Similarly, low-frequency repetitive transcranial magnetic stimulation (TMS), which is known to inhibit neural activity, improved autism-like social and behavioral outcomes (34). Further, pharmacologically enhancing excitation reduces social behaviors in mice (13), while enhancing inhibition in autism mouse models improves social, cognitive, and behavioral outcomes (15,35).

Electroencephalography (EEG) has long been utilized as an accessible and affordable tool to probe the relationship between brain and behavior, particularly in relation to psychiatric and neurodevelopmental conditions (36). Putative markers of excitation and/or inhibition include mismatch negativity (37–39), functional E/I ratio (3), and oscillatory activity in delta, theta and gamma bands (40–44). Recent developments in resting state and task-related EEG data analysis have revealed that the broadband aperiodic signal, in contrast to periodic activity within a narrowband frequency range (i.e., neuronal oscillations), might be a more accurate reflection of E/I balance (45–47). Aperiodic activity is the background, scale-free activity that is represented on a 1/f-like power distribution, where spectral power decreases as frequency increases (45,48). The aperiodic signal, which is comprised of both the exponent (spectral slope) and offset (broadband shift in power across frequencies), is thought to reflect the integration of underlying synaptic currents (49). These currents have a double-exponential shape in the time domain, which produces the 1/f-like spectrum (45,46). Excitatory currents have a faster time constant and produce constant power at lower frequencies before rapid decay, while power decays more slowly as a function of frequency

for inhibitory currents. Thus, a smaller exponent (flatter 1/f-like slope) reflects more excitation relative to inhibition, while a larger exponent (steeper slope) reflects the converse (45,46). Given its utility for examining E/I balance across the cortex, and relative affordability and simple acquisition, the aperiodic slope has been utilized to demonstrate E/I differences in neuro- and psycho-pathology (50–53). The aperiodic offset (i.e., the y-intercept of the spectrum) represents the uniform shift of broadband power across frequencies (45,54), and is thought to reflect neuronal population spiking (55,56) and fMRI BOLD signal (57). Aperiodic offset has been associated with age (45,48,58,59), sex (59), and working memory (45,47), but not selective attention and executive control in children (59). No work to date has applied aperiodic activity analyses to explore differences in E/I social communication in a non-clinical sample.

Due to the heterogeneity and spectrum nature of clinical conditions such as autism, it is important to investigate the extent to which brain processes relate to specific symptom domains relevant to specific conditions, rather than within the constraints of diagnostic boundaries. The broader autism phenotype reflects the notion that milder autistic traits are present in varying degrees across the non-clinical population globally (60,61), hence, extensive evidence supports the utility of non-clinical populations in investigating the behavioral and neurobiological characteristics of autism (25,62,63). The Research Domain Criteria (RDoC) highlight the limitations of diagnosis-specific research, and provides a rich framework for understanding psychiatric conditions in the context of varying degrees of symptomatology with neurobiological and/or genetic origins (64). Within the RDoC, the *social processes* domain includes social communication, affiliation and attachment with others, and perception and understanding of the self and others (65). Self-report measures of *social processes* recommended by RDoC include the Social Responsiveness Scale (SRS; 66), while objective measures include the Penn Emotion Recognition (PER; 67), and Reading the Mind in the Eyes (RME; 68) tasks. These measures have been utilized to assess social functioning in both clinical and non-clinical populations. Understanding the neurobiological mechanisms of *social processes*, at both clinical and non-clinical levels, is essential for broader global efforts to identify effective therapeutics to support those with social processing difficulties.

To that end, this study investigates the relationship between RDoC *social processes* and aperiodic activity in a young, non-clinical adults. It was hypothesized that *social processes* difficulties would be associated with a flatter aperiodic slope, indicating reduced global inhibitory tone (E<I). Given there has been no literature published to date on aperiodic activity and social difficulties, we also explored their relationship specifically in right and left temporo-parietal regions, which are known to be involved in social information processing (69–71).

2.   Methods and Materials
    2.1. Participants

A total of 40 young adults (24 female) aged 18-24 participated in this study (mean age=21.6 years, SD=1.9). Participants had an IQ>90 (mean=114.6, SD=10.15) and were excluded if they self-reported: a history of developmental, psychiatric genetic, or neurological conditions; a first-degree relative with a history of autism spectrum disorder, schizophrenia, bipolar disorder, or other psychiatric condition (not including depression and anxiety); a substance use disorder; regular smoking (> 1 cigarette/day); or taking any psychoactive medications. Participants were asked to abstain from caffeine and nicotine for 12 hours, and recreational drugs for 1 week, prior to testing; confirmation of compliance was made verbally on the day of testing. All participants provided written informed consent prior to any data collection for this study, and ethics approval was granted by Deakin University's Human Research Ethics Committee (2019-208) in accordance with the Declaration of Helsinki.

    2.2. Psychometric and Behavioral Measures

Social Responsiveness Scale Version-2 (SRS; 72): The SRS is a 65-item measure of symptom severity across five autism symptom domains: Social Awareness (8 items), Social Motivation (11 items), Social Cognition (12 items), Social Communication (22 items), and Repetitive Behaviors/Restricted Interests (12 items). Responses are measured on a 4-point scale from 1 (*not true*) to 4 (*almost always true*). The psychometric properties of the adult self-report measure of the SRS were assessed by Ingersoll et al. (2011), with good reliabilities across subscales reported in an adult sample (SRS total α=.95, Social Awareness α=.64, Social Cognition α=.78, Social Communication α=.88, Social Motivation α=.81, and Repetitive Behaviors/Restricted Interests α=.81). Within the RDoC framework, the SRS is registered as a measure of Social Communication - reception and production of non-facial communication.

Reading the Mind in the Eyes Task (RME; 68): The RME is a 36-item measure of perception and understanding of conspecific mental states through the assessment an actor's eyes. The computerized task presents consecutive photos of actor's eyes in the center of the screen, and four words (one in each corner of the screen) that might describe the emotional state of the eye. Participants are to select the word that best describes the actor's emotion via mouse click. Accuracy on the RME is scored as the proportion of correct responses (from 0 to 1). Because participants are provided a dictionary of the emotional labels presented, reaction time data are not analyzed. Within the RDoC framework, the RME is a measure of Perception and Understanding of Others – understanding mental states.

Penn Emotion Recognition Task (PER; Gur et al., 2002): The PER task includes 40 color photographs of static faces expressing 4 basic emotions (happiness, sadness, anger, or fear) and neutral expressions. Stimuli are balanced for poser's gender, age, and ethnicity, and for

each emotion category. Four high-intensity and four low-intensity expressions are included. Participants view one image at a time and choose the correct emotion label for each face via mouse click. A composite measure of accuracy and reaction time, the linear integrated speed-accuracy score (LISAS), is calculated according to the following formula: $LISAS = RT + (S_{RT} / S_{PE}) \times PE$, where RT is participant mean reaction time, PE is participant mean proportion of errors, S_{RT} is the standard deviation of the reaction time, and S_{PE} is the standard deviation of the proportion or errors (73). PER LISAS will be referred to as "PER performance" hereafter, with higher scores indicating better performance. Within the RDoC framework, the PER task is a measure of Social Communication – reception of facial communication.

2.3. Electroencephalography data acquisition

EEG data were recorded using a 64-channel HydroCel Geodesic Sensor Net (Electrical Geodesics, Inc, USA) containing silver/silver-chloride (Ag/AgCl) electrodes surrounded by electrolyte-wetted sponges. Electrical Geodesics Net Station EEG software (version 5.0) was used to acquire the data via a Net Amps 400 amplifier using a sampling rate of 1 KHz. Data were referenced online to electrode Cz, and electrode impedances were checked to ensure they were < 50 KOhms prior to data collection. Participants were seated in a quiet, dimly lit room and asked to sit still with their eyes open while looking at a fixation cross for 5 minutes, and then eyes closed for 5 minutes, while resting state data were collected. Electrode impedances were checked between eyes open and eyes closed blocks to ensure they remained < 50 KOhms. One participant withdrew from the study and did not undergo EEG.

2.4. EEG Pre-processing

EEG data were pre-processed in MATLAB (R2021a; The Mathworks, Massachusetts, USA) incorporating the EEGLAB toolbox (v2023.0; 74) and custom scripts. The Reduction of Electroencephalographic Artifacts (RELAX; v1.1.2) software (75,76) was used to clean each EEG dataset. The fully automated RELAX pipeline uses empirical approaches to identify and reduce non-neural artifacts, and includes the use of both multi-channel Wiener filters and wavelet enhanced independent component analysis (ICA). As part of the RELAX pipeline, the data were bandpass filtered between 0.5 and 80 Hz (fourth-order zero-phase Butterworth filter). A notch filter (47-53 Hz) was then applied to remove any line noise. Any bad channels were removed using a multi-step process incorporating the 'findNoisyChannels' function (PREP pipeline; 77). Multi-channel Wiener filtering (78) was used to initially clean blinks, muscle activity, and horizontal eye movement, followed by robust average re-referencing (77), and wavelet-enhanced ICA (78), with components for cleaning identified using the automated independent component (IC) classifier IClabel (79). Any rejected electrodes were then interpolated back into the data using spherical interpolation and all pre-processed data files were visually inspected prior to further analysis.

2.5. Electroencephalography data analysis

All pre-processed resting state data were visually inspected for quality assurance prior to inclusion in statistical analyses. Two participants were excluded from the study due to excessive line noise for both eyes open and eyes closed conditions, and an additional two participants' eyes open data were excluded due to excessive line noise. The power spectral density of the EEG signal was calculated using Welch's method (2-second Hamming window, 50% overlap) for all electrodes across all participants for both the eyes-open and eyes-closed datasets. The FOOOF toolbox (version 1.0.0; https://fooof-tools.github.io/fooof/; 45) was then used to parameterize the spectral data through separation of the periodic and aperiodic components of the signal. The aperiodic exponent and offset were extracted across 1 to 40 Hz using the 'fixed' aperiodic mode. Spectral parameterization settings for the algorithm were: peak width limits=[1, 12], maximum number of peaks=8, peak threshold=2, minimum peak height=0.05.

The performance of the FOOOF algorithm for fitting the data was assessed using two "goodness of fit" measures: R^2, explained variance; and mean absolute error (MAE), total error of the model fit (45,51). Model fits for both eyes open and eyes closed data were good (eyes open: R^2=.99, MAE=.047; eyes closed: R^2=.99, MAE=.054. Complete fit statistics are presented in the Supplementary Material). There was no difference in mean R^2 between eyes open and eyes closed conditions ($t(64.13)$=-.30, p=.765), while MAE for eyes open was significantly lower compared to eyes closed ($t(69.44)$=2.19, p=.032). Aperiodic exponent (slope) was significantly larger for eyes closed condition ($t(69.65)$=2.82, p=.006), indicating a steeper slope than for eyes open. Similarly, the offset (intercept) was larger for eyes closed than eyes open condition ($t(69.97)$=3.0, p=.004). There were strong, positive correlations between exponent and offset for eyes open (rho=.73, p<.001) and eyes closed (rho=.80, p<.001), and between eyes open and closed exponent (rho=.73, p>.001) and eyes open and closed offset (rho=.78, p<.001).

2.6. Statistical analyses

Statistical analyses were conducted in R (version 4.1.0). Multivariate linear models (MLMs) were designed to assess the extent to which aperiodic slope and offset with eyes open and eyes closed predicted *social processes* outcomes (SRS total, PER IES, and RME accuracy). Separate models were defined for exponent and offset (i.e., model 1 predictors included eyes open exponent and eyes closed exponent, model 2 included eyes open offset and eyes closed offset; see Supplementary material for full model deigns). Assumptions for all MLMs were assessed using the *olsrr* and *MVN* packages in R (80,81), including normality, linearity, homoscedasticity, and multicollinearity. All assumptions were met, except for a significant skew in RME and PER LISAS scores, which was subsequently normalized using the Yeo-Johnson transform in R (VGAM package; 82).

Exploratory MLMs were also designed to investigate the extent to which left and right social brain aperiodic activity predicted *social processes* outcomes. To calculate aperiodic slope and offset in social brain regions, electrodes over left and right temporo-parietal regions were averaged separately (left: P5 CP5 TP9 TP7 C5 T7 T9; right: P6 CP6 TP10 TP8 C6 T8 T10; see Figure 1). These electrodes cover brain regions including the superior, middle, and inferior temporal gyrus, supramarginal gyrus, and angular gyrus, all known to be involved in social information processing (69–71). False discovery rate was corrected for using the Benjamin Hochberg method in R (stats package; 83).

Figure 1: Electroencephalograph electrode placement. 64-channel Geodesic Sensor Net electrode placement (dark circles), with the cluster of electrodes selected to represent left (blue) and right (orange) social brain highlighted.

3. Results
    3.1. Descriptive statistics

Descriptive statistics for final participant sample characteristics (n=37), *social processes* outcomes, and aperiodic activity metrics are presented in Table 1. There were no significant age or biological sex effects on *social processes* measures (ps>.05), and no significant correlations between age and aperiodic activity outcomes (ps>.05). However, males compared to females had a significantly steeper exponent (t(24.20)=-2.56, p=.017, p_{FDR}=.033) and higher offset (t(29.20)=-2.08, p=.047, p_{FDR}=.047) during eyes open, as well as a steeper exponent (t(33.76)=-2.83, p=.008, p_{FDR}=.032) and higher offset during eyes closed (t(34.90)=-2.34, p=.025, p_{FDR}=.033; Figure 2).

Table 1: Descriptive statistics of psychometric, behavioral, and aperiodic activity measures

	Total (N=37)		Female (n=22)	Male (n=15)
	Mean (SD)	Min-Max	Mean (SD)	Mean (SD)
Age	22 (1.9)	18 – 25	21 (1.9)	22 (1.9)
IQ	115 (9.8)	91 – 148	114 (8.2)	117 (12)
Social Processes				
SRS total	52 (6.8)	42 – 69	51 (5.8)	54 (8.4)
SRS SCI	53 (6.9)	42 – 71	52 (5.6)	54 (8.6)
SRS RRB	53 (8.4)	40 – 73	51 (7.2)	55 (10)
RME accuracy	.81 (.08)	.64 – .94	.83 (.08)	.78 (.08)
PER RT	2.9 (.69)	2.0 – 4.9	2.8 (.48)	3.0 (.94)
PER accuracy	.89 (.05)	.78 – .97	.90 (.04)	.86 (.05)
PER performance	3.3 (.97)	2.2 – 7.0	3.1 (.65)	3.5 (1.3)
Aperiodic activity				
Whole brain				
EO exponent	1.2 (.17)	.96 – 1.7	1.2 (.14)	1.3 (.19)
EC exponent	1.3 (.17)	.95 – 1.7	1.3 (.17)	1.4 (.14)
EO offset	.21 (.38)	-.47 – 1.4	.10 (.36)	.36 (.38)
EC offset	.49 (.41)	-.63 – 1.3	.37 (.44)	.66 (.31)
Left social brain				
EO exponent	1.2 (.19)	.96 – 1.7	1.2 (.15)	1.3 (.21)
EC exponent	1.3 (.20)	.86 – 1.7	1.3 (.20)	1.4 (.15)
EO offset	.16 (.41)	-.50 – 1.5	.014 (.34)	.35 (.43)
EC offset	.42 (.45)	-.67 – 1.3	.25 (.45)	.66 (.35)
Right social brain				
EO exponent	1.2 (.20)	.67 – 1.7	1.2 (.18)	1.3 (.19)
EC exponent	1.3 (.20)	.77 – 1.6	1.2 (.21)	1.4 (.13)
EO offset	.18 (.42)	-.84 – 1.3	.026 (.37)	.38 (.41)
EC offset	.43 (.48)	-1.0 – 1.2	.26 (.50)	.67 (.34)

Notes: SD=standard deviation, SRS=Social Responsiveness Scale, SCI=Social Communication Index, RRB=Restricted Repetitive Behaviors, RME=Reading the Mind in the Eyes, PER=Penn Emotion Recognition Task, RT=reaction time, performance=linear integrated speed-accuracy score (LISAS), EO=eyes open, EC=eyes closed.

Figure 2: Sex differences in aperiodic activity. Sex differences in exponent (a) and offset (b) for eyes open and eyes closed conditions. $*p_{FDR}<.05$

3.2. Social processes outcomes

Spearman's rho correlations were conducted to test the relationship between the three *social processes* measures. There were no significant relationships between SRS total scores and RME accuracy or PER performance ($ps>.05$). The relationship between PER performance and RME accuracy was significant, with PER performance increasing with RME accuracy as would be expected ($rho=.391$, $p=.018$, $p_{FDR}=.026$).

3.3. Whole-brain aperiodic activity MLMs

Multivariate linear models were designed to test the extent to which aperiodic exponent and offset predicted *social processes* outcomes of SRS scores, RME accuracy, and PER performance. In the first model, the exponent values for both eyes open and eyes closed conditions were entered as predictors, however, these were not significant predictors of SRS total score ($R^2_{adj}=.058$, $F(2,31)=2.01$, $p=.150$), PER performance ($R^2_{adj}=-.057$, $F(2,31)=.12$, $p=.885$), or RME accuracy ($R^2_{adj}=.032$, $F(2,31)=1.55$, $p=.228$; Figure 3). In the second model, the offset values for both eyes open and eyes closed conditions were entered into the model as predictors of *social processes* outcomes. Again, offset was not a significant predictor of SRS total score ($R^2_{adj}=-.013$, $F(2,31)=.80$, $p=.460$), PER performance ($R^2_{adj}=-.044$, $F(2,31)=.31$, $p=.738$), or RME accuracy ($R^2_{adj}=.065$, $F(2,31)=2.14$, $p=.134$; Figure 3).

Figure 3: Scatter plots of aperiodic activity and Social Processes outcomes. Scatter plots of exponent (top[a-c]; model 1) and offset (bottom[d-e]; model 2) versus SRS total score (left[a,d]), PER performance (middle[b,e]) and RME accuracy (right[c,f]), in eyes open (green) and eyes closed (red) conditions.

3.4. Social brain aperiodic activity MLMs

Exploratory analyses were conducted to investigate whether aperiodic activity in social brain regions specifically might predict *social processes* outcomes (SRS, PER performance, RME accuracy). Therefore, measures of exponent and offset for both eyes open (EO) and eyes closed (EC) conditions were averaged over temporo-parietal clusters in right and left hemisphere (Figure 4), and two models were defined for exponent and offset separately with four predictors each (i.e., left STS EO, left STS EC, right STS EO, right STS EC) (see Supplementary material for models). None of the overall models for SRS, PER performance, and RME accuracy for both exponent and offset predictors were significant (*p*s>.1). When looking at the individual predictors in the models, left STS exponent (*ß*=-.042, *t*=-2.18, *p*=.037), and right STS offset (*ß*=.025, *t*=2.06, *p*=.049) predicted PER performance in the eyes closed condition (Figure 4). That is, when participants had their eyes closed, right offset increased (i.e., decreased E/I ratio) with increasing PER performance, while left STS exponent decreased with better performance (i.e., increased E/I ratio).

Figure 4: Scatter plots of aperiodic activity by hemisphere and Social Processes outcomes.
Scatter plots of exponent (top[a-c]; model 1) and offset (bottom[d-f]; model 2) versus SRS total score (left[a,d]), PER performance (middle[b,e]) and RME accuracy (right[c,f]), in eyes open (green) and eyes closed (red) conditions from the left (solid line) and right (dotted line) social brain regions.

4. Discussion

Investigations of aperiodic activity (exponent and offset) are increasingly reported across neurodevelopmental, cognitive, and psychiatric datasets as a putative measure of brain E/I ratios (50–53,84), with a steeper exponent (slope) indicating more brain inhibition (i.e., reduced E/I). This study was the first, to our knowledge, to investigate the extent to which resting state aperiodic activity predicts RDoC *social processes* outcomes in non-clinical young adults.

Given a shift in E/I towards increased excitation has been associated with social communication difficulties in clinical, non-clinical, and animal studies (13–15,25), it was predicted that reduced (i.e., flatter) aperiodic slope would be associated with more social communication and processing difficulties. Contrary to expectations, the aperiodic slope averaged across all electrodes (i.e., whole brain) did not predict *social processes* outcomes in either the eyes open or eyes closed condition. This may indicate that: 1) predictive power of exponent and offset is possibly region-specific, thus including all electrodes introduces substantial noise that effectively drowns out any signal; 2) there was insufficient power to

detect an effect; and/or 3) E/I measured by aperiodic brain activity does not predict the RDoC *social processes* measures utilized here.

To explore the regional specificity of aperiodic activity to social processes, two 'social brain' clusters were created left and right hemisphere that localized to superior, middle, and inferior temporal gyrus, supramarginal gyrus, and angular gyrus (71). These regions are known to be involved in social information processing (69,70). Results of the regression models suggested that aperiodic activity within these left and right social brain regions overall did not predict social processes outcomes in our non-clinical sample. When inspecting the individual regression coefficients within each model, however, there was some indication that better facial emotion recognition (PER performance) was associated with a flatter exponent (i.e., higher E/I ratio) in left social brain. In contrast, in right social brain, higher offset (i.e., shift in broadband power across frequencies) was associated with better facial emotion recognition. It is important to note that these effect sizes were very small and should therefore be interpreted cautiously.

Further investigations with larger samples are essential to support this finding, particularly given the relationship is in the opposite direction to what would be expected. Previous research suggests that social communication difficulties are associated with an increase in E/I (8,16), implying increased excitation and/or reduced inhibition. However, more recent research has suggested that the nature of the E/I imbalance, that is increased excitation or increased inhibition, is region specific (7,17,18), and regional specificity of increased inhibitory tone has been reported in mouse models (85). The increased inhibitory tone in social brain regions in conjunction with more social communication difficulties observed in this study may reflect underlying neurobiological mechanisms.

In line with previous literature, males had a significantly higher aperiodic offset and steeper slope than females in both eyes open and eyes closed conditions (48,59). Sex differences in a prior adolescent sample were thought to reflect an advanced neurophysiological maturity in females (86), which may be the case in this young adult sample as neurodevelopment continues into early adulthood for both males and females (87). Contrary to previous reports (45,48,58,59), exponent and offset were not associated with age, which may reflect that aperiodic activity is relatively stable between 18 and 25 years, or that the age range of the sample was too narrow to detect age effects. Also, in line with previous research in children and adolescents, steeper exponent and higher offset were recorded in the eyes closed compared to eyes open condition (48,59). This reflects a reduction in E/I when eyes are closed, which could be explained by the lack of visual stimulation, attention, and arousal, compared to when participants' eyes were open (88).

There were no significant relationships between RDoC *social processes* self-report (SRS) and behavioral (RME, PER) outcomes in this study. This was somewhat unexpected and possibly

problematic given the three measures, SRS, RME and PER, are proposed by the RDoC as gold standard measures of the *social processes* constructs. Although they appear to index different aspects of social processes, namely social communication (reception of facial communication [PER], and reception and production of non-facial [SRS]), and perception and understanding of others (understanding mental states [RME]), it is reasonable to expect that they would be related. The lack of interrelationship in this study may reflect the limited spread in performance/difficulties in the tasks in this non-clinical sample, insufficient data, or that the measures have poor construct validity, potentially due to poor insight when completing self-report measures.

The main limitations for this study were the modest sample size (N=37) and narrow spread of social processes difficulties, particularly for SRS scores. These issues combined may have been a significant contributor to the lack of statistically significant findings, given the small effect sizes reported herein. A more diverse spread of abilities within the sample, for example including clinical participants, may have revealed more compelling relationships between aperiodic activity and social processing abilities. Furthermore, EEG is limited in its ability to examine social brain areas specifically due to its relatively poor spatial resolution. Source localization using magnetoencephalography (MEG) and concurrent participant structural brain scans would allow for a more accurate analysis of social brain and social processes abilities.

Despite these limitations, this preliminary data shows promise for supporting the notion of an imbalance in brain E/I associated with social communication processing difficulties across clinical and non-clinical groups. Aperiodic exponent and offset were moderately associated with emotion recognition abilities, particularly in the left and right social brain regions. Future work should seek to examine this relationship in a larger sample across clinical and non-clinical populations. Furthermore, studies involving pharmacological interventions that modulate excitatory and/or inhibitory processes would provide valuable insights into the utility of the aperiodic slope as a putative neurobiomarker of social processes abilities.

Declarations of interest: none
Acknowledgements: Talitha Ford and this study were supported by a Dean's Postdoctoral Research Fellowship awarded by the Faculty of Health, Deakin University.

References

1. Zhou S, Yu Y. Synaptic E-I Balance Underlies Efficient Neural Coding. Frontiers in Neuroscience [Internet]. 2018 [cited 2023 Jul 11];12. Available from: https://www.frontiersin.org/articles/10.3389/fnins.2018.00046
2. Sohal VS, Rubenstein JLR. Excitation-inhibition balance as a framework for investigating mechanisms in neuropsychiatric disorders. Mol Psychiatry [Internet]. 2019 May 14 [cited 2019 Jul 1]; Available from: http://www.nature.com/articles/s41380-019-0426-0

3. Bruining H, Hardstone R, Juarez-Martinez EL, Sprengers J, Avramiea AE, Simpraga S, et al. Measurement of excitation-inhibition ratio in autism spectrum disorder using critical brain dynamics. Sci Rep. 2020 Dec;10(1):9195.

4. Dickinson A, Jones M, Milne E. Measuring neural excitation and inhibition in autism: Different approaches, different findings and different interpretations. Brain Research. 2016 Oct;1648:277–89.

5. Fatemi SH. The hyperglutamatergic hypothesis of autism. Progress in Neuro-Psychopharmacology & Biological Psychiatry. 2008 Apr 1;32(3):911, author reply 912-3.

6. Gogolla N, LeBlanc JJ, Quast KB, Südhof TC, Fagiolini M, Hensch TK. Common circuit defect of excitatory-inhibitory balance in mouse models of autism. J Neurodevelop Disord. 2009 Jun;1(2):172–81.

7. Nelson SB, Valakh V. Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism Spectrum Disorders. Neuron. 2015 Aug 19;87(4):684–98.

8. Rubenstein JLR, Merzenich MM. Model of autism: increased ratio of excitation/inhibition in key neural systems: Model of autism. Genes, Brain and Behavior. 2003 Oct 16;2(5):255–67.

9. Egerton A, Grace AA, Stone J, Bossong MG, Sand M, McGuire P. Glutamate in schizophrenia: Neurodevelopmental perspectives and drug development. Schizophrenia Research. 2020 Sep;223:59–70.

10. Marsman A, van den Heuvel MP, Klomp DWJ, Kahn RS, Luijten PR, Hulshoff Pol HE. Glutamate in Schizophrenia: A Focused Review and Meta-Analysis of 1H-MRS Studies. Schizophr Bull. 2013 Jan;39(1):120–9.

11. Bozzi Y, Provenzano G, Casarosa S. Neurobiological bases of autism-epilepsy comorbidity: a focus on excitation/inhibition imbalance. Eur J Neurosci. 2018 Mar;47(6):534–48.

12. Hanada T. Ionotropic Glutamate Receptors in Epilepsy: A Review Focusing on AMPA and NMDA Receptors. Biomolecules. 2020 Mar 18;10(3):464.

13. Bredewold R, Schiavo JK, van der Hart M, Verreij M, Veenema AH. Dynamic changes in extracellular release of GABA and glutamate in the lateral septum during social play behavior in juvenile rats: Implications for sex-specific regulation of social play behavior. Neuroscience. 2015 Oct;307:117–27.

14. Cochran DM, Sikoglu EM, Hodge SM, Edden RAE, Foley A, Kennedy DN, et al. Relationship among Glutamine, γ-Aminobutyric Acid, and Social Cognition in Autism Spectrum Disorders. J Child Adol Psychop. 2015 May;25(4):314–22.

15. Han S, Tai C, Jones CJ, Scheuer T, Catterall WA. Enhancement of inhibitory neurotransmission by GABAA receptors having alpha2,3-subunits ameliorates behavioral deficits in a mouse model of autism. Neuron. 2014 Mar 19;81(6):1282–9.

16. Yizhar O, Fenno LE, Prigge M, Schneider F, Davidson TJ, O'Shea DJ, et al. Neocortical excitation/inhibition balance in information processing and social dysfunction. Nature. 2011 Sep 8;477(7363):171–8.

17. Ajram LA, Pereira AC, Durieux AMS, Velthius HE, Petrinovic MM, McAlonan GM. The contribution of [1H] magnetic resonance spectroscopy to the study of excitation-inhibition in autism. Progress in Neuro-Psychopharmacology & Biological Psychiatry. 2019 Sep 21;

18. Ford TC, Crewther DP. A Comprehensive Review of the (1)H-MRS Metabolite Spectrum in Autism Spectrum Disorder. Frontiers in Molecular Neuroscience. 2016;9(14):1–27.

19. Brown MS, Singel D, Hepburn S, Rojas DC. Increased glutamate concentration in the auditory cortex of persons with autism and first-degree relatives: a (1)H-MRS study. Autism research : official journal of the International Society for Autism Research. 2013 Feb;6(1):1–10.

20. Gaetz W, Bloy L, Wang DJ, Port RG, Blaskey L, Levy SE, et al. GABA estimation in the brains of children on the autism spectrum: Measurement precision and regional cortical variation. Neuroimage. 2014 May 24;86(1):1–9.

21. Hegarty JP, Weber DJ, Cirstea CM, Beversdorf DQ. Cerebro-Cerebellar Functional Connectivity is Associated with Cerebellar Excitation–Inhibition Balance in Autism Spectrum Disorder. Journal of autism and developmental disorders. 2018;1–14.

22. Tebartz van Elst L, Maier S, Fangmeier T, Endres D, Mueller GT, Nickel K, et al. Disturbed cingulate glutamate metabolism in adults with high-functioning autism spectrum disorder: evidence in support of the excitatory/inhibitory imbalance hypothesis. Molecular psychiatry. 2014 Jul 22;

23. Pretzsch CM, Voinescu B, Lythgoe D, Horder J, Mendez MA, Wichers R, et al. Effects of cannabidivarin (CBDV) on brain excitation and inhibition systems in adults with and without Autism Spectrum Disorder (ASD): a single dose trial during magnetic resonance spectroscopy. Transl Psychiatry. 2019 Dec;9(1):313.

24. Pretzsch CM, Floris DL, Voinescu B, Elsahib M, Mendez MA, Wichers R, et al. Modulation of striatal functional connectivity differences in adults with and without autism spectrum disorder in a single-dose randomized trial of cannabidivarin. Molecular Autism. 2021 Dec;12(1):49.

25. Ford TC, Nibbs R, Crewther DP. Increased glutamate/GABA+ ratio in a shared autistic and schizotypal trait phenotype termed Social Disorganisation. NeuroImage: Clin. 2017;16:125–31.

26. Ford TC, Nibbs R, Crewther DP. Glutamate/GABA+ ratio is associated with the psychosocial domain of autistic and schizotypal traits. PLoS ONE. 2017;12(7):e0181961.

27. Aran A, Harel M, Cassuto H, Polyansky L, Schnapp A, Wattad N, et al. Cannabinoid treatment for autism: a proof-of-concept randomized trial. Molecular Autism. 2021 Dec;12(1):6.

28. da Silva Junior EA, Medeiros WMB, Torro N, Sousa JMM de, Almeida IBCM de, Costa FB da, et al. Cannabis and cannabinoid use in autism spectrum disorder: a systematic review. Trends Psychiatry Psychother. 2021 May 21;44:e20200149.

29. Hacohen M, Stolar OE, Berkovitch M, Elkana O, Kohn E, Hazan A, et al. Children and adolescents with ASD treated with CBD-rich cannabis exhibit significant improvements particularly in social symptoms: an open label study. Transl Psychiatry. 2022 Sep 9;12(1):375.

30. Frye RE. Clinical potential, safety, and tolerability of arbaclofen in the treatment of autism spectrum disorder. Drug Healthc Patient Saf. 2014 May 10;6:69–76.

31. Ajram LA, Horder J, Mendez MA, Galanopoulos A, Brennan LP, Wichers RH, et al. Shifting brain inhibitory balance and connectivity of the prefrontal cortex of adults with autism spectrum disorder. Translational Psychiatry. 2017 May 23;7(5):e1137.

32. Cifelli P, Ruffolo G, De Felice E, Alfano V, van Vliet EA, Aronica E, et al. Phytocannabinoids in Neurological Diseases: Could They Restore a Physiological GABAergic Transmission? International Journal of Molecular Sciences. 2020 Jan;21(3):723.

33. Parrella NF, Hill AT, Enticott PG, Barhoun P, Bower IS, Ford TC. A systematic review of cannabidiol trials in neurodevelopmental disorders. Pharmacology Biochemistry and Behavior. 2023 Sep;230:173607.

34. Tan T, Wang W, Xu H, Huang Z, Wang YT, Dong Z. Low-Frequency rTMS Ameliorates Autistic-Like Behaviors in Rats Induced by Neonatal Isolation Through Regulating the Synaptic GABA Transmission. Front Cell Neurosci [Internet]. 2018 Feb 28 [cited 2019 Jun 3];12. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5835518/

35. Stoppel LJ, Kazdoba TM, Schaffler MD, Preza AR, Heynen A, Crawley JN, et al. R-Baclofen Reverses Cognitive Deficits and Improves Social Interactions in Two Lines of 16p11.2 Deletion Mice. Neuropsychopharmacol. 2018 Feb;43(3):513–24.

36. Milovanovic M, Grujicic R. Electroencephalography in Assessment of Autism Spectrum Disorders: A Review. Frontiers in Psychiatry [Internet]. 2021 [cited 2023 Aug 1];12. Available from: https://www.frontiersin.org/articles/10.3389/fpsyt.2021.686021

37. Greenwood LM, Leung S, Michie PT, Green A, Nathan PJ, Fitzgerald P, et al. The effects of glycine on auditory mismatch negativity in schizophrenia. Schizophr Res. 2018 Jan;191:61–9.

38. Rowland LM, Summerfelt A, Wijtenburg SA, Du X, Chiappelli JJ, Krishna N, et al. Frontal Glutamate and gamma-Aminobutyric Acid Levels and Their Associations With Mismatch Negativity and Digit Sequencing Task Performance in Schizophrenia. JAMA Psychiatry. 2016 Feb;73(2):166–74.

39. Sehatpour P, Javitt DC, De Baun HM, Carlson M, Beloborodova A, Margolin DH, et al. Mismatch negativity as an index of target engagement for excitation/inhibition-based treatment development: a double-blind, placebo-controlled, randomized, single-dose cross-over study of the serotonin type-3 receptor antagonist CVN058. Neuropsychopharmacol. 2022 Feb;47(3):711–8.

40. Grent-'t-Jong T, Gross J, Goense J, Wibral M, Gajwani R, Gumley AI, et al. Resting-state gamma-band power alterations in schizophrenia reveal E/I-balance abnormalities across illness-stages. eLife. 2018 Sep 27;7:e37799.

41. Lisman J. Excitation, inhibition, local oscillations, or large-scale loops: what causes the symptoms of schizophrenia? Curr Opin Neurobiol. 2012 Jun;22(3):537–44.

42. Uhlhaas PJ, Singer W. Abnormal neural oscillations and synchrony in schizophrenia. Nature reviews Neuroscience. 2010 Feb;11(2):100–13.

43. Coben R, Clarke AR, Hudspeth W, Barry RJ. EEG power and coherence in autistic spectrum disorder. Clin Neurophysiol. 2008 May;119(5):1002–9.

44. Barttfeld P, Wicker B, Cukier S, Navarta S, Lew S, Sigman M. A big-world network in ASD: dynamical connectivity analysis reflects a deficit in long-range connections and an excess of short-range connections. Neuropsychologia. 2011 Jan;49(2):254–63.

45. Donoghue T, Haller M, Peterson EJ, Varma P, Sebastian P, Gao R, et al. Parameterizing neural power spectra into periodic and aperiodic components. Nat Neurosci. 2020 Dec;23(12):1655–65.

46. Gao R, Peterson EJ, Voytek B. Inferring synaptic excitation/inhibition balance from field potentials. NeuroImage. 2017 Sep;158:70–8.

47. Waschke L, Donoghue T, Fiedler L, Smith S, Garrett DD, Voytek B, et al. Modality-specific tracking of attention and sensory statistics in the human electrophysiological spectral exponent. Chait M, Shinn-Cunningham BG, Postle BR, Simon JZ, editors. eLife. 2021 Oct 21;10:e70068.

48. Hill AT, Clark GM, Bigelow FJ, Lum JAG, Enticott PG. Periodic and aperiodic neural activity displays age-dependent changes across early-to-middle childhood. Developmental Cognitive Neuroscience. 2022 Apr;54:101076.

49. Donoghue T, Schaworonkow N, Voytek B. Methodological considerations for studying neural oscillations. European Journal of Neuroscience. 2021;n/a(n/a):1–26.

50. Molina JL, Voytek B, Thomas ML, Joshi YB, Bhakta SG, Talledo JA, et al. Memantine Effects on Electroencephalographic Measures of Putative Excitatory/Inhibitory Balance in Schizophrenia. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging. 2020 Jun;5(6):562–8.

51. Ostlund B, Alperin BR, Drew T, Karalunas SL. Behavioral and cognitive correlates of the aperiodic (1/f-like) exponent of the EEG power spectrum in adolescents with and without ADHD. Developmental Cognitive Neuroscience. 2021 Apr 1;48:100931.

52. Robertson MM, Furlong S, Voytek B, Donoghue T, Boettiger CA, Sheridan MA. EEG power spectral slope differs by ADHD status and stimulant medication exposure in early childhood. Journal of Neurophysiology. 2019 Dec 1;122(6):2427–37.

53. Wilkinson CL, Nelson CA. Increased aperiodic gamma power in young boys with Fragile X Syndrome is associated with better language ability. Molecular Autism. 2021 Feb 25;12(1):17.

54. Ostlund B, Donoghue T, Anaya B, Gunther KE, Karalunas SL, Voytek B, et al. Spectral parameterization for studying neurodevelopment: How and why. Developmental Cognitive Neuroscience. 2022 Apr;54:101073.

55. Manning JR, Jacobs J, Fried I, Kahana MJ. Broadband shifts in local field potential power spectra are correlated with single-neuron spiking in humans. Journal of Neuroscience. 2009;29(43):13613–20.

56. Miller KJ, Honey CJ, Hermes D, Rao RP, denNijs M, Ojemann JG. Broadband changes in the cortical surface potential track activation of functionally diverse neuronal populations. NeuroImage. 2014 Jan;85:711–20.

57. Winawer J, Kay KN, Foster BL, Rauschecker AM, Parvizi J, Wandell BA. Asynchronous Broadband Signals Are the Principal Source of the BOLD Response in Human Visual Cortex. Current Biology. 2013 Jul 8;23(13):1145–53.

58. Cellier D, Riddle J, Petersen I, Hwang K. The development of theta and alpha neural oscillations from ages 3 to 24 years. Developmental Cognitive Neuroscience. 2021 Aug 1;50:100969.

59. McSweeney M, Morales S, Valadez EA, Buzzell GA, Fox NA. Longitudinal age- and sex-related change in background aperiodic activity during early adolescence. Developmental Cognitive Neuroscience. 2021 Dec;52:101035.

60. Austin EJ. Personality correlates of the broader autism phenotype as assessed by the Autism Spectrum Quotient (AQ). Personality and Individual Differences. 2005;38:451–60.

61. Hurley RSE, Losh M, Parlier M, Reznick JS, Piven J. The Broad Autism Phenotype Questionnaire. J Autism Dev Disord. 2007 Sep 20;37(9):1679–90.

62. Donaldson PH, Kirkovski M, Rinehart NJ, Enticott PG. Autism-relevant traits interact with temporoparietal junction stimulation effects on social cognition: a high-definition transcranial direct current stimulation and electroencephalography study. Eur J Neurosci. 2017;1–13.

63. Hill AT, Van Der Elst J, Bigelow FJ, Lum JAG, Enticott PG. Right anterior theta connectivity predicts autistic social traits in typically developing children. Biological Psychology. 2022 Nov;175:108448.

64. Insel TR. Research Domain Criteria (RDoC): Toward a New Classification Framework for Research on Mental Disorders. American Journal of Psychiatry. 2010;167(7):748–51.

65. Cuthbert BN, Insel TR. Toward the future of psychiatric diagnosis: the seven pillars of RDoC. BMC Med. 2013;11(1):126.

66. Constantino J, Gruber C. The Social Responsiveness Scale Manual. Los Angeles, CA: Western Psychological Services; 2005.

67. Gur RC, Sara R, Hagendoorn M, Marom O, Hughett P, Macy L, et al. A method for obtaining 3-dimensional facial expressions and its standardization for use in neurocognitive studies. Journal of Neuroscience Methods. 2002;115:137–43.

68. Baron-Cohen S, Wheelwright S, Hill J, Raste Y, Plumb I. The '"Reading the Mind in the Eyes"' Test Revised Version: A Study with Normal Adults, and Adults with Asperger Syndrome or High-functioning Autism. J Child Psychol Psychiat. 2001;42(2):241–51.

69. Blakemore SJ. The social brain in adolescence. Nat Rev Neurosci. 2008 Apr;9(4):267–77.

70. Jung M, Tu Y, Lang CA, Ortiz A, Park J, Jorgenson K, et al. Decreased structural connectivity and resting-state brain activity in the lateral occipital cortex is associated with social communication deficits in boys with autism spectrum disorder. NeuroImage. 2019 Apr;190:205–12.

71. Scrivener CL, Reader AT. Variability of EEG electrode positions and their underlying brain regions: visualizing gel artifacts from a simultaneous EEG-fMRI dataset. Brain Behav. 2022 Jan 18;12(2):e2476.

72. Constantino J, Gruber C. The Social Responsiveness Scale Manual, Second Edition (SRS-2). Los Angeles, CA: Western Psychological Services; 2012.

73. Vandierendonck A. A comparison of methods to combine speed and accuracy measures of performance: A rejoinder on the binning procedure. Behav Res. 2017 Apr;49(2):653–73.

74. Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. Journal of Neuroscience Methods. 2004 Mar;134(1):9–21.

75. Bailey NW, Biabani M, Hill AT, Miljevic A, Rogasch NC, McQueen B, et al. Introducing RELAX: An automated pre-processing pipeline for cleaning EEG data - Part 1: Algorithm and application to oscillations. Clinical Neurophysiology. 2023 May;149:178–201.

76. Bailey NW, Hill AT, Biabani M, Murphy OW, Rogasch NC, McQueen B, et al. RELAX part 2: A fully automated EEG data cleaning algorithm that is applicable to Event-Related-Potentials. Clinical Neurophysiology. 2023 May;149:202–22.

77. Bigdely-Shamlo N, Mullen T, Kothe C, Su KM, Robbins KA. The PREP pipeline: standardized preprocessing for large-scale EEG analysis. Front Neuroinform [Internet]. 2015 Jun 18 [cited 2023 Jun 19];9. Available from: http://journal.frontiersin.org/Article/10.3389/fninf.2015.00016/abstract

78. Castellanos NP, Makarov VA. Recovering EEG brain signals: Artifact suppression with wavelet enhanced independent component analysis. Journal of Neuroscience Methods. 2006 Dec;158(2):300–12.

79. Pion-Tonachini L, Kreutz-Delgado K, Makeig S. ICLabel: An automated electroencephalographic independent component classifier, dataset, and website. NeuroImage. 2019 Sep;198:181–97.

80. Aravind Hebbali. olsrr: Tools for Building OLS Regression Models [Internet]. 2020. Available from: https://CRAN.R-project.org/package=olsrr

81. Korkmaz S, Goksuluk D, Zararsiz G. MVN: An R Package for Assessing Multivariate Normality. The R Journal. 2014;6(2):151–62.

82. Yeo IK, Johnson RA. A New Family of Power Transformations to Improve Normality or Symmetry. Biometrika. 2000;87(4):954–9.

83. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Vienna, Austria; 2014.

84. Thuwal K, Banerjee A, Roy D. Aperiodic and Periodic Components of Ongoing Oscillatory Brain Dynamics Link Distinct Functional Aspects of Cognition across Adult Lifespan. eNeuro. 2021 Sep;8(5):ENEURO.0224-21.2021.

85. Gonçalves J, Violante IR, Sereno J, Leitão RA, Cai Y, Abrunhosa A, et al. Testing the excitation/inhibition imbalance hypothesis in a mouse model of the autism spectrum disorder: in vivo neurospectroscopy and molecular evidence for regional phenotypes. Molecular Autism. 2017 Sep 19;8(1):47.

86. Lenroot RK, Gogtay N, Greenstein DK, Wells EM, Wallace GL, Clasen LS, et al. Sexual dimorphism of brain developmental trajectories during childhood and adolescence. NeuroImage. 2007 Jul 15;36(4):1065–73.

87. Johnson SB, Blum RW, Giedd JN. Adolescent Maturity and the Brain: The Promise and Pitfalls of Neuroscience Research in Adolescent Health Policy. J Adolesc Health. 2009 Sep;45(3):216–21.

88. Barry RJ, Clarke AR, Johnstone SJ, Brown CR. EEG differences in children between eyes-closed and eyes-open resting conditions. Clinical Neurophysiology. 2009 Oct 1;120(10):1806–11.