The association between resting state aperiodic activity and Research Domain Criteria
Social Processes in young neurotypical adults

Talitha C. Ford*[1,2], Aron T. Hill[1], Nina-Francesca Parrella[1], Melissa Kirkovski[1,3], Peter Donaldson[1], Peter G. Enticott[1]

1. Cognitive Neuroscience Unit, School of Psychology, Deakin University, Burwood, Victoria, Australia
2. Centre for Mental Health and Brain Sciences, Swinburne University of Technology, Hawthorn, VIC, Australia
3. College of Sport, Health and Engineering, Victoria University, Melbourne, Victoria, Australia

*Corresponding author

Dr. Talitha Ford

Cognitive Neuroscience Unit, School of Psychology, Deakin University

221 Burwood Hwy, Burwood

Victoria 3125, Australia

Abstract:
The aperiodic exponent of the electrophysiological signal has been utilised to demonstrate
differences in brain excitation-inhibition in ageing, cognition, and neuro- and psycho-
pathology. Furthermore, excitation-inhibition imbalance has been associated with social
communication difficulties in clinical and non-clinical cohorts. No work to date, however,
has explored the association between aperiodic activity and social communication
difficulties. A total of 40 neurotypical young adults aged 18-24 (24 female; age mean = 21.6,
SD = 1.9) each underwent 5 minutes of eyes open and eyes closed resting state
electroencephalography using a 64-channel HydroCel Geodesic Sensor Net. Participants also
completed Research Domain Criteria *social processes* measures: Social Responsiveness
Scale, Reading the Mind in the Eyes task, and Penn Emotional Recognition Task. Sex
differences in aperiodic exponent and offset were observed, with larger exponent and
greater offset observed in males, suggestive of greater inhibitory tone. Whole-brain, as well
as left and right social brain, aperiodic exponent were moderately correlated with Social
Responsiveness Scale Social Communication Index, but not Reading the Mind in the Eyes or
Penn Emotional Recognition tasks. However, the correlations between aperiodic exponent
and Social Responsiveness Scale Social Communication Index did not survive multiple
comparisons correction, and regression models demonstrated that neither aperiodic
exponent or offset were significantly predictive of *social processes* outcomes. These
preliminary data provide some evidence that increased aperiodic activity may be associated
with social communication difficulties, however, larger samples are needed to validate
these initial findings.
23
24 **Key words:** aperiodic slope; aperiodic exponent; excitation-inhibition; social
25 communication; autism
26

1. Introduction

Maintaining the equilibrium of excitatory and inhibitory synaptic membrane currents is essential for neural development, information processing, and memory storage (Zhou & Yu, 2018). The brain's primary inhibitory neurotransmitter, gamma-aminobutyric acid (GABA), is responsible for regulating excitatory glutamate, with disturbance to inhibitory tone leading to hyper-excitation. Likewise, disruption to excitatory glutamatergic processes can result in hyper-excitation depending on the location and nature of the disturbance (Sohal & Rubenstein, 2019). An imbalance in brain excitation/inhibition (E/I) ratios biased toward over-excitation has been associated with several psychopathologies and neurodevelopmental conditions, such as autism spectrum disorder (Bruining et al., 2020; Dickinson et al., 2016; Fatemi, 2008; Gogolla et al., 2009; Nelson & Valakh, 2015; Rubenstein & Merzenich, 2003), schizophrenia (Egerton et al., 2020; Marsman et al., 2013), and epilepsy (Bozzi et al., 2018; Hanada, 2020). Due to the heterogeneity and spectrum nature of clinical conditions such as autism and schizophrenia, it is important to investigate the extent to which brain processes relate to specific symptom domains, rather than within the constraints of diagnostic boundaries. The broader autism phenotype reflects the notion that milder autistic traits are present in varying degrees across the non-clinical population (Austin, 2005; Hurley et al., 2007). Indeed, many of the behavioural and neurophysiological differences observed between autism and control groups appear to hold true in non-clinical high autistic trait groups (Abu-Akel et al., 2017; Ajram et al., 2019; Donaldson et al., 2017; Ford, Nibbs, et al., 2017a; Hill et al., 2025; Kirkovski et al., 2016). Thus, extensive evidence supports the utility of non-clinical populations in investigating the behavioural and neurobiological characteristics of autism (Donaldson et al., 2017; Ford, Nibbs, et al., 2017b; Hill, Van Der Elst, et al., 2022).

Both animal and human studies point firmly toward E/I imbalance as a putative marker of social communication difficulties in clinical and non-clinical populations (Bredewold et al., 2015; Cochran et al., 2015; Ford, Nibbs, et al., 2017b; Han et al., 2014; Yizhar et al., 2011). Human neuroscientific studies support the role of brain E/I in social communication (Ajram et al., 2019; Ford & Crewther, 2016), with increased glutamate and/or reduced GABA levels associated with social communication difficulties in clinical (Brown et al., 2013; Cochran et al., 2015; Gaetz et al., 2014; Hegarty et al., 2018; Pretzsch et al., 2019, 2021; Tebartz van Elst et al., 2014) and non-clinical samples (Ford, Nibbs, et al., 2017b, 2017a; Tebartz van Elst et al., 2014). Furthermore, psychopharmacological interventions that enhance brain inhibition, such as cannabidiol (CBD), riluzole, and R-Baclofen (STX209), have been shown to improve social functioning in children and adults with autism (Ajram et al., 2017; Aran et al., 2021; Cifelli et al., 2020; da Silva Junior et al., 2021; Frye, 2014; Hacohen et al., 2022; Parrella et al., 2023). In mice, optogenetic enhancement of excitation led to impairment in social behaviours, while systematically increasing inhibitory tone thereafter partially rescued these impairments (Yizhar et al., 2011). Similarly, low-frequency repetitive transcranial magnetic stimulation (TMS), which is known to inhibit neural activity, improved autism-like social and

behavioural outcomes (Tan et al., 2018). Further, pharmacologically enhancing excitation reduces social behaviours in mice (Bredewold et al., 2015), while enhancing inhibition in mouse models improves social, cognitive, and behavioural outcomes (Han et al., 2014; Stoppel et al., 2018). Taken together, these studies point firmly toward an E/I imbalance contributing to social communication difficulties across clinical lines.

Brain E/I imbalance can disrupt social communication abilities in several ways (Parrella et al., 2024), particularly in social brain regions such as the superior temporal sulcus (STS), temporo-parietal junction (TJP), and medial prefrontal cortex (mPFC) (Blakemore, 2008; Frith & Frith, 2007). E/I imbalance in social brain regions can affect connectivity between brain networks that are responsible for processing social cues (Gao & Penzes, 2015) and sensory sensitivities that heighten anxiety in social situations (Balasco et al., 2020; A. M. Gonçalves & Monteiro, 2023).

Electroencephalography (EEG) has long been utilised as an accessible and affordable tool to probe the relationship between brain and behaviour, particularly in relation to psychiatric and neurodevelopmental conditions (Milovanovic & Grujicic, 2021). Putative markers of excitation and/or inhibition include mismatch negativity (Greenwood et al., 2018; Rowland et al., 2016; Sehatpour et al., 2022), functional E/I ratio (Bruining et al., 2020), and oscillatory activity in delta, theta, and gamma bands (Barttfeld et al., 2011; Coben et al., 2008; Grent-'t-Jong et al., 2018; Lisman, 2012; Uhlhaas & Singer, 2010). Recent developments in resting state and task-related EEG data analysis have revealed that the broadband aperiodic signal, in contrast to periodic activity within a narrowband frequency range (i.e., neuronal oscillations), might be a more accurate reflection of E/I balance (Donoghue et al., 2020; Gao et al., 2017; Waschke et al., 2021). Aperiodic activity is the background, scale-free activity that is represented on a 1/f-like power distribution, where spectral power decreases as frequency increases (Donoghue et al., 2020; Hill, Clark, et al., 2022).

The aperiodic signal, which is comprised of both the exponent (spectral slope) and offset (broadband shift in power across frequencies), is thought to reflect the integration of underlying synaptic currents (Donoghue et al., 2021). These currents have a double-exponential shape in the time domain, which produces the 1/f-like spectrum (Donoghue et al., 2020; Gao et al., 2017). Excitatory currents have a faster time constant and produce constant power at lower frequencies before rapid decay, while power decays more slowly as a function of frequency for inhibitory currents. Thus, a smaller exponent (flatter 1/f-like slope) is thought to reflect more excitation relative to inhibition, while a larger exponent (steeper slope) reflects the converse (Donoghue et al., 2020; Gao et al., 2017). The aperiodic offset (i.e., the y-intercept of the spectrum) is thought to reflect neuronal population spiking (Manning et al., 2009; Miller et al., 2014), which is regulated by GABAergic interneurons (Tremblay et al., 2016). Although the neural underpinnings of the offset are not well

understood, offset and exponent are highly interrelated and thus correlated, suggesting that the offset, at least in part, is reflective of excitatory and/or inhibitory processes. Aperiodic exponent and offset have been associated with age (Cellier et al., 2021; Donoghue et al., 2020; Hill, Clark, et al., 2022; McSweeney et al., 2021), sex (McSweeney et al., 2021), and several cognitive processes such as working memory (Donoghue et al., 2020), visual attention (Waschke et al., 2021), and meta control (Zhang et al., 2023). It is important to note, however, that the exact physiological mechanisms that underlie changes in aperiodic activity are likely complex and remain subject to ongoing investigation. Therefore, interpretation of the functional relevance of aperiodic exponent and offset should be done with caution at this stage. For recent reviews of aperiodic activity within clinical and neurodevelopmental contexts see (Pani et al., 2022; Stanyard et al., 2024).

Given its utility for examining E/I balance across the cortex, and relative affordability and simple acquisition, the aperiodic exponent has been utilised to demonstrate E/I differences in autism (Arutiunian et al., 2024), schizophrenia (Molina et al., 2020), ADHD (Ostlund et al., 2021; Robertson et al., 2019), and Fragile X Syndrome (Wilkinson & Nelson, 2021). To our knowledge, three studies to date have investigated aperiodic exponent in relation to social communication: social-emotional development in toddlers born preterm, with increased aperiodic exponent (increased inhibition) associated with more social emotional problems (Shuffrey et al., 2022); toddlers with neurofibromatosis, with increased exponent associated with higher autistic traits for those with poor executive function (Carter Leno et al., 2022); and typically developing children with autistic traits associated with a larger exponent, when EEG was recorded during a facial emotion processing task (Hill et al., 2025). No work to date has applied aperiodic activity analyses to explore differences in E/I and social communication in a non-clinical adult population.

The Research Domain Criteria (RDoC) highlight the limitations of diagnosis-specific research, and provides a rich framework for understanding psychiatric conditions in the context of varying degrees of symptomatology with neurobiological and/or genetic origins (Insel, 2010). Within the RDoC, the *social processes* domain includes social communication, affiliation and attachment with others, and perception and understanding of the self and others (Cuthbert & Insel, 2013). Self-report measures of *social processes* recommended by RDoC include the Social Responsiveness Scale (SRS; Constantino & Gruber, 2005), while objective measures include the Penn Emotion Recognition (ER-40; Gur et al., 2002), and Reading the Mind in the Eyes (RME; Baron-Cohen et al., 2001) tasks. These measures have been utilised to assess social functioning in both clinical and non-clinical populations. Understanding the neurobiological mechanisms of *social processes*, at both clinical and non-clinical levels, is essential for broader global efforts to identify effective therapeutics to support those with social processing difficulties.

To that end, this study explores the extent to which aperiodic activity predicts RDoC *social*
processes outcomes in young non-clinical adults. In line with the brain E/I hypothesis in
autism, we hypothesised that a reduced aperiodic exponent (flatter slope), indicating
reduced global inhibitory tone (E>I), would predict more *social processes* difficulties. Given
there has been no literature published to date on aperiodic activity and social difficulties,
we also explored this hypothesis specifically in right and left temporo-parietal "social brain"
regions, which are known to be involved in social information processing (Blakemore, 2008;
Frith & Frith, 2007).
9
2. Methods and Materials
2.1. Participants
A total of 40 young adults (24 female, 16 male) aged 18-24 participated in this study (mean
age = 21.6 years, *SD* = 1.9). Participants had an IQ > 90 (range 91-148, mean = 114.6, *SD* =
10.15), measured using the Wechsler Abbreviated Scales of Intelligence (version 2; WASI-
III)(Wechsler, 2011), reflecting "average" (90-109) through to "superior" (>130) intelligence
within the sample. Education levels included 7 participants with secondary college, 29 with a
tertiary degree (undergraduate, diploma, vocational training), and 4 with postgraduate
degree (PhD, Masters). Represented ethnicities were: 30 European, 5 East Asian, 4 South
Asian, 1 Middle Eastern. Participants were excluded if they self-reported: a history of
developmental, psychiatric genetic, or neurological conditions; a first-degree relative with a
history of autism spectrum disorder, schizophrenia, bipolar disorder, or other psychiatric
condition (not including depression and anxiety); a substance use disorder; regular smoking
(> 1 cigarette/day); or taking any psychoactive medications. Participants were asked to
abstain from caffeine and nicotine for 12 hours, and recreational drugs for 1 week, prior to
testing; confirmation of compliance was made verbally on the day of testing. Participants
provided written informed consent prior to any data collection for this study and were
reimbursed $30AUD for their time participating in the study. Ethics approval was granted by
Deakin University's Human Research Ethics Committee (2019-208) in accordance with the
Declaration of Helsinki.
30
2.2. Psychometric and Behavioural Measures
The below measures were administered prior to EEG data collection.
Social Responsiveness Scale Version-2 (SRS; Constantino & Gruber, 2012)*:* The SRS is a 65-
item measure of symptom severity across five autism symptom domains: Social Awareness
(8 items), Social Motivation (11 items), Social Cognition (12 items), Social Communication
(22 items), and Repetitive Behaviours/Restricted Interests (12 items). Responses are
measured on a 4-point scale from 1 (*not true*) to 4 (*almost always true*), and t-scores are
calculated from raw scores to provide an indication of severity of autistic symptoms
("normal": <60, "mild": 60-65, "moderate": 66-75, "severe": >75). The psychometric
properties of the adult self-report measure of the SRS were assessed by Ingersoll et al.
(2011), with good reliabilities across subscales reported in an adult sample (SRS total α =

.95, Social Awareness α = .64, Social Cognition α = .78, Social Communication α = .88, Social
Motivation α = .81, and Repetitive Behaviours/Restricted Interests α = .81). Within the
RDoC framework, the SRS is a measure of Social Communication – reception and production
of non-facial communication. Given our focus on social communication, the SRS Social
Communication Index (SRS SCI) was utilised, which is a composite score of subscales: Social
Awareness, Social Motivation, Social Cognition, and Social Communication (Constantino &
Gruber, 2005, 2012).

Reading the Mind in the Eyes Task (RME; Baron-Cohen et al., 2001)*:* The RME is a 36-item
measure of perception and understanding of conspecific mental states through the
assessment an actor's eyes. The computerised task presents consecutive photos of actor's
eyes in the centre of the screen, and four words (one in each corner of the screen) that
might describe the emotional state of the eye. Participants are to select the word that best
describes the actor's emotion via mouse click. Accuracy on the RME is scored as the
proportion of correct responses (from 0 to 1). The RME is widely used, and has been shown
to exhibit acceptable internal consistency (α = .73)(Kittel et al., 2022). Because participants
are provided a dictionary of the emotional labels presented, reaction time data are not
analysed. Within the RDoC framework, the RME is a measure of Perception and
Understanding of Others – understanding mental states.

Penn Emotion Recognition Task (ER-40; Gur et al., 2002)*:* The ER-40 consists of 40 colour
photographs of actors expressing one of four emotions (happy, sad, anger, or fear) or
neutral expressions. Stimuli are balanced for actors' gender, age, and ethnicity, and for each
emotion category. Four high-intensity and four low-intensity expressions are included for
each emotion. Participants view one photograph at a time and choose the correct emotion
from a list to the right-hand side of the face via mouse click. The ER-40 has shown adequate
test-retest reliability (r = .753) and internal consistency (α = .645) in non-clinical adults
(Pinkham et al., 2016).
A composite measure of accuracy and reaction time, the linear integrated speed-accuracy
score (LISAS), is calculated according to the following formula: LISAS = RT + (S_{RT} / S_{PE}) x PE,
where RT is participant mean reaction time, PE is participant mean proportion of errors, S_{RT}
is the standard deviation of the reaction time, and S_{PE} is the standard deviation of the
proportion or errors (Vandierendonck, 2017). ER-40 LISAS will be referred to as "ER-40
performance" hereafter, with higher scores indicating better performance. Within the RDoC
framework, the ER-40 task is a measure of Social Communication – reception of facial
communication.

    2.3. Electroencephalography data acquisition
EEG data were recorded using a 64-channel HydroCel Geodesic Sensor Net (Electrical
Geodesics, Inc, USA) containing silver/silver-chloride (Ag/AgCl) electrodes surrounded by
electrolyte-wetted sponges. Electrical Geodesics Net Station EEG software (version 5.0) was

used to acquire the data via a Net Amps 400 amplifier using a sampling rate of 1 KHz. Data were referenced to electrode Cz and grounded to the common of the isolated amp circuit's power supply. Electrode impedances were checked to ensure they were < 50 KOhms prior to data collection. Participants were seated in a quiet, dimly lit room and asked to sit still with their eyes open while looking at a fixation cross for five minutes, and then eyes closed for five minutes, while resting state data were collected. Electrode impedances were checked between eyes open and eyes closed blocks to ensure they remained < 50 KOhms (considered low impedance for this system). One participant withdrew from the study and did not undergo EEG (remaining n = 39).

2.4. EEG pre-processing

EEG data were pre-processed in MATLAB (R2021a; The Mathworks, Massachusetts, USA) incorporating the EEGLAB toolbox (v2023.0; Delorme & Makeig, 2004) and custom scripts. The Reduction of Electroencephalographic Artifacts (RELAX; v1.1.2) software (Bailey, Biabani, et al., 2023; Bailey, Hill, et al., 2023) was used to clean each EEG dataset. The fully automated RELAX pipeline uses empirical approaches to identify and reduce non-neural artifacts, and includes the use of both multi-channel Wiener filters and wavelet enhanced independent component analysis (ICA). As part of the RELAX pipeline, the data were bandpass filtered between 0.5 and 80 Hz (fourth-order zero-phase Butterworth filter). A notch filter (47-53 Hz) was then applied to remove any line noise. Any bad channels were identified and removed using a multi-step process incorporating the default parameters of the 'findNoisyChannels' function (PREP pipeline; Bigdely-Shamlo et al., 2015). Multi-channel Wiener filtering (Castellanos & Makarov, 2006) was used to initially clean blinks, muscle activity, and horizontal eye movement, followed by robust average re-referencing (Bigdely-Shamlo et al., 2015), and wavelet-enhanced ICA (Castellanos & Makarov, 2006), with components for cleaning identified using the automated independent component (IC) classifier IClabel (Pion-Tonachini et al., 2019). Any rejected electrodes were then interpolated back into the data using spherical interpolation and all pre-processed data files were visually inspected prior to further analysis (Eyes closed data: mean rejected channels = 4.89, SD = 3.52; Eyes open data: mean rejected channels = 7.75, SD = 3.72).

2.5. Electroencephalography data analysis

All pre-processed resting state data were visually inspected for quality assurance prior to inclusion in statistical analyses. Two participants were excluded from the study due to excessive electrical line noise for both eyes open and eyes closed conditions (remaining n = 37) that persisted after cleaning, while an additional two participants' eyes open data were excluded due to excessive line noise (eyes open n = 35). The average length of continuous data (in seconds) after cleaning for the eyes open condition was 299.27 seconds (SD = 25.13), and for the eyes-closed condition was 304.70 seconds (SD = 32.29). The power spectral density of the EEG signal was calculated using Welch's method (2-second Hamming window, 50% overlap) for all electrodes across all participants for both the eyes-open and

eyes-closed datasets. The specparam (formally FOOOF) toolbox (version 1.0.0;

https://fooof-tools.github.io/fooof/; Donoghue et al., 2020) was then used to parameterise

the spectral data through separation of the periodic and aperiodic components of the signal.

The aperiodic exponent and offset were extracted across 1 to 40 Hz using the 'fixed'

aperiodic mode. Spectral parameterization settings for the algorithm were based on

recommendations on the specparam toolbox website (https://fooof-tools.github.io/fooof/)

and past literature (Ostlund et al., 2022): peak width limits = [1, 12], maximum number of

peaks = 8, peak threshold = 2, minimum peak height = .05.

The performance of the FOOOF algorithm for fitting the data was assessed using two

"goodness of fit" measures: R^2, explained variance; and mean absolute error (MAE), total

error of the model fit (Donoghue et al., 2020; Ostlund et al., 2021). Model fits for both eyes

open and eyes closed data were good (eyes open: R^2 = .99, MAE = .047; eyes closed: R^2 = .99,

MAE = .054. Complete fit statistics are presented in the Supplementary Table 1). There was

no difference in mean R^2 between eyes open and eyes closed conditions ($t(64.13)$ = -.30, p =

.765), while MAE for eyes open was significantly lower compared to eyes closed ($t(69.44)$ =

2.19, p = .032). Aperiodic exponent (slope) was significantly larger for the eyes closed

condition, compared to the eyes open condition ($t(69.65)$ = 2.82, p = .006). Similarly, the

offset (intercept) was larger for eyes closed than eyes open condition ($t(69.97)$ = 3.0, p =

.004). There were strong, positive correlations between exponent and offset values for eyes

open (*rho* = .73, p < .001) and eyes closed (*rho* = .80, p < .001) recordings, and between eyes

open and closed exponent (*rho* = .73, p < .001) and eyes open and closed offset (*rho* = .79, p

< .001).

To calculate aperiodic exponent and offset in brain regions associated with social

information processing, so called "social brain" (Blakemore, 2008; Moessnang et al., 2016),

electrodes over left and right temporo-parietal regions were averaged separately (left: P5

CP5 TP9 TP7 C5 T7 T9; right: P6 CP6 TP10 TP8 C6 T8 T10; see Figure 1). These electrodes

cover brain regions including the superior, middle, and inferior temporal gyrus,

supramarginal gyrus, and angular gyrus, all known to be involved in social information

processing (Blakemore, 2008; Jung et al., 2019; Moessnang et al., 2016).

1
2 **Figure 1: Electroencephalograph electrode placement.** 64-channel Geodesic Sensor Net
3 electrode placement (dark circles), with the cluster of electrodes selected to represent left
4 (blue) and right (orange) social brain highlighted.
5
2.6. Statistical analyses
Statistical analyses were conducted in R (version 4.1.0). Given this is the first study to
examine aperiodic activity in relation to social information processing, *a priori* power
analyses were not conducted. However, previous studies have reported significant
associations between brain E/I and social communication with sample sizes of 29 to 37 non-
clinical adults (Ford, Nibbs, et al., 2017a; Ford, Woods, et al., 2017; Tebartz van Elst et al.,
2014). Assumptions for all statistical analyses were assessed using the *olsrr* and *MVN*
packages in R (Aravind Hebbali, 2020; Korkmaz et al., 2014), including normality, linearity,
homoscedasticity, and multicollinearity. All assumptions were met, except for a significant
positive skew in RME and negative skew in ER-40 LISAS scores, which were subsequently
normalised using the Yeo-Johnson transform in R (VGAM package; Yeo & Johnson, 2000).
17
Initial bivariate Pearson's r correlation analyses were conducted to investigate the
association between *social processes* outcomes (SRS SCI score, RME accuracy, and ER-40
performance) and whole-brain aperiodic exponent and offset with eyes open and eyes
closed. Additional exploratory correlations analyses were conducted to investigate the
association between *social processes* outcomes and left and right social brain exponent and
offset. The significance level was set at p = .05, and false discovery rate (FDR) was corrected
for using the Benjamin and Hochberg method in R (Benjamini & Hochberg, 1995; stats
package; R Core Team, 2023).
26
27 Then, multivariate linear regressions (MLRs) were conducted to assess the extent to which
28 aperiodic exponent and offset with eyes open and eyes closed predicted *social processes*

outcomes. Due to the high correlation between EO and EC exponent and offset, four separate MLR models were defined (i.e., model 1 predictor: eyes open exponent; model 2 predictor: eyes closed exponent; model 3 predictor: eyes open offset; model 4 predictor: eyes closed offset). There were significant sex effects for exponent and offset in both conditions, therefore, sex was entered as a covariate for each model (see Supplementary Material for full model designs). Additional exploratory MLR models were defined to investigate the extent to which aperiodic activity in left and right social brain predicted *social processes* outcomes; again, sex was entered as a covariate (see Supplementary Material for full model designs). The significance level for each model was set at $p = .05$, and FDR was corrected for using the Benjamini and Hochberg method in R (Benjamini & Hochberg, 1995; stats package; R Core Team, 2023).

3. Results

    3.1. Descriptive statistics

Descriptive statistics for final participant sample characteristics ($n = 37$), *social processes* outcomes, and aperiodic activity metrics are presented in Table 1. There were no significant age or biological sex effects on *social processes* measures ($ps > .05$), and no significant correlations between age and aperiodic activity outcomes ($ps > .05$). However, males compared to females had a significantly larger exponent ($t(24.20) = -2.56$, $p = .017$, $p_{FDR} = .033$) and higher offset ($t(29.20) = -2.08$, $p = .047$, $p_{FDR} = .047$) during eyes open, as well as a larger exponent ($t(33.76) = -2.83$, $p = .008$, $p_{FDR} = .032$) and higher offset during eyes closed ($t(34.90) = -2.34$, $p = .025$, $p_{FDR} = .033$; Figure 2).

1   Table 1: Descriptive statistics of psychometric, behavioural, and aperiodic activity measures

	Total (n = 37)		Female (n = 22)	Male (n = 15)
	Mean (SD)	Min-Max	Mean (SD)	Mean (SD)
Age	22 (1.9)	18 – 25	21 (1.9)	22 (1.9)
WASI IQ	115 (9.8)	91 – 148	114 (8.2)	117 (12)
Social Processes				
SRS total t-score	52 (6.8)	42 – 69	51 (5.8)	54 (8.4)
SRS SCI t-score	53 (6.9)	42 – 71	52 (5.6)	54 (8.6)
SRS RRB t-score	53 (8.4)	40 – 73	51 (7.2)	55 (10)
RME accuracy	.81 (.08)	.64 – .94	.83 (.08)	.78 (.08)
ER-40 RT (sec)	2.9 (.69)	2.0 – 4.9	2.8 (.48)	3.0 (.94)
ER-40 accuracy	.89 (.05)	.78 – .97	.90 (.04)	.86 (.05)
ER-40 performance	3.3 (.97)	2.2 – 7.0	3.1 (.65)	3.5 (1.3)
Aperiodic activity				
Whole brain				
EO exponent (a.u.)	1.2 (.17)	.96 – 1.7	1.2 (.14)	1.3 (.19)
EC exponent (a.u.)	1.3 (.17)	.95 – 1.7	1.3 (.17)	1.4 (.14)
EO offset (μV^2)	.21 (.38)	-.47 – 1.4	.10 (.36)	.36 (.38)
EC offset (μV^2)	.49 (.41)	-.63 – 1.3	.37 (.44)	.66 (.31)
Left social brain				
EO exponent (a.u.)	1.2 (.19)	.96 – 1.7	1.2 (.15)	1.3 (.21)
EC exponent (a.u.)	1.3 (.20)	.86 – 1.7	1.3 (.20)	1.4 (.15)
EO offset (μV^2)	.16 (.41)	-.50 – 1.5	.014 (.34)	.35 (.43)
EC offset (μV^2)	.42 (.45)	-.67 – 1.3	.25 (.45)	.66 (.35)
Right social brain				
EO exponent (a.u.)	1.2 (.20)	.67 – 1.7	1.2 (.18)	1.3 (.19)
EC exponent (a.u.)	1.3 (.20)	.77 – 1.6	1.2 (.21)	1.4 (.13)
EO offset (μV^2)	.18 (.42)	-.84 – 1.3	.026 (.37)	.38 (.41)
EC offset (μV^2)	.43 (.48)	-1.0 – 1.2	.26 (.50)	.67 (.34)

2   Notes: SD = standard deviation, WASImIQ = Weschler Abbreviated Scale of Intelligence
3   Intelligence Quotient, SRS = Social Responsiveness Scale, SCI = Social Communication Index,
4   RRB = Restricted Repetitive Behaviours, RME = Reading the Mind in the Eyes, ER-40 = Penn
5   Emotion Recognition Task, RT = reaction time, performance = linear integrated speed-
6   accuracy score (LISAS), EO = eyes open, EC = eyes closed.

1
2 **Figure 2: Sex differences in aperiodic activity.** Sex differences in exponent (a) and offset (b)
3 for eyes open and eyes closed conditions, and power vs frequency plots by sex for eyes
4 open (c) and eyes closed (d) conditions showing the aperiodic component of the neural
5 signal. Thin lines represent the aperiodic signal for individual participants, thick lines
6 represent the average across all participants. *p_{FDR}<.05
7
3.2. Social processes outcomes
Pearson's r correlations were conducted to test the relationship between the three *social
processes* measures. There were no significant relationships between SRS SCI scores and
RME accuracy or ER-40 performance (p_{FDR} > .05). The relationship between ER-40
performance and RME accuracy was significant, with ER-40 performance increasing with
RME accuracy as would be expected (r = .434, p = .018, p_{FDR} = .025).
14
15   3.3. Correlations between aperiodic activity and *social processes* outcomes

Pearson's r correlations between *social processes* measures and whole-brain and social brain aperiodic activity are presented in Table 2. There were significant moderate correlations between SRS SCI score and whole brain exponent, left social brain exponent and offset, and right social brain exponent in the eyes open condition only (uncorrected $p <$ .05), with aperiodic activity increasing with more social communication difficulties. However, these correlations did not survive FDR correction ($p_{FDR} >$ .05). There were no significant correlations between RME accuracy and ER-40 performance and aperiodic activity across whole-brain or social brain regions (uncorrected $p >$ .05). Scatterplots for these associations are presented in Figure 3 and Figure 4.

Table 2: Pearson's r correlations between RDoC *social processes* measures and aperiodic activity.

	SRS SCI score	RME accuracy	ER-40 performance
Whole brain			
EO Exponent (a.u.)	.34[a]	-.29	-.09
EC Exponent (a.u.)	.09	-.20	-.04
EO Offset (μV^2)	.30	-.32	-.11
EC Offset (μV^2)	.10	-.21	.06
Left social brain			
EO Exponent (a.u.)	.35[a]	-.23	-.12
EC Exponent (a.u.)	.14	-.21	-.12
EO Offset (μV^2)	.34[a]	-.27	-.14
EC Offset (μV^2)	.14	-.21	-.01
Right social brain			
EO Exponent (a.u.)	.40[a]	-.22	-.04
EC Exponent (a.u.)	.14	-.17	.04
EO Offset (μV^2)	.33	-.24	-.10
EC Offset (μV^2)	.15	-.19	.13

Notes: SRS = Social Responsiveness Scale, SCI = Social Communication Index, RME = Reading the Mind in the Eyes, ER-40 = Penn Emotion Recognition Task, EO = eyes open, EC = eyes closed. [a]correlations significant at $p <$ .05 prior to correction.

3.4. Whole-brain aperiodic activity and social processes outcomes

Multivariate linear regression models were designed to test the extent to which aperiodic exponent and offset predicted *social processes* outcomes of SRS SCI scores, RME accuracy, and ER-40 performance. Four models were defined with exponent and offset for eyes open and eyes closed entered as separate predictors. None of the models were significant (uncorrected $p >$ .05) suggesting that neither exponent or offset predicted *social processes* outcomes in either eyes open or eyes closed conditions (see Table 3 for model statistics).

1 Figure 3 illustrates the relationships between each of predictor and outcome measures, and
2 suggests a visual trend toward increased exponent (i.e., increased inhibitory tone) and
3 offset with higher SRS SCI scores, and poorer RME and ER-40 performance, particularly in
4 the eyes open condition.
5
6 Table 3: Multivariate linear regression model results for whole-brain aperiodic activity and
7 RDoC social processes measures.

	R^2	Adj. R^2	F (df)	p_{uncorr}	p_{FDR}	Cohens f^2
Exponent						
Model 1: Eyes open						
SRS SCI	.119	.063	2.10 (2,31)	.139	.334	.119
RME accuracy	.131	.075	2.43 (2,31)	.113	.334	.151
ER-40 performance	.008	-.056	.23 (2,31)	.881	.956	.008
Model 2: Eyes closed						
SRS SCI	.038	-.020	.07 (2,33)	.530	.795	.039
RME accuracy	.115	.061	2.15 (2,33)	.133	.334	.130
ER-40 performance	.003	-.058	.05 (2,33)	.956	.956	.003
Offset						
Model 3: Eyes open						
SRS SCI	.103	.045	1.79 (2,31)	.185	.370	.115
RME accuracy	.148	.093	2.70 (2,31)	.083	.334	.174
ER-40 performance	.008	-.056	.12 (2,31)	.887	.956	.008
Model 4: Eyes closed						
SRS SCI	.039	-.020	.66 (2,33)	.522	.795	.040
RME accuracy	.119	.065	2.22 (2,33)	.124	.334	.135
ER-40 performance	.012	-.050	.21 (2,33)	.815	.956	.013

8 Notes: Adj = Adjusted, df = degrees of freedom, uncorr = uncorrected, FDR = false discovery
9 rate, SRS = Social Responsiveness Scale, SCI = Social Communication Index, RME = Reading
10 the Mind in the Eyes, ER-40 = Penn Emotion Recognition Task, RDoC = Research Domain
11 Criteria.
12
13

1
2 **Figure 3: Scatter plots of aperiodic activity and Social Processes outcomes.** Scatterplots of
3 depicting exponent (top row; model 1 & 2) and offset (bottom row; model 3 & 4) versus
4 Social Responsiveness Scale Social Communication Index score (SRS SCI; left panel), Reading
5 the Mind in the Eyes (RME) accuracy (middle panel), and Penn Emotion Recognition (ER-40)
6 performance (right panel) in eyes open (green) and eyes closed (red) conditions.
7
3.5. Social brain aperiodic activity and social processes outcomes
Exploratory analyses were conducted to investigate whether aperiodic activity in temporal-
parietal regions known to be involved in social information processing (Blakemore, 2008;
Jung et al., 2019; Moessnang et al., 2016) might predict *social processes* outcomes (SRS SCI,
RME accuracy, ER-40 performance). Given the lack of effects of the eyes closed condition in
the whole-brain analyses, only eyes open data were analysed here. Therefore, measures of
exponent and offset for eyes open were averaged over temporo-parietal clusters in left and
right hemisphere, and four models were defined for left and right exponent and offset
separately (i.e., left exponent, left offset, right exponent, right offset) (see Supplementary
Material for models). Figure 4 illustrates the relationships between *social processes*
outcomes and predictors. None of the overall models for SRS SCI, ER-40 performance, and
RME accuracy for both exponent and offset predictors were significant (uncorrected *p*s >
.05, see Table 4 for model statistics). Despite the non-significant finding, there was a trend
toward increased right social brain exponent (i.e., increased inhibitory tone) with higher SRS

SCI scores (p = .068), with a moderate effect size (f^2 = .189). When visualising the data in Figure 4, exponent and offset appeared to increase with higher SRS SCI scores and poorer RME accuracy over both social brain regions.

Table 4: Multivariate linear regression model results for left and right social brain aperiodic activity and RDoC social processes measures.

	R^2	Adj. R^2	F (df)	p_{uncorr}	p_{FDR}	Cohen's f^2
Exponent						
Model 1: Left						
SRS SCI	.129	.072	2.29 (2,31)	.118	.251	.148
RME accuracy	.114	.056	1.99 (2,31)	.154	.251	.128
ER-40 performance	.013	-.050	.208 (2,31)	.813	.976	.013
Model 2: Right						
SRS SCI	.159	.105	2.94 (2,31)	.068	.251	.189
RME accuracy	.109	.052	1.90 (2,31)	.167	.251	.123
ER-40 performance	.000	-.064	.00 (2,31)	.998	.998	.000
Offset						
Model 3: Left						
SRS SCI	.127	.064	2.13 (2,31)	.136	.251	.137
RME accuracy	.124	.067	2.20 (2,31)	.128	.251	.142
ER-40 performance	.020	-.043	.32 (2,31)	.729	.972	.021
Model 4: Right						
SRS SCI	.114	.057	2.00 (2,31)	.152	.251	.129
RME accuracy	.115	.057	2.01 (2,31)	.152	.251	.129
ER-40 performance	.003	-.061	.054 (2,31)	.948	.998	.003

Notes: Adj = Adjusted, df = degrees of freedom, uncorr = uncorrected, FDR = false discovery rate, SRS = Social Responsiveness Scale, SCI = Social Communication Index, RME = Reading the Mind in the Eyes, ER-40 = Penn Emotion Recognition Task, RDoC = Research Domain Criteria.

1
2 **Figure 4: Scatter plots of aperiodic activity by hemisphere and Social Processes outcomes.**
3 Scatter plots of exponent (top row; model 1 & 3) and offset (bottom; model 2 & 4) versus
4 Social Responsiveness Scale Social Communication Index score (SRS SCI; left panel), Reading
5 the Mind in the Eyes (RME) accuracy (middle panel), and Penn Emotion Recognition (ER-40)
6 performance (right panel) in left (red) and right (green) social brain regions.
7
4.   Discussion
Investigations of aperiodic activity (exponent and offset) are increasingly reported across
neurodevelopmental, cognitive, and psychiatric datasets as a putative measure of brain E/I
ratios (Molina et al., 2020; Ostlund et al., 2021; Robertson et al., 2019; Thuwal et al., 2021;
Wilkinson & Nelson, 2021), with a larger exponent (steeper aperiodic slope) typically
interpreted as reflecting increased inhibitory tone (i.e., reduced E/I). This study was the first,
to our knowledge, to investigate the extent to which resting state aperiodic activity predicts
RDoC *social processes* outcomes in non-clinical young adults.
16
Given a shift in E/I towards increased excitation has been associated with social
communication difficulties in clinical, non-clinical, and animal studies (Bredewold et al.,
2015; Cochran et al., 2015; Ford, Nibbs, et al., 2017b; Han et al., 2014), it was predicted that
reduced aperiodic exponent (i.e., flatter slope) would be associated with more social
communication and processing difficulties. There was a moderate correlation between SRS
SCI scores and the aperiodic exponent averaged across all electrodes (i.e., whole brain)

indicating that, contrary to expectations, increased inhibitory tone was associated with
more social communication difficulties. However, this association did not survive correction
for FDR. Furthermore, whole-brain aperiodic exponent did not significantly predict *social*
processes outcomes in either the eyes open or eyes closed condition. This may indicate that:
1) predictive power of exponent and offset is possibly region-specific, thus including all
electrodes introduces substantial noise that effectively drowns out any signal; 2) there was
insufficient power to detect an effect; and/or 3) E/I measured by aperiodic brain activity
does not predict the RDoC *social processes* measures utilised here.
9
To explore the regional specificity of aperiodic activity to *social processes*, two temporo-
parietal 'social brain' clusters were created for left and right hemisphere that localised to
superior, middle, and inferior temporal gyrus, supramarginal gyrus, and angular gyrus
(Scrivener & Reader, 2022). These regions are known to be involved in social information
processing (Blakemore, 2008; Jung et al., 2019), particularly in the right cortical hemisphere
(Abu-Akel et al., 2017; Ford, Nibbs, et al., 2017a; Lindell, 2006; von dem Hagen et al., 2011).
Pearson correlations indicated that increased left and right social brain exponent values
(more inhibitory tone) were associated with higher SRS SCI scores in our non-clinical sample
with a moderate effect size, however, when controlling for false discovery rate, this
relationship was no longer statistically significant. There were no associations with RME
accuracy or ER-40 performance. Results of the regression models suggested that aperiodic
activity within these left and right social brain regions did not significantly predict any of the
social processes outcomes. Although non-significant, aperiodic exponent in social brain
regions were better at predicting SRS SCI scores with moderate effect sizes compared to
small effect sizes for whole-brain analysis.
25
Although there were no significant findings, the trend toward increased exponent
(suggestive of greater inhibitory tone) and offset with more *social processes* difficulties
warrants further investigation in a larger, more diverse samples, including clinical
populations where social communication difficulties are a core feature. Only two studies to
date have utilised aperiodic activity to investigate the relationship between brain E/I and
social communication – both of which are in clinical young children. Therefore, this study is
the first to specifically probe social communication difficulties and E/I via aperiodic activity.
Extensive research suggests that social communication difficulties in non-clinical and clinical
groups are associated with an increase in E/I (Rubenstein & Merzenich, 2003; Yizhar et al.,
2011), implying increased excitation and/or reduced inhibition. However, more recent
research has suggested that the nature of the E/I imbalance may be region specific (Ajram et
al., 2019; Ford & Crewther, 2016; Nelson & Valakh, 2015), and regional specificity of
increased inhibitory tone has been reported in mouse models (J. Gonçalves et al., 2017). The
slightly increased aperiodic exponent, potentially reflecting a non-invasive index of greater
inhibitory tone in social brain regions, in conjunction with more social communication
difficulties observed in this study may reflect underlying neurobiological mechanisms. This is

in line with a study in children demonstrating increased exponent with more pronounced

autistic traits (Hill et al., 2025) and increased exponent for autistic children with IQ>85

compared to typically developing children and autistic children with IQ<85 (Manyukhina et

al., 2022). These findings together indicate that the aperiodic exponent in relation to social

communication difficulties should be further examined in larger samples to better

understand the nature of neurobiological mechanisms across the autism spectrum.

In line with previous literature, males had a significantly higher aperiodic offset and larger

exponent than females in both eyes open and eyes closed conditions (Hill, Clark, et al., 2022;

McSweeney et al., 2021). Sex differences in a prior adolescent sample were thought to

reflect an advanced neurophysiological maturity in females (Lenroot et al., 2007), which

may be the case in this young adult sample as neurodevelopment continues into early

adulthood for both males and females (Johnson et al., 2009). Contrary to previous reports

(Cellier et al., 2021; Donoghue et al., 2020; Hill, Clark, et al., 2022; McSweeney et al., 2021),

exponent and offset were not associated with age, which may reflect that aperiodic activity

is relatively stable between 18 and 25 years, or that the age range of the sample was too

narrow to detect age effects. Also in line with previous research in children and adolescents,

larger exponent and higher offset were recorded in the eyes closed compared to eyes open

condition (Hill, Clark, et al., 2022; McSweeney et al., 2021). This reflects a reduction in E/I

when eyes are closed, which could be explained by the lack of visual stimulation, attention,

and arousal, compared to when participants' eyes were open (Barry et al., 2009).

There were no significant relationships between RDoC *social processes* self-report (SRS SCI)

and behavioural (RME, ER-40) outcome measures themselves. This was somewhat

unexpected and possibly problematic given the three measures, SRS SCI, RME, and ER-40,

are proposed by the RDoC as gold standard measures of the *social processes* constructs.

Although they appear to index different aspects of *social processes*, namely social

communication (reception of facial communication [ER-40], and reception and production

of non-facial [SRS SCI]), and perception and understanding of others (understanding mental

states [RME]), it is reasonable to expect that they would be related. The lack of

interrelationship in this study may reflect the limited spread in performance/difficulties in

the tasks in this non-clinical sample, insufficient data, or that the measures have poor

construct validity, potentially due to poor insight when completing self-report measures. In

fact, a recent study demonstrated the RME may lack the psychometric properties to

accurately measure theory of mind, compared to the original validation study, and

therefore, the lack of associations identified in this study might be due to inadequate

measurement tools (Higgins et al., 2023).

The main limitations for this study were the modest sample size (*n* = 37) and narrow spread

of *social processes* difficulties, particularly for SRS scores (range = 42-69). These issues

combined may have been a significant contributor to the lack of statistically significant

findings, given the small effect sizes reported herein. A more diverse spread of abilities within the sample, for example including clinical participants, may reveal more compelling relationships between aperiodic activity and social processing abilities. Furthermore, EEG is limited in its ability to examine social brain areas specifically due to its relatively poor spatial resolution. Source localization using magnetoencephalography (MEG) and concurrent participant structural brain scans would allow for a more accurate analysis of social brain and *social processes* abilities.

Despite these limitations, these preliminary data show promise for supporting the utility of aperiodic activity in investigating the relationship between brain E/I and social communication difficulties in non-clinical, as well as clinical, populations. Future work should seek to examine this relationship in a larger sample across clinical and non-clinical groups. Furthermore, studies involving pharmacological interventions that modulate excitatory and/or inhibitory processes would provide valuable insights into the utility of the aperiodic exponent as a putative neurobiomarker of *social processes* abilities.

Acknowledgements: Thank you to Dr Kaila Bianco, Dr Pam Barhoun, and Mr Ji-Shen Loong for assistance in collecting data for this study.

Data availability: The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

Abu-Akel, A., Apperly, I. A., Wood, S. J., & Hansen, P. C. (2017). Autism and psychosis expressions diametrically modulate the right temporoparietal junction. *Social Neuroscience*, *12*(5), 506–518. https://doi.org/10.1080/17470919.2016.1190786

Ajram, L. A., Horder, J., Mendez, M. A., Galanopoulos, A., Brennan, L. P., Wichers, R. H., Robertson, D. M., Murphy, C. M., Zinkstok, J., Ivin, G., Heasman, M., Meek, D., Tricklebank, M. D., Barker, G. J., Lythgoe, D. J., Edden, R. A. E., Williams, S. C., Murphy, D. G. M., & McAlonan, G. M. (2017). Shifting brain inhibitory balance and connectivity of the prefrontal cortex of adults with autism spectrum disorder. *Translational Psychiatry*, *7*(5), e1137. https://doi.org/10.1038/tp.2017.104

Ajram, L. A., Pereira, A. C., Durieux, A. M. S., Velthius, H. E., Petrinovic, M. M., & McAlonan, G. M. (2019). The contribution of [1H] magnetic resonance spectroscopy to the study of excitation-inhibition in autism. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*. https://doi.org/10.1016/j.pnpbp.2018.09.010

Aran, A., Harel, M., Cassuto, H., Polyansky, L., Schnapp, A., Wattad, N., Shmueli, D., Golan,

D., & Castellanos, F. X. (2021). Cannabinoid treatment for autism: A proof-of-concept

randomized trial. *Molecular Autism*, *12*(1), 6. https://doi.org/10.1186/s13229-021-

00420-2

Aravind Hebbali. (2020). *olsrr: Tools for Building OLS Regression Models* (Version R Package

version 0.5.3.) [Computer software]. https://CRAN.R-project.org/package=olsrr

Arutiunian, V., Arcara, G., Buyanova, I., Fedorov, M., Davydova, E., Pereverzeva, D., Sorokin,

A., Tyushkevich, S., Mamokhina, U., Danilina, K., & Dragoy, O. (2024). Abnormalities

in both stimulus-induced and baseline MEG alpha oscillations in the auditory cortex

of children with Autism Spectrum Disorder. *Brain Structure and Function*, *229*(5),

1225–1242. https://doi.org/10.1007/s00429-024-02802-7

Austin, E. J. (2005). Personality correlates of the broader autism phenotype as assessed by

the Autism Spectrum Quotient (AQ). *Personality and Individual Differences*, *38*, 451–

460. https://doi.org/10.1016/j.paid.2004.04.022

Bailey, N. W., Biabani, M., Hill, A. T., Miljevic, A., Rogasch, N. C., McQueen, B., Murphy, O.

W., & Fitzgerald, P. B. (2023). Introducing RELAX: An automated pre-processing

pipeline for cleaning EEG data - Part 1: Algorithm and application to oscillations.

Clinical Neurophysiology, *149*, 178–201.

https://doi.org/10.1016/j.clinph.2023.01.017

Bailey, N. W., Hill, A. T., Biabani, M., Murphy, O. W., Rogasch, N. C., McQueen, B., Miljevic,

A., & Fitzgerald, P. B. (2023). RELAX part 2: A fully automated EEG data cleaning

algorithm that is applicable to Event-Related-Potentials. *Clinical Neurophysiology*,

149, 202–222. https://doi.org/10.1016/j.clinph.2023.01.018

Balasco, L., Provenzano, G., & Bozzi, Y. (2020). Sensory Abnormalities in Autism Spectrum

Disorders: A Focus on the Tactile Domain, From Genetic Mouse Models to the Clinic.

Frontiers in Psychiatry, *10*, 1016. https://doi.org/10.3389/fpsyt.2019.01016

Baron-Cohen, S., Wheelwright, S., Hill, J., Raste, Y., & Plumb, I. (2001). The '"Reading the

Mind in the Eyes"' Test Revised Version: A Study with Normal Adults, and Adults with

Asperger Syndrome or High-functioning Autism. *Journal of Child Psychology and*

Psychiatry, *42*(2), 241–251. https://doi.org/10.1111/1469-7610.00715

Barry, R. J., Clarke, A. R., Johnstone, S. J., & Brown, C. R. (2009). EEG differences in children
between eyes-closed and eyes-open resting conditions. *Clinical Neurophysiology*,
120(10), 1806–1811. https://doi.org/10.1016/j.clinph.2009.08.006
Barttfeld, P., Wicker, B., Cukier, S., Navarta, S., Lew, S., & Sigman, M. (2011). A big-world
network in ASD: dynamical connectivity analysis reflects a deficit in long-range
connections and an excess of short-range connections. *Neuropsychologia*, *49*(2),
254–263. https://doi.org/10.1016/j.neuropsychologia.2010.11.024
Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and
powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series*
B (Methodological), 289–300.
Bigdely-Shamlo, N., Mullen, T., Kothe, C., Su, K.-M., & Robbins, K. A. (2015). The PREP
pipeline: Standardized preprocessing for large-scale EEG analysis. *Frontiers in*
Neuroinformatics, *9*. https://doi.org/10.3389/fninf.2015.00016
Blakemore, S.-J. (2008). The social brain in adolescence. *Nature Reviews Neuroscience*, *9*(4),
267–277. https://doi.org/10.1038/nrn2353
Bozzi, Y., Provenzano, G., & Casarosa, S. (2018). Neurobiological bases of autism-epilepsy
comorbidity: A focus on excitation/inhibition imbalance. *European Journal of*
Neuroscience, *47*(6), 534–548. https://doi.org/10.1111/ejn.13595
Bredewold, R., Schiavo, J. K., van der Hart, M., Verreij, M., & Veenema, A. H. (2015).
Dynamic changes in extracellular release of GABA and glutamate in the lateral
septum during social play behavior in juvenile rats: Implications for sex-specific
regulation of social play behavior. *Neuroscience*, *307*, 117–127.
https://doi.org/10.1016/j.neuroscience.2015.08.052
Brown, M. S., Singel, D., Hepburn, S., & Rojas, D. C. (2013). Increased glutamate
concentration in the auditory cortex of persons with autism and first-degree
relatives: A (1)H-MRS study. *Autism Res*, *6*(1), 1–10.
https://doi.org/10.1002/aur.1260
Bruining, H., Hardstone, R., Juarez-Martinez, E. L., Sprengers, J., Avramiea, A.-E., Simpraga,
S., Houtman, S. J., Poil, S.-S., Dallares, E., Palva, S., Oranje, B., Matias Palva, J.,
Mansvelder, H. D., & Linkenkaer-Hansen, K. (2020). Measurement of excitation-
inhibition ratio in autism spectrum disorder using critical brain dynamics. *Scientific*
Reports, *10*(1), 9195. https://doi.org/10.1038/s41598-020-65500-4

Carter Leno, V., Begum-Ali, J., Goodwin, A., Mason, L., Pasco, G., Pickles, A., Garg, S., Green, J., Charman, T., Johnson, M. H., Jones, E. J. H., the EDEN, Vassallo, G., Burkitt-Wright, E., Eelloo, J., Gareth Evans, D., West, S., Hupton, E., Lewis, L., … Taylor, C. (2022). Infant excitation/inhibition balance interacts with executive attention to predict autistic traits in childhood. *Molecular Autism*, *13*(1), 46. https://doi.org/10.1186/s13229-022-00526-1

Castellanos, N. P., & Makarov, V. A. (2006). Recovering EEG brain signals: Artifact suppression with wavelet enhanced independent component analysis. *Journal of Neuroscience Methods*, *158*(2), 300–312. https://doi.org/10.1016/j.jneumeth.2006.05.033

Cellier, D., Riddle, J., Petersen, I., & Hwang, K. (2021). The development of theta and alpha neural oscillations from ages 3 to 24 years. *Developmental Cognitive Neuroscience*, *50*, 100969. https://doi.org/10.1016/j.dcn.2021.100969

Cifelli, P., Ruffolo, G., De Felice, E., Alfano, V., van Vliet, E. A., Aronica, E., & Palma, E. (2020). Phytocannabinoids in Neurological Diseases: Could They Restore a Physiological GABAergic Transmission? *International Journal of Molecular Sciences*, *21*(3), Article 3. https://doi.org/10.3390/ijms21030723

Coben, R., Clarke, A. R., Hudspeth, W., & Barry, R. J. (2008). EEG power and coherence in autistic spectrum disorder. *Clin Neurophysiol*, *119*(5), 1002–1009. https://doi.org/10.1016/j.clinph.2008.01.013

Cochran, D. M., Sikoglu, E. M., Hodge, S. M., Edden, R. A. E., Foley, A., Kennedy, D. N., Moore, C. M., & Frazier, J. A. (2015). Relationship among Glutamine, γ-Aminobutyric Acid, and Social Cognition in Autism Spectrum Disorders. *Journal of Child and Adolescent Psychopharmacology*, *25*(4), 314–322. https://doi.org/10.1089/cap.2014.0112

Constantino, J., & Gruber, C. (2005). *The Social Responsiveness Scale Manual*. Western Psychological Services.

Constantino, J., & Gruber, C. (2012). *The Social Responsiveness Scale Manual, Second Edition (SRS-2)*. Western Psychological Services.

Cuthbert, B. N., & Insel, T. R. (2013). Toward the future of psychiatric diagnosis: The seven pillars of RDoC. *BMC Medicine*, *11*(1), 126. https://doi.org/10.1186/1741-7015-11-126

da Silva Junior, E. A., Medeiros, W. M. B., Torro, N., Sousa, J. M. M. de, Almeida, I. B. C. M.

de, Costa, F. B. da, Pontes, K. M., Nunes, E. L. G., Rosa, M. D. da, & Albuquerque, K. L.

G. D. de. (2021). Cannabis and cannabinoid use in autism spectrum disorder: A

systematic review. *Trends in Psychiatry and Psychotherapy*, *44*, e20200149.

https://doi.org/10.47626/2237-6089-2020-0149

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial

EEG dynamics including independent component analysis. *Journal of Neuroscience*

Methods, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Dickinson, A., Jones, M., & Milne, E. (2016). Measuring neural excitation and inhibition in

autism: Different approaches, different findings and different interpretations. *Brain*

Research, *1648*, 277–289. https://doi.org/10.1016/j.brainres.2016.07.011

Donaldson, P. H., Kirkovski, M., Rinehart, N. J., & Enticott, P. G. (2017). Autism-relevant

traits interact with temporoparietal junction stimulation effects on social cognition:

A high-definition transcranial direct current stimulation and electroencephalography

study. *Eur J Neurosci*, 1–13. https://doi.org/10.1111/ejn.13675

Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A.

H., Wallis, J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing neural

power spectra into periodic and aperiodic components. *Nature Neuroscience*, *23*(12),

1655–1665. https://doi.org/10.1038/s41593-020-00744-x

Donoghue, T., Schaworonkow, N., & Voytek, B. (2021). Methodological considerations for

studying neural oscillations. *European Journal of Neuroscience*, *n/a*(n/a), 1–26.

https://doi.org/10.1111/ejn.15361

Egerton, A., Grace, A. A., Stone, J., Bossong, M. G., Sand, M., & McGuire, P. (2020).

Glutamate in schizophrenia: Neurodevelopmental perspectives and drug

development. *Schizophrenia Research*, *223*, 59–70.

https://doi.org/10.1016/j.schres.2020.09.013

Fatemi, S. H. (2008). The hyperglutamatergic hypothesis of autism. *Progress in Neuro-*

Psychopharmacology & Biological Psychiatry, *32*(3), 911, author reply 912-3.

https://doi.org/10.1016/j.pnpbp.2007.11.004

Ford, T. C., & Crewther, D. P. (2016). A Comprehensive Review of the (1)H-MRS Metabolite

Spectrum in Autism Spectrum Disorder. *Frontiers in Molecular Neuroscience*, *9*(14),

1–27. https://doi.org/10.3389/fnmol.2016.00014

Ford, T. C., Nibbs, R., & Crewther, D. P. (2017a). Glutamate/GABA+ ratio is associated with

the psychosocial domain of autistic and schizotypal traits. *PLoS ONE*, *12*(7),

e0181961. https://doi.org/10.1371/journal.pone.0181961

Ford, T. C., Nibbs, R., & Crewther, D. P. (2017b). Increased glutamate/GABA+ ratio in a

shared autistic and schizotypal trait phenotype termed Social Disorganisation.

NeuroImage: Clinical, *16*, 125–131. https://doi.org/10.1016/j.nicl.2017.07.009

Ford, T. C., Woods, W., & Crewther, D. P. (2017). Mismatch field latency, but not power,

may mark a shared autistic and schizotypal trait phenotype. *International Journal of*

Psychophysiology, *116*, 60–67. https://doi.org/10.1016/j.ijpsycho.2017.02.008

Frith, C. D., & Frith, U. (2007). Social Cognition in Humans. *Current Biology*, *17*(16), R724–

R732. https://doi.org/10.1016/j.cub.2007.05.068

Frye, R. E. (2014). Clinical potential, safety, and tolerability of arbaclofen in the treatment of

autism spectrum disorder. *Drug, Healthcare and Patient Safety*, *6*, 69–76.

https://doi.org/10.2147/DHPS.S39595

Gaetz, W., Bloy, L., Wang, D. J., Port, R. G., Blaskey, L., Levy, S. E., & Roberts, T. P. (2014).

GABA estimation in the brains of children on the autism spectrum: Measurement

precision and regional cortical variation. *Neuroimage*, *86*(1), 1–9.

https://doi.org/10.1016/j.neuroimage.2013.05.068

Gao, R., & Penzes, P. (2015). Common Mechanisms of Excitatory and Inhibitory Imbalance in

Schizophrenia and Autism Spectrum Disorders. *Current Molecular Medicine*, *15*(2),

146–167. https://doi.org/10.2174/1566524015666150303003028

Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance

from field potentials. *NeuroImage*, *158*, 70–78.

https://doi.org/10.1016/j.neuroimage.2017.06.078

Gogolla, N., LeBlanc, J. J., Quast, K. B., Südhof, T. C., Fagiolini, M., & Hensch, T. K. (2009).

Common circuit defect of excitatory-inhibitory balance in mouse models of autism.

Journal of Neurodevelopmental Disorders, *1*(2), 172–181.

https://doi.org/10.1007/s11689-009-9023-x

Gonçalves, A. M., & Monteiro, P. (2023). Autism Spectrum Disorder and auditory sensory

alterations: A systematic review on the integrity of cognitive and neuronal functions

related to auditory processing. *Journal of Neural Transmission*, *130*(3), 325–408.

https://doi.org/10.1007/s00702-023-02595-9

Gonçalves, J., Violante, I. R., Sereno, J., Leitão, R. A., Cai, Y., Abrunhosa, A., Silva, A. P., Silva, A. J., & Castelo-Branco, M. (2017). Testing the excitation/inhibition imbalance hypothesis in a mouse model of the autism spectrum disorder: In vivo neurospectroscopy and molecular evidence for regional phenotypes. *Molecular Autism*, *8*(1), 47. https://doi.org/10.1186/s13229-017-0166-4

Greenwood, L. M., Leung, S., Michie, P. T., Green, A., Nathan, P. J., Fitzgerald, P., Johnston, P., Solowij, N., Kulkarni, J., & Croft, R. J. (2018). The effects of glycine on auditory mismatch negativity in schizophrenia. *Schizophr Res*, *191*, 61–69. https://doi.org/10.1016/j.schres.2017.05.031

Grent-'t-Jong, T., Gross, J., Goense, J., Wibral, M., Gajwani, R., Gumley, A. I., Lawrie, S. M., Schwannauer, M., Schultze-Lutter, F., Navarro Schröder, T., Koethe, D., Leweke, F. M., Singer, W., & Uhlhaas, P. J. (2018). Resting-state gamma-band power alterations in schizophrenia reveal E/I-balance abnormalities across illness-stages. *eLife*, *7*, e37799. https://doi.org/10.7554/eLife.37799

Gur, R. C., Sara, R., Hagendoorn, M., Marom, O., Hughett, P., Macy, L., Turner, T., Bajcsy, R., Posner, A., & Gur, R. E. (2002). A method for obtaining 3-dimensional facial expressions and its standardization for use in neurocognitive studies. *Journal of Neuroscience Methods*, *115*, 137–143. https://doi.org/10.1016/S0165-0270(02)00006-7

Hacohen, M., Stolar, O. E., Berkovitch, M., Elkana, O., Kohn, E., Hazan, A., Heyman, E., Sobol, Y., Waissengreen, D., Gal, E., & Dinstein, I. (2022). Children and adolescents with ASD treated with CBD-rich cannabis exhibit significant improvements particularly in social symptoms: An open label study. *Translational Psychiatry*, *12*(1), 375. https://doi.org/10.1038/s41398-022-02104-8

Han, S., Tai, C., Jones, C. J., Scheuer, T., & Catterall, W. A. (2014). Enhancement of inhibitory neurotransmission by GABAA receptors having alpha2,3-subunits ameliorates behavioral deficits in a mouse model of autism. *Neuron*, *81*(6), 1282–1289. https://doi.org/10.1016/j.neuron.2014.01.016

Hanada, T. (2020). Ionotropic Glutamate Receptors in Epilepsy: A Review Focusing on AMPA and NMDA Receptors. *Biomolecules*, *10*(3), 464. https://doi.org/10.3390/biom10030464

Hegarty, J. P., Weber, D. J., Cirstea, C. M., & Beversdorf, D. Q. (2018). Cerebro-Cerebellar

    Functional Connectivity is Associated with Cerebellar Excitation–Inhibition Balance in

    Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, 1–14.

    https://doi.org/doi.org/10.1007/s10803-018-3613-y

Higgins, W. C., Ross, R. M., Langdon, R., & Polito, V. (2023). The "Reading the Mind in the

    Eyes" Test Shows Poor Psychometric Properties in a Large, Demographically

    Representative U.S. Sample. *Assessment*, *30*(6), 1777–1789.

    https://doi.org/10.1177/10731911221124342

Hill, A. T., Clark, G. M., Bigelow, F. J., Lum, J. A. G., & Enticott, P. G. (2022). Periodic and

    aperiodic neural activity displays age-dependent changes across early-to-middle

    childhood. *Developmental Cognitive Neuroscience*, *54*, 101076.

    https://doi.org/10.1016/j.dcn.2022.101076

Hill, A. T., Ford, T. C., Bailey, N. W., Lum, J. A. G., Bigelow, F. J., Oberman, L. M., & Enticott, P.

    G. (2025). EEG during dynamic facial emotion processing reveals neural activity

    patterns associated with autistic traits in children. *Cerebral Cortex*, *35*(2), bhaf020.

    https://doi.org/10.1093/cercor/bhaf020

Hill, A. T., Van Der Elst, J., Bigelow, F. J., Lum, J. A. G., & Enticott, P. G. (2022). Right anterior

    theta connectivity predicts autistic social traits in typically developing children.

    Biological Psychology, *175*, 108448.

    https://doi.org/10.1016/j.biopsycho.2022.108448

Hurley, R. S. E., Losh, M., Parlier, M., Reznick, J. S., & Piven, J. (2007). The Broad Autism

    Phenotype Questionnaire. *Journal of Autism and Developmental Disorders*, *37*(9),

    1679–1690. https://doi.org/10.1007/s10803-006-0299-3

Insel, T. R. (2010). Research Domain Criteria (RDoC): Toward a New Classification

    Framework for Research on Mental Disorders. *American Journal of Psychiatry*,

    167(7), 748–751. https://doi.org/10.1176/appi.ajp.2010.09091379

Johnson, S. B., Blum, R. W., & Giedd, J. N. (2009). Adolescent Maturity and the Brain: The

    Promise and Pitfalls of Neuroscience Research in Adolescent Health Policy. *The*

    Journal of Adolescent Health : Official Publication of the Society for Adolescent

    Medicine, *45*(3), 216–221. https://doi.org/10.1016/j.jadohealth.2009.05.016

Jung, M., Tu, Y., Lang, C. A., Ortiz, A., Park, J., Jorgenson, K., Kong, X.-J., & Kong, J. (2019).

    Decreased structural connectivity and resting-state brain activity in the lateral

occipital cortex is associated with social communication deficits in boys with autism spectrum disorder. *NeuroImage*, *190*, 205–212. https://doi.org/10.1016/j.neuroimage.2017.09.031

Kirkovski, M., Enticott, P. G., Hughes, M. E., Rossell, S. L., & Fitzgerald, P. B. (2016). Atypical Neural Activity in Males But Not Females with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, *46*(3), 954–963. https://doi.org/10.1007/s10803-015-2639-7

Kittel, A. F. D., Olderbak, S., & Wilhelm, O. (2022). Sty in the Mind's Eye: A Meta-Analytic Investigation of the Nomological Network and Internal Consistency of the "Reading the Mind in the Eyes" Test. *Assessment*, *29*(5), 872–895. https://doi.org/10.1177/1073191121996469

Korkmaz, S., Goksuluk, D., & Zararsiz, G. (2014). MVN: An R Package for Assessing Multivariate Normality. *The R Journal*, *6*(2), 151–162.

Lenroot, R. K., Gogtay, N., Greenstein, D. K., Wells, E. M., Wallace, G. L., Clasen, L. S., Blumenthal, J. D., Lerch, J., Zijdenbos, A. P., Evans, A. C., Thompson, P. M., & Giedd, J. N. (2007). Sexual dimorphism of brain developmental trajectories during childhood and adolescence. *NeuroImage*, *36*(4), 1065–1073. https://doi.org/10.1016/j.neuroimage.2007.03.053

Lindell, A. K. (2006). In Your Right Mind: Right Hemisphere Contributions to Language Processing and Production. *Neuropsychology Review*, *16*, 131–148. https://doi.org/doi.org/10.1007/s11065-006-9011-9

Lisman, J. (2012). Excitation, inhibition, local oscillations, or large-scale loops: What causes the symptoms of schizophrenia? *Curr Opin Neurobiol*, *22*(3), 537–544. https://doi.org/10.1016/j.conb.2011.10.018

Manning, J. R., Jacobs, J., Fried, I., & Kahana, M. J. (2009). Broadband shifts in local field potential power spectra are correlated with single-neuron spiking in humans. *Journal of Neuroscience*, *29*(43), 13613–13620. Scopus. https://doi.org/10.1523/JNEUROSCI.2041-09.2009

Manyukhina, V. O., Prokofyev, A. O., Galuta, I. A., Goiaeva, D. E., Obukhova, T. S., Schneiderman, J. F., Altukhov, D. I., Stroganova, T. A., & Orekhova, E. V. (2022). Globally elevated excitation–inhibition ratio in children with autism spectrum

disorder and below-average intelligence. *Molecular Autism*, *13*(1), 20.

https://doi.org/10.1186/s13229-022-00498-2

Marsman, A., van den Heuvel, M. P., Klomp, D. W. J., Kahn, R. S., Luijten, P. R., & Hulshoff

Pol, H. E. (2013). Glutamate in Schizophrenia: A Focused Review and Meta-Analysis

of 1H-MRS Studies. *Schizophrenia Bulletin*, *39*(1), 120–129.

https://doi.org/10.1093/schbul/sbr069

McSweeney, M., Morales, S., Valadez, E. A., Buzzell, G. A., & Fox, N. A. (2021). Longitudinal

age- and sex-related change in background aperiodic activity during early

adolescence. *Developmental Cognitive Neuroscience*, *52*, 101035.

https://doi.org/10.1016/j.dcn.2021.101035

Miller, K. J., Honey, C. J., Hermes, D., Rao, R. P., denNijs, M., & Ojemann, J. G. (2014).

Broadband changes in the cortical surface potential track activation of functionally

diverse neuronal populations. *NeuroImage*, *85*, 711–720.

https://doi.org/10.1016/j.neuroimage.2013.08.070

Milovanovic, M., & Grujicic, R. (2021). Electroencephalography in Assessment of Autism

Spectrum Disorders: A Review. *Frontiers in Psychiatry*, *12*.

https://www.frontiersin.org/articles/10.3389/fpsyt.2021.686021

Moessnang, C., Schäfer, A., Bilek, E., Roux, P., Otto, K., Baumeister, S., Hohmann, S.,

Poustka, L., Brandeis, D., Banaschewski, T., Meyer-Lindenberg, A., & Tost, H. (2016).

Specificity, reliability and sensitivity of social brain responses during spontaneous

mentalizing. *Social Cognitive and Affective Neuroscience*, *11*(11), 1687–1697.

https://doi.org/10.1093/scan/nsw098

Molina, J. L., Voytek, B., Thomas, M. L., Joshi, Y. B., Bhakta, S. G., Talledo, J. A., Swerdlow, N.

R., & Light, G. A. (2020). Memantine Effects on Electroencephalographic Measures of

Putative Excitatory/Inhibitory Balance in Schizophrenia. *Biological Psychiatry:*

Cognitive Neuroscience and Neuroimaging, *5*(6), 562–568.

https://doi.org/10.1016/j.bpsc.2020.02.004

Nelson, S. B., & Valakh, V. (2015). Excitatory/Inhibitory Balance and Circuit Homeostasis in

Autism Spectrum Disorders. *Neuron*, *87*(4), 684–698.

https://doi.org/10.1016/j.neuron.2015.07.033

Ostlund, B., Alperin, B. R., Drew, T., & Karalunas, S. L. (2021). Behavioral and cognitive

correlates of the aperiodic (1/f-like) exponent of the EEG power spectrum in

adolescents with and without ADHD. *Developmental Cognitive Neuroscience*, *48*, 100931. https://doi.org/10.1016/j.dcn.2021.100931

Ostlund, B., Donoghue, T., Anaya, B., Gunther, K. E., Karalunas, S. L., Voytek, B., & Pérez-Edgar, K. E. (2022). Spectral parameterization for studying neurodevelopment: How and why. *Developmental Cognitive Neuroscience*, *54*, 101073. https://doi.org/10.1016/j.dcn.2022.101073

Pani, S. M., Saba, L., & Fraschini, M. (2022). Clinical applications of EEG power spectra aperiodic component analysis: A mini-review. *Clinical Neurophysiology*, *143*, 1–13. https://doi.org/10.1016/j.clinph.2022.08.010

Parrella, N.-F., Hill, A. T., Dipnall, L. M., Loke, Y. J., Enticott, P. G., & Ford, T. C. (2024). Inhibitory dysfunction and social processing difficulties in autism: A comprehensive narrative review. *Journal of Psychiatric Research*, *169*, 113–125. https://doi.org/10.1016/j.jpsychires.2023.11.014

Parrella, N.-F., Hill, A. T., Enticott, P. G., Barhoun, P., Bower, I. S., & Ford, T. C. (2023). A systematic review of cannabidiol trials in neurodevelopmental disorders. *Pharmacology Biochemistry and Behavior*, *230*, 173607. https://doi.org/10.1016/j.pbb.2023.173607

Pinkham, A. E., Penn, D. L., Green, M. F., & Harvey, P. D. (2016). Social Cognition Psychometric Evaluation: Results of the Initial Psychometric Study. *Schizophrenia Bulletin*, *42*(2), 494–504. https://doi.org/10.1093/schbul/sbv056

Pion-Tonachini, L., Kreutz-Delgado, K., & Makeig, S. (2019). ICLabel: An automated electroencephalographic independent component classifier, dataset, and website. *NeuroImage*, *198*, 181–197. https://doi.org/10.1016/j.neuroimage.2019.05.026

Pretzsch, C. M., Floris, D. L., Voinescu, B., Elsahib, M., Mendez, M. A., Wichers, R., Ajram, L., Ivin, G., Heasman, M., Pretzsch, E., Williams, S., Murphy, D. G. M., Daly, E., & McAlonan, G. M. (2021). Modulation of striatal functional connectivity differences in adults with and without autism spectrum disorder in a single-dose randomized trial of cannabidivarin. *Molecular Autism*, *12*(1), 49. https://doi.org/10.1186/s13229-021-00454-6

Pretzsch, C. M., Voinescu, B., Lythgoe, D., Horder, J., Mendez, M. A., Wichers, R., Ajram, L., Ivin, G., Heasman, M., Edden, R. A. E., Williams, S., Murphy, D. G. M., Daly, E., & McAlonan, G. M. (2019). Effects of cannabidivarin (CBDV) on brain excitation and

inhibition systems in adults with and without Autism Spectrum Disorder (ASD): A single dose trial during magnetic resonance spectroscopy. *Translational Psychiatry*, *9*(1), 313. https://doi.org/10.1038/s41398-019-0654-8

R Core Team. (2023). *R: A Language and Environment for Statistical Computing* [Computer software]. R Foundation for Statistical Computing. https://www.R-project.org/

Robertson, M. M., Furlong, S., Voytek, B., Donoghue, T., Boettiger, C. A., & Sheridan, M. A. (2019). EEG power spectral slope differs by ADHD status and stimulant medication exposure in early childhood. *Journal of Neurophysiology*, *122*(6), 2427–2437. https://doi.org/10.1152/jn.00388.2019

Rowland, L. M., Summerfelt, A., Wijtenburg, S. A., Du, X., Chiappelli, J. J., Krishna, N., West, J., Muellerklein, F., Kochunov, P., & Hong, L. E. (2016). Frontal Glutamate and gamma-Aminobutyric Acid Levels and Their Associations With Mismatch Negativity and Digit Sequencing Task Performance in Schizophrenia. *JAMA Psychiatry*, *73*(2), 166–174. https://doi.org/10.1001/jamapsychiatry.2015.2680

Rubenstein, J. L. R., & Merzenich, M. M. (2003). Model of autism: Increased ratio of excitation/inhibition in key neural systems: Model of autism. *Genes, Brain and Behavior*, *2*(5), 255–267. https://doi.org/10.1034/j.1601-183X.2003.00037.x

Scrivener, C. L., & Reader, A. T. (2022). Variability of EEG electrode positions and their underlying brain regions: Visualizing gel artifacts from a simultaneous EEG-fMRI dataset. *Brain and Behavior*, *12*(2), e2476. https://doi.org/10.1002/brb3.2476

Sehatpour, P., Javitt, D. C., De Baun, H. M., Carlson, M., Beloborodova, A., Margolin, D. H., Carlton, M. B. L., Brice, N. L., & Kantrowitz, J. T. (2022). Mismatch negativity as an index of target engagement for excitation/inhibition-based treatment development: A double-blind, placebo-controlled, randomized, single-dose cross-over study of the serotonin type-3 receptor antagonist CVN058. *Neuropsychopharmacology*, *47*(3), 711–718. https://doi.org/10.1038/s41386-021-01170-8

Shuffrey, L. C., Pini, N., Potter, M., Springer, P., Lucchini, M., Rayport, Y., Sania, A., Firestein, M., Brink, L., Isler, J. R., Odendaal, H., & Fifer, W. P. (2022). Aperiodic electrophysiological activity in preterm infants is linked to subsequent autism risk. *Developmental Psychobiology*, *64*(4), e22271. https://doi.org/10.1002/dev.22271

Sohal, V. S., & Rubenstein, J. L. R. (2019). Excitation-inhibition balance as a framework for investigating mechanisms in neuropsychiatric disorders. *Molecular Psychiatry*. https://doi.org/10.1038/s41380-019-0426-0

Stanyard, R. A., Mason, D., Ellis, C., Dickson, H., Short, R., Batalle, D., & Arichi, T. (2024). *Aperiodic and Hurst EEG exponents across early human brain development: A systematic review* [Preprint]. Neuroscience. https://doi.org/10.1101/2024.02.02.578622

Stoppel, L. J., Kazdoba, T. M., Schaffler, M. D., Preza, A. R., Heynen, A., Crawley, J. N., & Bear, M. F. (2018). R-Baclofen Reverses Cognitive Deficits and Improves Social Interactions in Two Lines of 16p11.2 Deletion Mice. *Neuropsychopharmacology*, *43*(3), 513–524. https://doi.org/10.1038/npp.2017.236

Tan, T., Wang, W., Xu, H., Huang, Z., Wang, Y. T., & Dong, Z. (2018). Low-Frequency rTMS Ameliorates Autistic-Like Behaviors in Rats Induced by Neonatal Isolation Through Regulating the Synaptic GABA Transmission. *Frontiers in Cellular Neuroscience*, *12*. https://doi.org/10.3389/fncel.2018.00046

Tebartz van Elst, L., Maier, S., Fangmeier, T., Endres, D., Mueller, G. T., Nickel, K., Ebert, D., Lange, T., Hennig, J., Biscaldi, M., Riedel, A., & Perlov, E. (2014). Disturbed cingulate glutamate metabolism in adults with high-functioning autism spectrum disorder: Evidence in support of the excitatory/inhibitory imbalance hypothesis. *Mol Psychiatry*. https://doi.org/10.1038/mp.2014.62

Thuwal, K., Banerjee, A., & Roy, D. (2021). Aperiodic and Periodic Components of Ongoing Oscillatory Brain Dynamics Link Distinct Functional Aspects of Cognition across Adult Lifespan. *Eneuro*, *8*(5), ENEURO.0224-21.2021. https://doi.org/10.1523/ENEURO.0224-21.2021

Tremblay, R., Lee, S., & Rudy, B. (2016). GABAergic interneurons in the neocortex: From cellular properties to circuits. *Neuron*, *91*(2), 260–292. https://doi.org/10.1016/j.neuron.2016.06.033

Uhlhaas, P. J., & Singer, W. (2010). Abnormal neural oscillations and synchrony in schizophrenia. *Nature Reviews. Neuroscience*, *11*(2), 100–113. https://doi.org/10.1038/nrn2774

Vandierendonck, A. (2017). A comparison of methods to combine speed and accuracy measures of performance: A rejoinder on the binning procedure. *Behavior Research Methods*, *49*(2), 653–673. https://doi.org/10.3758/s13428-016-0721-5

von dem Hagen, E. A. H., Nummenmaa, L., Yu, R., Engell, A. D., Ewbank, M. P., & Calder, A. J. (2011). Autism Spectrum Traits in the Typical Population Predict Structure and Function in the Posterior Superior Temporal Sulcus. *Cerebral Cortex*, *21*(3), 493–500. https://doi.org/10.1093/cercor/bhq062

Waschke, L., Donoghue, T., Fiedler, L., Smith, S., Garrett, D. D., Voytek, B., & Obleser, J. (2021). Modality-specific tracking of attention and sensory statistics in the human electrophysiological spectral exponent. *eLife*, *10*, e70068. https://doi.org/10.7554/eLife.70068

Wechsler, D. (2011). *WASI-II: Wechsler abbreviated scale of intelligence*. PsychCorp.

Wilkinson, C. L., & Nelson, C. A. (2021). Increased aperiodic gamma power in young boys with Fragile X Syndrome is associated with better language ability. *Molecular Autism*, *12*(1), 17. https://doi.org/10.1186/s13229-021-00425-x

Yeo, I.-K., & Johnson, R. A. (2000). A New Family of Power Transformations to Improve Normality or Symmetry. *Biometrika*, *87*(4), 954–959.

Yizhar, O., Fenno, L. E., Prigge, M., Schneider, F., Davidson, T. J., O'Shea, D. J., Sohal, V. S., Goshen, I., Finkelstein, J., Paz, J. T., Stehfest, K., Fudim, R., Ramakrishnan, C., Huguenard, J. R., Hegemann, P., & Deisseroth, K. (2011). Neocortical excitation/inhibition balance in information processing and social dysfunction. *Nature*, *477*(7363), 171–178. https://doi.org/10.1038/nature10360

Zhang, C., Stock, A.-K., Mückschel, M., Hommel, B., & Beste, C. (2023). Aperiodic neural activity reflects metacontrol. *Cerebral Cortex*, *33*(12), 7941–7951. https://doi.org/10.1093/cercor/bhad089

Zhou, S., & Yu, Y. (2018). Synaptic E-I Balance Underlies Efficient Neural Coding. *Frontiers in Neuroscience*, *12*. https://www.frontiersin.org/articles/10.3389/fnins.2018.00046