A hybrid machine learning framework for functional annotation applied to mitochondrial glutathione metabolism and transport in cancers.

Luke Kennedy[1,2], Jagdeep K Sandhu[1,3], Mary-Ellen Harper[1,2] *†, and Miroslava Cuperlovic-Culf[1,4] *†

*Corresponding authors

†Co-senior authors

Affiliations:

[1]Department of Biochemistry, Microbiology and Immunology, Faculty of Medicine University of Ottawa, 451 Smyth Road, Ottawa, ON K1H 8M5, Canada.

[2]Ottawa Institute of Systems Biology, University of Ottawa, 451 Smyth Road, Ottawa, ON K1H 8M5, Canada.

[3]Human Health Therapeutics Research Centre, Bldg M54, National Research Council Canada, 1200 Montreal Road, Ottawa, ON K1A 0R6, Canada.

[4]Digital Technologies Research Centre, Bldg M50, National Research Council Canada, 1200 Montreal Road, Ottawa, ON K1A 0R6, Canada.

Correspondence:

Mary-Ellen Harper **mharper@uottawa.ca** or Miroslava Cuperlovic-Culf **miroslava.cuperlovic-culf@nrc-cnrc.gc.ca**

Abstract

Background

Alterations of metabolism, including changes in mitochondrial and glutathione (GSH) metabolism, are a well appreciated hallmark of many cancers. Mitochondrial GSH (mGSH) transport is a poorly characterized aspect of GSH metabolism, which we investigate in the context of cancer. Existing functional annotation approaches from machine (ML) or deep learning (DL) models based only on protein sequences are unable to annotate functions in biological contexts, meaning new approaches must be developed for this task.

Results

We develop a flexible ML framework for functional annotation from diverse feature data. This hybrid approach leverages cancer cell line multi-omics data and other biological knowledge data as features, to uncover potential genes involved in mGSH metabolism and membrane transport in cancers. This framework achieves an average AUROC across functional annotation tasks of 0.900 and can be effectively applied to annotate a range of biological functions. For our application, classification models predict the known mGSH transporter SLC25A39 but not SLC25A40 as being highly probably related to GSH metabolism in cancers. SLC25A24 and the orphan SLC25A43 are also predicted to be associated with mGSH metabolism by this approach and structural analysis of these proteins reveal similarities in potential substrate binding regions to the binding residues of SLC25A39.

Conclusion

These findings have implications for a better understanding of cancer cell metabolism and novel therapeutic targets with respect to GSH metabolism through potential novel functional annotations of genes. The hybrid ML framework proposed here can be applied to other biological function classifications or multi-omics datasets to generate hypotheses in various biological contexts. Code and a tutorial for generating models

52   and predictions in this framework are available at:

53   https://github.com/lkenn012/mGSH_cancerClassifiers.

54   **Keywords:** Mitochondria, glutathione, machine learning, cancer, protein function

55   annotation, transmembrane transport, SLC25, multi-omics, knowledge-based, gene

56   ontology

Background

Glutathione (GSH) is a highly abundant tripeptide antioxidant within cells, crucial for many biological processes with its major role in regulation of reactive oxygen species (ROS). Metabolic changes are one of the hallmarks of cancer[1], with well documented alterations in GSH metabolism foremost among them. These alterations of metabolism appear to benefit cancer cells, aiding in tumor proliferation and survival, through not yet fully understood mechanisms and interactions[2]. Alterations of GSH metabolism are essential for tumor proliferation in several cancers[3,4] either directly where they mitigate perturbations in redox homeostasis[5,6,7], or indirectly through ferroptosis[8] and metabolism of chemotherapeutics[8,9].

Central to the contributions of GSH in the metabolic characteristics of cancer cells is mitochondrial GSH (mGSH), which is found at millimolar concentrations in the organelle, at similar levels to those in the cytoplasm. Mitochondria are the primary source of ROS, largely as a byproduct of the oxidative phosphorylation system (OXPHOS), and alterations in redox balance and ROS signaling are central to the role of mitochondria in cancer cell proliferation[10]. Many GSH-utilizing enzymes with altered expression patterns in cancers are found in mitochondria, most notably glutathione peroxidase 4 (GPX4) which is a major regulator of the ferroptosis pathway. Also important are mitochondrial glutathione enzymes such as glutathione S-transferase, which conjugates and detoxifies xenobiotics and the peroxiredoxins, which lower ROS via GSH and have also been shown to promote cancer cell survival[11].

The processes involved in the uptake of GSH into mitochondria are poorly understood. While high concentrations of GSH are present in both mitochondria and the cytosol, the synthetic enzymes are exclusively within the cytosol[12]. SLC25A10 and SLC25A11, also known as the mitochondrial dicarboxylate and oxoglutarate carriers, respectively, were initially proposed as the proteins responsible for GSH transport into mitochondria over 25 years ago[13]. However, detailed functional experiments in lipid vesicle systems in 2014 convincingly showed that no GSH transport was mediated by these two proteins[14].

Recent evidence from two groups demonstrated mGSH transport roles for SLC25A39

and SLC25A40[15,16]. Specifically, Wang *et al.*[15] identified the transporters through

quantitative proteomics of mitochondria from GSH-depleted HeLa cells. Subsequently,

Shi *et al.*[16] leveraged CRISPR screening of gene and environment interactions to

demonstrate buffering interactions between SLC25A39 and the mitochondrial iron

transporter SLC25A37, revealing SLC25A39 as a candidate. The roles of these

transporters in neurodegenerative diseases and cancer have recently been explored[17].

However, it remains unclear if the remaining GSH import in A39 knockouts observed in

these studies is facilitated by SLC25A40, which is expressed at roughly one tenth the

level of SLC25A39, or if other transport mechanisms exist, such as those for other GSH

species[18,19]. Interactions affecting mGSH metabolism and transport, having secondary

effects on transport may also be relevant and are even less well understood.

With recent successes of AlphaFold[20] and RosettaFold[21] in the elucidation of protein

structure, there is a promising future for computational biology in the related problem of

protein function prediction. However, competitions such as Critical Assessment of

Functional Annotation[22] have not yet found a solution for *de novo* function prediction

from sequence. General function annotation models like DeepGO and DeepGOPlus[23,24]

represent major advances for the field. Along with sequence-based methods, models

that leverage non-sequence features for function annotation exist[25,26] as alternative

approaches to sequence-based approaches. In addition to the general function

prediction models, there are many sequence-based models designed for specific

protein feature annotation such as protein-protein interactions[27] and antibody design[28].

Omics-based methods for functional annotation are limited. Recent work by Kunc and

Kléma[29] utilize characteristics in co-expression networks constructed from genes to

predict shared KEGG pathways between genes. Similarly, Wekesa *et al.*[30] combine

differential gene expression information and knowledge of protein-protein interactions

through a neighbor-voting algorithm for prediction of shared functions between yeast

proteins. Finally, Wang *et al.*[31] predict gene-gene interactions by combining co-

expression features with prior biological knowledge features (e.g., subcellular

localization, homology, Reactome similarity).

The approaches of Wekesa *et al.* and Wang *et al.* can be classified as hybrid ML

approaches, which aim to capture "the best of both worlds" by combining the predictive

powers of data-driven approaches of ML with the interpretability of theory-based models

such as mechanistic or knowledge-based models. These types of models have been

applied to several domains, most notably in physics-informed ML models[32], and with

respect to computational biology, hybrid modeling has been applied through diverse

frameworks. Non-ML approaches have been used to uncover biological phenomena,

which would otherwise be missed; a recent example integrates behavioral,

transcriptomic, and network modeling to reveal the role of brain mitochondria and

organization in mouse behavior[33]. Alternatively, ML models have been integrated with

mechanistic models of metabolism to determine kinetic parameters and predict

downstream metabolic effects[34]. P-net uses a neural network architecture based on

biological knowledge of hierarchical gene-pathway-process interactions to predict

prostate cancer discovery from gene features, such as methylation and copy number[35].

On the other hand, AlphaFold[20] incorporates evolutionary information through multiple

sequence alignments and physical structural constraints to inform its protein structure

predictions. These hybrid models retain the predictive power of data-driven methods,

often performing better than, or comparable to, standard data-driven models, while

increasing the interpretability of predictions that are in-part based on biological

knowledge.

With this, we sought to develop a ML framework that leverages multi-omics data from

human cancer cells and existing knowledge to identify potential genes involved in

mGSH transport and potential interacting metabolic processes. Several ML classifier

models were developed that utilize cancer cell line encyclopedia (CCLE)[36]

transcriptomics features to predict gene ontology (GO) annotations for genes of relevant

GO terms. Specifically, three classifiers were developed for annotating to: glutathione

metabolic process (GO:0006749), mitochondrion (GO:0005739), and transmembrane

transporter activity (GO:0022857). Additionally, we developed hybrid models for this

task which include prior knowledge as features through MitoCarta[37] scores for

mitochondrial localization classification, TrSSP[38] scores for transporter classification, or

CCLE GSH & glutathione disulfide (GSSG) metabolomics data for glutathione

classification. We find that Random Forest (RF) classifiers perform the best from our
models tested, with hybrid models outperforming strictly transcriptomics models.

Methods

All pre-processing, models and other computational work were conducted using in-
house code written in Python (Python Software Foundation. Python Language
Reference, version 3.8. Available at www.python.org). All algorithms and methods used
in ML model building and training were implemented via Python's scikit-learn package[39]
unless stated otherwise. Figures for evaluations of classifier models were produced via
the matplotlib and seaborn libraries[40,41]; GO enrichment plots were produced using
ShinyGO 0.77[42]; and all other images were created with BioRender.com.

Data collection & pre-processing

Metabolomic & transcriptomic CCLE data

The Broad Institute's Cancer Cell Line Encyclopedia (CCLE) provides multi-omics data
across over 1,000 human cancer cell lines (CCLs). CCLE transcriptomics, and
metabolomics data were used in building out models[36,43]. Raw transcriptomics and
metabolomics data were downloaded from broadinstitute.org (available at the time of
writing via depmap.org/portal/download/).

Data were cleaned and imputed by removing all CCLs and transcripts/metabolites with
>30% missing values or standard deviation values of 0 across samples. Remaining
missing values were imputed using the k-nearest neighbors (k-NN) imputation algorithm
(using Euclidean distance with k=5 selected to minimize impact of the data structure on
the missing data imputation[44]). For metabolomics features, Spearman correlations were
generated across common CCLs between GSH, GSSG metabolomics data and
transcripts in CCLE transcriptomics data. Non-significant correlations (p-value ≥ 0.05,
Student's t-test with 2 degrees of freedom) were set to 0. Spearman rho values were
normalized by Fisher transformation[45].

Feature space reduction while selecting major variances in CCLE transcriptomics was

performed using principal component analysis (PCA). These principal components

(PCs) were used as features in classifier models (further described in Model

development and feature selection process).

Knowledge-based features

Mitochondrial localization scores for genes were based on MitoCarta 3.0[37]

(broadinstitute.org/mitocarta). MitoCarta scores are determined through a naive-Bayes

model which combines features from several independent domains from features used

in our analysis, such as homology, sequence domain, and tandem mass spectrometry

of purified mitochondria. CCLE genes were assigned MitoCarta scores according to

these data, with 1 indicating mitochondrial localization, and 0 indicating non-

mitochondrial localization. For genes that are not assigned scores in MitoCarta, a value

of -1 is assigned.

Transporter activity was based on TrSSP[38] values, obtained from Zhao lab webpage

(www.zhaolab.org/TrSSP). TrSSP utilizes support vector machine models to predict

membrane transport proteins and their substrate classes using primary protein

sequence features along with position-specific scoring matrices to predict transport

function and specificity. Like MitoCarta scores, TrSSP scores were incorporated into the

model as classifications for positive, negative, or no prediction by TrSSP (1, 0, -1,

respectively).

For use as features in our classifiers, these knowledge-based categorical scores were

transformed to Boolean features using one-hot encoding.

mGSH transporter classifier models

Training and test gene sets selection

197 For each classifier, training and test genes were selected based on existing knowledge

198 of gene functions. Gene ontology[46] terms were used to determine genes related to

199 mGSH metabolism: glutathione metabolic process GO:0006749 (65 genes),

mitochondrion GO:0005739 (1685 genes), transmembrane transporter activity GO:0022857 (1154 genes). These terms cover all classes of ontology terms: biological process, cellular component, and molecular function, respectively. Annotated genes for each term were identified via AmiGO 2[47] and those genes annotated based on only low confidence computational or inferred evidence were removed. This left 40 genes for GO:0006749, 781 genes for GO:0005739, and 700 genes for GO:0022857 annotated based on experimental evidence to create the final list of positive class genes for the classifier models. We found that there is little overlap among these gene sets; primarily, common genes are found between mitochondrial and transmembrane transporter activity genes (72 genes), with 5 genes overlapping the mitochondrial and glutathione metabolic process sets and only one gene found in both transmembrane transporter activity and glutathione metabolic process genes. There are no genes annotated by all three GO terms. AmiGO annotations based on sequence similarities were retained along with the experimental AmiGO annotations used for the other GO annotations to generate the 40 genes used in the final positive gene set for glutathione metabolic process term. These annotations were retained due to the small number of genes annotated by experimental evidence for the GO term, thus creating a larger positive gene set for model training. bioDBnet[48] was used to link gene symbols from AmiGO to Ensembl gene IDs found in the CCLE data.

The negative classification set of genes was generated by randomly undersampling from genes that were not included in the positive set based on GO annotations. By undersampling, positive and negative genes (samples) for model training and testing were balanced. However, training a model through this process is biased by the specific set of randomly selected negative class genes in the training set. To ensure robustness of the models, 100 iterations of random sampling were performed to create different sets of samples for model training. Final predictions and evaluations are average results over training and testing iterations on the 100 different data sets.

Model development and feature selection process

To determine the best methodology for our task, several common ML algorithms and

feature sets were tested. Random Forest (RF), Decision Tree (DT), Naive Bayes (NB),

and Support Vector Machine (SVM) were applied for classification tasks of

mitochondrial, transmembrane transporter activity, and GSH metabolic process GO

terms. Models were tested with several sets of features that combine experimental

features from CCLE data with knowledge-based features, or feature sets containing

only experimental features. Experimental feature sets include the first 5, 14, 30, or 50

CCLE transcriptomics PCs which capture 93%, 95%, 97%, or 98% of the variance,

respectively. For classifying the GSH metabolic process GO term, Spearman

correlations values between gene transcriptomics and GSH or GSSG metabolomics

were included as features as well. Features from existing biological knowledge were

included through MitoCarta scores for mitochondrial localization in the mitochondrial

classification task, and TrSSP prediction scores for membrane transport proteins in the

transporter classification task.

For comparisons of feature sets in models, either metabolomics or knowledge features

were replaced with additional PC features to create the transcriptomics-only models

with an equivalent number of features (i.e., a mitochondrial classifier with the first 5 PCs

and MitoCarta scores as features is comparable to a transcriptomics-only classifier

using the first 6 PCs as features).

Model evaluation

Model performance was analyzed from predicted class probabilities using common

evaluation metrics including receiver operating characteristic (ROC) and precision-recall

curves (PRC). These methods provide a balanced approach to evaluating classifier

performance, which considers the effects of true positives (TPs), true negative (TNs),

false positives (FPs), and false negatives (FNs). Models were evaluated and selected

based on values for the areas under ROC (AUROC) and PRC (AUPRC), and Matthew's

correlation coefficient (MCC)[49,50]. These evaluation metrics were calculated across all

test-set predictions from five-fold cross-validations of randomly sampled training/test

sets. Reported metrics are calculated as average values across 100 iterations of

training/testing using the different gene sets.

Protein structural analysis

SLC25 structures

Protein structures are obtained from the AlphaFold[20] structure database

(alphafold.ebi.ac.uk) for use in structural analyses. Additionally, homology models from

SWISS-MODEL[51] for both SLC25A39 and A40 are included in our analyses. Homology

based structures are modeled on the crystal structure of the bovine ADP/ATP carrier

(PDB:OKC1)[52] and cryo-electron microscopy determined structure of human UCP1

(PDB:8HBV)[53] for A39 and A40, respectively.

Protein structure alignments

Comparisons of 3D protein structures were conducted using sequence-independent

alignment through the TM-align algorithm[54] and quantified by the corresponding TM-

scores, which quantifies similarities through similarities in protein structure topology.

This method performs comparably to other common alignment methods and does not

consider sequence similarity for alignment, which could skew the alignments due to

conserved sequences within the SLC25 family. Alignments were implemented using the

TMalign module for PyMOL[55] and predicted mGSH transporter structures were

compared to known mGSH transporters (SLC25A39, A40).

The CAVER PyMOL plugin version 3.0.3[56] was used to identify tunnel residues for each

SLC25 structure. Tunnels were identified by initializing the tunneling at the base or

innermost point of the transporters tunnel with maximum starting point distance and

desired radius of 3Å for starting point optimization. From this base, the tunneling

algorithm probed outward to generate an interior tunnel channel. CAVER default

parameters were specified for the tunneling algorithm, using 12 approximating balls with

minimum probe radius of 1.5Å, shell depth of 20Å, shell radius of 7Å (or 10Å for

structures where a smaller radius fails to fill the tunnel space), and a clustering

threshold of 3.5. Default parameters were used for computation memory and speed,

and were adjusted as described, if necessary, for finding transporter tunnels. Relevant

tunnels were identified from the top-ranked tunnels identified by CAVER as those

tunnels which traversed and filled the length of the proteins' tunnel without escaping to

the protein surface through a side gap or by "overflowing" the interior tunnel.

In the same fashion as the full structure alignments, analysis of the tunnel regions for

the SLC25 family were performed by aligning the tunnel-interacting residues identified

from CAVER using TM-align.

Results

Classifier models for mGSH transporters in cancer

To identify top mGSH transporter candidates, three classifier models were developed to

classify genes based on our desired candidate characteristics: mitochondrial

localization, association with GSH metabolism, and transmembrane transport function

(see **Fig.** 1.a. for overview and **Methods** for details). Each model uses features from

CCLE transcriptomics data along with one or more features from other sources; GSH

and GSSG metabolomics correlations for GSH metabolic process classification,

MitoCarta scores for mitochondrial localization classification, and TrSSP scores for

transmembrane transporter activity classification (**Fig.** 1. b.).

303
Figure 1: Overview of mGSH transporter classifier model design and evaluation. (**a**) Schematic of the

workflow for ML model development (inspired by designs by Fellows Yates *et al.*[85]). In this diagram, lines

represent the flow of information from the initially collected raw data through several transformation steps

to the processed model features, and then through the ML training, evaluation, and selection steps to

obtain the final classification of our unknown genes. (**b**) Example feature vectors assigned to genes in this

framework, where transcriptomic features are PCs and model-specific features are MitoCarta scores, for

example. ROC curves are presented for mitochondrial (**c**), transporter (**d**) and glutathione (**e**) classifiers

with various feature sets. Models which incorporate transcriptomics and other feature data are colored

while transcriptomics-only models are gray. Differently shaded curves represent models with various

numbers of transcriptomics features (range: 5-50).

Our classifiers performed well in the tasks of mitochondrial, GSH, and transporter

annotation classification across our evaluation metrics, as shown in **Table 1**. Across all

models and feature selections, mean five-fold cross-validations shows AUROC is 0.820,

and RF classifiers being the best performer with mean AUROC values of 0.900.

Classifier performance varied across our three classifier types, with mitochondrial

classifiers performing the best and glutathione classifiers performing the worst, on

average, which may be attributed to two factors. First, the set of annotated genes

related to GSH metabolism is much smaller than the gene sets for the other classifiers

(~40 genes vs. ~700 genes), thus providing a smaller training set for these models.

Additionally, the GSH classifiers rely solely on CCLE multi-omics data unlike the other

two classifiers which also include data from existing knowledge-sources.

Algorithm	Metric		
	AUROC	AUPRC	MCC
Decision tree	0.7663	0.8270	0.5371
	$0.6896^{\dagger}$	$0.7692^{\dagger}$	$0.3840^{\dagger}$
Naive-Bayes	0.8553	0.8422	0.3517*
	$0.7433^{\dagger}$	$0.7334^{\dagger}$	$0.2028^{\dagger}$ *
Random forest	**0.9003**	**0.8941**	**0.6349**
	$\mathbf{0.8380^{\dagger}}$	$\mathbf{0.8156^{\dagger}}$	$\mathbf{0.5437^{\dagger}}$
Support vector machine	0.7614	0.7558	0.3487
	$0.7130^{\dagger}$	$0.7250^{\dagger}$	$0.3125^{\dagger}$

† scores from models with transcriptomics-only features.
* low MCC values relative to other evaluation metrics due to low positive class
accuracy.

Table 1. **Average performance of classifiers in mGSH transporter classification**. Scores are mean

values across each classifier type and all feature sets which include transcriptomic and other data.

Individual scores are calculated as averages across bootstrap iterations. The highest scoring model for

each evaluation metric is bolded.

Along with testing several ML algorithms in our classifiers, multiple feature sets were

explored to determine the best selection for our task. Specifically, variations of the

number of top PCs as linear combinations of transcriptomic features (described in

Methods) from 5-50, were tested. Increasing the number of these features in our models

did not notably improve performance by our metrics (**Supp. Fig.** 1). Based on these

results, a small number of the first several PCs are sufficient for effective classification.

However, specific gene classification probabilities are variable across models with

different numbers of PC features used. Since performance is similar across these

models, rather than selecting a specific model, we used mean gene classification

probabilities with standard error (SE) calculated from models with different numbers of

PC features (5, 14, 30, 50) for final classification results.

Comparison of hybrid models and transcriptomics-based models

Consistently across all tested models, classifiers that incorporate both experimental

data from CCLE transcriptomics data and data from other knowledge bases as features

outperform those models with an equivalent number of experimental data features alone

(**Fig**. 1.c.,d., **Table 1**). Although no knowledge component exists for the GSH classifiers,

transcriptomics-based models were compared to models which incorporate GSH and

GSSG metabolomics data from the CCLE along with the CCLE transcriptomics data for

features. Interestingly, the additional data source does not seem to similarly improve

performance in the GSH models, with performance of transcriptomics-only GSH

classifiers comparable to GSH classifiers using both transcriptomic and metabolomic

features (**Fig.** 1.e.).

Comparison of hybrid models to knowledge-based models

Like the previous analysis, we sought to evaluate the effects of adding transcriptomics

information to the existing predictive models from the literature. Classifications from our

mitochondrial and transporter models were compared to those of MitoCarta and TrSSP.

The combined transcriptomic and MitoCarta/TrSSP classifiers appear to only show

363   slight differences (**Fig.** 2.a.,b.). The mitochondrial classifier identifies slightly more

364   mitochondrial genes than MitoCarta alone, however, there are more false positives

365   (FPs) in this classifier (at a classification threshold of 0.75).

367
368   **Figure 2:** Confusion matrices of existing mitochondrial and transporter models (left) and models produced

369   here (right). (**a**) MitoCarta compared to our mitochondrial RF classifier. (**b**) TrSSP compared to our

370   transporter RF classifier. Matrices are colored based on the number of samples contained within each

371   quadrant. A classification threshold of 0.75 was used for the mitochondrial and transporter classifiers as it

provided the best balance of maximizing true positives and minimizing false positives. (**c**) Venn diagram of false positive mitochondrial genes predicted by MitoCarta 3.0 and by the mitochondrial RF classifier produced here. (**d**) Venn diagram of false positive transporter genes predicted by TrSSP and by the transporter RF classifier produced here. (**e-h**) GO enrichment analysis of genes predicted to localize to mitochondria by RF classifiers with no existing evidence in localization databases. Enrichment plots of false positive mitochondrial genes for KEGG pathways (**e**) cellular component (**f**), molecular function (**g**), and biological process (**h**) GO terms.

False positive mitochondrial and transporter genes

Interestingly, our classifiers predict more false positives compared to the results of MitoCarta and TrSSP, suggesting potential for identifying novel functions. To assess this, mitochondrial FP genes were compared to other sources of mitochondrial genes beyond the high confidence GO annotations used for classifier training. Low confidence GO annotations that were originally removed from the training set, Human Protein Atlas subcellular localization annotations from histology images[57], Swiss-Prot annotations[58], and integrated mitochondrial protein index (IMPI) annotations based on MitoMiner[59] were used for other sources of localization evidence. Of the 844 FP genes predicted by the mitochondrial RF classifier, 615 were predicted by both MitoCarta and RF, and 229 were predicted by RF only (**Fig.** 2.c.). 206 of these RF FP genes have no evidence in the other datasets, while the remaining have evidence in at least one source (**Supp. Data**). Enrichment analysis revealed that the 206 unannotated FPs without existing evidence are significantly enriched in KEGG pathways (**Fig.** 2.e.) and GO terms (**Fig.** 2.f.-h.) that are associated with degradation pathways, such as the proteasome, RNA stability, degradation and processing pathways, and are furthermore highly associated with one another based on high confidence evidence through STRING[60] (**Supp. Fig.** 4). The proteasome system is closely linked to mitochondrial remodeling in several physiological and disease states, including cancer[61,62].

Comparisons of FPs in TrSSP and our transporter RF classifier find less overlap between the two (**Fig.** 2.d.). 147 FPs are exclusive to TrSSP, 218 are exclusive to the

RF classifier, and 1477 are common to both models. In the same fashion as

mitochondrial genes, other evidence sources beyond the high confidence GO term

annotations were compared to the FPs predicted by transporter models. Low

confidence AMiGO annotations, Transporter Classifier Data Base annotations[63], and the

SLCAtlas[64] were used as alternate evidence sources. Swiss-Prot annotations were

excluded as these were used in TrSSP model training. For both TrSSP and our RF

classifier, only a small number of FP genes have evidence from other sources (2 for

TrSSP, and 8 for the RF classifier, **Supp. Data**). Unlike the mitochondrial FPs by our RF

classifier, transporter FP genes are not extensively enriched in specific GO terms

(**Supp. Fig.** 5), and no enrichment of KEGG pathways is observed. The most apparent

enrichment is in biological process terms related to cell-cell interactions, such as cell

adhesion, cell junctions, and synapses (**Supp. Fig.** 5.c.).

mGSH transporter candidates

The results of classifiers identify several potential mGSH transporters. Within the SLC25

family, the known transporter SLC25A39 is among the most probable GSH-related

genes (**Table 2**). Surprisingly, SLC25A40, the homolog of SLC25A39, has a very low

GSH probability by the RF classifier models and is not predicted to be related to GSH.

Amongst the top candidates are several well characterized proteins such as SLC25A1,

SLC25A10, and A11 as well as some orphan members like SLC25A43 and A50. To

understand the possible roles of these members in GSH transport, existing evidence

was considered and described below.

Rank	Gene Symbol	GSH probability
1	SLC25A1	0.7891 (0.0173)
2	SLC25A10	0.7800 (0.0256)
3	SLC25A13	0.7737 (0.0156)

Rank	Gene Symbol	GSH probability
1	SLC25A1	0.7891 (0.0173)
2	SLC25A10	0.7800 (0.0256)
3	SLC25A13	0.7737 (0.0156)
4	**SLC25A39**	**0.7716 (0.0230)**
5	SLC25A50	0.7620 (0.0222)
6	SLC25A43	0.7603 (0.0318)
7	SLC25A24	0.7577 (0.0345)
8	SLC25A37	0.7494 (0.0260)
9	SLC25A3	0.7410 (0.0489)
10	SLC25A11	0.7322 (0.0074)
37	**SLC25A40**	**0.4702 (0.0152)**

Table 2: Most probable GSH transporting SLC25 proteins by mean GSH RF classifier probability.
GSH probabilities are mean probabilities with standard error for RF classifiers over different feature sets.
The known GSH transporters SLC25A39 and SLC25A40 are in bold.

Validation of annotation framework

Several other metabolite classifier models were developed for the mGSH transporter
candidates identified by this approach. Like the GSH classifiers, RF models with 5-50
CCLE transcriptomic PCs and corresponding metabolomic correlations as features were
trained for the annotation of genes related to 2-oxoglutarate, glutamate, and carnitine.

Following the procedure for the GSH classification task, positively classified genes were
identified from GO terms for these metabolites to generate training and test gene sets
for classifiers. 2-oxoglutarate, glutamate, and carnitine were selected for this evaluation
because, out of metabolites found in the CCLE metabolomics set, they have a relatively
larger number of annotated genes, making classifier training possible. Additionally,
there is evidence for transport by each for one or more SLC25 proteins, which allows for
some measure of classifier validation. For each of these classifiers, performance was
comparable to the GSH classifier by our evaluation metrics (**Supp. Table** 1, **Table 3**).

Rank	Gene Symbol	Glutamate probability	Rank	Gene Symbol	2-oxoglutarate probability	Rank	Gene Symbol	Carnitine probability
1	SLC25A39	0.7998 (0.0409)	1	SLC25A39	0.8408 (0.0166)	1	SLC25A30	0.8204 (0.0361)
2	SLC25A1	0.7889 (0.0357)	**2**	**SLC25A11**	**0.8304 (0.0228)**	2	SLC25A29	0.8116 (0.0196)
3	SLC25A50	0.7784 (0.0322)	3	SLC25A1	0.8287 (0.0351)	3	SLC25A36	0.7682 (0.0131)
4	SLC25A5	0.7706 (0.0148)	4	SLC25A28	0.8060 (0.0145)	4	SLC25A25	0.7632 (0.0249)
5	SLC25A10	0.7617 (0.0248)	5	SLC25A10	0.8059 (0.0292)	5	SLC25A38	0.7618 (0.0211)
11	**SLC25A13**	**0.7378 (0.0305)**				**28**	**SLC25A20**	**0.6415 (0.0345)**
23	**SLC25A22**	**0.6847 (0.0304)**						
36	**SLC25A12**	**0.6139 (0.0469)**						
42	**SLC25A18**	**0.4663 (0.0178)**						

Table 3: **Top SLC25 candidates for non-GSH metabolites by RF classifiers**. Candidates ranked by
mean RF classifier probabilities (5-50 PC features) with standard error. Known metabolite transporters
are bolded.

Non-SLC25 mGSH-related proteins

Alongside the SLC25 family, the remaining genes in our dataset were considered for

roles in mGSH transport. Top candidates were identified by their mean GSH probability

across RF classifiers. Additionally, the mean mitochondrial and transporter classifier

scores were considered. To identify high confidence candidates, only genes with

probabilities greater than 0.80 across all classifications are selected (i.e., genes already

annotated with one of our GO terms are said to have probabilities of 1). This

thresholding removes 99.95% of the available 49309 genes in the datasets, leaving only

27 genes that meet our criteria (**Supp. Table** 2). Of these, 4 genes are already

annotated with the glutathione metabolic process GO term, 13 are annotated with the

mitochondrion GO term, and 5 with the transmembrane transporter activity term.

Structural comparisons

To further examine the SLC25 proteins most probably relevant to GSH transport, we

next conducted structural analyses. Experimental determinations of protein structures

for the SLC25 family are currently lacking. The first complete human SLC25 structure

was recently determined for UCP1 by cryo-electron microscopy[53]. Beyond this,

structures for orthologs in model species have been determined through x-ray

crystallography for the bovine[54] and *Thermothelomyces thermophila*[65] ADP/ATP carriers

(SLC25A4), and by NMR molecular fragment searching for the yeast UCP2

(SLC25A8)[66]. Predicted protein structures must be used for comparison of family

members with unknown structures. Particularly, we were interested in structural

comparisons between our candidates with the GSH transporters SLC25A39, and

SLC25A40 to reveal any similarities in structure and potential function. Pairwise

structural alignments were determined using TM-align (see Methods) for full 3D

structures ("global" alignments) or tunnel-interacting residues within structures ("local"

alignments). The residues for local alignments were identified via CAVER[56] (**Supp. Fig.**

6). Rather than considering the entire transporter structure, of more relevance to

transporter function and specificity is the interior tunnel of the protein surface, along

which substrates interact and move across the IMM[65,67]. By aligning only potentially

substrate binding residues we get a more relevant analysis of the SLC25 family with

respect to substrate binding as these structures are expected to be more variable and

better indicators of shared transport activities.

Pairwise global alignments showed high similarities (TM-scores close to 1) across the

transporter family (**Supp. Data**). This is expected due to common structural domains

across the family, namely the six transmembrane helices. TM-scores for local

alignments of tunnel residues are much lower relative to the global alignment scores

(mean scores excluding self-alignments 0.376 vs. 0.701) (**Supp. Fig.** 7) and more

variable (coefficient of variation of 0.350 vs. 0.172 for global alignments), indicating the

diversity of transport tunnels and functions of the SLC25 proteins. **Figure** 3.a. details

the TM-scores for local pairwise alignments between top mGSH candidate SLC25s as

well as SLC25A39 and SLC25A40.

489
Figure 3: Structural alignment by TM-align of CAVER-identified transporter tunnels in SLC25 family

members predicted to be associated with glutathione metabolic processes. (**a**) Heatmap of TM-scores for

pairwise alignments of tunnel-interacting regions identified by CAVER for candidate SLC25 members.

Values are TM-scores normalized to the sequence length of the column protein. Highest scoring

alignments for each column are bolded. (**b**) & (**c**) Pymol visualization of transporter tunnel alignments for

SLC25A39 (dark purple) and most similar SLC25 candidates by TM-score, SLC25A43 (green) and

SLC25A24 (orange). Indicated residues are those identified as relevant for GSH transport by SLC25A39

and corresponding residues in SLC25A43 and SLC25A24. Protein tunnels visualized from the

intermembrane space with tunnel-interacting residues identified by CAVER highlighted and non-tunnel

residues faded.

Amongst these results, relatively high scores between SLC25A24, SLC25A43, and

SLC25A39 indicate similarities in transporter tunnels and possible overlapping transport

functions of the proteins. Shi *et al.*[16] identify R225, D226, and K329 as important

residues for GSH transport in SLC25A39. Based on local structural alignments of tunnel

regions with SLC25A43 and SLC25A24 (**Fig.** 3.b. & c.), the position of the K329 residue

is conserved by K275 in SLC25A43 and by K453 in SLC25A24. However, the R225,

D226 motif of SLC25A39 appears less conserved in the aligned structures. In the

SLC25A43 alignment, residues G173 and A174 are in closest proximity to the A39

binding residues, and these non-polar residues are not identified by CAVER as tunnel-

interacting residues. Interestingly, the sidechains of SLC25A43 residues K226 and

R227 appear near R225 and D226 sidechains in SLC25A39, where the positively

charged K226 replaces the negative charge of D226 in SLC25A39. Like SLC25A43, the

closest aligned residues of SLC25A24 to the GSH binding residues of SLC25A39 are

the non-polar G353 and I354, which do not interact with the transporter tunnel. Where

K226 and R227 are found in SLC25A43, instead there is R410 and T411 in SLC25A24.

Comparison to DeepGOPlus

For a comparison to the models developed here, genes from our GO terms of interest

were annotated by DeepGOPlus. DeepGOPlus[24] is a general function annotation deep

learning model based on protein amino acid sequences. The model is constructed as a

convolutional neural network (CNN) in which protein sequences are taken as inputs and

the predicted GO term annotation and probabilities are returned. Gene ontology terms

with predicted annotation probabilities were obtained through the DeepGOWeb API[68]

(Version 1.0.13) using FASTA sequences obtained from canonical Ensembl transcript

IDs for the genes under each ontology term. All 39 high confidence glutathione

metabolic process annotations, and 150 randomly selected genes from each other

ontology term were annotated using DeepGOWeb. The true positive rates (TPRs) or

sensitivity measures of annotations were determined for model comparison (**Fig.** 4,

Supp. Table 3).

530
Figure 4: Evaluation of gene ontology annotation predictions by DeepGOPlus (dashed) and RF classifier

models (solid). Sensitivity, or TPR, are calculated at various classification thresholds.

Our classifiers perform better than DeepGOPlus for annotating all GO terms of interest.

This is most evident for the set of genes involved in glutathione metabolism, where

DeepGOPlus fails to annotate this set of genes with an average TPR of only 0.073 over

classification thresholds compared to 0.867 for the RF classifier. DeepGOPlus can

reasonably classify mitochondrial localization and transporter activity annotations with

mean TPRs of 0.525 and 0.722, respectively.

Discussion

The importance of mitochondria and GSH in cancers is widely appreciated.

Mitochondria have a major role in metabolic shifts, one of the hallmarks of cancer, while

mGSH is important in metabolism, ROS processing, post-translational protein

modifications and tumor drug resistance. An improved understanding of mitochondrial

transport mechanisms in cancer cells, including GSH transport mechanisms, is

important for advancing the cancer biology field. To this end we developed a hybrid ML

framework that utilizes multi-omics data and existing knowledge for annotating genes

with GO terms of interest. We applied this model to the problem of mitochondrial GSH

metabolism, specifically focusing on the aspect of transport. We propose several

potential candidates that are predicted to be related to, and possibly modulated by,

mGSH transport that are identified by this model to guide future experimentation in this

area of research.

Random forest classifiers were developed here for the annotation of mGSH transporter

characteristics: mitochondrial localization, relation to GSH metabolism, and transporter

function. Models combine CCLE transcriptomics and metabolomics data for annotation

of genes related to GSH metabolism, and knowledge from existing predictive models for

mitochondrial localization and transporter activity annotation. This framework resulted in

models with high performance (mean RF AUROC of 0.900) across each classification

task.

For comparison, we attempted similar classifications using DeepGOPlus[24,68] for

annotation of our terms of interest. This general GO annotation model performed worse,

as measured by sensitivity in correctly classifying annotated genes, than our framework

This is most evident in GSH metabolic process classification, likely due to the relatively

small set of annotated genes for this term which is underrepresented during training in

the DeepGOPlus framework, indicating a weakness of these general frameworks and a

strength of our approach. Furthermore, as a sequence-based approach, DeepGOPlus

cannot be applied to disease contexts as we do with our proposed method, in this case

using cancer cell transcriptomics and metabolomics data to focus analysis on a cancer

context.

As a further validation of our methodology, several other classifiers using our framework

were developed and tested for annotating other GO terms. Identical RF models to the

GSH classifier were developed which instead use genes annotated with GO terms for

glutamate, 2-oxoglutarate, and carnitine metabolic process for training. These models

show similar performance to the GSH classifier despite a small number of annotated

genes for training, indicating the robustness of this framework for annotating a variety of

biological functions. Furthermore, with respect to our interest in transport, these

classifiers correctly identify many previously known SLC25 transporters as their most

probable candidates.

Considering the SLC25 family of mitochondrial carriers, we find several predicted to be

related to GSH metabolic processes. Most notably, SLC25A39, the recently identified

GSH transporter is amongst our top hits (ranked 4th). Furthermore, several SLC25

which are known to be relevant to mGSH metabolism are amongst the top candidates:

SLC25A1[69,70] (ranked 1st), SLC25A10[71,72] (ranked 2nd), SLC25A13[73] (ranked 3rd), and

the iron transporter SLC25A37[16] (ranked 8th). Surprisingly, the homolog of SLC25A39,

SLC25A40, is not predicted to be involved in GSH metabolism, ranking 37th of the 53

SLC25 family proteins. This may be due to the very low expression of SLC25A40

across the CCLE cell lines (**Supp. Fig.** 2), suggesting that SLC25A40 is quantitatively

less relevant to mGSH transport in cancer cells. Other potential mitochondrial carriers of

interest include SLC25A43 (ranked 6th), SLC25A24 (ranked 7th), and the OMM

transporter SLC25A50 (ranked 5th). SLC25A43 is an orphan transporter which has

recently been found to affect redox homeostasis[74], but the other candidates have not

been, to our knowledge, associated with GSH metabolism.

Structural similarities in the candidates, specifically within the tunnel region of the

proteins, can inform potential roles of these transporters in mGSH metabolism. We

investigated this through sequence-independent structural alignment of SLC25

transporters using TMalign. We find that, aside from its homolog SLC25A40, SLC25A39

shows the most similarities to our candidates SLC25A43, and SLC25A24. Important

GSH binding residues in SLC25A39 are somewhat conserved in our candidates where,

specifically, SLC25A43 shows a shift in residue position within the transporter tunnel

and a substitution of an aspartate to a lysine residue, and SLC25A24 substitutes a

threonine. These findings indicate that despite overall similarities in transporter tunnels

between the GSH transporter and our candidates SLC25A43 and SLC25A24, substrate

binding and transport mechanism are likely somewhat different, e.g., binding to

alternate forms of GSH species such as GSSG or glutathione esters.

Expanding our search beyond the SLC25 family, 27 genes were classified with high

probability (>0.80) for all classification tasks. Of these, some of the most probable

candidates are discussed here for their possible roles in mGSH metabolism and

transport.

Pyruvate carboxylase (PC) catalyzes the conversion of pyruvate to oxaloacetate in

mitochondria. PC links glucose to GSH synthesis and plays roles in the control of redox

and oxidative stress[75]. Recently, pyruvate metabolism has been associated with GSH

metabolism and ferroptosis in lung cancer cells through the plasma membrane cysteine

transporter SLC7A11[76] (ranked 3[rd] by GSH metabolic process GO term annotation

probability in RF classifiers. PC is reported to specifically localize to the mitochondrial

matrix with little evidence supporting it being membrane-bound; thus, it may be that PC

interacts with the IMM-bound transporters to facilitate and modulate mGSH transport.

Mitochondrial calcium uniporter (MCU) is also a top hit in our models and a well-known

mitochondrial transporter. Associated with its role in Ca^{2+} homeostasis are roles in iron

and redox homeostasis, and cell death pathways[77]. With respect to GSH, MCU is

regulated via s-glutathionylation[77] and downregulation is associated with enriched GSH

metabolism in melanoma cells[78]. It is possible that MCU plays a greater role in

mitochondrial iron and GSH metabolism and transport in the context of cancer

metabolism than has been reported thus far.

ATP-binding cassette family B6 (ABCB6) is a transporter protein with evidence for both

plasma membrane and mitochondrial localization[79]. This protein transports the heme

precursor porphyrin[80], however the mechanism of transport has been shown to be GSH-

dependent, with significantly increased activity in the presence of GSH[81]. These suggest

GSH-mediated transport of substrates by ABCB6 but it is unclear if GSH merely

modulates transport, or if it is transported as well, likely through GSH-conjugated

species. Similarly, neudesin neurotrophic factor (NENF) is a mitochondrial localized

protein primarily involved in the differentiation/development of neuronal cells[82]. NENF

activity is modulated by the binding of heme[83], which presents a connection to GSH

that, to the best of our knowledge, has not yet been explored. Some evidence suggests

NENF is membrane-associated[82].

The results of our models find several genes that may play roles in mGSH metabolism

and transport, and thus could guide future experimental analyses. Our findings are

limited by our relatively small training dataset, in which more data would improve model

performance and confidence of predictions. Furthermore, our models are generalized

across cancer cell data from a range of tissue and cancer types. Future work,

leveraging larger datasets, or datasets of specific cancer types would provide promising

avenues for uncovering interactions of high relevance to the specific diseases. Another

limitation of this work is the use of gene ontology terms for identifying relevant genes.

Specific and manually curated databases for these functions, similar to MitoCarta, likely

provide a more accurate set of relevant genes; however, these databases are limited.

Using a database like gene ontology provides a resource for the annotation of many

functions, processes, and cell localizations as demonstrated here, which cannot be

replicated by specific databases.

Finally, the classifications from the individual models provide novel candidates for areas

beyond the scope in this work. For example, the classifiers for other metabolites here

were used for validation of the glutathione classifier, but predicted relevant genes for

those metabolites were not investigated. Nevertheless, we anticipate that our findings

here can be instrumental in the identification of potentially novel proteins involved in

mitochondrial glutathione metabolism and transport in cancer cells. The results here

and classifier code provide a tool to accelerate knowledge discovery and identify

potential target genes for other biological functions and contexts.

Abbreviations

GSH: Glutathione

mGSH: Mitochondrial glutathione

ML: Machine learning

DL: Deep learning

ROS: Reactive oxygen species

OXPHOS: Oxidative phosphorylation

KEGG: Kyoto Encyclopedia of Genes and Genomes

CCLE: Cancer Cell Line Encyclopedia

GO: Gene Ontology

TrSSP: The Transporter Substrate Specificity Prediction Server

GSSG: Glutathione disulfide

PCA: Principal component analysis

PC: Principal component

RF: Random Forest

DT: Decision Tree

NB: Naive Bayes

SVM: Support Vector Machine

ROC: Receiver operating characteristic

PRC: Precision-recall curve

TP: True positive

TN: True negative

FP: False positive

FN: False negative

AUROC: Area under receiver operating characteristic curve

AUPRC: Area under precision-recall curve

MCC: Matthew's correlation coefficient

CNN: Convolutional neural network

TPR: True positive rate

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

694   All data and code used and generated in this work are publicly available at

695   https://github.com/lkenn012/mGSH_cancerClassifiers[84].

Competing interests

697   The authors declare no competing interests.

Funding

699   This research was funded by the National Research Council Canada's New Beginnings

700   Initiative-Ideation Fund (MH, MCC), Canadian Institutes of Health Research (MH FDN

701   143278) and the NSERC-CREATE Metabolomics Advanced Training and International

702   Exchange Program (MATRIX).

Author contributions

704   L.K. designed and trained models and performed all other analyses. L.K. M.-E.H., and

705   M.C.-C. interpreted the results and wrote the manuscript. M.-E.H., and M.C.-C., jointly

706   supervised the study and acquired funding. L.K. M.-E.H., and M.C.-C. envisioned the

707   study. All authors provided feedback and suggestions to develop the research and

708   manuscript. All authors reviewed the manuscript.

References

1. Pavlova NN, Zhu J, Thompson CB (2022) The hallmarks of cancer metabolism: Still emerging. Cell Metab 34:355–377

2. Kennedy L, Sandhu JK, Harper M-E, Cuperlovic-Culf M (2020) Role of Glutathione in Cancer: From Mechanisms to Therapies. Biomolecules. https://doi.org/10.3390/biom10101429

3. Lien EC, Lyssiotis CA, Juvekar A, Hu H, Asara JM, Cantley LC, Toker A (2016) Glutathione biosynthesis is a metabolic vulnerability in PI(3)K/Akt-driven breast cancer. Nat Cell Biol 18:572–578

4. Harris IS, Treloar AE, Inoue S, et al (2015) Glutathione and thioredoxin antioxidant pathways synergize to drive cancer initiation and progression. Cancer Cell 27:211–222

5. Kong H, Chandel NS (2018) Regulation of redox balance in cancer and T cells. J Biol Chem 293:7499–7507

6. Piskounova E, Agathocleous M, Murphy MM, Hu Z, Huddlestun SE, Zhao Z, Leitch AM, Johnson TM, DeBerardinis RJ, Morrison SJ (2015) Oxidative stress inhibits distant metastasis by human melanoma cells. Nature 527:186–191

7. McGuirk S, Audet-Delage Y, Annis MG, et al (2021) Resistance to different anthracycline chemotherapeutics elicits distinct and actionable primary metabolic dependencies in breast cancer. Elife. 10:e65150.

8. Yang WS, SriRamaratnam R, Welsch ME, et al (2014) Regulation of Ferroptotic Cancer Cell Death by GPX4. Cell 156:317–331

9. Ballatori N, Krance SM, Marchan R, Hammond CL (2009) Plasma membrane glutathione transporters and their roles in cell physiology and pathophysiology. Mol Aspects Med 30:13–28

10. Wallace DC (2012) Mitochondria and cancer. Nat Rev Cancer 12:685–698

11. Whitaker HC, Patel D, Howat WJ, et al (2013) Peroxiredoxin-3 is overexpressed in prostate cancer and promotes cancer cell survival by protecting cells from oxidative stress. Br J Cancer 109:983–993

738   12.   Oestreicher J, Morgan B (2018) Glutathione: subcellular distribution and
739   membrane transport1. Biochem Cell Biol. https://doi.org/10.1139/bcb-2018-0189

740   13.   Chen Z, Lash LH (1998) Evidence for mitochondrial uptake of glutathione by
741   dicarboxylate and 2-oxoglutarate carriers. J Pharmacol Exp Ther 285:608–618

742   14.   Booty LM, King MS, Thangaratnarajah C, Majd H, James AM, Kunji ERS,
743   Murphy MP (2015) The mitochondrial dicarboxylate and 2-oxoglutarate carriers
744   do not transport glutathione. FEBS Lett 589:621–628

745   15.   Wang Y, Yen FS, Zhu XG, et al (2021) SLC25A39 is necessary for mitochondrial
746   glutathione import in mammalian cells. Nature 599:136–140

747   16.   Shi X, Reinstadler B, Shah H, et al (2022) Combinatorial GxGxE CRISPR screen
748   identifies SLC25A39 in mitochondrial glutathione transport linking iron
749   homeostasis to OXPHOS. Nat Commun 13:2483

750   17.   von Bohlen und Halbach O (2022) Controlling glutathione entry into
751   mitochondria: potential roles for SLC25A39 in health and (treatment of) disease.
752   Signal Transduction and Targeted Therapy 7:1–2

753   18.   Cooper AJL, Hanigan MH (2018) Metabolism of glutathione S-conjugates:
754   Multiple pathways. In: Comprehensive Toxicology. Elsevier, pp 363–406

755   19.   Ramsay EE, Dilda PJ (2014) Glutathione S-conjugates as prodrugs to target
756   drug-resistant tumors. Front Pharmacol 5:181

757   20.   Jumper J, Evans R, Pritzel A, et al (2021) Highly accurate protein structure
758   prediction with AlphaFold. Nature 596:583–589

759   21.   Baek M, DiMaio F, Anishchenko I, et al (2021) Accurate prediction of protein
760   structures and interactions using a three-track neural network. Science 373:871–
761   876

762   22.   Zhou N, Jiang Y, Bergquist TR, et al (2019) The CAFA challenge reports
763   improved protein function prediction and new functional annotations for hundreds
764   of genes through experimental screens. Genome Biol 20:1–23

765   23.   Kulmanov M, Khan MA, Hoehndorf R (2017) DeepGO: predicting protein
766   functions from sequence and interactions using a deep ontology-aware classifier.
767   Bioinformatics 34:660–668

768   24.   Kulmanov M, Hoehndorf R (2019) DeepGOPlus: improved protein function
769   prediction from sequence. Bioinformatics 36:422–429
770   25.   Zhou J, Xiong W, Wang Y, Guan J (2021) Protein Function Prediction Based on
771   PPI Networks: Network Reconstruction vs Edge Enrichment. Front Genet
772   12:758131
773   26.   Hu S, Luo Y, Zhang Z, Xiong H, Yan W, Jiang M, Zhao B (2022) Protein function
774   annotation based on heterogeneous biological networks. BMC Bioinformatics
775   23:1–16
776   27.   Lei Y, Li S, Liu Z, Wan F, Tian T, Li S, Zhao D, Zeng J (2021) A deep-learning
777   framework for multi-level peptide–protein interaction prediction. Nat Commun
778   12:1–10
779   28.   Shanehsazzadeh A, Bachas S, McPartlon M, et al (2023) Unlocking de novo
780   antibody design with generative artificial intelligence. bioRxiv 2023.01.08.523187
781   29.   Kunc V, Klema J (2022) On functional annotation with gene co-expression
782   networks. 2022 IEEE International Conference on Bioinformatics and
783   Biomedicine (BIBM). https://doi.org/10.1109/bibm55620.2022.9995542
784   30.   Wekesa JS, Luan Y, Meng J (2021) Predicting Protein Functions Based on
785   Differential Co-expression and Neighborhood Analysis. J Comput Biol 28:1
786   31.   Wang Y, Yang S, Zhao J, Du W, Liang Y, Wang C, Zhou F, Tian Y, Ma Q (2019)
787   Using Machine Learning to Measure Relatedness Between Genes: A Multi-
788   Features Model. Sci Rep 9:1–15
789   32.   Karpatne A, Kannan R, Kumar V (2022) Knowledge Guided Machine Learning:
790   Accelerating Discovery using Scientific Knowledge and Data. CRC Press
791   33.   Rosenberg AM, Saggar M, Monzel AS, et al (2023) Brain mitochondrial diversity
792   and network organization predict anxiety-like behavior in male mice. Nat
793   Commun 14:1–19
794   34.   Liu F, Heiner M, Gilbert D (2022) Hybrid modelling of biological systems: current
795   progress and future prospects. Brief Bioinform.
796   https://doi.org/10.1093/bib/bbac081
797   35.   Elmarakeby HA, Hwang J, Arafeh R, et al (2021) Biologically informed deep
798   neural network for prostate cancer discovery. Nature 598:348–352

36. Yu K, Chen B, Aran D, Charalel J, Yau C, Wolf DM, van 't Veer LJ, Butte AJ,
Goldstein T, Sirota M (2019) Comprehensive transcriptomic analysis of cell lines
as models of primary tumors across 22 tumor types. Nat Commun 10:1–11

37. Rath S, Sharma R, Gupta R, et al (2021) MitoCarta3.0: an updated mitochondrial
proteome now with sub-organelle localization and pathway annotations. Nucleic
Acids Res 49:D1541–D1547

38. Mishra NK, Chang J, Zhao PX (2014) Prediction of membrane transport proteins
and their substrate specificities using primary sequence information. PLoS One
9:e100278

39. Pedregosa F, Varoquaux G, Gramfort A, et al (2012) Scikit-learn: Machine
Learning in Python. arXiv. https://doi.org/10.48550/arXiv.1201.0490

40. Hunter JD (2007) Matplotlib: A 2D Graphics Environment. Comput Sci Eng 9:90–
95

41. Waskom M, Botvinnik O, O'Kane D, et al (2017) mwaskom/seaborn: v0.8.1
(September 2017). https://doi.org/10.5281/zenodo.883859

42. Ge SX, Jung D, Yao R (2020) ShinyGO: a graphical gene-set enrichment tool for
animals and plants. Bioinformatics 36:2628–2629

43. Li H, Ning S, Ghandi M, et al (2019) The landscape of cancer cell line
metabolism. Nat Med 25:850–860

44. Beretta L, Santaniello A (2016) Nearest neighbor imputation algorithms: a critical
evaluation. BMC Med Inform Decis Mak 16:197–208

45. Fisher RA (1915) Frequency Distribution of the Values of the Correlation
Coefficient in Samples from an Indefinitely Large Population. Biometrika 10:507–
521

46. Ashburner M, Ball CA, Blake JA, et al (2000) Gene ontology: tool for the
unification of biology. The Gene Ontology Consortium. Nat Genet 25:25–29

47. Carbon S, Ireland A, Mungall CJ, Shu S, Marshall B, Lewis S, AmiGO Hub, Web
Presence Working Group (2009) AmiGO: online access to ontology and
annotation data. Bioinformatics 25:288–289

48. Mudunuri U, Che A, Yi M, Stephens RM (2009) bioDBnet: the biological
database network. Bioinformatics 25:555

49.   Matthews BW (1975) Comparison of the predicted and observed secondary
831   structure of T4 phage lysozyme. Biochimica et Biophysica Acta (BBA) - Protein
832   Structure 405:442–451

833   50.   Chicco D, Jurman G (2020) The advantages of the Matthews correlation
834   coefficient (MCC) over F1 score and accuracy in binary classification evaluation.
835   BMC Genomics 21:1–13

836   51.   Waterhouse A, Bertoni M, Bienert S, et al (2018) SWISS-MODEL: homology
837   modelling of protein structures and complexes. Nucleic Acids Res 46:W296–
838   W303

839   52.   Pebay-Peyroula E, Dahout-Gonzalez C, Kahn R, Trézéguet V, Lauquin GJ-M,
840   Brandolin G (2003) Structure of mitochondrial ADP/ATP carrier in complex with
841   carboxyatractyloside. Nature 426:39–44

842   53.   Kang Y, Chen L (2023) Structural basis for the binding of DNP and purine
843   nucleotides onto UCP1. Nature 620:226–231

844   54.   Zhang Y, Skolnick J (2005) TM-align: a protein structure alignment algorithm
845   based on the TM-score. Nucleic Acids Res 33:2302–2309

846   55.   The PyMOL Molecular Graphics System, Version 2.4.1 Schrödinger, LLC.

847   56.   Chovancova E, Pavelka A, Benes P, et al (2012) CAVER 3.0: A Tool for the
848   Analysis of Transport Pathways in Dynamic Protein Structures. PLoS Comput
849   Biol 8:e1002708

850   57.   Thul PJ, Lindskog C (2018) The human protein atlas: A spatial map of the human
851   proteome. Protein Sci 27:233–244

852   58.   The UniProt Consortium, Bateman A, Martin M-J, et al (2022) UniProt: the
853   Universal Protein Knowledgebase in 2023. Nucleic Acids Res 51:D523–D531

854   59.   Smith AC, Robinson AJ (2019) MitoMiner v4.0: an updated database of
855   mitochondrial localization evidence, phenotypes and diseases. Nucleic Acids Res
856   47:D1225–D1228

857   60.   Szklarczyk D, Gable AL, Lyon D, et al (2019) STRING v11: protein-protein
858   association networks with increased coverage, supporting functional discovery in
859   genome-wide experimental datasets. Nucleic Acids Res 47:D607–D613

61.   Yue S, Wang L, DeMartino GN, Zhao F, Liu Y, Sieber MH (2022) Highly
conserved shifts in ubiquitin-proteasome system (UPS) activity drive
mitochondrial remodeling during quiescence. Nat Commun 13:1–14

62.   Sharma A, Nair R, Achreja A, et al (2022) Therapeutic implications of
mitochondrial stress-induced proteasome inhibitor resistance in multiple
myeloma. Sci Adv 8:eabq5575

63.   Saier MH, Reddy VS, Moreno-Hagelsieb G, et al (2020) The Transporter
Classification Database (TCDB): 2021 update. Nucleic Acids Res 49:D461–D467

64.   Gyimesi G, Hediger MA (2022) Systematic in silico discovery of novel solute
carrier-like proteins from proteomes. PLoS One 17:e0271062

65.   Ruprecht JJ, King MS, Zögg T, Aleksandrova AA, Pardon E, Crichton PG,
Steyaert J, Kunji ERS (2019) The Molecular Mechanism of Transport by the
Mitochondrial ADP/ATP Carrier. Cell 176:435–447.e15

66.   Berardi MJ, Shih WM, Harrison SC, Chou JJ (2011) Mitochondrial uncoupling
protein 2 structure determined by NMR molecular fragment searching. Nature
476:109–113

67.   Ruprecht JJ, Kunji ERS (2020) The SLC25 Mitochondrial Carrier Family:
Structure and Mechanism. Trends Biochem Sci 45:244–258

68.   Kulmanov M, Zhapa-Camacho F, Hoehndorf R (2021) DeepGOWeb: fast and
accurate protein function prediction on the (Semantic) Web. Nucleic Acids Res
49:W140–W146

69.   Fernandez HR, Gadre SM, Tan M, et al (2018) The mitochondrial citrate carrier,
SLC25A1, drives stemness and therapy resistance in non-small cell lung cancer.
Cell Death Differ 25:1239–1258

70.   Wadey AL, Muyderman H, Kwek PT, Sims NR (2009) Mitochondrial glutathione
uptake: characterization in isolated brain mitochondria and astrocytes in culture.
J Neurochem 109 Suppl 1:101–108

71.   Punzi G, Porcelli V, Ruggiu M, et al (2017) SLC25A10 biallelic mutations in
intractable epileptic encephalopathy with complex I deficiency. Hum Mol Genet
27:499–504

72.   Zhang D, Sun T, Li Y, et al (2022) Mitochondrial outer membrane protein
891   FUNDC2 promotes ferroptosis and contributes to doxorubicin-induced
892   cardiomyopathy. Proceedings of the National Academy of Sciences 119:

893   73.   Palmieri L, Pardo B, Lasorsa FM, et al (2001) Citrin and aralar1 are Ca(2+)-
894   stimulated aspartate/glutamate transporters in mitochondria. EMBO J 20:5060–
895   5069

896   74.   Zhang J, Zhao Y, Tian Y, Geng M, Liu Y, Zhang W, Shuai L (2022) Genome-
897   wide screening in the haploid system reveals Slc25a43 as a target gene of
898   oxidative toxicity. Cell Death Dis 13:1–11

899   75.   Fu A, van Rooyen L, Evans L, et al (2021) Glucose metabolism and pyruvate
900   carboxylase enhance glutathione synthesis and restrict oxidative stress in
901   pancreatic islets. Cell Rep 37:110037

902   76.   Zhao L, Deng H, Zhang J, et al (2023) A non-canonical function of LDHB
903   promotes SLC7A11-mediated glutathione metabolism and protects against
904   glutaminolysis-dependent ferroptosis in KRAS-driven lung cancer. bioRxiv
905   2023.02.12.525859

906   77.   Dong Z, Shanmughapriya S, Tomar D, et al (2017) Mitochondrial Ca2+ Uniporter
907   is a Mitochondrial Luminal Redox Sensor That Augments MCU Channel Activity.
908   Mol Cell 65:1014

909   78.   Stejerean-Todoran I, Zimmermann K, Gibhardt CS, et al (2022) MCU controls
910   melanoma progression through a redox-controlled phenotype switch. EMBO Rep
911   23:e54746

912   79.   Song G, Zhang S, Tian M, Zhang L, Guo R, Zhuo W, Yang M (2021) Molecular
913   insights into the human ABCB6 transporter. Cell Discovery 7:1–11

914   80.   Xie L-H, Fefelova N, Pamarthi SH, Gwathmey JK (2022) Molecular Mechanisms
915   of Ferroptosis and Relevance to Cardiovascular Disease. Cells 11:2726

916   81.   Wang C, Cao C, Wang N, Wang X, Wang X, Zhang XC (2020) Cryo-electron
917   microscopy structure of human ABCB6 transporter. Protein Sci 29:2363

918   82.   Kimura I, Nakayama Y, Zhao Y, Konishi M, Itoh N (2013) Neurotrophic effects of
919   neudesin in the central nervous system. Front Neurosci 7:54475

920 83. Moutaoufik MT, Malty R, Amin S, et al (2019) Rewiring of the Human

921   Mitochondrial Interactome during Neuronal Reprogramming Reveals Regulators

922   of the Respirasome and Neurogenesis. iScience 19:1114–1132

923 84. GitHub - lkenn012/mGSH_cancerClassifiers: Code, data, and other information

924   for mGSH transporter models. In: GitHub.

925   https://github.com/lkenn012/mGSH_cancerClassifiers. Accessed 18 Sep 2023

926 85. Fellows Yates JA, Lamnidis TC, Borry M, Valtueña AA, Fagernäs Z, Clayton S,

927   Garcia MU, Neukamm J, Peltzer A (2021) Reproducible, portable, and efficient

928   ancient genome reconstruction with nf-core/eager. PeerJ 9:e10947