Developmental effect of RASopathy mutations on neuronal network activity on a chip

Eva-Maria Weiss[1*], Debarpan Guhathakurta[1], Aneta Petrušková [1,2,3], Verena Hundrup[1], Martin Zenker[4], Anna Fejtová[1*]

[1]Department of Psychiatry and Psychotherapy, Universitätsklinikum Erlangen, Friedrich-Alexander Universität Erlangen-Nürnberg, Erlangen, Germany

[2]Charles University, Third Faculty of Medicine, Prague, CZ

[3]National Institute of Mental Health, Klecany, CZ

[4] Institute of Human Genetics, University Hospital Magdeburg, Medical Faculty, Otto-von-Guericke University, Magdeburg, Germany

* Correspondence:

Eva-Maria Weiss (Eva-Maria.Weiss@uk-erlangen.de) and Anna Fejtova (Anna.Fejtova@uk-erlangen.de)

Abstract

RASopathies are a group of genetic disorders caused by mutations in genes encoding components and regulators of the RAS/MAPK signaling pathway, resulting in an overactivation in signaling. RASopathy patients exhibit distinctive facial features, cardiopathies, growth and skeletal abnormalities, and varying degrees of developmental delay, neurocognitive impairment, intellectual disabilities, and attention deficits. At present, it is unclear how RASopathy mutations cause neurocognitive impairment and what their neuron-specific cellular and network phenotypes are. Such knowledge would be crucial for the development of specific therapies, which are still lacking for the treatment of the neurocognitive and behavioral symptoms in RASopathies. Here, we investigated the effect of RASopathy mutations on establishment and functional maturation of neuronal networks. We isolated cortical neurons from RASopathy mouse models and cultured them for several weeks in multi-well plates with implanted multi-electrode arrays. The arrays were used for longitudinal recordings of spontaneous activity in developing networks as well as for recordings of evoked responses in mature neurons. To facilitate the analysis of large and complex data sets resulting from long-term multi-electrode recordings we developed MATLAB-based tools for the data processing, analysis and statistical evaluation. Longitudinal analysis of spontaneous network activity revealed a convergent developmental phenotype in neurons carrying the gain-of-function RASopathy mutations $Ptpn11^{D61Y}$ and $Kras^{V14I}$. The phenotype was more pronounced at early time points and faded out over time suggesting compensatory effects during network maturation. Nevertheless, persistent differences in excitatory/inhibitory balance and network excitability were observed in mature networks. This study improves the understanding of the complex relationship between genetic mutations and clinical

manifestations in RASopathies by adding insights into functional network processes as an additional piece of the puzzle.

Keywords

neurodevelopmental disorder, multielectrode array, principle component analysis, rare diseases, mouse models, excitation/ inhibition balance, Noonan syndrome, network establishment and maturation

Acknowledgement

We are deeply grateful to all those who contributed to the success of this research project. First, we would like to thank our cooperation partner from stem cell biology department at the University Hospital Erlangen, particularly Prof. Dr. Beate Winner, Dr. Tania Rizo and Michaela Farrell for great technical support. Furthermore, we thank the group of Prof. Dr. Thomas Winkler at the department of biology at the Friedrich-Alexander University of Erlangen for their support. The study was funded by BMBF Project GeNeRARe: German Research Network for RASopathies (01GM1902B) and cofounded by EURAS (Grant Agreement No.: 101080580).

Disclaimer

Funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the Health and Digital Executive Agency. Neither the European Union nor the granting authority can be held responsible for them.

Introduction

RASopathies cover a group of genetic disorders associated with a characteristic pattern of anomalies including distinctive facial features, cardiopathies, growth and skeletal abnormalities and various degrees of developmental delay [43]. They are typically caused by germline mutations in genes encoding for components or regulators of the RAS/ mitogen-activated protein kinase (MAPK) signaling pathway. Common is an overactivation in signaling, either by activating mutations in the pathway components or by loss of function in negative modulators [39, 43]. The RAS/MAPK signaling pathway has a central role in cellular response to extracellular stimuli [27, 42]. In neurons, it's indispensable for proper differentiation, synapse formation and plasticity and is therefore critical for brain development, learning, memory and cognition [34]. Indeed, most RASopathies are associated with neurocognitive impairments and variable degree of intellectual disability [7, 18, 38]. While patients with Cardio-facio-cutaneous syndrome (CFC), Costello syndrome (CS) and SYNGAP1-related encephalopathy show moderate to severe intellectual disabilities and neurodevelopmental delay, epilepsy and autism spectrum disorder [1, 5, 30, 32, 36], patients with Noonan Syndrome (NS) and Neurofibromatosis type 1 are in general less severely affected. Common diagnoses are mild neurodevelopmental delay, impaired executive functioning, learning difficulties and attention deficit hyperactivity disorder symptomatology [2, 19, 21, 26, 29, 31, 37].

Interestingly, some of the neurocognitive features diminish with increasing age of the patients, while different emerge. For example, children with NS have shown more extensive cognitive problems concomitant with learning difficulties and intelligence impairment compared to adults [33].

Animal studies have not yet addressed the developmental aspect systematically. While most studies demonstrated electrophysiological and behavioral phenotypes in mice carrying different RASopathy mutations, most of them did not assess the respective phenotypes along the animal lifespan [3, 10, 11, 23]. An exception is a recent study that used mice where RASopathy-related $Kras^{G12V}$ mutation was expressed exclusively in neurons. In hippocampi of these animals an upregulation of RAS/MAPK signaling was seen during early phases of postnatal development, but not in the adult state [28]. Interestingly, the normalization of RAS/MAPK pathway activity was concomitant with enhanced GABAergic synaptogenesis indicating that maladaptive development of neuronal network might underly changes in RASopathy-induced neurocognitive effects observed during lifespan in patients. Importantly, the switch in the cellular mechanism underlying the network dysfunction might also have consequences on the selection of effective treatment strategy. In line with that, Papale and colleagues were able to improve the cognitive phenotypes in $Kras^{G12V}$ mice by treatment with RAS/MAPK inhibitors applied in juvenile animals. Strikingly, the same treatment was not effective if applied in older animals. Here, in line with increased GABAergic synaptogenesis, inhibition of GABAergic neurotransmission showed positive effects.

In this study, we follow up the hypothesis that a pathological activation of RAS/MAPK signaling potentially induces maladaptive processes leading to compensatory effects in developing neuronal networks. After all, neural circuits make up regulatory pathways for motion control, cognition, emotion, learning and memory in the brain. However, *in vivo* examination by real-time monitoring of neuronal activity and functional connectivity is restricted, since deep tissue and components of neural circuits are hard to access [6]. As an alternative opportunity to study functional connectivity and network parameters, we investigate neuronal cell cultures derived from RASopathy mouse models plated on multi-well MEA (mwMEA) plates in a longitudinal study. The disease models used in this study bear gain of function mutations in $Ptpn11^{D61Y}$ [3] and in $Kras^{V14l}$ [15, 17, 35]. To perform an extensive functional analysis of spontaneous network activity and electrically evoked activity, we used in-house developed analysis-tool written in MATLAB and routines for the application of principle component analysis (PCA) to reduce dimensionality in the data set for better data visualization and statistical evaluation. We observed significant effects of RASopathy mutation on network activity at early developmental stages that were compensated during network maturation. However, the mature RASopathy networks with apparently normal network activity still exhibited differences in excitatory/inhibitory balance and network excitability.

Material and Methods

Animals

For this study conditional $Ptpn11^{D61Yfloxed/wt}$ mice (B6.129S6-$Ptpn11$tm1Toa/Mmjax, Jackson Laboratories, RRID# MMRRC_032103-JAX) [8] and $Kras^{V14lfloxed/wt}$ mice [17] were used. In these strains a LOX-STOP-LOX cassette is cloned before the exon containing the RASopathy mutation. To generate RASopathy mouse model heterozygote mice bearing the mutation were crossed with mice homozygote for Emx1-cre alleles (B6.129S2-Emx1tm1(cre)Krj/J animals, Jackson Laboratories, RRID# IMSR_JAX:005628, genotyping according to provider's recommendations). This strain expresses cre recombinase in forebrain excitatory neurons and glia from embryonic day 10.5 on. [14]. All mouse strains were bred on C57BL/6N background for more than 5 generations in our laboratory. Mice were kept at 22 ± 2 °C 12 h light-dark cycle with food and water ad libitum. Breeding of animals and experiments using animal material were performed in accordance with local animal welfare officer (FAU: TS12/2016), and in accordance with the European Directive 2010/63/EU. Animal records were kept by Python-based Relational Animal Tracking Software (PYRAT-Scionics Computer Innovation GmbH, Dresden, Germany).

Animals were genotyped using published protocols. Specifically, EMX1-cre genotyping protocol provided by Jackson lab was followed. To detect $Ptpn11^{D61Y}$ allele, polymerase chain reaction (PCR) using TGGAGCTGTTACCCACATCA and GCACAGTTCAGCGGGTACTT primers followed by a melting point analysis using the High Resolution Melting and Gene scanning application on the LightCycler 480 (Roche Diagnostics) was performed according to [3]. To determine the genotype in $Kras^{V14l}$, PCR was performed using AGG GTA GGT GTT GGG ATA GC, CTC AGT CAT TTT CAG CAG GC, CTG CTC TTT ACT GAA GGC TC primers using S7 Fusion High-Fidelity DNA polymerase (Biozym, Cat#MD-S7-100) according to [17]. The protocol was generated to discriminate between 403 bp (wild type) and 621 bp (knock in mutation) alleles. Primers were purchased from Eurofins, Ebersberg, Germany.

Dissociated cell culture and plating on multi-well MEA plate or 18mm coverslips.

Chemical reagents are listed in Supplementary Material Table S 1. The day before dissection, wells of 48-well mwMEA plates (Cytoview Cat#MEA 48, M768-tMEA_48B, Axion Biosystems) and 18mm Menzel glass coverslips (#6311342, VWR International, Radnor, USA) were coated with poly-L-lysine (0.5 mg/ml) followed by overnight incubation at 37°C. The next day wells and coverslips were washed three times with sterile double-deionized water and left in HBSS-/-, which was removed directly before cell seeding. Prior to seeding of cells on 18 mm coverslips, they were coated with 100 µl of Neurobasal™ -A media (NBA) supplemented with 0.2 mM Glutamax, 2% (v:v) B27, 0.1 M Antibiotic-Antimycotic – together named NBA mix and 10% of fetal calf serum (FCS), as previously described [4], for better attachment of neurons.

Dissociated, neuronal cultures were prepared from individual brains of newborn mice (P0-P1) as previously described in [4]. Briefly, *Ptpn11*$^{D61Yfloxed/wt}$ (from here termed as *Ptpn11*^{D61Y}) or *Kras*$^{V14lfloxed/wt}$ (from here termed as *Kras*^{V14l}) mice and their siblings (from here termed as control) were decapitated, the brains were removed, the forebrains were separated and freed of meninges. Dissociation of cells took place chemically in Papain-protease-DNase mix (HBSS-/-, 0.01% Papain reconstituted in EBSS, 0.01% (w:v) DNase I, 0.1% (w:v) Dispase II) at 37°C for ten minutes followed by mechanical dissociation. This process was repeated 2 times. Cell suspension was passed through a 70 µm cell strainer. Cells were spun down and washed with NBA mix three times, afterwards resuspended in NBA. Cells were counted and diluted to plate 80.000 cells in a volume of 50 µL onto the coated wells of the mwMEA plate. Incubating at 37°C in 5% CO_2 atmosphere for 1 h allowed cells to settle, followed by adding 250 µL NBA mix for further culturing. Similarly, 200.000 cells in a volume of 100 µl were carefully seeded onto coverslips for 1 h, immediately after removing the NBA mix containing FCS droplet from them. They were then transferred to 12-well plates containing 1 ml of NBA mix per well and maintained at 37°C in 5% CO_2 until maturation. Genotyping was done post hoc from tail biopsies obtained from euthanized pups.

Immunocytochemistry and image acquisition

All used antibodies are listed in Supplementary Material Table S 1. Immunofluorescence staining of vesicular GABA transporter (VGAT) and vesicular glutamate transporter (VGLUT1) were performed as double staining in combination with integral synaptic vesicle protein synaptophysin. Neuronal cultures grown on coverslips for 21 days were fixed with 4% (w:v) paraformaldehyde in PBS for 4 min at room temperature (RT). Fixed neurons were then washed in PBS and then blocked and permeabilized in tandem with PBS solution containing 10% (v:v) FCS, 0.1% (v:v) glycine and 0.3% (v:v) TritonX-100 for 45min at RT (Supplementary Material Table S 1). Coverslips were incubated with primary antibodies at 4 degrees overnight, and after washing with PBS with secondary antibodies at RT for 1 h. All antibodies were applied in PBS containing 3% (v:v) FCS. Coverslips were mounted on glass slides with Fluoroshield. All experimental conditions investigated in each experiment were processed in parallel with identical antibodies, solutions and other chemicals. Image acquisition was performed by an epifluorescence microscope (Nikon Eclipse Ti, Nikon Corporation) equipped with an iXon EM+ 885 EMCCD Andor camera (Andor Technology) using a 60X/NA 1.2 objective (Plan APO VC Nikon CFI, Nikon Corporation) and controlled by VisiView software (Visitron Systen GmbH).

MEA recordings

The wells in a 48-well MEA plates were equipped with 16 PEDOT electrodes arranged in a 4 x 4 grid with electrode spacing of 340 µm, electrode diameter of 50 µm and recording area of 1.1 mm x 1.1 mm. Recordings were performed using a Maestro multi-well MEA recorder (Axion Biosystems) at 37°C in 5% CO_2 : 95% air atmosphere. Before recording, neuronal cultures were visually checked for vitality under a light microscope (Supplement Material Figure S 1A). Prior to each recording during time series

experiments, the MEA plate was kept in the MEA recorder for at least 5 min to let the neuronal cultures adjust to ambient conditions and to avoid bias by mechanical disturbances. To acquire data and for spike detection, we used the Axion Integrated Studio (AxIS) software (2.4.2.13). The heat map visualization in the software enabled a first control of neuronal activity online (Supplement Material Figure S 1B). Voltage potentials were recorded across all electrodes with a sampling rate of 12.5 kHz (Supplement Material Figure S 1C). Recordings were subdivided into 5 minutes time units, to avoid a huge amount of data at once (.raw file). The raw signal was filtered by a Butterworth filter of 200 Hz to 2.5 Hz band pass. The filtered signal was detected for neuronal events when the signal exceeds a threshold denoted as spikes (cf. Ch. data analysis, Supplement Material Figure S 1D). These were stored in a spike list (.csv file), containing information about time and amplitude of detected spikes chronologically.

Time series of spontaneous network activity

Multiple recordings of spontaneous activity in neuronal networks plated on the same mwMEA plate were longitudinally and non-invasively conducted in the time range 10 days in vitro (DIV) to 33 DIV. On a measuring day, spontaneous activity was recorded for 20 min in 5 min time units.

Disinhibition-induced stimulation experiments

The application of bicuculline (bic) to *Ptpn11*[D61Y] neuronal cultures was performed on 33 DIV. The baseline was recorded for 20 min in 5 min time units, then the plate was taken out to add bic (volume: 10 µL, end concentration 10 µM, diluted in H_2O from a 10 mM stock solution in DMSO). After treatment, the plate was placed back to the MEA recorder and kept there for further 10 min without recording, followed by 50 min post bic recordings in 5 min time units. (Supplementary Material Figure S 2). Values were normalized to the baseline (100%) and presented as percentage.

Electrical stimulation experiments

Electrical stimulations were applied on mature networks (three independent experiments on DIV29, DIV30 and DIV39). To provide electrical test stimulation (STIM) and tetanus stimulation (TET), individual electrodes from the MEA were triggered to deliver electrical pulses (Supplementary Material Figure S 3A). Protocols for STIM and TET were modified from [9] (Supplementary Material Figure 3C). STIM protocol consists of a train of pulses (0.2 Hz). The pulses were biphasic from +750 mV to -750 mV lasting for 250 µs per phase. During STIM, six electrodes per array (stimulation electrodes e1-6 as shown in Supplementary Material Figure 3A) were triggered sequentially for 5 min, also denoted as sessions 1-6 (Supplementary Material Figure 3D). Prior to each STIM, consequently lasting 30 min in total, spontaneous activity was recorded initially for 20 min (prior STIM1) and subsequently for 10 min (prior STIM2 and STIM3). (Supplementary Material Figure 3D). TET was delivered from one electrode and consisted of 20 bursts at 0.2 Hz containing 11 pulses at an intra-burst frequency of 20 Hz (Supplementary Material Figure 3 C).

Data analysis

Immunocytochemistry

Identical camera and illumination settings were applied during image acquisition for coverslips of all experimental conditions imaged on the same day. Data was stored as 16-bit images and analyzed using the in-house MATLAB routine SynEval [16]. Herein, the pictures were segmented according to puncta labelled by immunofluorescent staining for synaptophysin (region of interest, ROI). Subsequently, within ROI, puncta were identified by being positive for VGAT or VGLUT, respectively. In brief, the segmentation algorithm follows an iterative, water-shed based approach and the identification of puncta positive in co-staining bases on immunofluorescence intensity (IFI) threshold individually calculated for each picture and then applied on the averaged IFI value derived from ROI encompassed pixels. Inter alia, the program returns the fraction and the IFI of positive puncta in co-staining.

Processing MEA-based data

For further data analysis, customized software packages, coded in MATLAB, processed the spike lists (.csv) that were output by AxIS software. Each analysis runs in a batch mode to save time and avoid human bias. The packages enable an analysis for each experimental setting including spontaneous network activity in time series recordings plus network disinhibition experiments as well as electrically evoked activity. We extended the spontaneous network activity analysis for time series experiments with MATLAB programmed functions performing PCA commonly applied in machine learning to reduce dimensionality in data sets.

Spontaneous network activity

The software package for analyzing spontaneous network activity in longitudinal experiments is equipped with a graphical user interphase (GUI) (Supplementary material Figure S 4) to enable the user-friendly input of experiment-related relevant data (cf. with scheme in Supplementary Material Figure S 5A). In the first step, the user selects the spike list files (.csv) that had been output by Axion software after each measurement. The minimum spike rate to determine active electrodes (herein 0.1 Hz,) and the minimum number of active electrodes (herein, unless otherwise stated: 10) can be specified in the GUI.

Spike Calculation: Spikes, that define neuronal firing events, are detected when the filtered voltage signal crosses a threshold in the continuous data stream. In this study, the threshold was adaptively set as a multiple of the median noise level (5.5 of standard deviation (SD)) on an electrode (Supplementary material Figure S 1C). The mean firing rate (MFR) is defined as the total number of spikes divided by recording time, $MFR = n_{spikes}/s$. The weighted MFR is calculated as $wMFR = (e_{max}/e_{act}) \cdot MFR$. Herein, the maximum number of active electrodes (e_{max}) was determined at the culture age of maturity (DIV 21-24). e_{act} referred to the currently active number of active electrodes. An electrode was defined as active when its spiking rate was at least 0.1 Hz and then it was also described as

contributing channel. Respectively for disinhibition experiments, e_{max} was determined as the number of active electrodes during the baseline measurement prior to treatment. Analogously, wtMFR (mean firing rate weighted to total (t) number of electrodes) indicates the same calculation, but e_{max} refers to the total number of electrodes (herein 16 electrodes). Interspike interval is the mean time between spikes in s.

Burst detection and calculation: A burst was defined as neuronal activity that appears in high frequency spiking followed by a time period of quiescence on a single electrode level. Herein, burst detection performed along an interval interspike threshold detecting bursts as spiking events of no longer than 100 ms intervals between 5 consecutive spikes. The burst duration is calculated as the time from the first spike to the last spike within a burst. The mean bursting rate (MBR) is the total number of bursts within one well divided by recording time. The value is weighted analogously to wMFR (wMBR = (e_{act}/ e_{max})·MBR).

Network analysis: The analysis of synchronized spiking in network bursts (NB) and correlated spiking measured as spike time tiling coefficient requires the transformation of the spike list to a time stamp matrix, a binary array of sample points indicating spikes as ones and sample points without spike as zeros. With a sampling rate of 12.5 kHz, one sample point correlates to 0.08 ms. The assessment of network-wide coordinated activity requires the detection of spike clusters across multiple electrodes. Based on the time stamp matrix, a MATLAB coded routine detected NBs across the array as spike clusters containing a minimum number of 50 spikes with a maximum of 100 ms interspike interval and with at least 5 contributing channels. Then calculation of several NB describing parameters such as NB duration, number of NB per well, and number of spikes per NB occurred. The NB duration was calculated as the time from the first spike to last spike in a NB. To investigate changes in the network dynamic, the correlation of spiking between each pair of electrodes was analyzed by the spike time tiling coefficient (STTC) [12] that reveals the synchronization between pairs of electrodes by returning the unbiased correlation in firing. It is calculated as

$$STTC = \frac{1}{2}(\frac{P_A - T_B}{1 - P_A T_B} + \frac{P_B - T_A}{1 - P_B T_A})$$

where A and B represents two spike trains. P_A (P_B) is the proportion of spikes from A (B) that occurs within the time window ±Δt (100 ms) around any spike from B (A). T_A (T_B)is the proportion of recording time that lies within ±Δt of any spike from A (B). The pairwise computation of STTC between electrodes resulted in a correlation matrix. For network description and illustration, we calculated the mean and the skewness of STTC across all electrode pairs.

In the second part of the package (right side in GUI, compare Supplementary material Figure S 4c, d), functions are included that are capable to automatically group data according to a grouping file (.xlsx) that is provided by the user via the GUI.

Principle component analysis for time series recordings

For PCA, we developed functions in MATLAB that use the well-level data files obtained and saved after running the first part of the routine. Depending on the analysis, these functions created parameter arrays consisting of either 19-dimensional parameter vectors (Supplementary Material Table S 2) in order to quantify cluster separation between RASopathy-model and wild-type using principle component 1 and 2 (PC1/2 analysis). Furthermore, PCA was performed to create vectors of specific parameters describing the features spiking, bursting and network bursting (Supplementary Material Figure S 5B) following up projection in PC1 with time. After PCA, MATLAB functions grouped well-level data automatically according to genotype or animal/preparation and time. For PC1/2 analysis, PCA score plots visualized genotype clusters in a PC1-PC2 two-dimensional coordinates system. To quantify the separation between the genotype clusters, we followed the suggested quantification method presented in [13]. In brief, we calculated the Mahalanobis distance [24] resulting in the distance between the centroids between groups considering groups' data distribution. The calculations were performed by custom-written MATLAB functions that additionally calculated statistical significance with F-statistics.

Electrical evoked activity in stimulation experiments

For an automatized analysis of data derived from MEA-based electrical stimulation of neuronal cultures, we developed a software package in MATLAB finally returning the network wide evoked spiking rate upon STIM normalized to the spontaneous activity ($evMFR_{norm}$) (Data processing illustrated in Supplementary Material Figure S 6). Based on the initial measurement of spontaneous activity (SA1), pre-selection of valid wells was performed based on the condition of minimum number of active electrodes per array (here: 8 electrodes) and with an electrode defined as active whenever its spiking rate exceeds a threshold (herein 0.1 Hz). To calculate $evMFR_{norm}$, firstly the evoked activity $evMFR^{'}$ was determined as the number of spikes within a time period of 1 s upon an electrical pulse divided by this time (Supplementary Material Figure S 3b), followed by averaging across all pulses on one electrode during one session. For each electrode, $evMFR^{'}$ was then normalized to the spontaneous activity (MFR'_{SA}) of the same electrode resulting in $evMFR'_{norm}$. To evaluate the network wide evoked activity, $evMFR'_{norm}$ was averaged across all array's electrodes and across all sessions within STIM yielding in $evMFR_{norm}$. The software package returned a data array containing $evMFR_{norm}$, as well as the corresponding coefficient of variance and the skewness for each well and STIM. Based on a user provided excel file, the package automatically grouped the wells according to genotype.

Data cleaning and selection of valid wells

To generate reliable results and to reduce the variability due to hardly predictable environment and physiological changes, criteria for selection of valid wells were defined. Firstly, wells were selected according to their activity as measured by the minimum number of active electrodes (if not otherwise stated: 10) determined by minimum spiking rate (herein > 0.1 Hz). For the time series experiments,

this selection was done based on measurements taken at time of network maturity that we denote here as reference time point (DIV 21-24). For disinhibition-induced stimulation experiments the baseline measurement at fourth time unit was taken as reference. Wells that did not fulfil the criteria were discarded from the analysis. We implied a further quality criteria and discarded wells once they lost their integrity beyond the reference time point. Loss of integrity was measured as a considerable drop in the number of active electrodes (no. of contributing channels < 80% of no. of contributing channels at reference time point) or as a clear decrease of neuronal activity (wMFR < 70% $wMFR_{ref}$).

For stimulation experiments being more prone to interferences due to a measurement duration above 2h, additional exclusion criteria were applied to ensure that solely stable cultures were included to analyses. First, cultures had to exhibit stability in their spontaneous activity prior (SA1) and after (SA2) STIM1 (Supplementary Material Figure S 3d). For each well, MFR and SD of firing rate were computed and significant changes between SA1 and SA2 were identified by Welch's t-test to discard wells that got significantly altered. Secondly, of particular importance for the evaluation of the effect of tetanic stimulation on the network evoked activity, we discarded cultures getting pathologically affected by STIM sessions. To evaluate this, we regarded the relation between $evMFR_{norm}$ upon STIM2 and upon STIM1 and checked for outliers (Supplementary Material Table S 4).

Statistics

For statistical calculations we used the GraphPad Prism (version 8.3.0) Software and MS Excel. The distribution of all data sets was tested by normality and lognormality tests (Supplementary Material Table S 4 and Figure S 7).

To test for statistical differences within time series experiments, subsequent PCA and disinhibition-induced stimulation experiments, we performed a mixed-effects analysis with GraphPad Prism. For time series experiments, we assume an increased risk of type two error caused by a systematic effect by time on the whole population potentially superimposing the effect of genotype and therefore performed either an unpaired t-test for normal distributed data or Mann-Whitney U tests for non-normal distributed data without correction for multiple comparisons. Respectively, the effect size is indicated as r^2 for unpaired student's t-test [22] or r for Mann-Whitney U test according to Wendt [41](Figure 1+2). Standard benchmarks for r fit well with our set of data: <.3 small effect, .3-.5 moderate effect, > .5 strong effect (r^2 respectively: <.09 small effect, .09-.25 moderate effect, > .25 strong effect). A p-value < .05 was considered significant.

Projected data set on PC1 was shown to be approximately normally distributed (Supplementary material Figure S 9) with equal variances in all groups (Supplementary material Figure S 10), leading us to perform mixed-effects model with Sidak's test to correct for multiple comparison for an increased statistical power. Since concomitant with PCA, normalization of data values centers the data points around zero for each time point, the systematic increase of neuronal activity coming along with

network development was filtered out in all groups. Thus, the effect caused by mutation on spiking, bursting and network bursting phenotype was unveiled decreasing the probability of error type two. As well, for disinhibition-induced stimulation experiments Sidak's multiple comparisons test was performed.

To compare the evoked activity upon STIM, we used the non-parametric Mann-Whitney test due to non-normal distribution of the data set for MFR. Additionally, we investigated differences in distribution by Kolmogorov-Smirnov test. To analyze the effect of tetanic stimulation on networks, we calculated the relation $X = \frac{evMFR_{log}^{STIM3}}{evMFR_{log}^{STIM2}}$ between the logarithmic $evMFR_{norm}$ after (STIM3) and prior (STIM2) the tetanic stimulation and check for alterations between genotypes by a Welch's t-test [40]. Therefore, we calculated descriptive measures from the independent, individual experiments first, followed by Welch's t-test according to

$$ t = \frac{\bar{X}_{floxed/WT} - \bar{X}_{WT/WT}}{\sqrt{\frac{S_{floxed/WT}^2}{n_{floxed/WT}^2} + \frac{S_{WT/WT}^2}{n_{WT/WT}^2}}} $$

The mean value $\bar{X}$ is the average of X across all experiments for the indicated genotype. S^2 is given by the averaged variance of indicated genotype derived from the individual experiment and n is the total number of all wells over all experiments for the indicated genotype. To check effect of tetanus on networks, we additionally performed one-sample Wilcoxon test on the pooled data set of X from all experiments.

Results

Increase in population-wide neuronal activity in neuronal networks from *Ptpn11*[D61Y]mice.

To identify the effect of RASopathy mutations on functional development of neuronal network, we performed longitudinal recordings of neuronal network activity in cultured dissociated cortical neurons grown on mwMEA-plates. The neurons were derived from cerebral cortices of newborn mice. We used mice being heterozygote for floxed RASopathy associated mutation *Ptpn11*[D61Y] or *Kras*[V14l] and for Emx1-cre allele as RASopathy model and their littermates being homozygote for *Ptpn11* or *Kras* wild type allele and heterozygote for Emx1-cre allele as control. The spontaneous network activity of cortical cultures was recorded multiple times in cultures in a time range of DIV 11 until DIV 33 (time series). For one experiment, cultures were prepared from six to eight newborn mice (P0-P1) littermates.

Fig. 1 Analysis of spontaneous spiking and bursting activity in neuronal networks from RASopathy models *Ptpn11*[D61Y] and *Kras*[V14I].

(a-b) Parameters describing neuronal activity as a function of culture age as box plots indicating the median and interquartile range with whiskers extending from 5% to 95% confidence interval. **a** wMFR and interspike interval. **b** Burst duration and wMBR. Parameters are statistically evaluated by mixed-effects model approach. Below the graphs, p and F values for main and intermediate effects are shown. If F-test reveals significance in genotype, p values from Mann-Whitney U test (black) and effective size r according to Wendt (grey, cursive) are shown in the graphs. Stars indicate significant differences between genotypes for a time point: < 0.05 (*), < 0.01 (**). A data point represents the mean values across the wells related to one animal/preparation (preparation-level).

To analyze the parameters describing spontaneous activity in neuronal networks, we averaged the data derived from individual wells across preparation/animal and subsequently grouped the datapoints according to genotype. To assess differences between groups and time points, mixed-effects model was calculated. Herein, we accepted the violation of the distributional assumption of equal standard deviation within groups due to the robustness of the mixed effects model (Supplementary data Fig. S 8). The global evaluation revealed that all groups reliably exhibited spontaneous activity on DIV 11-13. A gradual increase in population-wide firing activity (assessed as spiking and bursting, Fig. 1a, b) and in functional connectivity/network synchronicity (assessed as network bursting and correlation in spiking, Fig. 2b ,c) was evident between DIV 19 and DIV 25, when the all parameters got stabilized presumably marking the network maturity. Sample sizes are listed in Fig. 2a.

To assess the effect of RASopathy mutations on spiking and bursting behavior, we followed up four parameters including wMFR, interspike interval, burst duration and wMBR (Fig. 1). *Ptpn11*[D61Y] networks were significantly different in nearly all parameters compared to their control according to F-test in mixed effects model. The spiking activity was clearly increased as reflected in significantly higher values in wMFR nearly throughout all observation time points (exception on DIV 19/20) and in a decreased interspike interval reaching significance at the early points in time (DIV 11-13, DIV 15/16, DIV 19/20) (Fig. 1a, left column). Furthermore, *Ptpn11*[D61Y] networks showed an increased bursting behavior compared to their control (Fig. 1b, left column). The appearance of spontaneous bursts during development is important for the formation of neuronal circuits [20]. *Ptpn11*[D61Y] networks exhibited significantly higher wMBR on DIV 15/16, DIV 19/20 and DIV 30-33. However, the burst duration remained unchanged. By contrast, we detected no significant differences in any parameter describing spiking and bursting behavior in *Kras*[V14I]. However, visualization of data indicates a conspicuous trend of higher median values in parameters reflecting neuronal activity (wMFR, wMBR, burst duration) (Fig. 1a, b, right column).

A fundamental feature of neuronal networks is their functional connectivity/network synchronicity describing the impact of the activity of other neurons within the same network on the activity of given neuron [25]. To assess functional connectivity during network development, we calculated the number of network bursts (Fig. 2b) and determined the correlation between all pairs of electrodes during NBs measured as STTC (Fig. 2c). *Ptpn11*[D61Y] networks globally demonstrated significantly higher functional connectivity compared to their control groups revealed by mixed effects model in all observed parameters (Fig. 2b, c, left side). In *Ptpn11*[D61Y] networks, the number of NBs was increased

at early time points (DIV 11-13, DIV 15/16 and DIV 19/20), but this effect was abolished at later time points. Moreover, compared to controls, $Ptpn11^{D61Y}$ networks indicated significant differences in mean and skewness of STTC throughout the whole observation time as tested by multiple comparison tests (Figure 2C, left side). In contrast, compared to control group, $Kras^{V14I}$ showed no alterations in the number of NBs and in mean STTC (Fig. 2b, c, right side). However, a significant global difference was calculated for the skewness of STTC according to F-test on main effect genotype on DIV 27/28. In general, we identified elevated predicted median values in the $Kras^{V14I}$ networks in most parameters, however the differences were not significant likely due to large variability, which is typical to MEA-derived data.

a

animals / wells per culture age (DIV) from 4 independent experiments / litters

DIV	11-13	15/16	19/20	21-23	25-27	30-33	
Ptpn11^{D61Y}	7/34	7/34	7/34	9/44	8/37	8/36	
control	10/43	10/43	10/43	14/60	14/59	13/56	
DIV	10-12	15/16	17-19	21/22	24/25	27/28	30/31
Kras^{V14I}	12/62	8/47	8/45	12/54	12/47	8/29	8/23
control	12/61	9/50	9/50	12/61	12/58	9/39	8/30

Fig. 2 Analysis of functional connectivity/synchronicity in networks with RASopathy mutations *Ptpn11*[D61Y] and *Kras*[V14I].
a Number of animals and wells analyzed at each culture age cohort for all parameters of spontaneous activity. **(b-c)** Parameters describing functional connectivity as a function of culture age as box plots indicating the median and interquartile range with whiskers extending from 5% to 95% confidence interval **b** Values of no. of network bursts. **c** Values of mean STTC and skewness STTC. Parameters are statistically evaluated by mixed-effects model approach. Below the graph, p and F values for main and intermediate effects are shown. If F-test reveals significance in genotype, in **b** p values from Mann-Whitney U test (black) and effective size r according to Wendt (grey, cursive) are shown, in **c** p values from unpaired t-test together with effect size r^2 are shown in the graphs. Stars indicate significant differences between genotypes for one time point < 0.05 (*), < 0.01 (**), < 0.001 (***), < 0.0001 (****). A data point represents the mean values across the wells related to one animal/preparation (preparation-level).

Principle Component analysis of MEA parameters reveals differences in genotype clusters and identifies differences in spiking, bursting and network bursting in both RASopathy models

Analysis of most individual parameters from the longitudinal recordings in *Ptpn11*[D61Y] neurons revealed higher effect sizes in younger cultures that reduced at later time points. Similar effects were observed in *Kras*[V14I] neurons, even though not significant. To increase interpretability of datasets, power of statistical testing and to allow more simple visualization we introduced PCA, that reduces data dimensionality. We performed PCA including 19 parameters obtained from custom-written MATLAB program (Supplementary material Table S 2). We conducted the PCA on well-level. In contrast to preparation-level we yielded a higher number of data points, resulting in a lower probability of error type two but decreased power.

We analyzed all age cohorts from the longitudinal recordings and visualized the transformed data points in a PC1 and PC2 coordination system (Fig. 3a, Fig. 4a). Sample numbers are shown in Fig. 3d and Fig. 4d. In both RASopathy disease models, the respective clusters and their control appear to converge with culture age, also indicated by the visualized distance between the cluster's centroids in the graphs. To quantify the separation between genotype clusters, we calculated the Mahalanobis distance that accounts for the correlation of the variables [24]. Statistical significance was evaluated by a Hotelling's two-sample T^2 related to an F-value as introduced earlier [13]. For both RASopathy model networks, Mahalanobis distance to the control group turned out being strongly significant at the early stages, but dramatically decreased with culture age (Fig. 3b, Fig. 4b). At the end of the observation period, *Ptpn11*[D61Y] networks still differed significantly from their control group according to the F-statistic on DIV 30-33, while for Kras[V14I] the clusters converge more closely and did not differ significantly on DIV 27/28. The reliability of the results is underpinned by the cumulative fraction of variance of PC1 and PC2 explaining around 70% of the variation in both data sets (Fig. 3c, Fig. 4c). This approach confirmed a developmental phenotype in spontaneous network activity in both RASopathy disease models.

Fig. 3 Principle component analysis of *Ptpn11*[D61Y] 19-dimensional parameter vector.
a Principle component PC1 versus PC2 scores plot in dependence of culture age (DIV). A solid line is drawn between the centroids of control cluster (triangle) and mutant cluster (rectangle). Each color-coded data point represents a recording from one well. **b** Cluster separation in dependence of culture age was measured by Mahalanobis distance (green dotted line), F-value shows the statistical significance (black solid line) ($F_{critical} = 3.1$). **c** Fraction of variance explained by each PC summed over all PCs. Cumulative fraction of variance is indicated for each evaluated culture age. **d** Sample size is given per culture age as number of animals and wells.

To further identify at which level the spontaneous network activity is particularly affected in RASopathy models, we performed a further PCA that specifically combines parameters describing the features spiking, bursting and network bursting behavior, respectively (Supplement Material Fig. S 5b). Therefore, we formed parameter vectors from the extracted parameters from the spontaneous activity analysis. The values were logarithmized for non-normal distributed data. This resulted in a 4-dimensional vector for spiking ($\log_{10}$(wMFR), $\log_{10}$(wtMFR), $\log_{10}$(interspike interval), no. of contributing electrodes), a 3-dimensional vector for bursting ($\log_{10}$(wMBR), $\log_{10}$(burst duration), $\log_{10}$(no. spikes per burst)) and a 4-dimensional parameter vector for network bursting (no. of network bursts, mean NB duration, no. of spikes per NB, no. of contributing channels), respectively. The PCA has resulted in a component projection on PC1 for each age cohort (Fig. 5). We visualized the projected parameter vector on PC1 for each neuronal age cohort (Fig. 5a-c). We were able to identify a

changed phenotype in spiking, bursting and network bursting for both RASopathy models compared to wild type.

Fig. 4 Principle component analysis of *Kras*^{V14I} 19-dimensional parameter vector.
a Principle component PC1 versus PC2 scores plot in dependence of culture age (DIV). A solid line is drawn between the centroids of control cluster (triangle) and mutant cluster (rectangle). Each color-coded data point represents a recording from one well. **b** Cluster separation in dependence of culture age was measured by Mahalanobis distance (green dotted line) between the clusters of genotypes, F-value shows the statistical significance (black solid line) (F$_{critical}$ = 3.1). **c** Fraction of variance explained by each PC summed over all PCs. Cumulative fraction of variance is indicated for each evaluated culture age. **d** Sample size is given per culture age as number of animals and wells.

On the spiking level, the mixed effects model revealed a strong significance between the group mean in the main effect genotype (Fig. 5a). Thereby multiple comparisons testing showed that the strong differences were primarily at early time points (*Ptpn11*^{D61Y}: DIV 11-13, DIV 15/16, DIV 19/20; *Kras*^{V14I}: DIV 15/16, DIV 17-19, DIV 20-22). Similarly, for network bursting the mixed effects model indicated a significant difference in genotype with strong differences on DIV 11-13 and on DIV 15/16 in *Ptpn11*^{D61Y} and *Kras*^{V14I}, respectively (Fig. 5c). In contrary, as assessed by multiple comparisons testing, for bursting merely *Ptpn11*^{D61Y} showed significant differences at later time points (DIV 21-23, DIV 30-33), when networks are expected to be mature. Interestingly, mixed effects model identified strong interaction effect over time and between genotypes in all parameter vectors describing the network activity for both RASopathy models (genotype x time, indicated in all graphs in Fig 5).

Fig. 5 Principle component analysis (PCA) of parameter vectors related to spiking, bursting and network bursting derived from *Ptpn11*[D61Y] (left) and *Kras*[V14I] (right).

(a-c) PC1 projection as a function of culture age as box plots indicating the median and interquartile range with whiskers extending from 5% to 95% confidence interval. **a** Quantification of spiking behavior calculated as a projection of 4-dimensional parameter vector (no. of spiking electrodes, wMFR, wtMFR, interspike interval). **b** Quantification of bursting behavior calculated as projection of 3-dimensional parameter vector (wMBR, bursting duration, no. of spikes per bursts). **c** Quantification of network bursting calculated as a projection of 4-dimensional parameter vector (no. of network bursts (NB), mean NB duration, number of spikes per NB, mean no. of contributing channels per NB). Statistical significance was assessed by mixed-effects model approach. p-values (in black) for each comparison were obtained by Sidak's multiple comparisons test and effect size (r^2, in grey, cursive) were derived from the unpaired t-test. Stars indicate significant differences of median values between genotypes for time point: .05 (*), .01 (**), .001(***), <.0001(****).

Taken together, we particularly observed an alteration in spiking and network bursting for both RASopathy models during network development, but there were also effects on bursting caused by RASopathy mutation. Data also indicate that differences are more pronounced during early development and shades out during network maturation.

Pharmacological disinhibition has a specific effect on network bursting in *Ptpn11*[D61Y]

Our experiments revealed a change in neuronal activity and functional connectivity during network development in both RASopathy models. Interestingly, the effect of mutations on network characteristics is attenuated over time pointing to possible adaptation.

Fig. 6 Effect of disinhibition on neuronal network activity in *Ptpn11*[D61Y] on DIV 33.

a Number of animals (from 3 independent experiments) and wells used for this analysis. **b** Representative raster plots showing spatiotemporal spiking activity of control and *Ptpn11*[D61Y] networks prior and 30 min after bicuculline application (bic). (**c-d**) Line graphs of wMFR (**c**) and MNBR (**d**) as a function of time prior and after application of bic showing an elevated spiking upon treatment in both genotypes. Values are represented in percent related to a baseline generated from values recorded over a time interval of 20 min prior to bic treatment, averaged and set to 100 %. Significant increase against baseline was statistically tested by one sample Wilcoxon. In (**d**), interaction effect between time and genotype was calculated by mixed effects model approach. Sidak's multiple comparison tests determine significance between genotypes, Stars indicate significant differences < 0.05 (*).

To assess a possible contribution of shift in excitatory/inhibitory (E/I) balance, previously suggested in [28], we decided to test an effect of pharmacological interference with inhibitory tone on network activity. Therefore, we applied bic, a potent antagonist of $GABA_A$ receptor, to cultured cortical neurons (DIV 33) from *Ptpn11*[D61Y] and their wild type littermates.

As expected, the bic treatment resulted in a reliable increase in spiking activity reflected in increased wMFR (Fig 6b, c), statistically confirmed by one sample Wilcoxon test for each well (Figure 6c).

While spiking and bursting behavior did not differ between genotypes upon bic treatment (Fig. 6c, Supplementary Material Fig. S 11), we detected an elevated network bursting in disinhibited *Ptpn11*[D61Y] networks compared to control. Mixed effects model revealed a global interaction effect of time and genotype (genotype x time $p < .0001$), leading us to perform Sidak's multi-comparison tests resulting in significant differences from minute 50 post bic (Fig. 6c, d). For an overall conclusion of the main effect genotype, the mixed effects model returns a p-value of .06 that is near to significance. This result revealed changes in inhibitory system in *Ptpn11*[D61Y] networks, which likely developed to counteract increased spontaneous network activity.

Less GABAergic synapses, but higher expression of vesicular transporters for GABA and glutamate were detected in the networks with *Ptpn11*[D61Y] mutation

To examine the altered inhibitory tone on a synaptic level, we visualized and quantified glutamatergic and GABAergic boutons in *Ptpn11*[D61Y] networks and their control. Therefore, neuronal cultures were immunolabeled with specific antibodies against vesicular GABA transporter (VGAT) or vesicular glutamate transporter (VGLUT1), respectively, together with antibodies against synaptophysin (Syp) to visualize all synapses (Fig. 7a, b). We used our in-house developed MATLAB based program SynEval to obtain Syp positive puncta (ROI) and to sort them as VGAT or VGLUT-positive in an automatized batch mode, which reduces human bias and time effort. The fraction of glutamatergic boutons was significantly higher in *Ptpn11*[D61Y] compared to the control group (Fig.7d), and accordingly the fraction of GABAergic boutons decreased (Fig. 7c). Interestingly, we noticed an increased synaptic IFI for VGAT or VGLUT indicating increased synaptic abundance of these neurotransmitter transporters (Fig. 7c, d). These findings further point towards changes in excitation-inhibition balance in *Ptpn11*[D61Y] networks.

Fig. 7 Immunolabeling of excitatory and inhibitory synapses in neuronal networks from *Ptpn11*[D61Y].

a Representative images of neurons (DIV 21) labeled using antibodies against VGAT (green) and Syp (red). **b** Representative images of neurons labeled using antibodies against VGLUT1 (green) and synaptophysin (Syp,red). **c** Quantification of VGAT IFI in Syp-positive synaptic puncta and of the fraction of VGAT positive synapses. **d** Quantification of VGLUT1 IFI in Syp-positive synaptic puncta and of the fraction of VGLUT1 positive synapses. Statistical significance was determined by Mann-Whitney test. Data points correspond to individual analyzed visual fields derived from two independent experiments. The scale bar is 5 μM.

Altered evoked activity in *Ptpn11*[D61Y] probed by electrical stimulation

The longitudinal analysis indicated a normalization of spontaneous network activity during maturation in the neuronal networks expressing RASopathy mutation. To assess, whether there are functional defects in processing of information in these apparently "normal" networks we analyzed network activity evoked by delivery of electrical stimulation. We recorded network activity evoked by single-pulse stimulation (test stimulus, STIM) delivered from different electrodes in *Ptpn11*[D61Y] networks grown for 33 days (DIV 33) (Supplement Material Fig. S 3a, d). Of note, we solely analyzed robust and stable networks, which was not affected by the application of STIM itself, i.e. activity evoked by STIM1 was same as activity evoked by STIM2 (cf. Material and Methods, Data cleaning and selection of valid wells). The data points plotted in Fig. 8a display the evMFR$_{norm}$ representing the evoked MFR induced by STIM on electrode level and normalized to recorded spontaneous activity on the same electrode, averaged across an array (cf. for detailed information on calculation Supplementary Material Fig. S 6). Interestingly, evMFR$_{norm}$ was found to be significantly decreased in *Ptpn11*[D61Y] networks compared to wild type (Fig. 8a, upper plot). The graph is showing the data for STIM2, similar results were obtained for STIM1 (Supplementary Material Fig. S 12). Kolmogorov-Smirnov test confirmed a different distribution of data between both genotypes. The dampened electrical

excitability of *Ptpn11*[D61Y] networks is in line with our finding of affected inhibitory tone and the decrease in synaptic activity in *Ptpn11*[D61Y] shown in [3].

Fig. 8 Assessment of evoked network activity in *Ptpn11*[D61Y] neurons.

a Evoked mean firing rate (evMFR$_{normed}$) before and after TET stimulation. Black lines indicate median values, data point correspond to one well. Significance was tested by Mann-Whitney (MW) to check differences in mean values and Kolmogorov-Smirnov (KS) to check differences in data distributions. **b** Relative change (ratio) of evMFR$_{normed}$ recorded before and after application of TET are shown from three independent experiments (Exp1-Exp3), Whiskers indicate SEM. p-value was calculated by Welch t-test; number of included wells for corresponding genotype are color-coded for each experiment. **c** Analysis of data pooled from all 3 experiments. Black lines indicate median values, one data point refers to one well. One-sample Wilcoxon test (WT) tested against hypothetical value of 1 (indicating no change in evMFR$_{normed}$ caused by TET) and Mann-Whitney test (MW) to calculate for statistical difference between genotypes.

Effect of high frequency electrical stimulation on evoked activity in *Ptpn11*[D61Y]

We observed that evoked neuronal activity upon STIM is potentially affected by RASopathy associated mutations and excitability is dampened. As next, to understand if neuronal plasticity might also be affected in *Ptpn11*[D61Y] networks, we tested effect of tetanic stimulus (TET) on evoked synaptic activity. The stimulation protocol described in Supplementary Material Fig. S 3c was previously suggested to be suitable for analysis of neuronal plasticity since it affected evoked activity in networks [9]. The TET stimulus applied on *Ptpn11*[D61Y] networks induced different effect compared to controls in all in three independent experiments individually plotted in Fig. 8b. Calculating the relative change (ratio) between post- and pre-tetanus logarithmic (log$_{10}$)evMFR$_{norm}$ showed the mean value in *Ptpn11*[D61Y] being higher than in control group for each experiment (Exp1-Exp3) with an overall significant difference confirmed by Welch's t-test (Mann-Whitney test for individual experiments: Exp1: p = 0.03, Exp2: ns, Exp3: ns). More precisely, the findings reveal that TET resulted in a significant increase in evMFR$_{norm}$ in *Ptpn11*[D61Y] networks while not inducing an effect in controls (Fig.

8c). That was shown by testing the pooled data set against the hypothetical value 1 (no change between pre and post-tetanus logarithmic $evMFR_{norm}$) by a one sample Wilcoxon test. Interestingly, the difference in evMFR_norm, demonstrated between $Ptpn11^{D61Y}$ and control pre TET, has been diminished post TET (Fig. 8a, bottom plot). According to Kolmogorov-Smirnov test also the difference in distribution fail significance after TET. Indeed, in contrast to control, $Ptpn11^{D61Y}$ demonstrated a shift between STIM2 and STIM3 curves in their relative frequency plots, particularly for higher $evMFR_{norm}$, pointing to altered distribution in data set (Supplementary Material Fig. S 12b, c).

Discussion

RASopathies are a group of rare genetic disorders with various symptoms including heart defects, skeleton anomalies, increased risk for tumors and variable degree of neurocognitive deficits. Clinically, severity of neurocognitive impairments in RASopathies are often varying during life time, often getting milder with age of the patients. Neurodevelopmental phenotypes were also described in RASopathy mouse models, where involvement of maladaptive changes in cortical networks was suggested [28].

To investigate, whether similar maladaptation occur in neuronal networks on a chip, we cultured cortical neurons carrying two distinct RASopathy mutations on mwMEAs and recorded their population-wide spontaneous network activity to monitor network establishment and maturation during the time period of 5 weeks. To facilitate the evaluation of large datasets acquired longitudinally with high sampling rate (12.5 kHz), we developed and implemented an analysis pipeline programmed in MATLAB. From the list of spikes per electrode, provided by acquisition software, it computes a custom set of parameters describing various features of neuronal spontaneous network activity, including spiking, bursting, network bursting and correlation in spiking. This custom-written software package permits fast and unbiased analysis of the large MEA-datasets while being flexibly applicable for a wide set of experimental designs in general. The provision of a GUI renders the package user-friendly. Furthermore, we developed an add-on package of MATLAB based functions that implement PCA to reduce complexity of analysis outcomes by projecting the resulting parameters into a two-dimensional system. The package allows the calculation of Mahalanobis distance and statistical testing to mathematically compare the network behavior between two groups derived in our case from mutant and wild type animals and determine if these constitute significantly distinguishable populations. In the next step it also computes, whether the difference between networks clusters can be contributed to one specific network function feature (spiking, bursting and network synchronicity) or whether changes in multiple features contribute to the clusters' differences. This approach turned out to be suitable to uncover effects in the MEA data that is inherently characterized by high complexity and variability.

On the basis of MEA-recorded data and supported by the MATLAB analysis pipeline, we identified a convergent developmental phenotype in both used the RASopathy models. Multiple parameters describing neuronal activity and functional connectivity were significantly different in neurons with *Ptpn11*[D61Y] mutation. The differences were more pronounced in younger networks and faded out as networks matured. In contrast, the neurons with *Kras*[V14l] mutation showed only a tendency towards similar changes in spontaneous network activity parameters as seen in *Ptpn11*[D61Y] neurons. Being aware of the high inherent variance of MEA-data, we performed the PCA analysis to obtain an overall description of network parameters for both models. This analysis revealed a significant separation between clusters representing neurons with *Ptpn11*[D61Y] or *Kras*[V14l] mutation and their respective controls. Moreover, for both mutations the cluster separation from control networks decreased throughout development and consecutive analysis identified major contribution of changes in spiking and network bursting to the PCA cluster separation. Thus, based on the analysis of spontaneous network activity we have clearly demonstrated similar developmental phenotypes for both mutations confirming that the effect of RASopathy mutations on neuronal network function converges. It is interesting to note that mutations' effect on network function correlates with the strength of their activation assessed biochemically in previous studies. While *Kras*[V14l] exhibits an attenuated biochemical activation compared to common cancer-associated KRAS alleles [56], strong activation was shown for *Ptpn11*[D61Y], which is even linked to juvenile myelomonocytic leukemia (JMML) in patients [38].

The normalization of network activity parameters in mature networks suggest that compensatory mechanisms develop to counteract the increased network activity observed in juvenile networks. A shift in the E/I balance might represent one such mechanism. Indeed, the balance between E/I within neuronal circuits is crucial for correct brain development and function and its disturbances contribute to numerous neurodevelopmental disorders like Autism Spectrum Disorders and epilepsy which indeed are common comorbidities in more severe RASopathies including CFC or CS [1, 30]. Moreover, the recent characterization of RASopathy animal model expressing strongly activating allele *Kras*[G12V] indicated that increased inhibitory drive contributes to the electrophysiological and behavioral abnormalities in this model [28]. In line with these assumptions, we demonstrated an increased network bursting in mature *Ptpn11*[D61Y] networks upon disinhibition mediated by pharmacological block of GABA receptors. The immunocytochemical analysis revealed an increased proportion of excitatory and decreased proportion of inhibitory synapses in *Ptpn11*[D61Y] networks as well as an increased expression of vesicular transporters for both glutamate and GABA. In line with a modified inhibitory tone, we have found that the application of electrical test stimuli in a low frequency elicited a reduced evoked response in the *Ptpn11*[D61Y] networks pointing towards maladaptive dampening of network activity, which leads to an attenuated response to stimulation. An application of TET stimulation that had no effect on evoked responses in control had a significant impact on responses in neurons with *Ptpn11*[D61Y] mutation. Interestingly, the TET stimulation

statistically restored the affected evoked activity in the RASopathy model. However, the change in distribution of data values between pre- and post-tetanus stimulations indicates that there is much higher variability in responses of individual tested networks expressing $Ptpn11^{D61Y}$ mutations compared to controls. At this timepoint it's not clear, which concrete parameters are involved in the RASopathy induced network adaptation. A better understanding of the circumstances, under which plasticity can be induced can potentially be used in therapeutic approaches to break the maladaptive circuits and use them for targeted treatment of RASopathies and in their comorbidities.

In summary our study reveals that different RASopathy mutations have convergent effects of neuronal network activity. We observed more severe differences in network characteristic in juvenile networks that were compensated in the late stages of network maturation, which is in good agreement with clinical data. Despite of apparent normalization of spontaneous network activity, we uncovered significant differences in E/I balance and in evoked responses in mature networks. Last but not least the cells on-a-chip approach developed here is well scalable, accessible for treatments and manipulations and it is also adaptable for stem cell technology-derived humanized disease modelling. Therefore, we assume that the analysis workflow developed here could be implemented by drug screening and testing platforms also for unrelated brain diseases.

Availability of code and data

All custom written source code used for analyzing experiments and plotting is available on a GitHub repository. Code for analyzing spontaneous network activity by functional analysis of neuronal activity and connectivity can be found at [Analysis of spontaneous network activity](#), functions to perform PCA, Mahalanobis distance calculation and corresponding statistics at [Principle Component Analysis](#) and for analyzing electrically evoked activity at [Analysis of electrically evoked activity.](#)

Data was obtained according to the material section. The datasets during and/or analyzed during the current study available from the corresponding author on reasonable request.

Abbreviations

AxIS	Axion Integrated Studio
bic	bicuculline
CFC	cardio-facio-cutaneous syndrome
CS	costello syndrome
DIV	days in vitro
e$_{act}$	currently active number of active electrodes
e$_{max}$	maximum number of active electrodes
FCS	fetal calf serum
GUI	graphical user interphase
IFI	immunofluorescence intensity

MAPK	mitogen-activated protein kinase
MBR	mean bursting rate
MEA	multielectrode array
MFR	mean firing rate
mwMEA	multi-well MEA
NB	network burst
NBA	Neurobasal^{TM} -A media
NS	Noonan syndrome
PC	principle component
PCA	*Principle component analysis*
PCR	polymerase chain reaction
ROI	region of interest
RT	room temperature
SA	spontaneous activity
SD	standard deviation
STIM	test stimulation
STTC	spike time tiling coefficient
TET	tetanus stimulation
VGAT	vesicular GABA transporter
VGLUT1	vesicular glutamate transporter
wMBR	weighted mean bursting rate
wtMFR	mean firing rate weighted to total number of electrodes

References

1. Adviento, B., Corbin, I.L., Widjaja, F., Desachy, G., Enrique, N., et al. (2014) *Autism traits in the RASopathies.* J Med Genet. 51(1):p. 10-20.DOI:10.1136/jmedgenet-2013-101951.
2. Alfieri, P., Piccini, G., Caciolo, C., Perrino, F., Gambardella, M.L., et al. (2014) *Behavioral Profile in RASopathies.* American Journal of Medical Genetics Part A. 164(4):p. 934-942.DOI:10.1002/ajmg.a.36374.
3. Altmuller, F., Pothula, S., Annamneedi, A., Nakhaei-Rad, S., Montenegro-Venegas, C., et al. (2017) *Aberrant neuronal activity-induced signaling and gene expression in a mouse model of RASopathy.* PLoS Genet. 13(3):p. e1006684.DOI:10.1371/journal.pgen.1006684.
4. Anni, D., Weiss, E.M., Guhathakurta, D., Akdas, Y.E., Klueva, J., et al. (2021) *Abeta1-16 controls synaptic vesicle pools at excitatory synapses via cholinergic modulation of synapsin phosphorylation.* Cell Mol Life Sci. 78(11):p. 4973-4992.DOI:10.1007/s00018-021-03835-5.
5. Armour, C.M. and Allanson, J.E. (2008) *Further delineation of cardio-facio-cutaneous syndrome: clinical features of 38 individuals with proven mutations.* J Med Genet. 45(4):p. 249-54.DOI:10.1136/jmg.2007.054460.
6. Bang, S., Hwang, K.S., Jeong, S., Cho, I.J., Choi, N., et al. (2021) *Engineered neural circuits for modeling brain physiology and neuropathology.* Acta Biomater. 132:p. 379-400.DOI:10.1016/j.actbio.2021.06.024.
7. Cesarini, L., Alfieri, P., Pantaleoni, F., Vasta, I., Cerutti, M., et al. (2009) *Cognitive profile of disorders associated with dysregulation of the RAS/MAPK signaling cascade.* Am J Med Genet A. 149A(2):p. 140-6.DOI:10.1002/ajmg.a.32488.
8. Chan, G., Kalaitzidis, D., Usenko, T., Kutok, J.L., Yang, W., et al. (2009) *Leukemogenic Ptpn11 causes fatal myeloproliferative disorder via cell-autonomous effects on multiple stages of hematopoiesis.* Blood. 113(18):p. 4414-24.DOI:10.1182/blood-2008-10-182626.
9. Chiappalone, M., Massobrio, P.and Martinoia, S. (2008) *Network plasticity in cortical assemblies.* Eur J Neurosci. 28(1):p. 221-37.DOI:10.1111/j.1460-9568.2008.06259.x.
10. Costa, R.M., Federov, N.B., Kogan, J.H., Murphy, G.G., Stern, J., et al. (2002) *Mechanism for the learning deficits in a mouse model of neurofibromatosis type 1.* Nature. 415(6871):p. 526-30.DOI:10.1038/nature711.
11. Cui, Y., Costa, R.M., Murphy, G.G., Elgersma, Y., Zhu, Y., et al. (2008) *Neurofibromin regulation of ERK signaling modulates GABA release and learning.* Cell. 135(3):p. 549-60.DOI:10.1016/j.cell.2008.09.060.
12. Cutts, C.S. and Eglen, S.J. (2014) *Detecting pairwise correlations in spike trains: an objective comparison of methods and application to the study of retinal waves.* J Neurosci. 34(43):p. 14288-303.DOI:10.1523/JNEUROSCI.2767-14.2014.
13. Goodpaster, A.M. and Kennedy, M.A. (2011) *Quantification and statistical significance analysis of group separation in NMR-based metabonomics studies.* Chemometr Intell Lab Syst. 109(2):p. 162-170.DOI:10.1016/j.chemolab.2011.08.009.
14. Gorski, J.A., Talley, T., Qiu, M., Puelles, L., Rubenstein, J.L., et al. (2002) *Cortical excitatory neurons and glia, but not GABAergic neurons, are produced in the Emx1-expressing lineage.* J Neurosci. 22(15):p. 6309-14.DOI:20026564.
15. Gremer, L., Merbitz-Zahradnik, T., Dvorsky, R., Cirstea, I.C., Kratz, C.P., et al. (2011) *Germline KRAS mutations cause aberrant biochemical and physical properties leading to developmental disorders.* Hum Mutat. 32(1):p. 33-43.DOI:10.1002/humu.21377.
16. Guhathakurta, D., Akdas, E.Y., Fejtova, A.and Weiss, E.M. (2022) *Development and Application of Automatized Routines for Optical Analysis of Synaptic Activity Evoked by Chemical and Electrical Stimulation.* Front Bioinform. 2:p. 814081.DOI:10.3389/fbinf.2022.814081.
17. Hernandez-Porras, I., Fabbiano, S., Schuhmacher, A.J., Aicher, A., Canamero, M., et al. (2014) *K-RasV14I recapitulates Noonan syndrome in mice.* Proc Natl Acad Sci U S A. 111(46):p. 16395-400.DOI:10.1073/pnas.1418126111.

18.   Kang, M. and Lee, Y.S. (2019) *The impact of RASopathy-associated mutations on CNS development in mice and humans.* Mol Brain. 12(1):p. 96.DOI:10.1186/s13041-019-0517-5.
19.   Kim, Y.E. and Baek, S.T. (2019) *Neurodevelopmental Aspects of RASopathies.* Mol Cells. 42(6):p. 441-447.DOI:10.14348/molcells.2019.0037.
20.   Kirkby, L.A., Sack, G.S., Firl, A.and Feller, M.B. (2013) *A role for correlated spontaneous activity in the assembly of neural circuits.* Neuron. 80(5):p. 1129-44.DOI:10.1016/j.neuron.2013.10.030.
21.   Krab, L.C., Aarsen, F.K., de Goede-Bolder, A., Catsman-Berrevoets, C.E., Arts, W.F., et al. (2008) *Impact of neurofibromatosis type 1 on school performance.* J Child Neurol. 23(9):p. 1002-10.DOI:10.1177/0883073808316366.
22.   Lakens, D. (2013) *Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs.* Front Psychol. 4:p. 863.DOI:10.3389/fpsyg.2013.00863.
23.   Lee, Y.S., Ehninger, D., Zhou, M., Oh, J.Y., Kang, M., et al. (2014) *Mechanism and treatment for learning and memory deficits in mouse models of Noonan syndrome.* Nat Neurosci. 17(12):p. 1736-43.DOI:10.1038/nn.3863.
24.   Mahalanobis, P.C. *On the generalized distance in statistics*. 1936. National Institute of Science of India.
25.   Makarov, V.A., Panetsos, F.and de Feo, O. (2005) *A method for determining neural connectivity and inferring the underlying network dynamics using extracellular spike recordings.* J Neurosci Methods. 144(2):p. 265-79.DOI:10.1016/j.jneumeth.2004.11.013.
26.   Mautner, V.F., Kluwe, L., Thakker, S.D.and Leark, R.A. (2002) *Treatment of ADHD in neurofibromatosis type 1.* Dev Med Child Neurol. 44(3):p. 164-70.DOI:10.1017/s0012162201001876.
27.   Molina, J.R. and Adjei, A.A. (2006) *The Ras/Raf/MAPK pathway.* J Thorac Oncol. 1(1):p. 7-9.
28.   Papale, A., d'Isa, R., Menna, E., Cerovic, M., Solari, N., et al. (2017) *Severe Intellectual Disability and Enhanced Gamma-Aminobutyric Acidergic Synaptogenesis in a Novel Model of Rare RASopathies.* Biol Psychiatry. 81(3):p. 179-192.DOI:10.1016/j.biopsych.2016.06.016.
29.   Payne, J.M., Hyman, S.L., Shores, E.A.and North, K.N. (2011) *Assessment of executive function and attention in children with neurofibromatosis type 1: relationships between cognitive measures and real-world behavior.* Child Neuropsychol. 17(4):p. 313-29.DOI:10.1080/09297049.2010.542746.
30.   Pierpont, E.I., Kenney-Jung, D.L., Shanley, R., Zatkalik, A.L., Whitmarsh, A.E., et al. (2022) *Neurologic and neurodevelopmental complications in cardiofaciocutaneous syndrome are associated with genotype: A multinational cohort study.* Genet Med. 24(7):p. 1556-1566.DOI:10.1016/j.gim.2022.04.004.
31.   Pierpont, E.I., Tworog-Dube, E.and Roberts, A.E. (2015) *Attention skills and executive functioning in children with Noonan syndrome and their unaffected siblings.* Dev Med Child Neurol. 57(4):p. 385-92.DOI:10.1111/dmcn.12621.
32.   Roberts, A., Allanson, J., Jadico, S.K., Kavamura, M.I., Noonan, J., et al. (2006) *The cardiofaciocutaneous syndrome.* J Med Genet. 43(11):p. 833-42.DOI:10.1136/jmg.2006.042796.
33.   Roelofs, R.L., Janssen, N., Wingbermuhle, E., Kessels, R.P.and Egger, J.I. (2016) *Intellectual development in Noonan syndrome: a longitudinal study.* Brain Behav. 6(7):p. e00479.DOI:10.1002/brb3.479.
34.   Samuels, I.S., Saitta, S.C.and Landreth, G.E. (2009) *MAP'ing CNS development and cognition: an ERKsome process.* Neuron. 61(2):p. 160-7.DOI:10.1016/j.neuron.2009.01.001.
35.   Schubbert, S., Zenker, M., Rowe, S.L., Boll, S., Klein, C., et al. (2006) *Germline KRAS mutations cause Noonan syndrome.* Nat Genet. 38(3):p. 331-6.DOI:10.1038/ng1748.
36.   Schulz, A.L., Albrecht, B., Arici, C., van der Burgt, I., Buske, A., et al. (2008) *Mutation and phenotypic spectrum in patients with cardio-facio-cutaneous and Costello syndrome.* Clin Genet. 73(1):p. 62-70.DOI:10.1111/j.1399-0004.2007.00931.x.

37. Sharland, M., Burch, M., McKenna, W.M.and Paton, M.A. (1992) *A clinical study of Noonan syndrome.* Arch Dis Child. 67(2):p. 178-83.DOI:10.1136/adc.67.2.178.

38. Tajan, M., Paccoud, R., Branka, S., Edouard, T.and Yart, A. (2018) *The RASopathy Family: Consequences of Germline Activation of the RAS/MAPK Pathway.* Endocr Rev. 39(5):p. 676-700.DOI:10.1210/er.2017-00232.

39. Tartaglia, M., Zampino, G.and Gelb, B.D. (2010) *Noonan syndrome: clinical aspects and molecular pathogenesis.* Mol Syndromol. 1(1):p. 2-26.DOI:10.1159/000276766.

40. Welch, B.L. (1947) *The Generalization of 'stundent's' problem when several different pupulation varlances are involved* Biometrika. 34(1-2):p. 28-35.DOI:10.1093/biomet/34.1-2.28.

41. Wendt, H.W. (1972) *Dealing with a common problem in Social science: A simplified rank-biserial coefficient of correlation based on the U statistic.* European Journal of Social Psychology. 2(4):p. 463-465.DOI:https://doi.org/10.1002/ejsp.2420020412.

42. Yoon, S. and Seger, R. (2006) *The extracellular signal-regulated kinase: multiple substrates regulate diverse cellular functions.* Growth Factors. 24(1):p. 21-44.DOI:10.1080/02699050500284218.

43. Zenker, M. and Kutsche, K. (2016) *RASopathies.* Medizinische Genetik. 28(1):p. 15-36.DOI:10.1007/s11825-016-0080-8.