Title: Inhibitory tagging both speeds and strengthens saccade target selection in the

superior colliculus during visual search

Author names and affiliations: Christopher Conroy[1], Rakesh Nanjappa[1,2], Robert M.

McPeek[1]

[1]Department of Biological and Vision Sciences, SUNY College of Optometry, New York,

NY, 10036, USA

[2]Department of Optometry, School of Medical & Allied Sciences, G.D. Goenka

University, Gurugram, 122103, India

Corresponding author contact: cconroy@sunyopt.edu

Conflict of interest statement: The authors declare no conflicts of interest.

Acknowledgements: This work was funded by National Institutes of Health (NIH)-

National Eye Institute (NEI) grant number R01-EY030669

ABSTRACT

It has been suggested that, during difficult visual-search tasks involving time pressure

and multiple saccades, inhibitory tagging helps to facilitate efficient saccade target

selection by inhibiting saccades to objects in the scene once they have been searched

and rejected. The superior colliculus (SC) is a midbrain structure that is involved in

target selection, and recent findings suggest an influence of inhibitory tagging on activity

in the SC. Precisely how, and by how much, inhibitory tagging affects target selection by

SC neurons, however, is unclear. The purpose of this study, therefore, was to

characterize and quantify the influence of inhibitory tagging on target selection in the

SC. Rhesus monkeys performed a relatively difficult visual-search task involving time

pressure and multiple saccades. During the task, target selection by SC neurons was

both more rapid and more robust—as quantified via shifts in selection metrics derived

from temporal receiver operating characteristic analyses—when an uninspected

saccade-target object was selected from among inspected, as opposed to uninspected,

objects in the display. This suggests that inhibitory tagging both speeds and strengthens

saccade target selection at the level of individual SC neurons. The results provide

constraints on models of target selection based on SC activity.

NEW AND NOTEWORTHY

An important aspect of efficient behavior during difficult, time-limited visual search is the efficient selection of sequential saccade targets. Recent findings suggest that inhibitory tagging may help to facilitate such efficient selection by modulating activity in the superior colliculus (SC). Here, we characterize and quantify the influence of this modulation and find that inhibitory tagging both speeds and strengthens saccade target selection at the level of individual SC neurons.

INTRODUCTION

Our daily lives are replete with examples of visual-search tasks that we need to

complete efficiently. Under time pressure, we execute saccade sequences, interleaved

with brief fixations, as a means to search the visual scene for a particular object of

interest. In such contexts, an important aspect of efficiency is the selection of

behaviorally relevant saccade targets, meaning objects in the visual scene that could

plausibly be the particular object of interest. It has been suggested that inhibitory

tagging may help to facilitate such efficient selection by inhibiting saccades to irrelevant

objects once they have been searched and rejected (1-7).

In a recent study, we reported evidence of an influence of inhibitory tagging on

the activity of individual neurons in the superior colliculus (SC) (8), a midbrain structure

that is involved in saccade target selection (9-26). More specifically, we reported

evidence that, during a difficult visual-search task involving time pressure and multiple

saccades, the visually evoked activity of many SC neurons was lower when the objects

in their response fields (RFs) had been previously fixated relative to when they had not.

The largest effects of inhibitory tagging, moreover, were observed in the subset of SC

neurons that had response properties consistent with a role in saccade target selection,

suggesting that a modulation of the selection process at the level of individual selection-

relevant neurons may provide the mechanism by which inhibitory tagging supports

efficient search behavior. Precisely how, and by how much, inhibitory tagging affects

target selection in the SC, however, is unclear, as we did not examine, much less

quantify, the selection process in our recent study of inhibitory tagging.

The purpose of this study, therefore, was to extend our previous work and to

characterize and quantify the influence of inhibitory tagging on target selection by SC

neurons. As an approach, we used receiver operating characteristic (ROC) analyses

based on signal detection theory (27), which have been used in numerous previous

studies of visual perception, target selection, and, in particular, target selection in the

SC (11-14, 18, 20, 21, 23, 26), to quantify the time course of target selection, as well as

the strength of selection signals, in individual SC neurons both before and after

inhibitory tagging. We reasoned that, if a modulation of the selection process at the level

of individual SC neurons provides the mechanism by which inhibitory tagging supports

efficient search behavior, inhibitory tagging should affect the dynamics of the selection

process and/or the strength of selection signals carried by the subset of SC neurons

that are involved in target selection. As will be seen, this is what we found.

METHODS

Details of the behavioral and physiological procedures can be found in our recent report

(8). Briefly, we used standard methods to record the activity of individual SC neurons in

two male rhesus monkeys (*Macaca mulatta*). While we recorded the activity of each

neuron, the monkey performed two behavioral tasks for liquid reward: a delayed-

saccade task and a visual-search task, in that order. Experimental protocols were

approved by the Institutional Animal Care and Use Committee at SUNY College of

Optometry and complied with the guidelines of the US Public Health Service policy on

Humane Care and Use of Laboratory Animals.

At the beginning of each trial of the delayed-saccade task, a red or green disc

appeared at the center of the monitor. The monkey was required to fixate this central

disc for 450-650 ms, at which point a red or green target disc appeared at a peripheral

location. Following the appearance of the target disc, the monkey was required to fixate

the central disc for an additional 500-700 ms, at which point the central disc

disappeared. The monkey was rewarded for making a saccade to the target disc within

70-400 ms. All discs subtended 0.5-2.0° of visual angle and had luminances of 12

cd/m^2. The luminance of the monitor's background was 0.3 cd/m^2. The color of the

target disc always differed from that of the central disc, was randomly selected at the

beginning of each recording session, and was fixed throughout each recording session.

The location of the target disc was systematically varied to determine the RF of the

neuron understudy. The location that yielded the highest level of visual and/or motor

activity was taken as the RF center.

At the beginning of each trial of the visual-search task, a red or green disc

appeared at the center of the monitor. The monkey was required to fixate this central

disc for 500 ms, at which point the search array appeared. The array consisted of

multiple red and green discs positioned at the intersection points of an invisible grid

centered on the central disc. One (or occasionally more than one) of the discs in the

array (never the central disc) held a reward, meaning that, when fixated, a liquid reward

was administered (immediately) to the monkey. The color of the reward-bearing disc(s)

always differed from the color of the central disc and, within a session, was the same as

the color of the target disc in the delayed-saccade task. Following the appearance of the

array, the monkey had eight seconds to locate the reward-bearing disc(s). If the monkey

fixated all reward-bearing discs during this eight-second window (for trials on which

there was more than one reward-bearing disc, a reward was administered only once

upon fixation of each disc), the trial ended. Otherwise, the monkey did not receive a

reward. For each neuron, the array was configured such that when the monkey fixated

one of the discs, a nearby disc was at the center of the RF. For most neurons, this

positioning scheme permitted the use of a five-by-five rectangular array, consisting of

nine potentially reward-bearing and 16 not potentially reward-bearing discs. For some

neurons with large RF eccentricities, the array consisted of fewer discs. For these

neurons, attempts were made to maintain the same ratio of potentially reward-bearing

to not-potentially reward-bearing discs. On each trial, the locations of the potentially

reward-bearing and not-potentially reward-bearing discs were randomly selected. The

sizes and luminances of the discs as well as the luminance of the monitor's background

were the same as in the delayed-saccade task.

Data analysis

As noted in the Introduction, the purpose of this study was to examine the influence of

inhibitory tagging on target selection by SC neurons. As such, we focused our analyses

on neurons that exhibited significant delay-period ("prelude" or "buildup") activity in the

delayed-saccade task, because SC neurons that are involved in saccade target

selection tend to exhibit this type of activity (9, 10, 11-14, 20, 23, 26, 27). To determine

if a particular neuron exhibited significant prelude activity in the delayed-saccade task,

we compared its activity during a 100-ms window beginning 300-ms after the onset of

the target (i.e., its delay-period activity) to its activity during a baseline 100-ms window

beginning 100-ms before the onset of the target. Only trials on which the target was

positioned at the RF center were considered. If the delay-period activity was

significantly greater than the baseline activity (Wilcoxon rank-sum test, α = 0.05), the

neuron was categorized as having significant prelude activity and was included in

subsequent analyses.

For the search task, a fixation was defined as the period between subsequent

saccades. The activity of each neuron was analyzed separately for three categories of

fixation, defined based on the fixation status of the disc at the center of the RF

[previously fixated (PF) or not previously fixated (NPF) on the same trial] and whether or

not it was selected as the target of the subsequent saccade: (1) fixations on which the

RF disc had not been previously fixated and was selected as the target of the

subsequent saccade (NPF-in-RF, saccade-towards fixations); (2) fixations on which the

RF disc had not been previously fixated and was not selected as the target of the

subsequent saccade (NPF-in-RF, saccade-away fixations); and (3) fixations on which

the RF disc had been previously fixated and was not selected as the target of the

subsequent saccade (PF-in-RF, saccade-away fixations). In addition to the requirement

of significant prelude activity, a neuron was only included in further analyses if its

activity was recorded on more than five fixations for each of these three categories. The

initial fixation on each trial did not count towards this criterion and, moreover, this

fixation was not included in any analysis. Schematic illustrations of the search task, the

search array, a typical saccade sequence, and the three categories of fixation are

shown in Fig. 1.

FIG. 1: Schematic illustrations of the search task, the search array, a typical saccade sequence, and the three categories of fixation. **A.** Each trial of the search task began with the appearance of a disc positioned at the center of the display, which the monkey was required to fixate for 500 ms. **B.** The search array then appeared on the screen. At fixation, the response field (RF) of the neuron under study, illustrated here and in panel A as a dotted gray circle, overlapped a nearby disc. One of the discs (and sometimes more than one of the discs) in the array held a reward. In the trial illustrated in this

figure, let us assume that the disc in the upper-right corner of the array held the reward.

The monkey's task was to search the array and fixate the reward-bearing disc. **C.**

Because the reward-bearing disc was only indicated by its color, which it shared with

many other discs in the array, fixation of the reward-bearing disc typically entailed a

sequence of multiple saccades. This panel schematizes a situation where fixation of the

reward-bearing disc was preceded by six saccades. Each arrow represents a saccade

vector. Our analyses focused on three categories of fixation: **D.** fixations on which the

disc in the RF had not been previously fixated on the same trial [i.e., not previously

fixated (NPF)] and was selected as the target of the subsequent saccade (sac) (NPF-in-

RF, sac towards), **E.** fixations on which the disc in the RF had not been previously

fixated on the same trial and was not selected as the target of the subsequent saccade

(NPF-in-RF, sac away), and **F.** fixations on which the disc in the RF had been

previously fixated (PF) on the same trial and was not selected as the target of the

subsequent saccade (PF-in-RF, sac away). In **D-F**, the arrow gives the vector of the

subsequent saccade, whereas the black and green circles give the RF during fixation.

Note that, in the actual experiment, the discs in the array were red and green (illustrated

here as different shades of gray), and the color of the central disc served to cue the

monkey to the color of the reward-bearing disc.

We recorded the activity of 113 neurons across the two monkeys. Of these, 65

met our inclusion criteria. For each neuron, spike-density functions were created by

convolving individual spikes with a Gaussian kernel (σ = 10 ms) and two ROC analyses

were performed. The first compared the activity on NPF-in-RF, saccade-towards

fixations to the activity on NPF-in-RF, saccade-away fixations. We refer to this as the

NPF-re-NPF analysis. The other compared the activity on NPF-in-RF, saccade-towards

fixations to the activity on PF-in-RF, saccade-away fixations. We refer to this as the

NPF-re-PF analysis. Note that the only difference between the two analyses was the

fixation status of the RF disc on saccade-away fixations and so any differences in the

results can be attributed to inhibitory tagging. For both analyses, activity was aligned on

both fixation onset and offset. An ROC curve was then computed at each ms by

comparing the distribution of activity (across fixations) on NPF-in-RF, saccade-towards

fixations to the distribution of activity on saccade-away fixations (NPF-in-RF or PF-in-

RF, depending on the analysis). An ROC area was computed at each ms by integrating

the area under the ROC curve. ROC areas were bounded between 0.0 and 1.0 and

provided an estimate of the degree to which the two distributions overlapped. An ROC

area of 0.5 indicated complete overlap, areas closer to 1.0 indicated greater levels of

activity on saccade-towards fixations, and areas closer to 0.0 indicated greater levels of

activity on saccade-away fixations. Thus, areas greater than 0.5 were suggestive of

saccade target selection, whereas those equal to or less than 0.5 were not.

For both ROC analyses and alignments, we obtained two selection metrics:

discrimination time and magnitude. Discrimination time was defined as the earliest time

at which saccade-towards activity was significantly greater than saccade-away activity

and stayed that way for a particular duration. Significance was determined by

performing a permutation test (2,000 iterations) on the ROC curve (11, 12, 18, 23, 29)

and was assessed at each ms beginning 50 ms before and ending 150 ms after fixation

onset for the fixation-onset alignment or beginning 150 ms before and ending with

fixation offset for the fixation-offset alignment. For the fixation-onset alignment,

discrimination time was taken as the first time point within the analysis window at which

the permutation test indicated that the ROC area was significantly different from 0.5 ($P <$

.05), the mean saccade-towards activity was greater than the mean saccade-away

activity, and both of these conditions were met for at least 20 consecutive ms. For the

fixation-offset alignment, discrimination time was taken as the first time point within the

analysis window at which the permutation test indicated that the ROC area was

significantly different from 0.5 and the mean saccade-towards activity was greater than

the mean saccade-away activity for all remaining time points until the onset of the

saccade. For each analysis and alignment, discrimination magnitude was taken simply

as the maximum (max) ROC area within the analysis window. Statistical comparisons of

discrimination times and magnitudes were accomplished using Wilcoxon signed-rank

tests ($\alpha = 0.05$).

RESULTS

The results for an example neuron are shown in Fig. 2. Focusing first on the activity

associated with RF objects that had not yet been searched nor rejected, it can be seen

that, for the example neuron, activity was higher on fixations on which the object at the

center of the RF was (Fig. 2A, solid-black trace), versus was not (Fig. 2A, dotted-black

trace), selected as the target of the subsequent saccade. The difference emerged

shortly following fixation onset (the fixation-onset-aligned discrimination time for the

NPF-re-NPF analysis was 53 ms relative to fixation onset; Fig. 2B, left column, vertical

dotted black line), well before fixation offset (the fixation-offset-aligned discrimination

time for the NPF-re-NPF analysis was -112 ms relative to fixation offset; Fig. 2B, right

column, vertical dotted black line), and persisted throughout the full fixation duration.

The discrimination magnitude was large as well (the fixation-onset- and -offset-aligned

max-ROC areas for the NPF-re-NPF analysis were 0.76 and 0.83, respectively; Fig. 2B,

solid-black traces). Thus, the pattern of activity exhibited by the example neuron was

consistent with the notion that this neuron played a role in saccade target selection, at

least when both the target and nontarget objects had not yet been searched nor

rejected.

FIG. 2: Activity and ROC area functions for an example SC neuron. **A.** Mean fixation-

onset- (left panel) and -offset- (right panel) -aligned spike-density functions. Different

traces show activity for different categories of fixation: the solid black trace shows the

activity on NPF-in-RF saccade-towards fixations, the dotted black trace shows the

activity on NPF-in-RF saccade-away fixations, and the dotted green trace shows the

activity on PF-in-RF saccade-away fixations. Each trace represents the mean activity

across all fixations in the relevant category, and the shaded region represents +/-1

standard error of the mean across fixations. **B.** Fixation-onset- (left panel) and -offset-

(right panel) -aligned ROC area functions for the same neuron. The black trace shows

the results of the NPF-re-NPF analysis, whereas the green trace shows the results of

the NPF-re-PF analysis. The vertical dotted black and green lines show the

discrimination (discrim) times for the NPF-re-NPF and NPF-re-PF ROC analyses,

respectively.

Moreover, the visually evoked activity of the example neuron was affected by the

placement of inhibitory tags (i.e., this neuron exhibited lower levels of activity on

saccade-away fixations on which there was a previously fixated, rather than not

previously fixated, object in its RF; Fig. 2A dotted green versus dotted-black traces; cf.

Conroy et al., 2023), and this influence, in turn, affected the dynamics and strength of

saccade target selection. That is, for the example neuron, target selection was both

more rapid and more robust (i.e., discrimination times were shorter and discrimination

magnitudes were larger) when an uninspected saccade-target object (i.e., a not

previously fixated object) was selected from among inspected (i.e., searched and

rejected) as opposed to uninspected objects in the display. As noted above, for the

fixation-onset-aligned NPF-re-NPF analysis, the discrimination time was 53 ms relative

to fixation onset, and the discrimination magnitude was 0.76. For the same alignment,

the discrimination time decreased to 22 ms and the discrimination magnitude increased

to 0.84 for the NPF-re-PF analysis. A decrease in discrimination time (from -112 ms to -

123 ms for the NPF-re-NPF to NF-re-PF analysis) and increase in discrimination

magnitude (from 0.83 to 0.89 for the NPF-re-NPF to NF-re-PF analysis) occurred for the

fixation-offset-aligned analysis as well.

A decrease in discrimination times and increase in discrimination magnitudes

after inhibitory tagging was evident for many of the neurons in our sample. Figure 3

shows the fixation-onset-aligned target-selection metrics for the 51 (of 65) neurons that

selected the saccade target during the fixation-onset-aligned analysis window (i.e., that

yielded a fixation-onset-aligned discrimination time for both the NPF-re-NPF and NPF-

re-PF analyses). For these neurons and alignment, the median discrimination time was

significantly shorter for the NPF-re-PF than NPF-re-NPF analysis (33 versus 49 ms

relative to fixation onset; $Z = -3.953$; $P < .001$; Fig. 3A-B). Indeed, for the NPF-re-PF

analysis, many neurons exhibited very early discrimination times, with a substantial

number selecting the saccade target even before fixation onset and/or the visual-

afferent delay. Such predictive saccade target selection has been reported previously

by SC neurons (27) and, in the case of the NPF-re-PF analysis, is consistent with a

remapping of inhibitory-tagging across saccades (cf. Refs 8, 30, 31). Note that, in our

analysis, the earliest-possible discrimination time was -50 ms relative to fixation onset

because that was the earliest time point at which an ROC curve was computed. As

such, inhibitory tagging may have an even greater influence on the dynamics of target

selection than is evident from Fig. 3.

For the same subset of 51 neurons, the median discrimination magnitude for the

fixation-onset alignment was higher for the NPF-re-PF than NPF-re-NPF analysis (0.79

versus 0.73; $Z = 5.521$; $P < .001$; Fig. 3C-D), and, for the 48 neurons that selected the

saccade target during the fixation-offset-aligned analysis window, we observed the

same basic effect of inhibitory tagging on the dynamics and strength of saccade target

selection: the median discrimination time was shorter (-113 versus -97 ms relative to

fixation offset; Z = -2.962; P = .003; Fig. 4A-B) and the median discrimination magnitude

was larger (0.83 versus 0.79; Z = 4.636 P < .001; Fig. 4C-D) for the NPF-re-PF than

NPF-re-NPF analysis. Moreover, the color of the RF disc—meaning, its color in the

context of the task: potentially reward bearing or not—did not influence this general

pattern of results. To confirm this, we performed the same ROC analyses on a restricted

dataset including only fixations on which the RF object was potentially reward bearing

on that trial [i.e., had the same color as the reward-bearing disc(s)] and found that, after

inhibitory tagging, there was a decrease in discrimination times and increase in

discrimination magnitudes for both the fixation-onset and -offset alignments. For the 45

neurons from this restricted dataset that selected the saccade target during the fixation-

onset-aligned analysis window discrimination times decreased from 62 ms to 27 ms

relative to fixation onset (Z = -4.632; P < .001) and discrimination magnitudes increased

from 0.74 to 0.80 (Z = 4.984; P < .001) for the NPF-re-NPF to NPF-re-PF analysis. For

the 41 neurons that selected the saccade target during the fixation-offset-aligned

analysis window, discrimination times decreased from -70 ms to -97 ms relative to

fixation offset (Z = -3.630; P < .001) and discrimination magnitudes increased from 0.81

to 0.84 (Z = 4.970; P < .001). We also performed the ROC analyses on a restricted

dataset including only fixations on which the RF object was not potentially reward

bearing and found a significant decrease in discrimination times (from 59 to 45 ms

relative to fixation onset for the NPF-re-NPF relative to the NPF-re-PF analysis; Z = -

2.245; P = .025) and increase in discrimination magnitudes (from 0.75 to 0.81 for the

NPF-re-NPF relative to the NPF-re-PF analysis; Z = 4.128; P < .001) for the 35 neurons

that selected the saccade target during the fixation-onset-aligned analysis window, a

significant increase in discrimination magnitudes (from 0.80 to 0.85 for the NPF-re-NPF

to NPF-re-PF analysis; *Z* = 3.259; *P* = .001) for the 35 neurons that selected the

saccade target during the fixation-offset-aligned analysis window, and clear trend

towards a decrease in discrimination times (from -74 to -91 relative to fixation offset for

the NPF-re-NPF relative to NPF-re-PF analysis; *Z* = -1.936; *P* = .053) for the same 35

neurons. Thus, taken together, the results suggest that inhibitory tagging both speeds

and strengthens saccade target selection at the level of individual SC neurons.

FIG. 3: Summary of the fixation-onset-aligned target-selection metrics (discrimination

times and magnitudes) for the 51 of 65 neurons in our sample that selected the saccade

target within the fixation-onset-aligned analysis window (-50-150 ms relative to fixation

onset) for both the NPF-re-NPF and NPF-re-PF analyses. **A.** Distributions of

discrimination times for the two ROC analyses. The black and green bars give the

distributions for the NPF-re-NPF and NPF-re-PF analyses, respectively. Note that -50

ms relative to fixation onset was the earliest possible discrimination time given the

analysis window. **B.** Relationship between NPF-re-PF and NPF-re-NPF discrimination

times for individual neurons (individual symbols). **C.** Distributions of discrimination

magnitudes for the NPF-re-NPF (black bars) and NPF-re-PF (green bars) analyses. **D.**

Relationship between NPF-re-PF and NPF-re-NPF discrimination magnitudes for

individual neurons (individual symbols).

FIG. 4: Summary of the fixation-offset-aligned target-selection metrics for the 48 of 65

neurons in our sample that discriminated the saccade target within the fixation-offset-

aligned analysis window (-150-0 ms relative to fixation offset) for both the NPF-re-NPF

and NPF-re-PF analyses. Otherwise, what is shown in each panel corresponds to what

is shown in Fig. 3 and the plotting conventions are the same. Note that, for the fixation-

offset-aligned discrimination times, however, the earliest possible discrimination time

was -150 ms.

DISCUSSION

Saccade target selection involves a competition between representations of potential

saccade plans on the SC visuomotor map. An inhibitory-tagging mechanism that seeks

to facilitate efficient selection would be expected to affect this competitive process by

reducing the activity associated with particular saccade plans—specifically, those that

would move the eyes towards objects in the scene that have already been searched

and rejected—and increasing the relative level of activity associated with others—

specifically, those that would move the eyes towards objects in the scene that have not

yet been searched nor rejected. Previously, we showed this basic pattern of

modulations after inhibitory tagging in the SC (8), and, here, we characterized and

quantified the influence of these modulations on target selection by SC neurons. Using

ROC analyses, We found that inhibitory tagging both speeds and strengthens saccade

target selection at the level of individual SC neurons, yielding significantly shorter target-

nontarget discrimination times and significantly larger target-nontarget discrimination

magnitudes in the subset of SC neurons that were likely involved in saccade target

selection. Thus, we conclude that modulations of the selection process at the level of

individual SC neurons may provide the mechanism by which inhibitory tagging supports

efficient search behavior. Models of target selection based on SC (or, more generally,

neural-priority-map) activity (e.g., Ref 32) may wish to take these modulations into

account to more accurately capture the dynamics of the selection process during

difficult, time-limited search.

REFERENCES

Adeli, H., Vitu, F., & Zelinsky, G. J. (2017). A model of the superior colliculus predicts

fixation locations during scene viewing and visual search. Journal of

Neuroscience, 37(6), 1453-1467.

Basso, M. A., & Wurtz, R. H. (1998). Modulation of neuronal activity in superior

colliculus by changes in target probability. Journal of Neuroscience, 18(18),

7519-7534.

Carello, C. D., & Krauzlis, R. J. (2004). Manipulating intent: evidence for a causal role of

the superior colliculus in target selection. Neuron, 43(4), 575-583.

Conroy, C., Nanjappa, R., & McPeek, R.M. (2023). Inhibitory tagging in the superior

colliculus during visual search. Journal of Neurophysiology.

doi:10.1152/jn.00095.2023

Green, D.M., & Swets, J.A. (1966). Signal Detection Theory and Psychophysics. New

York: Wiley.

Glimcher, P. W., & Sparks, D. L. (1992). Movement selection in advance of action in the

superior colliculus. Nature, 355(6360), 542-545.

Horwitz, G. D., & Newsome, W. T. (1999). Separate signals for target selection and

movement specification in the superior colliculus. Science, 284(5417), 1158-

1161.

Horwitz, G. D., & Newsome, W. T. (2001). Target selection for saccadic eye

movements: prelude activity in the superior colliculus during a direction-

discrimination task. Journal of Neurophysiology, 86(5), 2543-2558.

Kim, B., & Basso, M. A. (2008). Saccade target selection in the superior colliculus: a

signal detection theory approach. Journal of Neuroscience, 28(12), 2991-3007.

Kim, B., & Basso, M. A. (2010). A probabilistic strategy for understanding action

selection. Journal of Neuroscience, 30(6), 2340-2355.

Krauzlis, R. J., & Dill, N. (2002). Neural correlates of target choice for pursuit and

saccades in the primate superior colliculus. Neuron, 35(2), 355-363.

Klein, R. (1988). Inhibitory tagging system facilitates visual search. Nature, 334(6181),

430-431.

Klein, R. M., & MacInnes, W. J. (1999). Inhibition of return is a foraging facilitator in

visual search. Psychological Science, 10(4), 346-352.

McPeek, R. M., & Keller, E. L. (2002). Saccade target selection in the superior colliculus

during a visual search task. Journal of Neurophysiology, 88(4), 2019-2034.

McPeek, R. M., & Keller, E. L. (2004). Deficits in saccade target selection after

inactivation of superior colliculus. Nature Neuroscience, 7(7), 757-763.

Mirpour, K., Arcizet, F., Ong, W. S., & Bisley, J. W. (2009). Been there, seen that: a

neural mechanism for performing efficient visual search. Journal of

neurophysiology, 102(6), 3481-3491.

Mirpour, K., Bolandnazar, Z., & Bisley, J. W. (2019). Neurons in FEF keep track of items

that have been previously fixated in free viewing visual search. Journal of

Neuroscience, 39(11), 2114-2124.

Müller, H. J., & Mühlenen, A. V. (2000). Probing distractor inhibition in visual search:

inhibition of return. Journal of Experimental Psychology: Human Perception and

Performance, 26(5), 1591.

Ogawa, H., Takeda, Y., & Yagi, A. (2002). Inhibitory tagging on randomly moving

objects. Psychological Science, 13(2), 125-129.

Peterson, M. S., Kramer, A. F., Wang, R. F., Irwin, D. E., & McCarley, J. S. (2001).

Visual search has memory. Psychological Science, 12(4), 287-292.

Port, N. L., & Wurtz, R. H. (2009). Target selection and saccade generation in monkey

superior colliculus. Experimental brain research, 192, 465-477.

Posner, M. I., & Cohen, Y. (1984). Components of visual orienting. In H. Bouma and D.

Bowhuis (Eds.), Attention and Performance (pp. 531-556). New Jersey: Erlbaum.

Shen, K., & Paré, M. (2007). Neuronal activity in superior colliculus signals both

stimulus identity and saccade goals during visual conjunction search. Journal of

Vision, 7(5), 15-15.

Shen, K., Valero, J., Day, G. S., & Paré, M. (2011). Investigating the role of the superior

colliculus in active vision with the visual search paradigm. European Journal of

Neuroscience, 33(11), 2003-2016.

Shen, K., & Paré, M. (2014). Predictive saccade target selection in superior colliculus

during visual search. Journal of Neuroscience, 34(16), 5640-5648.

Takeda, Y., & Yagi, A. (2000). Inhibitory tagging in visual search can be found if search

stimuli remain visible. Perception & Psychophysics, 62(5), 927-934.