Title: Inhibitory tagging both speeds and strengthens saccade target selection in the

superior colliculus during visual search

Running head: Inhibitory tagging speeds and strengthens target selection

Author names and affiliations: Christopher Conroy[1], Rakesh Nanjappa[1,2], Robert M.

McPeek[1]

[1]Department of Biological and Vision Sciences, SUNY College of Optometry, New York,

NY, 10036, USA

[2]Department of Optometry, School of Medical & Allied Sciences, G.D. Goenka

University, Gurugram, 122103, India

Corresponding author contact: cconroy@sunyopt.edu

Conflict of interest statement: The authors declare no conflicts of interest.

Acknowledgements: This work was funded by National Institutes of Health (NIH)-

National Eye Institute (NEI) grant number R01-EY030669

ABSTRACT

It has been suggested that, during difficult visual search tasks involving time pressure

and multiple saccades, inhibitory tagging helps to facilitate efficient saccade target

selection by reducing responses to objects in the scene once they have been searched

and rejected. The superior colliculus (SC) is a midbrain structure involved in target

selection, and recent findings suggest an influence of inhibitory tagging on SC activity.

Precisely how, and by how much, inhibitory tagging influences target selection by SC

neurons, however, is unclear. The purpose of this study, therefore, was to characterize

and quantify the influence of inhibitory tagging on target selection in the SC. Rhesus

monkeys performed a visual search task involving time pressure and multiple saccades.

Early in the fixation period between saccades, a subset of SC neurons reliably

discriminated the stimulus selected as the next saccade goal, consistent with a role in

target selection. Discrimination occurred earlier and was more robust, however, when

unselected stimuli in the search array had been previously fixated on the same trial.

This indicates that inhibitory tagging both speeds and strengthens saccade target

selection in the SC during multisaccade search. The results provide constraints on

models of target selection based on SC activity.

NEW AND NOTEWORTHY

An important aspect of efficient behavior during difficult, time-limited visual search tasks

is the efficient selection of sequential saccade targets. Inhibitory tagging, i.e., a

reduction of neural activity associated with previously fixated objects, may help to

facilitate such efficient selection by modulating the selection process in the superior

colliculus (SC). In this study, we characterized and quantified this modulation and found

that, indeed, inhibitory tagging both speeds and strengthens target selection in the SC.

INTRODUCTION

Our daily lives are replete with examples of visual search tasks that we need to

complete efficiently. Under time pressure, we execute saccade sequences, interleaved

with brief fixations, to search the visual scene for objects of interest. Performance in

such contexts depends critically on efficient and appropriate selection of sequential

saccade targets. Inhibitory tagging, i.e., a reduction of neural activity associated with

previously fixated objects, may help to facilitate such efficient selection by increasing

the speed and strength with which uninspected objects can be discriminated from

previously fixated objects in the visual scene (1-6).

In a recent study, we reported evidence of an influence of inhibitory tagging on

activity in the superior colliculus (SC) (6), a midbrain structure that is involved in

saccade target selection (7-25). Specifically, we found that, during a time-limited,

multisaccade visual search task, the activity of many SC neurons was lower when the

stimuli in their response fields (RFs) had been previously fixated compared to when

they had not. Moreover, the largest effects of inhibitory tagging were observed in the

subset of SC neurons that had response properties consistent with a role in saccade

target selection. This suggests that modulation of the selection process at the level of

individual SC neurons may provide a mechanism by which inhibitory tagging supports

efficient search behavior. Precisely how, and by how much, inhibitory tagging influences

target selection in the SC, however, is unclear, as we did not examine nor quantify the

influence of inhibitory tagging on target selection in our previous study.

The purpose of this study, therefore, was to extend our previous work and to

characterize and quantify the influence of inhibitory tagging on target selection in the

SC. As an approach, we analyzed the ability of SC neurons to discriminate between

selected and unselected RF stimuli during the brief fixation period between saccades in

the multisaccade search task and compared the time course and strength of

discrimination with and without inhibitory tagging. More specifically, we first analyzed

the ability of SC neurons to discriminate between RF stimuli that were selected as the

saccade target and unselected stimuli in the absence of inhibitory tagging, i.e., when

neither the selected nor unselected stimuli had been previously fixated on the same

trial. We then compared the time course and strength of discrimination in this condition

with results obtained when the unselected stimuli had been tagged with inhibition due to

previous fixation on the same trial. We reasoned that if inhibitory tagging supports

efficient search behavior by modulating the selection process in the SC, we should

observe more rapid and robust discrimination when the unselected stimuli had been

tagged with inhibition. As will be seen, this is what we found, indicating that inhibitory

tagging both speeds and strengthens saccade target selection in the SC.

METHODS

Experimental protocols were approved by the Institutional Animal Care and Use

Committee at SUNY College of Optometry and complied with the guidelines of the US

Public Health Service policy on Humane Care and Use of Laboratory Animals. A

detailed description of the surgical, recording, and behavioral procedures can be found

in our recent report (6). Briefly, we used microelectrodes and standard methods to

record the activity of well-isolated single SC neurons in two male rhesus monkeys

(*Macaca mulatta*). During recording, monkeys performed two behavioral tasks for liquid

reward: a delayed-saccade task and a multisaccade search task. Eye movements were

measured using a video-based eye tracker (Eyelink 1000, SR Research).

The delayed-saccade task was used to determine the RF center for each neuron

and for neuronal classification. At the beginning of each trial of this task, a red or green

disc appeared at the center of a stimulus-display monitor. Monkeys were required to

fixate this central disc for 450-650 ms, at which point a red or green target disc

appeared at a peripheral location. Following the appearance of the target disc, monkeys

were required to maintain fixation for an additional 500-700 ms, at which point the

central disc disappeared. Monkeys were rewarded for making a saccade to the target

disc within 70-400 ms of the central disc's disappearance. All discs subtended 0.5-2.0°

of visual angle [M-scaled based on eccentricity to keep salience approximately constant

(cf. Ref. 26)] and had a luminance of 12 cd/m^2 on a 0.3 cd/m^2 background. The color of

the target disc always differed from that of the central disc, was randomly selected at

the beginning of each recording session and was fixed throughout each session.

Initially, the location of the target was manually varied to home in the neuron's RF

center, defined as the location yielding the largest visual and/or motor response.

Following identification of the RF center, the activity of each neuron was recorded in at

least ten trials of the delayed-saccade task with the target positioned at the RF center.

At the beginning of each trial of the search task, a red or green disc appeared at

the center of the monitor. Monkeys were required to fixate this disc for 500 ms, at which

point a search array appeared. The array consisted of multiple red and green discs

positioned at the intersection points of an invisible grid centered on the central disc. All

discs had the same size and luminance as the discs presented to the RF center in the

delayed-saccade task. One of the discs in the array (never the central disc) held a

reward, meaning that, when fixated, a liquid reward was immediately administered to

the monkey. The color of the reward-bearing disc always differed from the color of the

central disc and, within a session, remained constant from trial to trial. Thus, if the

central disc was green, the reward-bearing disc was red, and vice versa. No other

colors were tested. Each trial lasted for eight seconds or until the reward-bearing disc

had been fixated, whichever came first. For each neuron, the array was configured such

that when the monkey fixated one of the discs, a nearby disc was positioned at the RF

center. For most neurons, this positioning scheme permitted the use of a five-by-five

rectangular array and, for most of these neurons, the array consisted of nine potentially

reward bearing discs. For some neurons, typically with large RF eccentricities, the array

consisted of fewer discs, in which cases attempts were made to maintain approximately

the same ratio of potentially reward-bearing to not-potentially reward-bearing discs as in

the five-by-five rectangular array case (e.g., for one neuron with an RF center at 15°,

the array consisted of 17 discs, six of which were potentially reward bearing). The

locations of potentially reward-bearing and not-potentially reward-bearing discs within

the array were randomly selected on each trial. On a small proportion of all trials

(roughly 7%), all "potentially" reward-bearing discs were in fact reward bearing,

meaning that the monkey received a reward upon the first (and only the first) fixation of

each disc that had a color that differed from that of the central disc. This trial type was

used to increase the monkey's motivation. These trials, like the single-reward trials,

lasted for eight seconds or until all reward-bearing discs had been fixated, whichever

came first.

Data analysis

As noted in the Introduction, the purpose of this study was to examine the influence of

inhibitory tagging on target selection in the SC. Thus, we focused our analyses on

neurons exhibiting significant delay-period activity in the delayed-saccade task because

SC neurons involved in saccade target selection tend to exhibit this type of activity (cf.

7-12, 18, 21, 24, 25). To measure delay-period activity, we compared each neuron's

activity during a 100-ms window beginning 300-ms after the onset of the target to its

baseline activity during a 100-ms window beginning 100-ms before target onset. Only

trials in which the target was positioned at the RF center were considered. If delay-

period activity was significantly greater than baseline activity (Wilcoxon rank-sum test, α

= 0.05), the neuron was included in subsequent analyses.

For the search task, the activity of each neuron was analyzed during fixations

between sequential saccades. Activity was analyzed separately for three categories of

fixation, which differed in terms of the fixation status of the RF disc [previously fixated

(PF) or not previously fixated (NPF) in the same trial] and whether the RF disc was

selected as the next saccade goal: (1) fixations during which an NPF disc in the RF was

selected as the next saccade goal (NPF-in-RF, saccade-towards); (2) fixations during

which an NPF disc in the RF was not selected as the next saccade goal (NPF-in-RF,

saccade-away); and (3) fixations during which a PF disc in the RF was not selected as

the next saccade goal (PF-in-RF, saccade-away). A neuron was only included in the

analyses if its activity was recorded during more than five fixations for each category.

The initial fixation in each trial did not count towards this criterion and, moreover, was

not included in any analysis. Schematic illustrations of the search task, the search array,

an example saccade sequence, and the three fixation categories are shown in Fig. 1.

FIG. 1: Schematic illustrations of the search task. **A.** Trials began with a disc at display

center which monkeys fixated. **B.** The search array then appeared on the screen. At

fixation, a nearby disc was positioned at the center of the RF of the neuron under study

(dotted white circle). One disc differing in color from the central disc triggered a liquid

reward when fixated. The monkey's task was to fixate this reward-bearing disc. **C.**

Fixation of the reward-bearing disc was typically preceded by a sequence of multiple

saccades to other discs (see Ref. 6 for details on search behavior). This panel

illustrates a situation where fixation of the reward-bearing disc (in this example, the disc

in the upper right corner) was preceded by a sequence of six saccades, represented by

white arrows. Analyses focused on three fixation categories: **D.** fixations during which

an NPF disc in the RF was selected as the next saccade goal (NPF-in-RF, saccade-

towards), **E.** fixations during which an NPF disc in the RF was not selected as the next

saccade goal (NPF-in-RF, saccade-away), and **F.** fixations during which a PF disc in the

RF was not selected as the next saccade goal (PF-in-RF, saccade-away). In **D-F**, the

arrow illustrates the vector of the next saccade and the dotted-white circle illustrates the

RF center.

We recorded 113 neurons from two monkeys. Of these, 65 met our inclusion

criteria. For each neuron, spike-density functions were created by convolving individual

spikes with a Gaussian kernel (σ = 10 ms). We then applied two receiver operating

characteristic (ROC) analyses (27) to the activity of each neuron as a way to measure

the effects of inhibitory tagging on the target-selection process. Similar ROC analyses

have been used in previous studies of target selection in the SC and other areas (e.g.,

Refs. 9-12, 16, 18, 19, 21, 24, 25, 28-30). The first analysis measured the

discriminability of selected and unselected RF discs in the absence of inhibitory tagging

by comparing activity during NPF-in-RF, saccade-towards fixations to the activity during

NPF-in-RF, saccade-away fixations. We refer to this as the NPF-re-NPF analysis. The

other analysis measured how inhibitory tagging of unselected discs affects target

selection in the SC by comparing activity during NPF-in-RF, saccade-towards fixations

to activity during PF-in-RF, saccade-away fixations. We refer to this as the NPF-re-PF

analysis. We used the same NPF-in-RF, saccade-towards activity for both analyses

because, as noted in the Introduction, we were interested in understanding how

inhibitory tagging supports efficient search behavior by modulating SC target selection.

In this task, efficient search requires selection of NPF discs. Thus, if inhibitory tagging

supports efficient search behavior by modulating the selection process in the SC, it

should facilitate efficient selection of NPF discs in particular.

For both ROC analyses, activity was aligned on both fixation onset and the onset

of the subsequent saccade (fixation offset). For each alignment, an ROC curve was

computed at each ms by comparing the distributions of activity for selected and

unselected disc, and area under the ROC curve was computed. An ROC area of 0.5

indicates complete overlap between selected and unselected distributions, areas closer

to 1.0 indicate greater activity for selected discs, and areas closer to 0.0 indicate greater

activity for unselected discs. We analyzed both fixation- and saccade-onset-aligned

activity to accurately measure the time course of target discrimination relative to both

fixation and saccade onset, as has been standard in previous studies (e.g., 9-11, 16,

18, 24, 25, 28-30). Statistical significance was assessed using a permutation test (2,000

iterations) on ROC area (cf. 9, 10, 16, 21, 30) at each ms beginning 50 ms before and

ending 150 ms after fixation onset for the fixation-onset-aligned analysis and beginning

150 ms before and ending with saccade onset for the saccade-onset-aligned analysis.

For all ROC analyses, we obtained two selection metrics for each neuron:

discrimination time and magnitude (e.g., 18, 24, 25, 29). For the fixation-onset-aligned

analysis, discrimination time was defined as the earliest time point within the analysis

window at which ROC area was significantly different from 0.5, the mean selected

activity was greater than the mean unselected activity, and both of these conditions

were met for at least 20 consecutive ms. For the saccade-onset alignment,

discrimination time was defined as the earliest time within the analysis window at which

ROC area was significantly different from 0.5, the mean selected activity was greater

than the mean unselected activity, and both of these conditions were met at all time

points until saccade onset. Discrimination magnitude was defined as the maximum

ROC area within each analysis window. Thus, the discrimination-time metric provided

an estimate of the earliest time point at which each neuron reliably discriminated

selected from unselected discs in its RF and therefore an estimate of the time of the

completion of the selection process. We predicted that if inhibitory tagging supports

efficient search behavior by modulating the selection process in the SC, discrimination

of an NPF target disc should be faster when that target is embedded in PF discs (NPF-

re-PF analysis) than when it is embedded in NPF discs (NPF-re-NPF analysis). In other

words, we anticipated shorter discrimination times for the NPF-re-PF than NPF-re-NPF

analysis. Discrimination magnitude provided an estimate of the discriminability of

selected and unselected discs following completion of the selection process. We

predicted that if inhibitory tagging strengthens selection signals in the SC, it should

increase discriminability of selected and unselected discs when the unselected discs

have been previously fixated. Thus, we anticipated larger discrimination magnitudes for

the NPF-re-PF than NPF-re-NPF analysis. Statistical comparisons of selection metrics

were accomplished using Wilcoxon signed-rank tests (α = 0.05).

RESULTS

Many of the neurons in our sample exhibited activity during the search task consistent

with a role in saccade target selection, and, for many of these neurons, the time course

of the selection process, as well as the strength of selection signals, was influenced by

inhibitory tagging. This is illustrated in Fig. 2, which shows the results (activity and ROC

area versus time) for an example neuron. We focus first on the activity associated with

NPF RF discs and the results of the NPF-re-NPF analysis, which, again, characterized

the selection process without inhibitory tagging. Relatively early in the fixation period,

this neuron reliably discriminated selected from unselected NPF discs. The fixation- and

saccade-onset-aligned discrimination times for the NPF-re-NPF analysis were 53 ms

relative to fixation onset and -112 ms relative to saccade onset, respectively. Activity

associated with selected discs steadily increased and eventually culminated in a

saccadic burst time locked to the next saccade, whereas activity associated with

unselected discs was initially inhibited and then held at bay. The consequence of these

two activity patterns was robust selected-unselected discriminability (i.e., NPF-re-NPF

ROC areas > 0.5) throughout most of the fixation period following the initial phase of

discrimination and, in particular, around the time of saccade onset. Discrimination

magnitudes for the NPF-re-NPF analysis aligned on fixation and saccade onset, which,

it can be seen, reflected ROC areas around the time of saccade onset, were 0.76 and

0.83, respectively. Thus, before inhibitory tagging, this neuron reliably and robustly

discriminated selected from unselected NPF discs, consistent with a role in saccade

target selection.

FIG. 2: Activity and ROC area functions for an example neuron. **A.** Mean fixation- (left

panel) and saccade-onset- (right panel) -aligned spike-density functions. Different

traces show activity for different fixation categories: the solid black trace shows the

activity during NPF-in-RF, saccade-towards fixations, the dotted black trace shows the

activity during NPF-in-RF, saccade-away fixations, and the dotted green trace shows

the activity during PF-in-RF, saccade-away fixations. Each trace represents the mean

activity and the shaded region represents +/-1 standard error of the mean. In the left

column, the inset histogram shows the distribution of saccade-onset times for the NPF-

in-RF saccade-towards fixations. In the right column, it shows the distribution of fixation

onset times for the same fixation category. The y-axis in each panel gives the fixation

counts for the distributions [i.e., the numbers on the y axis give both spikes per second

(for the spike-density functions) and fixation counts (for the fixation-onset time

distribution)]. The downward pointing triangles show the means of the distributions. A

small number of fixations were longer than 250 ms and are not shown for visualization

purposes. **B.** Fixation- (left panel) and saccade-onset- (right panel) -aligned ROC area

functions for the same neuron. The black trace shows the results of the NPF-re-NPF

analysis, the green trace shows the results of the NPF-re-PF analysis, and the vertical

dotted lines show the discrimination times for the two analyses. The downward pointing

triangles replot the mean saccade onset (left column) and onset (right column) times for

NPF-in-RF, saccade-towards fixations from Fig. 2A.

Discrimination of selected NPF discs occurred earlier and was more robust,

however, following the placement of inhibitory tags on unselected discs in the search

array, as indicated by the decrease in discrimination times and increase in

discrimination magnitudes for the NPF-re-PF relative to the NPF-re-NPF analysis. For

the fixation-onset-aligned analysis, the discrimination time decreased from 53 ms for the

NPF-re-NPF condition to 22 ms for the NPF-re-PF condition, i.e., when the unselected

stimulus had been tagged with inhibition. Furthermore the discrimination magnitude

increased from 0.76 to 0.84 going from the NPF-re-NPF to NPF-re-PF analysis. A

decrease in discrimination time (from -112 ms to -123 ms for the NPF-re-NPF to NF-re-

PF analysis) and increase in discrimination magnitude (from 0.83 to 0.89 for the NPF-

re-NPF to NF-re-PF analysis) occurred for the saccade-onset-aligned analysis as well.

The discrimination magnitudes for the NPF-re-PF analysis also reflected ROC areas

around the time of saccade onset, and so the larger discrimination magnitude for the

NPF-re-PF than NPF-re-NPF analysis indicates that inhibitory tagging of unselected

discs was strong even around the time of saccade onset, potentially reducing the

probability that activity associated with unselected discs would disrupt or interfere with

the selected saccade (31, 32).

A decrease in discrimination times and increase in discrimination magnitudes

after inhibitory tagging was evident for many of the neurons in our sample. Figure 3

shows the fixation-onset-aligned target-selection metrics for the 51 of 65 neurons in our

sample that discriminated selected from unselected RF discs during the fixation-onset-

aligned analysis window (i.e., neurons that yielded a fixation-onset-aligned

discrimination time for both the NPF-re-NPF and NPF-re-PF analyses). For these

neurons and alignment, the median discrimination time was significantly shorter for the

NPF-re-PF than NPF-re-NPF analysis (33 versus 49 ms relative to fixation onset; Z = -

3.953; P < .001; Fig. 3A-B). Indeed, for the NPF-re-PF analysis, many neurons

exhibited very early discrimination times, with a substantial number discriminating

selected from unselected discs even before fixation onset and/or the visual-afferent

delay. Specifically, for the NPF-re-PF fixation-onset-aligned analysis, 19 of the 51

neurons that discriminated selected from unselected discs did so before fixation onset.

An additional 15 did so within the first 50 ms following fixation onset. Such early

discrimination times are consistent with the idea that the discrimination process began

before fixation onset (i.e., during the planning or execution of the previous saccade; cf.

Refs. 25, 33) and that there was a predictive remapping of inhibitory-tagging signals

preceding the onset of each saccade (cf. Refs 5, 6). Thus, inhibitory tagging may not

only facilitate the selection process during individual fixations in multisaccade

sequences but also may facilitate the efficient planning of multisaccade sequences as a

whole.

For the same subset of 51 neurons that discriminated selected from unselected

discs during the fixation-onset-aligned analysis window, the median discrimination

magnitude was higher for the NPF-re-PF than NPF-re-NPF analysis (0.79 versus 0.73;

$Z = 5.521$; $P < .001$; Fig. 3C-D). For the 48 neurons that discriminated selected from

unselected RF discs during the saccade-onset-aligned analysis window, we observed a

similar pattern: the median discrimination time was shorter (-113 versus -97 ms relative

to saccade onset; $Z = -2.962$; $P = .003$; Fig. 4A-B) and the median discrimination

magnitude was larger (0.83 versus 0.79; $Z = 4.636$ $P < .001$; Fig. 4C-D) for the NPF-re-

PF than NPF-re-NPF analysis. Thus, we conclude that inhibitory tagging both speeds

and strengthens saccade target selection at the level of individual SC neurons.

Additional analyses that bolster this conclusion are reported in the supplementary

material. Also in the supplementary material, we report analyses showing a relationship

between behavioral search efficiency during the SC recording sessions and the

improvement in selection metrics for the SC neurons recorded during those sessions.

This supports the view that the target-selection benefits shown here contributed to

efficient search behavior.

FIG. 3: Summary of the fixation-onset-aligned target-selection metrics for the 51 of 65

neurons in our sample that discriminated the saccade target within the fixation-onset-

aligned analysis window for both the NPF-re-NPF and NPF-re-PF analyses. **A.**

Distributions of discrimination times for the two analyses. The black and green bars

show the distributions for the NPF-re-NPF and NPF-re-PF analyses, respectively. Note

that -50 ms relative to fixation onset was the earliest possible discrimination time given

the analysis window. **B.** Relationship between NPF-re-PF and NPF-re-NPF

discrimination times for individual neurons (individual symbols). **C.** Distributions of

discrimination magnitudes for the NPF-re-NPF (black bars) and NPF-re-PF (green bars)

analyses. **D.** Relationship between NPF-re-PF and NPF-re-NPF discrimination

magnitudes for individual neurons (individual symbols).

FIG. 4: Summary of the saccade-onset-aligned target-selection metrics for the 48 of 65

neurons in our sample that discriminated the saccade target within the saccade-onset-

aligned analysis window (-150-0 ms relative to saccade onset) for both the NPF-re-NPF

and NPF-re-PF analyses. Otherwise, what is shown in each panel corresponds to what

is shown in Fig. 3 and the plotting conventions are the same. Note that, for the saccade-

onset-aligned discrimination times, however, the earliest possible discrimination time

was -150 ms.

DISCUSSION

Saccade target selection involves a competition between potential saccade goals on the

SC visuomotor map. An inhibitory-tagging mechanism that facilitates efficient visual

search would be expected to influence this competitive process by reducing activity

associated with previously fixated objects and thus biasing the competitive process

towards uninspected objects in the visual scene. Previously, we showed this basic

pattern of modulations after inhibitory tagging in the SC (6), perhaps reflecting the

influence of inhibitory-tagging signals originating in cortical areas (4, 5). Here, we

characterized and quantified the influence of these modulations on SC target selection.

Using ROC analyses, we found that inhibitory tagging both speeds and strengthens

saccade target selection in the SC, yielding significantly faster discrimination of selected

and unselected RF stimuli and significantly larger discrimination magnitudes. We also

found that the benefit of inhibitory tagging for target selection in these neurons was

related to the monkeys' behavioral search efficiency during the SC recording sessions.

Thus, we conclude that modulation of the selection process at the level of individual SC

neurons likely provides a mechanism by which inhibitory tagging supports efficient

search behavior. Models of target selection based on SC (or, more generally, priority-

map) activity (e.g., Ref. 34) may wish to take these modulations into account to more

accurately capture the dynamics of the selection process during time-limited,

multisaccade search.

SUPPLEMENTARY MATERIAL

Supplementary material: DOI: https://doi.org/10.6084/m9.figshare.24596952

GRANTS

This work was supported by National Institutes of Health (NIH) Grant R01-EY030669 (to

R.M.M.)

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

REFERENCES

1. Posner, M. I., & Cohen, Y. (1984). Components of visual orienting. In H. Bouma

and D. Bowhuis (Eds.), Attention and Performance (pp. 531-556). New Jersey:

Erlbaum.

2. Klein, R. (1988). Inhibitory tagging system facilitates visual search. Nature,

334(6181), 430-431.

3. Klein, R. M., & MacInnes, W. J. (1999). Inhibition of return is a foraging facilitator

in visual search. Psychological Science, 10(4), 346-352.

4. Mirpour, K., Arcizet, F., Ong, W. S., & Bisley, J. W. (2009). Been there, seen that:

a neural mechanism for performing efficient visual search. Journal of

Neurophysiology, 102(6), 3481-3491.

5. Mirpour, K., Bolandnazar, Z., & Bisley, J. W. (2019). Neurons in FEF keep track

of items that have been previously fixated in free viewing visual search. Journal

of Neuroscience, 39(11), 2114-2124.

6. Conroy, C., Nanjappa, R., & McPeek, R. M. (2023). Inhibitory tagging in the

superior colliculus during visual search. Journal of Neurophysiology, 130(4), 824-

837.

7. Glimcher, P. W., & Sparks, D. L. (1992). Movement selection in advance of

action in the superior colliculus. Nature, 355(6360), 542-545.

8. Basso, M. A., & Wurtz, R. H. (1998). Modulation of neuronal activity in superior

colliculus by changes in target probability. Journal of Neuroscience, 18(18),

7519-7534.

9. Horwitz, G. D., & Newsome, W. T. (1999). Separate signals for target selection

and movement specification in the superior colliculus. Science, 284(5417), 1158-

1161.

10. Horwitz, G. D., & Newsome, W. T. (2001). Target selection for saccadic eye

movements: prelude activity in the superior colliculus during a direction-

discrimination task. Journal of Neurophysiology, 86(5), 2543-2558.

11. Krauzlis, R. J., & Dill, N. (2002). Neural correlates of target choice for pursuit and

saccades in the primate superior colliculus. Neuron, 35(2), 355-363.

12. McPeek, R. M., & Keller, E. L. (2002). Saccade target selection in the superior

colliculus during a visual search task. Journal of Neurophysiology, 88(4), 2019-

2034.

13. Ratcliff, R., Cherian, A., & Segraves, M. (2003). A comparison of macaque

behavior and superior colliculus neuronal activity to predictions from models of

two-choice decisions. Journal of Neurophysiology, 90(3), 1392-1407.

14. McPeek, R. M., & Keller, E. L. (2004). Deficits in saccade target selection after

inactivation of superior colliculus. Nature Neuroscience, 7(7), 757-763.

15. Carello, C. D., & Krauzlis, R. J. (2004). Manipulating intent: evidence for a causal

role of the superior colliculus in target selection. Neuron, 43(4), 575-583.

16. Lee, K. M., & Keller, E. L. (2006). Symbolic cue-driven activity in superior

colliculus neurons in a peripheral visual choice task. Journal of Neurophysiology,

95(6), 3585-3595.

17. Ratcliff, R., Hasegawa, Y. T., Hasegawa, R. P., Smith, P. L., & Segraves, M. A.

(2007). Dual diffusion model for single-cell recording data from the superior

colliculus in a brightness-discrimination task. Journal of Neurophysiology, 97(2),

1756-1774.

18. Shen, K., & Paré, M. (2007). Neuronal activity in superior colliculus signals both

stimulus identity and saccade goals during visual conjunction search. Journal of

Vision, 7(5), 15-15.

19. Kim, B., & Basso, M. A. (2008). Saccade target selection in the superior

colliculus: a signal detection theory approach. Journal of Neuroscience, 28(12),

2991-3007.

20. McPeek, R. M. (2008). Reversal of a distractor effect on saccade target selection

after superior colliculus inactivation. Journal of Neurophysiology, 99(5), 2694-

2702.

21. Port, N. L., & Wurtz, R. H. (2009). Target selection and saccade generation in

monkey superior colliculus. Experimental brain research, 192, 465-477.

22. Kim, B., & Basso, M. A. (2010). A probabilistic strategy for understanding action

selection. Journal of Neuroscience, 30(6), 2340-2355.

23. Nummela, S. U., & Krauzlis, R. J. (2010). Inactivation of primate superior

colliculus biases target choice for smooth pursuit, saccades, and button press

responses. Journal of Neurophysiology, 104(3), 1538-1548.

24. Shen, K., Valero, J., Day, G. S., & Paré, M. (2011). Investigating the role of the

superior colliculus in active vision with the visual search paradigm. European

Journal of Neuroscience, 33(11), 2003-2016.

25. Shen, K., & Paré, M. (2014). Predictive saccade target selection in superior

colliculus during visual search. Journal of Neuroscience, 34(16), 5640-5648.

26. Rovamo, J., & Virsu, V. (1979). An estimation and application of the human

cortical magnification factor. Experimental Brain Research, 37, 495-510.

27. Green, D.M., & Swets, J.A. (1966). Signal Detection Theory and Psychophysics.

New York: Wiley.

28. Thompson, K. G., Hanes, D. P., Bichot, N. P., & Schall, J. D. (1996). Perceptual

and motor processing stages identified in the activity of macaque frontal eye field

neurons during visual search. Journal of Neurophysiology, 76(6), 4040-4055.

29. Thomas, N. W., & Paré, M. (2007). Temporal processing of saccade targets in

parietal cortex area LIP during visual search. Journal of Neurophysiology, 97(1),

942-947.

30. Song, J. H., & McPeek, R. M. (2010). Roles of narrow-and broad-spiking dorsal

premotor area neurons in reach target selection and movement production.

Journal of Neurophysiology, 103(4), 2124-2138.Mirpour et al., 2009

31. McPeek, R. M., Han, J. H., & Keller, E. L. (2003). Competition between saccade

goals in the superior colliculus produces saccade curvature. Journal of

Neurophysiology, 89(5), 2577-2590.

32. Port, N. L., & Wurtz, R. H. (2003). Sequential activity of simultaneously recorded

neurons in the superior colliculus during curved saccades. Journal of

Neurophysiology, 90(3), 1887-1903.

33. McPeek, R. M., & Keller, E. L. (2002). Superior colliculus activity related to

concurrent processing of saccade goals in a visual search task. Journal of

Neurophysiology, 87(4), 1805-1815.

34. Adeli, H., Vitu, F., & Zelinsky, G. J. (2017). A model of the superior colliculus

predicts fixation locations during scene viewing and visual search. Journal of

Neuroscience, 37(6), 1453-1467.