Tetraspanin CD81 promotes leukemia stem cell function and represents a new therapeutic vulnerability in acute myeloid leukemia

Fanny Gonzales, MD, PhD[1, 2,*], Pauline Peyrouze, MS[1,*], Thomas Boyer, PharmD, PhD[1, 3], Soizic Guihard, PhD[1], Francois Sevrin, MD[1], Djohana Laurent, MS[1], Adriana Plesa, MD[4], Adeline Barthelemy, PhD[1], Antonino Bongiovanni, PhD[5], Nicolas Pottier, PharmD, PhD[1], Claude Preudhomme, PharmD, PhD[1, 3], Nicolas Duployez, PharmD, PhD[1, 3], Céline Berthon, MD, PhD[1, 6], Christophe Roumier, PharmD[1, 3], and Meyling Cheok, PharmD, PhD[1]

[1] Univ. Lille, CHU Lille, IRCL, CNRS-UMR9020, Inserm-U1277 – CANTHER – Cancer Heterogeneity Plasticity and Resistance to Therapies, F-59000 Lille, France

[2] Pediatric Hematology Department, CHU Lille, Lille, France

[3] Laboratoire d'hématologie, Centre de Biologie-Pathologie, CHU Lille, France

[4] Hospices Civils de Lyon, Service d'hématologie biologique, Pierre Bénite, France

[5] Univ. Lille, CNRS UMR 8204, Inserm U1019, CHU Lille, Institut Pasteur de Lille - CIIL - Center for Infection and Immunity of Lille, Lille, France

[6] Hôpital Claude Huriez, Maladies du sang, CHU Lille, France

[*] These authors contributed equally

Running Title: Therapeutic targeting of CD81 in AML

Keywords: acute myeloid leukemia, CD81, tetraspanin, leukemia stem cell, xenograft, adhesion/migration, drug resistance

Counts: abstract: 192, main: 4046, 6 Figures, 71 References, 1 Supplement

*Correspondence to:
Meyling H. CHEOK, PharmD, PhD
UMR9020 CNRS - UMR-S1277 Inserm
59045 LILLE cedex, France

Abstract

Despite important progress over the last decade, acute myeloid leukemia (AML) is still associated with poor clinical outcome. Novel potent therapies ideally effective against AML stem cells (LSC), a major driver of leukemia initiation and progression, are urgently needed. In particular, targeting common AML-associated antigens at the stem and progenitor cell level represents an attractive therapeutic strategy to achieve deep long-term remissions and is currently the subject of intensive research efforts. In this study, we identified the tetraspanin CD81, a cell surface antigen frequently expressed on AML cells including LSC, as a new determinant of relapse and poor prognosis. CD81 expression was higher in AML cells compared to normal bone marrow cells, and more markedly expressed at relapse. We further showed that modulation of CD81 expression using gain- and loss-of-function approaches affected leukemia aggressiveness, tumor burden, LSC-homing and -xenoengraftment as well as mouse survival. Finally, anti-hCD81 monoclonal antibody-treatment combined with standard chemotherapy in mice with pre-established AML not only reduced leukemia burden but also prolonged relapse-free and overall survival. Collectively, these results identified a new efficacious and safe pharmacological strategy for targeting LSC, opening up novel therapeutic avenues to improve AML outcome.

Key points

- CD81 expression in AML including LSC is a new determinant of aggressive disease and poor prognosis.
- Anti-hCD81 monoclonal antibody-treatment of AML xenografts reduced leukemia burden and improved survival rates.

Introduction

Acute myeloid leukemia (AML) is the most common acute leukemia in adults with an annual incidence of approximately 3 to 4 cases per 100,000 persons worldwide.[1] Over the past few decades, the therapeutic strategy has mostly relied on intensive combination chemotherapy, notably cytarabine and anthracyclines, often followed by hematopoietic stem cell transplantation (HSCT). More recently, various targeted therapies have become available for specific patients,[2] but despite this progress, mortality in AML patients remains high and poor clinical outcome is mainly due to disease relapse or treatment failure, occurring in up to 40% of younger and 80% of older adult AML cases.[3–6] Even though several relapse risk factors are known,[7,8] a better understanding of the mechanisms leading to refractory or relapsed AML is still needed to develop more potent therapeutic strategies.

Therapy-resistant and quiescent leukemia stem cells (LSC), present within the leukemia bulk at very low frequency and functionally defined as capable to initiate AML when transplanted into immunocompromised mice,[9] are considered to be the major cause of disease progression and relapse.[10] Development of new potent treatment strategies to eradicate LSC has thus recently become the focus of intensive research.[11] While various distinct cell surface proteins such as CD33,[12] CD123,[13] CD44, CD47,[14,15] TIM-3,[16] CLL-1[17,18] and CD45RA[19] have been found enriched or absent such as CD90[20] on LSC, therapies targeting these markers are lagging behind expectations, mainly due to their overlapping expression with hematopoietic stem and progenitor cells (HSPC) as well as the lack of antigens broadly and consistently expressed on LSC in most AML subtypes. Therefore, the identification of new druggable and discriminating LSC markers remains essential for the successful development of a selective anti-LSC therapy.

CD81 refers to a cell surface protein, also known as "target of an antiproliferative antibody 1" (TAPA-1),[21] that belongs to the evolutionary conserved tetraspanin superfamily of proteins. CD81 is the structural component of specialized membrane microdomains known as tetraspanin-enriched microdomains,[22–24] and plays an important role in signal transduction by interacting with various proteins such as integrins and several distinct membrane

proteins.[25,26] In particular, CD81 has been shown to regulate B cell signaling by interacting

with CD19 and CD21 on B-lymphocytes[27,28] and is known to be overexpressed in several

hematological malignancies implicating B cells such as lymphoma[21,29] and acute

lymphoblastic leukemia.[30] Importantly, previous studies have demonstrated that

immunotargeting of CD81 represents a new effective treatment for B cell lymphoma.[29,31]

Nevertheless, despite its adverse role in B cell malignancies, its functional implication in AML

remains to be characterized.

In this study, using a combination of *in vitro* and *in vivo* approaches, we demonstrated

that CD81 is an important determinant of AML progression, drug resistance and leukemia

stemness. Finally, we provide proof of concept that targeting CD81 may represent a safe and

efficacious therapeutic strategy to eradicate LSC in AML.

Materials and Methods

Patients and cell line models

In this study, we included 11 normal bone marrow (BM) samples and 290 primary AML including 38 relapse samples recruited at Lille University Hospital (2009-2016, Table 1). All patients have signed the informed consent form and the Institutional Review Board approved these studies (ANA-FE-BMH 20221007). CD81 protein expression (N = 290, Supplemental Figure 1), CD81 mRNA expression (N = 53), LSC17 gene expression score (N = 107) and AML xenoengraftment (N = 59) were performed as described previously.[32–34] Diagnostic patient characteristics are given in Table 1, which excluded not intensively treated (N = 48), pediatric (N = 16) and patients with incomplete clinical data (N = 8).

AML cell lines U-937, OCI-AML3, HNT-34 (DSMZ), and BM stromal fibroblasts HS-5 (ATCC) were cultured as recommended. Authenticated AML cell lines (FTA Kit, LGC, ATCC-135-XV-20) were transduced using the luciferase containing lentiviral vector (pLenti PGK V5-LUC Neo (623-2), Addgene). U-937 cells were transfected with pCDM8 hCD81[21] (Addgene) using the Amaxa Nucleofactor II, (kit C, W001; Lonza) and CD81 positive cells sorted (ARIA III BD, BD Biosciences). HNT-34 and OCI-AML3 cells were transduced with shRNA-CD81 or non-targeting (NT) shRNA lentiviral vectors (TRCN0000300291[sh291], TRCN0000300293[sh293], TRCN0000300433[sh433] or TRC2 pLKO.5-puro; Sigma-Aldrich).

AML xenografts

Patient-derived xenografts (PDX) and cell-line-derived xenografts (CDX) were generated using NSG immunodeficient mice as previously described[33] (Supplemental Figure 2). We monitored CDX engraftment by *in vivo* imaging (luciferin 0.1 mg/kg IP, Xenogen IVIS 50, Perkin Elmer). We collected femurs and sternums from CDX when hCD45 was >1% in peripheral blood (PB) and from control mice injected with PBS. Bones were fixed with 4% paraformaldehyde-PBS overnight at 4°C, decalcified (DC1, Q Path) for 18h at 4°C, dehydrated and embedded in paraffin. Immunohistochemistry was performed on 4 µm

sections using the BenchmarkUltra (Roche), anti-hCD45 (M0701, Dako) and UltraView DAB

Detection Kit (Ventana). We evaluated AML cell homing by flow cytometry (FCM) analysis of

BM using a minimum of 1 x10^6 acquired events, 72h after IV injection of 12 x10^6 (OCI-AML3)

or 20 x10^6 (U-937) AML cells. French ethics committee approved all animal experiments

(APAFIS#33794-2021100815558055).

CD81 expression

CD81 cell surface protein expression was determined by FCM (LSR Fortessa X20,

BD Biosciences), 5 x10^5 cells were labeled with 1 µg anti-hCD81-PE clone 5A6 or 1 µg

isotype control-PE (349506, 400114; Biolegend)[32] and by Western blotting (4-12% SDS-

PAGE, clone JS-81, BD Biosciences). The FCM antibody panel and gating strategy are

detailed in Supplemental Figure 1.[33] CD81 mRNA was quantified by Taqman RT-PCR and

GAPDH normalized (CD81: Hs00174717_m1, GAPDH: Hs03929097_g1, StepOnePlus Real-

Time PCR System, ThermoFisher).

***In vitro* drug resistance**

We assessed drug resistance *in vitro* as previously described.[35] Drug concentration

ranges were 0.000042-3.33 µg/mL (daunorubicin [DNR]) and 0.0025-40 µg/mL (cytarabine

[AraC]) and drug exposure times were 72h (AML cell lines) and 96h (primary AML cells).

Cell adhesion, migration and invasion

We coated 96-well plates with fibronectin (1 µg per cm^2, S5171, Sigma-Aldrich) or

seeded 4 x10^4 HS-5 for 48h (70-80% confluence). Then, cells in 100 µL (U-937, OCI-AML3

[1 x10^6]; HNT-34 [0.5 x10^6]) were allowed to adhere in triplicates for 90 min (U-937, OCI-

AML3) or 30 min (HNT-34). Plates were washed three times with PBS and bioluminescence

measured (luciferin 0.3 mg/mL, Xenogen IVIS 50; Perkin Elmer). We used 24-well transwell

inserts and plates (5 µm 29442-118, Corning) to evaluate cell migration and invasion. Lower

chambers were seeded with 5 x10^5 HS-5 in 300 µL 24h prior to the experiment. Then, 3 x10^5

AML cells in 300 µL were loaded onto the upper chambers and cultured for 24h. Cells

migrated to the lower chambers were quantified by bioluminescence as described above. For

invasion experiments, the upper chambers were loaded with a gel matrix using 100 µL

Basement Membrane Extract (BME) at 0.1X per insert (Cultrex 5X BME Solution, Trevigen)

4h prior to AML cell loading. All experiments were performed in duplicates. Adhesion,

migration and invasion rates were determined as the percentage of cells *vs.* input control

(Supplemental Figure 3).

Confocal microscopy

AML cells (6×10^4 cells/cm^2) were seeded for 24h on poly-D-Lysine (Gibco) coated

Lab-Tek II Chamber Slides (Nunc) at a density of. Then, slides were rinsed with PBS and

cells fixed with 4% paraformaldehyde-PBS for 20 min and permeabilized with PBS-Triton X-

100 0.1% for 10 min at room temperature (RT). After blocking with PBS-BSA 2% for 45 min,

cells were stained with Alexa Fluor 488 phalloidin (1:40, Invitrogen) for 30 min at RT in the

dark, and nuclei were stained with DAPI (ThermoFisher) for 10 min. We generated confocal

microscopy images (LSM710, Zeiss) and estimated cellular diameter and circularity using

ImageJ, (https://imagej.nih.gov/ij/)[36].

Anti-hCD81 antibody-treatment

Cells were exposed to anti-hCD81 monoclonal antibody or isotype control for 4h with

1 µg per 1×10^6 cells in 100 µL. AML bearing mice were treated with anti-hCD81 antibody *vs.*

isotype control (JS-81, BD Biosciences) 100 µg, IP using three or five doses in OCI-AML3

CDX and PDX, respectively in combination with chemotherapy.[37] *In vivo* imaging of OCI-

AML3 CDX and FCM of PDX blood[33], was used to track leukemia engraftment and disease

progression.

Colony-Forming Unit (CFU) assays

CFU assays were performed according to the MethoCult™ H4034 Optimum protocol

(Stemcell Technologies). Briefly, 5 x10^4 and 2.5 x10^4 human BM mononuclear cells (BMNC,

Lonza) were plated in triplicate and cultured for 10 days. Colonies were counted and

normalized to the number of cells seeded on day 0.

Cell cycle and viability analyses

Cell cycling of BMNC was determined on 0.25 x10^6 BMNC washed in PBS and fixed

in cold ethanol 70% at -20°C for 2h. Then, cells were washed twice in PBS-FBS 1% and

stained with 20 µL of anti Ki-67 (FITC Mouse Anti-Ki-67 Set, BD Biosciences) for 30 min at

RT in the dark. Cells were washed in PBS-FBS 1%, incubated with RNaseA 2 µg/mL for 15

min at 4°C and stained with PI 35 µg/mL for 30 min before analysis by FCM. Viability of

BMNC (0.25 x10^6) was assessed by Annexin V and 7-AAD staining (APC Annexin V

Apoptosis Detection Kit 7-AAD, Biolegend) using FCM.

Statistics

We used Student's t-test and Pearson correlation statistics unless otherwise specified

and results included sample means, standard deviation and R^2. We evaluated time-to-

xenoengraftment using Cox proportional hazard model with *P*-values based on the log-rank

test. We censored time at transplantation date in the event of HSCT or at the date of last

follow-up and defined overall survival (OS) as the time from date of inclusion to date of death

from any cause. Relapse-free survival (RFS) was determined in patients achieving complete

remission (CR) and is defined as the time from date of CR to date of first relapse, or in the

case of no relapse, time was censored at the date of last follow-up. The alpha level was set

to 5% and all statistical tests were two-sided. We used GraphPad Software version 10

(Prism). Graphs and analyses were computed using Kaplan-Meier or Cumulative incidence

plots, hazard ratio (HR) and 95% confidence intervals (CI) were determined using R-

v.4.2.0.[38] CD81 mRNA expression and survival of publicly available Beat-AML dataset[39] were

obtained *via* the cBioPortal.[40,41] Similar to our cohort the analyses included AML (non-FAB-

M3, non-core-binding factor (CBF), age >50, chemotherapy).

Results

CD81 overexpression is associated with AML relapse and poor prognosis.

Previously, we determined that high CD81 expression prognostic using a cohort of 134 diagnostic AML patients.[32] These initial findings are now extended using a cohort of 252 diagnostic and 38 relapse samples. Since CBF-AML have well-known favorable prognosis, we first compared CD81 protein expression between CBF and non-CBF AML and found it to be lower in the CBF group (N = 28, Figure 1A). Therefore, we excluded CBF-AML to avoid selection bias and confirmed that higher CD81 protein expression adversely affected RFS and OS (N = 150, RFS: HR [CI] = 1.91 [1.12-3.28]; OS: HR [CI] = 1.88 [1.07-3.32]; Figure 1B). Interestingly, we also showed that higher CD81 protein expression negatively affected OS within *NPM1* mutated AML, a molecular subtype with heterogeneous clinical response (N = 54, OS: HR [CI] = 7.71 [1.00-59.23]; Figure 1C), independently of co-existing *FLT3-ITD* mutational status (data not shown) and a similar trend was obtained for relapse (RFS: P = 0.15; Figure 1C). As both CD81 mRNA expression and protein abundance were correlated in our cohort (P = 0.01, Figure 1D), we further characterized the detrimental effect of CD81 overexpression on AML prognosis independently using the publicly available mRNA data from the Beat AML dataset.[39] Consistent with our results, CBF-AML from the Beat-AML cohort also showed a significant lower CD81 mRNA expression compared to non-CBF AML (Figure 1E). Importantly, we confirmed the negative impact of high CD81 mRNA on OS not only within non-CBF AML but also within *NPM1* mutated AML (non-CBF AML: HR [CI] = 1.80 [1.00-3.26]; *NPM1* mutated AML: HR [CI] = 5.39 [1.58-18.36]; Figure 1F). Finally, given the prognostic effect of high CD81 expression, we hypothesized that blasts at diagnosis exhibit an increased expression of this protein and an even higher CD81 expression at relapse, a disease state usually refractory to treatment. As expected, CD81 was overexpressed in AML compared to BM obtained from healthy individuals (Figure 1G) and AML at relapse exhibited higher CD81 expression when compared to samples at diagnosis (Figure 1H) or using paired diagnostic-relapsed samples (67% *vs.* 19%, Figure 1I).

Collectively, these findings demonstrated that CD81 overexpression on AML blasts

adversely affects AML outcome and may contribute to the aggressiveness characterizing

blasts at relapse.

***In vitro* modulation of CD81 expression affects xenoengraftment and leukemia**

aggressiveness

To further strengthen the deleterious role of CD81 in AML, we modulated its

expression in human AML cell lines: OCI-AML3, HNT-34 and U-937. Overexpression of

CD81 was performed in U-937 cells, which have undetectable CD81 expression, whereas

silencing of CD81 was achieved using a shRNA approach in HNT-34 and OCI-AML3 cells,

which by contrast highly express CD81 (Supplemental Figure 4). Since we linked primary

CD81 overexpression to AML relapse, we first assessed whether modulating CD81

expression affected chemoresistance. We found that U937-CD81 overexpressing cells were

more resistant to daunorubicin, whereas CD81-depleted cells (OCI-AML3, HNT-34) were

more sensitive to daunorubicin and cytarabine (Figure 2A). Then, we tested engraftment

efficacy, a recognized indicator of AML aggressiveness,[42,43] evaluated by *in vivo* imaging of

leukemia cells exhibiting either high or low CD81 expression in immunocompromised NSG

mice. As shown in Figure 2B, xenoengraftment was more successful in U-937

overexpressing CD81 compared to control cells, whereas CD81-depleted OCI-AML3 cells

exhibited reduced xenoengraftment capacity. Consistent with this, AML cell homing was

similarly affected by CD81 modulation (Figure 2C). As AML engraftment implicates multiple

cell-based processes such as adhesion, migration and invasion,[44] we also examined the

contribution of CD81 on each of these cellular programs (Figure 3A). Of particular interest,

overexpression of CD81 in the U-937 model increased AML cell adhesion using fibronectin-

coated plates or HS-5 fibroblast co-cultures, whereas CD81 depletion in both OCI-AML3 and

HNT-34 cell models had the opposite effect. Similar results were obtained for cellular

migration and invasion of AML cells into a semisolid gel matrix. Confocal microscopy using

phalloidin staining additionally showed that CD81 overexpressing AML cells formed filopodia-

like membrane protrusions, a typical morphological change reminiscent of the cellular

processes investigated (Figure 3B). Finally, we evaluated whether CD81 modulation also

affected leukemia aggressiveness by assessing mouse survival and AML infiltration into

various tissues using hCD45 as a marker of leukemic cells. As shown in Figure 4A, mice

engrafted with U937-CD81 overexpressing cells have higher leukemia burden in blood as

well as in BM and its surrounding muscle tissue (Figure 4B) along with a worse survival (HR

[CI] = 3.42 [1.32-9.90], Figure 4C). By contrast, CD81-depleted cells induced a less

aggressive phenotype with reduced tissue infiltration (OCI-AML3, Figure 4D) and a better

survival (HR [CI] = 0.33 [0.12-0.09], Figure 4E).

Altogether, these results indicate that CD81 is an important determinant of leukemia

chemoresistance and aggressiveness.

CD81 is associated with LSC function in primary AML

As LSC are considered the main source of relapse[45–47] and given the increased CD81

expression in blasts at the time of relapse, we investigated whether CD81 also affects LSC

function. First, we assessed its expression within the AML LSC subpopulation, defined as

CD34+/CD38-/CD90-/CD123+. This showed that CD81 was expressed within this LSC

population and that CD81+ LSC fraction had increased at relapse when compared to

samples obtained at diagnosis (Figure 5A) or between paired relapse and diagnosis samples

(49% *vs.* 20%; Figure 5B). Similar results were found for the non-HSPC fraction defined as

CD34+/CD38-/CD90- (53% *vs.* 8%; Supplemental Figure 5A, B). Then we investigated

whether a high proportion of LSC positive for CD81 negatively affects AML outcome. Our

results showed that non-CBF AML patients with a higher proportion of CD81+LSC at

diagnosis were at increased risk of relapse and death (RFS: HR [CI] = 2.01 [1.16-3.48]; OS:

HR [CI] = 2.38 [1.32-4.29]; Figure 5C). Interestingly, the fraction of LSC positive for CD81

was correlated with bulk CD81 protein expression (Figure 5D), suggesting that bulk CD81

expression likely reflects the proportion of LSC positive for CD81.

To further characterize the relationship between CD81 overexpression and LSC

function, we performed serial AML xenoengraftment experiments using immunocompromised

NSG mice. As expected, patients whose AML xenoengrafted had worse outcome than those

whose AML failed to engraft (RFS: HR [CI] = 4.45 [1.31-15.16]; OS: HR [CI] = 2.74 [1.15-

6.51]; Figure 5E) and were more resistant to cytarabine and daunorubicin (Figure 5F-G).

Moreover, our results also showed that AML with higher CD81 expression were not only

associated with positive engraftment (Figure 5H) but also more successfully engrafted over

time compared to low CD81 expressing AML (HR [CI] = 4.53 [1.64-12.56]; Figure 5I).

Interestingly, PDX of AML with higher CD81 expression had larger spleens at sacrifice than

those with low CD81 expression (Figure 5J), suggestive of higher extra-medullary infiltrative

potential. Importantly, CD81 expression in non-CBF AML was correlated with the LSC17

gene signature score (Figure 2K), a known and reliable marker of functional LSC in AML.[34,48]

Taken together, these results unambiguously showed that high CD81 expression in

primary AML is associated with functional LSC.

CD81-targeting as a therapeutic strategy in AML.

Having established the deleterious role of CD81 in AML and its translational

relevance, we then investigated the therapeutic potential of an anti-hCD81 antibody. Firstly,

we evaluated whether *in vitro* exposure of AML cells with this immunotherapy is able to

inhibit their engraftment into NSG mice. Consistent with the data obtained using CD81-

depleted cell lines, antibody-treatment of OCI-AML3 cells effectively reduced their

engraftment into murine BM and PB and similarly reduced spleen infiltration, the preferential

infiltration site of OCI-AML3 cells (Figure 6A). Additionally, targeting of CD81 using this

therapeutic approach also increased *in vitro* sensitivity of OCI-AML3 to daunorubicin and

cytarabine compared to IgG control (Figure 6B). Then, we evaluated potential toxicity by

testing the *in vitro* effect of anti-hCD81 antibody on the viability of primary mononucleated

BM cells (BMNC) obtained from a healthy individual. Of great interest, antibody-treatment

neither affected BMNC viability at 4 and 24h nor altered BMNC cell cycle when compared to

IgG control (Figure 6C-D). More importantly, no sign of BMNC toxicity was detected for the

hematopoietic progenitors CFU-GEMM, CFU-GM, BFU-E or CFU-E (Figure 5E). Finally, we

applied a curative protocol (Figure 6F) consisting of IP administration of either the anti-

hCD81 antibody or control before, during and after intensive chemotherapy[29] in leukemia

bearing NSG mice using not only OCI-AML3 cells (Figure 6G), but also newly diagnosed

primary AML cells (Figure 6H) and assessed treatment response by *in vivo* imaging and

FCM, respectively. Our results showed that immunotargeting of CD81 strongly reduced

tumor burden as early as one week after the last injection of the antileukemic drugs (Figure

6G, Supplemental Figure 6A) when using OCI-AML3 cells and more importantly, significantly

reduced tumor burden after ten weeks off chemotherapy when using primary AML cells

(Figure 6H, Supplemental Figure 6B). Moreover, in both cases, leukemia bearing mice

treated with the anti-hCD81 antibody exhibited a less aggressive disease with delayed

relapse and subsequently improved survival (HR [CI] = 2.60 [1.06-6.34], Figure 6I and HR

PB-relapse [CI] = 5.228 [0.874-31.27], HR [CI] = 5.20 [0.53-50.6], Figure 6J).

In summary, immunotargeting of CD81 may represent a novel safe and effective

therapeutic strategy for AML.

Discussion

AML is a hematological stem cell malignancy originating from a rare subpopulation of LSC that is responsible for the accumulation of undifferentiated myeloid blast cells at the expense of the hematopoietic system.[49] Recent sequencing studies have revealed that AML is genetically complex with patients frequently harboring many distinct driver mutations whereas no commonly dysregulated oncogenic pathways has been identified yet.[50] This implies that specific targeting of any particular leukemogenic molecular alteration may not only be effective in just a small subset of patients but may also only eliminate a fraction of neoplastic cells, as multiple mutations frequently coexist within a leukemia. Therefore, it is increasingly evident that achieving long-term remissions or even curing AML would unlikely rely on a single targeted-therapy[51–53] but instead should ideally require the development of treatment strategies effective against LSC and across a wide range of AML subtypes, irrespective of their mutational status. In this regard, AML-specific cell surface proteins detectable in most patients have raised particular theranostic interest such as CD33, CD123, CLL-1, CD44, CD47 or TIM-3 and are currently under investigation.[11,49] Nevertheless, targeting these molecules is lagging behind expectations as severe toxicities caused by the lack of expression specificity and resistance frequently occur. For example, targeting CD33 (Siglec-3), a myeloid cell surface antigen highly expressed in most AML subtypes and LSC-enriched populations, using a recombinant humanized anti-hCD33 antibody conjugated to the cytotoxic calicheamicin (gemtuzumab ozogamicin)[54] has shown efficacy especially in favorable AML while inducing severe toxicity given its expression on normal hematopoietic cells.

In this study, we aimed to further assess the therapeutic potential of CD81 in AML, a cell surface glycoprotein expressed on immune cells and known to be essential for B cell maturation and differentiation.[55,56] Our results notably showed that CD81 is an independent prognostic marker of treatment response in AML, especially in non-favorable AML. Consistent with this, we functionally validated its deleterious role using gain and loss of function *in vitro* and *in vivo* approaches by demonstrating that CD81 expression affects

leukemia aggressiveness and tumor burden by enhancing cell adhesion, migration and

invasion processes.[57–59] Indeed, tetraspanins such as CD81 are major components of

specialized membrane microdomains known to interact with the extracellular matrix to

mediate cell adhesion and motility,[24,60] but a functional role of CD81 in myeloid cell adhesion

has not been reported yet.

As LSC are considered to play a pivotal role in AML relapse, eradicating these cells

by identifying a highly expressed if not specific therapeutic target remains critical to ultimately

cure AML while sparing other important cell populations such as normal HSPC clones.[61] As

cell adhesion and motility are fundamental for LSC interaction with the BM niche

microenvironment and given our results showing the importance of CD81 in these processes,

we reasoned that this cell surface protein may also influence LSC function. For this, we first

showed that CD81 is expressed within the AML-LSC-containing subpopulation commonly

defined as CD34+CD38-CD90-. Indeed, previous studies combining immunophenotyping of

AML samples and xenotransplantation have established that LSC are characterized by the

CD34+CD38-CD90- immunophenotype in more than 90% of AML cases.[19,47] Interestingly,

CD81 was also expressed in the CD34+CD38-CD90-CD123+ subset which included CD123

(IL-3Rα), a well-established LSC specific cell surface marker.[12,13,62] As targeting several

antigens may be necessary to eliminate leukemic cells that persist at remission,

combinatorial immunotargeting[63] of both CD81 and CD123 may represent a novel

therapeutic approach especially to circumvent antigen escape, a major cause of resistance

to immunotherapy. Nevertheless, whether CD81 expression persists in LSC and progenitor

populations of AML patients during clinical remission remains to be investigated. Consistent

with previous studies related to CD33[64] and CD123,[65] we showed that CD81 was

overexpressed on bulk tumor and LSC in most AML patients across different subtypes of

AML, and more importantly, that its expression increased from diagnosis to relapse,

suggesting a role for CD81 in AML treatment escape and disease progression. Therefore,

targeting CD81 may not only be effective against persistent leukemic cells but also as

second-line treatment when relapse occurs. Lastly, we demonstrated that CD81 expression

is associated with functional LSC[9] using serial xenoengraftment experiments as well as the

LSC17 score,[48] a known surrogate marker of LSC activity.

Finally, as targeting cell surface proteins are often associated with severe toxicity

related to their expression in hematopoietic tissues,[12,47] we evaluated whether anti-hCD81

antibody-treatment is a safe strategy. When exposed to normal BMNC, antibody-treatment

revealed no hematopoietic toxicity neither alone nor in combination with intensive

chemotherapy and was well tolerated by xenoengrafted mice. This is consistent with

previous reports showing that (i) CD81 expression is low in normal BM cells,[32] (ii) anti-

mCD81 antibody-treatment is well-tolerated in inflammatory bowel diseased mice[66] and more

importantly, (iii) an anti-hCD81 antibody developed as an inhibitor of HCV infection[67] and

recognizing both human and monkey CD81 is well-tolerated in primates using several

therapeutic doses over a prolonged time.[68] Lastly, murine knockout models of CD81 have

been established and no life threatening hematopoietic defects were observed except

reduced B cell activity,[55,69,70] mirroring the phenotype characterizing human autosomal

recessive CD81 deficiency.[71]

In conclusion, we provide proof of concept for CD81 as a new relevant therapeutic

target and diagnostic in AML. In particular, immunotargeting of CD81 combined with

standard chemotherapy may represent a novel efficacious treatment strategy to improve

AML outcome especially during post-remission therapy and at relapse given the importance

of CD81 for LSC function.

Table 1. Diagnostic parameters of the study cohort treated with standard

chemotherapy.

Parameter	Total	CD81 low	CD81 high	P-value
No. of patients (%)	150 (100)	68 (45)	82 (55)	-
Age [y], median (min-max)	55.6 [17.3-77.6]	53.6 (17.3-77.6)	56.3 (17.8-75.7)	0.71
Gender (M/F)	72/78	31/37	41/41	1.0
WBC [x10^9/L], median (min-max)	27.2 (1-405)	13.4 (1-249)	46.1 (1-405)	0.010*
BM blast [%], median (min-max)	70 (5-98)	61.5 (5-97)	72 (5-98)	0.114
Cytogenetics				0.86
Favorable, *N* (%)	-	-	-	
Normal, *N* (%)	85 (56.7)	41 (60.3)	44 (53.6)	
Abnormal, *N* (%)	35 (23.3)	13 (19.1)	22 (26.8)	
Adverse, *N* (%)	26 (17.3)	12 (17.6)	14 (17.1)	
NA, *N* (%)	4 (2.7)	2 (3)	2 (2.5)	
ELN2022 risk				0.73
Favorable, *N* (%)	26 (17.3)	15 (22.0)	11 (13.0)	
Intermediate, *N* (%)	94 (62.7)	39 (57.4)	55 (67.1)	
Adverse, *N* (%)	26 (17.3)	12 (17.6)	14 (17.1)	
NA, N (%)	4 (2.7)	2 (3)	2 (2.5)	

*P <0.05

Figure Legends

Figure 1. CD81 expression is associated with AML relapse and poor clinical outcome.

(A) Violin plot comparing CD81 expression in non-CBF *vs*. CBF-AML (*N* = 252). (B) Kaplan-

Meier curves showing worse RFS and OS of patients with higher CD81 expression in their

AML and similarly, (C) in a subset of patients with *NPM1* mutant AML. (D) Scatterplot

correlating CD81 mRNA and protein expression (*N* = 33, R^2 = 0.19, *P* = 0.019). (E) Violin plot

depicting that favorable CBF AML expressed lower levels of CD81 mRNA expression (Beat

AML dataset, *N* = 364). (F) Kaplan-Meier curves of patients stratified by median CD81 mRNA

at AML diagnosis, showing worse OS for patients with AML overexpressing CD81 overall

(left panel) and within *NPM1* mutant AML (right panel). (G) Violin plot indicating CD81

overexpression in AML compared to normal BMNC, (H) which increased at relapse

compared to leukemia diagnosis on all AML (*N* = 290) and (I) also in paired AML samples

(red: increase, blue: no change, green: decrease; *N* = 19; paired t-test). *P <0.05, **P

<0.005, ***P <0.0005.

Figure 2. AML models of high CD81 expression are chemoresistant and capable to engraft in immunodeficient mice.

(A) Daunorubicin (DNR, left) and cytarabine (AraC, right) drug-response curves and bar charts of IC_{50} using AML models with either CD81 overexpression (red, CD81+) or silenced CD81 (blue, CD81-) *vs.* respective controls U-937 [top], OCI-AML3 [middle], HNT-34 [bottom] evaluated by *in vitro* drug resistance (N >3). (B) Line graphs of luminescence and *in vivo* images of xenoengrafts demonstrating increased tumor burden of U-937 CD81 overexpressing (CD81+, red) cells compared to control (N =17, Supplemental Figure 8A). Conversely, OCI-AML3 CD81 silencing (CD81-, blue) decreased tumor burden compared to control (N =11, Supplemental Figure 8B). (C) AML cell homing by FCM analysis of BM cells 72h after IV injection indicate increased BM homing when CD81 was overexpressed (top panel, U-937, $N = 3$, MWU) whereas it decreased when CD81 was silenced (bottom panel, OCI-AML3, $N = 5$, MWU). *P <0.05, **P <0.005, ***P <0.0005.

Figure 3. CD81 overexpression modeling in AML reveals enhanced adhesion, migration and invasion capacity and filopodia-like membrane protrusion formation.

CD81 overexpression (CD81+, red) induced cell adhesion to fibronectin and HS-5 fibroblasts, migration and invasion compared to control (U-937, N >4, left panels). Conversely, CD81 silencing (CD81-, blue) decreased cell adhesion in both OCI-AML3 (N >3, middle panel) and HNT-34 (N >3, right panel) compared to control. (B) Confocal microscopy displaying filopodia-like cell membrane protrusions in high CD81 expressing AML models (top images, U-937, OCI-AML3, HNT-34) compared to their CD81 low expression counterparts exempt of membrane protrusions (bottom images). Corresponding bar charts summarize cell size and circularity detected by confocal microscopy (U-937, $N = 3$, 218/434 cells; OCI-AML3, $N = 2$, 295/170 cells; HNT-34, $N = 2$, 263/173 cells). *P <0.05, **P <0.005, ***P <0.0005.

Figure 4. CD81 overexpression models of AML induce leukemia burden and aggressive disease in xenoengrafted mice.

(A) CD81 overexpression (CD81+, red) induced leukemia infiltration in BM and PB in U-937

AML xenografts (N = 7). (B) Immmunohistochemical analysis of the epiphyseal bone of U-

937 AML xenografts stained with anti-hCD45 antibody indicate that CD81 overexpression

(CD81+, left panel) induced higher BM (green outline) and adjacent muscle tissue AML

infiltration (red outline) compared to control (CD81-, right panel). (C) Kaplan-Meier plot

illustrating worse survival of mice xenografted with U937 CD81 overexpressing cells (CD81+)

compared to control (N = 12). (D) In contrast, CD81 silencing in OCI-AML3 cells (CD81-,

blue) reduced leukemia infiltration in BM, PB and spleen of AML xenografts (N = 7) and (E)

prolonged survival in HNT-34 compared to control (N = 10). *P <0.05, **P <0.005, ***P

<0.0005.

Figure 5. CD81 overexpression in AML is associated to increased LSC function,

chemoresistance and worse clinical prognosis.

(A) Violin plot indicating that the LSC fraction with CD81 protein expression increased from

diagnosis to relapse (LSC defined as CD34+CD38-CD90-CD123+; N = 290) and (B) similar

was found for paired diagnosis and relapse samples (red: increase, blue: no change, green:

decrease; N = 19, paired t-test). (C) Kaplan-Meier plot showing worse RFS and OS in AML

patients with higher CD81 (red) compared to lower CD81 (blue) within the LSC fraction at

AML diagnosis (N = 150). (D) Scatterplot correlating CD81 protein expression within the LSC

compartment with CD81 expression of the leukemia bulk (N = 290; R^2 = 0.44, P <0.0001). (E)

Kaplan-Meier curves of RFS and OS for patients stratified by xenoengraftment capacity of

primary AML (N = 56), where patients with engrafting AML (red) had worse outcome

compared to those whose AML cannot engraft (blue). (F) Cytarabine- (N = 11) and (G)

daunorubicin (N = 14) drug-response curves and bar plots of IC_{50} tested at AML diagnosis

indicating higher resistance in engrafting (red) *vs.* not engrafting (blue) AML (MWU,

Supplemental Figure 9). (H) High CD81 protein expression is not only associated with

positive xenoengraftment (N = 40, MWU), but was also more successful as show by (I)

cumulative incidence stratified by median expression (N = 40). (J) Scatter plot of spleen

weights of xenografts comparing high *vs.* low median CD81 demonstrating that higher CD81

expression resulted in higher splenic infiltration ($N = 21$, MWU). (K) Scatterplot correlating

LSC17 gene score and CD81 protein expression in non-CBF AML ($N = 28$. $R^2 = 0.19$, $P = $

0.018). *$P < 0.05$, **$P < 0.005$, ***$P < 0.0005$.

Figure 6. Efficacy and toxicity of therapeutic targeting of CD81 in AML.

(A) Anti-hCD81 antibody-treatment decreased OCI-AML3 cell engraftment in BM, PB and

spleen (N >7) and (B) sensitized OCI-AML3 cells to chemotherapies: daunorubicin (DNR, *N*

= 6) and cytarabine (AraC, $N = 4$) compared to control. Anti-hCD81 antibody-treatment of

normal BMNC cells affected neither viability (C), cell cycle (D) nor hematopoietic progenitor

CFU (E) when compared to control. (F) Curative treatment protocols of OCI-AML3 CDX and

PDX assessing anti-hCD81 antibody-treatment efficacy. When AML engraftment was 1-5%

in PB, treatment was initiated with anti-hCD81 or control (black arrows) and chemotherapy

consisting of cytarabine (5 days, blue), doxorubicin (3 days, dark red) and gemtuzumab

ozogamicin for CDX (day 3 and 5, green). (G) Line graph (left panel) illustrates therapeutic

efficacy of anti-CD81 antibody (blue, $N = 11$) in reducing tumor burden compared to control

(red, $N = 10$) using OCI-AML3 CDX and corresponding *in vivo* images (right panel). (H) Line

graph illustrating reduced AML burden in PB (left panel) and lower cumulative incidence of

relapse (right panel) of PDX mice treated with anti-hCD81 antibody (blue, $N = 5$) compared to

control (red, $N = 3$). Kaplan-Meier plots of (I) CDX- and (J) PDX mice stratified by treatment

showing that prolonged OS with anti-hCD81 antibody *vs.* control. CTL: control, CFU: colony-

forming unit, *$P < 0.01$, **$P < 0.001$, ***$P < 0.0001$.

Acknowledgements

French National Cancer Institute INCa grants: SIRIC Oncolille 2014-17 (CP, MC), PRTK

2016 (CP, MC) and PLBio 2018 (MC), the Foundations Laurette Fugain 2012 (MC) and ARC

2014 (MC) and the Association Ligue contre le cancer 2016 (MC). The authors are grateful

for the participation of the patients, and clinical and administrative staff implicated in this study.

Disclosures

The authors declare that they have no competing interests.

Authors' contributions

FG, PP, NP, ND, MC wrote the manuscript; FG, PP, TB, SG, AB, FS, FD, AP, CR, MC performed experiments and analyzed results, CP, MC designed and financed experiments, CR, MC interpreted and discussed results, and CB is the referring physician of these patients.

References

1. Visser, O. *et al.* Incidence, survival and prevalence of myeloid malignancies in Europe. *European Journal of Cancer* **48**, 3257–3266 (2012).

2. Tiong, I. S. & Wei, A. H. New drugs creating new challenges in acute myeloid leukemia. *Genes, Chromosomes and Cancer* **58**, 903–914 (2019).

3. Döhner, H. *et al.* Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood* **129**, 424–447 (2017).

4. Röllig, C. *et al.* Long-term prognosis of acute myeloid leukemia according to the new genetic risk classification of the European LeukemiaNet recommendations: evaluation of the proposed reporting system. *J Clin Oncol* **29**, 2758–2765 (2011).

5. Dombret, H. & Gardin, C. An update of current treatments for adult acute myeloid leukemia. *Blood* **127**, 53–61 (2016).

6. Vardiman, J. W. *et al.* The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes. *Blood* **114**, 937–951 (2009).

7. Thol, F., Schlenk, R. F., Heuser, M. & Ganser, A. How I treat refractory and early relapsed acute myeloid leukemia. *Blood* **126**, 319–327 (2015).

8. O'Reilly, E., Zeinabad, H. A. & Szegezdi, E. Hematopoietic versus leukemic stem cell quiescence: Challenges and therapeutic opportunities. *Blood Reviews* **50**, 100850 (2021).

9. Lapidot, T. *et al.* A cell initiating human acute myeloid leukaemia after transplantation into SCID mice. *Nature* **367**, 645–648 (1994).

10. Shlush, L. I. *et al.* Tracing the origins of relapse in acute myeloid leukaemia to stem cells. *Nature* (2017) doi:10.1038/nature22993.

11. Pollyea, D. A. & Jordan, C. T. Therapeutic targeting of acute myeloid leukemia stem cells. *Blood* **129**, 1627–1635 (2017).

12. Taussig, D. C. Hematopoietic stem cells express multiple myeloid markers: implications for the origin and targeted therapy of acute myeloid leukemia. *Blood* **106**, 4086–4092 (2005).

13. Jordan, C. *et al.* The interleukin-3 receptor alpha chain is a unique marker for human acute myelogenous leukemia stem cells. *Leukemia* **14**, 1777–1784 (2000).

14. Majeti, R. *et al.* CD47 Is an Adverse Prognostic Factor and Therapeutic Antibody Target on Human Acute Myeloid Leukemia Stem Cells. *Cell* **138**, 286–299 (2009).

15. Jaiswal, S. *et al.* CD47 Is Upregulated on Circulating Hematopoietic Stem Cells and Leukemia Cells to Avoid Phagocytosis. *Cell* **138**, 271–285 (2009).

16. Kikushige, Y. *et al.* TIM-3 Is a Promising Target to Selectively Kill Acute Myeloid Leukemia Stem Cells. *Cell Stem Cell* **7**, 708–717 (2010).

17. van Rhenen, A. *et al.* The novel AML stem cell–associated antigen CLL-1 aids in discrimination between normal and leukemic stem cells. *Blood* **110**, 2659–2666 (2007).

18. Moshaver, B. *et al.* Identification of a Small Subpopulation of Candidate Leukemia-Initiating Cells in the Side Population of Patients with Acute Myeloid Leukemia. *Stem Cells* **26**, 3059–3067 (2008).

19. Goardon, N. *et al.* Coexistence of LMPP-like and GMP-like leukemia stem cells in acute

myeloid leukemia. *Cancer Cell* **19**, 138–152 (2011).

20. Blair, A., Hogge, D. E., Ailles, L. E., Lansdorp, P. M. & Sutherland, H. J. Lack of

Expression of Thy-1 (CD90) on Acute Myeloid Leukemia Cells With Long-Term

Proliferative Ability In Vitro and In Vivo. *Blood* **89**, 3104–3112 (1997).

21. Oren, R., Takahashi, S., Doss, C., Levy, R. & Levy, S. TAPA-1, the target of an

antiproliferative antibody, defines a new family of transmembrane proteins. *Mol Cell Biol*

10, 4007–15 (1990).

22. Shoham, T., Rajapaksa, R., Kuo, C.-C., Haimovich, J. & Levy, S. Building of the

Tetraspanin Web: Distinct Structural Domains of CD81 Function in Different Cellular

Compartments. *Mol Cell Biol* **26**, 1373–1385 (2006).

23. Yáñez-Mó, M., Barreiro, O., Gordon-Alonso, M., Sala-Valdés, M. & Sánchez-Madrid, F.

Tetraspanin-enriched microdomains: a functional unit in cell plasma membranes. *Trends*

in Cell Biology **19**, 434–446 (2009).

24. Hemler, M. E. Tetraspanin functions and associated microdomains. *Nat Rev Mol Cell Biol*

6, 801–811 (2005).

25. Umeda, R. *et al.* Structural insights into tetraspanin CD9 function. *Nat Commun* **11**, 1606

(2020).

26. Zimmerman, B. *et al.* Crystal Structure of a Full-Length Human Tetraspanin Reveals a

Cholesterol-Binding Pocket. *Cell* **167**, 1041-1051.e11 (2016).

27. Susa, K. J., Seegar, T. C., Blacklow, S. C. & Kruse, A. C. A dynamic interaction between

CD19 and the tetraspanin CD81 controls B cell co-receptor trafficking. *eLife* **9**, e52337

(2020).

28. Bradbury, L. E., Kansas, G. S., Levy, S., Evans, R. L. & Tedder, T. F. The CD19/CD21

signal transducing complex of human B lymphocytes includes the target of

antiproliferative antibody-1 and Leu-13 molecules. *J. Immunol.* **149**, 2841–2850 (1992).

29. Vences-Catalán, F. *et al.* CD81 is a novel immunotherapeutic target for B cell lymphoma.

J. Exp. Med. **216**, 1497–1508 (2019).

30. Quagliano, A., Gopalakrishnapillai, A., Kolb, E. A. & Barwe, S. P. CD81 knockout
promotes chemosensitivity and disrupts in vivo homing and engraftment in acute
lymphoblastic leukemia. *Blood Advances* **4**, 4393–4405 (2020).

31. Vences-Catalán, F. *et al.* Targeting the tetraspanin CD81 reduces cancer invasion and
metastasis. *Proc Natl Acad Sci U S A* **118**, (2021).

32. Boyer, T. *et al.* Tetraspanin CD81 is an adverse prognostic marker in acute myeloid
leukemia. *Oncotarget* (2016) doi:10.18632/oncotarget.11481.

33. Boyer, T. *et al.* Flow Cytometry to Estimate Leukemia Stem Cells in Primary Acute
Myeloid Leukemia and in Patient-derived-xenografts, at Diagnosis and Follow Up. *J Vis*
Exp (2018) doi:10.3791/56976.

34. Duployez, N. *et al.* The stem cell-associated gene expression signature allows risk
stratification in pediatric acute myeloid leukemia. *Leukemia* **33**, 348–357 (2019).

35. Nibourel, O. *et al.* Copy-number analysis identified new prognostic marker in acute
myeloid leukemia. *Leukemia* **31**, 555–564 (2017).

36. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* **9**, 671–675 (2012).

37. Zhang, C. C. *et al.* Gemtuzumab Ozogamicin (GO) Inclusion to Induction Chemotherapy
Eliminates Leukemic Initiating Cells and Significantly Improves Survival in Mouse Models
of Acute Myeloid Leukemia. *Neoplasia* **20**, 1–11 (2018).

38. R Core Team. R: A language and environment for statistical computing. (2022).

39. Tyner, J. W. *et al.* Functional genomic landscape of acute myeloid leukaemia. *Nature*
562, 526–531 (2018).

40. Cerami, E. *et al.* The cBio cancer genomics portal: an open platform for exploring
multidimensional cancer genomics data. *Cancer Discov* **2**, 401–404 (2012).

41. Gao, J. *et al.* Integrative analysis of complex cancer genomics and clinical profiles using
the cBioPortal. *Sci Signal* **6**, pl1 (2013).

42. Saland, E. *et al.* A robust and rapid xenograft model to assess efficacy of

chemotherapeutic agents for human acute myeloid leukemia. *Blood Cancer J* **5**, e297

(2015).

43. Shultz, L. D. *et al.* Human lymphoid and myeloid cell development in NOD/LtSz-scid IL2R

gamma null mice engrafted with mobilized human hemopoietic stem cells. *J. Immunol*

174, 6477–6489 (2005).

44. Dick, J. E. & Lapidot, T. Biology of Normal and Acute Myeloid Leukemia Stem Cells.

International Journal of Hematology **82**, 389–396 (2005).

45. Stelmach, P. & Trumpp, A. Leukemic stem cells and therapy resistance in acute myeloid

leukemia. *haematol* **108**, 353–366 (2023).

46. Vetrie, D., Helgason, G. V. & Copland, M. The leukaemia stem cell: similarities,

differences and clinical prospects in CML and AML. *Nat Rev Cancer* **20**, 158–173 (2020).

47. Thomas, D. & Majeti, R. Biology and relevance of human acute myeloid leukemia stem

cells. *Blood* **129**, 1577–1585 (2017).

48. Ng, S. W. K. *et al.* A 17-gene stemness score for rapid determination of risk in acute

leukaemia. *Nature* **540**, 433–437 (2016).

49. Mitchell, K. & Steidl, U. Targeting Immunophenotypic Markers on Leukemic Stem Cells:

How Lessons from Current Approaches and Advances in the Leukemia Stem Cell (LSC)

Model Can Inform Better Strategies for Treating Acute Myeloid Leukemia (AML). *Cold*

Spring Harb Perspect Med **10**, a036251 (2020).

50. Papaemmanuil, E. *et al.* Genomic Classification and Prognosis in Acute Myeloid

Leukemia. *N Engl J Med* **374**, 2209–2221 (2016).

51. Welch, J. S. *et al.* The Origin and Evolution of Mutations in Acute Myeloid Leukemia. *Cell*

150, 264–278 (2012).

52. Ding, L. *et al.* Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-

genome sequencing. *Nature* **481**, 506–510 (2012).

53. Parkin, B. *et al.* Clonal evolution and devolution after chemotherapy in adult acute

myelogenous leukemia. *Blood* **121**, 369–377 (2013).

54. Bross, P. F. *et al.* Approval summary: gemtuzumab ozogamicin in relapsed acute

myeloid leukemia. *Clin Cancer Res* **7**, 1490–6 (2001).

55. Maecker, H. T. & Levy, S. Normal lymphocyte development but delayed humoral immune

response in CD81-null mice. *J Exp Med* **185**, 1505–10 (1997).

56. Vences-Catalán, F. *et al.* A mutation in the human tetraspanin CD81 gene is expressed

as a truncated protein but does not enable CD19 maturation and cell surface expression.

J. Clin. Immunol. **35**, 254–263 (2015).

57. Paterson, E. K. & Courtneidge, S. A. Invadosomes are coming: new insights into function

and disease relevance. *FEBS J* **285**, 8–27 (2018).

58. Jacquemet, G., Hamidi, H. & Ivaska, J. Filopodia in cell adhesion, 3D migration and

cancer cell invasion. *Current Opinion in Cell Biology* **36**, 23–31 (2015).

59. Bari, R. *et al.* Tetraspanins regulate the protrusive activities of cell membrane.

Biochemical and Biophysical Research Communications **415**, 619–626 (2011).

60. Termini, C. M. & Gillette, J. M. Tetraspanins Function as Regulators of Cellular Signaling.

Front. Cell Dev. Biol. **5**, (2017).

61. Baum, C. M., Weissman, I. L., Tsukamoto, A. S., Buckle, A. M. & Peault, B. Isolation of a

candidate human hematopoietic stem-cell population. *Proc. Natl. Acad. Sci. U.S.A.* **89**,

2804–2808 (1992).

62. Vergez, F. *et al.* High levels of CD34+CD38low/-CD123+ blasts are predictive of an

adverse outcome in acute myeloid leukemia: a Groupe Ouest-Est des Leucemies Aigues

et Maladies du Sang (GOELAMS) study. *Haematologica* **96**, 1792–8 (2011).

63. Haubner, S. *et al.* Coexpression profile of leukemic stem cell markers for combinatorial

targeted therapy in AML. *Leukemia* **33**, 64–74 (2019).

64. Baer, M. R. High frequency of immunophenotype changes in acute myeloid leukemia at

relapse: implications for residual disease detection (Cancer and Leukemia Group B

Study 8361). *Blood* **97**, 3574–3580 (2001).

65. Bras, A. E. *et al.* CD123 expression levels in 846 acute leukemia patients based on

standardized immunophenotyping. *Cytometry* **96**, 134–142 (2019).

66. Hasezaki, T., Yoshima, T. & Mine, Y. Anti-CD81 antibodies reduce migration of activated

T lymphocytes and attenuate mouse experimental colitis. *Sci Rep* **10**, 6969 (2020).

67. Bailly, C. & Thuru, X. Targeting of Tetraspanin CD81 with Monoclonal Antibodies and

Small Molecules to Combat Cancers and Viral Diseases. *Cancers* **15**, 2186 (2023).

68. Vexler, V. *et al.* Target-mediated drug disposition and prolonged liver accumulation of a

novel humanized anti-CD81 monoclonal antibody in cynomolgus monkeys. *MAbs* **5**, 776–

786 (2013).

69. Miyazaki, T. Normal development but differentially altered proliferative responses of

lymphocytes in mice lacking CD81. *The EMBO Journal* **16**, 4217–4225 (1997).

70. Tsitsikov, E. N., Gutierrez-Ramos, J. C. & Geha, R. S. Impaired CD19 expression and

signaling, enhanced antibody response to type II T independent antigen and reduction of

B-1 cells in CD81-deficient mice. *Proc. Natl. Acad. Sci. U.S.A.* **94**, 10844–10849 (1997).

71. Van Zelm, M. C. *et al.* CD81 gene defect in humans disrupts CD19 complex formation

and leads to antibody deficiency. *J. Clin. Invest.* **120**, 1265–1274 (2010).