1
2
3

The Kinetic and Energetic Pull of Chemical Entropy

On Small Ensembles of Proteins

Josh E. Baker[1*]

[1]University of Nevada, Reno School of Medicine: Reno, NV 89521 USA

*Corresponding author. Email: jebaker@unr.edu

Abstract: Proteins in biological systems function at the interface of single molecule and bulk chemistry and thus provide novel insights into the basic physical chemistry of scaling and emergent phenomena. For example, a binary mechanical model that accounts for many energetic and mechanical aspects of muscle contraction provides a novel and testable framework for investigating the relationship between molecular mechanics and chemical thermodynamics in single molecules and small molecular ensembles. The problem addressed here is that the model and supporting experimental data require that entropic forces balance chemical reactions. While this is consistent with statistical mechanics, it is at odds with basic chemistry. Specifically, counter to a classic chemical activity analysis, the mere presence of molecules (chemical activity or mass action) does not physically push a reaction toward equilibrium; rather, the number of microstates, Ω, accessible in a chemical state pulls a reaction down an entropic funnel of increasing Ω toward equilibrium. Here I develop an entropic model of chemical thermodynamics and kinetics and compare it with conventional chemical activity models. I show that an a priori system reaction energy landscape fully describes the chemical kinetics of both equilibrium and non-equilibrium chemical reactions, formally justifying the thermodynamic rate constants required for binary mechanical models of muscle contraction.

28   **Keywords**: Entropy, Chemical Thermodynamics, Free Energy, Reaction Energy Landscape,

29   Microstates, Micropathways, Non-Equilibrium, Kinetics, Entropic Funnel

Introduction

Technological advances over the past 70 years have fueled a molecular revolution in the

biological sciences, providing remarkably detailed descriptions of the structure and function of a

myriad of biomolecules. The question remains, to what extent is biological function the sum of

these molecular parts versus an emergent property of systems of molecules? While this is

fundamentally a physical chemical question, insights from small molecular ensembles in

biological systems are informing new perspectives on this important chemical thermodynamic

problem. For example, a binary mechanical model that describes the mechanochemistry of

muscle contraction offers a new perspective on entropic forces in enzyme-catalyzed chemical

reactions [1,2]. This model provides a clear description of the mechanisms required to contain

entropic forces, providing an explicit solution to the Gibbs paradox [3]. The problem addressed

here is that the model and supporting experimental data require that chemical reactions are

balanced by entropic forces [2]. While this is consistent with the entropy of binary systems in

statistical mechanics, it is inconsistent with the conventional assumption in chemistry that

chemical reactions are balanced – not by entropy, but – by forces exerted by the mere physical

presence of molecules (through chemical activity or mass action). Here, I compare an entropic

model of chemical thermodynamics with the conventional chemical activity model and show that

while the reaction free energy equations have the same form, the physical meaning of the

"concentration" terms differ fundamentally. In an entropic model an a priori system reaction

energy landscape fully describes the chemical kinetics of both equilibrium and non-equilibrium

chemical reactions, formally justifying the thermodynamic rate equations required in a binary

mechanical model of muscle contraction [2].

Figure 1 illustrates a binary mechanical system [2]. The molecular mechanism is a

reversible molecular switch (Fig. 1A) that displaces an element external to the switch a distance,

d, with a transition from state A to B. If the element is stiff, the molecular switch generates force.

An equilibrium ensemble of molecular switches constitutes a binary mechanical system (Fig. 1B)

within which a Boltzmann distribution of chemical states, N_B/N_A, decreases exponentially with a

force, F, applied to the system as

$N_B/N_A = \exp[-(\Delta G° + F \cdot d)/k_B T]$ Eq. 1

where N_A and N_B are the number of molecules in states A and B, and $\Delta G°$ is the standard free

energy for a molecular transition from state A to B.

In muscle, the molecular switch is a myosin motor (oval) that generates force through a

discrete conformational change induced by actin binding (solid black line) and driven

energetically by the binding free energy, $\Delta G°$. We have directly observed in mechanics

experiments on systems ranging from single myosin motors to single muscle cells that myosin

motors function as reversible molecular switches [4–6]. We have also shown in muscle fibers

that the force-dependence of the distribution between the two states of the molecular switch

follows Eq. 1 [5,7]. Moreover we have shown that this binary mechanical model accurately

accounts for the basic mechanics and energetics of muscle contraction [1,2].

According to Eq. 1, when $\Delta G° = 0$,

$F = -k_B T \cdot \ln(N_B/N_A)/d.$ Eq. 2

Because the passive force exerted on binary mechanical systems (e.g., muscle, rubber, titin) is

entropic [2,8,9], $k_B T \cdot \ln(N_B/N_A)$ is entropic energy. While this is a classic statistical mechanic

description of a binary system [10], it is at odds with conventional chemistry where the term

$k_B T \cdot \ln(N_B/N_A)$ is assumed to be part of a chemical potential not entropy. Thus, statistical

mechanics and chemistry offer two fundamentally different descriptions of the term

$k_B T \cdot \ln(N_B/N_A)$. In conventional chemistry, N_A and N_B are numbers of molecules that physically

exist in states A and B (chemical activities), which through their mere presence physically push a

reaction (mass action) toward equilibrium. In an entropic analysis, N_A and N_B describe a system

reaction coordinate $\{N_A, N_B\}$ that is defined independent of the physical presence of molecules in

states A and B. Rather than being pushed by physical quantities (or a mass) of molecules, in an

entropic analysis the reaction is pulled toward states of higher system entropy.

While this problem is highlighted by the entropic force of a binary mechanical system

(Eq. 2), it is not limited to mechanical systems. Here, I show that even in non-mechanical ($F = 0$)

chemical systems, $k_B T \cdot \ln(N_B/N_A)$ is entropic energy not a chemical potential. This represents a

radical transformation of our basic understanding of chemical energetics and kinetics (the model

90   implies that the physical presence of molecules, N_A and N_B, has no effect on the kinetics or

91   energetics of a chemical reaction) and leads to an entropic formulation of chemical kinetics in

92   which non-equilibrium kinetics are fully defined by an a priori system reaction energy landscape.

Reaction Free Energy

In a given system state $\{N_A,N_B\} = \{\delta\eta,N–\delta\eta\}$, the free energy, G, for the reaction in Fig.

1B is expressed in terms of the extent of the reaction, $\delta\eta$, as

$G = (\mu_B – \mu_A)\cdot\delta\eta + F\cdot d\cdot\delta\eta – TS,$ Eq. 3

where $(\mu_B – \mu_A)$ is the chemical potential difference between states A and B, $\underline{F}\cdot d\cdot\delta\eta$ is the net

work performed by $\delta\eta$ displacements, d, of a system spring with a force, $\underline{F}$, and S is the system

entropy in state$\{\delta\eta,N–\delta\eta\}$. When one molecule transitions from state A to B the change, Δ, in

system energy is described by the reaction free energy

$\Delta_rG = \delta G/\delta\eta = (\mu_B – \mu_A) + F\cdot d – T\Delta S,$ Eq. 4

where ΔS is the reaction entropy, $\delta S/\delta\eta$. The terms on the right hand side of Eq. 4 are the

chemical forces that when $\Delta_rG < 0$ energetically drive a reaction from $\{N_A,N_B\}$ to $\{N_A–1,N_B+1\}$,

or when $\Delta_rG > 0$ energetically drive a reaction from $\{N_A–1,N_B+1\}$ to $\{N_A,N_B\}$, or when $\Delta_rG = 0$

balance the reaction.

Early chemists were averse to the concept of a reaction entropy, ΔS, in part because of

their limited mathematical backgrounds but also because they did not understand Gibbs [11].

Instead, with a more Newtonian worldview, they were convinced by the general arguments of

Waage and Guldberg in 1864 that the mere presence (number or mass) of molecules in a state i

physically pushes molecules out of that state through a mechanism often referred to as mass

action [11]. Because this mechanism needed an energetic basis, chemists defined the chemical

potential, μ_i, of chemical state, i, as a function of the number of molecules in that state,

$\mu_i = \mu°_i + k_BT\cdot\ln N_i$, such that when N_i increases μ_i increases, energetically pushing molecules out

state i. According to this analysis, $\mu_B – \mu_A = \Delta G° + k_BT\cdot\ln(N_B/N_A)$, where $\Delta G°$ is the standard

free energy, which when substituted into Eq. 4 describes an equilibrium force, $\underline{F}$, in Eq. 2 that is

balanced against the force of mass action, $\mu_B - \mu_A$, not entropy, ΔS.

While entropic contributions to the standard free energy, $\Delta G° = \Delta H° - T\Delta S°$, are often

considered, $\Delta S°$ differs from ΔS. The standard entropic change, $\Delta S°$, describes the entropic

change of a molecule with an A to B transition, whereas the reaction entropy, ΔS, describes the

entropic change of a system of molecules with a $\{N_A,N_B\}$ to $\{N_A-1,N_B+1\}$ transition. The reaction

entropy, ΔS, continues to be ignored by chemists despite the fact that the force exerted on a

binary system (Eq. 2) is entropic, $\Delta S = -k_B \cdot \ln(N_B/N_A)$.

In both a chemical activity and entropic analysis, the free energy equation (Eq. 4) is

$\Delta_r G = \Delta G° + F \cdot d + k_B T \cdot \ln(N_B/N_A).$ Eq. 5

The question at hand is whether $k_B T \cdot \ln(N_B/N_A)$ is a component of a chemical potential, $\mu_B - \mu_A$,

where N_A and N_B describe the physical presence (chemical activity or mass) of molecules in

states A and B, or whether $k_B T \cdot \ln(N_B/N_A)$ is a system entropic energy, $T\Delta S$, where N_A and N_B

describe an a priori system reaction coordinate $\{N_A,N_B\}$ that is defined independent of the

physical presence of molecules.

The free energy change, $\Delta G°$, for a molecular transition from state A to B

The standard free energy, $\Delta G°$, for a reaction is a molecular property. For example, $\Delta G°$

for the hydrolysis of ATP to ADP and P_i can be physically defined for one molecule as the

change in free energy associated with breaking a single phosphoanhydride bond.

Figure 2A illustrates an energy landscape for a single molecule that reversibly switches between

molecular states A and B with forward and reverse rate constants, $k°_+$ and $k°_-$, and a free energy

difference of $\Delta G°$ [12]. This energy landscape describes changes in free energy along the

reaction (or conformational) coordinate of a molecule [13]. According to kinetic theory, the

inverse time to diffuse in both forward, $k°_+$, and reverse, $k°_-$, directions over an activation energy

barrier separating A and B (Fig. 2A) are related to $\Delta G°$ as $k°_+/k°_- = \exp(-\Delta G°/k_B T)$ [14]. Thus,

for an isolated single molecule, the ratio of the time-averaged occupancy of molecular states B

and A is

$N_B/N_A = \exp(-\Delta G°/kT).$ Eq. 6

This equation describes an energetic balance between ΔG° and the energy term $-$

$k_B T \cdot \ln(N_B/N_A)$, which evident from this example does not describe a chemical activity (mass

action) since there is only one molecule diffusing in the molecular energy landscape. While this

energy term emerges from the molecular energy landscape, it is clearly not part of it. It is an

ensemble, system energy that is defined independent of the physical presence of molecules in

states A and B (Eq. 6). As proposed by T.L. Hill [15], when one molecule undergoes a transition

from A to B, the only contribution of that molecular transition to the system free energy

$(\mu_B - \mu_A)$ is ΔG° not $\Delta G^\circ + k_B T \cdot \ln(N_B/N_A)$.

Free energy changes for system transitions from chemical state $\{N_A, N_B\}$ to $\{N_A-1, N_B+1\}$

For an ensemble of N such molecules contained in a solution held at constant T, P, V, and

$F = 0$, the reaction occurs within a system energy landscape along a system reaction coordinate,

$\{N_A, N_B\}$ (Fig. 2B). System energy landscapes can be plotted as changes in the reaction free

energy of the system, $\Delta_r G$, (Eq. 4, Fig. 3A) or as changes in the free energy of the system, G,

(Eq. 3, Fig. 3B) both along the same system reaction coordinate, $\{N_A, N_B\}$ [10,16]. In addition to

the incremental molecular free energy changes, ΔG°, that contribute to the system free energy

with each molecular step from A to B, $\Delta_r G$ accounts for non-molecular (non-ΔG°) changes in the

free energy of the system with each chemical step from $\{N_A, N_B\}$ to $\{N_A-1, N_B+1\}$. These system

contributions to free energy include PV work, non-PV work ($F \cdot d$) and ΔS in Eq. 4.

Reaction (system) entropy, ΔS

Using elementary statistical mechanic arguments, the reaction entropy, ΔS , for any

reversible two-state reaction (Figs. 1 and 2) is shown below to be

$\Delta S = -k_B \cdot \ln[(N_B + 1))/(N_A)]$ for molecular transitions and

$\Delta S = -R \cdot \ln([B]/[A])$ for molar transitions,

resulting in molecular and molar reaction free energy equations ($\Delta_r G = \Delta G^\circ - T\Delta S$)

$\Delta_r G = \Delta G^\circ + k_B T \cdot \ln[(N_B + 1))/(N_A)]$ and   Eq. 7

$\Delta_r G = \Delta G^\circ + R T \cdot \ln([B]/[A])$.

While these equations are consistent with a conventional chemical activity analysis, here

N_A, N_B, [A], and [B] are unrelated to chemical activities. That is, they are not physical quantities

of molecules that through their mere presence (mass action) balance chemical reactions. Clear

from the analysis below, N_A, and N_B are system reaction coordinates that describe the reaction

entropy, $\Delta S(N_A,N_B)$ in chemical state $\{N_A,N_B\}$; and [A] and [B] are molar reaction coordinates

that describe the molar reaction entropy, $\Delta S([A],[B])$ in chemical state $\{[A],[B]\}$.

184   At equilibrium, $k_B T \cdot \ln(N_B/N_A) = -\Delta G°$ not because $N_A + N_B$ molecules equilibrate within

185   the energy landscape of one molecule [i.e., when $\mu_B - \mu_A = 0$], but because N_A and N_B describe

186   the point along the system reaction coordinate, $\{N_A,N_B\}$, at which the system equilibrates [i.e.,

187   the point at which $T\Delta S(N_A,N_B) = -\Delta G°$]. This is illustrated in Figs. 1B and 2 for small numbers

188   of molecules and derived in general as follows.

189   According to Boltzmann, the system entropy in state $\{N_A,N_B\}$ is $S = k_B \cdot \ln(\Omega_{NA,NB})$, where

190    $\Omega_{NA,NB}$ is the number of microstates available to the system in chemical state $\{N_A,N_B\}$.

191   Specifically, $\Omega_{NA,NB} = N!/(N_A!N_B!)$. The change in entropy with a single chemical step

192   from $\{N_A,N_B\}$ to $\{N_A-1,N_B+1\}$ is then

193            $\Delta S = -k_B \cdot \ln[((N_A - 1)!(N_B + 1)!)/(N_A!N_B!)] = -k_B \cdot \ln[(N_B + 1))/(N_A)]$.

195   The change in system entropy associated with $\delta\eta$ chemical steps from $\{N_A,N_B\}$ to

196    $\{N_A - \delta\eta,N_B+\delta\eta\}$ is

197            $\Delta S = -k_B \cdot \ln[((N_A - \delta\eta)!(N_B + \delta\eta)!)/(N_A!N_B!)] \approx -\delta\eta \cdot k_B \cdot \ln[(N_B + \delta\eta/2))/(N_A - \delta\eta/2)]$,

199   and so

200            $\Delta_r G = \delta\eta \cdot \Delta G° + \delta\eta \cdot k_B T \cdot \ln[(N_B + \delta\eta/2)/(N_A - \delta\eta/2)]$.   Eq. 8

202   Here the reaction is driven both by the change in molecular free energy, $\cdot \Delta G°$, summed

203   over $\delta\eta$ molecular transitions from A to B and by the average change in system entropy,

204    $k_B T \cdot \ln[(N_B + \delta\eta/2)/(N_A - \delta\eta/2)]$, summed over $\delta\eta$ chemical steps from $\{N_A,N_B\}$ to

205    $\{N_A - \delta\eta,N_B+\delta\eta\}$. The change in entropy, ΔS, relative to $S = 0$ occurs from $\{N/2,N/2\}$ to

206    $\{N/2-\delta\eta, N/2+\delta\eta\}$, or

208            $\Delta S = -\delta\eta \cdot k_B \cdot \ln[(N/2 + \delta\eta/2))/(N/2 - \delta\eta/2)$,

or

$\Delta S = -\delta\eta \cdot k_B \cdot \ln[(1 + \delta\eta/N)/(1 - \delta\eta/N)]$,   Eq. 9

where $\delta\eta/N$ is a fractional change in the extent of the reaction relative to unity. Assuming

$\delta\eta = 1$ mol, Eq. 9 can be written in its molar form

$\Delta S = -R \cdot \ln([B]/[A])$.

Entropic changes along a system reaction coordinate.

Figure 2B illustrates entropic changes along the reaction coordinate for a system of $N = 5$

molecules each occupying either molecular state A or B. When the system is in state {5,0},

$\Omega_{5,0} = 1$. Unlike in chemical activity models where $k_B \cdot \ln(0/5)$ is undefined, here the system

entropy, $k_B \cdot \ln(1)$, is zero. After one net forward step, the system enters state {4,1}, increasing the

number of microstates from 1 to 5 and increasing the system entropy from 0 to $k_B \cdot \ln(5)$. This

increase in the number of microstates physically pulls the reaction forward since there are five-

times more micropathways in the forward direction than in the reverse direction. This contrasts

with a chemical activity of molecules that purportedly pushes a reaction forward. When $\Delta G^o = 0$,

the reaction continues until a maximum system entropy is reached along the reaction coordinate,

equilibrating in system state {3,2} when $(N_B + 1)/N_A = \exp(0)$ (Eq. 7).

Figure 3A is a plot of the reaction free energy (Eq. 5) along the system reaction

coordinate, $\{N_A, N_B\}$, for $N = 10$ molecules. The reaction entropy, $k_B T \cdot \ln[N_B+1, N_A]$, increases

logarithmically with each step whereas the molecular free energy, ΔG^o, is a constant offset (it is

the same for each step). Figure 3A shows that when ΔG^o decreases from 0 to $-1.5 \ k_B T$ (down

arrow) the equilibrium state of the system changes from {5.5,4.5} to {2,8} (right arrow)

corresponding to a change from $(N_B + 1)/N_A = \exp(0)$ to $(N_B + 1)/N_A = \exp(1.5)$.

Because $\Delta_r G$ is the change in free energy, G, of the system with a chemical step (Eqs. 3 and 4),

the free energy, G, of the system can be obtained by integrating Eq. 4 over the reaction

coordinate. In Fig. 3B, the two plots in Fig. 3A are integrated and replotted as free energy, G.

These are the system free energy landscapes described by Gibbs within which the reaction

equilibrates at a point along the reaction coordinate $\{N_A, N_B\}$ where G is a minimum. Figure 3

clearly shows that entropic forces, $T\Delta S$, (Fig. 3A) drive a reaction down an entropic energy

funnel, TS (Fig. 3B) toward equilibrium.

Figure 3B illustrates how changes in $\Delta G°$ affect the free energy, G, landscape. Changes

in N_A and N_B, however, have no effect on the energy landscape because they are reaction

coordinates, not chemical activities. In this way, an entropic chemical analysis radically

transforms our understanding of chemical kinetics and energetics. Because chemical kinetics and

energetics are defined by free energy landscapes, the physical presence of molecules, N_A and N_B,

has no effect on the kinetics or energetics of a chemical reaction.

Equilibrium Energetics

At equilibrium ($\Delta_r G = 0$), the probability of finding the system in state $\{N_A+1,N_B–1\}$

relative to $\{N_A,N_B\}$ is $\exp[–\Delta G°/k_B T + \ln(\Omega_{NA-1,NB+1}/\Omega_{NA,NB})]$, where

$\ln(\Omega_{NA-1,NB+1}/\Omega_{NA,NB}) = T\Delta S/k_B T$. Thus, when $\Delta G° = 0$, $\Omega_{NA,NB}$ is the probability density

function, $P_{NA,NB}$. For comparison, according to chemical activity models the probability of

finding a molecule in state B relative to state A is $\exp[–\Delta G°/k_B T]$, and when $\Delta G° = 0$, $P_{NA,NB}$ is a

binomial distribution. The equilibrium constant, $K = (\Omega_{NA,NB}/\Omega_{NA-1,NB+1})_{eq} = \exp[–\Delta G°/k_B T]$ is

$K = (N_{B(eq)} + 1)/N_{A(eq)}$ and describes the point along the reaction coordinate $\{N_A,N_B\}_{eq}$ at which

the system equilibrates (Fig. 3). For comparison, in chemical activity models the equilibrium

constant $K = N_{B(eq)}/N_{A(eq)}$ describes the physical distribution of N molecules within the energy

landscape of a single molecule (Fig. 2A).

Non-Equilibrium Energetics

When the system is perturbed from equilibrium by a change in the internal energy of the

system, δE, [i.e., a change in $\Delta G°$ or $k_B T \cdot \ln[(\Omega_{NA-1,NB+1}/\Omega_{NA,NB})$], the free energy equation

becomes

$\Delta_r G = [\Delta G° – k_B T \cdot \ln(\Omega_{NA-1,NB+1}/\Omega_{NA,NB})]_{eq} + \delta E = \delta E,$

where δE is a non-equilibrium (ne) perturbation to any of the energy term on the right-hand side

of Eq. 5. Here,

$\delta E = [\Delta G° – k_B T \cdot \ln(\Omega_{NA-1,NB+1}/\Omega_{NA,NB})]_{ne}.$

As illustrated in Fig. 3, if the system is perturbed by a change in $\Delta G°$, the system relaxes

to a new equilibrium state, $\{N_A,N_B\}_{eq}$, as $k_B T \cdot \ln(\Omega_{NA-1,NB+1}/\Omega_{NA,NB})$ approaches the new $\Delta G°$.

Through this process, $\Delta_r G$ returns to zero and δE remains in the system in the form of a change

in system entropy. If the system perturbation is a change in $k_B T \cdot \ln(\Omega_{NA-1,NB+1}/\Omega_{NA,NB})$ [an

irreversible transfer of molecules between molecular states] the system relaxes back to the

original equilibrium state, $\{N_A, N_B\}_{eq}$, as $k_B T \cdot \ln(\Omega_{NA-1,NB+1}/\Omega_{NA,NB})_{ne}$ approaches ΔG°. Through

this process the entropic δE is lost from the system as heat. This analysis becomes more complex

when non-$P\Delta V$ work is performed, in which case, in addition to exchanges between

$k_B T \cdot \ln(\Omega_{NA-1,NB+1}/\Omega_{NA,NB})$ and ΔG°, δE can be exchanged with internal mechanical potentials;

and in addition to being lost from the system as heat or stored in the system as entropy, δE can

be lost from the system as work performed on the surroundings [2].

Equilibrium Kinetics

Chemical kinetics, like energetics, have both molecular and system components. The net

rate at which the system transitions from chemical state $\{N_A, N_B\}$ to chemical state $\{N_A-1, N_B+1\}$

is

$d\{N_A, N_B\}/dt = k_+ - k_-,$

where k_+ and k_- are the forward and reverse transition rates between these states. At equilibrium,

the probability of finding the system in state $\{N_A+1, N_B-1\}$ relative to state $\{N_A, N_B\}$ is

$k_{+(eq)}/k_{-(eq)} = \exp[(-\Delta G^\circ + k_B T \cdot \ln[\Omega_{NA-1,NB+1}/\Omega_{NA,NB}]_{eq})/k_B T]$ or

$k_{+(eq)}/k_{-(eq)} = (k^\circ_+/k^\circ_-) \cdot \exp[k_B T \cdot \ln(\Omega_{NA-1,NB+1}/\Omega_{NA,NB})_{eq}/k_B T].$

Here, k°_+ and k°_- are molecular contributions to k_+ and k_- and $\exp[-T\Delta S/k_B T]$ is the system

contribution to k_+ and k_-. The influence of $-\Delta S/k_B$ on kinetics can be understood energetically as

a tilt of the system landscape that adds to the tilt of the molecular landscape, ΔG°. It can also be

understood kinetically as

$k_{+(eq)}/k_{-(eq)} = (k^\circ_+/k^\circ_-) \cdot (\Omega_{NA-1,NB+1}/\Omega_{NA,NB})_{eq}$

where the factor $\Omega_{NA-1,NB+1}/\Omega_{NA,NB}$ describes the number of micropathways available for the

forward reaction relative to the reverse reaction. For example, in Fig. 2B there are five

micropathways from {5,0} to {4,1} and one micropathway back from {4,1} to {5,0}, which

means that the forward reaction is entropically five-fold more likely than the reverse reaction

($\Omega_{NA-1,NB+1}/\Omega_{NA,NB} = 5$). This implies forward and reverse rate constants of

$k_{+(eq)} = k°_{+} \cdot \Omega_{NA-1,NB+1(eq)}$

$k_{-(eq)} = k°_{-} \cdot \Omega_{NA,NB(eq)}.$

At equilibrium, $k_{+(eq)} = k_{-(eq)}$, or

$k°_{+} \cdot \Omega_{NA-1,NB+1(eq)} = k°_{-} \cdot \Omega_{NA,NB(eq)}.$

To demonstrate consistency with the chemical activity approach,

$k_{+(eq)} = k°_{+} \cdot N_{A(eq)}$

$k_{-(eq)} = k°_{-} \cdot (N_{B(eq)} + 1)$

where here again $N_{A(eq)}$ and $N_{B(eq)}$ are not physical, active numbers of molecules, they are the

system reaction coordinate $\{N_A,N_B\}_{eq}$ at which the system equilibrates.

Non-Equilibrium Kinetics

Chemical kinetics is fully determined from the molecular and system tilt of the system

energy landscape. It follows that non-equilibrium kinetics is fully determined by the non-

equilibrium tilt, δE, of these landscapes, where a non-equilibrium perturbation to $\Delta G°$ affects the

molecular rates $k°_{+}/k°_{-}$, and an irreversible transfer of molecules between molecular states affects

$\Omega_{NA,NB}/\Omega_{NA-1,NB+1}$. In general,

$k_{+}(t)/k_{-}(t) = (k°_{+}/k°_{-})(\Omega_{NA-1,NB+1}(t)/\Omega_{NA,NB}(t)) = \exp[-\delta E(t)/k_B T].$ Eq. 10

According to Eq. 10, when $\delta E(t) < 0$, $k_{+}(t) > k_{-}(t)$, consistent with the requirement that a

negative $\delta E(t)$ drives the reaction forward. This forward reaction increases $T\Delta S$ (Fig. 3) until

$\delta E(t) = 0$, at which point $k_+(t) = k_-(t)$. When $\delta E(t) > 0$, $k_+(t) > k_-(t)$, consistent with the

requirement that a positive $\delta E(t)$ drives the reaction backward. This reaction decreases $T\Delta S$ (Fig.

3) until $\delta E(t) = 0$, at which point $k_+(t) = k_-(t)$.

The non-equilibrium rate equation is the same as the equilibrium rate equation

$d\{N_A,N_B\}/dt = k°_+\cdot\Omega_{NA-1,NB+1} - k°_-\cdot\Omega_{NA,NB}(t)$.

For comparison with a chemical activity analysis, from above

$dN_A/dt = k°_+\cdot N_A(t) - k°_-\cdot(N_B(t) + 1)$ Eq. 11

where one last time, $N_A(t)$ and $N_B(t)$ describe the time dependence of steps along the system

reaction coordinate $\{N_A,N_B\}$, not the time-dependence of the physical numbers of molecules in

different states. Changes in $N_B(t) + 1$ are equal and opposite changes in $N_A(t)$ along the system

reaction coordinate, and so

$dN_A/dt = -N_A(t)\cdot[k°_- + k°_+]$

which means that $N_A(t)$ decreases exponentially with time as

$N_A(t) = N_{Ao}\cdot\exp(-(k°_+ + k°_-)t)$,

with a relaxation rate, $k°_+ + k°_-$, where N_{Ao} is the initial position on the reaction coordinate

$\{N_A,N_B\}_o$.

Conclusion

In summary, a molecular energy landscape (Fig. 2A) a priori defines $\Delta G°$ and rate

constants, $k°_+$ and $k°_-$, for an isolated molecule (Fig. 2A) independent of the physical presence of

that molecule [13]. Similarly, an entropic energy landscape (Fig. 2B) a priori defines system

energetics and rates, k_+ and k_-, for N molecules (Figs. 2B and 3) independent of the physical

presence of those molecules. Together $\Delta G°$ and system entropy create a system energy landscape

within which a reaction is energetically pulled down an entropic funnel (Fig. 3) until at

equilibrium entropic forces are balanced against $\Delta G°$ (Eq. 5).

I have shown that non-equilibrium chemical kinetic equations (Eq. 10) can be derived

from chemical thermodynamics, and that these equations have the same form as kinetic

equations shown to accurately describe non-equilibrium kinetics. Claims that non-equilibrium

kinetic equations cannot be derived from chemical thermodynamics are based on a chemical

activity analysis in which N_A and N_B are not coordinates along the reaction energy landscape but

are instead physical quantities of molecules that through their mere presence balance chemical

reactions.

In addition to ΔS, other non-molecular (non-$\Delta G°$) free energies that tilt the system

reaction free energy landscape are easily incorporated into Eq. 10. For example, we have added

the work term, $F(t)d$, to $\delta E(t)$ in Eq. 10, and have shown that simulations of a binary mechanical

model that incorporate these non-equilibrium kinetic rates accurately describe both the transient

and steady state mechanics of active muscle. Under near-equilibrium conditions, these

simulations converge with the thermodynamic relationship between $\Delta G°$, N_A, N_B, and F

described by Eq. 5 at $\Delta_r G = 0$. In short, the analysis presented herein provides a transformative

entropic interpretation of reaction free energies that unites chemical kinetics and

thermodynamics.

References and Notes

[1]   J. E. Baker and D. D. Thomas, *A Thermodynamic Muscle Model and a Chemical Basis for A.V. Hill's Muscle Equation*, J Muscle Res Cell Motil. **21**, 335 (2000).

[2]   J. E. Baker, *Thermodynamics and Kinetics of a Binary Mechanical System: Mechanisms of Muscle Contraction*, Langmuir **38**, 15905 (2022).

[3]   J. E. Baker, *Cells Solved the Gibbs Paradox by Learning How to Contain Entropic Forces*, Arxiv **arXiv:2305**, (2023).

[4]   J. E. Baker, I. Brust-Mascher, S. Ramachandran, L. E. LaConte, and D. D. Thomas, *A Large and Distinct Rotation of the Myosin Light Chain Domain Occurs upon Muscle Contraction.*, Proc. Natl. Acad. Sci. U. S. A. **95**, 2944 (1998).

[5]   J. E. Baker, L. E. W. LaConte, I. Brust-Mascher, and D. D. Thomas, *Mechanochemical Coupling in Spin-Labeled, Active, Isometric Muscle.*, Biophys. J. **77**, 2657 (1999).

[6]   J. E. Baker, C. Brosseau, P. B. Joel, and D. M. Warshaw, *The Biochemical Kinetics Underlying Actin Movement Generated by One and Many Skeletal Muscle Myosin Molecules.*, Biophys. J. **82**, 2134 (2002).

[7]   J. E. Baker and D. D. Thomas, *Thermodynamics and Kinetics of a Molecular Motor Ensemble.*, Biophys. J. **79**, 1731 (2000).

[8]   J. C. Phys and D. E. Hanson, *The Molecular Kink Paradigm for Rubber Elasticity : Numerical Simulations of Explicit Polyisoprene Networks at Low to Moderate Tensile Strains of Explicit Polyisoprene Networks at Low to Moderate Tensile Strains*, **135**, 054902 (2011).

[9]   M. S. Kellermayer, S. B. Smith, H. L. Granzier, and C. Bustamante, *Folding-Unfolding Transitions in Single Titin Molecules Characterized with Laser Tweezers*, **276**, 1112 (1997).

[10]   C. Kittel and H. Kroemer, *Thermal Physics* (1980).

[11]   J. Quílez, *On the Early Thermodynamic and Kinetic Deductions of the Equilibrium Constant Equation*, Found. Chem. **23**, 85 (2021).

[12]   L. Styer, *Biochemistry* (W.H. Freeman and Company, New York, 1995).

[13]   H. Frauenfelder, S. G. Sligar, P. G. Wolynes, H. Frauenfelder, S. G. Sligar, and P. G. Wolynes, *The Energy Landscapes and Motions of Proteins*, **254**, 1598 (1991).

[14]   J. Howard, *Mechanics of Motor Proteins and the Cytoskeleton*, 1st ed. (Sinauer Associates, 2001).

[15]   T. L. Hill, *Free Energy Transduction and Biochemical Cycle Kinetics* (Springer-Verlag,

New York, 1989).

[16]   J. Gibbs, *Elementary Principles in Statistical Mechanics Developed with Especial*

Reference to the Rational Foundation of Thermodynamics (1902).

Acknowledgments: I thank LEB, JWG, Julie, and my students, colleagues, and mentors who

have over many years inspired and guided this work. This work was funded by a grant from the

National Institutes of Health 1R01HL090938-01.

Funding: JEB was funded by a grant from the National Institutes of Health 1R01HL090938.

Competing interests: Author declares that they have no competing interests.

Data and materials availability: All data are available in the main text.

Fig. 1. Entropic Force in a Binary Mechanical System. (A) A reversible molecular switch undergoes a discrete conformational change from chemical state A to B displacing compliant elements external to the switch a distance, *d*. The switch occurs with forward and reverse rates, k°_{+} and k°_{-}. **(B)** An ensemble of molecular switches constitutes a binary mechanical system where, according to Le Chatelier's principle, a force applied to the system reverses the force generating chemistry of the molecular switches.

Fig. 2. The molecular and system contributions to the reaction free energy. (A) The

molecular contribution to the reaction free energy is described by a molecular energy landscape.

The free energy difference between the two metastable states, A and B, is ΔG°. The rate, k°_+, at

which a molecule transitions from molecular state A to B varies exponentially with $-G^\ddagger_+$, where

$G^\ddagger_+$ is the height of the activation barrier in the forward direction. The rate, k°_-, at which a

molecule transitions from B to A varies exponentially with $-G^\ddagger_-$, where $G^\ddagger_-$ is the height of the

activation energy barrier in the reverse direction. **(B)** The system contribution to the reaction free

energy is entropic. For $N = 5$, there are 6 system states, $\{N_A, N_B\}$, along the reaction coordinate

each with $\Omega_{NA,NB} = N!/(N_A! N_B!)$ microstates, illustrated with circles, and entropy $k_B \cdot \ln(\Omega_{NA,NB})$.

There are five micropathways (right arrows) by which a molecule in state $\{5,0\}$ can transition to

state $\{4,1\}$, and one micropathway (left arrow) by which a molecule in state $\{4,1\}$ can transition

to state $\{5,0\}$.

A

$G^{\ddagger}_{+}$

$G^{\ddagger}_{-}$

ΔG°

A k°_{+} B

k°_{-}

B

A:	5	4	3	2	1	0
B:	0	1	2	3	4	5
Ω:	1	5	10	10	5	1

Reaction Coordinate

Fig. 3. **Change in Free energy along a reaction coordinate. (A)** For N = 10, Eq. 3 is plotted for a $\Delta G°$ of 0 (green line) and -1.5 k_BT (blue line). For $\Delta G° = 0$, the reaction equilibrates ($\Delta_r G = 0$) at {5.5,4.5}. When $\Delta G°$ is decreased from 0 to -1.5 k_BT (down arrow) the system re-equilibrates (right arrow) at {2,8}. **(B)** The system free energy, G, (the integral of $\Delta_r G$ over the reaction coordinate) is calculated and plotted for $\Delta G°$ of 0 (green line) and -1.5 k_BT (blue line) values in panel A. The same trajectory in panel A describing re-equilibration following a decrease in $\Delta G°$ is illustrated in panel B (arrows).