1 **Enhancer-AAVs allow genetic access to oligodendrocytes and diverse populations of**
2 **astrocytes across species**
3

Authors: John K. Mich[1*], Smrithi Sunil[2], Nelson Johansen[1], Refugio A. Martinez[1], Jiatai Liu[1], Bryan B.
Gore[1], Joseph T. Mahoney[1], Mckaila Leytze[1], Yoav Ben-Simon[1], Darren Bertagnolli[1], Ravi Bhowmik[1],
Yemeserach Bishaw[1], Krissy Brouner[1], Jazmin Campos[1], Ryan Canfield[4], Tamara Casper[1], Nicholas P.
Donadio[1], Nadezhda I. Dotson[1], Tom Egdorf[1], Amanda Gary[1], Shane Gibson[4], Jeff Goldy[1], Erin L.
Groce[1], Kenta M. Hagihara[2], Daniel Hirschstein[1], Han Hou[2], Will D. Laird[1], Elizabeth Liang[1], Luke V.
Loftus[1], Nicholas Lusk[1], Jocelin Malone[1], Naomi X. Martin[1], Deja Monet[1], Josh S. Nagra[1], Dakota
Newman[1], Nhan-Kiet Ngo[1], Paul A. Olsen[1], Victoria Omstead[1], Ximena Opitz-Araya[1], Aaron Oster[1],
Christina A. Pom[1], Lydia Potekhina[1], Melissa Reding[1], Christine Rimorin[1], Augustin Ruiz[1], Adriana E.
Sedeño-Cortés[5], Nadiya V. Shapovalova[1], Michael Taormina[1], Naz Taskin[1], Michael Tieu[1], Nasmil J.
Valera Cuevas[1], Sharon W. Way[1], Natalie Weed[1], Vonn Wright[1], Zizhen Yao[1], Thomas Zhou[1], Delissa
A. McMillen[1], Michael Kunst[1], Medea McGraw[1], Bargavi Thyagarajan[1], Jack Waters[1], Trygve E.
Bakken[1], Nick Dee[1], Shenqin Yao[1], Kimberly A. Smith[1], Karel Svoboda[2], Kaspar Podgorski[2], Yoshiko
Kojima[3,6], Greg D. Horwitz[4,6], Hongkui Zeng[1], Tanya L. Daigle[1], Ed S. Lein[1,7,8], Bosiljka Tasic[1],
Jonathan T. Ting[1,4,6], Boaz P. Levi[1*]
18

Author Affiliations:
1 Allen Institute for Brain Science, Seattle WA, USA.
2 Allen Institute for Neural Dynamics, Seattle WA, USA.
3 Department of Otolaryngology Head and Neck Surgery, University of Washington, Seattle, WA, USA.
4 Department of Neurobiology and Biophysics, University of Washington, Seattle, WA, USA.
5 Division of Medical Genetics, University of Washington, Seattle, WA, USA.
6 Washington National Primate Research Center, University of Washington, Seattle, WA, USA.
7 Department of Neurological Surgery, Univ. of Washington, Seattle WA.
8 Department of Laboratory Medicine and Pathology, Univ. of Washington, Seattle WA
* Corresponding authors boazl@alleninstitute.org and johnmi@alleninstitute.org

Abstract: Proper brain function requires the assembly and function of diverse populations of neurons
and glia. Single cell gene expression studies have mostly focused on characterization of neuronal cell
diversity; however, recent studies have also revealed substantial diversity of glial cells, particularly
astrocytes. To better understand glial cell types and their roles in neurobiology, we built a new suite of
adeno-associated viral (AAV)-based genetic tools to enable genetic access to astrocytes and
oligodendrocytes. These oligodendrocyte and astrocyte enhancer-AAVs are highly specific (usually >
95% cell type specificity) with variable expression levels, and the astrocyte enhancer-AAVs show
multiple distinct expression patterns reflecting the spatial distribution of astrocyte cell types. To provide
the best glial-specific functional tools, several enhancer-AAVs were: optimized for higher expression
levels, shown to be functional and specific in rat and macaque, shown to maintain specific activity
across transgenes and in epilepsy where traditional promoters changed activity, and used to drive
functional transgenes in astrocytes including Cre recombinase and acetylcholine-responsive sensor
iAChSnFR. The astrocyte-specific iAChSnFR revealed a clear reward-dependent acetylcholine
response in astrocytes of the nucleus accumbens during reinforcement learning. Together, this
collection of glial enhancer-AAVs will enable characterization of astrocyte and oligodendrocyte
populations and their roles across species, disease states, and behavioral epochs.

Introduction

Glial cell types play critical roles in CNS development, function, and homeostasis[1,2]. Astrocytes provide trophic support for neurons[3,4], coordinate regional wiring patterns[5], respond to and regulate neurotransmission[6,7], and drive repair or pathology after traumatic injury[8–10]. Oligodendrocytes form myelin sheaths[11], strengthen circuits[12], secrete critical neurotrophic factors[13], and contribute to pathologic disease progression[14,15]. Transcriptomic characterization of glial cells has revealed an array of astrocyte and oligodendrocyte cell types, often with pronounced regional signatures[16–18]. Furthermore, species-specific features have been described[19], although the functional significance of these differences is unknown. Glial cell types have also been shown to play critical roles in CNS diseases ranging from epilepsy[20] to neurodegenerative diseases[21] to cancer[22,23]. To understand how glia differ between cell types, regions, species, and disease states, a set of tools is needed to grant targeted genetic access to these specific populations across species.

Adeno-associated virus (AAV) vectors are exceptionally useful tools for somatic transgenesis across mammalian species including human[24–29]. Short enhancer or promoter regulatory elements work effectively in AAV expression cassettes to drive cell-type selective gene expression in the brain or other organs. Recent work has shown that selective AAVs can be rationally designed by using enhancers identified from epigenetic datasets that are selectively active and fit in an AAV vector[30–40]. Enhancer-AAVs were recently developed to target different populations of excitatory and inhibitory neurons in the brain, and some enhancers have shown successful targeting of glial cell populations as well[35,41–43]. However, the field largely relies on glial promoters that have some undesirable characteristics, most notably loss of specificity or change in strength in different contexts as seen for astrocytic *GFAP* promoter fragments[44–46]. Furthermore, single cell genomics studies have revealed region-specific astrocyte cell types for which no current tools are available[16–18].

Here we present characterization of a collection of enhancer-AAVs that selectively target astrocytes and oligodendrocytes. Twenty-five astrocyte and 21 oligodendrocyte enhancer AAVs were identified from mouse and human neocortical epigenetic data that produced reporter expression that was highly specific for the intended populations, often labeling more than half of the intended cells in the area, and with a wide range of expression strengths. Multiple astrocyte-targeting vectors exhibited distinct CNS region-specific expression patterns, whereas oligodendrocyte-selective vectors generally drove expression throughout the entire CNS. Several enhancer-AAVs maintained selective expression for astrocytes or oligodendrocytes across rat and macaque, across epileptic disease status, and even for difficult transgenes. Lastly, several astrocyte tools were adapted to drive expression of functional transgenes like Cre or the detection of neurotransmitters to reveal the role of astrocytes in neurobiology. We used astrocyte-selective AAV expressing iAChSnFR[47] to measure the dynamics of acetylcholine in astrocytes of the nucleus accumbens during reinforcement learning. This collection of tools opens up new opportunities for selective labeling and functional interrogation of glial cell types across species and disease states, and could have translational applications via AAV-based therapeutics[48–50].

Results

Generation of astrocyte- and oligodendrocyte-specific enhancer-AAVs. We identified putative
enhancers specific for astrocytes and oligodendrocytes from single cell/single nucleus assay for
transposase-accessible chromatin (sc/snATAC-seq[33,34]) and single nucleus methyl-cytosine sequencing
(snmC-seq) studies from neocortex[51–54]. Thousands of astrocyte- and oligodendrocyte-selective
scATAC-seq peaks were identified previously in both human middle temporal gyrus (MTG) and mouse
primary visual cortex (VISp), averaging approximately 300-600 bp in size (**Figure 1A**). Additional
ATAC-seq datasets confirmed these peaks[55,56]. Generally, astrocyte and oligodendrocyte candidate
enhancers were accessible in non-neuronal cells but not in neuronal cells across the human
forebrain[33,55] (**Figure 1—figure supplement 1A-F**), and in the corresponding astrocyte or
oligodendrocyte subclasses across the mouse forebrain without strong cell type preferences[56] (**Figure
1—figure supplement 1G-L**).

We used three strategies to identify putative enhancers for testing: "high specificity", "high strength",
and "marker gene" (**Figure 1B-E**). The "high specificity" nomination criteria (gold square or star icons)
required enhancers and their orthologs to show accessibility specifically for both mouse and human
astrocytes or oligodendrocytes but not other cell types[33,34]. In addition, we required that these putative
enhancers not be detected in demethylated genomic regions in both mouse and human neuron
populations[52]. Some "high specificity" enhancers also showed specific demethylation in bulk human and
mouse glial cells[51] (marked by gold star icons). "High strength" putative enhancers were selected on the
basis of strong astrocyte-specific peaks using only mouse scATAC-seq data[34], with strength measured
by accessibility read count within peaks. Finally, "marker gene" putative enhancers showed specific and
strong accessibility near known astrocyte- and oligodendrocyte-specific marker genes.

We tested putative enhancer function in AAV vectors upstream of a minimal promoter driving the
reporter SYFP2 and evaluated expression throughout the mouse brain after systemic administration of
PHP.eB-serotyped AAVs. Enhancer-AAVs that showed anticipated reporter expression, were further
evaluated for specificity, completeness of expression, and cross-species activity (**Figure 1C** and **Table
S1**). All three strategies were effective, with approximately half of the candidates for both cell types
yielding astrocyte or oligodendrocyte-specific expression patterns during primary screening (**Figure 1D,
E**), which were confirmed on-target by immunohistochemistry (IHC) and/or scRNA-seq (see figures
below). Some of these vectors were also characterized in parallel large-scale efforts[35], all raw data from
which are available online at https://portal.brain-map.org/genetic-tools/genetic-tools-atlas, and the full
set of experimental mice described below are summarized in **Table S2**.

A collection of astrocyte-specific enhancer-AAVs. We screened 50 candidate astrocyte-specific
enhancer-AAVs, and 25 (50%) of them labeled astrocytes specifically with SYFP2 expression (**Figure
2—figure supplement 1**). Astrocyte-specific enhancer-AAVs showed a range of expression strengths
and patterns (**Figure 2A-O**) and vectors were categorized based on their labeling as: "Most of the
CNS", "Regional", "Scattered", "Weak" and "Mixed specificities". "Most of the CNS" astrocyte enhancer-
AAVs, including AiE0380h and the human *GFAP* promoter (GfaABC1D[57]), labeled cells with astrocyte
morphology in both brain and spinal cord (SpC, **Figure 2A-B**, **G-H**). Other examples in this category
include AiE0387m, AiE0390h, AiE0390m, and the synthetic element ProB12[41] (**Figure 2—figure**

supplement 1). "Regional" astrocyte enhancer-AAVs showed regionally restricted expression, such as
AiE0385m that labeled astrocytes primarily in the telencephalon (**Figure 2C**, **F, Figure 2—figure**
supplement 1). Other "Regional" enhancer-AAVs labeled astrocytes in subcortical domains but not in
the telencephalon, such as AiE0381h and AiE2160m (**Figure 2—figure supplement 1**); while
AiE0375m only labeled Bergmann glia, specialized astrocytes in the cerebellar cortex (CBX, **Figure 2—**
figure supplement 1). Interestingly, enhancers AiE2120m and AiE2160m labeled astrocytes in nearly
mutually exclusive regions (**Figure 2—figure supplement 2**). "Scattered" enhancer-AAVs labeled
astrocytes strongly but sparsely in most brain regions. These enhancer-AAVs include AiE0374m and its
ortholog AiE0374h (**Figure 2—figure supplement 1**). Enhancer-AAVs labeled as "Weak" gave
astrocyte-specific patterns with low expression of SYFP (e.g., AiE0373m and AiE0386m, **Figure 2—**
figure supplement 1). Last, we designated several enhancer-AAVs as "Mixed specificities" because
they labeled astrocytes and neurons. For example, AiE2118m labels many astrocytes strongly and
specifically within the telencephalon, but also labels neurons strongly in non-telencephalic structures
like midbrain (MB), deep cerebellar nuclei (CBN), and globus pallidus, external segment (GPe) (**Figure**
2—figure supplement 1).

We quantified the specificity of many of these astrocyte-specific enhancer-AAVs using multiple
independent techniques. First, we characterized SYFP2-expressing cells with immunohistochemistry
(IHC) for Sox9, a marker of astrocytes throughout the brain[58] (**Figure 2G-O**). Many of the astrocyte-
specific enhancer-AAV vectors show high specificity, which we define as >80% specificity for the target
cell population[33]. Astrocyte-specific enhancer-AAVs are usually >95% specific, and often >99% specific
in VISp for Sox9-expressing astrocytes (**Figure 2P**). Second, we also observed high specificity when
we isolated single SYFP2+ cells by flow cytometry and profiled them by scRNA-seq (**Figure 2P**).
Additionally, we assessed completeness of astrocyte labeling using IHC, and we observed that vectors
scored as "Most of CNS" often label >50% of astrocytes in VISp, but "Weak" or "Scattered" vectors
labeled many fewer astrocytes (**Figure 2P**). "Regional" vectors showed differing completeness across
brain regions as expected (**Figure 2P**). Whole-brain serial two-photon tomography (STPT) of mouse
brain transduced with astrocyte-specific enhancer-AAVs demonstrated distinct astrocyte morphologies
in multiple brain regions (**Figure 2Q**). Thus, this collection of astrocyte-specific enhancer-AAVs are
diverse with regard to the density of labeled cells, expression strength, and regionalization.

A collection of oligodendrocyte-specific enhancer-AAVs. We screened 43 candidate oligodendrocyte
enhancers, of which 21 (49%) gave oligodendrocyte-specific expression patterns (**Figure 3—figure**
supplement 1). Unlike the astrocyte collection, the oligodendrocyte enhancer-AAVs all produced
similar expression patterns throughout the gray matter and white matter tracts without any obvious
regional specificity (**Figure 3A-F**). Oligodendrocyte-specific enhancer-AAV vectors ranged in
expression from strong (for example AiE0410m, **Figure 3A**, AiE0641m, AiE0395h, and AiE0396h), to
moderate (for example AiE0409h, **Figure 3B**, and the Myelin Basic Protein (*MBP*) promoter[25,59], **Figure**
3—figure supplement 1), to weak (for example AiE0400h, **Figure 3C**). These vectors also labeled
oligodendrocytes throughout the spinal cord (**Figure 3D-F**). We confirmed molecular oligodendrocyte
characteristics of the vector-labeled cells by co-staining with CC1, a marker of oligodendrocytes[60],
which showed most vectors were highly specific across multiple brain regions (**Figure 3G-O**).
Quantification by immunohistochemistry and scRNA-seq on sorted SYFP2-expressing cells showed

>99% specificity and >45% completeness of labeling in VISp for multiple vectors (**Figure 3P**). STPT
demonstrated myelinating oligodendrocyte morphologies in multiple parts of mouse brain (**Figure 3Q**).
This collection of oligodendrocyte-specific enhancer-AAV vectors shows a diversity of expression
strengths, but appears to label a homogeneous population of oligodendrocytes.

Transcriptomic identities of astrocytes and oligodendrocytes. To investigate distinctions among
enhancer-AAV-transduced cells, we performed SMARTerV4 scRNA-seq on sorted SYFP2-expressing
cells. We characterized 2040 cells from 47 mice injected with 31 different enhancer-AAVs (1-2 mice per
enhancer-AAV). After removing low-quality single-cell transcriptomes and cells not expressing the
SYFP2 transcript, we focused the analysis on 1946 high-quality single cells. Astrocytes and
oligodendrocytes separated in the UMAP space, as did astrocytes sorted from the distinct brain regions
including VISp, midbrain/hindbrain (MB/HB), and CBX (**Figure 4A**). The molecular distinctions among
regional astrocyte populations agree with findings from recent whole-brain atlases[18]. Indeed, mapping
to a whole-brain taxonomic atlas indicates that, with high confidence, VISp-profiled astrocytes are
predominantly mapped to the *Gja1-* and *Gfap*-expressing cluster "5112 Astro-TE NN_3"[18], whereas
MB/HB-profiled astrocytes marked by AiE0381h and AiE2160m mapped primarily to the *Gja1-* and *Agt*-
expressing cluster "5109 Astro-NT NN_2"[18]. Likewise, the CBX-profiled Bergmann glia astrocytes
mapped primarily to cluster identity "5102 Bergmann NN" as expected (**Figure 4B-D**). In contrast,
labeled oligodendrocytes largely mapped to *Cldn11-* and *Mog*-expressing and most abundant
oligodendrocyte cluster "5158 MOL NN"[18] regardless of the enhancer used to label them (**Figure 4E**),
confirming that oligodendrocyte enhancer-AAVs label a largely homogeneous population of
oligodendrocytes.

To understand the molecular regulation of these astrocyte and oligodendrocyte-selective enhancer-
AAVs, we performed de novo motif detection on a collection of specific and strong astrocyte and
oligodendrocyte enhancers (n = 15 each) using MEME-CHIP[61]. This analysis yielded one motif
occurring in the majority of enhancers in each set (**Figure 4F**, **G**). Thes motifs had strong enrichments
as measured by MEME-CHIP E-values less than 0.01, corresponding to the expected number of
equally sized motifs of same or greater log likelihood ratio occurring in a set of random sequences of
equal nucleotide content. We mapped these motifs against known transcription factor (TF) motif
databases[62–64], which revealed top matches to the Zic family consensus motifs for astrocytes (JASPAR
accession numbers MA0697.2, MA1628.1, and MA1629.1; average of these three shown) and the Sox
family motif for oligodendrocytes (JASPAR accession number MA0442.1 [Sox10 shown], and also
Uniprobe accession numbers UP00030.1 and UP00062.1; **Figure 4F**, **G**). These analyses suggest that
Zic and Sox family transcription factors might be key determinants of astrocyte versus oligodendrocyte
identity in the CNS[65–68], confirming the results of independent analysis of these sequences[36]. Moreover,
Zic and Sox gene family members were differentially expressed between the profiled astrocytes and
oligodendrocytes (*Zic5* 32-fold mean difference, *Sox10* 455-fold mean difference, $p < 1e-16$ each;
Figure 4H). These results suggest Zic5 and Sox10 each play key roles in determining specificity of
these glial enhancer-AAVs.

Regional expression correlates with astrocyte cell type distribution. Using STPT imaging, we observed
astrocyte-specific enhancer-AAVs to have two distinct expression patterns within the basal ganglia
circuit. Several enhancer-AAVs showed elevated expression in astrocytes of the dorsolateral striatum

and depletion in the globus pallidus (GP; **Figure 4I-K**), and several other enhancer-AAVs drove
stronger transgene expression in astrocytes in the GP compared with those of the lateral striatum
(**Figure 4L-N, Figure 2—figure supplement 2**). To determine if these enhancer-AAV expression
patterns correspond to transcriptomically defined astrocyte cell types, we evaluated the spatial
distributions of all astrocyte cell types in the Allen Brain Cell Whole Mouse Brain taxonomy[18].
Interestingly, two closely related astrocyte cell types were strongly enriched in the dorsolateral striatum
and cortex ("5112 Astro-TE NN_3" and "5113 Astro-TE NN_3"), while another was strongly enriched in
the GP ("5109 Astro-NT NN_2"), demarcating the same boundaries observed with the collection of
astrocyte enhancer-AAVs (**Figure 4O-R**).

Measuring and optimizing enhancer strength. In some cases, enhancer-AAVs might not drive sufficient
levels of a transgene to functionally affect the target cell. We sought to boost the expression levels of
some enhancers by assembling concatemers of "core" sequences. These core sequences are
responsible for the selective expression patterns and are often found in the central third of the original
enhancer region identified by snATAC-seq[33] (that is, ~100-200 bp core from ~300-600 bp original
enhancer, **Figure 5A**). We observed that concatenation of the core can substantially increase
expression from the original enhancer, such as AiE0387m concatenated to AiE0387m_3xC1 (**Figure**
5B, **C**), AiE0390h concatenated to AiE0390h_3xC2 (**Figure 5D**, **E**), or AiE0390m concatenated to
AiE0390m_3xC2 (**Figure 5F**, **G**) while retaining similar expression patterns (**Figure 5H-J**) and cell type
specificity (**Figure 5K**). However, concatenation sometimes resulted in a less dramatic effect on
expression, (e.g., AiE0410m_3xC and AiE0641m_3xC; **Figure 3—figure supplement 1**).

We established single-cell measurements of reporter expression to compare enhancer strengths. We
found that single-cell reporter fluorescence by flow cytometry correlated with vector read counts from
SMARTerV4 scRNA-seq for both astrocytes and oligodendrocytes (astrocyte Pearson correlation
coefficient = 0.63, p < 0.001; oligodendrocyte Pearson correlation coefficient = 0.53, p < 0.05; **Figure**
5L, **M**, **Figure 5—figure supplement 1**). These measurements revealed that several concatenated
enhancer-AAVs, including AiE0390m_3xC2 and AiE0390h_3xC2, drove the strongest expression
among the vectors we have tested (**Figure 5L**), consistent with the microscopy results (**Figure 5B-J**).
Conversely, MB/HB (AiE0381h and AiE2160m) and Bergmann glia (AiE0375m) astrocyte enhancers
have among the weakest expression levels we have tested (**Figure 5L**), likely a consequence of
selecting glial enhancers from cortical accessibility datasets.

Enhancers can yield more robust specific expression across transgenes than classic promoters.
Previously it was demonstrated that the human *GFAP* promoter (GfaABC1D[57]) can change specificity
from astrocytes to neurons over time when driving certain difficult transgene cargos including NeuroD1-
mCherry fusion protein[44]. Accordingly we tested whether astrocyte-specific enhancers might be more
robust to transgenes than *GFAP* promoter. As demonstrated previously, we found that at seventeen
days post transduction, *GFAP*-driven NeuroD1-mCherry is expressed predominantly in NeuN+ neurons
in primary visual cortex (VISp, 90 ± 2% of labeled cells, mean ± standard deviation), unlike *GFAP*-
driven SYFP2 which is expressed with high Sox9+ astrocyte specificity (98 ± 2% of labeled cells,
Figure 6A-B, p < 0.001, see also **Figure 2**). In sharp contrast an optimized astrocyte-specific
enhancer, AiE0390m_3xC2 maintains high levels of astrocyte specificity whether driving SYFP2 (100 ±
0% of labeled cells) or NeuroD1-mCherry (95 ± 4% of labeled cells, **Figure 6A-B**, p < 1e-7, see also

Figure 5K). These results suggest that optimized astrocyte-specific enhancers may provide more
predictable expression for certain transgenes, and confirm the previous results that astrocytic NeuroD1-
mCherry expression alone is insufficient to drive astrocyte transdifferentiation to neurons[44].

Predictability of enhancer-AAV expression across tissues and disease states. Recent work suggests
that AAV-mediated transduction and high transgene expression in organs such as the liver and dorsal
root ganglia is associated with toxicity[29,69]. We tested if we could predict off-target activity from
enhancer accessibility profiles in a human body-wide epigenetic dataset[70]. Different astrocyte-specific
enhancers showed either moderate or low accessibility across many body organs (**Figure 7A**), since
this was not a factor in their selection for testing. We assessed off-target transgene expression in liver
after intravenous delivery since PHP.eB capsid transduces the liver[71]. We observed that astrocyte
enhancers with moderate levels of accessibility in liver (AiE0381h, AiE0371m, AiE0371h, and
AiE0386m) expressed SYFP2 in many hepatocytes, whereas the enhancers with little or no liver
accessibility (AiE0387m, AiE0375m, AiE0390h, and AiE0390m) expressed SYFP2 in many fewer
hepatocytes ($p < 0.001$, **Figure 7B-C**), demonstrating strong predictability of enhancer expression
profiles in off-target tissues. In contrast, the *GFAP* promoter can drive bright expression in few
hepatocytes, which is not predictable from any epigenetic or transcriptomic atlases.

GFAP expression can change expression in the context of disease or injury[9], and the synthetic *GFAP*
promoter can change specificity when delivering different transgenes[44–46], suggesting this might be a
poor tool for genetic access to astrocytes in disease. We compared *GFAP* promoter and one of the
best enhancer-AAVs (AiE0390m) in *Scn1a*[+/-] Dravet syndrome model mice since they have strong
epilepsy-associated reactive astrogliosis[72]. We injected SYFP2-expressing enhancer-AAVs into these
mice prior to the Dravet syndrome critical period of high susceptibility to seizures and mortality at P21
and analyzed tissues at P42 (**Figure 8A**). Significant hippocampal gliosis was seen in Dravet syndrome
model mice, revealed by elevated endogenous GFAP immunoreactivity in all hippocampal layers
(**Figure 8B**). Concomitant with this gliosis, the *GFAP* promoter-driven AAV reporter changed its
expression pattern. Normally, this promoter drives moderate levels of astrocyte-specific reporter
expression. However, in the Dravet mice experiencing epilepsy, expression strength in astrocytes was
considerably elevated, and ectopic expression was observed in many dentate gyrus granule cell
neurons (DGCs, 12 ± 6% of DGCs labeled, $p < 0.01$, **Figure 8B, C**). In sharp contrast, the AiE0390m
enhancer-AAV vector maintains astrocyte specificity at moderate levels despite the profound reactive
gliosis in these diseased animals (0 ± 0% of DGCs labeled, **Figure 8B, C**).

Astrocyte specific sensing of cholinergic signals in the nucleus accumbens during behavior. We next
asked whether the vectors would drive sufficient expression to obtain functional signals in a cell-type
specific manner. We created a vector driving astrocyte-specific expression of the acetylcholine indicator
iAChSnFR[47] (**Figure 9A,B**), to detect extracellular acetylcholine fluctuations in the nucleus accumbens
(NAc) in an awake and behaving animal using fiber photometry. After stereotaxic injection, we
implanted optical fibers above the injection site to perform fiber photometry. We trained mice to perform
a dynamic foraging reinforcement learning task while we recorded photometry signals to assess bulk
acetylcholine fluctuations in the NAc. In the task, water-restricted mice chose freely between two lick
ports for a water reward after an auditory cue. Reward probabilities of the two lick ports were changed
in a block-design manner, which resulted in both rewarded and unrewarded trials (**Figure 9C**). During

these trials, the astrocyte-specific iAChSnFR vector drove sufficient expression to observe fluorescence
intensity fluctuations (**Figure 9D**) which can be seen to differ during individual rewarded and
unrewarded licks (**Figure 9D**, bottom left). Both rewarded and unrewarded trials showed an increase in
fluorescent signal at the time of choice, followed by a deviation in signal depending on whether the trial
was rewarded (**Figure 9E**). Astrocyte acetylcholine signals decreased more in rewarded trials than in
unrewarded trials (**Figure 9E**). In summary, these results indicate that glial-selective enhancer-AAVs
can be applied to measure functional acetylcholine dynamics in the NAc.

Astrocyte- and oligodendrocyte-specific AAV-Cre vectors. AAVs that can selectively drive Cre
recombinases are valuable tools for mouse genetics since the AAV can be delivered somatically for cell
type-specific recombination of floxed alleles. We used some of the best astrocyte- and oligodendrocyte-
specific enhancers to express an attenuated R297T mutant Cre recombinase in *Ai14* reporter mice[73,74]
(**Figure 10A**). AiE0390m_3xC2-driven Cre vector drove astrocyte-specific recombination in many parts
of mouse brain, including specific (95 ± 3%) and complete (98 ± 1%) labeling in VISp gray matter
astrocytes (**Figure 10B-D**). Similarly, an AiE0410m-driven Cre vector also incorporating three miR126
binding sites to prevent unwanted endothelial cell recombination[75] drove oligodendrocyte-specific
recombination in many parts of mouse brain, including specific (97 ± 3%) and complete (81 ± 0.1%)
labeling in VISp gray matter oligodendrocytes (**Figure 10E-G**). Additional vectors utilizing AiE0390m
and AiE0387m also showed specific and complete astrocyte labeling in VISp (**Figure 10H-I**). These
tools may permit fine interrogation of the roles of astrocyte and oligodendrocyte gene expression in
brain biology.

Cross-species genetic access to astrocytes and oligodendrocytes. We tested whether several glial-
selective enhancer-AAV vectors could maintain specific expression across species. We first tested
conservation in neonatal rats after ICV administration (**Figure 11A**). We observed that an optimized
AiE0390m_3xC2 vector containing 4X2C miRNA binding sites to prevent any unwanted expression in
excitatory neurons[76] labeled rat cortical astrocytes with high specificity (98% specific, **Figure 11B**, **C**).
We also found that an AiE0641m-driven AAV vector labeled rat cortical oligodendrocytes with high
specificity (91% specific, **Figure 11D**). Note that astrocyte labeling completeness and spread to caudal
brain structures could not be assessed since ICV administered virus resulted in uneven spread. Thus,
some vectors identified in the mouse screen maintained specificity in rats after ICV injection into
neonates, consistent with the results from other enhancer-AAV collections[38].

We extended these cross-species tests to primate, using multisite intraparenchymal injections into
Macaca nemestrina (pig-tailed macaque, **Figure 11E**). We administered AiE0410m AAV vector into
motor (MO) cortex and observed cell morphologies of myelinating oligodendrocytes throughout the
cortical column (**Figure 11F-G**). Interestingly, we also observed SYFP2-expressing cells with a different
morphology: one to three processes that spiral around stretches of tubular structures approximately 15-
20 microns in diameter often running perpendicular to the cortical pial surface (**Figure 11H**). These
tubular structures have not yet been defined and were not observed in mouse or rat testing, but both
morphological types of SYFP2-expressing cells co-expressed the oligodendrocyte marker *SOX10* by
mFISH with high specificity (94%, **Figure 11I**).

We also injected the somatosensory cortex (SS) with the astrocyte-specific AiE0390m AAV vector and
observed many SYFP2-expressing cells with astrocyte morphology throughout the cortical column
(**Figure 11J**; note a small number of large layer 5 pyramidal neurons labeled as well, which we did not
observe in mouse testing). SYFP2-expressing astrocytes co-expressed GFAP either in parenchyma
(**Figure 11K**) or in apposition to a large blood vessel (**Figure 11L**), and some showed fibrous
morphology in white matter (**Figure 11—figure supplement 1**). Enhancer-AAV-labeled astrocytes also
expressed the astrocyte-specific transcript *FGFR3* with high specificity (92%) and about half of the
FGFR3+ gray matter astrocytes were labeled through the whole cortical depth near the injection site
(51%) (**Figure 11M**). In an additional animal we also tested AiE0390m_3xC2 vector containing 4X2C
miRNA binding sites in MO, and we observed 99% specificity for SOX9+ astrocytes and no layer 5
pyramidal neuron labeling (**Figure 11N-O**), and a final injection into a third animal also confirmed 98%
specificity of AiE0390h-driven vector for SOX9+ astrocytes in SS (**Figure 11P-Q**). These studies
suggest that enhancer-AAV vectors can provide specific and dependable genetic access to astrocytes
and oligodendrocytes across multiple species and reveal morphological glial features not observed in
the mouse.

Discussion

Flexible and dependable tools to target glial cell populations will be essential to understand their diverse roles in brain biology. Here we report a collection of astrocyte- and oligodendrocyte-specific enhancers that can be used in AAV vectors and applied across species. Most of these enhancer-AAVs generated highly specific labeling of astrocytes or oligodendrocytes, and often substantial completeness of labeling. Detailed characterization revealed the enhancers showed a range of expression strengths, and the astrocyte enhancers frequently exhibited regional enrichment and differences in labeling densities. We demonstrate that this enhancer-AAV toolset can be applied: 1) across species in mouse, rat and macaque, 2) in epileptic mice where gliosis is occurring without losing specificity, and 3) to deliver Cre selectively to many astrocytes and oligodendrocytes, 4) irrespective of transgene delivered, and 5) to measure circuit dynamics with a neurotransmitter sensor *in vivo*. As a result, these glial enhancer-AAVs will be useful for interrogating the roles of these glial cell types in health and disease.

Lesson learned from screening. Several lessons were learned through the process of screening for astrocyte and oligodendrocyte enhancer-AAVs. First, multiple selection criteria can identify strong glial enhancer-AAVs. Excellent functional enhancers were derived from genome-wide peak selection across mouse and human datasets of distinct epigenetic modalities, peak selection from one mouse dataset based on peak strength, or peak selection only near marker genes. Second, screening one enhancer-AAV at a time can be efficient for the identification of useful enhancer-AAVs. Nearly half of the candidate enhancers proved to be specific for the targeted type, and this created a large and diverse library of new enhancer-AAVs that labeled astrocytes or oligodendrocytes in different ways. Third, while enhancers were identified from neocortex at the subclass level, brain-wide patterns match well to whole brain cell gene expression atlas patterns[18]. Specifically, AAV-labeled oligodendrocytes appear homogeneous across the brain, while astrocytes show prominent regional enrichments and differences in cell density. Fourth, body-wide specificity of enhancer-AAV expression can be predicted based on body-wide epigenetic datasets[70]. Thus, off-target expression can be predicted and limited during the enhancer selection process, and enhancers can be identified that label different cell types throughout the body. These attributes will make enhancer-AAVs a valuable tool for precision gene therapy where only the cell population of interest is expressing the therapeutic transgene.

Enhancer-AAVs show diverse expression patterns. We tested a large collection of astrocyte-selective enhancer-AAVs and saw a diversity of expression patterns. Some astrocyte enhancer-AAVs predominantly labeled in telencephalic structures, some in MB/HB structures, and others showed sparse "Scattered" but uniform expression. Within the forebrain, we also observed multiple enhancer-AAVs that showed mutually exclusive enriched or depleted expression in dorsolateral striatum or the globus pallidus. These regional differences were reflected by region-specific astrocyte transcriptomic profiles in agreement with recent results[18]. The GPe astrocytes express high levels of GABA uptake gene *Slc6a11*[16,77,78] while striatal and cortical (but not GPe) astrocytes express high levels of glutamate uptake gene *Slc1a2*[16,78,79]. This suggests that the astrocytes help selectively maintain glutamate or GABA tone depending on the local neurotransmitter balance. The functional roles of astrocytes in different brain regions will require additional experiments that this collection of enhancer-AAVs may

facilitate. Similarly, it is not yet clear what produces "Scattered" astrocyte reporter expression, and the
answer will await future experiments.

Enhancer-AAVs can be optimized to improve vector function. Achieving functional levels of transgene
expression is critical for applying enhancer-AAV tools to learn new biology and deliver gene therapies.
We show that two astrocyte enhancers could be optimized through generation of a triple concatenated
core to produce significantly higher levels of transgene expression without sacrificing astrocyte
specificity. One such optimized tool was used to detect acetylcholine activity in the NAc. The NAc has
been implicated in reward-related reinforcement learning and receives inputs from dopaminergic and
serotonergic neurons, in addition to local cholinergic signaling[80–83]. We used enhancer-AAVs to deliver
the acetylcholine indicator iAChSnFR to astrocytes within the NAc, and measured dynamics with fiber
photometry. This experiment showed that the astrocyte enhancer AiE0390m_3xC2 maintained faithful
astrocyte specificity after direct injection into NAc and expressed sufficient iAChSnFR for sensing of
acetylcholine in live awake animals. It also showed that iAChSnFR expressed by astrocytes could
readily detect acetylcholine dynamics in NAc, and that astrocytes are a good cellular compartment for
the sensor.

Different optimization efforts enabled the generation of a functional Cre-AAV that was specific for
astrocytes in most parts of the brain. Cre recombinases can lose specificity when expressed from
enhancer-AAVs, possibly due to low-level expression of this potent enzyme[30]. We were able to
successfully generate an astrocyte-selective Cre AAV using an attenuated Cre recombinase with the
R297T mutation[73,74], in combination with the AiE0390m_3xC2 enhancer. As a result we produced a
highly specific somatic astrocyte Cre. Similarly we were able to produce a highly specific
oligodendrocyte Cre AAV using the AiE0410m enhancer along with miR126 binding sites. We
anticipate these vectors may unlock many types of functional experiments dissecting the roles of these
important glial cell populations.

Enhancers-AAVs can be identified that have conserved specificity from mouse to macaque. Some
astrocyte and oligodendrocyte enhancer-AAVs are active and selective across species from mouse
through macaque. This property will allow these tools to be applied somatically in multiple organisms
besides mice. Testing enhancers in macaques revealed interesting morphological differences
compared to mouse. Abundant mature oligodendrocytes with large dendritic arbors were labeled in
both mouse and primate tissues with AiE0410m, but this enhancer-AAV also labeled *SOX10*+ cells
wrapping around radial tubes (presumably blood vessels) in primate but not mouse. Also, the astrocyte
enhancer AiE0390m labeled abundant protoplasmic astrocytes in the gray matter in mouse and
macaque, but also labeled several large fibrous astrocytes that were not observed in the mouse
experiments. The ability to function across species and label cells that are not obviously represented in
mouse tissue, makes this collection of enhancer-AAVs a powerful toolset to better understand new
biology. It also makes a compelling case that some of these enhancer-AAVs could be suitable for use
in human gene therapy where astrocyte or oligodendrocyte expression selectivity is required.

Enhancer-AAVs can be cell state-independent. AAVs that drive expression using promoters can cause
gene expression changes in a state dependent fashion. We showed that the *GFAP* promoter changed
expression strength and specificity in the context of epilepsy-induced gliosis using an *Scn1a*

haploinsufficiency model. This is not surprising since the *GFAP* gene is known to change in response to

disease and injury[8,9]. Additionally, *GFAP* promoter-driven expression changed specificity from

astrocytes to neurons with the difficult transgene NeuroD1-mCherry, as seen in other reports[44].

Enhancer-AAVs utilizing AiE0390m, on the other hand, did not show a change of expression in either

epileptic gliosis conditions or due to differing transgenes. This could be due to the enhancer being

selected based on core cell type identity, and most properties of cell types have not been seen to

change character dramatically in the context of disease[20,84,85]. Enhancers can be selected that do not

change activity during development, aging, or disease, in a similar way to avoid selecting enhancers

predicted to have activity in off-target tissues. As epigenetic datasets are expanding to cover these

axes of disease, development, and aging, it is becoming feasible to select only putative enhancers with

the desired activity profiles.

Conclusion. We have characterized a large collection of enhancer-AAV vectors for targeting astrocytes

and oligodendrocytes. These vectors will provide researchers with the ability to mark and manipulate

these critical cell types in a variety of species, genetic backgrounds, ages, and disease contexts, and

could also enable delivery of therapeutics. Combined with other recently discovered AAV-based

tools[25,26,31–38,40,43,71], this glial-targeting toolbox will help to advance the understanding of the roles of

glial cell types in brain biology, make the complex cellular anatomy of the brain more experimentally

tractable, and advance the development of AAV-based therapeutics for human CNS disorders.

Methods

Epigenetic analysis and enhancer nomination. We identified candidate astrocyte- and oligodendrocyte-specific enhancers from cortical epigenetic datasets. We used the following datasets: human middle temporal gyrus snATAC-seq[33], mouse primary visual cortex scATAC-seq[34], human frontal cortex snmC-seq[52], mouse frontal cortex snmC-seq[52], human frontal cortex bulk mC-seq[51], and mouse frontal cortex bulk mC-seq[51]. A single cell glial snmC-seq dataset[53,54] became available only after initial identification of most of the enhancers described in this study. From the single nucleus/cell ATAC-seq datasets, we aggregated reads according to cell subclass as in the references, and then called peaks using Homer findPeaks (http://homer.ucsd.edu/homer/) with the -region flag, yielding typically tens of thousands of peaks per subclass, sized approximately 300-600 bp, as previously described[33]. To find differentially methylated regions (DMRs) we either used the published regions by Luo et al. 2017 in Extended Data Tables 5 and 6[52], and aggregated by subclass and then to all neurons using bedtools merge (https://bedtools.readthedocs.io/en/latest/). Alternatively for bulk non-neuronal DMRs we used methylpy DMRfind with minimum differentially methylated sites set to 1 on the dataset of Lister 2013, as previously described[33,52]. To convert mouse and human peak or DMR regions to each other's genomic coordinates for direct intersectional analysis, we used liftOver (https://genome.ucsc.edu/cgi-bin/hgLiftOver) with minMatch parameter set to 0.6. All peak regions described in this manuscript successfully liftOver from human to mouse, and vice-versa, except AiE0733m which does not have an obvious human ortholog via liftOver.

To automatically identify peaks and DMRs genome-wide that are astrocyte- or oligodendrocyte-specific within each dataset, we used a series of bedtools intersectBed (https://bedtools.readthedocs.io/en/latest/) operations to filter for regions that are only detected in astrocytes or oligodendrocytes. For the "high specificity" criteria, we found peaks that were specifically detected in both human and mouse cortical astrocytes/oligodendrocytes[33,34], but did not overlap DMRs from either human or mouse neurons of any subclass[52], and these candidate enhancers are marked by gold square icons in **Figure 1**. These criteria yielded a set of 87 candidate astrocyte-specific enhancers and 112 candidate oligodendrocyte-specific enhancers, and the top 17 (Astrocyte) or 16 (Oligodendrocyte) candidate enhancers were chosen from this list as ranked by Homer findPeaks score. Homer findPeaks score is a measure of peak significance relative to local background, not peak strength. Additionally, a small number of these "high specificity" criteria candidate enhancers also overlapped with DMRs from both human and mouse non-neuronal cells[51] (3 Astrocyte and 4 Oligodendrocyte), and these are marked by gold star icons in **Figure 1** and **Figure 1—figure supplement 1**.

For the "high strength" criteria we found peaks that were specifically detectable in cortical astrocytes, using mouse scATAC-seq data only[34], and agnostic to detection in human and methylation datasets. This analysis yielded 2119 (astrocyte) and 3940 (oligodendrocyte) candidate enhancers, which were ranked by read counts within the region, and the top 7 candidate enhancers for astrocytes from this list were chosen. This ranking led to nomination of peaks that are overall stronger and longer, and these candidate enhancers are marked with a purple triangle in **Figure 1B**, but accessibility profiles were not always conserved in human tissue, as shown in **Figure 1—figure supplement 1A,C**.

Some candidate enhancers were identified manually in the vicinity of known astrocyte or
oligodendrocyte marker genes by visual inspection of ATAC-seq read pileups on UCSC browser
(marked as "*M*" for Marker genes in **Figure 1B**). Methylation data was not visualized in this manual
nomination process. Importantly we found that both automatic and manual approaches can identify
peaks with high strength and specificity, as shown in **Figure 1D-E**.

Additionally, enhancer AiE2160m was initially identified as a candidate enhancer for pericytes in cortex
using the data of Graybuck et al.[34], but it was found in the course of this study to instead label
mid/hindbrain astrocytes.

To model enhancer screening results as a generalized linear model, we confined analysis to 50
screened enhancers where we observed a clear yes/no screening result for both itself and its cross-
species ortholog. These candidate enhancers were AiE0371, 372, 373, 375, 377, 379, 380, 382, 383,
384, 388, 393, 394, 398, 399, 401, 406, 407, 408, 374, 376, 390, 395, 409, and 410, both the m and h
orthologs for each. For each of these genomic regions we calculated candidate enhancer strength
(read CPM within either astrocytes or oligodendrocytes), candidate enhancer specificity (defined as the
proportion of astrocyte or oligodendrocyte enhancer strength relative to the summed strength in all
populations, using the data of Mich et al. or Graybuck and Daigle et al.[33,34]), candidate enhancer length
in base pairs, region-segmented PhyloP using the previous method[33], and tabulated whether each
candidate enhancer's partner ortholog worked (binary yes [1] or no [0]). We fit a logistic generalized
linear model of testing results from these predictors using glm() in R with the following command:

```
503  glm( Screen_result_01 ~ Length + PhyloP + Specificity + Strength_cpm +
504  Ortholog_result_01, family=binomial(link='logit'), data = data)
```

The significance of each coefficient to predict the screening result was determined from the coefficients
of the model, using the data as provided in **Table S1**. Although high peak specificity and strength were
important criteria for candidate enhancer identification, these metrics each had little predictive power to
explain success or failure of screening collection testing as evidenced by coefficients of fit to a logistic
linear model (**Table S1**; strength z-value = 1.43, p = 0.15; specificity [defined as proportional strength
within target cell subclass] z-value = -1.20, p = 0.23), similar to enhancer length (z-value = -0.13, p =
0.89), enhancer sequence conservation measured by PhyloP (z-value = 0.87, p = 0.38), and the
presence of a functional ortholog in testing (z-value = 1.77, p = 0.077), which suggests that there are
additional undiscovered elements that determine successes versus failures in AAV-based enhancer
screening. Overall, the null deviance was 67.3 on 49 degrees of freedom, and the residual deviance
was 57.7 on 44 degrees of freedom, again indicating little power of these features to predict the
screening results.

Cloning and packaging enhancer-AAVs. With candidate enhancers chosen, we next found their
predicted DNA sequence from genomic reference sequence using Bioconductor package Bsgenome[86].
We extracted the sequence and padded 50 bp to each side of the enhancer to provide room for forward
and reverse primer binding sites that capture the entire enhancer. From these padded sequences, we
used automatic primer design in Geneious to identify primer pairs within the 50 bp pads to specifically
amplify each enhancer, and append a constant 5' homology arm to each enhancer for automatic
Gibson assembly into reporter-AAV plasmid. We amplified the regions from C57Bl/6 tail snip DNA or

from human male genomic DNA (Promega catalog # G1471) using FastPhusion 2x Master Mix

(Thermo Fisher catalog # F548L), and >90% of the PCR reactions were successful on the first try. In

some cases we redesigned primers to attempt a second amplification.

We cloned into reporter backbone CN1244 (Addgene plasmid #163493) using the sites MluI/SacI and

the 5' primer homology arms F: TTCCTGCGGCCGCACGCGT and R:

GACTTTTATGCCCAGCCCGAGCTC, using Infusion kit (Takara catalog # 638949). For some

enhancers we instead cloned into a next generation reporter vector backbone that includes a SYFP2-

P2A-3xFLAG-H2B reporter for detection of cytosolic SYFP2 and nuclear FLAG for simultaneous

expression analysis and snRNA-seq, using the same cut sites and homology arms (see **Table S1**). We

transformed infusion reactions into Mix N' Go (Zymo Research catalog # T3001) chemically competent

Stbl3 E. coli (Thermo Fisher catalog # C737303) and selected on 100 µg/mL carbenicillin plates. We

cultured individual clones at 32 C, verified them by Sanger sequencing, maxiprepped them with 100

µg/mL ampicillin, and saved them as frozen glycerol stocks at -80°C.

We used maxiprep DNA for packaging into PHP.eB AAV particles. For routine enhancer-AAV screening

by intravenous delivery in mouse we generated small-scale crude AAV preps by transfecting 15 µg

maxiprep enhancer-reporter DNA,15 µg PHP.eB cap plasmid, and 30 µg pHelper plasmid into one 15-

cm dish of confluent HEK-293T cells using PEI-Max (Polysciences Inc. catalog # 24765-1). After

transfection the next day we changed the medium to 1% FBS, and after 5 days the cells and

supernatant were collected, freeze-thawed 3x to release AAV particles, treated with benzonase (1 µL)

for 1 hr to degrade free DNA, then clarified (3000g 10min) and then concentrated to approximately 150

µL by using an Amicon Ultra-15 centrifugal filter unit at 5000g for 30-60 min (NMWL 100 kDa, Sigma

#Z740210-24EA), yielding a titer of approximately 3-5 E13 genome copies (gc)/mL. For large-scale

gradient preps for intraparenchymal injection into macaque or mouse or ICV injection into rat, we

transfected 10 15-cm plates of cells, and also purified preps by iodixanol gradient centrifugation. We

assessed viral titer for both crude and gradient AAV preps by digital droplet PCR on a BioRad QX200

system. All vectors showing specific expression patterns are available through Addgene.

Optimizing enhancer strength through concatemerization. For some native enhancers that showed

specific expression patterns, we sought to boost their expression levels through concatemerization. To

concatemerize, we segmented the enhancer (typically approximately 400-600 bp) into approximately

thirds with approximately 25 bp of overlaps at the junctions (each a candidate "core" of approximately

200 bp), then designed a tandem array of three cores in series (approximately 600 bp). These synthetic

tandem array sequences were gene synthesized by Azenta/GeneWiz PRIORITYGene synthesis

service with flanking MluI/SacI sites for restriction enzyme digestion and ligation into corresponding

sites in CN1244. We then packaged and tested concatemerized PHP.eB enhancer-AAVs as above.

Mice and injections. All mouse experimentation was approved by Allen Institute Institutional Animal

Care and Use Committee (IACUC) as part of protocol #2020-2002. In these studies, we purchased

C57Bl/6J mice from The Jackson Laboratory (Stock # 000664). For enhancer screening these C57Bl/6J

mice were injected with AAVs in the retro-orbital sinus at age P21 with 5E11 genome copies of

AAV/PHP.eB viral vectors with brief isoflurane anesthesia. For enhancer validation studies (IHC) mice

were injected the same way but between ages P42 to P56. Tissues from mice were harvested at 3 to 4

weeks post injection for analysis, except in the case of the NeuroD1-mCherry experiments (**Figure 1F**)
where mice were analyzed at 17 days post injection as previously reported[44], and in the liver
experiments (**Figure 6C**) in which 1-2 mice for each vector were analyzed at 3 weeks and 1-2 mice
were analyzed at 36 weeks to confirm durability across age. We perfused animals with saline then
4%PFA in PBS, and harvested brains or other tissues and post-fixed in 4%PFA overnight, before
rinsing and cryoprotecting in 30% sucrose solution before sectioning at 30 micron thickness on sliding
microtome with a freezing stage. PFA used in these experiments was prepared by dissolving solid PFA
(VWR Chemicals #28794.295) in PBS by heating, which was then stored in frozen aliquots. For
enhancer screening we counterstained with DAPI and propidium iodide and mounted in Vectashield
Vybrance, and imaged on either a Nikon Ti-Eclipse or Nikon Ti-Eclipse 2 epifluorescent microscope,
Olympus FV-3000 confocal microscope, or Leica Aperio slide scanner. In some experiments where
noted, we tested enhancer-AAVs after bilateral intracerebroventricular (ICV) injection at age P2 using
the technique of Kim et al.[87] These ICV-injected pups were harvested for tissue analysis at age P21.
For whole brain imaging of expression pattern, we performed sequential blockface imaging of brains
using the TissueCyte 1000 serial two-photon tomography system[88].

For testing in Dravet syndrome model mice, 129S1/SvImJ -*Scn1a^{em1Dsf/J}* mice (strain # 034129) were
purchased from Jackson Laboratories and bred to C57Bl/6J mice to create *Scn1a^{R613X/+}* pups on a F1
hybrid C57Bl/6J:129S1/SvImJ background, and these pups were injected retro-orbitally at P21 with
tissue analysis at P42. Additionally, we also tested enhancer-AAV vectors in *CMV-Cre;Scn1a^{A1783V/+}*
pups on C57Bl/6J background, which were generated from crossing B6(Cg)-*Scn1a^{tm1.1Dsf/J}* male mice
(The Jackson Laboratory, strain #:026133) with homozygous CMV-Cre female mice (*B6.C-Tg(CMV-*
cre)1Cgn/J, The Jackson Laboratory, strain # 006054). Data from both lines appeared similar, so we
combined them for analysis.

Mouse immunohistochemistry (IHC). For IHC and ISH, we transcardially perfused mice with ice-cold 25
mL HBSS (Thermo Fisher Scientific # 14175079) containing 0.25 mM EDTA (Thermo Fisher Scientific
AM9260G), followed by 12 mL of ice-cold 4% paraformaldehyde in 1x PBS, freshly prepared from
16% PFA (Electron Microscopy Sciences #15710). We dissected brains and other tissues from
carcasses and post-fixed them at 4 degrees overnight, and the next morning we rinsed the tissues with
fresh PBS, and then transferred to 30% sucrose solution in PBS for cryoprotection. For sectioning on
Leica CM3050 cryostat, we then embedded tissues in OCT cryo-compound (Tissue-Tek # 4583) at
room temperature at least 3 hours, then froze the blocks on dry ice and stored at -80°C until sectioning
at 25 micron thickness. Alternatively, we sectioned half-brains at 25 µm thickness on frozen 30%
sucrose solution slabs on a sliding microtome (Leica SM2000R) equipped with freezing stage. Sections
were stored at 4 degrees in PBS containing 0.1% sodium azide until analysis. IHC of liver was
performed as for brain, focusing on the frontal lobe.

For IHC we used the following antibodies: chicken anti-GFP (Aves # GFP-1010, 1/3000), rabbit anti-
Sox9 (Cell Signaling clone D8G8H, # 82630S), mouse CC1 antibody (Abcam # ab16794, 1/1000),
mouse anti-GFAP (Millipore Sigma clone G-A-5, # G3893, 1/1000), with 5% normal goat serum
(Thermo Fisher Scientific # 31872) and 0.1% Triton X-100 (VWR 97062-208) for blocking and
permeabilization, and appropriate Alexa Fluor-conjugated secondary antibodies for detection.

Flow cytometry and single cell transcriptomics. We prepared cell suspensions for flow cytometry and
single cell RNA-seq from brain tissue as previously described[89]. Briefly, for flow cytometry, we perfused
mice transcardially under anesthesia with ACSF.1. We harvested the brains, embedded in 2% agarose
in PBS, then sliced thick 350 micron sections using a compresstome with blockface imaging, then
picked the sections containing the region of interest (VISp, or mid- and hindbrain, or cerebellar cortex),
and dissected out the regions of interest. We then treated dissected tissues with 30U/mL papain
(Worthington LK003176) in ACSF.1 containing 30% trehalose (ACSF.1T) in a dry oven at 35°C for 30
minutes. After papain treatment we quenched digestion with ACSF.1T containing 0.2% BSA, triturated
sequentially using fire-polished glass pipettes with 600, 300, and 150 micron bores, filtered the
released cell suspensions into ACSF.1T containing 1% BSA, centrifuged cells at 100g for 10 min, then
resuspended cells in ACSF.1T containing 0.2% BSA and 1 μg/mL DAPI prior to flow cytometry and
sorting on a FACSAria III (Becton-Dickinson). SYFP2 reporter brightness was measured as the ratio of
positive cell population mean fluorescence intensity, divided by the low mean fluorescence intensity of
autofluorescence in non-expressing cells. This measure of reporter brightness is more consistent than
positive cell population mean fluorescence intensity alone, due to differences in raw signal across days,
cytometers, and cytometer settings.

For single cell RNA-seq, we sorted single SYFP2+ cells into tubes and processed them via SMARTer
v4 using the workflow described previously[89], on 47 enhancer-AAV-injected mice. In each experiment
from one mouse injected with one single enhancer-AAV we sorted and profiled up to 48 cells per
experiment, and each measurement was taken from a distinct individual cell. After retroorbital
injections, enhancer-AAV SYFP2-expressing cells consisted of on average 7% of the positive brain
cells (range 0.1-20.1% of cells, n = 47 experiments). We sequenced single cell-derived SMARTer
libraries at 659996 ± 199038 (mean ± standard deviation) reads per library on an Illumina NovaSeq
instrument at Broad Institute (Cambridge, MA) or on an Illumina NextSeq instrument at Allen Institute
(Seattle, WA). We aligned the libraries to mm10 genome using STAR
(https://github.com/alexdobin/STAR), and also aligned them to the synthetic AAV transgene reference
construct using bowtie2 (https://bowtie-bio.sourceforge.net/bowtie2/index.shtml). From 2040 initial cells,
we excluded from analysis the libraries with poor library quality metrics, consisting of: firstly low-quantity
or degraded libraries (judged as less than 65% percentage of cDNA library sized greater than 400 bp,
consisting of 71 [3.4%] libraries in this study), and secondly those that lacked AAV transgene-mapping
reads (likely mis-sorted events, 23 [1.1%] of remaining libraries). Applying these filtering criteria yielded
a dataset for analysis of 1946 high-quality AAV transgene-expressing cells, with alignment rates of 92 ±
3% to mm10 genome and 4654 ± 1285 genes detected per cell (mean ± standard deviation). To assess
enhancer specificity within the cortex we mapped the high quality transgene-expressing SMARTer cells
to the SMARTer-based VISp cellular taxonomy generated by Tasic et al. using bootstrapped
hierarchical approximate nearest neighbor mapping[18,89], and quantified the specificity as the percentage
of positively sorted cells that mapped to the expected cell subclass (astrocytes or oligodendrocytes). To
test for significance of correlation of brightness by flow cytometry with expression levels by scRNA-seq,
we calculated Pearson's product-moment correlation coefficient by cor.test() function in R.

To understand different characteristics of different regional astrocyte populations we utilized
scrattch.mapping (https://github.com/AllenInstitute/scrattch) from the Allen Institute. To accomplish this

we first transformed these cells by principal component analysis and performed UMAP dimensionality
reduction on the first 40 principal components for visualization using the default scanpy parameters,
which clearly separated oligodendrocytes and regional groupings of astrocytes. For clustering
astrocytes we subset the dataset to astrocytes only, then identified the top 2000 genes ranked by
variance among them, recomputed UMAP projections from these high-variance genes, then performed
Leiden clustering[90] which identified VISp, MB/HB, and CBX astrocyte clusters as expected, and finally
identified differential genes among them (differential gene expression threshold false discovery rate
less than 5% and $\log_2$-fold change greater than 0.5) using scanpy
(https://scanpy.readthedocs.io/en/stable/). In doing so we detected two major subgroups of VISp
astrocytes that are distinguished by presence or absence of immediate-early gene markers (for
example, *Fos*, *Fosl2*, *Nr4a1*, *Irs2*, *Pde10a*, and *Pde7b*). This distinction may be an artifact of the cell
dissociation process for scRNA-seq; for the purposes of this study we collapse these cortical astrocyte
clusters. In order to understand the different regional characteristics of astrocyte populations we
mapped cells to whole brain taxonomy we mapped to the best-correlated mean-aggregated taxonomic
cluster from the ABC-WMB atlas[18] with 100 bootstrapped iterations using the top 10% of high-variance
genes and omitting a variable number of genes (10-50%) each round. We interpret the frequency of
correct mapping rounds as the mapping confidence. We also used CELLxGENE for single cell
visualization (https://github.com/chanzuckerberg/cellxgene). Spatial transcriptomic analysis was
performed as described in the recent whole brain transcriptomic taxonomy study[18], and cell type
location data was visualized using Cirrocumulus
(https://cirrocumulus.readthedocs.io/en/latest/index.html).

For determination of *Zic5* and *Sox10* differential gene expression between astrocytes and
oligodendrocytes, we used two-sided ANOVA on expression measurements from individually profiled
cells from all the experiments with no exclusion, and no covariates were tested. Testing for normality by
the Shapiro-Wilk test revealed that *Zic5* and *Sox10* expression are not normally distributed (*Zic5* W =
0.539, p-value < 2.2e-16; *Sox10* W = 0.958, p-value = 5.1e-16), so we used a non-parametric Wilcoxon
rank-sum test. No significance thresholds adjustments were made for multiple comparisons since only
one comparison was performed. For the comparison of *Sox10* versus *Zic5* expression (mean counts
per million +/- standard deviation, n cells): astrocyte *Zic5* expression 32 ± 71, n = 864; astrocyte *Sox10*
expression 0.3 ± 5, n = 864; oligodendrocyte *Zic5* expression 0.6 ± 6, n = 964, oligodendrocyte *Sox10*
expression 456 ± 276, n = 964.

Motif analysis. We performed de novo motif discovery from the sets of astrocyte and oligodendrocyte
enhancers that showed specific and strong expression patterns, excluding those enhancers scored as
weak. For astrocytes this list consisted of: AiE0375m, AiE0376h, AiE0376m, AiE0377m, AiE0380h,
AiE0381h, AiE0385h, AiE0385m, AiE0380h, AiE0390h, AiE0390m, AiE2120m, AiE2122m, AiE2160m,
and ProB12. For oligodendrocytes this list consisted of: AiE0361h, AiE0395h, AiE0395m, AiE0396h,
AiE0397m, AiE0398h, AiE0400m, AiE0401h, AiE0403h, AiE0407h, AiE0409h, AiE0409m, AiE0410h,
AiE0410m, and AiE0641m. We used MEME-CHIP[61] to identify recurrent de novo motifs in these sets of
sequences, using the parameter -meme-maxw 12, and comparing to a background set of random
sequence with the same nucleotide content. This analysis revealed one strong motif in each set of
sequences, as measured by its E-value, which is an estimate of the number of motifs expected by

chance to have as strong a log likelihood ratio as itself within the given sequences. These de novo

motifs where then mapped to known sequences using TomTom[61] which revealed several possible

matches to known motifs at significant p-values, but the strongest motif match (lowest p-value) in each

case is shown. In the case of the Zic family transcription factors, for simplicity we averaged together the

highly correlated strongest hits in the Zic family (JASPAR accession numbers MA0697.2, MA1628.1,

and MA1629.1 covering Zic1, Zic2, and Zic3), since Zic5 itself is not present in databases. In the case

of Sox10, the highly correlated Sox family members Sox4 and Sox11 (Uniprobe accession numbers

UP00062.1 and UP00030.1) showed slightly stronger motif match p-values than Sox10 (JASPAR

accession number MA0442.1), but these were excluded from analysis due to lack of expression in

almost all oligodendrocytes as observed by Tasic et al.[89] and in this study.

Enhancer-AAV testing in rat. The Allen Institute Institutional Animal Care and Use Committee (IACUC)

approved the following in vivo testing experiments in rat under protocol 2010. We procured timed-

pregnant female Sprague-Dawley rats from Charles River laboratories. We tattooed and injected ice-

anesthetized neonatal pups at P1 with 1.5e11 viral genomes of enhancer-AAV virus, diluted with PBS

to a total volume of 10 µL, unilaterally into the forebrain lateral ventricle (ICV delivery) with a 31-gauge,

4 point, 12º bevel 1 inch needle (custom ordered from Hamilton) and 25 µL capacity removable needle

syringe (Hamilton, 7636-01). Between injections we washed the needle and syringe with 100% ethanol,

and then nuclease-free water. We targeted the ICV space at 2 mm posterior to bregma, 2 mm lateral to

the anterior-posterior midline, and at a depth of 2 mm perpendicular to the surface of the skull. We

injected into the ventricle slowly over approximately 30 seconds. After injection, we held the needle in

place for approximately 10 seconds to prevent viral leakage, then slowly withdrew the needle at the

same relative angle as injection and then placed the animal onto a prewarmed heating pad in a clean

cage. We sacrificed pups at 18 days post injection, prior to weaning, and transcardially perfused with

1X PBS and then 4% PFA in PBS. We hemisected each brain and cryoprotected in 30% sucrose in

deionized water for a minimum of 24 hours before sectioning. We sectioned each brain at 30 µm

thickness using a sliding microtome (Leica part number SM2000R) on a leveled mount of Tissue-Tek®

O.C.T. Compound, collecting 3 sagittal planes separated by approximately 500 µm. We counterstained

sections with 1 µg/mL DAPI and 2.5 µg/mL propidium iodide (Thermo catalog # P1304MP) overnight at

4ºC and mounted in Vectashield HardSet Antifade Mounting Medium (Vector Laboratories # H-1400-

10) prior to imaging by epifluorescence.

Macaque enhancer-AAV testing. Macaca nemestrina (pig-tailed macaque) animals were housed and

injected at the Washington National Primate Center according to NIH guidelines and as approved by

the University of Washington Animal Care and Use Committee under UW IACUC protocol #41-6701.

These animals received several intraparenchymal injections under general anesthesia at spatially

distinct sites located at least ~1cm apart throughout the brain. During injection, over the course of 10

minutes we expelled a total of approximately 1e11 gc iodixanol gradient-purified PHP.eB-packaged

viral vectors in a total volume of 5 µL at 10 depths ranging from 200 to 2000 microns deep in the

animals. After injection the animal rested for 10 minutes between injections. These numbers are

approximate and timing, volume, and depths may be adjusted according to animal anatomy and

surgical considerations. The experiments described here result from four injection sites in three *Macaca*

nemestrina animals (two males, one female). We harvested tissue from these animals after necropsy at
38, 47, and 113 days post injection.

After locating the injection sites and cutting out tissue blocks about 1-2cm on each side surrounding the
injection sites, we fixed these tissue blocks in 4% PFA for 24 hrs. Then we rinsed the blocks with PBS,
cut 350 µm thick slices on the sliding microtome, and postfixed the slices in 4% PFA for 2 hours at
room temperature (RT), and then washed three times in PBS for 10 min each. For ISH analysis using
HCR we transferred slices to 70% EtOH at 4°C for a minimum of 12 hours, and up to 30 days. To stain
we first incubated the slices in 8% SDS in PBS at RT for two hours with agitation, then washed the
slices at RT with 5X sodium chloride sodium citrate (SSC) for three hours, exchanging with fresh 5X
SSC every hour. Next we performed HCR v3.0 using reagents and a modified protocol from Molecular
Technologies and Molecular Instruments[91]. We first incubated slices in pre-warmed 30% probe
hybridization buffer (30% formamide, 5X sodium chloride sodium citrate (SSC), 9 mM citric acid pH 6.0,
0.1% Tween 20, 50 µg/mL heparin, 1X Denhardt's solution, 10% dextran sulfate) at 37°C for 5 min.
Then we exchanged hybridization buffer for hybridization buffer containing probes added at a
concentration of 2 nM. Molecular Instruments designed the probes using the following accession
numbers: SLC17A7 – XM_011768126.1, GAD1 – XM_011744029.1, FGFR3 – XM_011744842.2,
SOX10 – XM_011712410.2. Hybridization proceeded overnight at 37°C, and afterwards we washed the
tissue thrice with 5X SSC for 10 minutes each (total 30 minutes), then 30% probe wash buffer (30%
formamide, 5X SSC, 9 mM citric acid pH 6.0, 0.1% Tween 20, 50 µg/mL heparin) for one hour at 37°C.
Then we exchanged probe wash buffer with 5X SSC, then amplification buffer (5X SSC, 0.1% Tween
20, 10% dextran sulfate) for 5 min at room temperature. Meanwhile we pooled even and odd
amplification hairpins for each of the three genes and snap-cooled them by heating to 95°C for 90
seconds then cooling to room temperature for 30 min, and afterwards we added the snap-cooled
hairpins to amplification buffer at a final concentration of 60 nM, and finally centrifuged at 18000g for 1
minute. Then we incubated tissue slices in amplification solution containing amplification hairpins for 4
hours at room temperature, followed by staining in DAPI (10µg/mL in 2X SSC) for 1 hour at room
temperature, and finally washing twice for 30 min in 2X SSC at room temperature before imaging. We
prepared a fresh aliquot of 67% 2,2'-Thiodiethanol (TDE) solution for use as a clearing and immersion
fluid by mixing ≥99% TDE (Sigma-Aldrich) with deionized water to create a 67% TDE solution with a
refractive index of 1.46. We transferred slices to 67% TDE and allowed them to equilibrate for at least 1
hour at room temperature prior to imaging on a confocal microscope (Olympus FV-3000). For IHC
analysis of injected NHP tissue we subsectioned 350 µm thick slices to 30 µm and stained as for
mouse tissue.

Stereotaxic injection and fiber implant surgery. Virus injection and optic fiber implantation surgery was
performed in C57BL/6J mice (The Jackson Laboratory, #000664) at around P60. Mice were
anesthetized with isoflurane and monitored throughout the surgery using breathing rate and tail pinch.
The skin above the skull surface was removed to make room for the fiber implant and headframe. After
leveling the skull, a craniotomy was drilled above the injection and fiber coordinates (AP: 1.2 mm, ML: -
1.3 mm, DV: 4.1 mm). First, a glass pipette positioned at the injection coordinates was lowered through
the craniotomy and virus injection was performed (100 nl, titer: 4e13 gc/mL). Once the injection was
complete, the pipette was slowly raised, and the optic fiber probe was positioned at the same AP and

ML coordinates as the injection. The tip of the fiber was then lowered to 100 μm above the injection site
and glued in place where the base of the fiber ferrule contacts the skull. A custom headframe was then
glued to the skull to allow head-fixed behavior and imaging. After surgery, the mouse was returned to
the home cage and allowed to recover for at least two weeks prior to start of water restriction for
behavior and imaging.

Dynamic foraging reinforcement learning task. Water-restricted and head-restrained mice were trained
to perform a reinforcement learning task where they freely choose between two lick ports that delivered
a water reward with nonstationary probabilities. This is a variation on the task described in Bari et al.[92].
The base reward probability of both lick ports summed to 0.6 where the probabilities of the two lick
ports were selected from two sets of ratios (0.53/0.07, 0.51/0.09). Block lengths that corresponded to
each ratio lasted for about 30 trials (min trials per block: 40, max trials per block: 60). Each trial began
with an auditory "go cue" that signaled the start of a trial. The mouse was free to choose between the
left or right lick port immediately after the "go cue". The trials were separated by a variable inter-trial-
interval (range between 1-7 seconds). The data shown in this study was from a two-hour behavior
session that consisted of 438 trials (170 rewarded trials).

Fiber photometry and analysis. Fiber photometry was performed using a commercially available
photometry system (Neurophotometrics LLC, FP3002). A 470 nm LED was used to excite the
iAChSnFR fluorophore, Venus, and the emitted fluorescence signals were collected using a CMOS
camera. The 470 nm excitation was interleaved with a 415 nm LED as an isosbestic control to remove
motion artifacts. Bonsai acquisition software was used to record the photometry signals as well as the
behavior trigger signals events (go-cue, left and/or right lick choices, reward/no reward) for offline
alignment of imaging data to behavioral events. Prior to start of acquisition, an ROI was drawn over the
fiber image seen on the camera, and fluorescence intensity within this ROI was averaged for real-time
signal visualization and offline analysis. First, the fiber photometry acquisition was started, following
which the behavior task was initialized. Photometry signals were analyzed using custom python scripts.
First, the fluorescence signal was detrended for photobleaching using a fourth order polynomial
function and then corrected for motion using the control signal from the 415 nm excitation using
standard photometry analysis techniques[93]. Acetylcholine signal changes were calculated as a change
in fluorescence intensity over the mean fluorescence (ΔF/F as a percentage). The photometry signals
were then aligned to behavior events using simultaneously acquired TTL readouts of behavior events
(go-cue, left and/or right lick choices, reward/no reward) using a NI USB card. These behavior events
were then used to calculate trial averaged traces of rewarded and unrewarded signals.

Data Availability:

All AAV viral vector plasmids are freely available for research use at Addgene (addgene.org/). Mouse scRNA-seq generated from this study are available at GEO with the accession number GSE235987 (https://www.ncbi.nlm.nih.gov/geo/). Mouse primary testing data are available at the Allen Genetic Tools Atlas (https://portal.brain-map.org/genetic-tools/genetic-tools-atlas). Serial two photon tomography datasets will be made available through the Brain Imaging Library (https://www.brainimagelibrary.org/). All other data will be made available upon request.

Ethics Declarations:

Competing interests

Acknowledgements

We would like to thank the Washington National Primate Center and staff for animal care, as well as the supporting grant P51OD010425 from the National Institutes of Health to support primate research. The WaNPRC SPF *M. nemestrina* colony is supported by grant U42OD011123 from the NIH Office of Research Infrastructure Programs. We would like to acknowledge Kathryn Gudsnuk for programmatic support. This work was supported by the following grants: RF1MH114126-01 from the National Institute of Mental Health to BPL, JTT, and ESL; UG3MH120095-01, -02, -03 from the National Institute of Mental Health to BPL, JTT, ESL, and FKK, UF1MH128339-01 from the National Institute of Mental Health to BT, TB, TLD, BPL, and JTT, RF1MH121274-01 from the National Institute for Mental Health to BT and U19MH114830 to HZ. We also would like to acknowledge the estate of Paul G. Allen for his vision, encouragement, and support.

Figures and Figure Legends

825 **Figure 1: Astrocyte and oligodendrocyte enhancer discovery from single cell epigenetics.**
826

(**A**) Example astrocyte- and oligodendrocyte-specific peaks near the loci of astrocyte-specific gene
AQP4 and oligodendrocyte-specific gene *OPALIN*, identified in human MTG snATAC-seq data[33].

(**B**) Differing approaches to identify candidate enhancers. Specific accessibility peaks are depicted as
peaks, and specifically demethylated regions are depicted as troughs. Schemes not utilizing a particular
data modality are shown as "Agnostic". Marker gene selection criteria can use accessibility from either
mouse or human. Icons represent identification schemes; gold star candidate enhancers undergo more
stringent criteria than those with gold squares (see Methods for details).

(**C**) Workflow for enhancer cloning, packaging, screening, and validation. Enhancers are cloned into a
pAAV plasmid upstream of a minimal human beta-globin promoter and SYFP2 reporter, and plasmids
are packaged into PHP.eB AAVs. Enhancer-AAVs are injected intravenously into retro-orbital sinus,
and expression is assessed by imaging. Promising enhancer-AAVs then go on to secondary validation
experiments consisting of cross-species validation, molecular characterization by IHC and/or
multiplexed FISH, and flow cytometry for single cell RNA-seq.

(**D-E**) Candidate identification schemes and summarized screening results for astrocyte-specific (**D**)
and oligodendrocyte-specific enhancers (**E**). Overall 50% (25/50) of candidate astrocyte enhancers,
and 49% (21/43) of candidate oligodendrocyte enhancers, show the intended specificity after
intravenous delivery of PHP.eB AAVs. Testing result bar: Y = yes, enhancer-AAV gives strong or
moderate on-target expression pattern; N = no, enhancer-AAV fails to express; W = weak on-target
expression pattern; Mx = mixed expression pattern consisting of on-target cells plus unwanted neuronal
populations; Off = off-target expression pattern; ND = no data. Note both enhancers giving
strong/moderate ("Y") and weak ("W") specific expression are grouped here for overall success rate
analysis.

A AiE0380h — Most of CNS

B *GFAP* promoter — Most of CNS

C AiE0385m — CH — Regional — Native SYFP2 fluorescence

1 mm

D 500 µm — SpC

E

F

G VISp — SYFP2 / Sox9 — 50 µm

H

I

J CBX — 50 µm

K

L

M Pons — 50 µm

N

O

P

Most of CNS | Scattered | Weak | Mixed spec. | Regional

Astrocyte specificity (%)
- IHC (Sox9+)
- scRNA-seq

Completeness by IHC (%)
- >50%
- 10-50%
- 1-10%
- <1%

Astrocyte specificity (%) y-axis: 0, 20, 40, 60, 80, 100

Bar labels:
- AiE0380h: 317/521 (n=2); 76/77 (n=2)
- AiE0390m: 807/807 (n=3); 78/82 (n=2)
- AiE0390h: 452/454 (n=2); 86/86 (n=2)
- AiE0387m: 473/473 (n=3); 44/46 (n=1)
- proB12: 449/466 (n=3); 40/40 (n=1)
- *GFAP* promoter: 411/419 (n=2); 85/87 (n=2)
- AiE0376h: 165/170 (n=2); 35/37 (n=1)
- AiE0371h: 2/2 (n=1); nd
- AiE0374m: 75/95 (n=2); 60/82 (n=2)
- AiE0373m: nd; 12/13 (n=1); 12/14 (n=1)
- AiE0386m: 85/99 (n=1); 42/44 (n=1)
- AiE2117m: 153/156 (n=1); nd
- AiE2118m: 244/246 (n=2)
- AiE2121m: 236/268 (n=2); nd
- AiE0385m: VISp 312/319 (n=2), 74/76 (n=2); Pons 17/42 (n=2), nd
- AiE2120m: VISp 353/357 (n=2), nd; Pons 63/100 (n=2), nd
- AiE2122m: VISp 122/123 (n=1), nd; Pons 1/2 (n=1), nd
- AiE0381h: VISp 0/33 (n=3), nd; Pons 204/208 (n=3), 69/74 (n=2)
- AiE0375m: VISp 2/3 (n=3), nd; CBX 460/461 (n=3), 79/80 (n=2)

VISp | VISp Pons | VISp Pons | VISp Pons | VISp Pons | VISp CBX

Q AiE0387m — dSTR, CA1, CBX — Most of CNS — Native SYFP2 fluorescence

1 mm

dSTR — 50 µm | CA1 | CBX

Figure 2: A collection of astrocyte-specific enhancer-AAV vectors with varying regional specificities and expression densities.

(**A-B**) Astrocyte-specific enhancer-AAVs marking many astrocytes throughout most of the CNS. AiE0380h (**A**, n = 8 animals tested) and *GFAP* promoter (**B**, n = 3) mark many astrocytes throughout gray matter in FB, MB, HB, and CBX.

(**C**) AiE0385m shows a regional pattern labeling astrocytes in cerebrum (CH) but not in MB, HB, or CBX (n = 6).

(**D-F**) Astrocyte-specific enhancer-AAVs labeling astrocytes in lumbar SpC. AiE0380h (**D**) and *GFAP* promoter (**E**) label many astrocytes in SpC gray matter, but AiE0385m (**F**) does not label SpC astrocytes.

(**G-O**) Positive confirmation of molecular astrocyte identity across brain regions. SYFP2+ astrocytes are colabeled with anti-Sox9 immunoreactivity in VISp, CBX, and Pons. Images in G-O represent n = 2 animals each assessed by IHC.

(**P**) Quantification of specificity for astrocytes by astrocyte enhancer-AAVs. Specificity and completeness for astrocyte labeling by enhancer-AAVs was quantified by costaining with anti-Sox9 antibody in VISp, Pons, and CBX. Specificity is defined as the number of SYFP2+Sox9+ / total SYFP2+ cells x 100%. Completeness is defined as the number of SYFP2+Sox9+ / total Sox9+ cells x 100%. Brains from one to three mice per condition were analyzed, with range 131-827 cells counted (median 311) per brain region analyzed. AiE0375m-labeling was only quantified in the Purkinje cell layer of CBX, not in the granule or molecular layers. Specificity was also quantified by scRNA-seq, defined as the percentage of sorted SYFP2+ cells mapping as astrocytes within the VISp molecular taxonomy[89]. Overall, specificity is high for many astrocyte-specific vectors, with "Scattered" and "Weak" vectors showing low completeness, and "Regional" vectors showing more completeness in certain regions. One to three animals were analyzed per vector per modality, as indicated. The same IHC specificity data for *GFAP* promoter-SYFP2 are repeated in **Figure 6B.**

(**Q**) Distinct astrocyte morphologies throughout the brain with AiE0387m enhancer-AAV targeting "Most of CNS". Images were acquired on a serial two-phtoton tomography blockface imaging platform (STPT, TissueCyte, n = 4 tested).

Abbreviations: CH cerebrum, dSTR dorsal striatum, CA1 cornu ammonis 1, CBX cerebellar cortex, SpC spinal cord, VISp primary visual cortex.

Figure 3: A collection of oligodendrocyte-specific enhancer-AAV vectors with varying levels of expression.

(**A-C**) Oligodendrocyte enhancer-AAVs marking many oligodendrocytes throughout most of the CNS. AiE0410m (**A**, n = 7 animals tested), AiE0409h (**B**, n = 5), and AiE0400h (**C**, n = 4) label many oligodendrocytes throughout FB, MB, HB, and CBX, but at differing expression levels.

(**D-F**) Oligodendrocyte enhancer-AAVs marking oligodendrocytes in lumbar SpC. AiE0410m (**D**), AiE0409h (**E**), and AiE0400h (**F**) mark oligodendrocytes in gray and white matter of SpC, but at different intensities.

(**G-O**) Positive confirmation of molecular oligodendrocyte identity across brain regions. SYFP2+ oligodendrocytes are colabeled with CC1 immunoreactivity in VISp, CBX, and Pons. Two to three animals per enhancer-AAV were analyzed.

(**P**) Quantification of specificity for oligodendrocytes by oligodendrocyte enhancer-AAVs. Specificity and completeness for oligodendrocyte labeling by enhancer-AAVs was quantified by costaining with CC1 antibody in VISp. Specificity is defined as the number of SYFP2+CC1+ / total SYFP2+ cells x 100%. Completeness is defined as the number of SYFP2+CC1+ / total CC1+ cells x 100%. Brains from one to three mice per condition were analyzed, with range 101-332 cells counted (median 147) per brain region analyzed. Specificity was also quantified by scRNA-seq, defined as the percentage of sorted SYFP2+ cells mapping as oligodendrocytes within the VISp molecular taxonomy[89]. One to three animals were analyzed per vector per modality, as indicated. Overall, specificity is high for many oligodendrocyte-specific vectors, with "Weak" vectors showing low completeness.

(Q) Myelinating oligodendrocyte morphologies throughout the brain with AiE0410m. Sections were visualized with STPT (n = 2 tested).

Abbreviations: SpC spinal cord, VISp primary visual cortex, CBX cerebellar cortex, LSX lateral septal complex, MY medulla.

Figure 4: Transcriptomic identities of prospectively targeted astrocytes and oligodendrocytes.

(**A-C**) Groups of transcriptomically profiled single cells, as visualized by UMAP. Single cells labeled by various astrocyte- and oligodendrocyte-specific enhancer-AAVs (n = 1946 quality-filtered cells) were profiled from 47 brains in 47 independent experiments by SMARTerV4[89], one to three animals per enhancer. Libraries were aligned to mm10 and transformed into UMAP space for visualization, with coloring by enhancer (**A**), mapped taxonomic cell type cluster (**B**), and taxonomic mapping confidence (**C**). Overall CTX astrocytes group away from CTX oligodendrocytes as expected, and MB/HB astrocytes and Bergmann glia astrocytes group away from CTX astrocytes, consistent with recent results[18]. Note that AiE0381h- and AiE2160m-labeled astrocytes were dissected from MB/HB region, and AiE0375m-labeled Bergmann glia were dissected from CBX region, but the remainder of the cells were dissected from VISp.

(**D-E**) Quantifications of taxonomic cell type cluster mapping by enhancer vector. Prospectively labeled astrocytes from all enhancer-AAV vectors dissected from VISp predominantly map to cluster "5112 Astro-TE NN_3", whereas those from MB/HB dissections (AiE0381h and AiE2160m) predominantly map to cluster "5109 Astro-NT NN_2", and AiE0375m-labeled astrocytes from CBX dissections predominantly map to cluster "5102 Bergmann NN". In contrast, all prospectively labeled oligodendrocytes predominantly map to cluster "5158 MOL NN". Cluster identities are from a recent whole mouse brain taxonomy study[18].

(**F-H**) De novo motif detection in astrocyte- and oligodendrocyte-specific enhancer sequences using MEME-CHIP[61] identifies one strong consensus motif in each set of sequences (top). These de novo motifs were mapped against databases of known TF motifs using TomTom (bottom), which identified the top hits as the Zic family consensus motifs for astrocytes, and Sox family motif for oligodendrocytes (Sox10 shown). These TFs (Zic5 and Sox10) show highly specific expression differences between astrocytes and oligodendrocytes from prospective scRNA-seq profiling (**H**). *** non-parametric Wilcoxon rank-sum test W = 577624 (*Zic5*) or W = 9838 (*Sox10*), p < 1e-16 each.

(**I-N**) Intrinsic SYFP2 expression from the indicated enhancer-AAVs after retro-orbital administration. Images were generated by STPT. Boxes in I and L correspond to K and N, respectively. Scale in I and K is 500 μm.

(**O-R**) MERFISH data showing the distribution of three astrocyte cell types revealed by single cell gene expression from the whole mouse brain[18].

Abbreviations: CTX cerebral cortex, STR striatum, GPe globus pallidus, external segment.

Figure 5: Optimizing astrocyte and oligodendrocyte enhancer strength.

(**A**) Native Enhancer and for Enhancer_3xC2 vector designs. The central approximate third of the enhancer (the "Core2" element) is marked by dark hatches, and this element is triply concatemerized in the Enhancer_3xC2 vector. Alternatively, the first or third segment ("Core1" or "Core3") may be concatemerized (determined empirically).

(**B-J**) Dramatic increase in expression levels while maintaining specificity using Enhancer_3xC1/2 vector designs. Brains from mice injected with the Enhancer or Enhancer_3xC1/2 vectors were processed and imaged in parallel in these experiments. (**H-J**) Zoom in view of AiE0390m- and AiE0390m_3xC2-injected mouse VISp shows high specificity for morphological astrocytes throughout cortical layers in both cases. Data represent one to two animals per vector with parallel tissue processing and imaging.

(**K**) Quantification of specificity for astrocytes by concatemer astrocyte enhancer-AAVs within VISp by IHC and scRNA-seq as described in **Figure 2P**. The same IHC data for AiE0390m_3xC2 are repeated in **Figure 6B**.

(**L-M**) Direct correlated quantification of enhancer strength by flow cytometry and scRNA-seq, for both astrocyte- (**L**) and oligodendrocyte-specific (**M**) enhancer-AAVs. The left (blue) y-axis represents the log-transformed vector transgene reads per million in individual sorted scRNA-seq-profiled cells. The right (brown) y-axis represents the log-transformed SYFP2 signal intensity of positively gated vector-expressing cells observed on the flow cytometer, quantified as the fold signal of positive cells normalized to non-expressing cell autofluorescence (taken as background). Points represent individual cells observed by scRNA-seq and by flow cytometry, visualized also as violins, and with the horizontal bar representing mean expression levels across all cells expressing that enhancer-AAV, across one to three replicate experiments per vector. Across all experiments, we observe significant correlation between mean expression intensity at the RNA level by scRNA-seq, and mean SYFP2 reporter expression by signal intensity (astrocytes: n = 26 experiments, Pearson correlation coefficient [PCC] 0.63, t = 3.97, df = 24, p = 0.00057; oligodendrocyte n = 22 experiments, PCC 0.53, t = 2.82, df = 20, p = 0.011; correlation t-tests). Furthermore, 3xC astrocyte enhancers are among the strongest enhancers we have characterized, typically several fold stronger than their native counterparts. One to three animals were analyzed per enhancer-AAV vector (the same dataset described in **Figure 4A**).

Abbreviations: VISp primary visual cortex.

Figure 6: Robust specificities of astrocyte enhancer-AAVs with a difficult transgene cargo.

(**A-B**) Specific delivery of a difficult transgene cargo with an optimized astrocyte-specific transgene.
Mice received the indicated enhancer-AAVs driving either SYFP2 or NeuroD1-mCherry by intravenous
PHP.eB AAVs (5e11 gc per animal), and expression analyzed in brain by IHC with NeuN neuronal
marker or Sox9 astrocyte marker. Specificity for neurons and astrocytes was quantified as
(Reporter+[NeuN or Sox9]+ cells) / (All Reporter+ cells) in primary visual cortex (VISp, Layer 5 cells
shown, n=2-3 animals per condition). The same IHC specificity data for *GFAP* promoter-SYFP2 and
AiE0390m_3xC2-SYFP2 are repeated in **Figures 2P** and **5K**, respectively. Two to three animals were
analyzed per condition (each dot represents one animal). *** ANOVA with Tukey's post hoc adjustment
for multiple comparisons, F = 647, p < 1e-7, for *GFAP* promoter-driven NeuroD1-mCherry compared to
every other condition.

990
991

A Predictable expression across body organs

Human chromatin accessibility atlas (Zhang et al., 2021)

Normalized accessibility

Predicted liver expression

Strong
$ Little or none
? No prediction

B Whole liver transverse section

500 μm

AiE0371h

AiE0387m

C

Liver SYFP2 expression

100 μm

***

Figure 7: Predictability of astrocyte enhancer-AAV expression patterns across body organs.

(**A**) Accessibility profiles of astrocyte-specific enhancers in the human whole-body accessibility atlas[70]. Single-cell profiles were grouped within each tissue into pseudo-bulk aggregates, then normalized according to the signal (reads in peaks) within the dataset. Accessibility profiles are likely to predict enhancer activities within each tissue. Focusing on liver, some astrocyte-specific enhancers are predicted to have "Strong" expression, and some are predicted to have very "Little or none" expression. In contrast, accessibility atlases do not predict expression of *GFAP* promoter across tissues.

(**B**) Whole livers from mice injected intravenously with AiE0371h- and AiE0387m-enhancer-AAV vectors, stained with anti-GFP antibody. AiE0371h has high liver accessibility, is predicted to have high liver expression, and shows many strong SYFP2-expressing hepatocytes throughout the liver as predicted. In contrast, AiE0387m has little liver accessibility and so is predicted to have little liver expression, and in fact shows few positive SYFP2-expressing hepatocytes as predicted.

(**C**) Agreement between liver expression predictions and liver expression measurements across several astrocyte-specific enhancer-AAV vectors. AiE0371m, AiE0371h, AiE0381h, and AiE0386m all show many SYFP2-expressing hepatocytes as predicted. AiE0390m, AiE0390h, AiE0375m, AiE0387m, and AiE0380h all show few weak SYFP2-expressing hepatocytes as predicted. *GFAP* promoter shows few but strongly expressing hepatocytes, which was not predictable from the accessibility atlases. Liver images in **B** and **C** represent two to four mice analyzed for each vector (each dot represents one mouse). *** non-parametric Wilcoxon rank sum test, W = 153, p < 1e-4 for the comparison of Strong versus "Little or none"-predicted enhancers.

A

Predictable expression
across disease status

Astrocyte enhancer-AAVs delivered IV

Scn1a[+/-] or littermate controls

↓

Tissue analysis at P42

B

| | *Scn1a*[+/+]; AiE0390m | *Scn1a*[+/-]; AiE0390m | *Scn1a*[+/+]; *GFAP* promoter | *Scn1a*[+/-]; *GFAP* promoter |

Anti-GFP   200 µm

Anti-GFAP

Anti-GFP
Anti-GFAP   25 µm

ML

GCL

PoL

C

DGC neruons labeled (%)

Scn1a[+/+] AiE0390m

Scn1a[+/-] AiE0390m

Scn1a[+/+] *GFAP* pr.

Scn1a[+/-] *GFAP* pr.   **

Figure 8: Predictability of astrocyte enhancer-AAV expression patterns across disease states.

(**A-C**) Testing fidelity of enhancer-AAV expression across disease states. We used Dravet syndrome model $Scn1a^{+/-}$ mice to induce epilepsy-associated hippocampal gliosis, injected enhancer-AAVs prior to the critical period, and analyzed tissue for expression patterns after the critical period (**A**). We assessed hippocampal gliosis with anti-GFAP antibody and enhancer-AAV expression with anti-GFP antibody (**B**). AiE0390m maintained specific expression and similar levels in hippocampal astrocytes regardless of epileptic gliosis. In contrast, *GFAP* promoter expression strongly increased in gliotic astrocytes, and also was observed in dentate gyrus granule cells. Red dashed rectangles indicate the position of the expanded zoomed view, and the curved red arrows indicate a rotated view for one zoom. (**C**) Quantification of DGC labeling reveals a much greater number of DGC neurons labeled with GFAP promoter in Dravet syndrome model mice. Two to four animals were analyzed per condition (each dot represents one mouse). ** ANOVA with Tukey's post hoc adjustments, F = 16, p < 0.01 for the *GFAP* promoter in $Scn1a^{+/-}$ mice compared to each other condition.

Abbreviations: ML molecular layer, GCL granule cell layer, PoL polymorphic layer.

Figure 9: Astrocyte specific sensing of cholinergic signals in the nucleus accumbens during behavior.

(**A**) AiE0390m_3xC2 driving expression of iAChSnFR. Enhancer vector is cloned into a pAAV plasmid and packaged into PHP.eB AAVs.

(**B**) Coronal section showing stereotaxic injection of enhancer virus expressing iAChSnFR in the nucleus accumbens (injection coordinates: AP: 1.2, ML: 1.3, DV: 4.1).

(**C**) Behavior and imaging experiment setup. Top: dynamic foraging behavior task schematic. Bottom: Fiber photometry instrumentation schematic and fiber location in a coronal section.

(**D**) Fiber photometry signals of acetylcholine fluctuations during task performance. Top: ~30 min segment of a 2-hour session of dynamic foraging. Black dots represent the auditory cue, red dots represent time of first lick, blue dots represent water reward delivery. Bottom: 100 second (980-1080 seconds) zoom in on above session with 6 individual trials (4 rewarded and 2 unrewarded trials). Data from one mouse shown, representative of three mice assayed.

(**E**) Trial-averaged signals of rewarded and unrewarded trials aligned to time of first lick (mean±sem).

Figure 10: Astrocyte- and oligodendrocyte-specific Cre AAVs.

(**A**) Design and testing of astrocyte- and oligodendrocyte-specific attenuated Cre-expressing enhancer-AAVs.

(**B-D**) Specific Cre recombination in astrocytes driven by AiE0390m_3xC2. Ai14 reporter recombination is observed in astrocytes throughout the mouse brain (**B**). Recombination within VISp is highly specific (**C**), as evidenced by highly specific and complete recombination in Sox9+ astrocytes (**D**).

(**E-G**) Specific Cre recombination in astrocytes driven by AiE0410m. This vector also includes three 3'UTR mir126 binding sites to eliminate unwanted expression in mir126-expressing endothelial cells[75]. Ai14 reporter recombination is observed in oligodendrocytes throughout the mouse brain (**E**). Recombination within VISp is highly specific (**F**), as evidenced by highly specific and complete recombination in CC1+ oligodendrocytes (**G**).

(**H-I**) Quantification of recombination specificity (**H**) and completeness (**I**). Specificity is defined as (tdTomato+Sox9 or CC1+ cells) / (all tdTomato+ cells) x 100%. Completeness is defined as (tdTomato+Sox9 or CC1+ cells) / (all Sox9 or CC1+ cells) x 100%. Two to three animals were analyzed for each condition (each dot represents one mouse).

A

Neonatal rat ICV injections

Neonatal rat ICV injections

B L1   L23   L4   L5   L6   200 µm

AiE0390m_3xC2, 4x2C miRNA BS

SYFP2   Sox9   50 µm

Rat astrocyte specificity 98%

C AiE0390m_3xC2, 4x2C miRNA BS   2 mm

SYFP2 in rat brain

D L1   L23   L4   L5   L6   200 µm

AiE0641m

mTFP1   CC1   50 µm

Rat oligodendrocyte specificity 91%

E

Multisite intraparenchymal macaque brain injections

Multisite intraparenchymal NHP injections

F AiE0410m   Pia   500 µm

NHP MO CTX

G SYFP2   Nissl   50 µm

H SYFP2   NeuN

I SYFP2   25 µm

White matter

SOX10   Specificity 94%   Completeness 36%

J AiE0390m   500 µm   Pia

White matter

NHP SS CTX

K SYFP2   NeuN   GFAP   50 µm

L SYFP2   NeuN   GFAP

M SYFP2   25 µm

FGFR3   Specificity 92%   Completeness 51%

N AiE0390m_3xC2 4x2C miRNA BS   1 mm   **O** Merge w/ DAPI

Pia

White matter

NHP MO CTX   Specificity 99% Completeness 45%

Anti-GFP

Anti-SOX9   50 µm

P AiE0390h   500 µm   **Q** Merge w/ DAPI

Pia

White matter

NHP SS CTX   Specificity 98% Completeness 66%

Anti-GFP

Anti-SOX9   50 µm

1068 **Figure 11: Genetic targeting of astrocytes and oligodendrocytes across species.**

1069

(**A**) Testing enhancer-AAV vectors by neonatal rat ICV injections.

(**B-C**) Validation of astrocyte-specific enhancer-AAV vectors in rat by IHC with Sox9 antibody.
AiE0390m_3xC2 vector also incorporates 4X2C 3'UTR miRNA binding sites to prevent any off-target
labeling in excitatory neurons[76]. Each image represents one animal tested.

(**D**) Validation of oligodendrocyte-specific enhancer-AAV vector in rat. AiE0641m shows specific
expression in CC1+ VISp oligodendrocytes. Each image represents one animal tested.

(**E**) Multiple stereotactic intraparenchymal injections into macaque brain.

(**F-I**) Prospective labeling of macaque oligodendrocytes in vivo. AiE0410m enhancer-AAV vector gives
widespread labeling of oligodendrocytes throughout the depth of motor cortex (**F**). Most labeled
macaque oligodendrocytes exhibit multipolar ramified morphology indicative of local axon myelination
(**G**). Some labeled oligodendrocytes exhibit morphologies suggesting wrapping around wider tubular
structures highlighted with dashed white lines (**H**). SYFP2-expressing cells of both morphological types
express the oligodendrocyte/OPC marker *SOX10* with high specificity (**I**). One animal tested.

(**J-Q**) Prospective labeling of macaque astrocytes in vivo. AiE0390m enhancer-AAV vector gives
widespread labeling of astrocytes throughout the depth of somatosensory cortex (**J**). A few large Layer
5 pyramidal neurons are also labeled. Labeled astrocytes show the expected bushy morphology and
GFAP immunoreactivity of astrocytes in parenchyma (**K**) and sometimes reside near walls of large-
diameter tubular structures (**L**). SYFP2-expressing astrocytes express the astrocyte marker *FGFR3*
with high specificity (**M**). In another animal, an injection of AiE0390m_3xC2 vector with 4X2C 3'UTR
miRNA binding sites gives high specificity (99%, **N**) for SOX9+ astrocytes in MO (**O**), without any large
Layer 5 pyramidal neurons labeled. In another animal, AiE0390h vector gives high specificity (98%) of
labeling for SOX9+ astrocytes (**P-Q**). Each vector represents one animal tested.

Abbreviations: VISp primary visual cortex, MO motor cortex, SS somatosensory cortex.

1093

Figure 1—figure supplement 1: Epigenetic characterization of candidate enhancers in additional chromatin accessibility datasets.

(**A-D**) Accessibility profiles of all tested candidate human astrocyte-specific (**A**, **C**) and human oligodendrocyte-specific (**B**, **D**) enhancers. Human enhancer regions are characterized in the datasets of Mich et al.[33] (**A-B**), who performed snATAC-seq on neurosurgical MTG samples, and of Fullard et al.[94] (**C-D**), who performed bulk ATAC-seq on neuronal (sorted NeuN$^+$) and non-neuronal (sorted NeuN$^-$) nuclei from dissections spanning multiple regions of human postmortem forebrain. Overall, many candidate astrocyte- and oligodendrocyte-specific enhancers show accessibility specific to non-neuronal cells across much of the human forebrain. For each genomic region we show their peak nomination scheme matching to **Figure 1B**, enhancer name.

(**E-F**) Screening results from testing human candidate enhancers (repeated from **Figure 1D-E**, provided again for visualization). Testing result bar: Y = yes, enhancer-AAV gives strong or moderate on-target expression pattern; N = no, enhancer-AAV fails to express; W = weak on-target expression pattern; Mx = mixed specificities consisting of on-target cells plus unwanted neuronal populations; Off = off-target expression pattern, ND = no data.

(**G-J**) Accessibility profiles for all tested candidate mouse astrocyte-specific (**G**, **I**) and oligodendrocyte-specific enhancers (**H**, **J**). Mouse enhancer regions are characterized in the VISp scATAC-seq dataset of Graybuck et al.[34] (**G-H**) and in the full mouse cerebrum dataset of Li et al.[56] (**I-J**).

(**K-L**) Screening results from testing mouse candidate enhancers (repeated from **Figure 1D-E**, provided again for visualization).

Figure 2—figure supplement 1: Full screening results of all candidate enhancer-AAVs targeting astrocytes.

We injected mice with the indicated enhancer-AAV vectors between P42 and P56, then after 3-4 weeks we harvested brains, sliced them on a sliding microtome with freezing stage at 30 µm thickness, co-stained the sections with DAPI, then mounted them with Vectashield Vybrance. Insets show a full cortical column from VISp (primary visual cortex), and in some cases also the labeling in MB (midbrain) or HB (hindbrain) or CBX (cerebellar cortex) is also shown. Astrocyte-specific enhancer-AAV vectors are broadly grouped by expression pattern into the following categories: "Most of CNS astrocytes", "Regional" meaning present at medium-to-high levels in one or more broad brain regions but not all, "Scattered" meaning a few astrocytes are strongly labeled throughout the brain, "Weak" meaning many astrocytes throughout the brain are labeled at low level, "Mixed specificities" meaning one or more off-target neuron populations are also labeled in addition to astrocytes, and "No astrocyte expression" meaning failure to detect any clear astrocytes in these whole-brain sagittal images. These epifluorescent screening images represent n = 1 to 27 animals tested per enhancer (median 2 animals), and were sometimes taken on multiple different microscopes (see **Table S2** for full summary of screened animals).

1136 **Figure 2—figure supplement 2: Distinct astrocyte-specific expression domains of AiE2120m and**
1137 **AiE2160m.**
1138
We injected mice with the indicated astrocyte-specific SYFP2-expressing enhancer-AAVs and
performed whole-brain serial two-photon tomography using the TissueCyte platform[88]. These vectors
display largely non-overlapping zones of astrocyte expression: AiE2120m is expressed in astrocytes
within multiple forebrain structures including CTX, STR, OB, LSX, HPF, and TH, as well as MB,
whereas AiE2160m is expressed in MB, CBX, and HB structures as well as complementary forebrain
structures including HY, MSC, and GPe, and OB. In the OB AiE2120m is expressed in astrocytes within
the granule cell layer, internal plexiform layer, and periglomerular cell layer, whereas AiE2160m is
expressed in a complementary pattern of astrocytes within the external plexiform layer. Each image
series shows one animal representative of two to three animals tested.
1148
1149 Abbreviations: CTX cerebral cortex, STR striatum, OB olfactory bulb, LSX lateral septal complex, HPF
1150 hippocampal formation, TH thalamus, MB midbrain, CBX cerebellar cortex, HB hindbrain, HY
1151 hypothalamus, MSC medial septal complex, GPe globus pallidus, external layer.
1152

Strong oligodendrocytes

AiE0395h · AiE0395m · AiE0396h · AiE0397m · AiE0401h
AiE0409m · AiE0410m · AiE0410m_3xC · AiE0641m · AiE0641m_3xC

Moderate

AiE0361h · AiE0391h · AiE0398h · AiE0400m · AiE0403h
AiE0407h · AiE0409h · AiE0410h · MBP_promoter

Weak

AiE0396m · AiE0397h · AiE0400h · AiE0402m · AiE0405m

Mixed spec.

AiE0364h · AiE0403m · AiE0404h

No oligodendrocyte expression

AiE0391m · AiE0392h · AiE0393h · AiE0393m · AiE0394h
AiE0394m · AiE0398m · AiE0399h · AiE0399m · AiE0401m
AiE0402h · AiE0404m · AiE0405h · AiE0406h · AiE0406m
AiE0407m · AiE0408h · AiE0408m · AiE0733m

1 mm

100 μm

 Figure 3—figure supplement 1: Full screening results of all candidate enhancer-AAVs targeting
oligodendrocytes.

We injected mice with the indicated enhancer-AAV vectors between P42 and P56, then after 3-4 weeks
we harvested brains, sliced them on a sliding microtome with freezing stage at 30 μm thickness, co-
stained the sections with DAPI, and mounted them with Vectashield Vybrance. Oligodendrocyte-
specific enhancer-AAV vectors are broadly grouped by expression pattern into the following categories:
"Strong oligodendrocytes", "Weak" meaning many oligodendrocytes throughout the brain are labeled at
low level, "Mixed specificities" meaning several off-target neuron or astrocyte populations are also
present in addition to oligodendrocytes, and "No oligodendrocyte expression" meaning failure to detect
any clear oligodendrocytes in these whole-brain sagittal images. These epifluorescent screening
images represent n = 1 to 20 animals tested per enhancer (median 2 animals), and were sometimes
taken on multiple different microscopes (see **Table S2** for full summary of screened animals).

1166   **Figure 5—figure supplement 1: Sorting enhancer-AAV-labeled astrocytes and oligodendrocytes.**
1167
1168   Example gating strategies for sorting AiE0390h_3xC2-labeled astrocytes and AiE0396h-labeled
1169   oligodendrocytes from mouse VISp. Each flow gating strategy represents one animal tested for that
1170   vector (n = 26 total animals for astrocytes and n = 21 for oligodendrocytes).

1171

1172

AiE0390m in NHP somatosensory cortex

A SYFP2
250 µm
L1
Pia
L2
B
L3
L4
L5
L6
White matter
C

B 50 µm
Gray matter protoplasmic astrocytes

C 50 µm
White matter fibrous astrocyte

D 25 µm
White matter fibrous astrocyte **SYFP2**

FGFR3

1173 **Figure 11—figure supplement 1: Diverse morphologies of macaque astrocytes labeled by**
1174 **enhancer-AAVs.**

1175

(**A-C**) Labeling of both gray matter protoplasmic astrocytes and white matter fibrous astrocytes by
AiE0390m enhancer-AAV. We show full cortical column of a somatosensory cortex injection site in A,
with expanded insets to show protoplasmic astrocytes in gray matter (**B**) and fibrous astrocytes in white
matter (**C**). Images represent one animal tested.

(**D**) Confirmation of astrocyte identity by mFISH. Fibrous astrocytes in white matter express the
astrocyte marker *FGFR3*, similar to gray matter protoplasmic astrocytes (**Figure 8M**).

1182

1183   **Table S1.**

Enhancer name	Identification scheme	Genome	Chr	Start	Stop	Enhancer_group	PhyloP	Length	Specificity	Strength (CPM)	Screen result	Expression pattern	Included in GLM model
eHGT_361h	Marker_gene	hg38	chrX	103780509	103781455	Human_Oligo	1.823	855	0.635	11.72	Yes	Oligodendrocytes_moderate	No
eHGT_361m	Marker_gene	mm10	chrX	136826512	136827460	Mouse_Oligo	1.488	807	0.879	12.03	No_data	No_data	No
eHGT_364h	Marker_gene	hg38	chr21	33051808	33052323	Human_Oligo	2.5	458	0.614	2.29	Mixed_spe	Mixed_specificity_oligodendrocy	No
eHGT_364m	Marker_gene	mm10	chr16	91246204	91246699	Mouse_Oligo	1.99	444	0.945	9.54	No_data	No_data	No
eHGT_371h	High_specificity	hg38	chr3	13485182	13485473	Human_Astro	0.967	269	0.454	3.81	Yes	Scattered_astrocytes	Yes
eHGT_371m	High_specificity	mm10	chr6	91160424	91160736	Mouse_Astro	0.87	241	0.216	4.28	No	No_astrocyte_expression	Yes
eHGT_372h	High_specificity	hg38	chr7	121914818	121915080	Human_Astro	0.33	216	0.668	2.6	No	No_astrocyte_expression	Yes
eHGT_372m	High_specificity	mm10	chr6	22910076	22910291	Mouse_Astro	0.194	187	0.742	2.85	No	No_astrocyte_expression	Yes
eHGT_373h	High_specificity	hg38	chr5	97134713	97135080	Human_Astro	0.537	329	0.72	3.81	No	No_astrocyte_expression	Yes
eHGT_373m	High_specificity	mm10	chr17	17409626	17409965	Mouse_Astro	0.531	319	0.909	11.78	Yes	Most_astrocytes_weak	Yes
eHGT_374h	High_specificity	hg38	chr15	64845370	64845672	Human_Astro	1.618	266	0.424	2.86	Yes	Scattered_astrocytes	Yes
eHGT_374m	High_specificity	mm10	chr9	65576810	65577101	Mouse_Astro	1.091	259	0.542	5.35	Yes	Scattered_astrocytes	Yes
eHGT_375h	High_specificity	hg38	chr13	99961720	99962046	Human_Astro	4.977	268	0.659	2.51	No	No_astrocyte_expression	Yes
eHGT_375m	High_specificity	mm10	chr14	122455334	122455655	Mouse_Astro	3.82	268	0.867	5.71	Yes	Bergmann_glia_astrocytes	Yes
eHGT_376h	High_specificity	hg38	chr3	188076136	188076690	Human_Astro	1.341	457	0.646	4.24	Yes	Most_astrocytes_strong	Yes
eHGT_376m	High_specificity	mm10	chr16	24326000	24326508	Mouse_Astro	1.083	435	0.865	11.78	Yes	Most_astrocytes_strong	Yes
eHGT_377h	High_specificity	hg38	chr10	80406003	80406485	Human_Astro	0.687	453	0.782	4.41	No	No_astrocyte_expression	Yes
eHGT_377m	High_specificity	mm10	chr14	41015471	41016233	Mouse_Astro	0.541	747	0.656	7.85	Yes	Most_astrocytes_strong	Yes
eHGT_378h	High_specificity	hg38	chr11	120182373	120182964	Human_Astro	2.312	559	0.681	6.32	No	No_astrocyte_expression	No
eHGT_378m	High_specificity	mm10	chr9	43268975	43269581	Mouse_Astro	1.699	559	0.911	12.49	Weak	Most_astrocytes_weak	No
eHGT_379h	High_specificity	hg38	chr3	64837170	64837760	Human_Astro	0.151	496	0.659	4.93	No	No_astrocyte_expression	Yes
eHGT_379m	High_specificity	mm10	chr6	93093752	93094309	Mouse_Astro	0.216	460	0.939	8.21	No	No_astrocyte_expression	Yes
eHGT_380h	High_specificity	hg38	chr6	44045566	44046145	Human_Astro	0.591	527	0.633	10.82	Yes	Most_astrocytes_strong	Yes
eHGT_380m	High_specificity	mm10	chr17	45765782	45766284	Mouse_Astro	0.536	479	0.727	7.14	No	No_astrocyte_expression	Yes
eHGT_381h	High_specificity	hg38	chr7	131400918	131401149	Human_Astro	2.166	173	0.4	1.9	Weak	Mid/hindbrain_astrocytes	No
eHGT_381m	High_specificity	mm10	chr6	31434638	31434879	Mouse_Astro	1.8	172	0.927	5.35	No	No_astrocyte_expression	No
eHGT_382h	High_specificity	hg38	chr10	113047655	113048169	Human_Astro	0.544	435	0.685	3.98	No	No_astrocyte_expression	Yes
eHGT_382m	High_specificity	mm10	chr19	55825471	55825949	Mouse_Astro	0.439	424	0.694	5.71	No	No_astrocyte_expression	Yes
eHGT_383h	High_specificity	hg38	chr2	220959419	220959789	Human_Astro	1.026	312	0.858	3.38	No	No_astrocyte_expression	Yes
eHGT_383m	High_specificity	mm10	chr1	76942759	76943103	Mouse_Astro	0.81	315	0.902	11.42	No	No_astrocyte_expression	Yes
eHGT_384h	High_specificity	hg38	chr6	121897357	121897797	Human_Astro	1.695	351	0.342	2.77	No	No_astrocyte_expression	Yes
eHGT_384m	High_specificity	mm10	chr10	56893293	56893720	Mouse_Astro	1.576	366	0.846	9.64	No	No_astrocyte_expression	Yes
eHGT_385h	High_specificity	hg38	chr9	123741938	123742543	Human_Astro	6.233	563	0.655	7.88	Weak	Telencephalon_astrocytes	No
eHGT_385m	High_specificity	mm10	chr2	38111566	38112219	Mouse_Astro	4.798	572	0.714	14.27	Yes	Telencephalon_astrocytes	No
eHGT_386h	High_specificity	hg38	chr1	15736012	15736372	Human_Astro	0.595	287	0.417	2.51	Weak	Most_astrocytes_weak	No
eHGT_386m	High_specificity	mm10	chr4	141623748	141624075	Mouse_Astro	1.042	231	0.527	5	Yes	Most_astrocytes_weak	No
eHGT_387h	High_specificity	hg38	chr1	219881964	219882484	Human_Astro	0.601	467	0.729	6.06	No_data	No_data	No
eHGT_387m	High_specificity	mm10	chr1	185493675	185494167	Mouse_Astro	0.633	425	0.959	15.35	Yes	Most_astrocytes_strong	No
eHGT_388h	High_specificity	hg38	chr12	20657459	20657845	Human_Astro	2.912	369	0.754	3.98	No	No_astrocyte_expression	Yes
eHGT_388m	High_specificity	mm10	chr6	141495577	141496010	Mouse_Astro	2.283	362	0.879	16.77	No	No_astrocyte_expression	Yes
eHGT_389h	High_specificity	hg38	chr20	63252174	63252906	Human_Astro	0.003	694	0.648	9.52	Yes	Scattered_astrocytes	No
eHGT_389m	High_specificity	mm10	chr2	180952658	180953361	Mouse_Astro	0.351	667	0.695	8.56	Weak	Most_astrocytes_weak	No
eHGT_390h	High_specificity	hg38	chr7	42152720	42153410	Human_Astro	7.463	634	0.699	8.91	Yes	Most_astrocytes_strong	Yes
eHGT_390m	High_specificity	mm10	chr13	15543638	15544333	Mouse_Astro	5.544	639	0.869	16.42	Yes	Most_astrocytes_strong	Yes
eHGT_391h	High_specificity	hg38	chr11	729607	730042	Human_Oligo	-0.132	419	0.688	7.49	Weak	Oligodendrocytes_moderate	No
eHGT_391m	High_specificity	mm10	chr7	141365022	141365541	Mouse_Oligo	0.135	474	0.901	13.2	No	No_oligodendrocyte_expression	No
eHGT_392h	High_specificity	hg38	chr4	48787039	48787541	Human_Oligo	1.369	431	0.776	6.88	No	No_oligodendrocyte_expression	No
eHGT_392m	High_specificity	mm10	chr5	73264044	73264568	Mouse_Oligo	1.225	448	0.832	4.25	No_data	No_data	No
eHGT_393h	High_specificity	hg38	chr18	66603993	66604511	Human_Oligo	2.036	451	0.755	7.23	No	No_oligodendrocyte_expression	Yes
eHGT_393m	High_specificity	mm10	chr1	110977271	110977865	Mouse_Oligo	1.391	548	0.593	6.16	No	No_oligodendrocyte_expression	Yes
eHGT_394h	High_specificity	hg38	chr3	171776470	171776980	Human_Oligo	0.518	444	0.782	5.2	No	No_oligodendrocyte_expression	Yes
eHGT_394m	High_specificity	mm10	chr3	27977667	27978413	Mouse_Oligo	0.254	661	0.738	12.62	No	No_oligodendrocyte_expression	Yes
eHGT_395h	High_specificity	hg38	chr9	128374105	128374728	Human_Oligo	2.782	595	0.742	9.96	Yes	Oligodendrocytes_strong	Yes
eHGT_395m	High_specificity	mm10	chr2	29830874	29831537	Mouse_Oligo	2.002	632	0.881	24.5	Yes	Oligodendrocytes_strong	Yes
eHGT_396h	High_specificity	hg38	chr11	117295457	117295798	Human_Oligo	1.384	302	0.561	3.88	Yes	Oligodendrocytes_strong	No

1184

eHGT_396m	High_specificity	mm10	chr9	45854220	45854539	Mouse_Oligo	1.185	288	0.824	7.33	Weak	Oligodendrocytes_weak	No
eHGT_397h	High_specificity	hg38	chr15	44188767	44189188	Human_Oligo	1.986	396	0.591	4.67	Weak	Oligodendrocytes_weak	No
eHGT_397m	High_specificity	mm10	chr2	121799010	121799455	Mouse_Oligo	1.661	385	0.759	6.16	Yes	Oligodendrocytes_strong	No
eHGT_398h	High_specificity	hg38	chr16	57270420	57270919	Human_Oligo	1.042	475	0.793	7.49	Yes	Oligodendrocytes_moderate	Yes
eHGT_398m	High_specificity	mm10	chr8	94684265	94684766	Mouse_Oligo	1.199	446	0.806	11.74	No	No_oligodendrocyte_expression	Yes
eHGT_399h	High_specificity	hg38	chr2	36902653	36902934	Human_Oligo	4.206	253	0.628	2.73	No	No_oligodendrocyte_expression	Yes
eHGT_399m	High_specificity	mm10	chr17	78684296	78684576	Mouse_Oligo	3.818	249	0.885	4.11	No	No_oligodendrocyte_expression	Yes
eHGT_400h	High_specificity	hg38	chr4	114636222	114636638	Human_Oligo	2.948	372	0.605	3.88	Weak	Oligodendrocytes_weak	No
eHGT_400m	High_specificity	mm10	chr3	125898885	125899302	Mouse_Oligo	2.414	373	0.819	7.63	Yes	Oligodendrocytes_moderate	No
eHGT_401h	High_specificity	hg38	chr1	44592226	44592529	Human_Oligo	1.081	256	0.55	3.17	Yes	Oligodendrocytes_strong	Yes
eHGT_401m	High_specificity	mm10	chr4	117326998	117327312	Mouse_Oligo	1.024	271	0.778	3.96	No	No_oligodendrocyte_expression	Yes
eHGT_402h	High_specificity	hg38	chr17	75697915	75698299	Human_Oligo	0.976	320	0.626	4.67	No	No_oligodendrocyte_expression	No
eHGT_402m	High_specificity	mm10	chr11	115956504	115956883	Mouse_Oligo	0.956	326	0.919	13.94	Weak	Oligodendrocytes_weak	No
eHGT_403h	High_specificity	hg38	chr11	67410338	67410731	Human_Oligo	0.524	382	0.588	4.5	Yes	Oligodendrocytes_moderate	No
eHGT_403m	High_specificity	mm10	chr19	4183762	4184154	Mouse_Oligo	1.019	346	0.586	8.51	Mixed_spe	Mixed_specificity_oligodendrocy	No
eHGT_404h	High_specificity	hg38	chr2	163943542	163943945	Human_Oligo	1.738	370	0.721	5.2	Mixed_spe	Mixed_specificity_oligodendrocy	No
eHGT_404m	High_specificity	mm10	chr2	64325556	64326248	Mouse_Oligo	1.562	599	0.933	12.18	No	No_oligodendrocyte_expression	No
eHGT_405h	High_specificity	hg38	chr3	185175495	185175886	Human_Oligo	0.607	362	0.49	4.05	No	No_oligodendrocyte_expression	No
eHGT_405m	High_specificity	mm10	chr16	21724337	21724697	Mouse_Oligo	0.444	335	0.91	8.36	Weak	Oligodendrocytes_weak	No
eHGT_406h	High_specificity	hg38	chr2	88735239	88735593	Human_Oligo	-0.058	304	0.518	3.26	No	No_oligodendrocyte_expression	Yes
eHGT_406m	High_specificity	mm10	chr6	70774375	70774694	Mouse_Oligo	0.091	256	0.696	5.87	No	No_oligodendrocyte_expression	Yes
eHGT_407h	High_specificity	hg38	chr6	46098744	46099007	Human_Oligo	0.581	253	0.614	2.47	Yes	Oligodendrocytes_moderate	Yes
eHGT_407m	High_specificity	mm10	chr17	44121834	44122162	Mouse_Oligo	0.545	271	0.847	7.48	No	No_oligodendrocyte_expression	Yes
eHGT_408h	High_specificity	hg38	chr8	60867028	60867631	Human_Oligo	3.002	574	0.53	6.61	No	No_oligodendrocyte_expression	Yes
eHGT_408m	High_specificity	mm10	chr4	8867749	8868420	Mouse_Oligo	2.361	596	0.769	11.88	No	No_oligodendrocyte_expression	Yes
eHGT_409h	High_specificity	hg38	chr14	67539037	67539541	Human_Oligo	2.042	452	0.564	4.5	Yes	Oligodendrocytes_moderate	Yes
eHGT_409m	High_specificity	mm10	chr12	79035271	79035759	Mouse_Oligo	1.603	437	0.908	13.5	Yes	Oligodendrocytes_strong	Yes
eHGT_410h	High_specificity	hg38	chr9	81717541	81717791	Human_Oligo	2.177	202	0.579	1.41	Yes	Oligodendrocytes_moderate	Yes
eHGT_410m	High_specificity	mm10	chr4	72233773	72234067	Mouse_Oligo	1.838	203	0.817	9.68	Yes	Oligodendrocytes_strong	Yes
eHGT_495h	Marker_gene	hg38	chr19	33217923	33218346	Human_Astro	1.481	364	0.302	0.95	No_data	No_data	No
eHGT_495m	Marker_gene	mm10	chr7	35192340	35192802	Mouse_Astro	0.977	396	0.818	7.85	No	No_astrocyte_expression	No
eHGT_496h	Marker_gene	hg38	chr19	33225554	33226354	Human_Astro	1.18	749	0.516	4.76	No_data	No_data	No
eHGT_496m	Marker_gene	mm10	chr7	35185913	35186648	Mouse_Astro	0.955	686	0.734	15.35	No_data	No_data	No
eHGT_497h	Marker_gene	hg38	chr3	55237236	55237718	Human_Astro	1.648	437	0.037	0.09	No_data	No_data	No
eHGT_497m	Marker_gene	mm10	chr14	28781601	28782057	Mouse_Astro	1.444	410	0.966	7.49	No	No_astrocyte_expression	No
eHGT_641h	Marker_gene	hg38	chr21	33116993	33117641	Human_Oligo	4.073	557	0.755	16.66	No_data	No_data	No
eHGT_641m	Marker_gene	mm10	chr16	91305334	91305956	Mouse_Oligo	3.233	529	0.92	23.32	Yes	Oligodendrocytes_strong	No
eHGT_642h	Marker_gene	hg38	chr17	44539855	44540164	Human_Astro	2.026	230	0.463	0.26	No_data	No_data	No
eHGT_642m	Marker_gene	mm10	chr11	102579463	102579768	Mouse_Astro	1.655	228	0.86	11.06	Mixed_spe	Non-specific_astrocytes	No
eHGT_733m	Marker_gene	mm10	chr2	29273794	29274249	Mouse_Oligo	0.127	366	0.876	5.72	No	No_oligodendrocyte_expression	No
MGT_E117h	High_strength	hg38	chr7	23230417	23231661	Human_Astro	1.333	1213	0.197	4.15	No_data	No_data	No
MGT_E117m	High_strength	mm10	chr6	49020855	49022070	Mouse_Astro	1.202	1174	0.852	49.25	Mixed_spe	Non-specific_astrocytes	No
MGT_E118h	High_strength	hg38	chr13	25910516	25911691	Human_Astro	3.328	1146	0.062	0.26	No_data	No_data	No
MGT_E118m	High_strength	mm10	chr14	59736768	59737951	Mouse_Astro	2.384	1143	0.889	47.11	Mixed_spe	Mixed_specificity_astrocytes	No
MGT_E119h	High_strength	hg38	chr18	59386813	59388237	Human_Astro	4.406	1421	0.532	6.58	No_data	No_data	No
MGT_E119m	High_strength	mm10	chr18	66022617	66024006	Mouse_Astro	3.435	1386	0.926	38.9	Mixed_spe	Mixed_specificity_astrocytes	No
MGT_E120h	High_strength	hg38	chr2	134072857	134074292	Human_Astro	0.875	1436	0.333	2.16	No_data	No_data	No
MGT_E120m	High_strength	mm10	chr1	127159651	127161393	Mouse_Astro	0.697	1661	0.896	37.83	Yes	Forebrain_midbrain_astrocytes	No
MGT_E121h	High_strength	hg38	chr17	68641683	68642415	Human_Astro	0.531	733	0.139	0.17	No_data	No_data	No
MGT_E121m	High_strength	mm10	chr11	109753663	109754656	Mouse_Astro	0.322	994	0.911	38.9	Mixed_spe	Mixed_specificity_astrocytes	No
MGT_E122h	High_strength	hg38	chr4	175988170	175989315	Human_Astro	1.134	1121	0.675	8.65	No_data	No_data	No
MGT_E122m	High_strength	mm10	chr8	54791804	54792949	Mouse_Astro	1.078	1117	0.922	37.47	Yes	Forebrain_cerebellum_astrocyte	No
MGT_E123h	High_strength	hg38	chr14	80206581	80208078	Human_Astro	2.747	1497	0.701	12.12	No_data	No_data	No
MGT_E123m	High_strength	mm10	chr12	90733627	90735117	Mouse_Astro	2.427	1490	0.808	36.76	Mixed_spe	Mixed_specificity_astrocytes	No
MGT_E160h	None	hg38	chr2	48184996	48185538	Human_Astro	0.314	538	0	0	No_data	No_data	No
MGT_E160m	None	mm10	chr17	88297369	88297914	Mouse_Astro	0.347	516	0.282	1.07	Yes	Mid/hindbrain_astrocytes	No

1185

1186   **Table S1: Genomic coordinates, sequence characterization, and mouse screening results of all**
1187   **tested astrocyte and oligodendrocyte enhancers.**

1188   Calculations of parameters are as described in Methods section.

1189

Table S2.

Mouse ID	Genotype	Enhancer ID	Vector ID	Vector Full Name	Data Modality	Targeted Cell Population	Fluor	Delivery Method	Publication
642124	C57BL/6J	AiE0361h	AiP12979	pAAV-AiE0361h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
567931	C57BL/6J	AiE0361h	AiP12979	pAAV-AiE0361h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
637260	C57BL/6J	AiE0361h	AiP12979	pAAV-AiE0361h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
621157	C57BL/6J	AiE0361h	AiP12979	pAAV-AiE0361h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
567932	C57BL/6J	AiE0364h	AiP12980	pAAV-AiE0364h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621158	C57BL/6J	AiE0364h	AiP12980	pAAV-AiE0364h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
636670	C57BL/6J	AiE0371h	AiP12141	pAAV-AiE0371h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
725447	C57BL/6J	AiE0371h	AiP12141	pAAV-AiE0371h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
743804	C57BL/6J	AiE0371h	AiP12141	pAAV-AiE0371h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
567934	C57BL/6J	AiE0371h	AiP12141	pAAV-AiE0371h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
615690	C57BL/6J	AiE0371h	AiP12141	pAAV-AiE0371h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
623007	C57BL/6J	AiE0371h	AiP12141	pAAV-AiE0371h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671755	CD-1_IGS	AiE0371h	AiP12141	pAAV-AiE0371h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
499663	C57BL/6J	AiE0371m	AiP12082	pAAV-AiE0371m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
636669	C57BL/6J	AiE0371m	AiP12082	pAAV-AiE0371m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
725446	C57BL/6J	AiE0371m	AiP12082	pAAV-AiE0371m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
743802	C57BL/6J	AiE0371m	AiP12082	pAAV-AiE0371m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
513373	C57BL/6J	AiE0372h	AiP12142	pAAV-AiE0372h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
499664	C57BL/6J	AiE0372m	AiP12083	pAAV-AiE0372m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
722734	C57BL/6J	AiE0372m	AiP12083	pAAV-AiE0372m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
616877	C57BL/6J	AiE0372m	AiP12083	pAAV-AiE0372m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
551395	C57BL/6J	AiE0373h	AiP12504	pAAV-AiE0373h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
499665	C57BL/6J	AiE0373m	AiP12084	pAAV-AiE0373m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
616737	C57BL/6J	AiE0373m	AiP12084	pAAV-AiE0373m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623015	C57BL/6J	AiE0373m	AiP12084	pAAV-AiE0373m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
679835	C57BL/6J	AiE0373m	AiP12084	pAAV-AiE0373m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
550273	C57BL/6J	AiE0374h	AiP12493	pAAV-AiE0374h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
550277	C57BL/6J	AiE0374h	AiP12493	pAAV-AiE0374h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623008	C57BL/6J	AiE0374h	AiP12493	pAAV-AiE0374h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
647376	C57BL/6J	AiE0374h	AiP12493	pAAV-AiE0374h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615013	C57BL/6J	AiE0374m	AiP12098	pAAV-AiE0374m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
625965	C57BL/6J	AiE0374m	AiP12098	pAAV-AiE0374m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
636674	C57BL/6J	AiE0374m	AiP12098	pAAV-AiE0374m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
614201	C57BL/6J	AiE0374m	AiP12098	pAAV-AiE0374m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
630481	C57BL/6J	AiE0374m	AiP12098	pAAV-AiE0374m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
567935	C57BL/6J	AiE0374m	AiP12098	pAAV-AiE0374m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623016	C57BL/6J	AiE0374m	AiP12098	pAAV-AiE0374m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615691	C57BL/6J	AiE0374m	AiP12098	pAAV-AiE0374m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671754	CD-1_IGS	AiE0374m	AiP12098	pAAV-AiE0374m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
637359	Scn1a-R613X	AiE0374m	AiP14081	pAAV-AiE0374m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B*	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
637362	Scn1a-R613X	AiE0374m	AiP14081	pAAV-AiE0374m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B*	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
637358	Scn1a +/+	AiE0374m	AiP14081	pAAV-AiE0374m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B*	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
637356	Scn1a +/+	AiE0374m	AiP14081	pAAV-AiE0374m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B*	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
513374	C57BL/6J	AiE0375h	AiP12143	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
499666	C57BL/6J	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
615009	C57BL/6J	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
621237	C57BL/6J	AiE0375m	AiP13073	pAAV-AiE0375m-minBG-H2B-nsSYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
625958	C57BL/6J	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
627902	C57BL/6J	AiE0375m	AiP13073	pAAV-AiE0375m-minBG-H2B-nsSYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
637136	C57BL/6J	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
745309	C57BL/6J	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
745315	C57BL/6J	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
614198	C57BL/6J	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
651221	C57BL/6J	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
616878	C57BL/6J	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
623017	C57BL/6J	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671752	CD-1_IGS	AiE0375m	AiP12085	pAAV-AiE0375m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
615010	C57BL/6J	AiE0376h	AiP12144	pAAV-AiE0376h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
621239	C57BL/6J	AiE0376h	AiP13074	pAAV-AiE0376h-minBG-H2B-nsSYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
625960	C57BL/6J	AiE0376h	AiP12144	pAAV-AiE0376h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
627903	C57BL/6J	AiE0376h	AiP13074	pAAV-AiE0376h-minBG-H2B-nsSYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
614199	C57BL/6J	AiE0376h	AiP12144	pAAV-AiE0376h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
514392	C57BL/6J	AiE0376h	AiP12144	pAAV-AiE0376h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623009	C57BL/6J	AiE0376h	AiP12144	pAAV-AiE0376h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671756	CD-1_IGS	AiE0376h	AiP12144	pAAV-AiE0376h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
567936	C57BL/6J	AiE0376m	AiP12120	pAAV-AiE0376m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
615688	C57BL/6J	AiE0376m	AiP12120	pAAV-AiE0376m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
623018	C57BL/6J	AiE0376m	AiP12120	pAAV-AiE0376m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
514393	C57BL/6J	AiE0377h	AiP12145	pAAV-AiE0377h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
576009	C57BL/6J	AiE0377m	AiP13060	pAAV-AiE0377m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623019	C57BL/6J	AiE0377m	AiP13060	pAAV-AiE0377m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615686	C57BL/6J	AiE0377m	AiP13060	pAAV-AiE0377m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
554425	C57BL/6J	AiE0378h	AiP12676	pAAV-AiE0378h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
560757	C57BL/6J	AiE0378m	AiP12827	pAAV-AiE0378m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
740048	C57BL/6J	AiE0378m	AiP12827	pAAV-AiE0378m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
740048	C57BL/6J	AiE0378m	AiP12827	pAAV-AiE0378m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
740048	C57BL/6J	AiE0378m	AiP12827	pAAV-AiE0378m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
740048	C57BL/6J	AiE0378m	AiP12827	pAAV-AiE0378m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
550274	C57BL/6J	AiE0379h	AiP12494	pAAV-AiE0379h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
550278	C57BL/6J	AiE0379h	AiP12494	pAAV-AiE0379h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
499667	C57BL/6J	AiE0379m	AiP12086	pAAV-AiE0379m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
615012	C57BL/6J	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
625962	C57BL/6J	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
644030	C57BL/6J	AiE0380h	AiP14080	pAAV-AiE0380h-minBG-SYFP2-P2A-3XFLAG-10aa-H2B*	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
743809	C57BL/6J	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
745313	C57BL/6J	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
614200	C57BL/6J	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
651225	C57BL/6J	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
572497	C57BL/6J	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv

623010	C57BL/6J	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
616741	C57BL/6J	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
634487	C57BL/6J	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671753	CD-1_IGS	AiE0380h	AiP13098	pAAV-AiE0380h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
567937	C57BL/6J	AiE0380m	AiP12121	pAAV-AiE0380m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
615014	C57BL/6J	AiE0381h	AiP12146	pAAV-AiE0381h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
636671	C57BL/6J	AiE0381h	AiP12146	pAAV-AiE0381h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
725448	C57BL/6J	AiE0381h	AiP12146	pAAV-AiE0381h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
743805	C57BL/6J	AiE0381h	AiP12146	pAAV-AiE0381h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
615516	C57BL/6J	AiE0381h	AiP12146	pAAV-AiE0381h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
651222	C57BL/6J	AiE0381h	AiP12146	pAAV-AiE0381h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
567938	C57BL/6J	AiE0381h	AiP12146	pAAV-AiE0381h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
615689	C57BL/6J	AiE0381h	AiP12146	pAAV-AiE0381h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
632617	C57BL/6J	AiE0381h	AiP12146	pAAV-AiE0381h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
641421	C57BL/6J	AiE0381h	AiP12146	pAAV-AiE0381h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
501914	C57BL/6J	AiE0381m	AiP12100	pAAV-AiE0381m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
514394	C57BL/6J	AiE0382h	AiP12147	pAAV-AiE0382h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
501915	C57BL/6J	AiE0382m	AiP12101	pAAV-AiE0382m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
514395	C57BL/6J	AiE0383h	AiP12148	pAAV-AiE0383h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
501912	C57BL/6J	AiE0383m	AiP12097	pAAV-AiE0383m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
567939	C57BL/6J	AiE0384h	AiP12149	pAAV-AiE0384h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
500565	C57BL/6J	AiE0384m	AiP12087	pAAV-AiE0384m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
550275	C57BL/6J	AiE0385h	AiP12495	pAAV-AiE0385h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623011	C57BL/6J	AiE0385h	AiP12495	pAAV-AiE0385h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
647375	C57BL/6J	AiE0385h	AiP12495	pAAV-AiE0385h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615015	C57BL/6J	AiE0385m	AiP12122	pAAV-AiE0385m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
635183	C57BL/6J	AiE0385m	AiP14074	pAAV-AiE0385m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B*	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
637137	C57BL/6J	AiE0385m	AiP12122	pAAV-AiE0385m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
615515	C57BL/6J	AiE0385m	AiP12122	pAAV-AiE0385m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
651224	C57BL/6J	AiE0385m	AiP12122	pAAV-AiE0385m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
567940	C57BL/6J	AiE0385m	AiP12122	pAAV-AiE0385m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623021	C57BL/6J	AiE0385m	AiP12122	pAAV-AiE0385m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615687	C57BL/6J	AiE0385m	AiP12122	pAAV-AiE0385m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
551390	C57BL/6J	AiE0386h	AiP12496	pAAV-AiE0386h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623012	C57BL/6J	AiE0386h	AiP12496	pAAV-AiE0386h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
636672	C57BL/6J	AiE0386m	AiP12088	pAAV-AiE0386m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
725449	C57BL/6J	AiE0386m	AiP12088	pAAV-AiE0386m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
743803	C57BL/6J	AiE0386m	AiP12088	pAAV-AiE0386m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
616738	C57BL/6J	AiE0386m	AiP12088	pAAV-AiE0386m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
500566	C57BL/6J	AiE0386m	AiP12088	pAAV-AiE0386m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623022	C57BL/6J	AiE0386m	AiP12088	pAAV-AiE0386m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
679856	C57BL/6J	AiE0387h	AiP12150	pAAV-AiE0387h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
660181	C57BL/6J	AiE0387h	AiP12150	pAAV-AiE0387h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
749615	Ai14(RCL-tdT	AiE0387m	AiP13999	pAAV-AiE0387m-minBG-iCre(R297T)-BGHpA	EPI, IHC	Astro	iCre(R297	RO	Ben-Simon, et al. 2025 Cell
751717	Ai14(RCL-tdT	AiE0387m	AiP13999	pAAV-AiE0387m-minBG-iCre(R297T)-BGHpA	IHC	Astro	iCre(R297	RO	Mich et al., 2025 bioRxiv
751911	Ai14(RCL-tdT	AiE0387m	AiP13999	pAAV-AiE0387m-minBG-iCre(R297T)-BGHpA	IHC	Astro	iCre(R297	RO	Mich et al., 2025 bioRxiv
542087	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
608407	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
621234	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
625959	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
636673	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
648493	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
648494	C57BL/6J	AiE0387m	AiP14309	pAAV-TelN-LITR-AiE0387m-minBG-SYFP2-P2A-3XFLAG-:	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
648495	C57BL/6J	AiE0387m	AiP14309	pAAV-TelN-LITR-AiE0387m-minBG-SYFP2-P2A-3XFLAG-:	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
743807	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
745311	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
593373	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
502935	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
630972	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
629424	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
634482	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
639178	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
629424	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
634482	C57BL/6J	AiE0387m	AiP12102	pAAV-AiE0387m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
610229	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
610229	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
652697	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
652697	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
631110	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	EPI	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
631110	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	EPI	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
631110	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	EPI	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
631110	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
652697	C57BL/6J	AiE0387m	AiP13635	pAAV-AiE0387m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
634491	C57BL/6J	AiE0387m_3xC	AiP13568	pAAV-AiE0387m_3xC1-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
630973	C57BL/6J	AiE0387m_3xC	AiP13568	pAAV-AiE0387m_3xC1-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
634492	C57BL/6J	AiE0387m_3xC	AiP13569	pAAV-AiE0387m_3xC2-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
514396	C57BL/6J	AiE0388h	AiP12151	pAAV-AiE0388h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
502936	C57BL/6J	AiE0388m	AiP12103	pAAV-AiE0388m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
567941	C57BL/6J	AiE0389h	AiP12152	pAAV-AiE0389h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623013	C57BL/6J	AiE0389h	AiP12152	pAAV-AiE0389h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
576010	C57BL/6J	AiE0389m	AiP13061	pAAV-AiE0389m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623023	C57BL/6J	AiE0389m	AiP13061	pAAV-AiE0389m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
717604	Ai14(RCL-tdT	AiE0390h	AiP13998	pAAV-AiE0390h-minBG-iCre(R297T)-BGHpA	EPI	Astro	iCre(R297	RO	Ben-Simon, et al. 2025 Cell
750307	Ai14(RCL-tdT	AiE0390h	AiP13998	pAAV-AiE0390h-minBG-iCre(R297T)-BGHpA	STPT, IHC	Astro	iCre(R297	RO	Ben-Simon, et al. 2025 Cell
640524	Ai14(RCL-tdT	AiE0390h	AiP13998	pAAV-AiE0390h-minBG-iCre(R297T)-BGHpA	IHC	Astro	iCre(R297	RO	Mich et al., 2025 bioRxiv
738664	Ai14(RCL-tdT	AiE0390h	AiP13998	pAAV-AiE0390h-minBG-iCre(R297T)-BGHpA	IHC	Astro	iCre(R297	RO	Mich et al., 2025 bioRxiv
615011	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
621240	C57BL/6J	AiE0390h	AiP12822	pAAV-AiE0390h-minRho-SYFP2-10aa-H2Bmod-WPRE3-(	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
625961	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv

743808	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
745312	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
584420	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
584420	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
514397	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
639176	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
630974	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
639176	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
623014	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
630974	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
639176	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
630974	C57BL/6J	AiE0390h	AiP12153	pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671751	CD-1_IGS	AiE0390h		pAAV-AiE0390h-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
630975	C57BL/6J	AiE0390h_3xC	AiP13571	pAAV-AiE0390h_3xC1-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
651220	C57BL/6J	AiE0390h_3xC	AiP13572	pAAV-AiE0390h_3xC2-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
647378	C57BL/6J	AiE0390h_3xC	AiP13572	pAAV-AiE0390h_3xC2-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
630976	C57BL/6J	AiE0390h_3xC	AiP13572	pAAV-AiE0390h_3xC2-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
630977	C57BL/6J	AiE0390h_3xC	AiP13573	pAAV-AiE0390h_3xC3-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
630977	C57BL/6J	AiE0390h_3xC	AiP13573	pAAV-AiE0390h_3xC3-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615008	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
621236	C57BL/6J	AiE0390m	AiP12818	pAAV-AiE0390m-minRho-SYFP2-10aa-H2Bmod-WPRE3-	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
625964	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
629468	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
627901	C57BL/6J	AiE0390m	AiP12818	pAAV-AiE0390m-minRho-SYFP2-10aa-H2Bmod-WPRE3-	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
743806	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
745310	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
616740	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
556485	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
500567	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623024	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
678872	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
574637	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
678872	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
574637	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
678872	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
574637	C57BL/6J	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671887	CD-1_IGS	AiE0390m	AiP12089	pAAV-AiE0390m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
636220	Scn1a +/+	AiE0390m	AiP14038	pAAV-390m-minRho-SYFP2-P2A-3XFLAG-10aa-H2B*-W	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
636222	Scn1a-R613X	AiE0390m	AiP14038	pAAV-390m-minRho-SYFP2-P2A-3XFLAG-10aa-H2B*-W	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
637351	Scn1a-R613X	AiE0390m	AiP14038	pAAV-390m-minRho-SYFP2-P2A-3XFLAG-10aa-H2B*-W	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
637352	Scn1a-R613X	AiE0390m	AiP14038	pAAV-390m-minRho-SYFP2-P2A-3XFLAG-10aa-H2B*-W	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
637350	Scn1a +/+	AiE0390m	AiP14038	pAAV-390m-minRho-SYFP2-P2A-3XFLAG-10aa-H2B*-W	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
637354	Scn1a +/+	AiE0390m	AiP14038	pAAV-390m-minRho-SYFP2-P2A-3XFLAG-10aa-H2B*-W	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
751716	Ai14(RCL-tdT	AiE0390m_3xC	AiP14801	pAAV-AiE0390m_3xC2-minBG-iCre(R297T)-BGHpA	IHC	Astro	iCre(R297	RO	Mich et al., 2025 bioRxiv
751909	Ai14(RCL-tdT	AiE0390m_3xC	AiP14801	pAAV-AiE0390m_3xC2-minBG-iCre(R297T)-BGHpA	IHC	Astro	iCre(R297	RO	Mich et al., 2025 bioRxiv
749614	Ai14(RCL-tdT	AiE0390m_3xC	AiP14801	pAAV-AiE0390m_3xC2-minBG-iCre(R297T)-BGHpA	EPI, IHC	Astro	iCre(R297	RO	Ben-Simon, et al. 2025 Cell
626921	C57BL/6J	AiE0390m_3xC	AiP12558	pAAV-AiE0390m_3xC2-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
629462	C57BL/6J	AiE0390m_3xC	AiP12558	pAAV-AiE0390m_3xC2-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
651223	C57BL/6J	AiE0390m_3xC	AiP12558	pAAV-AiE0390m_3xC2-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
630480	C57BL/6J	AiE0390m_3xC	AiP12558	pAAV-AiE0390m_3xC2-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
574636	C57BL/6J	AiE0390m_3xC	AiP12558	pAAV-AiE0390m_3xC2-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
678873	C57BL/6J	AiE0390m_3xC	AiP12558	pAAV-AiE0390m_3xC2-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
639179	C57BL/6J	AiE0390m_3xC	AiP12558	pAAV-AiE0390m_3xC2-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622082	C57BL/6J	AiE0391h	AiP12505	pAAV-AiE0391h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
626924	C57BL/6J	AiE0391h	AiP12505	pAAV-AiE0391h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
617306	C57BL/6J	AiE0391h	AiP12505	pAAV-AiE0391h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
551396	C57BL/6J	AiE0391h	AiP12505	pAAV-AiE0391h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
634957	C57BL/6J	AiE0391h	AiP12505	pAAV-AiE0391h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
620746	C57BL/6J	AiE0391h	AiP12505	pAAV-AiE0391h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
500568	C57BL/6J	AiE0391m	AiP12090	pAAV-AiE0391m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
514398	C57BL/6J	AiE0392h	AiP12154	pAAV-AiE0392h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
514399	C57BL/6J	AiE0393h	AiP12155	pAAV-AiE0393h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
502937	C57BL/6J	AiE0393m	AiP12104	pAAV-AiE0393m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
567942	C57BL/6J	AiE0394h	AiP12156	pAAV-AiE0394h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
567943	C57BL/6J	AiE0394m	AiP12497	pAAV-AiE0394m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
622083	C57BL/6J	AiE0395h	AiP12157	pAAV-AiE0395h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621238	C57BL/6J	AiE0395h	AiP12823	pAAV-AiE0395h-minRho-SYFP2-10aa-H2Bmod-WPRE3-(	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
626925	C57BL/6J	AiE0395h	AiP12157	pAAV-AiE0395h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
629464	C57BL/6J	AiE0395h	AiP12157	pAAV-AiE0395h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
617307	C57BL/6J	AiE0395h	AiP12157	pAAV-AiE0395h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
562430	C57BL/6J	AiE0395h	AiP12157	pAAV-AiE0395h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
641432	C57BL/6J	AiE0395h	AiP12157	pAAV-AiE0395h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
515295	C57BL/6J	AiE0395h	AiP12157	pAAV-AiE0395h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
620747	C57BL/6J	AiE0395h	AiP12157	pAAV-AiE0395h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
634958	C57BL/6J	AiE0395h	AiP12157	pAAV-AiE0395h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
636677	C57BL/6J	AiE0395m	AiP12498	pAAV-AiE0395m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
567944	C57BL/6J	AiE0395m	AiP12498	pAAV-AiE0395m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
616882	C57BL/6J	AiE0395m	AiP12498	pAAV-AiE0395m-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
621446	C57BL/6J	AiE0395m	AiP12498	pAAV-AiE0395m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622084	C57BL/6J	AiE0396h	AiP12158	pAAV-AiE0396h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
626926	C57BL/6J	AiE0396h	AiP12158	pAAV-AiE0396h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
622118	C57BL/6J	AiE0396h	AiP12158	pAAV-AiE0396h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
567945	C57BL/6J	AiE0396h	AiP12158	pAAV-AiE0396h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
622617	C57BL/6J	AiE0396h	AiP12158	pAAV-AiE0396h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
601540	C57BL/6J	AiE0396m	AiP12099	pAAV-AiE0396m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
501913	C57BL/6J	AiE0396m	AiP12099	pAAV-AiE0396m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621447	C57BL/6J	AiE0396m	AiP12099	pAAV-AiE0396m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
515296	C57BL/6J	AiE0397h	AiP12159	pAAV-AiE0397h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
622618	C57BL/6J	AiE0397h	AiP12159	pAAV-AiE0397h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
642125	C57BL/6J	AiE0397m	AiP12123	pAAV-AiE0397m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv

570672	C57BL/6J	AiE0397m	AiP12123	pAAV-AiE0397m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
616881	C57BL/6J	AiE0397m	AiP12123	pAAV-AiE0397m-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
621448	C57BL/6J	AiE0397m	AiP12123	pAAV-AiE0397m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
637261	C57BL/6J	AiE0397m	AiP12123	pAAV-AiE0397m-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
515297	C57BL/6J	AiE0398h	AiP12160	pAAV-AiE0398h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
622619	C57BL/6J	AiE0398h	AiP12160	pAAV-AiE0398h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
500569	C57BL/6J	AiE0398m	AiP12091	pAAV-AiE0398m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621449	C57BL/6J	AiE0398m	AiP12091	pAAV-AiE0398m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
570673	C57BL/6J	AiE0399h	AiP12161	pAAV-AiE0399h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
502938	C57BL/6J	AiE0399m	AiP12105	pAAV-AiE0399m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
622085	C57BL/6J	AiE0400h	AiP12162	pAAV-AiE0400h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
626927	C57BL/6J	AiE0400h	AiP12162	pAAV-AiE0400h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
622119	C57BL/6J	AiE0400h	AiP12162	pAAV-AiE0400h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
570674	C57BL/6J	AiE0400h	AiP12162	pAAV-AiE0400h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621151	C57BL/6J	AiE0400h	AiP12162	pAAV-AiE0400h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
616880	C57BL/6J	AiE0400h	AiP12162	pAAV-AiE0400h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671883	CD-1_IGS	AiE0400h	AiP12162	pAAV-AiE0400h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
601541	C57BL/6J	AiE0400m	AiP12106	pAAV-AiE0400m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
502939	C57BL/6J	AiE0400m	AiP12106	pAAV-AiE0400m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621836	C57BL/6J	AiE0400m	AiP12106	pAAV-AiE0400m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
616884	C57BL/6J	AiE0400m	AiP12106	pAAV-AiE0400m-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
637264	C57BL/6J	AiE0400m	AiP12106	pAAV-AiE0400m-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
626928	C57BL/6J	AiE0401h	AiP12677	pAAV-AiE0401h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
629465	C57BL/6J	AiE0401h	AiP12677	pAAV-AiE0401h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
619575	C57BL/6J	AiE0401h	AiP12677	pAAV-AiE0401h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
570675	C57BL/6J	AiE0401h	AiP12677	pAAV-AiE0401h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
670101	C57BL/6J	AiE0401h	AiP12677	pAAV-AiE0401h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
616879	C57BL/6J	AiE0401h	AiP12677	pAAV-AiE0401h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671884	CD-1_IGS	AiE0401h	AiP12677	pAAV-AiE0401h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
574340	C57BL/6J	AiE0401m	AiP12124	pAAV-AiE0401m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
575095	C57BL/6J	AiE0402h	AiP12163	pAAV-AiE0402h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
500570	C57BL/6J	AiE0402m	AiP12092	pAAV-AiE0402m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621837	C57BL/6J	AiE0402m	AiP12092	pAAV-AiE0402m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622086	C57BL/6J	AiE0403h	AiP12499	pAAV-AiE0403h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
629467	C57BL/6J	AiE0403h	AiP12499	pAAV-AiE0403h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
619576	C57BL/6J	AiE0403h	AiP12499	pAAV-AiE0403h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
551391	C57BL/6J	AiE0403h	AiP12499	pAAV-AiE0403h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621152	C57BL/6J	AiE0403h	AiP12499	pAAV-AiE0403h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
634959	C57BL/6J	AiE0403h	AiP12499	pAAV-AiE0403h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
575092	C57BL/6J	AiE0403m	AiP12125	pAAV-AiE0403m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
515298	C57BL/6J	AiE0404h	AiP12164	pAAV-AiE0404h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621153	C57BL/6J	AiE0404h	AiP12164	pAAV-AiE0404h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
650133	C57BL/6J	AiE0404h	AiP12164	pAAV-AiE0404h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
576008	C57BL/6J	AiE0404m	AiP12500	pAAV-AiE0404m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
575096	C57BL/6J	AiE0405h	AiP12165	pAAV-AiE0405h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
601542	C57BL/6J	AiE0405m	AiP12107	pAAV-AiE0405m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
502940	C57BL/6J	AiE0405m	AiP12107	pAAV-AiE0405m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
616885	C57BL/6J	AiE0405m	AiP12107	pAAV-AiE0405m-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
621838	C57BL/6J	AiE0405m	AiP12107	pAAV-AiE0405m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
575097	C57BL/6J	AiE0406h	AiP12166	pAAV-AiE0406h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
502941	C57BL/6J	AiE0406m	AiP12108	pAAV-AiE0406m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
515299	C57BL/6J	AiE0407h	AiP12167	pAAV-AiE0407h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621154	C57BL/6J	AiE0407h	AiP12167	pAAV-AiE0407h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
637257	C57BL/6J	AiE0407h	AiP12167	pAAV-AiE0407h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
616883	C57BL/6J	AiE0407h	AiP12167	pAAV-AiE0407h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
575093	C57BL/6J	AiE0407m	AiP12126	pAAV-AiE0407m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
515300	C57BL/6J	AiE0408h	AiP12168	pAAV-AiE0408h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
575094	C57BL/6J	AiE0408m	AiP12127	pAAV-AiE0408m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
622087	C57BL/6J	AiE0409h	AiP13062	pAAV-AiE0409h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
629466	C57BL/6J	AiE0409h	AiP13062	pAAV-AiE0409h-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
619577	C57BL/6J	AiE0409h	AiP13062	pAAV-AiE0409h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
641433	C57BL/6J	AiE0409h	AiP13062	pAAV-AiE0409h-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
577081	C57BL/6J	AiE0409h	AiP13062	pAAV-AiE0409h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621155	C57BL/6J	AiE0409h	AiP13062	pAAV-AiE0409h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
576143	C57BL/6J	AiE0409h	AiP13062	pAAV-AiE0409h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622126	C57BL/6J	AiE0409h	AiP13062	pAAV-AiE0409h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
501908	C57BL/6J	AiE0409m	AiP12093	pAAV-AiE0409m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
577082	C57BL/6J	AiE0410h	AiP13063	pAAV-AiE0410h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
573437	C57BL/6J	AiE0410h	AiP13063	pAAV-AiE0410h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
621156	C57BL/6J	AiE0410h	AiP13063	pAAV-AiE0410h-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622127	C57BL/6J	AiE0410h	AiP13063	pAAV-AiE0410h-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
708281	C57BL/6J	AiE0410h_3xC2	AiP14256	pAAV-AiE0410h_3xC2-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
717603	Ai14(RCL-tdT)	AiE0410m	AiP13997	pAAV-AiE0410m-minBG-iCre(R297T)-BGHpA	EPI	Oligo	iCre(R297	RO	Ben-Simon, et al. 2025 Cell
717603	Ai14(RCL-tdT)	AiE0410m	AiP13997	pAAV-AiE0410m-minBG-iCre(R297T)-BGHpA	EPI	Oligo	iCre(R297	RO	Ben-Simon, et al. 2025 Cell
738666	Ai14(RCL-tdT)	AiE0410m	AiP14945	pAAV-AiE0410m-minBG-iCre(R297T)-3XmiR126TS-BGH	STPT, IHC	Oligo	iCre(R297	RO	Ben-Simon, et al. 2025 Cell
717606	Ai14(RCL-tdT)	AiE0410m	AiP14945	pAAV-AiE0410m-minBG-iCre(R297T)-3XmiR126TS-BGH	EPI, IHC	Oligo	iCre(R297	RO	Ben-Simon, et al. 2025 Cell
720915	Ai14(RCL-tdT)	AiE0410m	AiP14946	pAAV-AiE0410m-minBG-iCre(R297T)-3XmiR221TS-BGH	EPI	Oligo	iCre(R297	RO	Ben-Simon, et al. 2025 Cell
608408	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-|	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621235	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-|	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621241	C57BL/6J	AiE0410m	AiP12820	pAAV-AiE0410m-minRho-SYFP2-10aa-H2Bmod-WPRE3-	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
626922	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
629463	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
637135	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
647358	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
647359	C57BL/6J	AiE0410m	AiP14295	pAAV-TelN-LITR-AiE0410m-minBG-SYFP2-P2A-3XFLAG-:	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
648488	C57BL/6J	AiE0410m	AiP14295	pAAV-TelN-LITR-AiE0410m-minBG-SYFP2-P2A-3XFLAG-:	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
617304	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
556486	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
556486	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
502942	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv

574070	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
639180	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
711350	C57BL/6J	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
693219	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
610230	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
693219	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
610230	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
693219	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
693219	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
693219	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
610230	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
693219	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
610230	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
610230	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
610230	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
693219	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
610230	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
631111	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'EPI	Oligo	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
693219	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
610230	C57BL/6J	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671886	CD-1_IGS	AiE0410m	AiP12109	pAAV-AiE0410m-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
676096	CD-1_IGS	AiE0410m	AiP13636	pAAV-AiE0410m-minBG-SYFP2-P2A-3XFLAG-10aa-H2B-'EPI	Oligo	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
619578	C57BL/6J	AiE0410m_3xC2	AiP12556	pAAV-AiE0410m_3xC2-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
621842	C57BL/6J	AiE0410m_3xC2	AiP12556	pAAV-AiE0410m_3xC2-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
711351	C57BL/6J	AiE0410m_3xC2	AiP12556	pAAV-AiE0410m_3xC2-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
579922	C57BL/6J	AiE0410m_3xC2	AiP12556	pAAV-AiE0410m_3xC2-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
524230	C57BL/6J	AiE0495m	AiP12268	pAAV-AiE0495m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
591408	C57BL/6J	AiE0496m	AiP12939	pAAV-AiE0496m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
623025	C57BL/6J	AiE0496m	AiP12939	pAAV-AiE0496m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
575098	C57BL/6J	AiE0497m	AiP12243	pAAV-AiE0497m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
674240	C57BL/6J	AiE0497m	AiP12243	pAAV-AiE0497m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
674240	C57BL/6J	AiE0497m	AiP12243	pAAV-AiE0497m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622081	C57BL/6J	AiE0641m	AiP12845	pAAV-AiE0641m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
626923	C57BL/6J	AiE0641m	AiP12845	pAAV-AiE0641m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
601543	C57BL/6J	AiE0641m	AiP12845	pAAV-AiE0641m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
617305	C57BL/6J	AiE0641m	AiP12845	pAAV-AiE0641m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
563436	C57BL/6J	AiE0641m	AiP12845	pAAV-AiE0641m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
580147	C57BL/6J	AiE0641m	AiP12845	pAAV-AiE0641m-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
621841	C57BL/6J	AiE0641m	AiP12845	pAAV-AiE0641m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671885	CD-1_IGS	AiE0641m	AiP12845	pAAV-AiE0641m-minBG-SYFP2-WPRE3-BGHpA	STPT	Oligo	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
656791	C57BL/6J	AiE0641m_3xC	AiP14004	pAAV-AiE0641m_3xC2-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
563437	C57BL/6J	AiE0642m	AiP12846	pAAV-AiE0642m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
572498	C57BL/6J	AiE0733m	AiP13099	pAAV-AiE0733m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Mich et al., 2025 bioRxiv
570706	C57BL/6J	AiE0733m	AiP13099	pAAV-AiE0733m-minBG-SYFP2-WPRE3-BGHpA	EPI	Oligo	SYFP2	RO	Ben-Simon, et al. 2025 Cell
637134	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
622133	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622133	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622133	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622133	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
609039	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
609039	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
609039	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
609039	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622133	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
609039	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
609039	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622133	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622133	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
609039	C57BL/6J	AiE2117m	AiP1346	pAAV-AiE2117m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
637140	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
642136	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
689258	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604287	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703995	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604287	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604287	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604287	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703995	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
689258	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
689258	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604287	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703995	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604287	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
689258	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604287	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell

1195

638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703995	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703995	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703995	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604287	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
638242	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604287	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
617533	C57BL/6J	AiE2118m	AiP1347	pAAV-AiE2118m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
753053	C57BL/6J	AiE2118m	AiP14191	pAAV-AiE2118m-minBG-SYFP2-4XmiR124T-WPRE3-BGH	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703996	C57BL/6J	AiE2118m	AiP14698	pAAV-AiE2118m-minBG-SYFP2-4XmiR204T-WPRE3-BGH	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703996	C57BL/6J	AiE2118m	AiP14698	pAAV-AiE2118m-minBG-SYFP2-4XmiR204T-WPRE3-BGH	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703996	C57BL/6J	AiE2118m	AiP14698	pAAV-AiE2118m-minBG-SYFP2-4XmiR204T-WPRE3-BGH	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703996	C57BL/6J	AiE2118m	AiP14698	pAAV-AiE2118m-minBG-SYFP2-4XmiR204T-WPRE3-BGH	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
703997	C57BL/6J	AiE2118m	AiP14699	pAAV-AiE2118m-minBG-SYFP2-4XmiR708T-WPRE3-BGH	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613554	C57BL/6J	AiE2119m	AiP1421	pAAV-AiE2119m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613554	C57BL/6J	AiE2119m	AiP1421	pAAV-AiE2119m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600132	C57BL/6J	AiE2119m	AiP1421	pAAV-AiE2119m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613554	C57BL/6J	AiE2119m	AiP1421	pAAV-AiE2119m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613554	C57BL/6J	AiE2119m	AiP1421	pAAV-AiE2119m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600132	C57BL/6J	AiE2119m	AiP1421	pAAV-AiE2119m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600132	C57BL/6J	AiE2119m	AiP1421	pAAV-AiE2119m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613554	C57BL/6J	AiE2119m	AiP1421	pAAV-AiE2119m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600132	C57BL/6J	AiE2119m	AiP1421	pAAV-AiE2119m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600132	C57BL/6J	AiE2119m	AiP1421	pAAV-AiE2119m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
642132	C57BL/6J	AiE2120m	AiP1349	pAAV-AiE2120m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
642137	C57BL/6J	AiE2120m	AiP1349	pAAV-AiE2120m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
622134	C57BL/6J	AiE2120m	AiP1349	pAAV-AiE2120m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
606959	C57BL/6J	AiE2120m	AiP1349	pAAV-AiE2120m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671881	CD-1_IGS	AiE2120m	AiP1349	pAAV-AiE2120m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
642133	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
642135	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
604282	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
615684	C57BL/6J	AiE2121m	AiP1457	pAAV-AiE2121m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
637139	C57BL/6J	AiE2122m	AiP1351	pAAV-AiE2122m-minBG-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
706183	C57BL/6J	AiE2122m	AiP1351	pAAV-AiE2122m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622135	C57BL/6J	AiE2122m	AiP1351	pAAV-AiE2122m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622135	C57BL/6J	AiE2122m	AiP1351	pAAV-AiE2122m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
706183	C57BL/6J	AiE2122m	AiP1351	pAAV-AiE2122m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
706183	C57BL/6J	AiE2122m	AiP1351	pAAV-AiE2122m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
609040	C57BL/6J	AiE2122m	AiP1351	pAAV-AiE2122m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622135	C57BL/6J	AiE2122m	AiP1351	pAAV-AiE2122m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
609040	C57BL/6J	AiE2122m	AiP1351	pAAV-AiE2122m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622135	C57BL/6J	AiE2122m	AiP1351	pAAV-AiE2122m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600471	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600471	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600471	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600471	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600471	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600471	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600471	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600471	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600471	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell

613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
600471	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
613555	C57BL/6J	AiE2123m	AiP1422	pAAV-AiE2123m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
651228	C57BL/6J	AiE2160m	AiP1389	pAAV-AiE2160m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
651229	C57BL/6J	AiE2160m	AiP1389	pAAV-AiE2160m-minBG-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Mich et al., 2025 bioRxiv
622138	C57BL/6J	AiE2160m	AiP1389	pAAV-AiE2160m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
671882	C57BL/6J	AiE2160m	AiP1389	pAAV-AiE2160m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	ICV	Ben-Simon, et al. 2025 Cell
606960	C57BL/6J	AiE2160m	AiP1389	pAAV-AiE2160m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622138	C57BL/6J	AiE2160m	AiP1389	pAAV-AiE2160m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
606960	C57BL/6J	AiE2160m	AiP1389	pAAV-AiE2160m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
622138	C57BL/6J	AiE2160m	AiP1389	pAAV-AiE2160m-minBG-SYFP2-WPRE3-BGHpA	STPT	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
606960	C57BL/6J	AiE2160m	AiP1389	pAAV-AiE2160m-minBG-SYFP2-WPRE3-BGHpA	EPI	Astro	SYFP2	RO	Ben-Simon, et al. 2025 Cell
629461	C57BL/6J	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	EPI, IHC	Astro	SYFP2	RO	Lee et al., 2008 Glia
627900	C57BL/6J	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	EPI	Astro	SYFP2	RO	Lee et al., 2008 Glia
636676	C57BL/6J	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	EPI, IHC	Astro	SYFP2	RO	Lee et al., 2008 Glia
745308	C57BL/6J	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	EPI, IHC	Astro	SYFP2	RO	Lee et al., 2008 Glia
745314	C57BL/6J	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	EPI, IHC	Astro	SYFP2	RO	Lee et al., 2008 Glia
630479	C57BL/6J	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	scRNA-seq	Astro	SYFP2	RO	Lee et al., 2008 Glia
651226	C57BL/6J	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	scRNA-seq	Astro	SYFP2	RO	Lee et al., 2008 Glia
637364	Scn1a +/-	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	EPI, IHC	Astro	SYFP2	RO	Lee et al., 2008 Glia
643123	Scn1a +/-	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	EPI, IHC	Astro	SYFP2	RO	Lee et al., 2008 Glia
637366	Scn1a +/+	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	EPI, IHC	Astro	SYFP2	RO	Lee et al., 2008 Glia
643124	Scn1a +/+	gfaABC1D	AiP13972	pAAV-GfaABC1D-SYFP2-P2A-3XFLAG-10aa-H2B*-WPRE	EPI, IHC	Astro	SYFP2	RO	Lee et al., 2008 Glia
637138	C57BL/6J	MBP promoter	AiP14084	pAAV-mMbp_promoter-SYFP2-P2A-3XFLAG-10aa-H2B*-	EPI, IHC	Oligo	SYFP2	RO	Gow et al., 1992 J. C. B.
642134	C57BL/6J	MBP promoter	AiP14084	pAAV-mMbp_promoter-SYFP2-P2A-3XFLAG-10aa-H2B*-	EPI, IHC	Oligo	SYFP2	RO	Gow et al., 1992 J. C. B.
644029	C57BL/6J	MBP promoter	AiP14084	pAAV-mMbp_promoter-SYFP2-P2A-3XFLAG-10aa-H2B*-	EPI, IHC	Oligo	SYFP2	RO	Gow et al., 1992 J. C. B.
638254	C57BL/6J	MBP promoter	AiP14084	pAAV-mMbp_promoter-SYFP2-P2A-3XFLAG-10aa-H2B*-	scRNA-seq	Oligo	SYFP2	RO	Gow et al., 1992 J. C. B.
651227	C57BL/6J	MBP promoter	AiP14084	pAAV-mMbp_promoter-SYFP2-P2A-3XFLAG-10aa-H2B*-	scRNA-seq	Oligo	SYFP2	RO	Gow et al., 1992 J. C. B.
622080	C57BL/6J	proB12	AiP12214	pAAV-ProB12-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Juttner et al., 2019 Nature Neuroscience
625963	C57BL/6J	proB12	AiP12214	pAAV-ProB12-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Juttner et al., 2019 Nature Neuroscience
636675	C57BL/6J	proB12	AiP12214	pAAV-ProB12-SYFP2-WPRE3-BGHpA	EPI, IHC	Astro	SYFP2	RO	Juttner et al., 2019 Nature Neuroscience
616739	C57BL/6J	ProB12	AiP12214	pAAV-ProB12-SYFP2-WPRE3-BGHpA	scRNA-seq	Astro	SYFP2	RO	Juttner et al., 2019 Nature Neuroscience

Table S2: List of mice used for characterizing the activity of astrocyte and oligodendrocyte-specific enhancer-AAV reporter vectors.

"Mouse ID" is a unique six-digit identifier for each mouse. "Genotype" describes the genotype. "Enhancer ID" is a unique label for each enhancer or promoter element tested (grouped by colors). "Vector ID" is a unique label for the enhancer-AAV DNA packaged into PHP.eB for testing. "Vector Full Name" is a descriptive name of the elements in each enhancer-AAV. "Data Modality" refers to the data collected from each mouse: EPI epifluorescence, IHC immunohistochemistry, STPT serial two-photon tomography, scRNA-seq single cell RNA-seq (SMARTerV4). "Targeted Cell Population" is Astrocytes or Oligodendrocytes. "Fluor" is SYFP2 or iCre(R297T) used to recombine tdTomato in Ai14 mouse line. "Delivery Method" is retro-orbital (RO) or intracerebroventricular (ICV). "Publication" refers to this work (Mich et al.), or if the mice were also reported on in a previous work (Ben-Simon et al.[35]), or whether the enhancer or promoter was previously described and used here as a positive control (Lee et al.[57], Gow et al.[59], Jüttner et al.[41]).

References

References

1.   Khakh, B. S. & Sofroniew, M. V. Diversity of astrocyte functions and phenotypes in
neural circuits. *Nat. Neurosci.* **18**, 942–952 (2015).
2.   Zuchero, J. B. & Barres, B. A. Glia in mammalian development and disease. *Dev.*
Camb. Engl. **142**, 3805–3809 (2015).
3.   Volkenhoff, A. *et al.* Glial Glycolysis Is Essential for Neuronal Survival in Drosophila.
Cell Metab. **22**, 437–447 (2015).
4.   García-Cáceres, C. *et al.* Astrocytic Insulin Signaling Couples Brain Glucose Uptake
with Nutrient Availability. *Cell* **166**, 867–880 (2016).
5.   Molofsky, A. V. *et al.* Astrocyte-encoded positional cues maintain sensorimotor circuit
integrity. *Nature* **509**, 189–194 (2014).
6.   Lioy, D. T. *et al.* A role for glia in the progression of Rett's syndrome. *Nature* **475**, 497–
500 (2011).
7.   Wang, X. *et al.* Astrocytic Ca2+ signaling evoked by sensory stimulation in vivo. *Nat.*
Neurosci. **9**, 816–823 (2006).
8.   Sabelström, H. *et al.* Resident Neural Stem Cells Restrict Tissue Damage and Neuronal
Loss After Spinal Cord Injury in Mice. *Science* **342**, 637–640 (2013).
9.   Liddelow, S. A. *et al.* Neurotoxic reactive astrocytes are induced by activated microglia.
Nature **541**, 481–487 (2017).
10.   Hasel, P., Rose, I. V. L., Sadick, J. S., Kim, R. D. & Liddelow, S. A. Neuroinflammatory
astrocyte subtypes in the mouse brain. *Nat. Neurosci.* **24**, 1475–1487 (2021).
11.   Zuchero, J. B. *et al.* CNS Myelin Wrapping Is Driven by Actin Disassembly. *Dev. Cell*
34, 152–167 (2015).
12.   Steadman, P. E. *et al.* Disruption of Oligodendrogenesis Impairs Memory Consolidation
in Adult Mice. *Neuron* **105**, 150-164.e6 (2020).
13.   Wilkins, A., Majed, H., Layfield, R., Compston, A. & Chandran, S. Oligodendrocytes
Promote Neuronal Survival and Axonal Length by Distinct Intracellular Mechanisms: A Novel
Role for Oligodendrocyte-Derived Glial Cell Line-Derived Neurotrophic Factor. *J. Neurosci.* **23**,
4967–4974 (2003).
14.   Knowles, J. K. *et al.* Maladaptive myelination promotes generalized epilepsy
progression. *Nat. Neurosci.* **25**, 596–606 (2022).
15.   Zhang, P. *et al.* Senolytic therapy alleviates Aβ-associated oligodendrocyte progenitor
cell senescence and cognitive deficits in an Alzheimer's disease model. *Nat. Neurosci.* 1
(2019) doi:10.1038/s41593-019-0372-9.
16.   Boisvert, M. M., Erikson, G. A., Shokhirev, M. N. & Allen, N. J. The Aging Astrocyte
Transcriptome from Multiple Regions of the Mouse Brain. *Cell Rep.* **22**, 269–285 (2018).
17.   Endo, F. *et al.* Molecular basis of astrocyte diversity and morphology across the CNS in
health and disease. *Science* **378**, eadc9020 (2022).
18.   Yao, Z. *et al.* A high-resolution transcriptomic and spatial atlas of cell types in the whole
mouse brain. 2023.03.06.531121 Preprint at https://doi.org/10.1101/2023.03.06.531121
(2023).
19.   Hodge, R. D. *et al.* Conserved cell types with divergent features in human versus
mouse cortex. *Nature* **573**, 61–68 (2019).
20.   Chen, Z.-P. *et al.* Lipid-accumulated reactive astrocytes promote disease progression in
epilepsy. *Nat. Neurosci.* **26**, 542–554 (2023).

21.   Wang, C. *et al.* Selective removal of astrocytic APOE4 strongly protects against tau-
mediated neurodegeneration and decreases synaptic phagocytosis by microglia. *Neuron*
(2021) doi:10.1016/j.neuron.2021.03.024.
22.   Monje, M. *et al.* Hedgehog-responsive candidate cell of origin for diffuse intrinsic
pontine glioma. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 4453–4458 (2011).
23.   Alcantara Llaguno, S. *et al.* Malignant Astrocytomas Originate from Neural
Stem/Progenitor Cells in a Somatic Tumor Suppressor Mouse Model. *Cancer Cell* **15**, 45–56
(2009).
24.   Deverman, B. E. *et al.* Cre-dependent selection yields AAV variants for widespread
gene transfer to the adult brain. *Nat. Biotechnol.* **34**, 204–209 (2016).
25.   Chan, K. Y. *et al.* Engineered AAVs for efficient noninvasive gene delivery to the central
and peripheral nervous systems. *Nat. Neurosci.* **20**, 1172–1179 (2017).
26.   Nonnenmacher, M. *et al.* Rapid evolution of blood-brain-barrier-penetrating AAV
capsids by RNA-driven biopanning. *Mol. Ther. - Methods Clin. Dev.* **20**, 366–378 (2021).
27.   De, A., El-Shamayleh, Y. & Horwitz, G. D. Fast and reversible neural inactivation in
macaque cortex by optogenetic stimulation of GABAergic neurons. *eLife* **9**, e52658 (2020).
28.   El-Shamayleh, Y., Kojima, Y., Soetedjo, R. & Horwitz, G. D. Selective Optogenetic
Control of Purkinje Cells in Monkey Cerebellum. *Neuron* **95**, 51-62.e4 (2017).
29.   Mendell, J. R. *et al.* Single-Dose Gene-Replacement Therapy for Spinal Muscular
Atrophy. *http://dx.doi.org/10.1056/NEJMoa1706198*
https://www.nejm.org/doi/10.1056/NEJMoa1706198?url_ver=Z39.88-
2003&rfr_id=ori%3Arid%3Acrossref.org&rfr_dat=cr_pub%3Dwww.ncbi.nlm.nih.gov (2017)
doi:10.1056/NEJMoa1706198.
30.   Dimidschstein, J. *et al.* A viral strategy for targeting and manipulating interneurons
across vertebrate species. *Nat. Neurosci.* **19**, 1743–1749 (2016).
31.   Vormstein-Schneider, D. *et al.* Viral manipulation of functionally distinct interneurons in
mice, non-human primates and humans. *Nat. Neurosci.* 1–8 (2020) doi:10.1038/s41593-020-
0692-9.
32.   Hrvatin, S. *et al.* A scalable platform for the development of cell-type-specific viral
drivers. *eLife* **8**, e48089 (2019).
33.   Mich, J. K. *et al.* Functional enhancer elements drive subclass-selective expression
from mouse to primate neocortex. *Cell Rep.* **34**, 108754 (2021).
34.   Graybuck, L. T. *et al.* Enhancer viruses for combinatorial cell-subclass-specific labeling.
Neuron **109**, 1449-1464.e13 (2021).
35.   Ben-Simon, Y. *et al.* A suite of enhancer AAVs and transgenic mouse lines for genetic
access to cortical cell types. *Cell* (2025) doi:10.1016/j.cell.2025.05.002.
36.   Johansen, N. J. *et al.* Evaluating methods for the prediction of cell-type-specific
enhancers in the mammalian cortex. *Cell Genomics* 100879 (2025)
doi:10.1016/j.xgen.2025.100879.
37.   Kussick, E. *et al.* Enhancer AAVs for targeting spinal motor neurons and descending
motor pathways in rodents and macaque. *Cell Rep.* 115730 (2025)
doi:10.1016/j.celrep.2025.115730.
38.   Hunker, A. C. *et al.* Enhancer AAV toolbox for accessing and perturbing striatal cell
types and circuits. *Neuron* **113**, 1507-1524.e17 (2025).
39.   Furlanis, E. *et al.* An enhancer-AAV toolbox to target and manipulate distinct
interneuron subtypes. *Neuron* **113**, 1525-1547.e15 (2025).

40.   Li, L. *et al.* Identification and application of cell-type-specific enhancers for the macaque
brain. *Cell* **0**, (2025).
41.   Jüttner, J. *et al.* Targeting neuronal and glial cell types with synthetic promoter AAVs in
mice, non-human primates and humans. *Nat. Neurosci.* **22**, 1345–1356 (2019).
42.   Lin, C.-H. *et al.* Identification of cis-regulatory modules for adeno-associated virus-
based cell-type-specific targeting in the retina and brain. *J. Biol. Chem.* **298**, (2022).
43.   Gleichman, A. J., Kawaguchi, R., Sofroniew, M. V. & Carmichael, S. T. A toolbox of
astrocyte-specific, serotype-independent adeno-associated viral vectors using microRNA
targeting sequences. *Nat. Commun.* **14**, 7426 (2023).
44.   Wang, L.-L. *et al.* Revisiting astrocyte to neuron conversion with lineage tracing in vivo.
Cell **184**, 5465-5481.e16 (2021).
45.   Xie, Y., Zhou, J. & Chen, B. Critical examination of Ptbp1-mediated glia-to-neuron
conversion in the mouse retina. *Cell Rep.* **39**, 110960 (2022).
46.   Le, N., Appel, H., Pannullo, N., Hoang, T. & Blackshaw, S. Ectopic insert-dependent
neuronal expression of GFAP promoter-driven AAV constructs in adult mouse retina. *Front.*
Cell Dev. Biol. **10**, (2022).
47.   Borden, P. M. *et al.* A fast genetically encoded fluorescent sensor for faithful in vivo
acetylcholine detection in mice, fish, worms and flies. 2020.02.07.939504 Preprint at
https://doi.org/10.1101/2020.02.07.939504 (2020).
48.   Fröhlich, D. *et al.* Dual-function AAV gene therapy reverses late-stage Canavan disease
pathology in mice. *Front. Mol. Neurosci.* **15**, 1061257 (2022).
49.   Hillen, A. E. J. *et al.* In vivo targeting of a variant causing vanishing white matter using
CRISPR/Cas9. *Mol. Ther. - Methods Clin. Dev.* **25**, 17–25 (2022).
50.   Hagemann, T. L. *et al.* Antisense therapy in a rat model of Alexander disease reverses
GFAP pathology, white matter deficits, and motor impairment. *Sci. Transl. Med.* **13**, eabg4711
(2021).
51.   Lister, R. *et al.* Global Epigenomic Reconfiguration During Mammalian Brain
Development. *Science* **341**, 1237905 (2013).
52.   Luo, C. *et al.* Single-cell methylomes identify neuronal subtypes and regulatory
elements in mammalian cortex. *Science* **357**, 600–604 (2017).
53.   Lee, D.-S. *et al.* Simultaneous profiling of 3D genome structure and DNA methylation in
single human cells. *Nat. Methods* **16**, 999–1006 (2019).
54.   Liu, H. *et al.* DNA methylation atlas of the mouse brain at single-cell resolution. *Nature*
598, 120–128 (2021).
55.   Fullard, J. F. *et al.* Open chromatin profiling of human postmortem brain infers functional
roles for non-coding schizophrenia loci. *Hum. Mol. Genet.* (2019) doi:10.1093/hmg/ddy229.
56.   Li, Y. E. *et al.* An atlas of gene regulatory elements in adult mouse cerebrum. *Nature*
598, 129–136 (2021).
57.   Lee, Y., Messing, A., Su, M. & Brenner, M. GFAP promoter elements required for
region-specific and astrocyte-specific expression. *Glia* **56**, 481–493 (2008).
58.   Sun, W. *et al.* SOX9 Is an Astrocyte-Specific Nuclear Marker in the Adult Brain Outside
the Neurogenic Regions. *J. Neurosci.* **37**, 4493–4507 (2017).
59.   Gow, A., Friedrich, V. L., Jr & Lazzarini, R. A. Myelin basic protein gene contains
separate enhancers for oligodendrocyte and Schwann cell expression. *J. Cell Biol.* **119**, 605–
616 (1992).

60.   Bin, J. M., Harris, S. N. & Kennedy, T. E. The oligodendrocyte-specific antibody 'CC1'
binds Quaking 7. *J. Neurochem.* **139**, 181–186 (2016).
61.   Bailey, T. L. *et al.* MEME Suite: tools for motif discovery and searching. *Nucleic Acids*
Res. **37**, W202–W208 (2009).
62.   Hume, M. A., Barrera, L. A., Gisselbrecht, S. S. & Bulyk, M. L. UniPROBE, update
2015: new tools and content for the online database of protein-binding microarray data on
protein–DNA interactions. *Nucleic Acids Res.* **43**, D117–D122 (2015).
63.   Kulakovskiy, I. V. *et al.* HOCOMOCO: towards a complete collection of transcription
factor binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids*
Res. **46**, D252–D259 (2018).
64.   Castro-Mondragon, J. A. *et al.* JASPAR 2022: the 9th release of the open-access
database of transcription factor binding profiles. *Nucleic Acids Res.* **50**, D165 (2022).
65.   Furushima, K., Murata, T., Kiyonari, H. & Aizawa, S. Characterization of Opr deficiency
in mouse brain: Subtle defects in dorsomedial telencephalon and medioventral forebrain. *Dev.*
Dyn. **232**, 1056–1061 (2005).
66.   Inoue, T., Ota, M., Ogawa, M., Mikoshiba, K. & Aruga, J. Zic1 and Zic3 Regulate Medial
Forebrain Development through Expansion of Neuronal Progenitors. *J. Neurosci.* **27**, 5461–
5473 (2007).
67.   Hornig, J. *et al.* The Transcription Factors Sox10 and Myrf Define an Essential
Regulatory Network Module in Differentiating Oligodendrocytes. *PLOS Genet.* **9**, e1003907
(2013).
68.   Turnescu, T. *et al.* Sox8 and Sox10 jointly maintain myelin gene expression in
oligodendrocytes. *Glia* **66**, 279–294 (2018).
69.   Hinderer, C. *et al.* Severe Toxicity in Nonhuman Primates and Piglets Following High-
Dose Intravenous Administration of an Adeno-Associated Virus Vector Expressing Human
SMN. *Hum. Gene Ther.* **29**, 285–298 (2018).
70.   Zhang, K. *et al. A Cell Atlas of Chromatin Accessibility across 25 Adult Human Tissues*.
2021.02.17.431699 https://www.biorxiv.org/content/10.1101/2021.02.17.431699v1 (2021)
doi:10.1101/2021.02.17.431699.
71.   Goertsen, D. *et al.* AAV capsid variants with brain-wide transgene expression and
decreased liver targeting after intravenous delivery in mouse and marmoset. *Nat. Neurosci.* **25**,
106–115 (2022).
72.   Martín-Suárez, S., Abiega, O., Ricobaraza, A., Hernandez-Alcoceba, R. & Encinas, J.
M. Alterations of the Hippocampal Neurogenic Niche in a Mouse Model of Dravet Syndrome.
Front. Cell Dev. Biol. **8**, (2020).
73.   Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization
system for the whole mouse brain. *Nat. Neurosci.* **13**, 133–140 (2010).
74.   Hartung, M. & Kisters-Woike, B. Cre Mutants with Altered DNA Binding Properties*. *J.*
Biol. Chem. **273**, 22884–22891 (1998).
75.   Wang, S. *et al.* The Endothelial-Specific MicroRNA miR-126 Governs Vascular Integrity
and Angiogenesis. *Dev. Cell* **15**, 261–271 (2008).
76.   Sayeg, M. K. *et al.* Rationally Designed MicroRNA-Based Genetic Classifiers Target
Specific Neurons in the Brain. *ACS Synth. Biol.* **4**, 788–795 (2015).
77.   Cho, F. S. *et al.* Enhancing GAT-3 in thalamic astrocytes promotes resilience to brain
injury in rodents. *Sci. Transl. Med.* **14**, eabj4310 (2022).

78. Hasel, P., Aisenberg, W. H., Bennett, F. C. & Liddelow, S. A. Molecular and metabolic heterogeneity of astrocytes and microglia. *Cell Metab.* **35**, 555–570 (2023).

79. Aida, T. *et al.* Astroglial glutamate transporter deficiency increases synaptic excitability and leads to pathological repetitive behaviors in mice. *Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol.* **40**, 1569–1579 (2015).

80. Li, Z. *et al.* Cell-Type-Specific Afferent Innervation of the Nucleus Accumbens Core and Shell. *Front. Neuroanat.* **12**, (2018).

81. Beier, K. T. *et al.* Circuit Architecture of VTA Dopamine Neurons Revealed by Systematic Input-Output Mapping. *Cell* **162**, 622–634 (2015).

82. Ren, J. *et al.* Anatomically Defined and Functionally Distinct Dorsal Raphe Serotonin Sub-systems. *Cell* **175**, 472-487.e20 (2018).

83. Ren, J. *et al.* Single-cell transcriptomes and whole-brain projections of serotonin neurons in the mouse dorsal and median raphe nuclei. *eLife* **8**, e49424 (2019).

84. Gabitto, M. I. *et al.* Integrated multimodal cell atlas of Alzheimer's disease. 2023.05.08.539485 Preprint at https://doi.org/10.1101/2023.05.08.539485 (2023).

85. Kamath, T. *et al.* Single-cell genomic profiling of human dopamine neurons identifies a population that selectively degenerates in Parkinson's disease. *Nat. Neurosci.* **25**, 588–595 (2022).

86. Pagès, Hervé. BSgenome: Software infrastructure for efficient representation of full genomes and their SNPs. R package version 1.62.0. (2021).

87. Kim, J.-Y., Grunke, S. D., Levites, Y., Golde, T. E. & Jankowsky, J. L. Intracerebroventricular Viral Injection of the Neonatal Mouse Brain for Persistent and Widespread Neuronal Transduction. *JoVE J. Vis. Exp.* e51863 (2014) doi:10.3791/51863.

88. Ragan, T. *et al.* Serial two-photon tomography for automated ex vivo mouse brain imaging. *Nat. Methods* **9**, 255–258 (2012).

89. Tasic, B. *et al.* Shared and distinct transcriptomic cell types across neocortical areas. *Nature* **563**, 72 (2018).

90. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-connected communities. *Sci. Rep.* **9**, 5233 (2019).

91. Choi, J.-H. *et al.* Optimization of AAV expression cassettes to improve packaging capacity and transgene expression in neurons. *Mol. Brain* **7**, 17 (2014).

92. Bari, B. A. *et al.* Stable Representations of Decision Variables for Flexible Behavior. *Neuron* **103**, 922-933.e7 (2019).

93. Akam, T. & Walton, M. E. pyPhotometry: Open source Python based hardware and software for fiber photometry data acquisition. *Sci. Rep.* **9**, 3521 (2019).

94. Fullard, J. F. *et al.* An atlas of chromatin accessibility in the adult human brain. *Genome Res.* **28**, 1243–1252 (2018).