DEAD box RNA helicase 5 is a new pro-viral host factor for Sindbis virus infection

Mélanie Messmer [1], Louison Pierson [1], Charline Pasquier [1], Nikola Djordjevic [1], Johana

Chicher [2], Philippe Hammann [2], Sébastien Pfeffer [1], Erika Girardi [1]

[1] Université de Strasbourg, Architecture et Réactivité de l'ARN, Institut de Biologie

Moléculaire et Cellulaire du CNRS, 2 allée Konrad Roentgen, 67084 Strasbourg France.

[2] Université de Strasbourg, Institut de Biologie Moléculaire et Cellulaire du CNRS,

Plateforme Protéomique Strasbourg – Esplanade, 2 allée Konrad Roentgen, 67084 Strasbourg

France.

* To whom correspondence should be addressed: e.girardi@ibmc-cnrs.unistra.fr.

Abstract

Background

RNA helicases are emerging as key factors regulating host-virus interactions. The DEAD-box

ATP-dependent RNA helicase DDX5, which plays an important role in many aspects of

cellular RNA biology, was also found to either promote or inhibit viral replication upon

infection with several RNA viruses. Here, our aim is to examine the impact of DDX5 on

Sindbis virus (SINV) infection.

Methods

We analysed the interaction between DDX5 and the viral RNA using imaging and RNA-

immunoprecipitation approaches. The interactome of DDX5 in mock- and SINV-infected

cells was determined by mass spectrometry. We validated the interaction with DDX17 and the

viral capsid by co- immunoprecipitation in the presence or absence of an RNase treatment.

We determined the subcellular localization of DDX5, its cofactor DDX17 and the viral capsid

protein by co-immunofluorescence. Finally, we investigated the impact of DDX5 depletion

and overexpression on SINV infection at the viral protein, RNA and infectious particle

accumulation level. The contribution of DDX17 was also tested by knockdown experiments.

Results

In this study we demonstrate that DDX5 interacts with the SINV RNA during infection.

Furthermore, the proteomic analysis of the DDX5 interactome in mock and SINV-infected

HCT116 cells identified new cellular and viral partners and confirmed the interaction between

DDX5 and DDX17. Both DDX5 and DDX17 re-localize from the nucleus to the cytoplasm

upon SINV infection and interact with the viral capsid protein. We also show that DDX5

depletion negatively impacts the viral replication cycle, while its overexpression has a pro-

viral effect. Finally, we observed that DDX17 depletion reduces SINV infection, an effect

which is even more pronounced in a DDX5-depleted background, suggesting a synergistic

pro-viral effect of the DDX5 and DDX17 proteins on SINV.

Conclusions

These results not only shed light on DDX5 as a novel and important host factor to the SINV

life cycle, but also expand our understanding of the roles played by DDX5 and DDX17 as

regulators of viral infections.

Keywords: DDX5, DDX17, helicase, RNA, Sindbis virus

1. Background

DEAD box RNA helicases play crucial molecular functions in virtually all aspects of RNA metabolism, including transcription, pre-mRNA splicing, RNA transport, rRNA biogenesis and RNA decay (1). These enzymes can structurally remodel short RNA duplex molecules and rearrange RNA-protein interactions in an ATP-dependent fashion. The DEAD-box RNA helicase family is characterized by the presence of specific highly conserved motifs, including the Asp–Glu–Ala–Asp (D-E-A-D) box within the catalytic core (2) from which this family gets its name. Recently, DEAD box RNA helicases have emerged as pivotal host factors upon viral infection that act either by modulating antiviral signalling pathways or by directly recognizing viral RNA and acting at different steps of the viral life cycle (3–5).

The multifunctional DEAD-box RNA helicase DDX5 (also known as p68) is an essential splicing factor (6,7) involved in many other aspects of cellular RNA metabolism (8). Within the DDX gene family members, DDX5 exhibits the highest degree of homology with DDX17 (also known as p72) (9). These two proteins can form homodimers but are preferentially associated in hetero-dimers (10). DDX5 and DDX17 exhibit partially overlapping functions, including involvement in transcriptional regulation (11), alternative splicing modulation (12) and ribosomal biogenesis (13).

A growing body of evidence suggests that DDX5 and DDX17 play crucial roles during viral infection (14,15). Despite their known function as a heterodimer, these two proteins exhibit distinct functions in response to specific viral infections. For instance, depletion of DDX17 but not of DDX5 increases infection of Rift Valley Fever virus (RVFV), a segmented, negative-sense RNA virus. In particular, DDX17 binding to a stem-loop structure in the viral RNA is sufficient to trigger the antiviral effect (16).

Moreover, the same protein can have either a positive or a negative effect on different viruses. While DDX5 is antiviral in response to DNA viruses such as hepatitis B virus (HBV) (17) or

myxoma virus (MYXV) (18), it acts as a positive cofactor of the Human Immunodeficiency

Virus 1(HIV-1) proteins Rev and Tat facilitating viral replication (19,20). Interestingly,

several positive, single-stranded RNA viruses with a cytoplasmic replication cycle have also

evolved to hijack DDX5 and favour their own replication. DDX5 was also shown to be pro-

viral for hepatitis C virus (HCV) (21,22), severe acute respiratory syndrome coronavirus

(SARS-CoV) (23), japanese encephalitis virus (JEV) (24) and porcine reproductive and

respiratory syndrome virus (PRRSV) (25). In most cases, DDX5 promotes viral replication

through direct interaction with specific viral proteins and/or viral RNA.

In a previous study we identified DDX5 as one of the factors associated with the Sindbis virus

(SINV) replication complex by dsRNA-IP coupled to mass spectrometry (DRIMS) and

observed that DDX5 knockdown (KD) significantly reduces GFP expression in HCT116 cells

infected with a GFP-expressing SINV (26). SINV is a small, enveloped, arthropod-borne

virus, which belongs to the *Alphavirus* genus from the *Togaviridae* family. Alphaviruses are

widely distributed viruses (27) that pose a re-emerging threat to human health due to their

potential to cause severe arthritogenic and neurological diseases and for which approved

vaccines are not yet available. Most of our understanding on the alphavirus life cycle derives

from research on the prototypical model SINV. The virus possesses a positive, single-stranded

genomic RNA of approximately 11 kb, capped and polyadenylated, organized into two

distinct open reading frames (ORFs). The first ORF encodes the non-structural proteins (nsP1,

nsP2, nsP3 and nsP4) necessary for viral RNA replication. An antigenomic RNA accumulates

in the cytoplasm and serves as a template for the synthesis of both genomic and sub-genomic

RNA. The second ORF is translated from the sub-genomic RNA and gives rise to the

structural proteins (Capsid, E3, E2, 6K and E1) needed for viral particle assembly (28).

In this study we investigated the impact of DDX5 on SINV. Initially, we observed the co-

localization and interaction between DDX5 and the viral RNA in the cytoplasm of infected

cells by imaging and RNA IP approaches. We then conducted a proteomic analysis to

determine the interactome of DDX5 under both uninfected and SINV-infected conditions.

DDX17, a previously well-known DDX5 partner, emerged among the most significant hits.

Interestingly, we demonstrated that DDX5 and DDX17 re-localize to the cytoplasm upon

SINV infection and identified a novel RNA-independent interaction between DDX5 and viral

capsid proteins.

Functionally, our study demonstrated that reducing DDX5 levels by either knock-down or

knock-out approaches diminishes SINV replication in human HCT116 cells. Moreover, stable

overexpression of DDX5 in *DDX5* knock-out cells restored the capsid protein expression and

the viral particle production. Finally, we showed that knockdown of DDX17 has a negative

impact on SINV viral cycle with a more pronounced decrease in cells lacking both DDX17

and DDX5, underscoring the importance of both DDX5 and DDX17 in promoting SINV

infection.

2. Methods

2.1. Cell lines and viruses

HCT116, BHK21 and Vero E6 cells were maintained at 37°C in a humidified atmosphere containing 5% CO_2 and cultured in Dulbecco's Modified Eagle medium (DMEM; Invitrogen) supplemented with 10% FBS. Viral stocks were prepared from infectious clones (kindly provided by Dr Carla Saleh, Institut Pasteur, Paris, France) carrying either the wild-type or a GFP containing version of the SINV genomic sequence as previously described in (29). Briefly, the plasmids were linearized by XhoI digestion and used as a substrate for *in vitro* transcription using the mMESSAGE mMACHINE capped RNA transcription kit (Ambion, Thermo Fisher Scientific Inc.). In vitro synthesized viral RNA was transfected in BHK21 cells, viral infectious particles were collected 48 hpt and used to infect BHK21 for the full viral production. Titers were measured by standard plaque assay in Vero E6 cells. Cells were infected at a MOI of 1, 10^{-1} or 10^{-2} as indicated in the figure legends.

2.2. Plaque assay

For plaque assay, 1 million Vero E6 cells were seeded in a 96 well plate and infected for 1 hour with viral supernatants prepared in 10-fold dilution cascade. After removal of the viral inoculum, 2.5% carboxymethyl cellulose diluted in DMEM-FBS10% was added and cells were incubated at 37ºC in a humidified atmosphere of 5% CO2. Plaques were counted manually under the microscope after 72 h.

2.3. DDX5 CRISPR/Cas9 knockout

Guide RNA (gRNA) sequences targeting the human DDX5 gene were designed using the CRISPOR tool (http://crispor.tefor.net) and cloned into the pX459-V2 vector (Addgene

Plasmid #62988 was a gift from Feng Zhang (30)). Briefly 0.25 µM of the annealed oligos

and 100 ng of digested vector were ligated and transformed into DH5 alpha cells.

guideRNA#2 DDX5 sense: 5' CACCGATAATAGGGTGTTCATAGGT 3'

guideRNA#2 DDX5 antisense: 5' AAACACCTATGAACACCCTATTATC 3'

guideRNA#1 DDX5 sense: 5' CACCGCCCTACTTCCTCCAAATCG 3'

guideRNA#1 DDX5 antisense: 5' AAACCGATTTGGAGGAAGTAGGGC 3'

HCT116 cells were transfected with 2 plasmids containing the sequence of gRNA#1 and #2,

respectively. Twenty-four hours post-transfection, cells were treated with 1 µg/mL puromycin

for 48h. Surviving cells were diluted to obtain 0.5 cells/ well in 96 well plates. Two weeks

later, cellular genomic DNA was extracted from individual colonies. Cells were lysed in

genomic DNA extraction buffer (50 mM Tris-HCl [pH 8.0]; 100 mM EDTA [pH 8.0], 100

mM NaCl, 1% SDS) containing 0.1 mg of proteinase K and incubated overnight at 55°C.

Then, 50 ng of genomic DNA were amplified with the GoTaq (Promega) using specific

primers (IDT) to detect the deletion (DDX5 FW: 5'-ATAAATCCCCGGCTTCCGAC-3';

DDX5 RV: 5'-AGAGGGGGTAGGTGGAAACAA-3'). Wild type genomic DNA was used

as control template. PCR reactions were loaded on a 1 % agarose gel and the obtained

amplicons were gel purified and sequenced by Sanger sequencing.

2.4. SiRNA-based knockdown

For the knockdown experiments, 0.6 millions of cells HCT116 cells were reverse-transfected

with 20nM of ON-TARGETplus Human siRNAs (Horizon) against DDX5 (L-003774-00-

0005), DDX17 (L-013450-01-0005) or non-targeting Control Pool (D-001810-10-05) using

the Lipofectamine 2000 transfection reagent (Thermo Fisher Scientific, #11668019)

according to the manufacturer's instructions in 6 well plates. After 24 hours, cells were

transfected again with 20nM of the same mix of siRNAs and incubated overnight. Cells were

then infected with SINV-GFP at a MOI of 0.1 for 24 hours. Supernatant were collected for

plaque assay experiment and proteins and RNA were collected for western blot and RT-qPCR

analyses, respectively.

2.5. Lentivirus production and generation of stable cell lines

Lentiviral particles were produced by transfecting seven millions of HEK293T cells with 12

µg of either pLV-DDX5-V5 or pLV-BFP viral vector (Vector Builder), 2.4 µg and 9,6µg of

the packaging plasmids pCMV-VSV-G (Addgene #8454) and psPAX2 (Addgene #12260)

respectively, using Lipofectamine 2000 reagent (Invitrogen, Fisher Scientific) according to

the manufacturer's protocol. The viral supernatants were collected after 48 hours and filtered

using a 0.45µm PES filter.

DDX5 KO HCT116 cells were transduced in 6-well plates using the lentiviral supernatant

diluted in fresh medium supplemented with 4µg/mL of polybrene (Merck, Sigma-Aldrich).

The medium was changed after 8 h and the selection with the appropriate antibiotic was

initiated after 48 h. Polyclonal V5-DDX5-*DDX5* KO HCT116 (V5-DDX5) and BFP-*DDX5*

KO HCT116 (BFP) cells were maintained under selection for a minimum of 10 days before

analysis. In detail, two-hundred thousand of V5-DDX5 or BFP-DDX5 stable cells were

infected with SINV-GFP for 24 hours (MOI of 0.1). before collecting the supernatant for

plaque assay experiment, as well as proteins and RNA for western blot and RT-qPCR

analyses, respectively.

2.6. Live-cell imaging

Two-hundred thousand WT, *DDX5* KO, BFP or V5-DDX5- HCT116 cells were seeded in 12

well plate and infected with SINV-GFP at the indicated MOI. Uninfected cells were used as

control. GFP fluorescence and phase contrast were observed using a CellcyteX live-cell

imaging system (Discover Echo). Four to six images per well (10X objective) were acquired

every 3 or 6 hours for 72hrs and were analysed with the Cellcyte Studio software to determine

cell confluency and GFP relative intensity.

2.7. Cloning

Amplicons containing the coding sequence for the individual viral proteins were generated by

PCR from the plasmid carrying the SINV genomic sequence (kindly provided by Dr. Carla

Saleh, Institut Pasteur, Paris, France) and cloned into the pDONOR221 vector (Invitrogen) by

recombination using the Gateway BP clonase (Invitrogen) according to the manufacturer

instructions. The pDEST-FLAG-HA- nsP1, pDEST-FLAG-HA- nsP2, pDEST-FLAG-HA-

nsP3, pDEST-FLAG-HA- nsP4 and pDEST-FLAG-HA- capsid vectors were obtained by

recombination using Gateway LR clonase (Invitrogen) in the pDEST- FLAG-HA vector

according to the manufacturer instructions, as in (31).Primers used for the PCR are listed

below:

pDONOR NSP1 FW

5' GGGGACAAGTTTGTACAAAAAAGCAGGCTTCGAGAAGCCAGTAGTAAAC-3'

pDONOR NSP1 REV

5'-GGGGACCACTTTGTACAAGAAAGCTGGGTCTTATTTGCTCCGATGTCCG-3'

pDONOR NSP2 FW

5'GGGGACAAGTTTGTACAAAAAAGCAGGCTTCGCATTAGTTGAAACCCCG-3'

pDONOR NSP2 REV

5'GGGGACCACTTTGTACAAGAAAGCTGGGTCTTATTGGCTCCAACTCCATCTC-3'

pDONOR NSP3 FW

5'-GGGGACAAGTTTGTACAAAAAAGCAGGCTTCGCGCCGTCATACCGCACC-3'

pDONOR NSP3 REV

5'-GGGGACCACTTTGTACAAGAAAGCTGGGTCTTATTGTATTCAGTCCTCC-3'

pDONOR NSP4 FW

5'-GGGGACAAGTTTGTACAAAAAAGCAGGCTTCCTAACCGGGGTAGGTGGGTAC

3'

pDONOR NSP4 REV

5'GGGGACCACTTTGTACAAGAAAGCTGGGTCTTACTATTTAGGACCACCGTAGAG

-3'

pDONOR capsid FW

5'-GGGGACAAGTTTGTACAAAAAAGCAGGCTTCAATAGAGGATTCTTTAAC-3'

pDONOR capsid REV

5'-GGGGACCACTTTGTACAAGAAAGCTGGGTCTTATTCCACTCTTCTGTCC-3'

2.8. Flag-HA SINV protein co-IP

For the experiment, 0.6 millions of HCT 116 cells were seeded in 6 well plates. The following

day 3µg of the pDEST-HA-FLAG described above were transfected using the Lipofectamine

2000 transfection reagent as per manufacturer's recommendations. 48 hours later, the cells

were washed once with 1xPBS and lysed in 600 µL of Miltenyi lysate buffer (50 mM Tris-

HCl pH 7.5, 140 mM NaCl, 1.5 mM $MgCl_2$, 0.1% NP-40), supplemented with Complete-

EDTA-free Protease Inhibitor Cocktail (Merck). Cells were lysed for 30 min incubation on

ice and debris were removed by 10 min centrifugation at 10,000g at 4°C. An aliquot of the

cleared lysates (50 µL) was kept aside as protein Input. Samples were divided into equal parts

(250 µL each) and incubated with 50 µL of magnetic microparticles coated with monoclonal

HA or MYC antibodies (MACS purification system, Miltenyi Biotec) at 4°C for 1 hour under

rotation (10 rpm). Samples were passed through µ Columns (MACS purification system,

Miltenyi Biotec). The µ Columns were then washed 4 times with 200 µL of WASH buffer1

and 1 time with 200 µL of WASH buffer 2. To elute the immunoprecipitated proteins, 70µL

of 95°C pre-warmed 2x Western blot loading buffer (10% glycerol, 4% SDS, 62.5 mM Tris-

HCl pH 6.8, 5% (v/v) 2-β-mercaptoethanol, Bromophenol Blue) was passed through the µ

Columns. Proteins were analysed by western blotting.

2.9. DDX5 and DDX17 co-immunoprecipitation and RNA immunoprecipitation

DDX5 and DDX17 immunoprecipitations (IP) were performed as in (26) with some

modifications. Briefly, 4 millions of mock or SINV-GFP infected HCT116 cells (MOI 0.1, 24

hpi) were lysed using RIP immunoprecipitation buffer (50mM tris-HCL [pH 7.5], 150mM

NaCl, 5 mM EDTA, 0.05% SDS, 1% triton, 1 tablet of commercial protease inhibitor cocktail

(Merck)). RNase treatment was performed on the lysates by adding 1µL of RNase A at 10

mg/mL (Thermo Fisher) and incubating for 25 min at 37°C. DNase treatment was performed

on the lysates by adding 1µL of RNase-free DNase I at 1U/µL (Thermo Fisher Scientific),

10mM MgCl2, 5mM Cacl2 and 1µL of Ribolock (Thermo Fisher Scientific) to the 1 mL

lysate and incubating for 20 min at 37°C. Lysates were centrifuged for 10 min at 4°C.

Supernatants were collected and pre-cleared for 1h at room temperature with Dynabeads

protein G magnetics beads (Invitrogen) blocked with yeast tRNA (Invitrogen). The efficiency

of the RNase treatment was evaluated by RNA analysis of the input samples on a 1% agarose

gel.

Lysates were incubated over night at 4°C with 40 µL of Dynabeads protein G beads

conjugated with 2 µg rabbit anti-DDX5 antibody (ab21696; Abcam), or rabbit anti-DDX17

antibody (19910-1-AP, Proteintech) or 2µg rabbit IgG antibody (2729; Cell Signaling

Technology).

Beads were washed 3 times with RIP buffer containing 150mM NaCl, twice with the same

buffer supplemented with 50mM of NaCl, and 3 times 3 times with RIP buffer containing

150mM NaCl. 30% of the immunoprecipitated protein and 70% of the associated RNA were

eluted with Laemmli 1X at 95°C for 10 min or with 1 mL TriReagent (Thermo Fisher

Scientific), respectively. Proteins were analysed by western blot or by mass spectrometry,

while RNA was extracted and analysed by RT-qPCR.

2.10.   Western Blot

Cells were collected in 300 to 500 µL of lysis buffer (50mM tris-HCL [pH 7.5], 150mM

NaCl, 5 mM EDTA, 0.05% SDS, 1% triton, 1 tablet of commercial protease inhibitor cocktail

(ROCHE)) and incubated for 30 min on ice. Cell lysates were then collected and protein

concentration was determined using the Bradford method (Bio-Rad). For total protein

analysis, 20µg of protein samples were heated in 1X Laemmli buffer at 95°C for 5 min and

loaded on pre-cast 4-20% SDS-polyacrylamide electrophoresis gel (Bio-Rad). For the RIP

experiment, 0.5 % of input and 12,5% of immunoprecipitated protein were used. After

migration, the proteins were transferred onto 0.45 µm nitrocellulose membranes (GE

healthcare). The membranes were blocked with 5% non-fat milk diluted in PBS 1X

(Euromedex) complemented with 0.2% Tween-20 (PBS-T) for 1 h at room temperature and

then were incubated at 4°C overnight or 1 h at room temperature with the following specific

primary antibodies: anti- DDX5 (mouse- horseradish peroxidase(HRP), sc-365164; Santa

Cruz Biotechnology) or anti-DDX5 (rabbit, ab21696; Abcam), anti-DDX17 (mouse, sc-

271112; Santa Cruz Biotechnology), anti-tubulin HRP (ab21058; Abcam), anti-GAPDH-HRP

(mouse, ab9482; Abcam), anti-capsid (rabbit, kind gift of Diane Griffin), anti-GFP (mouse,

11814460001, Roche). Membranes were washed 3 times with PBS-T for 5 min and incubated

with mouse-HRP (A4416; Merck) and rabbit-HRP (GENA9640V, Merck) secondary

antibodies for 1 h at room temperature. Proteins were detected using the Chemiluminescence

substrate (Supersignal West Pico; Pierce) and analysed with the Fusion FX imaging system

(Vilber).

2.11.   RNA extraction and RT-qPCR

Total RNA and RNA enriched upon specific protein immunoprecipitation (RIP) were

extracted using TriReagent (Invitrogen, Thermo Fisher Scientific) according to the

manufacturer's instructions. For total RNA analysis, 1 µg of RNA was reverse transcribed

with the SuperScript VILO Master Mix (Invitrogen, Thermo Fisher Scientific) according to

the manufacturer's instructions. For the RIP experiment, 1 µL of samples before and after IP

was used for retro-transcription using the same protocol. RT-qPCR was performed on cDNA

using the Maxima SYBR Green qPCR Master Mix K0253; Thermo Fisher Scientific) on the

CFX96 touch Real-Time PCR machine (Bio-Rad) using the following primers at an annealing

temperature of 60°C:

SINV genome FW: `

5$^{'}$-CCACTACGCAAGCAGAGACG-3$^{'}$;

SINV genome RV:

5$^{'}$-AGTGCCCAGGGCCTGTGTCCG-3$^{'}$;

GAPDH FW:

5$^{'}$-CTTTGGTATCGTGGAAGGACT-3$^{'}$;

GAPDH RV:

5$^{'}$-CCAGTGAGCTTCCCGTTCAG-3'

SINV sub-genome-genome FW

5'CCACAGATACCGTATAAGGCA 3'

SINV sub-genome-genome RV

5' TGCAGGTAATGTACTCTTGG 3'

2.12.   Immunostaining

Mock or SINV WT- infected cells were plated on 8 wells LabTek slide (Merck Millipore),

were fixed with 4% formaldehyde (Merck) diluted in PBS 1X for 10 min at room temperature

and then washed 3 times with PBS 1X. Cells were blocked in blocking buffer (0.1% Triton X-

100; PBS 1 X; 5% normal goat serum) for 1 h. The following primary antibodies were diluted

1:400 in blocking buffer and incubated over night at 4°C: mouse anti-dsRNA J2 (RNT-SCI-

10010200; Jena bioscience), mouse anti- DDX5 (67025 Proteintech) or rabbit anti-DDX5

(ab21696; Abcam), anti-DDX17 (mouse, sc-27112; Santa Cruz Biotechnology), anti-capsid

(rabbit, kind gift of Diane Griffin). Cells were washed with PBS 1X-Triton 0.1%. and

incubated for 1 h at room temperature with secondary antibodies (goat anti-mouse Alexa 594

(A11032, Invitrogen) or goat anti-rabbit Alexa 488 (A11008, Invitrogen) secondary

antibodies) diluted at 1:1000 in PBS 1X-Tween 0.1%. DAPI staining was performed for 5

min in PBS 1X to reveal the nuclei (D1306, Invitrogen, Thermo Fisher Scientific). Slides

were mounted on coverslips with Fluoromount-G mounting media (Invitrogen, Thermo Fisher

Scientific) and observed by confocal microscopy (LSM780, Zeiss) with a 40X or 63X

objective. Images were analysed using Image J software and fluorescence intensity profiles

were obtained.

2.13. Sequential immunostaining and FISH

Mock or SINV-WT infected (MOI 1, 24hpi) HCT116 cells were grown on 18 mm round cover glass in 12-well cell culture plates.

Cells were fixed with 3.7 % formaldehyde diluted in PBS 1X for 10 min at room temperature.

Cells were then permeabilized in 1mL of 0.1% Triton X-100 in 1X PBS for 5 min at room temperature and incubated with anti-DDX5 primary antibody (rabbit, ab21696; Abcam) diluted to 1:400 in PBS 1X for 1 hour. Goat anti-rabbit Alexa 488 (A11008, Invitrogen) secondary antibody diluted to 1:1000 was added on the cells for 1 hour at room temperature.

Cells were fixed again with 3.7 % formaldehyde (Biosearch technologies) diluted in PBS 1X for10 min at room temperature and incubated over night at room temperature with the SINV genome specific LGC Biosearch Technologies' Stellaris® RNA FISH Probe diluted in RNA FISH hybridization buffer (Stellaris, Biosearch technologies). DAPI staining was performed for 30 min to reveal the nuclei (D1306, Invitrogen, Thermo Fisher Scientific). Slides were mounted on coverslips with Fluoromount-G mounting anti-fading media (Invitrogen, Thermo Fisher Scientific) and observed by confocal microscopy (LSM780, Zeiss). Images were analysed using Image J software.

2.14. Mass spectrometry analyses

For LC-MS/MS analyses, proteins were prepared as described in a previous study (26). Proteins eluted from the beads were washed with 2 sequential overnight precipitations with glacial 0.1 M ammonium acetate in 100% methanol (5 volumes) followed by 3 washes with glacial 0.1 M ammonium acetate in 80% methanol. Proteins were then solubilized in 50 mM ammonium bicarbonate for a reduction-alkylation step (dithiothreitol 5 mM – iodoacetamide 10 mM) and an overnight digestion with 300ng of sequencing-grade porcine trypsin (Promega, Fitchburg, MA, USA). Digested peptides were resuspended in 0.1% formic acid

(solvent A) and injected on an Easy-nanoLC-1000 system coupled to a Q-Exactive Plus mass

spectrometer (Thermo Fisher Scientific, Germany). One fourth of each sample was loaded on

a C-18 precolumn (75 µm ID × 20 mm nanoViper, 3µm Acclaim PepMap; Thermo-Fisher

Scientific) and separated on an analytical C18 analytical column (75 µm ID × 25 cm

nanoViper, 3µm Acclaim PepMap) with a 160 minutes gradient of solvent B (0.1% of formic

acid in acetonitrile).

Qexactive data were searched with Mascot algorithm (version 2.6.2, Matrix Science) against

the Swissprot database with *H.sapiens* taxonomy (release 2020_05) as well as the sequences

of GFP and of SINV virus (20 394 sequences) with a decoy strategy. The resulting .dat

Mascot files were then imported into Proline v2.0 software (32) to align the identified

proteins. Proteins were then validated on Mascot pretty rank equal to 1, 1% FDR on both

peptide spectrum matches (PSM) and protein sets (based on Mascot score).

For statistical analyses of the mass spectrometry data, spectral counts (SpC) of the identified

proteins were stored in a local MongoDB database and subsequently analysed through a Shiny

Application built upon the R packages msmsEDA (Gregori J, Sanchez A, Villanueva J

(2014). msmsEDA: Exploratory Data Analysis of LC-MS/MS data by spectral counts.

R/Bioconductor package version 1.22.0) and msmsTests (Gregori J, Sanchez A, Villanueva J

(2013). msmsTests: LC-MS/MS Differential Expression Tests. R/Bioconductor package

version 1.22.0). Differential expression tests were performed using a negative binomial

regression model. The P-values were adjusted with FDR control by the Benjamini–Hochberg

method and the following criteria were used to define differentially expressed proteins: an

adjusted P-value <0.05, a minimum of 5 SpC in the most abundant condition, and a minimum

absolute Log2 fold change of 1.

2.15.   **Statistical analysis**

Statistical analyses were performed using GraphPad Prism 9 software. Generally, a paired

Student's t test was performed for comparisons between two samples for plaque assays and a

one-sample t test was performed to determine the significance of the fold change on qPCR

results in which the control sample was set to 1. The number of replicates per experiment and

the specific statistical tests used are stated in the figure legends.

3. Results

3.1. DDX5 interacts with SINV viral RNA in the cytoplasm of SINV -infected cells.

It was previously reported that DDX5 is primarily localized in the nucleus (8), while the SINV replication cycle is strictly cytoplasmic (28). To determine whether DDX5 can affect SINV infection *via* a direct interaction with the viral components, we first verified whether DDX5 co-localizes with SINV RNA in infected cells. Despite the predominantly nuclear localization in uninfected cells, confocal microscopy analyses showed that DDX5 re-localizes to the cytoplasm in the presence of the virus and co-localizes with the viral genomic RNA at 24 hours post infection (**Figure 1A**), suggesting that DDX5 is recruited to the viral RNA. We subsequently performed RNA immunoprecipitation (RIP) on the endogenous DDX5 in mock and SINV GFP-infected HCT116 cells. Efficient DDX5 immunoprecipitation was assessed at the protein level by western blot (**Figure 1B)** and the associated RNA was analysed by RT-qPCR**.** We found that the viral genomic RNA is specifically enriched in the DDX5-RIP compared to the control (IgG-RIP) condition (**Figure 1C**). We also performed dot blot analyses using a J2 antibody, which detects dsRNA longer than 40 bp (33,34), and observed an enrichment of dsRNAs in the DDX5-IP compared to the IgG-IP in the infected samples (**Figure 1D**). Of particular note, we also observed enrichment of dsRNA in the mock condition, albeit to a lesser degree than upon SINV infection. This suggests that DDX5 binds to endogenous dsRNA molecules even in the absence of viral RNA. Whilst dsRNA can be found associated with DDX5 by dot blot analyses, confocal co-immunostaining analyses using the J2 antibody as a proxy for dsRNA do not demonstrate a clear co-localization of DDX5 and dsRNA upon SINV-infection (**Figure 1E**), but rather indicate a proximity between the protein and dsRNA.

Overall, these results suggest that DDX5 can bind to viral RNAs in the cytoplasm of SINV

infected cells and localizes in close proximity to cytoplasmic viral dsRNA structures.

3.2. DDX5 interacts with host and viral proteins in SINV-infected conditions.

Once we determined that DDX5 could be recruited to the viral RNA, we aimed to test

whether any interaction between DDX5 and the viral proteins could also occur to modulate

this association upon SINV infection. We immunoprecipitated DDX5 from mock and SINV-

infected HCT116 cells at MOI 0.1 for 24 hours and eluates were analysed by liquid

chromatography coupled to tandem mass spectrometry (LC-MS/MS). An IgG antibody was

used as a negative control. Significantly enriched proteins were determined as having an

FDR<1%, at least 5 spectral counts in the most abundant condition, a fold change of at least 2

and an adjusted p-value lower than 0.05.

Differential expression analysis revealed a total of 206 human proteins enriched in uninfected

DDX5-IP over IgG-IP conditions (**Figure 2A** and **Table S1**), while 177 proteins were

enriched in DDX5 IP over IgG IP control in SINV-infected samples (**Figure 2B and Table**

S2).

Comparison of the DDX5 enrichment profiles in mock and in SINV conditions showed that

156 human proteins overlapped in the mock and SINV DDX5 interactomes, suggesting an

overall stable DDX5 interactome core before and after infection (**Figure 2C**). Among them,

we identified the polyribonucleotide nucleotidyltransferase 1 (PNPT1), an enzyme

predominantly localized in the mitochondrial intermembrane space which acts as a 3'-to-5'

exoribonuclease and regulates mitochondrial RNA metabolism thereby modulating innate

immunity (35). Even though PNPT1 and DDX5 were both enriched in our previous DRIMS

experiments upon SINV infection (26), their close interaction has not been reported so far,

Several RNA helicases such as DHX9, DHX30, DHX15, DDX3X, DDX21, DDX24, DDX54,

DDX18, DDX10, DDX52 and DDX56, were also found in the DDX5 interactome. In

particular, one of the most enriched human proteins in both mock and SINV-infected DDX5

IP was its well-known interactor DDX17 (10).

The viral capsid and nsP2 proteins were the most significantly enriched proteins in DDX5 IP

SINV over DDX5 IP mock (**Figure 2D** and **Table S3**) and the only viral proteins significantly

identified in DDX5 IP over IgG IP upon SINV infection (**Figure 2B and Table S2).** We thus

decided to assess whether these association between DDX5 and these SINV proteins was

specific and whether it relied on the binding to the viral RNA.

3.3. DDX5 interaction with the viral capsid is RNA-independent.

To further characterise the interaction between DDX5 and the viral proteins, we chose to test

all four SINV non-structural proteins (nsP1, nsP2, nsP3, nsP4) required for viral RNA

synthesis and the structural capsid protein (CP) which associates with the viral genomic RNA

for its packaging (**Figure 3A**). We individually cloned each of their coding sequence in order

to express them with an N-terminal Flag-HA tag and verified their expression upon transient

transfection in HCT116 cells by western blot analysis using an anti-Flag antibody (**Figure 3B,**

upper panel). We next investigated whether the endogenous DDX5 co-immunoprecipitated

with any of the overexpressed viral proteins in uninfected HCT116 cells in the absence of

viral RNA or any other viral protein. While we did not confirm the interaction between nsP2

and DDX5 in the absence of viral RNA, we observed that the capsid protein associated with

DDX5 independently of the viral RNA (**Figure 3B, lower panel**).

We further validated the interaction of DDX5 and the viral capsid by co-immunoprecipitation

experiments upon SINV infection (**Figure 3C, left panels**). Reciprocal co-

immunoprecipitation of DDX17 confirmed its interaction with DDX5 and the viral capsid

(**Figure 3C, right panels**), suggesting that the three proteins may be in close proximity in

infected conditions. Of note, the interaction between DDX5 and capsid as well as the

association with DDX17 are mostly maintained upon RNase A treatment. (**Figure 3D-E**).

We then performed co-immunostaining analyses to verify whether DDX5 co-localizes with

the viral capsid in SINV-infected HCT116 cells. Confocal microscopy analysis demonstrated

that endogenous DDX5 co-localizes with the capsid protein in SINV-infected cells supporting

the possibility that the two proteins interact upon infection (**Figure 3F**).

Interestingly, while DDX5 and its cofactor DDX17 colocalize in the nucleus in uninfected

conditions, both proteins re-localize to the cytoplasm in discrete perinuclear foci upon SINV

infection (**Figure 3G**) most likely in proximity of viral replication sites (see **Figure 1A**).

Overall, these results confirm the interaction between the viral capsid and DDX5 in infected

cells and indicate that DDX17 interaction with DDX5 and capsid can occur in the cytoplasm

of the infected cells.

3.4. DDX5 depletion reduces SINV infection in HCT116 cells.

To formally determine the impact of DDX5 on SINV replication and infection, we first

knocked it down by siRNA treatment in HCT116 cells, which were then infected for 24h with

SINV-GFP at an MOI of 0.1. We observed a reduction in viral capsid protein level by western

blot upon DDX5 depletion (siDDX5) compared to the control condition (siCTRL) (**Figure

4A**). In addition, we found that the viral genomic (g)RNA level (**Figure 4B**) and the viral titer

(**Figure 4C**) were also reduced in siDDX5 compared to siCTRL-treated cells.

To further establish the negative effect of DDX5 depletion on SINV infection, we generated

homozygous *DDX5* knockout (KO) HCT116 cells by CRISPR/Cas9 (**Figure 4D**). The *DDX5*

KO clone was characterized at the genomic DNA level (**Figure 4E**) and the absence of DDX5

protein was verified by western blot (**Figure 4F**).

Upon SINV-GFP infection for 24h at an MOI of 0.1 we observed a strong reduction in viral

capsid protein in *DDX5* KO HCT116 compared to WT HCT116 cells by western blot (**Figure**

4F). The viral genomic RNA level (**Figure 4G**) and the viral titer (**Figure 4H-I**) were also

significantly reduced in the absence of DDX5.

We then measured the relative fluorescence of GFP to quantify the impact of *DDX5* knockout

during a time course of SINV infection using a CellcyteX automated cell counter and

analyser. A robust decrease in GFP fluorescence signal was observed in SINV-GFP infected

DDX5 KO compared to WT HCT116 cells at three MOIs tested (MOI of 0.01, 0.1 and 1) over

72 hours (**Figure 4J**). These results indicate that DDX5 depletion strongly reduces the

infection over time and independently on the MOI used already at the earliest step of the viral

replication cycle. Taken together our results point to a negative impact of DDX5 depletion on

SINV infection in our experimental conditions suggesting that DDX5 is a pro-viral host factor

for SINV infection.

3.5. DDX5 re-expression increases capsid levels in SINV-infected HCT116 cells.

To better assess the pro-viral function of DDX5 on SINV, we stably re-expressed a V5-DDX5

or BFP protein (control) in *DDX5* KO HCT116 cells and verified that a partial DDX5

expression could be restored in our experimental conditions compare to WT HCT116 cells

(**Figure 5A**). We then analysed the effect of the ectopic DDX5 expression on the virus at

24hpi at an MOI of 0.1 and observed an increase in viral capsid protein level by western blot

(**Figure 5B**) in V5-DDX5 cells compared to the control BFP one infected with SINV-GFP. In

contrast, we found that the viral genomic (g)RNA, sub-genomic (sg)RNA level (**Figure 5C-**

D) and the viral titer (**Figure 5E**) were not significantly affected in V5-DDX5 compared to

BFP expressing HCT116 KO cells at this specific time point.

Finally, while the effect was not as pronounced as in WT cells, the quantification of the

relative GFP fluorescence during a time course of SINV-GFP infection (MOI 0.1) revealed a

partial rescue of the GFP signal in the V5-DDX5 cells compared to the BFP and the *DDX5*

KO ones over time (**Figure 5F**).

Taken together, these observations suggest that the ectopic DDX5 expression has a positive

impact on SINV-GFP infection indicating an overall pro-viral effect on the virus. However,

the impact of DDX5 overexpression on the virus at 24hpi is mostly observed as an increase of

the viral capsid protein level and does not induce a significant increase on viral RNA and

infectious particle production, although we could observe a tendency for an increase in the

presence of DDX5. Nonetheless, the kinetic of GFP fluorescence measurement clearly

indicates that the phenotype of the DDX5 knock-out can be complemented by its re-

expression.

3.6. Combined DDX5 and DDX17 depletion has a synergic detrimental impact on

SINV.

Once established that DDX5 depletion has a negative impact on SINV and that its re-

expression in a *DDX5* KO background has a positive impact on the virus, we set to determine

whether DDX17 plays a role together with or independently of DDX5 during SINV infection.

We first knocked down DDX17, then infected HCT116 cells with SINV-GFP. We validated

the knockdown efficiency by western blot compared to an siRNA control and observed that

viral capsid levels were reduced upon siDDX17 treatment (**Figure 6A**). In addition, SINV

genomic and sub-genomic RNA levels (**Figure 6B-C**) as well as SINV viral particle

production (**Figure 6D**) were also significantly reduced in DDX17 knockdown conditions

compared to siCTRL. These results suggested that just like DDX5, DDX17 has a positive

effect on SINV infection.

We proceeded to investigate whether DDX17 acts in synergy or competition with DDX5 by

analysing the impact of DDX17 knockdown in *DDX5* KO HCT116 cells where viral infection

is already weakened (see **Figure 4D-J**). As previously reported (13), the protein levels of both

DDX7 isoforms (p72 and p82) increased in *DDX5* KO HCT116 compared to WT HCT116

cells and such an effect was independent on the infection (**Figure 6E**). However, upon SINV-

GFP infection, we observed a further reduction of the viral capsid expression in siDDX17

treated *DDX5* KO HCT116 cells compared to the control condition by western blot (**Figure**

6F). SINV genomic and sub-genomic RNA levels (**Figure 6G-H**), as well as SINV viral

particle production (**Figure 6I**) were significantly diminished in DDX17 knockdown

conditions compared to siCTRL in *DDX5* KO HCT116 cells. Overall, these results indicate

that DDX5 and DDX17 may have independent pro-viral functions during SINV infection

because their individual depletion causes reduction of viral infection at the protein, RNA and

viral production levels and their joint depletion has a cumulative negative effect on SINV

infection.

4. Discussion

The human DEAD-box RNA helicase DDX5 plays a pivotal role in virus–host interaction being mostly pro-viral for several viral families. In this study, we investigated and extended the function of DDX5 as a cellular cofactor of SINV replication in HCT116 cells. Despite being an uncommon model to study alphavirus infection, the HCT116 cell line was chosen because it can be easily infected with SINV and can be used for CRISPR-Cas9 gene editing procedures (36).

We found that DDX5 interacts with viral RNA at the cytoplasmic viral replication sites, which may be important for its pro-viral activity. RNA helicases recognize the RNA structure rather than their sequence and they have shown little substrate specificity *in vitro* (1). This lack of specificity is counterbalanced by their association with specific interactors *in vivo*. For instance, DDX5 interaction with either viral non-structural or structural proteins is essential for its pro-viral role across different RNA virus infections (14).

Our proteomic analysis revealed the interaction between DDX5 and SINV capsid, as well as with the nsP2 protein. The cytoplasmic SINV capsid protein is essential for genomic RNA packaging and the first steps of viral morphogenesis. In contrast, SINV nsP2 is the protease and helicase of the viral replication complex and it is the only viral protein known to re-localise to the nucleus to inhibit host transcription (37). The interaction between DDX5 and nsP2 may thus occur both in the nucleus and in the cytoplasm of the infected cell. The association between the viral capsid and DDX5 was further supported by co-immunoprecipitation experiments and imaging analyses. This interaction is RNA-independent since it occurs both in uninfected cells expressing a tagged version of the capsid protein and in the presence of an RNase treatment in SINV infected cells.

We showed that DDX5 depletion has a detrimental effect on viral replication and production in infected cells. Furthermore, we established a stable cell line that ectopically expressed

DDX5 in a *DDX5* KO background to assess its functional impact on the virus. Despite its

modest overexpression, the presence of DDX5 was sufficient to partially rescue the GFP

levels in a time course of SINV-GFP infection. Although we did not observe a significant

increase of viral gRNA, sgRNA or titers, we noted an enhanced accumulation of the capsid

protein in DDX5 KO cells upon re-expression of a tagged DDX5. This partial effect might

depend on the relatively low expression levels of the V5-DDX5 or on the choice of the

inappropriate time point of infection.

Indeed, we observed a partial rescue of fluorescence in cells expressing V5-DDX5 compared

to the controls when we monitored GFP expression from the virus in real time over 72 hours.

Taken together, these results support the pro-viral role of DDX5 on SINV.

The DDX5 interactome analysis also revealed associations with a number of cellular RNA

binding proteins that could have a function during SINV infection. These include several

RNA helicases of the DEAD/H family and other interesting factors such as PNPT1.

Interestingly, we previously showed that DDX5, DDX17 and PNPT1 are an integral part of

the SINV dsRNA associated proteome (26). Further investigations are needed to better

characterize if the association of those novel DDX5 interactors have functional consequences

on SINV infection. In this study we focused on DDX17 which was previously known to

associate with DDX5.

We demonstrated by co-immunoprecipitation experiments that DDX17 interacts with DDX5

and the viral capsid in infected cells. It is now established that the replication of positive

strand RNA viruses in the cytoplasm can induce re-localization of nuclear host proteins and

promote infection (38). Of note, we showed that DDX5 re-localizes from the nucleus to the

cytoplasm with DDX17 upon SINV infection where they can both interact with the viral

capsid protein. It is thus conceivable that the cofactor DDX17 operates with DDX5 within the

same complex, carrying out a synergistic or additive function during infection. A previous

641 study showed that Rm62, the fly orthologue of DDX17 and DDX5, did not affect SINV

642 infection in *D. melanogaster*, and that DDX17 knockdown showed no antiviral effect on the

643 accumulation of SINV capsid protein in human U2OS cells (16). In contrast, we demonstrated

644 that the single depletion of DDX17 reduces SINV infection in HCT116 cells and its combined

645 depletion with DDX5 in the KO background further decreases viral levels, suggesting a

646 synergism between the two proteins in promoting viral infection.

647 These experiments also ruled out the possibility that the observed reduction in SINV

648 replication upon DDX5 depletion was due to an antiviral effect of compensatory DDX17

649 overexpression.

650 As both DDX5 and DDX17 inhibition reduces virus levels, further investigation is required to

651 better characterize these factors as potential targets for antiviral therapies. It will also be

652 important to expand our functional observations to other cellular models more relevant for

653 alphavirus infection in the future.

654 Additional analyses will be needed to formally assess whether DDX5 direct action supports

655 viral replication and if its binding to viral RNA and helicase activity is necessary for its

656 function. Being involved in many aspects of cellular RNA biology, one cannot formally

657 exclude the contribution of DDX5 indirect effects on infection. For instance, DDX5 activity

658 as a regulator of N6-methyladenosine levels on the DHX58 and NF-κB mRNAs could

659 negatively impact antiviral innate immunity and therefore enhance viral infection (39).

660 Finally, it could be of interest to determine whether the helicase activity of DDX5 is required

661 for its pro-viral effect on SINV production and which protein domains are important for the

662 interaction with the capsid viral protein.

5. Conclusion

In conclusion, this study uncovers DDX5 as a novel positive host factor for the prototype alphavirus SINV and significantly advances our understanding of the role played by RNA helicases in viral infections. Our results demonstrate the interaction of DDX5 with viral RNA, suggesting a potential impact on early stages of viral replication. We discovered a re-localization of DDX5 together with its partner DDX17 to the cytoplasm upon SINV infection and their association with the viral capsid protein. Depletion of both DDX5 and DDX17 strongly impacts SINV infection. Of note, the detrimental effect caused by DDX5 depletion on the virus is partially offset by DDX5 ectopic expression, providing evidence for the pro-viral function of DDX5 on SINV.

Figure 1: DDX5 interacts with the viral RNA. A) Confocal microscopy analysis of SINV

(+) RNA and DDX5 protein localization in mock and infected HCT116 cells at 24 hpi by

RNA fluorescence in situ hybridization (FISH) (in magenta) combined with protein

immunostaining (in green). DAPI staining (in blue) and merge of the different channels are

shown. Scale bar, 10µm. Representative fluorescence intensity profiles of DDX5 (green

curve) and SINV RNA (magenta curve) along the yellow line represented on the merge panel

and normalized to the maximum value are shown on the right. **B**) Anti DDX5 western blot on

total lysate (INPUT), IgG-IP or DDX5-IP samples in both mock and SINV-GFP infected

HCT116 cells. **C**) RT-qPCR on SINV genomic (g) RNA upon DDX5 RIP or IgG RIP.

Results are expressed as percentage of Input (total RNA) and represent the mean ± standard

deviation (SD) of three biological replicates (n = 3). **D**) Anti-dsRNA dot blot assay on serial

dilutions of the total RNA (INPUT) and the undiluted RNA samples from DDX5-RIP or IgG-

RIP, in mock and SINV infected conditions. J2 antibody was used to detect dsRNAs. **E**)

Confocal co-immunofluorescence analysis was performed in mock and SINV infected

HCT116 cells using anti-DDX5 rabbit antibody (in green) or anti-J2 mouse antibody (in

magenta). Nuclei were stained with DAPI (in blue). Scale bar 10µm. Representative

fluorescence intensity profiles of DDX5 (green curve) and J2 (magenta curve) along the

yellow line represented on the merge panel and normalized to the maximum value are shown

on the right.

Figure 2: DDX5 interactome in mock and SINV-infected cells. A, B and D) Volcano plots

showing the global protein enrichment in (**A**) mock-DDX5-IP versus mock-IgG-IP, (**B**)

SINV-DDX5-IP versus mock-DDX5-IP and (**D**) SINV-DDX5-IP versus SINV-IgG-IP using

data from three independent replicates. The cells were infected with SINV-GFP during for

24h at an MOI 0.1. Significantly enriched or depleted proteins are represented in red or blue,

respectively. Cut off values: FDR<1%, at least 5 spectral counts in the most abundant

condition, fold change > 2, adjusted p-value lower than 0.05. Viral proteins and selected host

proteins are circled in black and orange, respectively. **C**) Venn diagram showing the number

of overlapping and unique significantly enriched proteins identified in DDX5 IP compared to

IgG in mock and SINV infected proteomic analyses.

Figure 3: DDX5 interacts with the SINV capsid protein. A) Schematic illustration of SINV

genomic organization. The 5´ cap and poly(A) tail are depicted. The non-structural proteins

(np1, nsp2, nsp3, nsp4) from the first ORF (in green) and the structural proteins (capsid, E3,

E2, 6K, E1) from the second ORF (in yellow and grey) are shown. The sequences coding for

nsp1, nsP2, nsP3, nsP4 and capsid proteins were individually cloned and fusd with an N-

terminal FLAG-HA tag. **B**) Western blot on total lysate (INP) or anti-HA IP (IP) from

transiently transfected HCT116 cells using anti-FLAG and anti-DDX5 antibodies. Position

and molecular weight (KDa) of protein size markers are shown on the left. **C**) Western blot on

total lysate (INPUT), IgG-IP (IgG), DDX5-IP (DDX5) or DDX17-IP (DDX17) samples in

both mock and SINV-GFP infected lysates, using antibodies against DDX5, DDX17 and viral

capsid. **D)** Representative agarose gel showing the total RNA before and after RNase

treatment. **E)** Western blot on total lysate (INPUT), IgG-IP (IgG) or DDX5-IP (DDX5)

samples in both mock and SINV-GFP infected lysates, treated or not with RNase A, using

antibodies against DDX5, DDX17 and viral capsid. **F-G**) Confocal co-immunofluorescence

analysis performed in mock and SINV- infected HCT116 cells using specific primary

antibodies. Rabbit anti-capsid (in green) and mouse anti-DDX5 antibody (in magenta) are

shown in (**F**). Rabbit anti-DDX5(in green) and mouse anti-DDX17 antibodies (in magenta)

are shown in (**G**). DAPI was used to stain the nuclei (in blue). Scale bar 10µm. Representative

fluorescence intensity profiles of capsid (green curve) and DDX5 (magenta curve) **(F)** or

DDX5 (green curve) and DDX17 (magenta curve) **G**) along the yellow line represented on the

merge panel and normalized to the maximum value are shown on the right.

Figure 4: SINV infection is reduced in DDX5-depleted HCT116 cells. A) Western blot on

HCT116 mock or SINV-GFP infected cells upon siCTRL and siDDX5 transfection. DDX5

and viral capsid were detected by specific primary antibodies. GAPDH was used as loading

control. Position and molecular weight (KDa) of protein size markers are shown on the left.

B) RT-qPCR of SINV genomic (g) RNA in SINV-GFP infected HCT116 cells (MOI 0.1, 24

hpi), upon siDDX5 and siCTRL transfection. Results are expressed as relative to GAPDH and

normalized to the siCTRL condition. **C**) SINV-GFP viral titers from infected HCT116 cells

transfected with siCTRL or siDDX5 was quantified by plaque assay. Results in **B-C** represent

the mean ± standard deviation (SD) of three biological replicates (n = 3). * p ≤ 0.05, (**B**) one-

sample t test or (**C**) paired Student's t test. **D**) Schematic diagram of the *DDX5* CRIPR/Cas9

KO strategy. Two gRNAs targeting *DDX5* exon 2 and intron 2 respectively were used to

delete the 5' splice site (ss) region (5'ss is shown in green, #1 target region in blue and #2

target region in magenta). The purple arrow represents an indel in *DDX5* exon 2 for the KO

allele 1. The grey rectangle represents a 378 nucleotide (nt) deletion in KO allele 2. **E**)

Agarose gel showing the PCR fragments amplified from cellular genomic DNA,

corresponding to the WT and KO alleles, respectively, which were gel purified and

sequenced. Amplicon size is shown: the upper band corresponds to KO allele 1, the lower

band corresponds to the KO allele 2. **F)** Western blot on mock or SINV-GFP infected cells in

WT or *DDX5* KO HCT116 cells. Antibodies against DDX5 and viral capsid were used.

GAPDH was used as loading normalizer. **G**) Level of SINV genomic RNA relative to

GAPDH monitored by RT-qPCR in WT or *DDX5* KO HCT116 cells upon SINV-GFP

infection at MOI 0.1 for 24 hours. Results are plotted as fold change relative to WT cells. **H**)

Viral particles of SINV-GFP from the supernatant of SINV-GFP infected WT or *DDX5* KO

HCT116 cells were quantified by plaque assay on Vero cells by plaque assay. The results

from three independent experiments are presented here. **I**) Representative plaque assay

images after crystal violet staining of Vero E6 cells infected with SINV-GFP supernatant at

the indicates dilutions (10^{-3} and 10^{-4}) from infected WT or *DDX5* KO HCT116 cells. **J**)

SINV-GFP infection kinetics in WT and *DDX5* KO HCT116 cells. The relative GFP

fluorescence area (expressed in percentage) of WT and *DDX5* KO HCT116 cells as a function

of time was measured after SINV GFP infection at an MOI of 0.01 (left panel), 0.1 (middle

panel) and 1 (right panel) every 6 hours for 72 hours with a CellcyteX automated cell counter

and analyser. WT HCT116 cells, black; *DDX5* KO HCT116 cells, red. Error bars in **G-H** and

I represent the mean ± standard deviation (SD) of three independent biological experiments. *

$p \leq 0.05$, **$p \leq 0.01$, (**G**) one- sample t test or (**H**) paired Student's t test.

Figure 5: Effect of DDX5 re-expression on SINV in *DDX5* KO HCT116 cells. A) Western

blot on WT, *DDX5* KO HCT116 and *DDX5* KO HCT116 stably expressing either V5-DDX5

(V5-DDX5) or BFP (BFP). Antibodies directed against DDX5 and GAPDH antibody were

used. **B)** Western blot on SINV-GFP infected *DDX5* KO HCT116 overexpressing V5-DDX5

or BFP (24 hpi, MOI 0.1). Antibodies directed against DDX5 and capsid were used. Tubulin

antibody was used as loading control. (**C-D**) RT-qPCR of SINV genomic (g) RNA (**C**) and

SINV sub-genomic (sg) RNA (**D**) in SINV-GFP infected *DDX5* KO HCT116 stably

expressing V5-DDX5 or BFP (MOI 0.1, 24 hpi). Results are expressed as relative to GAPDH

and normalized over the siCTRL condition. **E**) SINV-GFP viral titers from SINV-GFP

infected *DDX5* KO HCT116 stably expressing V5-DDX5 or BFP (MOI 0.1, 24 hpi),

quantified by plaque assay. Results in (**C-D-E**) represent the mean ± standard deviation of

three biological replicates (n = 3). ns = non significant, (**C-D**) one sample t test or (**E**) paired

Student's t test. **F)** SINV-GFP infection kinetics in WT, *DDX5* KO HCT116 and *DDX5* KO

HCT116 stably expressing either V5-DDX5 (V5-DDX5) or BFP (BFP). The relative GFP

fluorescence area (expressed in percentage) of the infected cells as a function of time was

measured after SINV GFP infection at an MOI of 0.1, every 3 hours for 72 hours with a

CellcyteX automated cell counter and analyser. WT HCT116 cells, black; *DDX5* KO HCT116

cells, red; V5-DDX5 cells, purple; BFP cells, blue.

Figure 6: Effect of DDX17 depletion on SINV in WT and *DDX5* KO HCT116 cells. A)

Western blot on SINV-GFP infected HCT116 cells, treated with siCTRL and siDDX17.

Antibodies directed against DDX5, DDX17 and capsid were used. Tubulin antibody was used

as loading control. (**B- C**) RT-qPCR of SINV genomic (g) RNA (**B**) and SINV sub-genomic

(sg) RNA (**C**) in siRNA treated SINV-GFP infected HCT116 cells (MOI 0.1, 24 hpi). Results

are expressed as relative to GAPDH and normalized over the siCTRL condition. **D)** SINV-

GFP viral titers from infected HCT116 cells transfected with the indicated siRNAs, quantified

by plaque assay. **E)** Western blot on mock SINV-GFP infected WT and *DDX5* KO HCT116

cells. Antibodies directed against DDX5, DDX17 and capsid were used. Tubulin antibody

was used as loading control. **F)** Western blot of samples from SINV-GFP infected *DDX5* KO

HCT116 cells upon siCTRL or siDDX17 transfection. Specific primary antibodies against

DDX5, DDX17 and viral capsid were used. Tubulin was used as loading control. Position and

molecular weight (KDa) of protein size markers are shown on the left. **G-H)** RT-qPCR of

SINV genomic (g) RNA (**G**) and SINV sub-genomic (sg) RNA (**H**) in SINV-GFP infected

DDX5 KO HCT116 cells (MOI 0.1, 24 hpi), upon siRNA transfection. Results are expressed

as relative to GAPDH and normalized to the siCTRL condition. **I)** SINV-GFP viral titers from

infected *DDX5* KO HCT116 cells transfected with the indicated siRNAs, quantified by plaque

assay. Results in (**B-C- D** and **G-H-I)** represent the mean ± standard deviation of three

biological replicates (n = 3). * p ≤ 0.05, **p ≤ 0.01, (**B-C** and **G-H**) one sample t test or (**D**

and **I**) paired Student's t test.

Funding

This work received financial support from the Agence Nationale de la Recherche (EndoDRAI project ANR-22-CE15-0011-01 (to E.G.) and was supported by the Idex 2022 Attractivité grant under the framework of the IdeX University of Strasbourg (to E.G.) and from the French Minister for Higher Education, Research and Innovation (PhD contract to C.P.). This work of the Interdisciplinary Thematic Institute IMCbio+, as part of the ITI 2021-2028 program of the University of Strasbourg, CNRS and Inserm, was supported by IdEx Unistra (ANR-10-IDEX-0002), by SFRI-STRAT'US project (ANR-20-SFRI-0012), EUR IMCBio (IMCBio ANR-17-EURE-0023) under the framework of the French Investments for the Future Program. S.P. also receive funding from ANR (Project RIFLAVIRAM ANR-21-CE35-0018-01). The mass spectrometry instrumentations were funded by the University of Strasbourg, IdEx "Equipement mi-lourd" 2015.

Contributions

MM: Conceptualization, Methodology, Validation, Investigation, Visualization, Writing - Original Draft. **LP**: Investigation. **CP**: Investigation. **ND**: Investigation. **JC**: Investigation, Validation. **PH**: Investigation, Validation. **SP**: Conceptualization., Writing - Review & Editing, Funding acquisition. **EG**: Conceptualization, Methodology, Validation, Investigation, Visualization, Writing - Review & Editing, Funding acquisition, Supervision.

Acknowledgements

857 We thank Béatrice Chane-Woon-Ming for her help with the proteomic analyses, Dr. Paula

858 Lopez for her initial support with plaque assays and Agathe Hunckler for technical assistance.

859 We also thank Dr. Carla Saleh for providing the SINV WT and GFP infectious clones and Dr.

860 Diane Griffin for providing the CAPSID antibody. We thank Dr. Cyril Bourgeois, Dr. Richard

861 Patryk Ngondo and members of the Pfeffer labs for critical reading of the manuscript.

863 Ethics declarations

864 Ethics approval and consent to participate

865 Not applicable.

866 Consent for publication

867 Not applicable.

868 Competing interests

869 The authors declare that there are no conflicts of interest.

870 Availability of data and materials

871 The mass spectrometry proteomics datasets supporting the conclusions of this article are

872 available and have been deposited to the ProteomeXchange Consortium via the PRIDE (40)

873 partner repository with the dataset identifier PXD044867 and 10.6019/PXD044867.

References

1. Tanner NK, Linder P. DExD/H Box RNA Helicases. Molecular Cell. 2001 Aug;8(2):251–62.

2. Linder P, Jankowsky E. From unwinding to clamping — the DEAD box RNA helicase family. Nat Rev Mol Cell Biol. 2011 Aug;12(8):505–16.

3. Meier-Stephenson V, Mrozowich T, Pham M, Patel TR. DEAD-box helicases: the Yin and Yang roles in viral infections. Biotechnology and Genetic Engineering Reviews. 2018 Jan 2;34(1):3–32.

4. Taschuk F, Cherry S. DEAD-Box Helicases: Sensors, Regulators, and Effectors for Antiviral Defense. Viruses. 2020 Feb 5;12(2):181.

5. Bonaventure B, Goujon C. DExH/D-box helicases at the frontline of intrinsic and innate immunity against viral infections. Journal of General Virology [Internet]. 2022 Aug 25 [cited 2022 Oct 25];103(8). Available from: https://www.microbiologyresearch.org/content/journal/jgv/10.1099/jgv.0.001766

6. Liu ZR. p68 RNA Helicase Is an Essential Human Splicing Factor That Acts at the U1 snRNA-5′ Splice Site Duplex. Molecular and Cellular Biology. 2002 Aug 1;22(15):5443–50.

7. Lin C, Yang L, Yang JJ, Huang Y, Liu ZR. ATPase/Helicase Activities of p68 RNA Helicase Are Required for Pre-mRNA Splicing but Not for Assembly of the Spliceosome. Molecular and Cellular Biology. 2005 Sep 1;25(17):7484–93.

8. Xing Z, Ma WK, Tran EJ. The DDX5/Dbp2 subfamily of DEAD‐box RNA helicases. WIREs RNA [Internet]. 2019 Mar [cited 2023 Aug 9];10(2). Available from: https://onlinelibrary.wiley.com/doi/10.1002/wrna.1519

9. Janknecht R. Multi-talented DEAD-box proteins and potential tumor promoters: p68 RNA helicase (DDX5) and its paralog, p72 RNA helicase (DDX17). Am J Transl Res. 2010 May 5;2(3):223–34.

10. Ogilvie VC, Wilson BJ, Nicol SM, Morrice NA, Saunders LR, Barber GN, et al. The highly related DEAD box RNA helicases p68 and p72 exist as heterodimers in cells. Nucleic Acids Res. 2003 Mar 1;31(5):1470–80.

11. Giraud G, Terrone S, Bourgeois CF. Functions of DEAD box RNA helicases DDX5 and DDX17 in chromatin organization and transcriptional regulation. BMB Rep. 2018 Dec 31;51(12):613–22.

12. Dardenne E, Polay Espinoza M, Fattet L, Germann S, Lambert MP, Neil H, et al. RNA Helicases DDX5 and DDX17 Dynamically Orchestrate Transcription, miRNA, and Splicing Programs in Cell Differentiation. Cell Reports. 2014 Jun;7(6):1900–13.

13. Jalal C, Uhlmann-Schiffler H, Stahl H. Redundant role of DEAD box proteins p68 (Ddx5)
and p72/p82 (Ddx17) in ribosome biogenesis and cell proliferation. Nucleic Acids
Research. 2007 May 7;35(11):3590–601.

14. Cheng W, Chen G, Jia H, He X, Jing Z. DDX5 RNA Helicases: Emerging Roles in Viral
Infection. IJMS. 2018 Apr 9;19(4):1122.

15. Hu M, Zheng H, Wu J, Sun Y, Wang T, Chen S. DDX5: an expectable treater for viral
infection- a literature review. Ann Transl Med. 2022 Jun;10(12):712–712.

16. Moy RH, Cole BS, Yasunaga A, Gold B, Shankarling G, Varble A, et al. Stem-Loop
Recognition by DDX17 Facilitates miRNA Processing and Antiviral Defense. Cell. 2014
Aug;158(4):764–77.

17. Zhang H, Xing Z, Mani SKK, Bancel B, Durantel D, Zoulim F, et al. RNA helicase
DEAD box protein 5 regulates Polycomb repressive complex 2/Hox transcript antisense
intergenic RNA function in hepatitis B virus infection and hepatocarcinogenesis.
Hepatology. 2016 Oct;64(4):1033–48.

18. Rahman MM, Bagdassarian E, Ali MAM, McFadden G. Identification of host DEAD-box
RNA helicases that regulate cellular tropism of oncolytic Myxoma virus in human cancer
cells. Sci Rep. 2017 Nov 16;7(1):15710.

19. Zhou X, Luo J, Mills L, Wu S, Pan T, Geng G, et al. DDX5 Facilitates HIV-1 Replication
as a Cellular Co-Factor of Rev. Vartanian JP, editor. PLoS ONE. 2013 May
31;8(5):e65040.

20. Sithole N, Williams CA, Abbink TEM, Lever AML. DDX5 potentiates HIV-1
transcription as a co-factor of Tat. Retrovirology. 2020 Mar 30;17(1):6.

21. Goh PY, Tan YJ, Lim SP, Tan YH, Lim SG, Fuller-Pace F, et al. Cellular RNA Helicase
p68 Relocalization and Interaction with the Hepatitis C Virus (HCV) NS5B Protein and
the Potential Role of p68 in HCV RNA Replication. J Virol. 2004 May 15;78(10):5288–
98.

22. Ríos-Marco P, Romero-López C, Berzal-Herranz A. The cis-acting replication element of
the Hepatitis C virus genome recruits host factors that influence viral replication and
translation. Sci Rep. 2016 May 11;6(1):25729.

23. Chen JY, Chen WN, Poon KMV, Zheng BJ, Lin X, Wang YX, et al. Interaction between
SARS-CoV helicase and a multifunctional cellular protein (Ddx5) revealed by yeast and
mammalian cell two-hybrid systems. Arch Virol. 2009 Mar;154(3):507–12.

24. Li C, Ge L ling, Li P peng, Wang Y, Sun M xia, Huang L, et al. The DEAD-box RNA
helicase DDX5 acts as a positive regulator of Japanese encephalitis virus replication by
binding to viral 3′ UTR. Antiviral Research. 2013 Nov;100(2):487–99.

25. Zhao S, Ge X, Wang X, Liu A, Guo X, Zhou L, et al. The DEAD-box RNA helicase 5
positively regulates the replication of porcine reproductive and respiratory syndrome virus
by interacting with viral Nsp9 in vitro. Virus Research. 2015 Jan;195:217–24.

26. Girardi E, Messmer M, Lopez P, Fender A, Chicher J, Chane-Woon-Ming B, et al.
Proteomics-based determination of double-stranded RNA interactome reveals known and
new factors involved in Sindbis virus infection. RNA. 2023 Mar;29(3):361–75.

27. Weaver SC, Barrett ADT. Transmission cycles, host range, evolution and emergence of
arboviral disease. Nat Rev Microbiol. 2004 Oct;2(10):789–801.

28. Griffin D. Alphaviruses, p 1023–1067. In: Fields virology, 5th ed Lippincott Williams &
Wilkins, Philadelphia, PA. 2007.

29. López P, Girardi E, Mounce BC, Weiss A, Chane-Woon-Ming B, Messmer M, et al.
High-Throughput Fluorescence-Based Screen Identifies the Neuronal MicroRNA miR-
124 as a Positive Regulator of Alphavirus Infection. Williams BRG, editor. J Virol. 2020
Feb 26;94(9):e02145-19, /jvi/94/9/JVI.02145-19.atom.

30. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using
the CRISPR-Cas9 system. Nat Protoc. 2013 Nov;8(11):2281–308.

31. Montavon TC, Baldaccini M, Lefèvre M, Girardi E, Chane-Woon-Ming B, Messmer M,
et al. Human DICER helicase domain recruits PKR and modulates its antiviral activity.
Horner SM, editor. PLoS Pathog. 2021 May 13;17(5):e1009549.

32. Bouyssié D, Hesse AM, Mouton-Barbosa E, Rompais M, Macron C, Carapito C, et al.
Proline: an efficient and user-friendly software suite for large-scale proteomics.
Bioinformatics. 2020 Feb 25;

33. Schonborn J, Oberstraß J, Breyel E, Tittgen J, Schumacher J, Lukacs N. Monoclonal
antibodies to double-stranded RNA as probes of RNA structure in crude nucleic acid
extracts. Nucl Acids Res. 1991;19(11):2993–3000.

34. Bonin M, Oberstraß J, Lukacs N, Ewert K, Oesterschulze E, Kassing R, et al.
Determination of preferential binding sites for anti-dsRNA antibodies on double-stranded
RNA by scanning force microscopy. RNA. 2000 Apr;6(4):563–70.

35. Dhir A, Dhir S, Borowski LS, Jimenez L, Teitell M, Rötig A, et al. Mitochondrial double-
stranded RNA triggers antiviral signalling in humans. Nature. 2018 Aug;560(7717):238–
42.

36. Petitjean O, Girardi E, Ngondo RP, Lupashin V, Pfeffer S. Genome-Wide CRISPR-Cas9
Screen Reveals the Importance of the Heparan Sulfate Pathway and the Conserved
Oligomeric Golgi Complex for Synthetic Double-Stranded RNA Uptake and Sindbis
Virus Infection. Greber UF, editor. mSphere. 2020 Dec 23;5(6):e00914-20.

37. Akhrymuk I, Kulemzin SV, Frolova EI. Evasion of the Innate Immune Response: the Old
World Alphavirus nsP2 Protein Induces Rapid Degradation of Rpb1, a Catalytic Subunit
of RNA Polymerase II. J Virol. 2012 Jul;86(13):7180–91.

38. Lloyd RE. Nuclear proteins hijacked by mammalian cytoplasmic plus strand RNA
viruses. Virology. 2015 May;479–480:457–74.

39. Xu J, Cai Y, Ma Z, Jiang B, Liu W, Cheng J, et al. The RNA helicase DDX5 promotes
viral infection via regulating N6-methyladenosine levels on the DHX58 and NFκB

transcripts to dampen antiviral innate immunity. Horner SM, editor. PLoS Pathog. 2021
Apr 28;17(4):e1009530.

40. Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan
S, et al. The PRIDE database resources in 2022: a hub for mass spectrometry-based
proteomics evidences. Nucleic Acids Research. 2022 Jan 7;50(D1):D543–52.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6