Natural variation of immune epitopes reveals intrabacterial antagonism

Danielle M. Stevens[1,2], Alba Moreno-Pérez[2], Alexandra J. Weisberg[3], Charis Ramsing[2], Judith Fliegmann[4], Ning Zhang[5,6], Melanie Madrigal[2], Gregory Martin[5,6], Adam Steinbrenner[7], Georg Felix[4], and Gitta Coaker[2]*

[1]Integrative Genetics and Genomics Graduate Group, University of California, Davis, Davis CA 95616, USA
[2]Department of Plant Pathology, University of California, Davis, Davis CA 95616, USA
[3]Department of Botany and Plant Pathology, Oregon State University, Corvallis OR, USA
[4]Center for Plant Molecular Biology (ZMBP), University of Tübingen, 72074 Tübingen, Germany
[5]Boyce Thompson Institute for Plant Research, Ithaca NY, USA
[6]Plant Pathology and Plant-Microbe Biology Section, School of Integrative Plant Science, Cornell University, Ithaca NY, USA
[7]University of Washington, Department of Biology, Box 351800, Seattle, WA, 98195, USA

*Corresponding Author: Gitta Coaker
Email: glcoaker@ucdavis.edu

Author Contributions: D.M.S. and G.C. conceived and designed the study. D.M.S. performed most bioinformatic analyses and experiments. A.M.P. performed callose deposition, seedling growth inhibition, MAPK, an independent assessment of ROS, and elf18 antagonism. A.J.W. assisted with computational analyses and performed selection analysis. C.R. analyzed synteny and qPCR in *Clavibacter.* J.F. performed ethylene measurement assays on a subset of csp22 variants under the guidance of G.F. G.F. and A.S. guided biochemical analyses of CSPs. N.Z. developed the *core* lines under the guidance of G.M. M.M. aided in genome curation. D.M.S. and G.C. wrote the manuscript, and all authors were involved in editing.

Keywords: comparative genomics, microbe-associated molecular patterns, natural variation, plant innate immunity, intrabacterial antagonism.

Abstract

Plants and animals detect biomolecules termed Microbe-Associated Molecular Patterns (MAMPs) and induce immunity. Agricultural production is severely impacted by pathogens which can be controlled by transferring immune receptors. However, most studies use a single MAMP epitope and the impact of diverse multi-copy MAMPs on immune induction is unknown. Here we characterized the epitope landscape from five proteinaceous MAMPs across 4,228 plant-associated bacterial genomes. Despite the diversity sampled, natural variation was constrained and experimentally testable. Immune perception in both *Arabidopsis* and tomato depended on both epitope sequence and copy number variation. For example, Elongation Factor Tu is predominantly single copy and 92% of its epitopes are immunogenic. Conversely, 99.9% of bacterial genomes contain multiple Cold Shock Proteins and 46% carry a non-immunogenic form. We uncovered a new mechanism for immune evasion, intrabacterial antagonism, where a non-immunogenic Cold Shock Protein blocks perception of immunogenic forms encoded in the same genome. These data will lay the foundation for immune receptor deployment and engineering based on natural variation.

Significance Statement

Plants recognize pathogens as non-self using innate immune receptors. Receptors on the cell surface can recognize amino acid epitopes present in pathogen proteins. Despite many papers investigating receptor signaling, the vast majority use a single epitope. Here, we analyzed the natural variation across five different epitopes and experimentally characterized their perception in plants. We highlight the importance of analyzing all epitope copies within a pathogen genome. Through genetic and biochemical analyses, we revealed a mechanism for immune evasion, intrabacterial antagonism, where a non-immunogenic epitope blocks perception of immunogenic forms encoded in a single genome. These data can directly inform disease control strategies by enabling prediction of receptor utility and deployment for current and emerging pathogens.

Main Text
Introduction

Plants contain hundreds of innate immune receptors capable of recognizing diverse pathogens. Biotic organisms carry microbe-associated molecular patterns (MAMPs), fragments of larger biomolecules such as proteins, carbohydrates, and lipids, that are recognized by surface-localized pattern recognition receptors (PRRs) (1-3). PRRs include receptor-like kinases (RLKs) and receptor-like proteins (RLPs) that recognize conserved MAMPs or damage-associated molecular patterns, resulting in PRR-triggered immunity (PTI) (3,4). Activation of PTI induces multiple defense responses including: the production of reactive oxygen species (ROS) and ethylene, apoplast alkalization, activation of mitogen-activated protein kinase (MAPK) cascades, transcriptional reprogramming, and callose deposition at the cell wall culminating in disease resistance (4,5).

The most well-studied PRR is FLAGELLIN-SENSING 2 (FLS2), which recognizes the 22-amino acid epitope, flg22, from bacterial flagellin FliC (6-9). Unlike most PRRs, *FLS2* is conserved throughout the plant kingdom (7,9,10). Since the flg22-FLS2 discovery, researchers have uncovered other epitope-receptor pairs restricted to certain plant families. For instance, the Elongation Factor Tu (EF-Tu)-derived epitope elf18 interacts with the EFR receptor, which is present in the Brassicaceae family (11,12). The 22-amino acid epitope of cold shock protein (CSP), csp22, interacts with CORE and a second flagellin epitope flgII-28 interacts with FLAGELLIN-SENSING 3 (FLS3), both solanaceous RLKs (13-16). Finally, the necrosis-and-ethylene inducing peptide 1 (Nep1)-like protein (NLP) epitope nlp20 interacts with RLP23 found in Brassicaceae (17-19). These receptors recognize certain bacterial MAMPs, including those present in pathogens and commensals.

Proteins that carry MAMP sequences are thought to be important for microbial survival and fitness. For example, bacterial flagellin enables bacterial swimming and swarming and is critical for the colonization of certain hosts (20). Different regions of flagellin monomer, FliC, are recognized by different receptors in plants and mammals despite the flagellin apparatus requiring multiple

components (21). CSPs are an ancient protein family first described in *Escherichia coli* with roles in RNA chaperoning in cold environments (22). This can include unwinding RNA, maintaining RNA stability, limiting internal cleavage, and enhancing expression via antitermination. Recent genetic work on other bacterial systems has shown additional post-translational regulatory roles related to virulence, stress tolerance, pili formation, and biofilm development (23-28). CSPs are composed of two critical motifs, RNP-1 and RNP-2, near the N-terminus of the small beta-barrel like structure that binds broadly to nucleic acids (22). While the regulatory function of CSPs appear conserved across bacteria, their targets and sequences are highly variable. NLPs are cross-kingdom phytotoxic virulence factors (17,19). Flagellin, CSPs, and NLPs are considered expendable, though their loss is predicted to confer fitness costs. Conversely, EF-Tu is a highly abundant, essential protein that transports aminoacyl-tRNAs to the ribosome during translation (29). MAMP-encoded gene function and abundance likely have an impact on their evolution in the context of plant immune interactions.

Using flg22 perception as a model, three general epitope outcomes have been described: immune activation, evasion, and antagonism. Many Gram-negative plant pathogens carry immunogenic epitopes, though some can evade perception (7,30-32). MAMP evasion, also known as masking, occurs by accumulating sequence variation that prevents epitope binding (33,34). MAMPs can act antagonistically and block subsequent perception of immunogenic epitopes from other bacteria by binding to the primary receptor and inhibiting proper signaling complex formation (34,35). Sequence variation within the perceived epitope is thought to be constrained by protein functionality, though this has only been studied in the flg22 epitope and FliC protein (8). How polymorphism affects protein function versus immune perception has not been investigated for other MAMP-encoded proteins. In addition, how MAMP copy number variation (CNV) encoded within a single genome impacts immune outcomes remains elusive. Therefore, many questions remain regarding how natural epitope variation interplays with protein function and plant immune perception.

To broadly understand how natural evolution impacts immune outcomes across five different MAMPs, we mined 34,262 MAMP epitopes from 4,228 whole bacterial genomes. Each MAMP displayed substantial copy number and sequence variation. While the theoretical number of MAMP variants is astronomically large, natural variation is constricted, making it experimentally testable to characterize MAMP evolution. We focused on characterizing the immunogenic outcomes of the EF-Tu (elf18) and CSP (csp22) MAMPs for sequence and CNV. Elf18 displays minimal sequence variation and gene expansion, with most variants inducing strong immune responses. In contrast, csp22 displays considerable variation in epitope sequence, CNV, and immune outcomes. Using a combination of phylogenetics, genetics, and biochemistry, we revealed conserved non-immunogenic CSPs in a subset of bacterial genera, some of which antagonize perception of immunogenic forms encoded in the same genome. We then characterized an actinobacterial CSP that acts as an intrabacterial antagonist of the CORE receptor, which enables immune evasion.

Results
Evolutionary trajectories depend on the immunogenic feature and bacterial genera

Here, we sought to assess the presence and diversity of bacterial proteinaceous MAMPs. Across 13 genera, we identified plant-associated bacterial genomes with representing members from alpha-, beta-, and gamma-proteobacteria as well as Gram-positive actinobacteria (Dataset S1). We extracted features from the following proteins and their corresponding MAMP epitopes: bacterial flagellin (flg22 and flgII-28), cold-shock protein (csp22), EF-Tu (elf18), and Nep1-like protein (nlp20). A computational pipeline was developed that extracts peptides using a modified BlastP protocol and local protein alignment, polymorphic ends and off-target correction, and clonality filtering (Fig. 1A). This enabled mining of epitopes in a gene description-dependent and -independent manner allowing for comprehensive genome-derived epitope comparisons (SI Appendix, Table S1).

Differences in sequence and CNV were detected in a MAMP-dependent manner. From the
4,228 genomes, 34,262 epitopes were extracted and their abundance was plotted on a 74 gene
maximum-likelihood tree (Fig. 1B). Notably, there are different patterns of epitope CNV in a lineage-
specific manner (Fig. 1B, SI Appendix, Fig. S1A). The MAMPs elf18, flg22, flgII-28, and nlp20 are
primarily encoded by single copy genes, whereas csp22 variants displayed expansion and are
encoded by 1 to 15 genes (Fig. 1C, SI Appendix, Fig. S1A). Next, we investigated CNV across 13
genera (SI Appendix, Fig. S1A). Strains had an average of four CSPs, with variation in a genera
and species-dependent manner. For example, CSP CNV was increased in *Agrobacterium* and
Streptomyces (average of eight and seven paralogs, respectively) compared to *Dickeya*,
Pectobacterium, *Xanthomonas*, *Ralstonia*, *Curtobacterium*, *Rathayibacter*, and *Clavibacter*
(average of three to five copies) (SI Appendix, Fig. S1A). Although *fliC* is primarily a single copy
gene in 80% (2921/3631) of the genomes, additional copies can be observed in *Erwinia*,
Pectobacterium, and *Agrobacterium* (SI Appendix, Fig. S1A). Similarly, EF-Tu is primarily a single
copy gene in 84% (2825/3346) of analyzed genomes, with additional copies predominantly found
in Gram-negative bacteria apart from *Streptomyces* (Fig. 1B-C, SI Appendix, Fig. S1A). Nlp20 was
only found in 13% (565/4228) of the genomes and predominantly in necrotic bacteria such as
Streptomyces, *Pectobacterium*, and *Dickeya* (SI Appendix, Fig. S3A-B).
We investigated MAMP conservation compared to the 'consensus' sequence commonly
used in the literature for immune assays. When all genera were analyzed together, each MAMP
exhibited different distributions compared to the consensus epitope (Fig. 1D). The cumulative
distributions of flg22 and flgII-28 variants closely mirror each other and are derived from different
regions of the same gene, FliC (Fig. 1D, SI Appendix, Fig. S1B, Fig. S3C). However, a lower
number of flgII-28 epitopes were detected (3661) compared to flg22 (5042), likely due to higher
sequence diversity in the flgII-28 region (S1 Appendix, Fig S2). Elf18 and nlp20 variants are the
most and least conserved, respectively (Fig. 1D, SI Appendix, Fig. S3C).
When analyzing MAMP distributions in individual genera, interesting trends are identified
(SI Appendix, Fig. S1A). Diversification of each FliC epitope in Gram-negative bacteria does not
always mirror each other. *Dickeya* and *Pectobacterium* exhibit similar amino acid similarity and
CNV for both flg22 and flgII-28 (SI Appendix, Fig S1A). However, *Agrobacterium* has a multimodal
distribution for flg22 and unimodal distribution for flgII-28 (SI Appendix, Fig S1A). Many Gram-
positives exhibit similar csp22 diversification, indicating ancient CSP paralog emergence (SI
Appendix, Fig. S1A). We also compared epitope variation independent of a 'consensus' sequence
by calculating amino acid similarity in an all-by-all manner (SI Appendix, Fig. S1B). These results
were complementary to the comparison to the 'consensus' epitope, with different patterns of
diversity for each MAMP. Overall, we observed that epitope abundance and sequence
diversification evolved in both a MAMP-based and genera-derived manner.
Genes that encode MAMP epitopes are postulated to be either essential or conditionally-
essential for bacterial survival. Considering their ancient origin, we assessed the total degree of
epitope variation, which we found was constrained. Except for nlp20, the total number of MAMP
epitopes detected ranged from 3,661 to 20,542 (Fig. 1E). However, the total number of unique
epitope variants was much lower with csp22 exhibiting the highest number (622) and elf18
displaying the lowest number (56, Fig. 1E). When considering the frequency of epitope variants,
there was substantially fewer that occurred at least 10 or 100 times (between 8 and 151, Fig. 1E).
Collectively, these data indicate it is possible to experimentally test the impact of natural variation
on immune outcomes across thousands of bacteria.

Most elf18 variants are immunogenic

To characterize the functional diversity of elf18, 25 variants were synthesized and
assessed for their immunogenicity using ROS, an early, quantitative output of immune induction
(5,35). ROS production by NADPH oxidases is required for biological processes, stress responses,
and the induction of systematic acquired resistance, providing broad-spectrum resistance (36). The
25 elf18 epitopes (100 nM) were screened for their ability to induce ROS on *Arabidopsis thaliana*
Col-0 and the *EFR* receptor mutant line. Water and the *E. coli*-derived 'consensus' elf18 epitope

were used as controls (SI Appendix, Fig. S4A). Of the 25 variants tested, 76% (19/25) displayed
equal ROS output in comparison to the consensus, 16% were weakly immunogenic (4/25), and 8%
(2/25) were non-immunogenic (SI Appendix, Fig. S4A). No measurable ROS was produced in the
efr mutant line (SI Appendix, Fig. S4A). To confirm our ROS screen was effective in classifying
immunogenicity, a blinded, independent evaluation displayed the same conclusions as the initial
screen (SI Appendix, Fig. S4B). Interestingly, all non-immunogenic variants and half of the weakly
immunogenic variants were derived from a second EF-Tu locus in *Streptomyces*, that clustered
separately (Fig. 2A). While all epitopes were derived from annotated EF-Tu loci, the second
Streptomyces EF-Tu copy displayed the most divergent elf18 sequence (Fig. 2A, SI Appendix, Fig.
S4A).
Next, we characterized mid- to late-stage immune outputs including MAPK induction (100
nM), callose deposition (1 uM), and seedling growth inhibition (100 nM) for a subset of epitopes
that exhibited different ROS immunogenicity. All secondary immune outputs were congruent with
each other. All immune outputs by any elf18 epitope variant were abolished in the *efr* mutant (SI
Appendix, Fig. S4C-E). Epitopes EF-77 and EF-81, immunogenic by ROS, inhibited seedling
growth, induced MAPK activation, and displayed high amounts of callose deposition (Fig. 2B-D). In
contrast, the weakly immunogenic and non-immunogenic variants EF-91, 92, 96, and 97 failed to
induce mid- to late-stage immune responses, mirroring water or mock controls (Fig. 2B-D). Elf18
variants EF-91 and EF-96 can uncouple their immune outputs, potentially enabling pathogen
evasion of strong immune response. Similarly, EF-80 is weakly immunogenic by ROS and has
been previously shown to not induce robust callose deposition, a late-stage immune output (5,37).
To understand the relationship between immunogenic outcomes and protein evolution, we
developed a maximum-likelihood tree of EF-Tu and plotted genera, the percent amino acid
similarity of each epitope to the consensus, and immunogenicity by ROS (Fig. 2A). The EF-Tu tree
structure was indicative of taxonomic origin (Fig. 2A). The 25 experimentally tested epitope variants
enabled the determination of immunogenicity of 98.8% (3757/3801) of elf18 variants across 3346
plant-associated bacteria (Fig. 2E). Most elf18 encoded variants (92.2%, 3504/3801) induce strong
immunity when *EFR* was present (Fig. 2E). Furthermore, of the genomes that encode for more than
one EF-Tu locus, only 35 have one immunogenic copy and second non-immunogenic copy (Fig
2F). While immune outcomes of weakly-immunogenic variants from *Streptomyces, Xanthomonas,*
and *Pseudomonas* are comparable, their polymorphisms are unique and their respective EF-Tu
proteins do not cluster phylogenetically, indicating convergent evolution (Fig. 2A, SI Appendix, Fig.
S4A) (37). Across all elf18 variants, polymorphisms were position-dependent, showcasing epitope
variation may be constrained by EF-Tu function (SI Appendix, Fig. S4A).

Diversification of CSPs contributes to differential immunogenic responses

Unlike EF-Tu and elf18, CSP and csp22 exhibited CNV and epitope diversification (Fig.
1C-D, SI Appendix, Fig. S2A). This provided an opportunity to characterize the functional diversity
of a second MAMP with a distinct evolutionary trajectory. Therefore, 65 csp22 epitope variants (200
nM) were screened for their ability to induce ROS on Rio Grande tomatoes. Water and the
Micrococcus luetus-derived 'consensus' csp22 epitope were used as controls (SI Appendix, Fig.
S6A) (13). We tested all variants with one of two independent *core*-deficient lines developed in the
Rio Grande cultivar via CRISPR-cas9 (SI Appendix, Fig. S5 and S6A). Each *core* line failed to
perceive csp22 but could still perceive flg22 and flgII-28 (SI Appendix, Fig. S5B). Across all 65
variants, no ROS was produced in the *core* mutant line, demonstrating *CORE* specificity (SI
Appendix, Fig. S6A). Only 54% (35/65) of csp22 epitope variants displayed equal ROS output in
comparison to the consensus, while 11% (7/65) were weakly immunogenic and 25% (23/65) were
non-immunogenic (SI Appendix, Fig. S6A). A subset of the csp22 variants (25) were also assessed
for their ability to induce ethylene production (1 uM). Though the resolution was qualitative, most
epitope variants were consistent with our ROS screen (SI Appendix, Fig. S7).
Csp22 variation was explored across diverse CSP-domain loci. The CSP domain ranges
from 65-75 amino acids in length and carries RPN-1 and RPN-2 nucleic acid binding motifs (22).
While the CSP domain was conserved, considerable diversity in gene sequence length was found

(SI Appendix, Fig. S6B). We identified CSPs carrying extra domains including calcium binding and
DUF domains (SI Appendix, Fig. S6B). We identified an *Agrobacterium* CSP containing two cold
shock protein domains fused together with a linker (Fig. S6B). Classical CSPs (<75 amino acids,
no additional domains) predominantly encoded immunogenic epitopes. However, several weakly-
or non-immunogenic epitopes were encoded within CSPs with unique domain architecture (SI
Appendix, Fig. S6). Some CSPs were conserved across multiple genera, while others were
restricted to a single genus (SI Appendix, Fig. S6).
To understand the relationship between immunogenic outcomes and protein evolution, we
developed a maximum-likelihood tree of the cold-shock domain and plotted genera, the percent
amino acid similarity of each epitope to the consensus, and immunogenicity by ROS (Fig. 3A).
Unlike EF-Tu, CSP domains display intricate clade structure (Fig. 3A). While over 19,000 cold-
shock domain-containing proteins were extracted, the 65 csp22 variants tested still managed to
capture 75% (14,587/19,423) of the immunogenetic landscape (Fig. 3A-B). Three conserved
clades contained divergent CSPs carrying non-immunogenic csp22 epitopes. Non-immunogenic
CSPs within Clade 1 were conserved across 62% (781/1261) of *Xanthomonas,* Clade 2 was
conserved across 83% (394/475) of *Pectobacterium* and *Erwinia* with a distant relative in
Agrobacterium, and Clade 3 comprised of 95% (580/612) of actinobacteria (Fig. 3C, right
panel).Unlike EF-Tu, almost all genomes carried more than one CSP and of those, 45.7% had at
least one non-immunogenic form (Fig. 3C, left panel).
Observing the differences in immunogenic outcomes and sequence diversity, we then
assessed conservation and selection of specific residues from the conserved non-immunogenic
CSP. Orthologous CSPs were classified using a combination of phylogeny, protein clustering, and
motif classification (SI Appendix, Materials and Methods, Fig. S8A-B). We then used an all-by-all
BlastP approach to confirm orthology (SI Appendix, Fig. S8C). Next, d_N/d_S was calculated for non-
immunogenic and immunogenic CSPs from the same bacterial genera. For non-immunogenic and
immunogenic CSP loci, selection was assessed and codon sites that were considered significantly
negative or positive were based on a set posterior probability (SI Appendix, Fig. S9A). Among
representative immunogenic and non-immunogenic CSP members from the three clades, most
sites are either negatively selected (purifying) or neutral with only a very small number of codons
displaying positive (diversifying) selection (SI Appendix, Fig. S9A). For all tested immunogenic,
weakly-immunogenic, and non-immunogenic epitopes, the RNP-1 motif is highly conserved and
under purifying selection (SI Appendix, Fig. S9A-B). Near the N-terminus (highlighted in grey), a
subset of residues between positions four and nine may be critical for strong immunity based on
the residue changes, R-group chemistry, and immunogenic outcomes (SI Appendix, Fig. S9A). The
epitope signature for each clade is unique and derived from distinct paralogs (Fig. 3A, SI Appendix,
Fig. S9B).

Conserved non-immunogenic cps22 variants reduce CORE immune perception

We observed three non-immunogenic CSP clades across different bacterial genera (Fig.
3A, 3C). These non-immunogenic CSPs may be maintained for their role in inhibiting immune
perception. To test this, we modified our high-throughput ROS assay to assess epitope
antagonism. Briefly, leaf disks were floated on one of two solutions: water or increasing
concentrations of the candidate antagonist. After overnight incubation, all liquid was removed, an
100 nM of agonist was used to elicit immunity, and ROS was measured (SI Appendix, Fig. S10A).
We selected at least one member from each non-immunogenic clade and used consensus csp22
as the agonist. Most non-immunogenic csp22 epitope variants reduced ROS produced by
consensus csp22. The variant conserved in the *Clavibacter* genus, *Cm* csp22-3, displayed the
strongest ROS reduction at concentrations which were 2.5x and 5x the respected agonist
concentration (Fig. 3D).
Clavibacter contains three CSPs with different immunogenic outcomes, immunogenic
cspA1 (*Cm* csp22-1), weakly immunogenic cspA2 (*Cm* csp22-2), and non-immunogenic cspB (*Cm*
csp22-3) (Fig. 3A, SI Appendix, Fig. S6A). We repeated the antagonism assay with agonist *Cm*
csp22-1 alongside two negative controls: elf18 and a scrambled version of *Cm* csp22-3 (designated

s-csp22-3) (SI Appendix, Fig. S10). Antagonism was observed via decreased ROS production after
incubation with *Cm* csp22-3 but not for the other negative control peptides (SI Appendix, Fig. S10).
We additionally tested *Cm* csp22-3 antagonism using BY-2 tobacco cell cultures by measuring
alkalization, an output of membrane depolarization through ion fluxes and inhibition of H+-ATPases
at the plasma membrane (Fig. 3E) (38,39). As expected, *Cm* csp22-1 (10 nM) was immunogenic
and able to induce a pH shift, while *Cm* csp22-3 (10 nM) was not. When 5x *Cm* csp22-3 was added
three minutes before *Cm* csp22-1, it was able to decrease the pH shift, showcasing its antagonistic
nature (Fig. 3E). Collectively, these data indicate that nonimmunogenic CSP epitope variants can
antagonize perception of immunogenic forms encoded in the same genome, which we term
intrabacterial antagonism (Fig. 3F).

We also observed that *Streptomyces* carried two copies of EF-Tu with distinct epitope
variants (Fig. 2A, SI Appendix, Fig. S12A). One non-immunogenic elf18 variant was conserved
across multiple *Streptomyces* genomes (EF-92), with a second non-immunogenic variant found in
one genome (EF-97). Using the same ROS antagonism assay, we tested both non-immunogenic
epitopes against two different agonists, consensus elf18 and the immunogenic *Streptomyces*
variant*,* EF-77. We tested elf12, a truncated version of consensus elf18, which has been previously
reported as a weak antagonist (11). As expected, elf12 was able to weakly reduce ROS induction
for either agonist. However, neither EF-92 nor EF-97 displayed consistent ROS reduction at any
concentration (SI Appendix, Fig. S12B). Therefore, the capacity for intrabacterial antagonism is
MAMP-dependent.

Intrabacterial antagonism of conserved actinobacterial CSP

Antagonism has been predominantly characterized using peptide assays. Therefore, we
investigated intrabacterial antagonism with full-length CSP proteins. The *Clavibacter* genus
comprises eight pathogenic species (40,41). Pathogens in the *Clavibacter* genus predominantly
colonize the xylem vasculature and upon systemic infection, colonize additional tissues (40). First,
conservation and synteny of each *Clavibacter* CSP was analyzed across multiple species. Both
immunogenic *cspA1* and non-immunogenic *cspB* were found in all *Clavibacter* genomes, whereas
the weakly immunogenic *cspA2* was found in 85% (74/87) (Fig. 4A). Across several species, each
gene was highly syntenic (Fig. 4A). Expression of each *CSP* was assessed via qPCR for the tomato
pathogen *C. michiganensis* in xylem-mimicking media and compared to a previously published
housekeeping gene, *bipA,* in TBY rich medium. While all CSPs are expressed, *cspA2* and *cspB*
exhibited higher expression than *cspA1* at both 6 and 24 hours with *cspB* displaying between 2.65
and 3.63x fold change with *cspA1* (Fig. 4B). Furthermore, when recombinant protein CspA1 and
CspB were mixed in varying concentrations, we observed a strong reduction in ROS at 2.5x
concentrations differences, a realistic difference as expression and protein abundance are
correlated in bacteria (Fig. 4C-D) (44).

Next, we wanted to investigate the ability of *C. michiganensis* (*Cm*) cell lysates to suppress
CORE-dependent immunity from another actinobacteria *Streptomyces spp.* ND05-3B (*St*) as well
as the proteobacteria *Pseudomonas syringae* pv. *tomato* DC3000Δ*flaA* (*Pto*). Both *St* and *Pto* carry
immunogenic CSPs and lack *cspB* (SI Appendix, Fig. S6A, S11). Both *Clavibacter* and
Streptomyces are non-flagellated and lack *fliC*; we used a *Pto fliC* deletion mutant (Δ*flaA*). On
Arabidopsis, *Cm* lysates strongly induce ROS while *St* and *Pto* weakly induce ROS in an *EFR*-
dependent manner (SI Appendix, Fig. S11). *Cm* lysates were unable to induce ROS in tomato cv.
Rio Grande or the *core* mutant line (Appendix, Fig. S11). However, *St* and *Pto* were able to induce
ROS in a *CORE*-dependent manner. Furthermore, we able to observe reduced ROS production in
Rio Grande when *Cm* lysates were mixed with either immunogenic *St* or *Pto* lysates at a 2.5 to 5x
concentration (Fig. 4E-F). Collectively, these data demonstrate *Cm* can antagonize *CORE*-
mediated perception of *St* and *Pto.*

We were unable to generate a *Clavibacter* knockout of *cspB* using a variety of approaches
likely due to the region's high GC content, ranging from 73 to 78% (42,43). In order to functionally
assess intrabacterial CspB antagonism *in planta*, we expressed codon-optimized *cspB* in the
tomato pathogen *Pto* DC3000 (Fig. 5). The five CSPs found in the DC3000 genome are

immunogenic and expressed *in planta* (Fig. 5A, SI Appendix, Fig. S5A, S11) (44). CSPs are known
to act as translational chaperones and antiterminators during environmental conditions such as
cold and stress (22). *In vitro* expression of *cspB* was confirmed via western blot (Fig. 5C). *In vitro*
growth was not significantly different between *P. syringae* expressing *cspB* or empty vector,
indicating that *cspB* expression does not grossly impact fitness (Fig. 5D).
In order to assess CspB antagonism, we used DC3000Δ*avrPto*Δ*avrPtoB*Δ*flaA* (referred to
as DC3000ΔΔΔ) to prevent AvrPto/AvrPtoB effector and flg22 recognition (45). After inoculation
with DC3000ΔΔΔ, both *core* mutant lines exhibited increased disease symptoms and bacterial titers
compared to the wild-type Rio Grande control (SI Appendix, Fig. S5C-D). These data demonstrate
that tomato *CORE* impacts *P. syringae* colonization. Next, we characterized bacterial titer
development between three- and five-days post infiltration for DC3000ΔΔΔ expressing *cspB.* In Rio
Grande, DC3000ΔΔΔ expressing *cspB* exhibited higher bacterial titers at four- and five-days post-
inoculation compared to DC3000ΔΔΔ carrying empty vector (Fig. 5E). It is likely we detected a late
increase in bacterial titer because cells require lysis to release CSPs and a minimum concentration
of CspB is required to overcome immunogenic CSPs from *P. syringae.* Importantly, when either
strain was inoculated on the *core* mutant line, there were no significant differences in bacterial titers
between DC3000ΔΔΔ expressing *cspB* and DC3000ΔΔΔ carrying empty vector over the course of
five days (Fig. 5F). Thus, intrabacterial expression of CspB is sufficient to inhibit CORE perception.
Collectively, these data demonstrate that *Clavibacter* CspB is an intrabacterial antagonist.

Discussion
334
Natural epitope variation showcases outcomes which can serve to inform evolution,
mechanistic interactions, and improve engineering approaches. We surveyed thousands of plant-
associated bacteria and found variability in epitope sequence and CNV in a MAMP- and genera-
dependent manner. Since natural variation was relatively low, characterizing the immune
landscape for MAMP perception across thousands of bacteria is possible. Our work showcases the
predictable nature of epitope evolution, enabling future rapid prediction of MAMP immunogenic
outcomes. We uncovered a new mechanism of immune evasion, intrabacterial antagonism, which
demonstrates that all genome-encoded MAMPs need to be considered when characterizing
bacterial-plant outcomes.
Millions of years of evolution led to bacterial divergence into different phyla. Recent work
has assessed sequence variation for the flagellin epitope, flg22, revealing diverse yet constrained
variation (32,35). Colaianni et al. identified 268 unique variants from 627 *Arabidopsis*-associated
genomes. We also identified a similar number of flg22 epitope variants (254) from 4,228 genomes,
highlighting the constrained nature of epitope variation across plant-associated bacteria (35). In
contrast, 1,059 flg22 variants were identified from 1,414 genomes from representative
proteobacterial genomes across diverse lifestyles (32). These studies revealed differential host
immunogenic outcomes based on specific residue positions. In flg22, sites 14 and 15 can have a
direct impact on immune perception but consequently impair motility (8,32,46). Therefore, the
prevalence of some mutations may be constrained due to negative fitness effects.
Epitope diversification may shift when multiple copies are present (35,47). Gene expansion
can impact both protein function and epitope variation. For some *Pseudomonas* strains, FliC-1 is
critical for both plant immune perception and motility, while FliC-2 is poorly expressed and only
weakly induces immune perception (47). For FliC-2, EF-Tu and CSPs, certain residues changes in
their respective epitope sequences were specific to additional copies (Fig. 2, 3) (47). With the
exception of *Streptomyces,* most other bacterial genera carry a single EF-Tu locus and
polymorphisms were highly constrained and predictable in position, likely reflective of the protein's
critical function to bacterial life (SI Appendix, Fig. S4A) (48). In contrast, CSPs are predominantly
multi-copy genes, which likely influences their expanded variation (Fig. 1, 3). Individual CSPs can
have minimal effects on bacterial survival when knocked out, although they can affect pathogen
virulence (22,27,28,49).
Some MAMP-encoded genes have multiple epitopes which can be recognized by different
receptors (50-52). Different epitopes of FliC can be detected from distinct PRRs in *Arabidopsis*,

tomato, rice, and vertebrates (15,21,53). Each plant genome contains hundreds of candidate
PRRs; thus, it is possible that linage specific, convergent evolved receptors is common and the
broad conservation of FLS2 is the exception (54). While *CORE* is restricted to the Solanaceae,
other genotypes in the Vitaceae and Rutaceae families respond to csp22 variants (10,55).
Considering CSP diversity, it is likely other convergently evolved PRRs recognize and respond to
csp22 variants.
MAMP-derived genes have been rarely assessed beyond the first copy. One study
assessed how a small number of *Pseudomonas* strains carrying two *FliC* genes exhibited
differential immune outputs when individually assessed (47). For bacteria carrying one *FliC*,
antagonism has been demonstrated against other genera (35,47). Here, we discovered
actinobacteria contain multiple encoded CSPs where one copy blocks perception of additional
immunogenic forms, thus functioning as intrabacterial antagonist (Fig. 3–5). Gram-positive
pathogens lack the type three secretion system and cannot deliver immune suppressing effectors
directly into host cells (40). Therefore, intrabacterial CSP antagonism provides an additional
mechanism of immune suppression. Intrabacterial MAMP antagonism may also influence
community dynamics or mixed infections.
While total variation differs between MAMPs, csp22 epitopes exhibited the most variation
(SI Appendix, Fig. S2). MAMP epitopes are proposed to undergo an arms race and are presumed
to be under diversifying selection (56,57). However, EF-Tu is under purifying selection (57).
Furthermore, across multiple CSP orthologs present in genera with both immunogenic and non-
immunogenic forms, most codons displayed either neutral or purifying selection (SI Appendix, Fig.
S9A). There are many CSP copies per genome with potentially different functions, frequently
ranging from three to nine, thus the selection pressure on any one CSP may be less compared to
other loci (Fig. 3A, SI Appendix, Fig. S2). Additionally, perception of CSPs is thought to require cell
lysis for perception, which may impact diversifying selection.
By characterizing the consequence of natural epitope variation on immune perception,
alternate immune outcomes have been revealed. While reduction in ROS production by some
epitope variants was measurable, we failed to characterize any elf18 or csp22 epitopes which
displayed statistically significantly higher ROS production compared to the consensus controls (SI
Appendix, Fig. S4A, Fig. S6A). This may reveal that a minimal number of residues at specific
positions are sufficient for strong complex formation and immune induction. Some flg22 variants,
called deviant peptides, can uncouple immune outputs; they are able to induce early-stage ROS
production but fail to induce late-stage immune responses such as SGI (35). This phenomenon of
uncoupled immune outputs was also found for some elf18 variants (Fig. 2). Two weakly
immunogenic variants by ROS, one from *Xanthomonas* and one from *Streptomyces,* were unable
to induce SGI, callose deposition, or MAPK phosphorylation. It is likely MAMPs beyond flg22 and
elf18 may also act as deviants, representing another strategy to reduce strong immune outputs.
We envision two possible explanations for deviant peptide evolution. One, we may be capturing a
snapshot of peptide evolution toward either maskers or antagonists. Two, these deviant peptides
may occupy the space of the receptor complex, limiting robust immune outputs by other
immunogenic peptides. These strategies are not unique to pathogens alone; beneficial microbial
communities have been shown to be enriched in antagonistic FliC variants, which block perception
of FLS2 (35).
A common strategy to confer disease resistance is to transfer receptors between plant
genotypes. *EFR* has been transferred to tomato, citrus, wheat, and apple (58-61). In each case,
EFR transgenic plants are able to significantly restrict pathogen titers and disease progression.
This conclusion is congruent with our elf18 screen since most EF-Tu epitopes (92%) induce strong
immune responses in *A. thaliana* (Fig. 2). In contrast, prevalent non-immunogenic epitopes exist
from FliC and CSPs, of which some block receptor recognition (Fig. 3, SI Appendix, Fig. S2)
(30,32,35,62). Therefore, careful consideration of pathogen epitope variation should be considered
for receptor transfer and subsequent engineering (63). We encourage future work using this dataset
to inform which receptors may be optimal for pathogen control. In concert, large epitope variant
databases may enable new synthetic receptor engineering via protein modeling of receptor

structure. Rational receptor design focused on contact with low-polymorphic ligand residues may
delay evolution of pathogen evasion.

422
423 **Materials and Methods**

424
Bacterial genomes were pulled from NCBI's Refseq and bacterial epitopes were mined via
BlastP, local alignment, and custom R scripts for filtering (Fig. 1A, Dataset S1) (6,11,13,16,17).
Phylogenetic trees for bacteria relatedness were built using GToTree and protein trees were built
using MAFFT for sequence alignment and IQ-TREE tree building. Epitopes of interests that were
assessed *in planta* were chosen based on abundance, epitope sequence, and gene annotation.
ROS, MAPK, seedling growth inhibition, callose deposition, and alkalinization were measured
similarly as previously described with some minor modifications (13, 31, 35). Gene structure was
assessed in *C. michiganensis* using BlastN and Clinker. Measurement of *C. michiganensis* CSP
expression was conducted similarly as previously described (64). Collection of bacterial lysates
were cultured in liquid media or on plates, collected, and processed in lysis buffer via a Emusiveflex-
C3 High-Pressure Homogenizer. Recombinant protein was expressed in *E. coli* and collected using
standard techniques (7). Bacterial titers of *P. syringae* in tomato was conducted similarly as
described before (14). Detailed *Materials and Methods* are found in SI Appendix.

438
439
Acknowledgments

This research was supported by NIH 1R35GM136402, US-Israel BARD IS-5499-22 to G.C.
and USDA-NIFA 2021-67034-35049, UC Davis Henry A. Jastro Scholarship, and UA Local 290
Scholarship to D.M.S. A.M.P. was supported by Fundación Alfonso Martín Escudero Postdoctoral
fellowship. A.J.W. was supported by startup funding from the Department of Botany and Plant
Pathology at Oregon State University. C.R. was supported by UC Davis Dean's Distinguished
Graduate Fellowship. We thank Marta Bjornson and Jeff Dangl for their thoughtful discussions of
this work and Emily Fucarino for plant growth. Jianfeng Xu (Arkansas Biosciences Institute)
provided tobacco BY-2 callus, Imran Khan and Karen MacDonald (UC Davis) and Fumi Fukada
(Okayama University) provided guidance with maintaining cell cultures, Jeff Chang (Oregon State
University) provided the helper plasmid strains pRK2013 and pRK600, Anjali Iyer-Pascuzzi (Purdue
University) provided the flgII-28 peptide, and Christopher Clarke (USDA Agricultural Research
Service, Beltsville, MD) for the *Streptomyces* strain ND05-3B. We thank Nitzan Shabek and Shahid
Siddique (UC Davis) for the usage of the EmusiveFlex and Biotek plate reader, respectively.

454
Data, Materials, and Software Availability

Accessions of genomes used in this study can be found Dataset S1. Plasmids pDSK519-
cspB (#207162), pRSET-Cm-CspA1 (#215398), and pRSET-Cm-CspB (#215400) can be found in
Addgene. All raw data can be found in Zenodo (Doi: 10.5281/zenodo.10724865). Complete details
and code can be found in the following GitHub repository:
DanielleMStevens/Mining_Known_MAMPs. An HTML file with all MAMPs mined can be found in
Zenodo (Doi: 10.5281/zenodo.10724865).

462
463
464 **References**

465 1. D. C. Lewis, D. M. Stevens, H. Little, G. L. Coaker, R. M. Bostock, Overlapping Local and
466   Systemic Defense Induced by an Oomycete Fatty Acid MAMP and Brown Seaweed Extract in
467   Tomato. *Mol. Plant-Microbe Interact.* 36, 359–371 (2023).

468
469   2. B. P. M. Ngou, P. Ding, J. D. G. Jones, Thirty years of resistance: Zig-zag through the plant
470   immune system. *The Plant Cell* 34, 1447–1478 (2022).
471

472   3. D. Ge, I.-C. Yeo, L. Shan, Knowing me, knowing you: Self and non-self recognition in plant
473   immunity. *Essays Biochem* 66, 447–458 (2022).
474

475   4. S. Lolle, D. Stevens, G. Coaker, Plant NLR-triggered immunity: from receptor activation to
476   downstream signaling. *Curr Opin Immunol* 62, 99–105 (2020).
477

478   5. X. Yu, B. Feng, P. He, L. Shan, From Chaos to Harmony: Responses and Signaling Upon
479   Microbial Pattern Recognition. *Annu Rev Phytopathol* 55, 1–29 (2016).
480

481   6. G. Felix, J. D. Duran, S. Volko, T. Boller, Plants have a sensitive perception system for the
482   most conserved domain of bacterial flagellin. *Plant J* 18, 265–276 (1999).
483

484   7. Y. Wei, *et al.,* An immune receptor complex evolved in soybean to perceive a polymorphic
485   bacterial flagellin. *Nat Commun* 11, 3763 (2020).
486

487   8. K. Parys, *et al.,* Signatures of antagonistic pleiotropy in a bacterial flagellin epitope. *Cell Host*
488   *Microbe* 29, 620-634.e9 (2021).
489

490   9. L. Wu, H. Xiao, L. Zhao, Q. Cheng, CRISPR/Cas9-mediated generation of fls2 mutant in
491   Nicotiana benthamiana for investigating the flagellin recognition spectrum of diverse FLS2
492   receptors. *Plant Biotechnol J* 20, 1853–1855 (2022).
493

494 10. J. Trinh, *et al.,* Variation in microbial feature perception in the Rutaceae family with immune
495   receptor conservation in citrus. *Plant Physiol.* (2023) doi:10.1093/plphys/kiad263.
496

497 11. G. Kunze, *et al.,* The N Terminus of Bacterial Elongation Factor Tu Elicits Innate Immunity in
498   Arabidopsis Plants. *Plant Cell Online* 16, 3496–3507 (2004).
499

500 12. C. Zipfel, *et al.,* Perception of the Bacterial PAMP EF-Tu by the Receptor EFR Restricts
501   Agrobacterium-Mediated Transformation. *Cell* 125, 749–760 (2006).
502

503 13. G. Felix, T. Boller, Molecular Sensing of Bacteria in Plants THE HIGHLY CONSERVED RNA-
504   BINDING MOTIF RNP-1 OF BACTERIAL COLD SHOCK PROTEINS IS RECOGNIZED AS
505   AN ELICITOR SIGNAL IN TOBACCO. *J Biol Chem* 278, 6201–6208 (2003).
506

507 14. L. Wang, *et al.,* The pattern-recognition receptor CORE of Solanaceae detects bacterial cold-
508   shock protein. *Nat Plants* 2, 16185 (2016).
509

510 15. S. R. Hind, *et al.,* Tomato receptor FLAGELLIN-SENSING 3 binds flgII-28 and activates the
511   plant immune system. *Nat. Plants* 2, 16128 (2016).
512

513 16. R. Cai, *et al.,* The Plant Pathogen Pseudomonas syringae pv. tomato Is Genetically
514   Monomorphic and under Strong Selection to Evade Tomato Immunity. *PLoS Pathog.* 7,
515   e1002130 (2011).
516

517 17. H. Böhm, *et al.,* A Conserved Peptide Pattern from a Widespread Microbial Virulence Factor
518   Triggers Pattern-Induced Immunity in Arabidopsis. *Plos Pathog* 10, e1004491 (2014).
519

520 18. I. Albert, *et al.,* An RLP23–SOBIR1–BAK1 complex mediates NLP-triggered immunity. *Nat*
521   *Plants* 1, nplants2015140 (2015).

19. M. F. Seidl, G. V. den. Ackerveken, Activity and Phylogenetics of the Broadly Occurring Family of Microbial Nep1-Like Proteins. *Annu Rev Phytopathol* 57, 367–386 (2019).

20. F. F. V. Chevance, K. T. Hughes, Coordinating assembly of a bacterial macromolecular machine. *Nat. Rev. Microbiol.* 6, 455–465 (2008).

21. Y. Rossez, E. B. Wolfson, A. Holmes, D. L. Gally, N. J. Holden, Bacterial Flagella: Twist and Stick, or Dodge across the Kingdoms. *PLoS Pathog.* 11, e1004483 (2015).

22. Y. Zhang, C. A. Gross, Cold Shock Response in Bacteria. *Annu. Rev. Genet.* 55, 1–24 (2021).

23. B. Schmid, *et al.,* Role of Cold Shock Proteins in Growth of Listeria monocytogenes under Cold and Osmotic Stress Conditions. *Appl. Environ. Microbiol.* 75, 1621–1627 (2009).

24. L. P. Burbank, D. C. Stenger, A Temperature-Independent Cold-Shock Protein Homolog Acts as a Virulence Factor in Xylella fastidiosa. *Mol. Plant-Microbe Interact.* 29, 335–344 (2016).

25. A. K. Eshwar, C. Guldimann, A. Oevermann, T. Tasara, Cold-Shock Domain Family Proteins (Csps) Are Involved in Regulation of Virulence, Cellular Aggregation, and Flagella-Based Motility in Listeria monocytogenes. *Front. Cell. Infect. Microbiol.* 7, 453 (2017).

26. A. Catalan-Moreno, *et al.,* One evolutionarily selected amino acid variation is sufficient to provide functional specificity in the cold shock protein paralogs of Staphylococcus aureus. *Mol. Microbiol.* 113, 826–840 (2020).

27. W. Wei, T. Sawyer, L. Burbank, Csp1, a Cold Shock Protein Homolog in Xylella fastidiosa Influences Cell Attachment, Pili Formation, and Gene Expression. *Microbiol. Spectr.* 9, e01591-21 (2021).

28. L. Wu, L. Ma, X. Li, Z. Huang, X. Gao, Contribution of the cold shock protein CspA to virulence in Xanthomonas oryzae pv. oryzae. *Mol. Plant Pathol.* 20, 382–391 (2019).

29. K. L. Harvey, V. M. Jarocki, I. G. Charles, S. P. Djordjevic, The Diverse Functional Roles of Elongation Factor Tu (EF-Tu) in Microbial Pathogenesis. *Front Microbiol* 10, 2351 (2019).

30. S. Wang, *et al.,* Rice OsFLS2-Mediated Perception of Bacterial Flagellins Is Evaded by Xanthomonas oryzae pvs. oryzae and oryzicola. *Mol. Plant* 8, 1024–1037 (2015).

31. Y. Wei, *et al.,* The Ralstonia solanacearum csp22 peptide, but not flagellin-derived peptides, is perceived by plants from the Solanaceae family. *Plant Biotechnol J* 16, 1349–1362 (2018).

32. J. H. T. Cheng, M. Bredow, J. Monaghan, G. C. diCenzo, Proteobacteria Contain Diverse flg22 Epitopes That Elicit Varying Immune Responses in Arabidopsis thaliana. *Mol Plant-microbe Interactions* 34, 504–510 (2021).

33. U. Fürst, *et al.,* Perception of Agrobacterium tumefaciens flagellin by FLS2XL confers resistance to crown gall disease. *Nat. Plants* 6, 22–27 (2020).

34. P. Buscaill, R. A. L. van der. Hoorn, Defeated by the nines: nine extracellular strategies to avoid microbe-associated molecular patterns recognition in plants. *Plant Cell* 33, koab109 (2021).

35. N. R. Colaianni, *et al.,* A complex immune response to flagellin epitope variation in commensal communities. *Cell Host Microbe* 29, 635-649.e9 (2021).

577　36. B. Castro, *et al.,* Stress-induced reactive oxygen species compartmentalization, perception
578　　　and signalling. *Nat. Plants* 7, 403–412 (2021).
579

580　37. F. Yang, G. Li, G. Felix, M. Albert, M. Guo, Engineered Agrobacterium improves
581　　　transformation by mitigating plant immunity detection. *New Phytol* (2022)
582　　　doi:10.1111/nph.18694.
583

584　38. M.M. Atkinson, J.-S. Huang, and K.A. Knopp, The Hypersensitive reaction of tobacco to
585　　　Pseudomonas syringae pv. Pisi: Activation of a plasmalemma K+/H+ Exchange Mechanism.
586　　　*Plant Physiol.* 79, 843-847 (1985).
587

588　39. J. M. Elmore, G. Coaker, The Role of the Plasma Membrane H+-ATPase in Plant–Microbe
589　　　Interactions. *Mol Plant* 4, 416–427 (2011).
590

591　40. S. P. Thapa, *et al.,* The Evolution, Ecology, and Mechanisms of Infection by Gram-Positive,
592　　　Plant-Associated Bacteria. *Annu Rev Phytopathol* 57, 341–365 (2019).
593

594　41. D. Arizala, S. Dobhal, A. M. Alvarez, M. Arif, Elevation of Clavibacter michiganensis subsp.
595　　　californiensis to species level as Clavibacter californiensis sp. nov., merging and re-
596　　　classification of Clavibacter michiganensis subsp. chilensis and Clavibacter michiganensis
597　　　subsp. phaseoli as Clavibacter phaseoli sp. nov. based on complete genome in silico
598　　　analyses. *Int. J. Syst. Evol. Microbiol.* 72, (2022).
599

600　42. L. Chalupowicz, *et al.,* Differential contribution of Clavibacter michiganensis ssp.
601　　　michiganensis virulence factors to systemic and local infection in tomato. *Mol Plant Pathol* 18,
602　　　336–346 (2017).
603

604　43. D. M. Stevens, A. Tang, G. Coaker, A Genetic Toolkit for Investigating Clavibacter Species:
605　　　Markerless Deletion, Permissive Site Identification, and an Integrative Plasmid. *Mol Plant-
606　　　microbe Interactions* 34, 1336–1345 (2021).
607

608　44. T. Nobori, *et al.,* Multidimensional gene regulatory landscape of a bacterial pathogen in
609　　　plants. *Nat. Plants* 6, 883–896 (2020).
610

611　45. B. H. Kvitko, *et al.,* Deletions in the Repertoire of Pseudomonas syringae pv. tomato DC3000
612　　　Type III Secretion Effector Genes Reveal Functional Overlap among Effectors. *Plos Pathog* 5,
613　　　e1000388 (2009).
614

615　46. K. Naito, *et al.,* Amino Acid Sequence of Bacterial Microbe-Associated Molecular Pattern flg22
616　　　Is Required for Virulence. *Mol. Plant-Microbe Interact.* 21, 1165–1174 (2008).
617

618　47. Y. Luo, J. Wang, Y.-L. Gu, L.-Q. Zhang, H.-L. Wei, Duplicated Flagellins in Pseudomonas
619　　　Divergently Contribute to Motility and Plant Immune Elicitation. *Microbiol. Spectr.* 11, e03621-
620　　　22 (2023).
621

622　48. P. J. P. Teixeira, N. R. Colaianni, C. R. Fitzpatrick, J. L. Dangl, Beyond pathogens: microbiota
623　　　interactions with the plant immune system. *Current Opinion in Microbiology* 49, (2019).
624

625　49. Y. Liu, *et al.,* The cold shock family gene cspD3 is involved in the pathogenicity of Ralstonia
626　　　solanacearum CQPS-1 to tobacco. *Microb. Pathog.* 142, 104091 (2020).
627

50. T. Furukawa, H. Inagaki, R. Takai, H. Hirai, F.-S. Che, Two Distinct EF-Tu Epitopes Induce
Immune Responses in Rice and Arabidopsis. *Mol Plant-microbe Interactions* 27, 113–124
(2014).

51. Y. Katsuragi, *et al.,* CD2-1, the C-Terminal Region of Flagellin, Modulates the Induction of
Immune Responses in Rice. *Mol Plant-microbe Interactions* 28, 648–658 (2015).

52. M. de O. Andrade, *et al.,* Suppression of citrus canker disease mediated by flagellin
perception. *Mol. Plant Pathol.* 24, 331–345 (2023).

53. J. Fliegmann, G. Felix, Immunity: Flagellin seen from all sides. *Nat. Plants* 2, 16136 (2016).

54. B. P. M. Ngou, R. Heal, M. Wyler, M. W. Schmid, J. D. G. Jones, Concerted expansion and
contraction of immune receptor gene repertoires in plant genomes. *Nat. plants* 8, 1146–1152
(2022).

55. L. P. Burbank, J. Ochoa, Evidence for Elicitation of an Oxidative Burst in Vitis vinifera by Xylella
fastidiosa Cold Shock Protein Peptide csp20. *PhytoFrontiers* 2, 339–341 (2022).

56. H. C. McCann, H. Nahal, S. Thakur, D. S. Guttman, Identification of innate immunity elicitors
using molecular signatures of natural selection. *Proc National Acad Sci* 109, 4215–4220
(2012).

57. N. Eckshtain-Levi, A. J. Weisberg, B. A. Vinatzer, The population genetic test Tajima's D
identifies genes encoding pathogen-associated molecular patterns and other virulence-related
genes in Ralstonia solanacearum. *Molecular Plant Pathology* 19, (2018).

58. S. Lacombe, *et al.,* Interfamily transfer of a plant pattern-recognition receptor confers broad-
spectrum bacterial resistance. *Nat Biotechnol* 28, 365 (2010).

59. H. Schoonbeek, *et al.,* Arabidopsis EF-Tu receptor enhances bacterial disease resistance in
transgenic wheat. *New Phytol* 206, 606–613 (2015).

60. L. K. Mitre, *et al.,* The Arabidopsis immune receptor EFR increases resistance to the bacterial
pathogens Xanthomonas and Xylella in transgenic sweet orange. *Plant Biotechnol J* 19, 1294–
1296 (2021).

61. S. Piazza, *et al.,* The Arabidopsis pattern recognition receptor EFR enhances fire blight
resistance in apple. *Hortic Res* 8, 204 (2021).

62. C. Pfund, *et al.,* Flagellin Is Not a Major Defense Elicitor in Ralstonia solanacearum Cells or
Extracts Applied to Arabidopsis thaliana. *Mol. Plant-Microbe Interact.* 17, 696–706 (2004).

63. A. Schultink, A. D. Steinbrenner, A playbook for developing disease-resistant crops through
immune receptor identification and transfer. *Curr. Opin. Plant Biol.* 62, 102089 (2021).

64. N. Jiang, *et al.,* Evaluation of suitable reference genes for normalization of quantitative reverse
transcription PCR analyses in Clavibacter michiganensis. *Microbiology open* 8, e928 (2019).

Figures and Legends

Fig. 1. Epitopes from diverse plant-associated bacteria exhibit different evolutionary trajectories.
(*A*) Pipeline to mine for MAMP epitopes from 4,228 plant-associated bacterial genomes (see
Methods Section for details). (*B*) Maximum-likelihood phylogeny built from 74 housekeeping genes
with tips labeled by genera. Each bar below presents the number of epitopes (csp22, elf18, flg22,
flgII-28, nlp20) present in each genome. (*C*) A violin plot showing the number of MAMP-encoded
genes from each genome separated by MAMP type. Tukey's boxplots are plotted on top. The
number of epitopes assessed are listed at the top. (*D*) A violin plot of percent amino acid (AA)

similarity of each epitope variant in comparison to each respective consensus sequence across all
bacteria sampled. Tukey's boxplots are plotted on top. (*E*) Lollipop plot displaying the number of
unique MAMP epitopes and their occurring frequencies.

688
Fig. 2. elf18 exhibits minimal diversification and most variants are immunogenic. (*A*) Maximum-
likelihood phylogenetic tree of EF-Tu. Tips were colored by genera, second bar represents percent
amino acid similarity to consensus, and third bar indicates immunogenicity by ROS from
Supplemental Figure 4. (*B*) *Arabidopsis* seedling growth inhibition after treatment with water or
elf18 peptides (100 nM). One point = 1 plant. Eight plants per biological replicate. A one-way
ANOVA and Tukey's mean comparison was used to determine significance, p<0.0001. (*C*)
Quantified callose deposition in Col-0 after treatment with water and elf18 variants (1 µM). Values
are from one representative experiment which includes an average of at least two images per leaf
from two leaves per plant, three plants per treatment. Statistical analyses were performed as
described in (*B*), p<0.0001. (*D*) Induction of MAPK by elf18 variants (100 nM) in Col-0 at zero- and
15-minutes post infiltration. CBB = protein loading. All experiments were repeated at least three
times with similar results. (*E*) Summary of immunogenic outcomes of EF-Tu-encoded elf18
epitopes from Supplemental Figure 4. (*F*) Summary of immunogenic outcomes with respect to EF-
Tu copy number.

Fig. 3 Convergently evolved non-immunogenic CSPs reduce CORE perception. (*A*) Maximum-likelihood phylogenetic tree of CSP domains. Tips were colored by genera, second bar represents percent amino acid similarity to consensus, and third bar indicates immunogenicity based on ROS experiments in Supplemental Figure 6 (200 nM). (*B*) Summary of immunogenic outcomes of csp22 epitopes tested by ROS. (*C*) Summary of immunogenic outcomes in respect to CSP copy number (Left). Summary of immunogenic outcomes from bacteria present across each non-immunogenic Clade in A and their relatives in the same genera (Right). (*D*) ROS screen for immune antagonism in Rio Grande tomato. Leaf disks were incubated overnight with candidate antagonist followed by elicitation with 200 nM agonist peptide. Control denotes positive control (untreated immunogenic agonist) and concentrations listed are of the candidate non-immunogenic antagonist. Maximum RLU averages were adjusted to a scale of 0 to 100,000 based on the controls (water and untreated agonist). Candidate antagonists, labeled by genera, were selected across the three conserved non-immunogenic-derived CSPs from Figure 5. A one-way ANOVA and Tukey's mean comparison was used to determine significance in respect to the untreated agonist control (denoted *: $p < 0.05$). (*E*) Alkalization of *Nicotiana tabacum* cv. Bright Yellow (BY-2) suspension cells after MAMP treatment

719   (10 nM *Cm* csp22-1; 10 nM *Cm* csp22-3). Antagonism was assessed by first treating with 50 nM
720   *Cm* csp22-3 for three minutes, then treating with 10 nM agonist *Cm* csp22-1. Two cell aliquots were
721   tested per treatment and the experiment was repeated at least two times. (*F*) Diagram of
722   *Clavibacter* CSP intrabacterial antagonism.
723

724
Fig. 4. *Clavibacter* CspB is conserved, expressed, and enables immune evasion of CORE. (*A*)
Conservation and syntenic gene structure of CSPs in representative *Clavibacter* genomes (*C.*
michiganensis: NCPPB382 and Z001; *C. capsici:* PF008; *C. insidiosus:* ATCC10253; *C.*
nebraskensis: NCPPB2581; *C. tessellarius:* ATCC33566 and CFBP8017). (*B*) Expression of CSPs
from *Clavibacter michiganensis* NCPBB382 in xylem-mimicking media compared to expression of
the housekeeping gene, *bipA,* in rich media TBY. Three technical replicates for two independent
biological cultures are plotted. The experiment was repeated three times with similar results. Error
bars = SEM. (*C*) Coomassie stained SDS-PAGE gel of CSP recombinant proteins purified from *E.*
coli. CspA1 is 13 kDa and CspB is 17kDa. (*D*) Maximum ROS production in tomato Rio Grande.
Recombinant CSP proteins were mixed in the indicated concentrations at the same time and
applied to leaf tissue. Concentrations were 100 nM for *Cm* CspA1 and different concentrations of
recombinant non-immunogenic *Cm* CspB (denoted in red font). (*E*) ROS induction of *Streptomyces*
sp. ND05-3B (1 ug/mL, *St*) and *C. michiganensis* NCPPB382 bacterial lysates (2.5 and 5 ug/mL,
Cm) in Rio Grande tomato. (*F*) ROS induction of *Pseudomonas syringae* pv. *tomato* DC3000Δ*flaA*
(1 ug/mL) and *C. michiganensis* NCPPB382 bacterial lysates (2.5 and 5 ug/mL) in Rio Grande
tomato. Lysates were mixed in the indicated concentrations at the same time and applied to leaf
tissue in (*E*) and (*F*). In (*D-F*) the Max. RLUs include an average of four punches per plant, 12

plants per treatment. Error bars = SEM. A one-way ANOVA and Tukey's mean comparison was
used to determine significance, $p < 0.05$.

Fig. 5. Intrabacterial transfer of CspB antagonizes CORE perception in a Gram-negative tomato
pathogen. (*A*) Induction of *Pseudomonas syringae* pv. *tomato* (*Pto*) DC3000 CSPs based on raw
count proteome data from *Nobori et al*., 2020.[44] Variants are labeled by their immunogenicity based
on Figure 3 and Supplemental Figure 6. (*B*) Diagram of *Pto* DC3000 transformed with pDSK519
carrying codon-optimized *cspB* where bacterial titers were tested in tomato. (*C*) *In vitro* expression
of pDSK519 carrying codon-optimized *cspB* (hereafter pDSK519-*cspB*) in *Pto*
DC3000Δ*avrPto*Δ*avrPtoB*Δ*flaA* (ΔΔΔ) (OD$_{600}$ = 1). WT = Wildtype strain DC3000ΔΔΔ. (*D*)
Representative plot of DC3000ΔΔΔ *in vitro* growth with pDSK519-ngfp (ev) and with pDSK519-
cspB in NYGB medium. (*E* and *F*) Bacterial titers of DC3000ΔΔΔ carrying pDSK519-ngfp (ev) or
pDSK519-*cspB* in Rio Grande (RG) and *core* mutants three to five days post infiltration. Two leaves
were sampled per plant with four plants per strain. The experiment was repeated three times. Error
bars = SEM. A one-way ANOVA and Tukey's mean comparison was used to determine
significance, p<0.01.