Assessing the link between tissue elasticity and plant growth

Valentin Laplaud[1], Natalia Demidova[1,2], Elise Muller[1], Stéphanie Drevensek[1], Arezki Boudaoud[1]

[1] LadHyX, Département de Mécanique, CNRS - Ecole Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau Cedex, France

[2] Department of Chemical and Physical Sciences, University of Toronto at Mississauga, L5L1C6 Mississauga, Canada

Corresponding author: Arezki Boudaoud

Email: arezki.boudaoud@polytechnique.edu

Author Contributions:
Conceptualization: VL, EM, SD, AB
Methodology: VL, EM, SD
Software: VL
Validation: VL
Formal Analysis: VL, EM, AB
Investigation: VL, ND
Data Curation: VL
Writing - Original Draft Preparation: VL, AB
Writing - Review & Editing: VL, ND, EM, SD, AB
Visualization: VL
Supervision: AB
Project Administration: AB
Funding Acquisition: AB

Competing Interest Statement: No conflict of interest.

Keywords: growth, mechanical model, microfluidics, Marchantia polymorpha

This PDF file includes:

Abstract, Main Text, Figures 1 to 4, Supplementary figures 1 to 4, Supplementary tables 1 to 2

Abstract:

Multicellular organisms grow and acquire their shapes by controlling the differential expansion and deformations of their cells. Recent research has addressed the role of the mechanical properties of cells and tissues in these processes. In the case of plants, it is generally believed that growth rate is a function of the mechanical stress exerted on the cell wall, the thin polymeric layer surrounding cells, involving an effective viscosity. Nevertheless, recent work has pointed to the relevance of cell wall elasticity in setting growth rate. To extensively assess this relevance, we chose to work on the genetically identical small plants produced in *Marchantia polymorpha* and we developed a microfluidic device to study the early growth and elastic properties of individual plants in a controlled environment with a high-throughput. We characterized the effect of osmotic treatment and of abscisic acid on plant growth and mechanics. Overall, we show that the instantaneous growth rate of individuals is significantly correlated to their bulk elastic modulus, with a correlation coefficient that changes with treatments. Our results are consistent with a framework in which growth rate is determined by elasticity of the wall and a chemical rate associated with remodeling of the cell wall or with changes in its composition. Accordingly, we propose that a tight coupling between mechanics and chemistry of the cell wall is key to setting growth patterns during morphogenesis.

 Main Text

 Introduction

Plant growth and morphogenesis are complex multiscale phenomena where both (bio)chemistry and (bio)mechanics play a fundamental role. While there is little questioning that mechanical

forces are involved in growth (Boudaoud2010, Huang2018, Echevin2019, Trinh2021, Coen2023), the relevance of different mechanical properties in growth is still subject to diverging

views, notably the role of elastic deformations (Coen 2023). Related to this question, modeling attempts fall into different frameworks, most of them considering either stress (force) or strain

(deformation) as the main determinant of growth rate (Smithers2019). Stress is a measure of the physical forces applied on a material, corresponding to a force per unit surface applied to

the sample. The strain is a measure of said material's deformation under stress, corresponding to a relative increase of dimensions of the sample. In the case of an elastic material, the elastic

modulus is the proportionality constant between stress and strain. In this study we provide experimental evidence that elastic strain and material elasticity are key to understanding growth

regulation.

The basic mechanism of plant growth involves the resistance of the cell wall, the polymeric thin layer that surrounds cells, to cell pressure (Boudaoud2010, Huang2018, Echevin2019,

Trinh2021, Coen2023). The high osmolite concentration inside plant cells creates a differential of water chemical potential that drives water into the cell, creating a high mechanical pressure

called turgor that is resisted by the rigid cell wall. The wall's ability to deform under pressure, characterized by its elasticity or its viscosity, the remodeling of its architecture, the change of its

composition, controls the growth dynamics along with the regulation of turgor pressure.

In most modeling frameworks, the strain rate, or expansion rate, of the cell wall depends on stress in the wall. This is typically the case of the Lockhart model (Lockhart1965) where the

growth rate G is proportional to stress σ in excess of a threshold Y (yield stress), the proportionality factor being the extensibility Φ which can be physically interpreted as the inverse

of a viscosity: $G = \Phi(\sigma\text{-}Y)$. This seminal description spawned a whole class of viscoplastic models of plant growth (Veytsman1998, Dumais2006, Dyson & Jensen 2010, Barbacci2013). Some of those models include the elastic modulus in their mechanical parameters but their final formulations make the modulus dispensable to describe steady state growth, ending in description where viscosity (or extensibility) alone is relevant.

Experimental work was conducted to assess the validity of Lockhart's model (Green1974, Cosgrove1987, Proseus1999) as well as to understand further the biological basis of the relevant parameters (McQueen-Mason1995, Cosgrove2000) and get an estimation of their value (Cosgrove1985, Cosgrove1987). However, it is important to note that the results of the aforementioned work could also be interpreted outside of this stress derived framework, because stress is proportional to elastic strain, a measure of the reversible deformation of the sample under stress.

Notably fewer theoretical studies considered elastic strain as the relevant mechanical variable to set growth rate (Rojas2011, Boudon2015). Nevertheless, mechanistic considerations on remodeling of the cell wall led to chemo-mechanical models of cell wall expansion that are compatible with such strain-based model of growth (Rojas2011, Ali2016). In this framework cell wall expansion rate is proportional to elastic strain and so is inversely proportional to elastic modulus (for a given turgor pressure). The proportionality prefactor is a time scale that would be related to the chemistry of the cell wall. Recent experimental work led to progress in the measurement of the cell wall mechanical properties and in particular of its elastic modulus (Milani2013). Some of these studies hint at the importance of elasticity and elastic strain in the regulation of growth in the shoot apical meristem of flowering plants. Organ outgrowth was found to be dependent on cell wall remodeling (pectin methyl-esterification in this case), which was associated with a reduction in elastic modulus (Peaucelle2011, Braybrook2013). Within the shoot apex, domains growing at a slower rate were found to have smaller elastic strain based on osmotic treatments (Kierzkowski2012) and higher elastic modulus based on atomic force microscopy (Peaucelle2011, Milani2011, Milani2014). In the roots of various species the fast

growing regions were measured to have a lower modulus (Petrova2023). Here we investigate experimentally whether stress or strain is more relevant to set growth rate of plants.

The difference in modeling between a viscous approach (based on extensibility) and an elastic approach (based on elastic modulus and a chemistry derived time constant) may seem small as they end up with the same mathematical description of growth dynamics. They however differ greatly in terms of underlying mechanisms. Overall, available experimental results do not discriminate between elastic strain and stress as the main determinant of growth rate. This is in part due to the difficulty of measuring mechanical properties and growth characteristics of the same individuals with a high throughput.

In order to overcome this difficulty, we chose *Marchantia polymorpha* as a model organism, using a microfluidic system to grow small plants in the same conditions. While Marchantia plants provide large numbers of genetically identical progeny by vegetative propagation, microfluidic chips enable the observation of a high number of individuals over time (Busch2012, Sakai2019, Yanagisawa2021). We developed a microfluidic-based approach to characterize mechanical properties, giving us access to growth rate and elastic modulus of the same individual plant with high throughput.

Results

> ➢ High-throughput characterization of growth in controlled conditions

To observe in parallel a large number of individuals in the same conditions, we designed a microfluidic chip with V-shaped traps (Fig. 1-A) to immobilize 50-100 small *Marchantia* plants (gemmae) during 30h. Gemmae were immersed in liquid growth medium (see methods for details), which was continuously renewed by slow injection in the chip during the experiments. We first optimized the collection of gemmae from the mother plant and ensured that growth of the gemmae was robust to the choice of experimental conditions (Fig. S1). We tracked gemmae growth with a microscope (Fig. 1-B) and retrieved their area using image analysis (see methods).

We observed an early phase with almost no growth, followed by a phase of approximately
138   exponential growth (Fig. 1-C). We characterized this dynamics with two parameters T_s and G,
obtained by a delayed exponential fit of each growth curve up to 15h of exponential growth (see
140   methods). T_s is the time between the immersion of plants in solution (a timing called imbibition
for seeds) and the start of exponential growth, G is the increase rate of the exponential.
142   In total we quantified growth for n = 282 gemmae in N = 4 independent experiments (Fig. 1-EF).
The median growth start time T_s over all experiments is 5.20 hours (4.93 to 5.38 hours for
144   individual experiments medians). The variability (expressed as relative average absolute
deviation, see methods for more details on statistical analysis) is of 28% over the complete
146   distribution (from 16% to 40% in individual experiments) showing a highly variable time of growth
start. The growth rate G has a median of 0.78 days^{-1} (0.74 to 0.82 days^{-1}) with a variability of
148   19% (17% to 20%) indicating that the growth rate of gemmae is also quite variable. For both
measurements the variability in single experiments is higher than the difference across
150   experiment medians (~10%), underlining the fact that the overall variability is representative of
the variation between gemmae and not an artifact of data pooling. Such level of variability within
152   a single sample is reminiscent of the germination time of seeds that can be largely variable even
for seeds placed in identical conditions (Bassel2016).

156   ➢ Measurement of bulk elasticity of individual plants

The microfluidic system that we built also allows for rapid (~1min) change of the medium in which
158   the gemmae are developing. We took advantage of this to design a method to measure the
mechanical properties of gemmae. This is based on osmotic steps during which the control
160   growth medium is replaced with a hyperosmotic medium (addition of 100mM mannitol to the
control medium). This drives water out of plants and leads to the contraction of the gemmae by
162   a few percent in area that can be observed and measured through image analysis (Fig.2-AB).
The reverse step is applied 15 minutes later by refilling the chip with the control medium and the
164   inflation of the gemmae is observed and quantified as well. From the curve associated with these
area deformations it is possible to extract a bulk elasticity modulus from the deflation and to

quantify deformation irreversibility (see methods for details). The first step appears to stop gemmae growth during the 15 minutes of treatment but for the return step and inflation we were led to account for growth of the gemmae to get a better measurement of recovery strain (Fig. 2-B and methods for details on the fitting). We can thus quantify the elastic modulus (Fig.2-C) as well as the irreversibility of the deformation between the two steps (Fig.2-D). In this experiment (n = 63) we obtained a median bulk elastic modulus of 16.6 MPa with 14% of variability; this value of modulus is in the lower range of moduli measured for plants (Boudaoud2010). The distribution of deformation irreversibility has a mean of 107.7% (p = 5.0e-10 in comparison to 100% that corresponds to full irreversibility). This slight irreversibility could be ascribed to potential damage to the cell walls caused by the initial step and contraction, or to fast adaptation of the plant during the 15 minutes between the osmotic steps.

To get a better quantification of gemmae elasticity and its variability we examined this experiment (Fig.2-CD) together with the controls of the different treatments discussed hereafter (Fig.4), yielding N = 6 experiments with n = 222 individual plants. The median bulk elastic modulus is 13.95 MPa (11.9 to 16.6 MPa in individual experiments) and the variability is 16% (11% to 14% in individual experiments). Contrary to what we observed for T_s and G, here the global variability comes from the difference between the experiments more than from internal variability. Regarding deformation irreversibility, individual experiments go from 101.7 to 107.7% of irreversibility, and are statistically different from 100% in 3/6 experiments. The standard error of those distributions is between 1.0 and 2.8 % of irreversibility.

Overall these measurements are very consistent and show that our method is robust. The observed low level of deformation irreversibility also supports the definition of a bulk elastic modulus for gemmae.

➢ Relation between growth rate and elasticity

We then asked if there was a relationship between the growth rate of a gemmae and its mechanical properties. The system we designed is well suited to answer such a question, as osmotic steps can be performed on a gemmae population after observation of its growth. It is thus possible to obtain a bulk elastic modulus for gemmae whose growth dynamic has previously

been characterized. When first looking at the correlation between G and E in controlled condition with experiment #2, we found no particular relationship (Fig.S2-B). As the obtained measure of elasticity is from a fast deformation (a few minutes) compared to the time scale over which the area doubling time is computed (several hours), we wanted a more 'instantaneous' characterization of growth as close as possible in time to the measurement of elasticity. For this we measured the instantaneous growth rate $\frac{1}{A}\frac{dA}{dt}$ from the growth curve and took its average value on the last 4 time points (~2 hours, see Fig. S2-A) to define G_{inst} as the 'instantaneous' growth rate just before the step. We found a correlation coefficient of -0.5 (Spearman's rho, p = 1.1e-4) (Fig.2-E) between G_{inst} and the modulus E. We note that this correlation is based on gemmae for which both growth and elasticity are properly quantified, reducing the number of gemmae to n = 55, lower than those observed growing (n = 83, Fig.1-DE) and those characterized mechanically (n=63, Fig.2-C). This is due to the fact that performing osmotic steps implies a fast flow of medium that can push away some gemmae from the observation field.

In the instantaneous case we see a clear decreasing relationship between elastic modulus and the growth rate G_{inst}, in opposition with the absence of correlation with the average growth rate G (Fig.2-E and S2-B). This significant correlation (p<0.05) shows that being softer goes together with faster growth, without indicating any causality of effect between those two parameters. The fact that the correlation is absent when considering average growth rate suggests that the mechanical properties and instantaneous growth rate change together in time during gemmae growth.

> Growth in perturbed conditions

To further probe the relation between modulus and growth rate, we used our system and analysis pipeline to quantify variations due to classical treatments. We considered the change of growth dynamics caused by external physico-chemical constraints or perturbed signaling. For each condition we used two microfluidic chambers in parallel, one control and one perturbed condition, with a common gemmae pool split between them, and repeated the experiment N = 3 times. The

aggregated results are presented in Fig.3 and the detailed results per experiment are available in Fig.S3.

We started with growth in higher osmoticum because it classically reduces growth. Gemmae were observed in control medium permanently supplemented with 100mM of mannitol (Man, Fig.3-A). This treatment significantly reduces the growth rate of gemmae, with a relative change RC = -20% when compared to control (p < 1e-10) and an effect size (see methods) ES = -1.2, as expected. Another effect of growing in a hyper osmotic environment (Fig. 3-AB) is a largely delayed growth start with a relative change RC = 71% (p < 1e-10) and an effect size ES = 3.6 indicating a strong change.

We then performed chemical treatment using a medium complemented with the abscisic acid (ABA) phytohormone. ABA is produced when water is lacking and enables plants to adapt to water stress (Sakata2013, Kuromori2018). The response to ABA was proposed to induce a delay in dormancy exit of Marchantia gemmae based on the observation of rhizoids emergence (Eklund2018). Based on these results, we used a concentration of 3.5µM ABA. Unexpectedly, this treatment (Fig. 3-CD) had a small, non-significant decreasing effect on the growth start time (RC = -10%, ES = -0.3, p = 0.071). However it had a similar effect on growth rate reduction as the hyper-osmotic treatment did (RC = -25%, ES = -1.2, p = 1.0e-6), consistent with ABA treated plants being smaller in Eklund et al.

Overall those perturbations lead to changes that are easily quantifiable with our system, in particular thanks to the facility to observe large numbers of individuals in identical conditions.

➢ Mechanics of plants in perturbed conditions

We then moved on to quantify the effect of those treatments on the mechanical properties of the gemmae (Fig.4). We performed osmotic steps on gemmae that grew 30h either in +100mM hyperosmotic medium or with 3.5µM ABA.

For the gemmae that grew in hyperosmotic medium (Fig.4-AB & S4-AB, N = 3, n = 95 & 80 for control and treated sample, respectively) we observed a decrease of the elastic modulus when

248 compared to their control (RC = -29%, ES = -2.0, p < 1e-10). In addition, we saw a small decrease of deformation irreversibility in the treated sample (RC = -4%, ES = -0.7, p = 2.0e-4)
250 (Fig.S4-A). A similar correlation between instantaneous growth rate and elasticity was observed in both the control and treated cases (Fig.4-B & Table S1).

252 Regarding the effect of ABA treatment (Fig.4-CD & S4-CD, N = 2, n = 64 & 78) we can see a small, non-significant, decrease of elasticity (RC = -6%, ES = -0.2, p = 0.067). The irreversibility
254 is unaffected by ABA treatment (Fig.S4-C). The correlation between elasticity and growth rate is still present for ABA treated gemmae although with a lower correlation coefficient (Fig.4-D and
256 Table S2).

Those results underline the complexity of the link between growth and mechanics. A clear
258 correlation between elastic properties of the gemmae and their instantaneous growth rate was established in reference conditions, and this correlation is conserved with the different
260 treatments. However two treatments that affect the average growth rate in similar ways have a different outcome on changes in mechanical properties. Nevertheless, this complexity can be
262 rationalized, see Discussion.

Discussion

We developed a microfluidic system to study the early growth of *Marchantia* gemmae in a controlled environment and with a high throughput. The environmental control given by microfluidics allowed us to perform osmotic steps on the growing gemmae and thus directly link their mechanical properties to their growth rate. In the course of 12 independent experiments we quantified the growth of 715 individuals and the mechanical properties of 426 individuals. For 284 among those we had access to both measurements, which allowed us to show that there is a correlation between bulk elasticity of the gemmae and their instantaneous growth rate, in line with previous reports of stiffer regions of the shoot apical meristem of tomato growing slower (Kierzkowski2012) and softer regions of roots growing faster (Petrova2023)

When subjecting the gemmae to water-stress by the means of a permanent hyper-osmotic environment, a delay in growth start as well as a reduced growth rate were observed, consistent with classical observations on plant growth in water-stressed conditions (Neumann1988, Slama2015). We also found gemmae to be softer after having grown in hyperosmotic medium. This was observed for the unicellular algae *Chlorella emersonii* (Munns1983) and is in line with early reports of osmotic treatment-induced extensibility variation in maize (Acevedo1971, Wakabayashi1997), where the increase of extensibility could be interpreted as a decrease of elastic modulus. Our results significantly strengthen those observations thanks to a tight control of experimental conditions and a large number of individuals probed. The correlation between instantaneous growth rate and bulk elasticity still holds after growth in a medium with higher osmolarity. In addition, given the observed variability of growth rate in time (see Fig.S2-C for a representative curve), and the variability of mechanical properties across experiments, our results suggest that the regulation of instantaneous growth rate is directly linked to mechanical properties of the plant.

When gemmae were treated with abscisic acid (ABA), we observed a reduction in growth rate, in line with previous observation of ABA treatment on plant growth (Kutschera1986). Surprisingly ABA treatment had no visible effect on the timing of growth start. Because ABA treatment has

previously been reported to delay dormancy exit in *Marchantia polymorpha*, as measured by the first appearance of rhizoids (Eklund2018), this result raises the question of the appropriate criterion for dormancy exit definition. We measured a growth start around 5h while the time of rhizoids appearance is rather at the timescale of a day. We can postulate that the delay in rhizoid appearance would rather be due to a slower growth that would make them reach the gemmae periphery later, than with a delayed dormancy exit. Regarding mechanical properties, growth in ABA medium did not affect the bulk elasticity of gemmae, and conserved the correlation between growth rate and elasticity, albeit with a different correlation coefficient. This shows that even though ABA plays a role in water stress regulation it is not involved in the mechanical response seen when growing gemmae in hyper osmotic medium. If taken together with the observation that ABA reduces cell wall extensibility (Kutschera1986), this suggests that ABA rather affects the dynamics of wall remodeling.

Our results on growth and mechanics taken together with previous work support a modeling framework where the growth rate is given by (1): $G = \chi P/E$ (Rojas2011), with E is the elastic modulus, P the turgor pressure and χ is a rate corresponding to cell wall remodeling. It can also be expressed as (2): $G = \chi \varepsilon$ with $\varepsilon = P/E$ the elastic strain. In (1) χ/E is the analog of the extensibility Φ in Lockhart's model, but with a different mechanistic interpretation. Instead of viewing it as the inverse of a viscosity, as Ortega did when he derived his augmented growth equation from Maxwell linear viscoelastic model (Ortega1985), we can interpret it as the capacity of the wall to remodel itself at a rate χ when under an elastic deformation ε defined by P and E, as the expression in (2) suggests. We would thus expect a growth rate anticorrelated with the elastic modulus, but with a prefactor depending on chemistry of the cell wall and turgor pressure. This result would not be affected by considering a threshold stress or strain for the cell wall to expand.

As stated before this leads to a new interpretation of plant growth modulation by osmotic and ABA treatment: osmotic treatment reduces the modulus E (Acevedo1971) and thus increases extensibility Φ, while ABA treatment reduces remodeling rate so Φ decreases as well

(Kutschera1986), assuming other parameters constant. This framework is also in better
agreement with studies associating elastic strain and growth rate (Kierzkowski2012).

In conclusion, we developed an experimental system to look at growth and mechanics in small plants with the ability to control the growth environment. Through our different treatments we have affected both turgor and chemical regulation and shown that the correlation between elastic modulus and growth rate still holds. Altogether, we believe that our experimental approach and results will help overcome the challenges raised by modeling growth of cells and organisms (Goriely).

Materials and Methods

❖ Plant material

The Tak-1 wild-type accession of *M. polymorpha* was used in this study. Plants were grown on medium (0.2%, Gamborg's B5 media w/ vitamins, Duchefa) mixed with agar (1.2% plant agar, Duchefa). Plant growth takes place in a phytotron (Aralab) with 24h lighting (4600 - 5100 lux) at 22°C. The gemmae used for the experiments are taken from branch order 2 gemmae cups (see Fig. S1-A (iv)) of 3-5 weeks old plants.

❖ Chip fabrication

The chip used to hold the gemmae during the experiment is made of PDMS (Sylgard 184) and molded in a brass mold. The mold was fabricated by micro-drilling (50μm drill) to allow non uniform height of the chamber. The PDMS mix (1:10 ratio) is poured in the mold, degassed for 15 minutes in a vacuum chamber and cured at 70°C for 90 minutes. Once cut out from the mold the chip's inlet and outlet are created using a biopsy puncher. Chip thickness varies from 1mm (inlet and outlet) to 300μm (trap section) and the total chip volume is 500μL. The gradient in

350 thickness allows the gemmae to enter the chip in any orientation without clogging and to be gently guided to a horizontal position in the trap section for observation.

❖ Experiment preparation

The PDMS chip is bonded to a 76x52x1 mm glass slide using a plasma oven (Harrick Plasma,
356 PDC-002). The inlets are connected to glass syringes (10 & 25mL, SGE Trajan) using PTFE tubing of 1.06 mm inner diameter (Adtech) to allow for the passage of gemmae without clogging.
358 The chip is then filled with medium using syringe pumps (Nemesys low pressure module, CETONI). Gemmae are collected in 1.5 mL tube containing medium with biocompatible
360 surfactant (1/1000 dilution, TWEEN 20, Sigma) to avoid aggregation and sticking to the collection tube and tubing during preparation. They are then transferred to the chip manually using a 5 mL
362 syringe (SGE Trajan). The initial medium is washed away after a constant perfusion of medium is put in place with a flow rate of 8.33µL/min (500µL/hour) and is maintained throughout the
364 experiment. For the osmotic step, the flow rate used is 500 µL/min to ensure a fast replacement of the medium in the whole chip (~1 min).

❖ Media and treatments

368 Control medium is 0.2% Gamborg's B5 media w/ vitamins (Duchefa) in water with pH adjusted to 5.8. Hyper-osmotic medium is composed of control medium with the addition of 100mM diluted
370 solid D-Mannitol (Sigma). The ABA medium is composed of control medium with 3.5µM Abscisic acid. The ABA powder (Merck) is first diluted in DMSO (Sigma) at 30mM concentration before
372 being used to make the ABA medium.

❖ Image acquisition

Images are obtained at a 16X magnification using a Zeiss Axiozoom V16 and the Zen 3.3
software. Images are taken every 30 minutes for growth experiments, and every 30 seconds
during osmotic steps. For growth experiments Z-stacks of 3 images, separated by 150 μm, are
acquired at each position. The Z-stacks are reduced to a single Z image using the Zen post
processing function 'Extended depth of focus' and the 'variance' method with default parameters.
For Osmotic steps experiments there is no Z-stack acquisition to have a faster acquisition rate.

❖ Analysis

• Image analysis

Videos (tiff image stacks) of individual gemmae are cropped from the original field of view using
ImageJ. The rest of the analysis is performed using Python, the code is available on demand.
The gemmae are detected in each image in several steps: First the white balance of each video
is adjusted. Second a 3 channel thresholding is applied on the HSV (hue, saturation, value)
representation of the original RGB image. The threshold values are adjusted manually for each
experiment, but are the same for all gemmae analyzed within each experiment. Third
morphological operations (binary closing with 5 μm circular elements, followed by binary opening
with a 30 μm circular element) are used on the created binary mask to recover the shape of the
gemmae. Finally, areas of the gemmae are measured on each image from this reconstituted
binary mask.

• Data fitting

From the image analysis we obtain experimental curves of the gemmae area in time for both the
growth and osmotic step experiments (Fig. 1-C and 2-B). For the growth curve parameter
extraction we fit the experimental data with a delayed exponential of the form:

$$A(t < T_{start}) = A_0$$

$$A(t > T_{start}) = A_0 e^{(t-T_{start})G}$$

It takes into account the delayed start and the exponential nature of growth. As we want to
406 compare growth rate for a similar time period for each gemmae, we use an iterative approach.
First the whole area curve is fitted to obtain a first value of T_s. Then the fit is repeated on a time
408 window stopping 15h after the T_s previously computed. This yields a new value for T_s that can
be used to define a new time window ending 15h after it. This process is iterated until the value
410 obtained for T_s converges with a precision of 0.1 % or it reaches a maximum 10 iteration in which
case the fit is not validated. This ensures a robust definition of T_s and a value of G representative
412 of the initial 15h of growth for all gemmae.

414 For the osmotic steps, each part of the curve (deflation and inflation) is fitted with an evolution
function of the form:

416
$$A\big(t < T_{step}\big) = A_0$$

418
$$A\big(t > T_{step}\big) = (A_0 - A_{eq}).\,e^{\frac{T_{step}-t}{\tau}} + A_{eq} + (t - T_{step}) * B$$

420 Representing an exponential transition between the two equilibrium states with the addition of a
linear component to take into account the growth during the step.
422

The elastic modulus is computed for the hyperosmotic step using the formula:

$$E = \frac{A_0}{|A_0 - A_{eq}|} \delta\pi$$

With $\delta\pi$ the change in external osmotic pressure.

The deformation irreversibility is computed as:

$$Irreversibility = \frac{|A_0 - A_{eq}|_{Second\ step}}{|A_0 - A_{eq}|_{First\ step}}$$

For the 'instantaneous' growth rate during growth we computed

$$GR(t) = \frac{1}{A(t)} \frac{dA}{dt}(t)$$

from the experimental data using a first order discretization:

440
$$\frac{dA}{dt}\left(\frac{t_{n+1}+t_n}{2}\right) = \frac{A(t_{n+1}) - A(t_n)}{t_{n+1} - t_n} \text{ and } \frac{1}{A\left(\frac{t_{n+1}+t_n}{2}\right)} = \frac{2}{A(t_{n+1}) + A(t_n)}$$

442 Where the t_n are the experimental time points.

- Statistical analysis

The variability for data in figures 1 and 2 as stated in the main text are computed as relative average absolute deviation, this means the average absolute deviation (AAD) of the sample divided by its median:

$$AAD = \frac{1}{n} \sum_{i=1}^{n} |x_i - \underline{x}|$$

$$Variability = \frac{AAD}{Median}$$

To assess the significance (p-value) of the irreversibility level when compared to 100% we used a one-sided t-test. For statistical analysis and significance (p-values) when comparing two samples the statistics used are non-parametric given the non-normal nature of data distribution in general. When 'p=0' is displayed it means that $p < 1e{-}10$. The paired sample comparison in figures S1, S2, 3, S3, 4, and S4 are done using Wilcoxon ranksum test. For correlations in figures S1, 2, and 4 we used the Spearman correlation coefficient as we cannot assume linear relationships between the compared quantities *a priori*.

In addition to p-values we chose to display the Relative Change (RC) and the Effect Size (ES) for each sample comparison. The RC is simply the difference between the samples median divided by the control median:

$$RC = \frac{Median_{Treated} - Median_{Ctrl}}{Median_{Ctrl}}$$

We computed a non-parametric ES as the difference between the two median divided by the average absolute deviation of the control condition:

$$ES = \frac{Median_{Treated} - Median_{Ctrl}}{AAD_{Ctrl}}$$

For example, an ES of 0.7 means that the median value of the treated sample is closer to the control median than the average point in the control sample. This gives a way to quantify the difference between two samples while taking into account the variability of the measurements. Computationally we can also understand the ES as the RC between the samples normalized by the variability of the control as defined above:

$$ES = \frac{RC}{Variability_{Ctrl}}$$

Acknowledgments

This work was funded by Agence Nationale de la Recherche (HydroField, grant # ANR-20-CE13-0022-03) and by Ecole polytechnique.

The microfluidic device was developed at the microfabrication facility of LadHyx. The authors are grateful to Caroline Frot for her help as the microfluidic engineer of LadHyX.

References

Boudaoud, A. (2010). An introduction to the mechanics of morphogenesis for plant biologists. *Trends in Plant Science*, *15*(6), 353-360. https://doi.org/10.1016/j.tplants.2010.04.002

488 Huang, C., Wang, Z., Quinn, D., Suresh, S., & Hsia, K. J. (2018). Differential growth and shape formation in plant organs. *Proceedings of the National Academy of Sciences*, *115*(49), 490 12359-12364. https://doi.org/10.1073/pnas.1811296115

492 Echevin, E., Le Gloanec, C., Skowrońska, N., Routier-Kierzkowska, A.-L., Burian, A., & Kierzkowski, D. (2019). Growth and biomechanics of shoot organs. *Journal of Experimental* 494 *Botany*, *70*(14), 3573-3585. https://doi.org/10.1093/jxb/erz205

496 Trinh, D.-C., Alonso-Serra, J., Asaoka, M., Colin, L., Cortes, M., Malivert, A., Takatani, S., Zhao, F., Traas, J., Trehin, C., & Hamant, O. (2021). How Mechanical Forces Shape Plant Organs. 498 *Current Biology*, *31*(3), R143-R159. https://doi.org/10.1016/j.cub.2020.12.001

500 Coen, E., & Cosgrove, D. J. (2023). The mechanics of plant morphogenesis. *Science*, *379*(6631), eade8055. https://doi.org/10.1126/science.ade8055
502

Smithers, E. T., Luo, J., & Dyson, R. J. (2019). Mathematical principles and models of plant 504 growth mechanics : From cell wall dynamics to tissue morphogenesis. *Journal of Experimental Botany*, *70*(14), 3587-3600. https://doi.org/10.1093/jxb/erz253
506

Lockhart, J. A. (1965). An analysis of irreversible plant cell elongation. *Journal of Theoretical* 508 *Biology*, *8*(2), 264-275. https://doi.org/10.1016/0022-5193(65)90077-9

Veytsman, B. A., & Cosgrove, D. J. (1998). A Model of Cell Wall Expansion Based on Thermodynamics of Polymer Networks. *Biophysical Journal*, *75*(5), 2240-2250. 512 https://doi.org/10.1016/S0006-3495(98)77668-4

Dumais, J., Shaw, S. L., Steele, C. R., Long, S. R., & Ray, P. M. (2006). An anisotropic-viscoplastic model of plant cell morphogenesis by tip growth. *The International Journal of* 516 *Developmental Biology*, *50*(2-3), 209-222. https://doi.org/10.1387/ijdb.052066jd

Dyson, R. J., & Jensen, O. E. (2010). A fibre-reinforced fluid model of anisotropic plant cell growth. *Journal of Fluid Mechanics*, *655*, 472-503. https://doi.org/10.1017/S002211201000100X

Barbacci, A., Lahaye, M., & Magnenet, V. (2013). Another Brick in the Cell Wall : Biosynthesis Dependent Growth Model. *PLoS ONE*, *8*(9), e74400. 522   https://doi.org/10.1371/journal.pone.0074400

Green, P. B., & Cummins, W. R. (1974). Growth Rate and Turgor Pressure : Auxin Effect Studies with an Automated Apparatus for Single Coleoptiles. *Plant Physiology*, *54*(6), 863-869. 526   https://doi.org/10.1104/pp.54.6.863

Cosgrove, D. J. (1987). Wall relaxation in growing stems : Comparison of four species and assessment of measurement techniques. *Planta*, *171*(2), 266-278. 530   https://doi.org/10.1007/BF00391104

Proseus, T. E., Ortega, J. K. E., & Boyer, J. S. (1999). Separating Growth from Elastic Deformation during Cell Enlargement1. *Plant Physiology*, *119*(2), 775-784. 534   https://doi.org/10.1104/pp.119.2.775

McQueen-Mason, S. J., & Cosgrove, D. J. (1995). *Expansin Mode of Action on Cell Walls'*. *107*. Cosgrove, D. J. (2000). Loosening of plant cell walls by expansins. *Nature*, *407*(6802), 321-326. 538   https://doi.org/10.1038/35030000

Cosgrove, D. J. (1985) Cell Wall Yield Properties of Growing Tissuel. *Plant Physiol. 78, 347-356*

Boudon, F., Chopard, J., Ali, O., Gilles, B., Hamant, O., Boudaoud, A., Traas, J., & Godin, C. (2015). A Computational Framework for 3D Mechanical Modeling of Plant Morphogenesis with 544   Cellular Resolution. *PLoS Computational Biology*, *11*(1), e1003950. https://doi.org/10.1371/journal.pcbi.1003950

Rojas, E. R., Hotton, S., & Dumais, J. (2011). Chemically Mediated Mechanical Expansion of the Pollen Tube Cell Wall. *Biophysical Journal*, *101*(8), 1844-1853. https://doi.org/10.1016/j.bpj.2011.08.016

Ali, O., & Traas, J. (2016). Force-Driven Polymerization and Turgor-Induced Wall Expansion. *Trends in Plant Science*, *21*(5), 398-409. https://doi.org/10.1016/j.tplants.2016.01.019

Milani, P., Braybrook, S. A., & Boudaoud, A. (2013). Shrinking the hammer : Micromechanical approaches to morphogenesis. *Journal of Experimental Botany*, *64*(15), 4651-4662. https://doi.org/10.1093/jxb/ert169

Peaucelle, A., Braybrook, S. A., Le Guillou, L., Bron, E., Kuhlemeier, C., & Höfte, H. (2011). Pectin-Induced Changes in Cell Wall Mechanics Underlie Organ Initiation in Arabidopsis. *Current Biology*, *21*(20), 1720-1726. https://doi.org/10.1016/j.cub.2011.08.057

Braybrook, S. A., & Peaucelle, A. (2013). Mechano-Chemical Aspects of Organ Formation in Arabidopsis thaliana : The Relationship between Auxin and Pectin. *PLoS ONE*, *8*(3), e57813. https://doi.org/10.1371/journal.pone.0057813

Kierzkowski, D., Nakayama, N., Routier-Kierzkowska, A.-L., Weber, A., Bayer, E., Schorderet, M., Reinhardt, D., Kuhlemeier, C., & Smith, R. S. (2012). Elastic Domains Regulate Growth and Organogenesis in the Plant Shoot Apical Meristem. *Science*, *335*(6072), 1096-1099. https://doi.org/10.1126/science.1213100

Milani, P., Gholamirad, M., Traas, J., Arnéodo, A., Boudaoud, A., Argoul, F., & Hamant, O. (2011). In vivo analysis of local wall stiffness at the shoot apical meristem in Arabidopsis using atomic force microscopy : Measuring wall stiffness in meristems with AFM. *The Plant Journal*, *67*(6), 1116-1123. https://doi.org/10.1111/j.1365-313X.2011.04649.x

Petrova, A., Ageeva, M., & Kozlova, L. (2023). Root growth of monocotyledons and dicotyledons is limited by different tissues. *The Plant Journal*, tpj.16440. https://doi.org/10.1111/tpj.16440

Milani, P., Mirabet, V., Cellier, C., Rozier, F., Hamant, O., Das, P., & Boudaoud, A. (2014). Matching Patterns of Gene Expression to Mechanical Stiffness at Cell Resolution through Quantitative Tandem Epifluorescence and Nanoindentation. *Plant Physiology*, *165*(4), 1399-1408. https://doi.org/10.1104/pp.114.237115

Sakata, Y., Komatsu K., Takezawa D., (2013), ABA as a Universal Plant Hormone., Progress in Botany 75, 57–96

Kuromori, T., Seo, M., & Shinozaki, K. (2018). ABA Transport and Plant Water Stress Responses. *Trends in Plant Science*, *23*(6), 513-522. https://doi.org/10.1016/j.tplants.2018.04.001

Eklund, D. M., Kanei, M., Flores-Sandoval, E., Ishizaki, K., Nishihama, R., Kohchi, T., Lagercrantz, U., Bhalerao, R. P., Sakata, Y., & Bowman, J. L. (2018). An Evolutionarily Conserved Abscisic Acid Signaling Pathway Regulates Dormancy in the Liverwort Marchantia polymorpha. *Current Biology*, *28*(22), 3691-3699.e3. https://doi.org/10.1016/j.cub.2018.10.018

Neumann, P. M., Van Volkenburgh, E., & Cleland, R. E. (1988). Salinity Stress Inhibits Bean Leaf Expansion by Reducing Turgor, Not Wall Extensibility. *Plant Physiology*, *88*(1), 233-237. https://doi.org/10.1104/pp.88.1.233

Slama, I., Abdelly, C., Bouchereau, A., Flowers, T., & Savouré, A. (2015). Diversity, distribution and roles of osmoprotective compounds accumulated in halophytes under abiotic stress. *Annals of Botany*, *115*(3), 433-447. https://doi.org/10.1093/aob/mcu239

Munns, R., Greenway, H., Setter, T. L., & Kuo, J. (1983). Turgor Pressure, Volumetric Elastic Modulus, Osmotic Volume and Ultrastructure of Chlorella emersonii Grown at High and Low External NaCl. *Journal of Experimental Botany*, *34*(2), 144-155. https://doi.org/10.1093/jxb/34.2.144

606   Acevedo, E., Hsiao, T. C., & Henderson, D. W. (1971). Immediate and Subsequent Growth Responses of Maize Leaves to Changes in Water Status. *Plant Physiology*, *48*(5), 631-636.
608   https://doi.org/10.1104/pp.48.5.631

610   Wakabayashi, K., Hoson, T., & Kamisaka, S. (1997). Osmotic Stress-Induced Growth Suppression of Dark-Grown Wheat (Triticum aestivum L.) Coleoptiles. *Plant and Cell*
612   *Physiology*, *38*(3), 297-303. https://doi.org/10.1093/oxfordjournals.pcp.a029166

614   Kutschera, U., & Schopfer, P. (1986). Effect of auxin and abscisic acid on cell wall extensibility in maize coleoptiles. *Planta*, *167*(4), 527-535. https://doi.org/10.1007/BF00391229
616
        Ortega, J. K. E. (s. d.). *Augmented Growth Equation for Cell Wall Expansion'*.
618   Goriely, A., The mathematics and mechanics of biological growth. DOI 10.1007/978-0-387-87710-5. Hardcover ISBN 978-0-387-87709-9.

Figure 1. Growth in a microfluidic chip

622   **A. (i)** Microfluidic chip design with V-shaped traps holding several gemmae (small Marchantia plants) for parallel observation. Scale bar: 5 mm. **(ii)** Zoom on traps. Scale bar: 1mm. **B.** Growth

624   of a gemma in time. The dashed lines serve as a marker of initial size. Scale bar: 250μm. **C.** Areal growth of the gemma represented in B (blue). The magenta dashed line is a delayed

626   exponential fit. Fitted points (15h after growth start) are shown in light blue. **D.** Values of growth start delay for 4 independent experiments. **E.** Values of average growth rate on 15h for the same

628   experiments as (D).

Figure 2. **Measuring mechanical properties with osmotic steps**

A. Illustration of the deformation of a gemmae subjected to an osmotic step (100mM mannitol). We show here a portion of the gemmae; the black line represents the contour of the gemmae at t=0 to underline the deformation. At t = 2 and 3 minutes one can observe a thin white line between the black line and the gemmae, corresponding to the contracted area. **B.** Dynamics of the area of a gemmae when subjected to an osmotic step followed by a return to control medium. The light blue dots are fitted by the blue dashed line and the mauve dots are fitted with the red line to obtain the elastic modulus and irreversibility. **C.** Distributions of bulk moduli. **D.** Distribution of irreversibility levels. The red dashed line illustrate the complete reversibility to which the distribution mean is compared. **E.** Relationship between the bulk moduli of a gemmae and its 'instantaneous' growth rate G_{inst} measured over the two hours preceding the step

 Figure 3. **Growth in perturbed conditions**

A. Area vs. time curves illustrating growth in hyperosmotic medium — reference medium supplemented with 100mM mannitol (shortened Man) — compared to reference medium (control, Ct). Each curve represents average and standard error for all individuals pooled from N = 3 experiments. The area curves are normalized by the initial (t=0) value for better comparison between conditions, see Fig. S3 for detail by experiment **B.** Quantification of growth start time (i) and growth rate (ii) for N=3 experiments in hyperosmotic medium. **C.** Area vs. time curves illustrating growth with abscisic acid (ABA, 3.5µM) treatment compared to reference medium (control, Ct). Each curve represents average and standard error for all individuals pooled from N = 3 experiments. The area curves are normalized by the initial (t=0) value for better comparison between conditions, see Fig. S3 for detail by experiment **D.** Quantification of growth start time (i) and growth rate (ii) for N=3 experiments with ABA.

Figure 4. Mechanics in perturbed conditions

A. Gemmae elastic moduli for control (Ct) and mannitol (Man) treated gemmae. **B.** Correlation between instantaneous growth rate and elasticity for control **(i)** and mannitol treated gemmae **(ii)**. The two stars each indicate one outlier that is out of the x-axis limits. **C.** Gemmae elastic moduli for control (Ct) and abscisic acid (ABA) treated gemmae **D.** Correlation between instantaneous growth rate and elasticity for control **(i)** and ABA treated gemmae **(ii)**.

Figure S1. Growth in controlled conditions
A. Comparison of growth parameters for gemmae coming from old (order 1) or young (order 3) cups on the mother plant. **(i)** Areas at t=0 are similar. **(ii-iii)** Time of growth start is larger and growth rate is lower for older cups. **(iv)** Illustration of the different branch orders of cups. Light yellow are order 1 (oldest) and yellow are order 3 (young) cups. For all experiments other than the one in this figure (S1) gemmae are from order 2 (red) cups. **B.** Comparison of growth parameters for different flow rates in the chip. **(i-ii)** Both time of growth start (T_s) and growth rate (G) are similar between conditions. **C.** Comparison of growth parameters for growth in degassed medium. **(i-ii)** Both time of growth start (T_s) and growth rate (G) are similar between conditions. **D.** Illustration of gemmae position in the chip **(i)** and its correlation with T_s **(ii)** and G **(iii)** for the data in Fig.1-DE. The position of the gemmae in the chip doesn't affect the growth parameters measured (Spearman's rho < 0.15 for both).

A

B

Figure S2. Osmotic steps and growth rates

A. Representative instantaneous growth rate curve. The red dashed line represents the 'instantaneous' growth rate G_{inst} (average over 2h) before step used for correlation with the modulus. **B.** Correlation between the elastic modulus of gemmae and the average growth rate G

<u>**Figure S3.**</u>**Growth in perturbed conditions –Detailed data**

A. Growth curves per experiment for100mM mannitol treatment (Man) and control (Ct) **B.** Quantification of the growth parameters corresponding to curves in (A) **C**. Growth curves per experiment for treatment with abscisic acid (ABA, 3.5μM) and control (Ct) **D.** Quantification of the growth parameters corresponding to curves in (C)

Figure S4. Mechanics in perturbed conditions – supplementary data
A. Irreversibility levels comparison for the pooled data presented in Fig.4-A for control and mannitol treated gemmae. **B.** Detail of individual experiments pooled in Fig.4-A (Ct: control, Man: mannitol-treated). **C.** Irreversibility levels comparison for the pooled data presented in Fig.4-C for control and ABA treated gemmae. **D.** Detail of individual experiments pooled in Fig.4-C (Ct: control, ABA: abscisic acid-treated.

Table S1. Growth rate vs elastic modulus correlation for individual Mannitol experiments

Experiment	#5_Ct	#5_Man	#6_Ct	#6_Man	#7_Ct	#7_Man
# of gemmae	26	9	35	31	12	16
CC	-0,359	-0,517	-0,485	-0,506	-0,650	-0,094
P-value	0,071	0,154	0,003	0,004	0,022	0,729

Table S2. Growth rate vs elastic modulus correlation for individual ABA experiments

Experiment	#10_Ct	#10_ABA	#11_Ct	#11_ABA
# of gemmae	24	33	25	31
CC	-0,689	-0,443	-0,521	-0,018
P-value	0,000	0,010	0,008	0,945